

MONOGRAPHIA CHALCIDITUM.

BY FRANCIS WALKER.

" —— the green myriads in the peopled grass."

VOL. II.

SPECIES COLLECTED BY C. DARWIN, ESQ.

L O N D O N :

HYPPOLITE BAILLIERE, 219, REGENT STREET ;

PARIS :—J. B. BAILLIERE, 13, RUE DE L'ÉCOLE DE MÉDECINE;

LEIPSIG :—J. & G. WEIGEL.

M D C C C X X X I X .

MONOGRAPHIA CHALCIDITUM.

VOL. II.

PART I.—AUSTRALIA.

GENUS.—EURYTOMA, *Illiger.*

Mas.—Corpus convexum, fere cylindricum, subnitens, punctatum, parce hirtum: caput transversum, breve, thorace paullo latius: antennæ 13-articulatæ, nodosæ, thorace non longiores; articulus 1^{us}. fusiformis; 2^{us}. cyathiformis; 3^{us}. et 4^{us}. minimi; 5^{us}. et sequentes nodosi, discreti, verticillato-pilosæ; clava fusiformis, articulo 10^o. duplo longior: thorax postice angustior: prothorax subquadratus, bene determinatus, latitudine vix longior: mesothoracis scutum transversum, sat magnum; parapsidum suturæ bene determinatæ, postice mutuo accedentes; scutellum obconicum: metathorax magnus, declivis: petiolus sat longus: abdomen compressum, altum, nitens, læve, fere glabrum, apice truncatum, thorace multo brevius: pedes hirti, simplices, subæquales; protarso cuique apice spina longior validior, subarcuata: alæ mediocres; hirsuties ad apices densior, sub nervum vix ulla; nervus humeralis gracilis ulnari longior, ulnaris crassus, radialis brevis, cubitalis radiali paullo brevior; stigma mediocre oblongum furcam emittens ad costam propensus; metalis nervus ulnaris humerali brevior, apice setis 3 hamatis armatus.

Fem.—Caput thoracis latitudine: antennæ 12-articulatæ, clavatæ, thorace breviores; articuli 3^o. ad 10^{um}. latescentes, transversi, approximati; clava ovata, articulo 9^o. latior et plus duplo longior: petiolus brevis: abdomen ovatum, acuminatum, thorace paullo brevius.

Sp. 1. Eur. Pidyes. Mas et Fem. *Atra, antennæ nigræ, pedes nigri fulro cincti, alæ limpidae.*

Mas et Fem.—*Atra*: oculi et ocelli rufi: antennæ nigræ: mari pedes fulvi; coxae nigræ; tarsi apice fusci; meso- et metapendulum femora nigra, tibiae piceo-cinetae: *fem.* pedes obscuriores; tibiae plerumque nigræ: alæ limpidae; squamulae piceæ; nervi picei, basi pallidiores. (Corp. long. lin. $\frac{3}{4}$ — $1\frac{1}{4}$; alar. lin. $1\frac{1}{4}$ — $1\frac{2}{3}$.)

Hobart Town, Van Diemen's Land; and in March at King George's Sound, Australia.

Fem.—Corpus convexum, longum, sublineare, fere cylindricum, obscurum, punctatum, parce hirtum: caput transversum, breve, thoracis latitudine: antennæ clavatae, graciles, thorace paullo breviores; articulus 1^{us}. gracilis, sublinearis: petiolus brevis: abdomen longiovatum, subcompressum, nitens, læve, fere glabrum, apice acuminatum, thorace vix brevius; segmenta subæqualia: alæ mediocres.

Sp. 2. Eur. Tellis. Fem. *Atra, antennæ nigræ basi fulvæ aut piceæ, pedes nigri fulro-cincti, tarsi flavi, alæ sublimpidæ.*

Atra, pilis albis vestita: oculi et ocelli rufi: antennæ nigræ; articulus 1^{us}. fulvus, apice piceus; 2^{us}. apice fulvus: oviductus fulvus: pedes nigri; trochanteres fulvi; tibiae piceæ, basi et apice fulvæ; tarsi flavi, apice fusci; protarsi fulvi: alæ sublimpidæ; squamulae piceæ; nervi picei, basi pallidiores. (Corp. long. lin. 1; alar. lin. $1\frac{2}{3}$.)

Var. β.—Antennis articulus 1^{us}. piceus.

Sydney, New South Wales.

Fem.—Corpus convexum, sublineare, punctatum, obscurum, parce hirtum: caput thorace vix latius: antennæ clavatae, robustæ, pilosæ, thorace breviores; articulus 1^{us}. gracilis, sublinearis; 2^{us}. longicyathiformis; 3^{us}. et 4^{us}. brevissimi; 5^{us}. et sequentes breves, usque ad 9^{us}. latescentes; clava longi-conica, articulo 9^o. latior et plus duplo longior: petiolus brevis: abdomen ovatum, subcompressum, nitens, læve, fere glabrum, thorace non longius; segmentum 1^{us}. maximum, 2^{us}. et sequentia brevissima: alæ mediocres.

Sp. 3. Eur. Volux. Fem. *Atra, antennæ nigræ apice piceæ, pedes nigri fulvo-cincti, alæ limpidæ.*

Atra: oculi et ocelli rufi: antennæ nigræ, apice piceæ: pedes nigri; trochanteres picei; genua fulva; tibiæ apice fulvæ; tarsi pallide fulvi, apice fusci; propedum tibiæ fulvæ, tarsi fusci: alæ limpidæ; squamulæ piceæ; nervi proalis picei, metalis fusci. (Corp. long. lin. 1½; alar. lin. 2.)

Hobart Town, Van Diemen's Land; and in March at King George's Sound, Australia.

Mas.—Corpus convexum, breve, punctatum, obscurum, parce hirtum: caput thorace paullo latius: antennæ nodosæ, verticillato-pilosæ, thorace paullo longiores: petiolus longus: abdomen ovatum, compressum, nitens, læve, fere glabrum, thorace multo brevius et angustius: alæ latae.

Fem.—Caput thoracis latitudine: antennæ clavatæ, pilosæ, thorace breviores: abdomen longiovatum, subcompressum, apice acuminatum, thorace non brevius.

Sp. 4. Eur. Aretheas. *Mas et Fem. Atra, antennæ nigræ, pedes nigri, tarsi fulvi, alæ limpidæ.*

Atra: oculi et ocelli rufi: antennæ nigræ: pedes nigri; trochanteres picei; genua fulva; tibiæ apice fulvæ; tarsi fulvi; propedum tibiæ fulvæ, tarsi obscuriores: alæ limpidæ; squamulæ piceæ; nervi proalis picei, metalis fusci. (Corp. long. lin. ¾—1; alar. lin. 1½—1¾.)

March, King George's Sound, Australia.

Mas.—Corpus convexum, sublineare, punctatum, obscurum, parce hirtum: caput thorace vix latius: antennæ graciles, moniliformes, verticillato-pilosæ, thorace paullo longiores: petiolus sat longus: abdomen ovatum, compressum, nitens, læve, fere glabrum, thorace brevius et angustius: alæ latae.

Sp. 5. Eur. Eleuther. *Mas. Atra, antennæ nigræ apice piceæ, pedes fulvi piceo et fusco cincti, alæ limpidæ.*

Mas.—*Atra*: oculi et ocelli rufi: antennæ nigræ, apice piceæ: pedes fulvi; coxæ piceæ; femora picea, apice fulva; meso- et metatibiæ fusco cinctæ; protarsi fusci: alæ limpidæ; squamulæ piceæ; proalis nervi picei, metalis fusci. (Corp. long. lin. ½; alar. lin. 1¾.)

Hobart Town, Van Diemen's Land; and in March at King George's Sound, Australia.

Fem.—Corpus longum, angustum, cylindricum, punctatum, parum nitens, parce hirtum: caput thorace paullo latius: antennæ subclavatæ, graciles, corpore breviores: petiolus brevis: abdomen fusiforme, nitens, læve, fere glabrum, thorace vix brevius: alæ angustæ.

Sp. 6. Eur. Olbus. *Fem.* *Atra, antennæ piceæ basi fulvæ, pedes fulvi, femora piceo cincta, alæ limpidæ.*

Atra: oculi et ocelli rufi: antennæ piceæ; articulus 1^{us}. fulvus: pedes fulvi; coxae piceæ; femora piceo cincta; tarsi apice fusci: alæ limpidæ; squamulæ fulvæ; nervi fulvi. (Corp. long. 1; alar. lin. 1½.)

Sydney, New South Wales.

GENUS.—*ISOSOMA*, *Walker*.

Mas.—Corpus convexum, breve, sublineare, obscurum, punctatum, parce hirtum: caput thorace paullo latius: antennæ moniliformes, sublineares, verticillato-pilosæ, thorace paullo longiores: petiolus sat longus: abdomen ovatum, compressum, nitens, læve, fere glabrum, thorace brevius et angustius: alæ latæ.

Sp. 1. *Iso.* *Ravola.* *Mas.* *Atrum, antennæ nigræ, frons flava, prothoracis latera flavo maculata, pedes nigro-picei, tarsi fulvi, alæ limpidæ.*

Atrum: capitis frons flava: oculi et ocelli rufi: antennæ nigræ: prothoracis latera antica flavo maculata: pedes nigri; trochanteres picei; genua fulva; tibiae piceæ, apice fulvæ; protarsi fulvi, apice fusci; protibiæ fulvæ, fusco semicinctæ: alæ limpidæ; squamulæ piceæ; proalis nervi picei, metalis fusci. (Corp. long. lin. $\frac{4}{3}$; alar. lin 1½.)

Hobart Town, Van Diemen's Land.

Fem.—Corpus convexum, rude punctatum, obscurum, parce hirtum: caput transversum, breve, thorace vix latius; vertex latus; frons excavata, abrupte declivis: oculi mediocres, non extantes; ocelli vertice triangulum fingentes, medius perparum antepositus: antennæ clavatæ, 11-articulatæ, pubescentes, corporis dimidio breviores; articulus 1^{us}. longifusiformis; 2^{us}. et 3^{us}. subrotundi; 4^{us}. longi-cyathiformis; 5^{us}. et sequentes transversi, discreti,

usque ad 8^{um}. curtantes et latescentes; clava longiconica, articulo 8^o. plus duplo longior: thorax longiovatus: prothorax magnus, transversus, subquadratus: mesothoracis scutum longitudine multo latius; parapsidum suturæ optime determinatæ; scutellum subrotundum: metathorax parvus, transversus: pedes validi, simplices, subæquales; tarsi breves: alæ mediocres; nervus humeralis ulnari plus duplo longior, radialis ulnari brevior, cubitalis radiali vix brevior; stigma sat magnum.

Sp. 2. Iso. Oritias. Fem. *Nigro-aeneum*, *antennæ nigræ*, *prothoracis latera fulvo maculata*, *pedes fulvi*, *femora ænea*, *alæ limpidæ*.

Nigræ-aeneum: oculi et ocelli rufi: antennæ nigræ; articuli 1^{us}. et 2^{us}. nigro-ænei: prothorax utrinque fulvo maculatus: pedes fulvi; coxae æneæ; femora ænea, apice fulva; tarsi flavi, apice fusci; metatibiæ fusco cinctæ: alæ limpidæ; squamulæ piceæ; nervi proalis fusci, metalis fulvi. (Corp. long. lin. 1½; alar. lin. 2½.)

March; King George's Sound, Australia.

GENUS.—MEGASTIGMUS, *Dalman*.

Sp. 1. Meg. Drances. Fem. *Fulcus nigro-varius*, *antennæ nigræ basi fulvæ*, *oviductus vaginæ abdomine multo longiores*, *pedes fulvi*, *alæ limpidæ*.

Fulcus, sub-nitens, scite punctatus, fere glaber: capitis vertex niger: os flavum: oculi et ocelli rufi: antennæ nigræ; articulus 1^{us}. fulvus: thorax nigro per medium vittatus: metathorax niger: abdomen læve nitens; segmenta basi nigra: oviductus fulvus; vaginæ nigræ, hirtæ, abdomine multo longiores: pedes fulvi; coxae piceæ; tarsi apice fusci; metafemora subtus piceo maculata: alæ limpidæ; squamulæ fulvæ; nervi fusci, basi pallidiores; stigma piceum, rotundum, sat magnum. (Corp. long. lin. 1½; alar. lin. 2½.)

Hobart Town, Van Diemen's Land.

Sp. 2. Meg. Borus. Fem. *Flarus fulvo et piceo varius*, *antennæ piceæ basi fulvæ*, *metathorax niger*, *oviductus vaginæ abdomine non longiores*, *pedes flavi*, *alæ limpidæ*.

Flavus, sub-nitens, scite punctatus, fere glaber: caput postice nigrum; vertex fulvus: oculi et ocelli rufi: antennæ piceæ; articulus 1st. fulvus: thoracis segmentorum disci fulvi; suturæ piceæ: metathorax niger: abdomen lœve, nitens; discus piceus, fulvo-fasciatus: oviductus fulvus; vaginæ nigræ, hirtæ, abdome non longiores: pedes flavi; tarsi apice fusci: alæ limpidæ; squamulæ flavæ; nervi fusci, basi pallidiores; stigma piceum, rotundum, sat magnum. (Corp. long. lin. 1½; alar. lin. 2½.)

Hobart Town, Van Diemen's Land.

Sp. 3. Meg. Iamenus. Fem. *Fulrus piceo-varius, antennæ piceæ basi fulvæ, oviductus vaginæ corporis longitudine, pedes fulvi, alæ limpidæ.*

Fulvus, subnitens, scite punctatus, fere glaber: oculi et ocelli rufi: antennæ piceæ, basi fulvæ: thoracis segmentorum suturæ piceæ: abdominis discus piceo fasciatus: oviductus fulvus; vaginæ nigræ, hirtæ, corporis longitudine: pedes fulvi; tarsi apice fusci: alæ limpidæ; squamulæ fulvæ; nervi fulvi; stigma piceum, rotundum, magnum. (Corp. long. lin. 1; alar. lin. 1¾.)

Hobart Town, Van Diemen's Land.

GENUS.—CALLIMOME, *Spinola.*

Mas.—Corpus breve, parvum, convexum, sublineare, nitens, scissimè squameum, parce hirtum: caput thorace paullo latius: antennæ filiformes, validæ, hirtæ, thorace non longiores: thorax longiovatus: prothorax brevis: mesothoracis scutum latitudine fere longius; parapsidum suturæ conspicuæ; paraptera et epimera bene determinata; scutellum ovatum: metathorax mediocris: petiolus crassus, brevissimus: abdomen sublineare, fere planum, thorace brevius et angustius: alæ sat latæ.

Sp. 1. Call. Osinius. *Mas. Viridis æneo-varius, antennæ nigræ, pedes fulvo-virides, alæ limpidæ.*

Viridis: oculi et ocelli rufi: antennæ nigræ; articulus 1st. viridis: thorax subtus æneo-varius: abdomen viridi-æneum; discus æneo-cupreus: pedes æneo-virides; tibiae fulvæ, viridi-cinctæ;

tarsi fulvi, apice fuscæ; alæ limpidæ; squamulæ piceæ; nervi fuscæ. (Corp. long. lin. $\frac{3}{4}$; alar. lin. $1\frac{1}{3}$.)

Var. β.—*Thoracis discus viridi-aeneus.*

March; King George's Sound, Australia.

Thorax longiovatus, convexus, nitens, scitissime squameus, parce hirtus: prothorax transversus, parvus, antice angustior: mesothoracis scutum longitudine paullo latius; parapsidum suturæ bene determinatæ; scutellum ovatum: metathorax transversus, mediocris, humilis, postice angustior: pedes simplices, subæquales: alæ amplæ; nervus humeralis ulnari paullo longior, radialis brevissimus, cubitalis adhuc brevior; stigma parvum.

Sp. 2. Call. Vibidia. *Cupreus viridi-varius, pedes flavi, femora viridia, alæ limpidæ.*

Thorax cupreus viridi-varius: prothorax viridis: pedes flavi; coxæ virides; femora viridia; meso- et metatarsi pallide flavi, apice fulvi: alæ limpidæ, fulvo vix tinctæ; squamulæ piceæ; nervi proalis fulvi, metalis flavi. (Alar. lin. $2\frac{1}{2}$.)

Sydney, New South Wales.

GENUS.—PALMON, *Dalmon.*

Fem.—Corpus longum, angustum, convexum, scitissime squameum, parum nitens, parce hirtum: caput transversum, breve, thorace paullo latius; vertex sat latus; frons excavata, abrupte declivis: oculi mediocres, non extantes: ocelli vertice triangulum fingentes: antennæ clavatae, crassæ, apice quasi truncatae, thorace non longiores; articulus 1st. sublinearis; 2nd. longicyathiformis; 3rd. et 4th. brevissimi; 5th. et sequentes breves, transversi, usque ad clavam latescentes: thorax sublinearis: prothorax transversus, subquadratus, sat magnus: mesothoracis parapsidum suturæ non bene determinatæ: petiolus brevis: abdomen compressum, laeve, nitens, thorace multo brevius et angustius: oviductus vaginæ pubescentes, corpore longiores: metapedum coxæ magnæ, femora maxima ovata subtus dentata, tibiæ arcuatæ: alæ mediocres.

Sp. 1. Pal. Olenus. Fem. *Aeneo-viridis, antennæ fulvæ, abdomen subtus fulvum, pedes fusco-fulti, femora viridia, alæ limpidæ.*

Æneo-viridis : oculi et ocelli rufi : antennæ fulvæ : abdomen cyaneo et cupreo-varium, subtus fulvum : oviductus rufus ; vaginæ nigræ : pedes fulvi ; coxae virides ; femora viridia ; tibiae fuscae : alæ limpidæ. (Corp. long. lin. 2 ; alar. lin. 2½.)

Sydney, New South Wales.

GENUS.—HOCKERIA, *De Laporte.*

Mas.—Corpus sublineare, convexum, rude punctatum, obscurum, parce hirtum : caput thorace vix latius ; frons profunde excavata : antennæ filiformes, corpore paullo breviores ; pedicellus longus ; articulus 1^{us}. longissimus, arcuatus ; 2^{us}. et sequentes lineares, arcte applicati : prothorax transversus, sat magnus : mesothoracis scutum longitudine latius ; parapsidum suturæ bene determinatæ ; scutellum obconicum : metathorax parvus, declivis : petiolus brevissimus : abdomen ovatum, nitens, læve, basi glabrum, thorace brevius et angustius : metapedum coxae magnæ, femora ovata maxima, tibiae arcuatæ : alæ mediocres ; proalis nervus humeralis ulnari triplo longior, radialis longus, cubitalis brevissimus.

Sp. 1. Hoc. Proxenus. *Mas. Atra, tarsi picei aut ferruginei, proalæ fusco semifasciatæ.*

Ater : oculi et ocelli picei : antennæ nigræ : pedes nigri ; genua picea ; tarsi picei aut ferruginei : alæ subsuscæ ; squamulæ piceæ ; nervi proalis picei, metalis fulvi ; proalæ fusco semifasciatæ. (Corp. long. lin. 1½ ; alar. lin. 2.)

Hobart Town, Van Diemen's Land ; and Sydney, New South Wales.

GENUS.—EUCHARIS, *Latreille.*

Mas.—Corpus convexum, nitens, glabrum, rude et profunde punctatum : caput transversum, brevissimum, thorace paullo latius ; vertex latus, non impressus ; frons abrupte declivis, ad os angusta et fere lævis : oculi parvi, subrotundi : ocelli remoti, medius perparum antepositus : antennæ 13-articulatæ, subfiliformes, biramosæ, graciles, dense pubescentes, thorace vix longiores ; articulus 1^{us}. longus, gracilis, linearis ; 2^{us}. brevi-cyathiformis ; 3^{us}. obconicus, 1°. longior ; 5^{us}. et sequentium omnis usque ad

12^{um}. breves, sublineares, ramulum longum subclavatum utrinque emittentes: thorax quasi obconicus, latus, altus, gibbus: prothorax minimus, supra non conspicuus: mesothoracis scutum magnum, transversum; parapsidum suturæ bene determinatae, postice mutuo accedentes; paraptera et epimera conspicua, hæ inter scutum et scutellum occurrentia; scutellum mediocre, subrotundum: metathorax magnus, abrupte declivis: petiolus longus, gracillimus, nitens, lævis: abdomen fusiforme, perangustum, convexum, compressum, nitens, læve, subtus carinatum; segmentum unicum dorsum occupans et ventralia fere obtegens: pedes gracieles, simplices, subæquales; coxae parvae; tarsis articuli 1°. ad 4^{um}. curtantes, 5^{us} 4°. paullo longior; unguis et pulvilli minuti: alæ amplæ; nervus ulnaris humerali multo brevior, radialis sat longus, cubitalis brevissimus stigma fingens punctiforme; metalis nervus unicus simplex costæ dimidio longior.

Sp. 1. Euch. Volusus. Mas. *Cuprea viridi varia, abdomen viridi-æneum, antennæ nigræ, pedes ferruginei, femora viridia, alæ limpidæ.*

Mas.—*Cuprea*: capitis vertex viridis: oculi et ocelli obscure rufi: antennæ nigræ; articuli 1^{us}. et 2^{us}. nigro-cuprei: thorax utrinque et subtus viridi-varius: abdomen viridi-æneum: pedes ferruginei; coxae virides; femora viridia, apice ferruginea; protarsi obscuriores; unguis et pulvilli fusci: alæ limpidæ; squamulæ ferrugineæ; nervi fulvi; stigma fuscum. (Corp. long. lin. 2½; alar. lin. 3½.)

"Antennæ 13-jointed, 1st. joint linear, 2^d. very short, cup-shaped, 3^d. obconic, longer than the 1st., 4th. very short, producing above towards the outside a long arched subclavate pilose branch, the following joints to the 12th. inclusive becoming gradually longer and producing a similar branch from the apex, and the exterior ones also a shorter branch towards the inner side; the last five branches of this series are sufficiently distinct, but that of the 8th. joint is reduced to a short tooth. The 13th. joint is as long as the longer branch of the preceding, and like it, subclavate and pilose."

—Haliday.

March; King George's Sound, Australia.

Mas.—Corpus convexum, nitens, scabre punctatum, pubescens: caput mediocre, transversum, brevissimum, thorace vix latius; vertex latus, non impressus; frons abrupte declivis, ad os angus-

tior : oculi parvi, subrotundi : ocelli approximati, vertice triangulum fingentes : thorax ovatus, gibbus : prothorax brevissimus, supra non conspicuus : mesothoracis scutum latitudine longius : parapsidum suturæ bene determinatae ; paraptera et epimera magna ; scutellum breve, prominulum, apex abrupte declivis : metathorax magnus, declivis : petiolus longus, gracilis : abdomen ovatum, gibbum, nitens, lœve, glabrum, thorace brevius et angustius, segmento unico obtectum : pedes graciles, simplices, pubescentes, subæquales ; coxæ parvæ ; tarsis articuli 1°. ad 4^{um}. curtantes, 5^{um}. 4°. longior ; unguis et pulvilli minuti : alæ amplæ ; nervus ulnaris humerali paullo brevior, radialis longus, cubitalis brevissimus ; nervus unicus simplex costæ dimidio longior.

Sp. 2. Euch. Fausta. Mas. *Viridis cupreo et cyaneo varia*, *pedes fulvi, femora viridia, alæ limpidæ*.

Viridis, cupreo-varia : capitis frons cyaneo-viridis : oculi et ocelli obscure rufi : thorax subtus et postice cyaneo-viridis : abdomen cupreum : pedes fulvi ; coxae virides ; femora viridia, apice fulva ; tarsi apice fusci : alæ limpidæ ; squamulae piceæ ; nervi fulvi, apice obscuriores. (Corp. long. lin. 1 $\frac{1}{2}$; alar. lin. 2 $\frac{3}{4}$.)

Hobart Town, Van Diemen's Land.

Mas.—Corpus convexum, parum nitens, rude punctatum, parce pubescens : caput sat magnum, breve, transversum, thorace latius ; vertex latus, non impressus ; frons abrupte declivis : oculi parvi, subrotundi, extantes : ocelli approximati, vertice triangulum fingentes : antennæ 12-articulatae, filiformes, pubescentes, thorace paullo longiores ; articulus 1^{um}. longus, gracilis, sublinearis ; 2^{um}. cyathiformis ; 3^{um}. longus ; 4^{um}. et sequentes breves, lineares, usque ad 9^{um}. curtantes ; 10^{um}. 11^{um}. et 12^{um}. arcte applicati clavam fingentes fusiformem 9°. duplo longiorem : thorax ovatus : segmentis suturæ bene determinatae : prothorax brevissimus, supra vix conspicuus : mesothoracis scutum latitudine latius ; parapsides conspicuae, extantes ; scutellum ovatum, valde convexum : metathorax mediocris, declivis : petiolus perlongus, sat validus : abdomen ovatum, convexum, nitens, lœve, glabrum, thorace brevius, apice non truncatum ; segmentum 1^{um}. magnum, dorsi plus dimidium occupans ; 2^{um}. et sequentia brevia : ventralia vix conspicua : pedes simplices, subæquales ; coxæ mediocres ; femora gracilia ; tibiæ rectæ ; tarsis articuli 1°.

ad 4^{um}. curtantes, 5^{us}. 4°. paullo longior : ungues et pulvilli minuti : alæ amplæ ; ulnaris humerali paullo brevior, radialis brevis, cubitalis brevissimus stigma fingens punctiforme.

Sp. 3. Euch. Valgius. Mas. *Viridis, antennæ nigræ, abdomen nigro-viride, pedes fulvi, femora viridia, alæ subfuscæ.*

Viridis : oculi et ocelli obscure rufi : antennæ nigræ ; articuli 1^{us}. et 2^{us}. virides : abdomen nigro-viride : pedes fulvi ; coxae virides ; femora viridia, apice fulva ; metatibiae pallide fuscæ, apice et basi fulvæ ; tarsi flavi, apice fusti ; protarsi obscuriores : alæ subfuscæ ; squamulae virides ; nervi fusti. (Corp. long. lin. 1 $\frac{1}{2}$. ; alar. lin. 2 $\frac{1}{2}$.)

Sydney, New South Wales.

Fem.—Corpus convexum, nitens, rude punctatum, parce pubescens : caput sat magnum, breve, transversum, thorace latius ; vertex latus, non impressus ; frons abrupte declivis : oculi parvi, subrotundi, extantes : ocelli approximati, vertice in triangulum dispositi : antennæ 12-articulatae, extrosum crassiores, pubescentes, thorace fere breviores ; articulus 1^{us}. longus, gracilis, sublinearis ; 2^{us}. cyathiformis ; 3^{us}. et sequentes transversi, usque ad 9^{um}. curtantes ; 10^{us}. 11^{us}. et 12^{us}. arcte applicati, clavam fingentes fusiformem 9°. duplo longiorum : thorax ovatus : segmentis suturæ bene determinatae : prothorax brevissimus, supra vix conspicuus : mesothoracis scutum longitudine latius ; parapsides conspicuae, extantes ; scutellum ovatum, valde convexum : metathorax mediocris, declivis : petiolus longus, sat validus : abdomen ovatum, convexum, nitens, lœve, glabrum, thorace brevius, apice non truncatum ; segmentum 1^{um}. magnum, dorsi plus dimidium occupans ; 2^{um}. et sequentia brevia : ventralia vix conspicua : pedes simplices, subæquales ; coxae mediocres ; femora gracilia ; tibiae rectæ ; tarsis articuli 1°. ad 4^{um}. curtantes, 5^{us}. 4°. paullo longior : ungues et pulvilli minuti : alæ amplæ, ulnaris humerali paullo brevior, radialis brevis, cubitalis brevissimus stigma fingens punctiforme.

Sp. 4. Euch. Theocles. Fem. *Viridis, antennæ ferrugineæ, abdomen nigro-viride, pedes fulvi, femora viridia, alæ limpidæ.*

Viridis : oculi et ocelli obscure rufi : antennæ ferrugineæ : abdomen nigro-viride : pedes fulvi ; coxae virides ; femora viridia, apice et

basi fulva; tarsi apice fusci; protarsi pallide fusci: alæ limpidæ; squamulæ virides; nervi fusci. (Corp. long. lin. 1½.; alar. lin. 2½.)

Sydney, New South Wales.

Mas.—Corpus convexum, parum nitens, punctatum, pubescens: caput sat magnum, transversum, breve, thorace paullo latius; vertex convexus, non impressus; frons abrupte declivis: oculi parvi, subrotundi: ocelli approximati, vertice triangulum fingentes: antennæ 11-articulatæ, longi-fusiformes, crassæ, pubescentes, thorace vix longiores; articulus 1st. et 2nd. brevissimi; 3rd. longus, obconicus; 4th. et sequentes subrotundi, approximati, usque ad clavam decrescentes; 10th. et 11th. minutus, clavam brevem suborbicularis 9th. fingentes: thorax ovatus: prothorax brevissimus, supra non conspicuus: mesothoracis scutum longitudine latius, parapsidum suturæ optime determinatæ; paraptera et epimera magna; scutellum ovatum, vix convexum: metathorax magnus, declivis: petiolus longus, gracilis: abdomen fusiforme, subcompressum, nitens, scitissime punctatum, parce pubescens, thorace brevius et multo angustius; segmentum unicum dorsum omne occupans; ventralia vix conspicua: pedes validi, recti, simplices, subæquales, pubescentes; coxae parvae; tarsi breves, articulus 1st. longus dilatatus, 2nd. et sequentes minutus brevissimi; ungues et pulvilli parvi: alæ mediocres; nervus humeralis ulnari paullo brevior, radialis longus, cubitalis brevissimus: metalis nervus unicus simplex costæ dimidio longior.

"The mandibles of this species are very strong. The usual structure of the antennæ may still be traced in it, and the gradual development of the two basal joints seen in *E. Zalates* (which retains some of the development of the metatarsus observable in this) until the 1st. becomes the longest joint, as in the other subfamilies, and in some of the *Eucharidæ*, as *E. Xenia* and *E. fuscata*."—*Haliday*.

Sp. 5. Euch. Iello. *Mas. Viridis, antennæ nigræ, abdomen nigrum, pedes nigro-virides, tarsi picei, alæ sublimpidæ.*

Mas.—Obscure viridis: frons viridi-cyanea: oculi et ocelli obscure rufi: os ferrugineum: antennæ nigræ; articuli 1st. et 2nd. nigro-virides: abdomen nigrum: pedes nigro-virides; trochanteræ ferruginei; genua ferruginea; tarsi picei: alæ sublimpidæ;

squamulæ piceo-virides; nervi fusci. (Corp. long. lin. 1 $\frac{1}{2}$; alar. lin. 2 $\frac{1}{2}$.)

Hobart Town, Van Diemen's Land.

Mas et Fem.—Corpus convexum, parum nitens, hirtum, rude punctatum: caput transversum, brevissimum, thorace fere latius; vertex latus, non impressus; frons abrupte declivis: oculi parvi, subrotundi, extantes: ocelli 3, medius perparum antepositus: antennæ 11-articulatae, sub-moniliformes, pubescentes, thorace paullo longiores; articuli exteriores *mari* distincti; 1st. mediocris, gracilis; 2nd. cyathiformis; 3rd. longissimus, subclavatus; 4th. et sequentes subcyathiformes, usque ad 10th. curtantes; 11th. ovatus, 10°. longior; 9th. 10th. et 11th. *fem.* clavam fingentes ovatam: thorax ovatus, altus, gibbus: prothorax minimus, supra non conspicuus: mesothoracis scutum magnum; parapsidum suturæ bene determinatæ; paraptera et epimera conspicua; scutellum latum, supra brevi-obconicum, postice angulum fingens rectum: metathorax sat magnus, declivis: petiolus brevis, validus: abdomen fusiforme, subcompressum, lœve, nitens, hirtum, thorace angustius vix longius; segmentum unicum dorsum omne occupans et ventralia prorsus obtegens: pedes graciles, simplices, pubescentes; coxae parvae; tibiæ rectæ; tarsis articulus 1st. longissimus, 2nd. brevis, 3rd. et 4th. brevissimi, 5th 2nd. vix longior; unguis et pulvilli minuti; metapedes propedibus multo longiores: alæ mediocres, pubescentes; nervus ulnaris humerali multo brevior, radialis longus, cubitalis brevissimus stigma minutum punctiforme fingens: metalis nervus unicus simplex costæ dimidio longior, apice setis uncinatis praeditus.

Sp. 6. Euch. Zalates. *Mas et Fem. Nigra, antennæ nigro-piceæ, pedes picei, tarsi fulvi, alæ limpidæ.*

Mas et Fem.—Nigra: oculi et ocelli obscure rufi: antennæ nigro-piceæ: pedes picei; genua fulva; tibiæ apice fulvæ; tarsi fulvi; unguis et pulvilli fusci; metatarsi basi fusci: alæ limpidæ; squamulæ fuscae; nervi fusci. (Corp. long. lin. 2; alar. lin. 2 $\frac{1}{2}$.)

Sydney, New South Wales; and in March, at King George's Sound, Australia.

Mas.—Corpus convexum, parum nitens, rude punctatum, fere glabrum: caput mediocre, transversum, brevissimum, thorace vix latius; vertex latus, non impressus; frons abrupte declivis, ad os

angustior : oculi parvi, subrotundi, extantes : ocelli vertice triangulum fingentes : antennæ subfiliformes, graciles, pubescentes, thorace vix longiores ; articulus 1st. longus, gracilis, sublinearis ; 2nd. cyathiformis ; 3rd. et sequentes ad 9th. breves, lineares, approximati : thorax ovatus : prothorax brevissimus, supra vix conspicuus : mesothoracis scutum latitudine longius ; parapsidum suturæ bene determinatæ ; paraptera et epimera magna ; scutellum subrhombiforme : metathorax magnus, declivis : petiolus brevis, validus : abdomen ovatum, gibbum, nitens, pubescens, scitissime punctatum, thorace multo brevius et angustius ; segmentum 1st. maximum ; 2nd. et sequentia brevissima : pedes mediocres, simplices, subæquales ; coxæ parvæ ; femora gracilia ; tibiæ rectæ ; tarsis articuli 1st. ad 4th. curtantes, 5th. 4th. paullo longior ; unguis et pulvilli parvi : alæ mediocres ; nervus ulnaris humerali brevior, radialis longus, cubitalis brevissimus ; metalis nervus unicus simplex costæ dimidio longior.

Fem.—Caput thorace paullo angustius : petiolus brevissimus : abdomen convexum, thoracis latitudine et longitudine.

Sp. 7. Euch. Eribotes. *Mas et Fem. Viridis, antennæ nigrae, pedes fulvi, femora viridia, alæ sublimpidæ.*

Viridis : capitis frons cyaneo-viridis : oculi et ocelli rufi : antennæ nigrae ; articuli 1st. et 2nd. virides : pedes fulvi ; coxæ virides ; trochanteres picei ; femora viridia, apice fulva ; tarsi pallide fusci : alæ sublimpidæ ; squamulae virides ; nervi picei. (Corp. long. lin. 1 $\frac{3}{4}$; alar. lin. 2 $\frac{3}{4}$.)

Hobart Town, Van Diemen's Land ; and Sydney, New South Wales.

Mas.—Corpus convexum, parum nitens, rude punctatum, parce pubescens : caput sat magnum, breve, transversum, thorace latius ; vertex latus, non impressus ; frons abrupte declivis : oculi parvi, subrotundi, extantes : ocelli approximati, vertice in triangulum dispositi : antennæ 12-articulatae, filiformes, pubescentes, thorace paullo longiores ; articulus 1st. longus, gracilis, sublinearis ; 2nd. cyathiformis ; 3rd. longus ; 4th. et sequentes breves, lineares, usque ad 9th. curtantes ; 10th. 11th. et 12th. arcte applicati clavam fingentes fusiformem 9°. duplo longiorem : thorax ovatus ; segmentis suturæ bene determinatæ : prothorax brevissimus, supra vix conspicuus : mesothoracis scutum longitudine latius ; parapsides conspicuae, extantes ; scutellum ovatum, valde convexum :

metathorax mediocris, declivis: petiolus longus, sat validus: abdomen ovatum, convexum, nitens, lœve, glabrum, thorace multo brevius et angustius, apice quasi truncatum; segmentum 1^{um}. magnum, dorsi plus dimidium occupans; 2^{um}. et sequentia brevia: ventralia vix conspicua: pedes simplices, subæquales; coxae mediocres; femora gracilia; tibiae rectæ; tarsis articuli 1^o. ad 4^{um}. curtantes, 5^m. 4^o. paullo longior: unguis et pulvilli minuti: alæ amplæ; nervus ulnaris humerali paullo brevior, radialis brevis, cubitalis brevissimus stigma fingens punctiforme.

Sp. 8. Euch. Xeniades. Mas. *Ænea-viridi varia, antennæ nigræ, abdomen nigro-æneum, pedes ferruginei, femora viridia, alæ sublimpidæ.*

Mas. — *Ænea*: capitis frons viridis: oculi et ocelli obscure rufi: antennæ nigræ; articuli 1^o. et 2^o. nigro-ænei: thorax subitus viridis: abdomen nigro-æneum: pedes ferruginei; coxae virides; femora viridia, apice ferruginea; tibiae piceæ, apice et basi ferrugineæ; tarsi apice fusci; protarsi fusci: alæ sublimpidæ; squamulæ piceo-virides; nervi fusci. (Corp. long. lin. 1½; alar. lin. 2½.)

Var. β. — Capitis vertex et thorax virides æneo-varii.

Sydney, New South Wales.

Mas. — Corpus parvum, breve, crassum, convexum, nitens, lœve, parce hirtum: caput transversum, breve, parvum, thoraci arcte applicatum; vertex latus; frons abrupte declivis: oculi mediocres, non extantes: antennæ graciles, submoniliformes, corporis dimidio longiores: thorax breviovatus, altus: petiolus brevis: abdomen ovatum, fere planum, thorace brevius et angustius: pedes graciles, simplices, subæquales: alæ amplæ.

Sp. 9. Euch. Democles. Mas. *Æneo-atra, antennæ piceæ, abdomen purpureum cupreo-varium, pedes picei, alæ limpidæ.*

Æneo-atra: oculi et ocelli rufi: antennæ piceæ: abdomen purpureum, cupreo-varium: pedes picei: alæ limpidæ; squamulæ piceæ; nervi fulvi. (Corp. long. lin. ¾; alar. lin. 1½.)

Sydney, New South Wales.

GENUS.—*PERILAMPUS*, *Latreille*.

Corpus parvum, breve, gibbum, nitens, rude punctatum, parce hirtum : thorax breviovatus : prothorax transversus, brevissimus : mesothoracis scutum longitudine latius ; parapsides fere laeves, suturæ bene determinatæ ; scutellum brevi-obconicum, extans, postice retractum : metathorax magnus, obconicus, declivis : petiolus brevissimus : abdomen rhombiforme, laeve, glabrum, contractum, thorace brevius et angustius : pedes pubescentes, simplices, subæquales : alæ mediocres ; nervus humeralis ulnari plus duplo longior, radialis ulnari vix brevior, cubitalis brevissimus ; stigma minutum, subrotundum.

Sp. 1. Peri. Saleius. Mas. *Viridis æneo-varius*, *abdomen atrum*, *pedes fulvi*, *femora viridia*, *alæ limpidæ*.

Viridis æneo-varius : abdomen atrum : pedes fulvi ; coxae virides ; femora viridia ; unguis et pulvilli fuscus : alæ limpidæ ; squamulae piceæ ; nervi proalis fuscus, metalis fulvi. (Corp. long. lin. $\frac{1}{6}$; alar. lin. $1\frac{1}{2}$.)

March ; King George's Sound, Australia.

GENUS.—*MISCOGASTER*, *Walker*.

Fem.—*Callimomi similis* : corpus angustum, convexum, nitens, scitissime squameum, parce hirtum : caput transversum, breve, thorace paulo latius ; vertex latus ; frons impressa, abrupte declivis : oculi mediocres, non extantes : thorax longiovatus : prothorax transversus, brevis : mesothoracis scutum longitudine paulo latius ; parapsidum suturæ bene determinatæ ; scutellum obconicum : metathorax transversus, mediocris, postice angustior : petiolus brevissimus : abdomen compressum, laeve, supra ovatum, planum, subtus valde carinatum, thorace paulo brevius : oviductus exertus ; vaginæ abdominis dimidio breviores : pedes graciles, simplices, subæquales : alæ amplæ ; nervus humeralis ulnari multo longior, radialis ulnari paulo brevior, cubitalis radiali multo brevior ; stigma parvum.

Sp. 1. Mis. Ciron. Fem. *Viridi-cyanea*, *antennæ nigræ*, *pedes flavi*, *femora viridi-cincta*, *alæ limpidæ*.

Viridi-cyanea : oculi et ocelli rufi : antennæ nigræ ; articuli 1st. et 2nd. virides : oviductus flavus ; vaginæ nigræ : pedes flavi ; coxae

virides; femora viridi-cincta; tarsi apice fusi: alae limpidæ; squamulæ piccæ; nervi proalis fusi, metalis fulvi. (Corp. long. lin. 1; alar. lin. 1 $\frac{1}{2}$.)

Hobart Town, Van Diemen's Land.

Fem.—Corpus breve, crassum, convexum, nitens, læve, hirtum: caput transversum, breve, thorace latius; vertex latus; frons abrupte declivis: oculi mediocres, non extantes: antennæ clavatæ, corporis dimidio vix longiores; articulus 1^o. longus, gracilis, linearis: thorax ovatus, parce punctatus: prothorax transversus, sat magnus, antice angustior; latera rotundata: mesothoracis scutum brevissimum; parapsides extantes, suturæ optime determinatae; scutellum parvum, obconicum: metathorax magnus, obconicus, declivis: petiolus brevis, gracilis: abdomen ovatum, fere glabrum, thorace brevius et angustius; segmenta 3 subæqualia dorso conspicua: oviductus vaginæ abdominis dimidio vix longiores: pedes simplices, subæquales: alæ mediocres; nervus humeralis ulnari plus duplo longior, radialis longus, cubitalis rectus; stigma mediocre, subfurcatum.

Sp. 2. Mis. Nelo. Fem. *Atra, antennæ nigræ, pedes picei, tarsi fulvi, alæ subfuscæ.*

Atra: oculi et ocelli picei: antennæ nigræ: oviductus fulvus: vaginæ nigræ: pedes picei; genua fulva; tibiæ apice fulvæ; tarsi fulvi, apice fusi: alæ subfuscæ; squamulæ piceæ; nervi proalis picei, metalis fusi. (Corp. long. lin. $\frac{2}{3}$; alar. lin. 1 $\frac{1}{4}$.)

March; King George's Sound.

Fem.—Corpus parvum, breve, convexum, nitens, scitissime squameum, parce hirtum: thorax ovatus: prothorax transversus, brevissimus: mesothoracis scutum longitudine multo latius; parapsidum suturæ bene determinatae; scutellum obconicum: metathorax transversus, parvus: petiolus brevis: abdomen ovatum, planum, læve, subtus vix carinatum, thorace paullo brevius et angustius: pedes simplices, subæquales: alæ mediocres; nervus humeralis ulnari longior, radialis ulnari brevior, cubitalis adhuc brevior; stigma mediocre.

Sp. 3. Mis. Nicon. Fem. *Viridi-ænea, abdomen cupreum, pedes flavi, alæ limpidæ.*

Viridi-aenea : mesothoracis scutellum viride, micans : abdomen cupreum : pedes flavi; coxae virides; tarsi apice fusi: alae limpidæ; squamulæ piceæ; nervi fulvi. (Corp. long. lin. $\frac{2}{3}$; alar. lin. $1\frac{1}{4}$.)

Sydney, New South Wales.

Mas.—Corpus convexum, hirtum, parum nitens, scite squameum: caput transversum, breve, thoracis latitudine; vertex latus; frons abrupte declivis: oculi mediocres, non extantes: antennæ clavatae, thorace breviores; articulus 1^{us}. gracilis, sublinearis; 2^{us}. longicyathiformis; 3^{us}. et 4^{us}. minimi; 5^{us}. et sequentes transversi, usque ad 10^{us}. curtantes; clava conica, compressa, articulo 1^o. latior et plus duplo longior: thorax ovatus: prothorax transversus, sat magnus: mesothoracis scutum longitudine latius; parapsidum suturæ vix conspicuæ; scutellum obconicum: metathorax brevis, parvus: petiolus sat longus: abdomen sublineare, planum, lœve, nitens, thorace brevius et multo angustius: metapedes longi; coxae magnæ; tibiæ subarcuatae, cujusque apici spina: alæ mediocres; nervus humeralis ulnari multo longior, radialis ulnari paullo brevior cubitali paullo longior, cubitalis longus ad alæ apicem propensus vix declivis; stigma minutum.

Sp. 4. Mis. Thera. Mas. *Viridis, antennæ fuscæ, abdomen cupreum, pedes fulvi, metapedes obscuriores, proalæ fusco bifasciatæ.*

Obscure viridis: oculi et ocelli rufi: antennæ fuscæ: abdomen cupreum: pedes fulvi; coxae virides; femora basi viridia; tibiæ fusco-vittatae: metapedes virides; trochanteres picei; genua ferruginea; tibiis spinæ fulvæ; tarsi nigri: alæ limpidæ; squamulæ piceæ; nervi proalis picei, metalis fusi; proalæ fusco-bifasciatæ, fascia 1^o. angusta ad nervi humeralis apicem, 2^o. lata ad nervum cubitalem. (Corp. long. lin. $\frac{4}{3}$; alar. lin. $1\frac{2}{5}$.)

Hobart Town, Van Diemen's Land.

Mas.—Corpus sublineare, convexum, parum nitens, scitissime squameum, parce hirtum: caput transversum, breve, convexum, thorace paullo latius; vertex latus; frons impressa, abrupte declivis: oculi mediocres, non extantes: antennæ filiformes, hirtæ, graciles, corpore paullo breviores; articulus 1^{us}. gracilis, sublinearis; 2^{us}. longicyathiformis; 3^{us}. et 4^{us}. minimi; 5^{us}. et sequentes lineares, longi, discreti, usque ad 12^{us}. paullatim curtantes: thorax longiovatus, postice lœvis nitens: prothorax