

XXX.—Descriptions of Chalcidites discovered in the Isle of Chonos by C. Darwin, Esq. By FRANCIS WALKER, Esq., F.L.S.

Lamprotatus Numitus, Mas. Niger, abdomen æneum, antennæ nigrae, pedes fulvi, femora nigra, alæ sublimpidæ.

Corpus breve, convexum, nigrum, parum nitens, scitissime squameum, parce hirtum: caput transversum, breve, thoracis latitudine; vertex latus; frons impressa, abrupte declivis: oculi rufi, mediocres, non extantes: antennæ nigrae, graciles, filiformes, thorace non longiores; articulus 1^{us} longus, gracilis; 2^{us} longicyathiformis; 3^{us} et 4^{us} minimi; 5^{us} et sequentes breves, usque ad 10^{um} paullo curtantes: thorax longi-ovatus: prothorax transversus, brevis, antice angustus: mesothoracis scutum longitudine latius; parapsidum suturæ bene determinatæ, postice approximatae; paraptera et epimera magna; scutellum subconicum: metathorax brevis, declivis, postice angustus: petiolus brevis: abdomen æneum, brevi-ovatum, nitens, læve, glabrum, thorace multo brevius; segmentum 1^{um} magnum; 2^{um} et sequentia brevissima: pedes fulvi, simplices, subæquales; coxæ nigrae; femora nigra, apice fulva; tarsi apice fusci: alæ sublimpidæ; squamulæ piceæ; nervi fusci; nervus humeralis ulnari fere duplo longior, radialis ulnari non brevior, cubitali multo longior; stigma mediocre. (Corp. long. lin. 1 $\frac{1}{3}$; alar. lin. 2 $\frac{1}{4}$.)

Pteromalus? Oxyntes, Fem. Ater, antennæ nigrae, pedes nigri, tarsi fusci, alæ subfusca.

Corpus angustum, convexum, atrum, nitens, subtilissime squameum, parce hirtum: caput transversum, breve, thorace vix latius; vertex latus; frons impressa, abrupte declivis: oculi picei, mediocres, non extantes: antennæ nigrae, subclavatae, graciles, thorace non longiores; articulus 1^{us} longus, linearis; 2^{us} longicyathiformis; 3^{us} et 4^{us} minimi; 5^{us} et sequentes breves, usque ad 10^{um} curtantes; clava longiconica, acuminata, articulo 10^o plus duplo longior: thorax ovatus: prothorax transversus, brevissimus: mesothoracis scutum longitudine latius; parapsidum suturæ non bene determinatæ; paraptera et epimera magna; scutellum subconicum: metathorax brevis, declivis, postice angustus: petiolus brevissimus: abdomen fusiforme, læve, supra planum, subtus carinatum, apice acuminatum, thorace multo longius: pedes nigri, simplices, subæquales; trochanteres picei; genua picea; tarsi fusci, apice picei: alæ subfusca; squamulæ piceæ; nervi fusci; nervus humeralis ulnari multo longior, radialis ulnari non brevior, cubitali multo longior; stigma parvum. (Corp. long. lin. 1; alar. lin. 1 $\frac{1}{2}$.)

Entedon Ufens, Fem. Viride, abdomen nigro-purpureum, antennæ nigrae, pedes fulvi fusco-cincti, femora viridia, alæ sublimpidæ.

Corpus sublineare, convexum, viride, nitens, scitissime squameum, parce hirtum: caput transversum, breve, thoracis latitudine; vertex latus; frons impressa, abrupte declivis: oculi rufi, mediocres, non extantes: antennæ nigrae: thorax ovatus: prothorax transversus, brevis: mesothoracis scutum longitudine latius; parapsidum suturæ bene determinatæ, postice approximatae; scutellum subconicum: metathorax mediocris, obconicus, declivis: petiolus brevis: abdomen brevi-ovatum, læve, supra depressum, subtus carinatum, apice acuminatum, thorace multo brevius; discus nigro-purpureus; segmentum 1^{um} sat magnum; 2^{um} et sequentia breviora, subæqualia: pedes fulvi, simplices, subæquales; coxæ virides; femora viridia; tarsi apice fusci; mesopedum et metapedum tibiæ apice fusca: alæ sublimpidæ; squamulæ piceæ; nervi fulvi; nervus ulnaris humerali longior, radialis ulnari brevior, cubitalis brevissimus; stigma minutum. (Corp. long. lin. $\frac{3}{4}$; alar. lin. 1 $\frac{1}{4}$.)

Closterocerus Pelor, Mas. Viridis, abdomen purpureum, antennæ nigrae, pedes fulvo-fulvi, femora nigra, alæ sublimpidæ.

Corpus angustum, sublineare, convexum, viride, nitens, scitissime squameum, parce hirtum : caput transversum, brevissimum, impressum, thorace paullo latius ; vertex latus ; frons abrupte declivis : oculi picci, mediocres, non extantes : antennæ nigrae, subfiliformes, hirtæ, graciles, thorace non longiores ; clava fusiformis, acuminata, articulo præcedente plus duplo longior : thorax ovatus : prothorax brevissimus, supra vix conspicuus : mesothoracis scutum longitudine latius ; parapsidum suturæ bene determinatæ, postice approximatæ ; scutellum subconicum : metathorax mediocris, declivis, obconicus : petiolus brevissimus : abdomen sublineare, depressum, læve, purpureum, basi viride, thorace paullo brevius et angustius : pedes fulvi, graciles, subæquales ; coxæ nigrae ; femora nigra, apice fulva ; tarsi flavi, apice fuscii : alæ sublimpidæ ; squamulæ picæ ; nervi fulvi ; nervus ulnaris humerali duplo longior, radialis vix ullus, cubitalis brevissimus in alæ discum abrupte declivis ; stigma minutum. (Corp. long. lin. $\frac{1}{2}$; alar. lin. 1.)

XXXI.—*Descriptions of Chalcidites discovered in Coquimbo by C. Darwin, Esq. By FRANCIS WALKER, Esq., F.L.S.*

Lamprotatus Tubero, Fem. Ater, abdomen nigro-purpureum, antennæ fusca, pedes fulvi, femora picea, alæ limpidæ.

Corpus breve, robustum, convexum, atrum, nitens, scitissime squameum, parce hirtum : caput transversum, breve, thorace latius ; vertex latus ; frons impressa, abrupte declivis : oculi picci, mediocres, non extantes : antennæ fusca, extrorsum crassiores, thorace non longiores ; articulus 1^{us} niger, longus, gracilis ; 2^{us} piccus, longicyathiformis ; 3^{us} et 4^{us} minimi ; 5^{us} et sequentes usque ad 10^{um} breves, approximati ; clava conica, acuminata, articulo 10^o duplo longior : thorax ovatus : prothorax transversus, mediocris, antice angustus : mesothoracis scutum longitudine multo latius ; parapsidum suturæ sat bene determinatæ ; scutellum subconicum ; paraptera et epimera magna : metathorax magnus, obconicus, declivis : petiolus sat longus : abdomen subrhombiforme, læve, nigro-purpureum, supra planum, subtus carinatum, thoracis dimidio vix longius : pedes fulvi, simplices, subæquales ; coxæ nigrae ; femora picea, apice basique fulva ; tarsi apice fuscii : alæ limpidæ ; squamulæ picæ ; nervi fulvi ; nervus humeralis ulnari multo longior, radialis ulnari vix brevior, cubitali multo longior ; stigma minutum. (Corp. long. lin. $1\frac{1}{4}$; alar. lin. $2\frac{1}{4}$.)

Var. β. Antennis articulus 1^{us} viridis : abdomen æneum : femora nigra ; tibiæ picæ.

Lamprotatus? Nævolus, Mas? Viridis, antennæ piceæ, pedes piceo-virides, tarsi flavi, alæ limpidæ.

Corpus angustum, convexum, viride, nitens, scitissime squameum, parce hirtum : caput transversum, breve, thorace latius ; vertex latus ; frons impressa, abrupte declivis : antennæ subfiliformes, graciles, picæ, thorace paullo longiores ; articulus 1^{us} longus, linearis, nigro-viridis ; 2^{us} longicyathiformis ; 3^{us} et 4^{us} minimi ; 5^{us} et sequentes usque ad 10^{um} breves, approximati ; clava longiconica, acuminata, articulo 10^o plus duplo longior : thorax ovatus : prothorax transversus, mediocris, antice angustus : mesothoracis scutum longitudine latius ; parapsidum suturæ bene determinatæ, postice approximatæ ; scutellum subconicum : metathorax mediocris, obconicus, declivis : abdomen ademptum : pedes virides, graciles, subæquales ; trochanteres picci ; genua flava ; tibiæ piceo-virides ; tarsi flavi, apice fuscii : alæ limpidæ ; squamulæ picæ ; nervi fuscii ; nervus humeralis ulnari multo