THE BOTANY

OF

THE ANTARCTIC VOYAGE

OF

H.M. DISCOVERY SHIPS EREBUS AND TERROR IN THE YEARS 1839—1843,

UNDER THE COMMAND OF

CAPTAIN SIR JAMES CLARK ROSS, Kt., R.N., F.R.S. &c.



BY

JOSEPH DALTON HOOKER, M.D., R.N., F.L.S.,

ASSISTANT SURGEON OF THE "EREBUS" AND BOTANIST TO THE EXPEDITION.



Victoria Barrier and Land. Lat. 78° S. Mount Erebus active Volcano, and Mount Terror.

Published under the Authority of the Lords Commissioners of the Admiralty.

LONDON;

REEVE, BROTHERS, KING WILLIAM STREET, STRAND.

1844.

[Auckland and

and in the little island of Trinidad, in lat. 20° S., where we effected a landing with considerable difficulty, in a rocky cove which was cut off by precipices from all other parts of the island, I found the Ferns at the level of the sea in the proportion of 2–3 to the phænogamic plants, and the species were the most common Brazilian ones. This remarkable disparity between the vegetable productions of two islands so contiguous as St. Helena and Ascension, and both so remote from any other land whatever, has some analogy to what obtains in the islands of another isolated group, also situated within the Tropics, though in another ocean—the Galapagos. From the examination of an excellent herbarium formed by Mr. Darwin in three of these islands, and of some of the plants from a fourth island, as well as of those collected hy Mr. Douglas, Dr. Scouler, Mr. Macrae, and Mr. Cuming, in the localities also visited by Mr. Darwin, it would appear not only that the plants of that little archipelago differ widely from those of the main land of S. America, but that its several islets possess in some cases different genera, and more often representative species. The Ferns there bear but a small proportion to the whole Flora, though a more considerable one to that of the two islands in which they are most abundant, and they are rather the common forms of the West Indies than of the neighbouring coasts of Columbia, Peru, or of Mexico.

The *Aspidium venustum*, as it grows in the low woods of Lord Auckland's group, is, for its size, among the most ornamental of Ferns, the larger tree-ferns alone excepted. In one respect it even excels those of more majestic growth, for its feathery fronds are spread out below the level of the eye, so that the beautiful symmetry of the crown, with its rich velvetty crosier-formed young leaves in the centre, is thus fully displayed.

3. ASPLENIUM, L.

1. ASPLENIUM obtusatum, Forst. Prodr. n. 430. Lab. Fl. Nov. Holl. v. 2. p. 93. t. 242. f. 2. Brown, Prodr. p. 150. Schkuhr, Fil. v. 1. p. 6. t. 68. Hombr. et Jacq. in Voy. au Pole Sud, Bot. Monocot. Crypt. t. 1. A. (sine descript.).

Far. β. obliquum;—A. obliquum, Forst. Prodr. v. 429. Labillard. l. c. t. 242. f. 1. Schkuhr, l. c. t. 71. A. chondrophyllum, Bertero in Herb. Hook. A. apicidentatum*, Hombr. et Jacq. l. c. t. 1. A. (sine descript.).

HAB. Lord Auckland's group and Campbell's Island; very common on the rocks near the sea and at the margins of the woods.

All the various stages between the A. obliquum, Forst., and A. oblusatum, Forst., exist in Lord Anckland's group, and probably in other islands of which this plant is an inhabitant; one of the specimens indeed, is intermediate between the excellent delineations of the two given by Schkuhr. MM. Hombron and Jacquinot have also figured both the states (from Lord Anckland's group), retaining them under the name of "oblusatum," and added to the plate a representation of another, under the name of A. apicidentatum, which is equally abundant with the others, and I have been unable to distinguish it even as a variety; the production of the apex of the pinna into a tooth, not affording a constant character. I have not quoted the Flora of Mr. Cunningham, or of M. A. Richard, the former not having gathered this species at the time of the publication of his Prodromus, and the latter author, considering it identical with A. lucidum, Forst., leaves it doubtful whether he knew both species. Besides the greater size, different texture, and shining surface of the A. lucidum, its involueres are always very much narrower and longer in proportion to the breadth of the frond. Both are common to many parts of the sonthern hemisphere, and are particularly frequent in the Pacific Islands.

I have retained the name of *obtusatum* for this species, that variety being the more frequent of the two described by Forster.

* *A. apicidentatum*, Homb. and Jacq.; this name probably applies to the pinnæ being terminated by *a* tooth; but all the pinnæ being serrated throughout their whole margin, I presume the term is not used in its ordinary acceptation.

America ; it is of an irregularly four-sided figure, bounded on the north by the strait of Magalhaens, and on the east and west respectively by the South Atlantic and South Pacific Oceans, whilst its southern shores are washed by the Antarctic Sea ; the main body of land lies between the 53rd and 56th parallels of latitude and the 64th and 70th degrees of west longitude, and its greatest extension is from east to west, indicated by a diagonal of 500 miles. The general appearance of the whole has been aptly compared, by Mr. Darwin, to what would be presented by a partially submerged chain of mountains. These islands are, in fact, formed by the southern termination of the great Cordillera that traverses both Americas, which here trends to the eastward, and whose further extension is probably indicated by South Georgia in the same latitude ; and possibly also by Prince Edward's Island, the Crozets, and Kerguelen's Land still more to the east, situated though these be in another ocean. The natural features of Fuegia have been admirably described by various voyagers, and more particularly by Cook, King, Fitzroy, and Darwin, to whose writings I would refer for more particular information. The exposed mountain-tops rise to a height of 7000 feet above the level of the sea, and the lower limit of perpetual snow is reckoned at 3500–4000 feet.

The botanical features exhibited by this country are not circumscribed by its geographical limits; along the north-east shores the very distinct Flora of East Patagonia accompanies the geological formation prolonged there from the Patagonian plains. On the south-west and south sides again, the vegetation is a continuation of that of West Patagonia, and is characteristic of the western flank of the Cordillera, from South Chili to Cape Horn. Thus it is that we find the Andes dividing two botanical regions from the North Polar almost to the Antarctic circle. The greater part of Fuegia is formed by the Andes alone; but the plants of the northeast portion, where the granitic formation of Patagonia introduces a change in the vegetation foreign to that of Tierra del Fuego, will be necessarily included in the present Flora.

The Deciduous Beech (*Fagus antarctica*), is the most distinguishing botanical production of this country. In company with the Evergreen Beech (*F. Forsteri*), it covers the land, especially on the west coasts, as far north as the Chonos Archipelago, in latitude 45° south. It is hardly seen in the north-east portions of Fuegia proper, northward of Staten Land, and though abundant on the west flanks of the Andes, through fourteen degrees of latitude, is unknown on the Atlantic side of Patagonia[†]. I have assumed therefore the shores of the

shores : and again, that the botany of the North Atlantie Islands, the Azores, Madeira, and the Canaries, though these groups are situated in the westerly winds, contain a large proportion of European species. The violence of the perennial westerly gales to the southward of 45° is proverbial amongst sea-faring men; such winds earried II.M.S. 'Chanticleer' from Cape Horn to the Cape of Good Hope, a distance of four thousand miles, in twenty-seven days, and have enabled an occanie fowl, the Cape pigeon, to maintain its position close to a ship during the whole of that distance; but still I am not inclined to attribute the prevalence of the Fuegian Flora over so vast an area to their influence, when exerted against many other opposing agents.

† Trees allied to these seem to have characterized the ancient or fossil flora of Fuegia, for I owe to Mr. Darwin's kindness impressions of the leaves of three apparently distinct species of deciduous Beech, and which are mentioned in that gentleman's journal.

Jacq. Foy. an Pole Sud, Bot. Dicot. t. 10. f. T. Perezia Magellanica, Lagasc. Amæn. vol. i. p. 31. Cassini, Opuse. vol. ii. p. 164. Hook. et Arn, in Comp. Bot. Mag. vol. ii. p. 42. Perdicium Magellanicum, Linn. fil. Suppl. p. 376. Vahl, in Skrivt. Nat. Selsk. vol. i. p. 10. t. 4. P. sinuatum, Banks et Sol. MSS. in Bibl. Banks. cum icone. (TAB. CXI.)

HAB. South Chili and Fuegia. Cape Tres Montes, alt. 2,000 feet, C. Darwin, Esq. Port Famine, Capt. King. Hermite Island, Cape Horn, J. D. H. Staten Land, Dr. Eights and Mr. Webster.

Very variable in size, from two inches to a span or upwards.

19. HOMOIANTHUS, DC.

1. HOMOIANTHUS echinulatus, Cass. in Dict. Sc. Nat. vol. xxxviii. p. 458. DC. Prodr. vol. vii. p. 65. Hook. Ic. Plant. t. 491. Homanthis echinulata, Homb. et Jacq. Foy. au Pole Sud, Bot. Dicot. t. 10. f. S. Perezia recurvata, Lessing, in Linnæa, vol. v. p. 21. Synops. p. 412. P. Doniana, Less. Synops. p. 412. Perdicium recurvatum, Fahl, in Skrivt. Nat. Selsk. vol. i. p. 13. t. 7. Gaud. in. Ann. Sc. Nat. vol. vi. p. 103. et in Freye. Voy. Bot. p. 135. D'Urville in Mém. Soc. Linn. Paris, vol. iv. p. 611. non Don, et Pappig. Chætanthera recurvata, Spreng. Syst. Veg. vol. iii. p. 503. Clarionea recurvata, Don, in Linn. Soc. Trans. vol. xvi. p. 206.

HAB. Strait of Magalhaens, Commerson. Port Famine and Port Gregory, Capt. King. Falkland Islands, very abundant, Gaudichaud, Capt. Sulivan, and all succeeding voyagers.

Rather a variable plant in size, in the glandular pubescence, in the number and size of the spinulæ on the leaves, which are in a single or double row, in the sharpness or bluntness of the leaves, and somewhat also in the form of the involucral scales. It is one of the most interesting plants of the Falkland Islands from the very sweet scent of its large pale-blue flowers, which has been compared to Jessamine and to Violets; it generally grows near the sea in rocky places, and has also been found on the S.E. coast of Patagonia. The *H. Beckii (Perezia*, Hook, et Arn.) of Patagonia is very nearly allied to this, but readily distinguished by the longer spinulose apex of the narrower leaves, and the recurved lower scales of the involucre. The leaves of the latter are exceedingly variable, sometimes wholly without marginal spinulæ, at others crested with white equidistant spinules much longer than those of *H. echinulatus*.

2. HOMOIANTHUS Magellanieus, DC., Prodr. vol. vii. p. 65. Aster Magellanieus, Lam. Illust. Gen. t. 681. f. 3. Perdicium lævigatum, Banks et Sol. MSS. in Mus. Banks. cum icone.

Var. β, lactucoides, duplo major, foliis paulo angustioribus. Perdicium lactucoides, Vahl, in Skrivt. Nat. Selsk. vol. i. p. 11. t. 5. Clarionea lactucoides, Don, in Linn. Soc. Trans. vol. xvi. p. 206. C. glaberrima, Cass. Opusc. vol. ii. p. 165. Perezia lactucoides, Lessing, Synops. p. 413.

HAB. Strait of Magalhaens, Commerson. Port Famine, Capt. King. Cape Negro, C. Darwin, Esq. Good Success Bay, Banks and Solander.

Two plants of very different stature have been brought together by De Candolle under the name of *H. Magel*lanicus; except however in size, I am unable to distinguish them. Lamarck's figure is highly characteristic of the smaller variety, and Cassini's and Lessing's descriptions of the larger. The variety β alone is in Mr. Darwin's Herbarium, the other collections contain both. Sir J. Banks' specimen of the largest state is upwards of two feet high.

20. ACHYROPHORUS, Scop.

1. ACHYROPHORUS tenuifolius, DC.; glabriusculus v. subaraneosus, caule simplici, foliis gramineis omnibus radicalibus filiformibus v. angustissime lineari-spathulatis lineari-lanecolatisve integerrimis sinuatis piunatifidisve segmentis patentibus remotis linearibus, scapo monocephalo, involucri ovato-campanulati squamis linearibus lineari-lanecolatisve acuminatis plus minusve araneo-tomentosis basi sparse hispido-pilosis. A. tenuifolius, *DC. Prodr.* vol. vii. p. 94. Seriola tenuifolia, *Hook. et Arn. in Comp. Bot. Mag.* vol. i. p. 31. S. incana, *Hook. et Arn. l. c.* vol. ii. p. 42. Oreophila tenuifolia, *Don, MSS.*

HAB. Strait of Magalhaens; Port Gregory, Capt. King. Elizabeth Island, C. Darwin, Esq.

A very variable species in the foliage, which is narrow and grass-like. Mr. Darwin has gathered a variety at Port St. Julian on the Patagonian coast, with rather larger capitula, but which does not appear otherwise distinct; it is *Seriola incana*, H. and A. It has also been collected by Capt. King at Cape Fairweather.

2. ACHYROPHORUS arenarius, Gaud.; parce hispido-pubescens v. glabriusculus, radice elongata collo 1-3-cephalo, foliis omnibus radicalibus lineari-obovato-lanceolatis interdum anguste lineari-elongatis longe petiolatis obtusis acuminatisve sinuato-dentatis pinnatifidisve, scapo foliis longiore monocephalo nudo foliisve 1-2 aucto, involucri campanulati squamis araneo-tomentosis glabratisve exterioribus parce hispido-pilosis. A. arenarius, *DC. Prodr.* vol. vii. p. 95. Hypochœris arenaria, *Gaud. in Ann. Sc. Nat.* vol. v. p. 103, *et in Freyc. Voy. Bot.* p. 134 et 461. *D'Urv. in Mém. Soc. Linn. Paris,* vol. iv. p. 609. H. minima ? *Willd. D'Urv. l.c.* Seriola apargioides, *Less. Hook. et Arn. in Comp. Bot. Mag.* vol. ii. p. 42. (TAB. CXII. *Dissection.*)

HAB. Strait of Magalhaens, Port Gregory, Capt. King. Falkland Islands, Gaudichaud, D'Urville, C. Darwin, Esq., J. D. H.

This again is a highly variable plant, the majority of the Falkland Island specimens scarcely agreeing with Gaudichaud's description (as given in Freycinet's Voyage), in which the peduncles are said to be clongated and branched, though in the notes on the species, M. Gaudichaud states that they are either branched or simple. Small specimens entirely coincide with D'Urville's character of *H. minima*? Willd. The other species of this genus, as *H. apargioides*, and *H. taraxaeoides*, are, however, so variable that the character of the single or many-flowered peduncle loses its value as a mark whereby to distinguish them.

PLATE CXII., middle dissections. Fig. 1, plumose pappus ; fig. 2, ripe achænium, transversely rugose :- both magnified.

21. TARAXACUM, Hall.

1. TARAXACUM dens-leonis, Desf.; Leontodon Taraxacum, Linn. Sp. Pl. n. 1122.

Var. lævigatum. T. lævigatum, DC. Cut. Hort. Monsp. p. 149. Prodr. vol. vii. p. 146. D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 604. Gaud. in Freyc. Voy. Bot. p. 134. Leontodon Lycodon, Banks et Sol. MSS. in Mus. Banks. cum icone. (TAB. CXII.)

HAB. Fuegia, Good Success Bay, Banks and Solander. Falkland Islands, D'Urville, J. D. H.

This variety has also been collected at Port St. Julian on the Patagonian coast, by Mr. Darwin.

PLATE CXII., right hand figure. Fig. 1, floret ; fig. 2, stamen ; fig. 3. seta of pappus ; fig. 4, ripe achænium : ---all magnified.

22. MACRORHYNCHUS, Less.

1. MACRORHYNCHUS pumilus, DC.; parce villo albido hirsutus, foliis anguste lineari-elongatis subgramineis integerrimis sinuatis runcinato-pinnatifidisve, scapo foliis longiore, involueri squamis linearilanceolatis foliaceis extus glanduloso-hispidis. M. pumilus? DC. Prodr. vol. vii. p. 152. Taraxacum pumilum et T. coronopifolium, Gaud. in Ann. Sc. Nat. vol. v. p. 103, et in Freye. Voy. Bot. p. 461. D'Ure. in Mém. Soc. Linn. Paris, vol. iv. p. 609. Macrorhynchus Chilensis, Hook. et Arn. in Comp. Bot. Mag. vol. ii. p. 42. Ixeris monocephala, Cass. in Diet. Sc. Nat. vol. xxxix. p. 389. Leontodon pubescens, Banks et Sol. MSS. in Mus. Banks. cum icone. (TAB. CXII. sub nom. M. coronopifolius.)

HAB. Falkland Islands, grassy places near the sea; Gaudichaud, D'Urville, C. Darwin, Esq., J. D. H.

This, again, appears a very Protean plant in the foliage, which is entire, sinuato-pinnatifid, or deeply pinnatifid with linear spreading segments. The plant varies from two to six inches long, and bears one or many seapes, all the parts being more or less clothed with a soft subtomentose pubescence; it has also been found at Cape Fairweather by Capt. King.

PLATE CXII., left hand figure. Fig. 1, receptacle; fig. 2, floret; fig. 3, stamens; fig. 4, achænium:-all magnified.

23. SONCHUS, L.

1. Sonchus oleraceus, Linn. Sp. Pl. n. 1116.

HAB. Chonos Archipelago, C. Darwin, Esq.

Most probably migrated thither since the discovery of South America, from the adjacent coast. It is also naturalized in several parts of Patagonia.

24. HIERACIUM, L.

1. HIERACIUM Antarcticum, D'Urv.; stolomibus nullis, foliis radicalibus lanceolato-spathulatis obtusis subacutisve basi in petiolum attenuatis obscure sinuato-dentatis glaberrimis v. parcissime pubescentibus caulinis paucis linearibus dentatis, caule nudiusculo patentim glanduloso-piloso superne subvilloso 2-3-floro, pedicellis obscure araneosis, involucri campanulati squamis linearibus pilis atris elongatis dense vestitis. II. Antarcticum, D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 608. Gaud. in Freye. Voy. Bot. p. 134.

HAB. Falkland Islands, D'Urville; rocky places near the sea, J. D. H.

Folia exemplaribus Falklandieis uncialia, Patagonieis 3-4-pollicaria. Caulis 4-6 unc. longus. Involucrum $\frac{1}{2}$ unc. longum.

I have described this species partly from my own specimens, which arc very imperfect, and partly from others gathered in Patagonia (Cape Fairweather) by Capt. King, where a second species occurs of which a diagnosis is subjoined.*

* HIERACIUM Patagonicum, Hook.fil.; totum pilis patentibus hirtum, stolonibus nullis, foliis radicalibus oblongolanceolatis subaeutis integerrimis in petiolum attenuatis caulinis paucis sessilibus angustioribus obscure et remote dentatis, caule crecto subnudo apice paniculatim ramoso, peduneulis pedicellis squamisque iuvolueri linearibus pilis atris rigidis patentibus subsetosis.

HAB. Patagonia; Cape Fairweather, Capt. King.

Planta pedalis. Folia pauca, 6-uncialia. Panicula 6-8-flora. Involucra $\frac{1}{2}$ unc. longa.—H. gracili, Hook., Americae borealis, affinis.

XXVI. STYLIDIEÆ, Juss.

1. FORSTERA, L.

1. FORSTERA muscifolia, Willd., Sp. Pl. vol. iv. p. 148. DC. Prodr. vol. vii. p. 338. F. uliginosa, Homb. et Jacq. in Voy. an Pole Sud, Bot. Phan. Dicot. t. 16 D. Phyllachne uliginosa, Forster, Comm. Goett. vol. ix. p. 24. Swartz in Schrad. Journ. vol. ii. p. 173. t. 1, et in Koenig and Sims Annals of Bot. vol. i. p. 286. t. 5. Lamarck Illust. Gen. t. 741. Journ. Hist. Nat. p. 190. t. 10. f. 2. Stibas, Commerson, MSS.

HAB. Strait of Magalhaens, Commerson, MM. Hombron et Jacquinot; Fuegia, Good Success Bay, Banks and Solander, Forster, C. Darwin, Esq.; Port Famine, Capt. King; Hermite Island, Cape Horn, J. D. H.

For remarks upon this species, see Part 1. p. 39 of the present work. Like the *Donatia*, a plant which, from the nature of the soil, climate, and vegetation of the Falklands, might be expected to have been met with there, accompanying the *Cattha appendiculata* and *Astelia pumila*.

XXVII. LOBELIACEÆ, Juss.

1. PRATIA, Gaud.

1. PRATIA repens, Gaud. vid. ante Part 1. p. 42. in note.

HAB. Fuegia, Staten Land, Dr. Eights; Falkland Islands, Gaudichaud, D'Urville, and all succeeding voyagers.

Since the publication of the synopsis of this genus, in the first part of the present work, I have examined a new species from the Straits of Magalhaens, also inhabiting the eastern side of the Andes of Chili, specimens of which, from the latter locality, were then considered to be the true *P. repens*, which, so far as I am aware, is a native of the Falkland Islands, Staten Land, and Valparaiso only.

2. PRATIA *longiflora*, Hook. fil.; glaberrima, caule breviusculo repente subsimplici, foliis paucis erectis carnosis longe petiolatis ovatis obtusis integerrimis v. obscure sinuatis, pedunculis fere terminalibus folio æquilongis ebracteatis, calycis segmentis ovatis acutis, corollæ tubo cylindraceo elongato lobis patentibus triplo longiore.

HAB. Strait of Magalhaens; Cape Negro, C. Darwin, Esq.

Herba laxe cæspitosa. Caulis diametro pennæ passerinæ, 1 unc. longus, repens, nodosus, apice ascendente. Petioli basi vaginantes, $\frac{1}{2}$ -1 unc. longi, crassiusculi, crecti. Folia magnitudine varia, $\frac{1}{4} - \frac{3}{4}$ uncialia, subcoriacea, enervia. Pedunculi ex axillis supremis orti, validi, infra florem gradatim incrassati. Ovarium late oblongum, gibbosum. Calycis dentes erecti sub $\frac{1}{2}$ lin. longi.

Very nearly allied to the former, but differing in the short stems and much smaller and narrower foliage, and most materially in the narrow cylindrical tube of the corolla, which is far longer than the segments, and nearly four times as long as broad. Mr. Bridges has gathered specimens in the marshes of El Valle de las Cuevas, on the eastern side of the Andes of Chili.

XXVIII. GESNERIACEÆ, Nees.

1. MITRARIA, Cav.

1. MITRARIA coecinea, Cavanilles, Icones, vol. vi. p. 67. t. 579. DC. Prodr. vol. vii. p. 537. HAB. Chonos Archipelago, C. Darwin, Esq.

XXIX. ERICEAE, Br.

1. PERNETTYA, Gaud.

1. PERNETTYA mucronata, Gaud. in Ann. Sc. Nat. vol. v. p. 102. in note. DC. Prodr. vol. vii. p. 587. Hombr. et Jacq. in Voy. au Pole Sud, Bot. Phan. Dicot. t. 22. X. Y. Z. Arbutus mucronata, Linn. fil. Suppl. 239. Forst. Comm. Goett. vol.ix. p. 31. Lamarck, Illust. t. 366. f. 7. Graham, in Bot. Mag. t. 3093. Lindley, Bot. Reg. t. 1675. Lodd. Bot. Cab. t. 1848. A. rigida, Banks et Sol. MSS. in Bibl. Banks. cum icone.

HAB. Strait of Magalhaens, Commerson; Fuegia, Banks and Solander, and found by all succeeding voyagers, throughout that country.

One of the most abundant of Fuegian plants, exceedingly variable in the size of its foliage. Owing, apparently, to the puncture of an insect, the apices of the ramuli in the present and following species frequently assume the form of cones, being covered with densely imbricated leaves so metamorphosed as exactly to resemble the scales of an *Abies*.

Though Protean in its foliage, this species is very confined in its geographical limits, advancing no further north than Cape Fairweather, on the east coast of Patagonia.

2. PERNETTYA *pumila*, Hook.; humilis, glaberrima, subcæspitosa, ramosa, caulibus prostratis vel suberectis, foliis imbricatis sessilibus ovatis acutis obtusisve concavis subter carinatis marginibus subtilissime cartilagineo-serrulatis, pedicellis axillaribus arcuatis folio æquilongis longioribusve 1-floris basi bracteolatis.

Var. a, minor, foliis densius imbricatis obtusis. P. pumila, Hook. Ic. Plant. t. 9. DC. Prodr. vol. vii.
p. 586. Homb. et Jacq. in Voy. au Pole Sud, Bot. Dicot. t. 22. S et T. Arbutus pumila, Linn. fil. Suppl.
n. 239. Forst. Comm. Goett. vol. ix. p. 32. Andromeda humilis, Banks et Sol. MSS. in Bibl. Banks. cum icone.

Var. β, empetrifolia, foliis laxe imbricatis angustioribus subacutis obtusisve. P. empetrifolia, Gaud. in Ann. Sc. Nat. vol. v. p. 102. Freye. Toy. Bot. p. 454. t. 67. D'Urville in Mém. Soc. Linn. Paris, vol. iv. p. 607. DC. Prodr. vol. vii. p. 586. Andromeda empetrifolia, Lamk. Encycl. vol. i. p. 155. Arbutus empetrifolia, Linn. fil. Suppl. v. 239. Bruyère à feuilles pointues," Pernetty, Voy. t. 2. p. 64.

HAB. Var. a. From Cape Tres Montes (Patch Cove, alt. 2,000 feet), on the west coast of South Chili to Cape Horn, and in the Falkland Islands, *Commerson*, *Banks and Solander*, *Forster*, and all succeeding voyagers. Var. β. South part of Tierra del Fuego, *Forster*, C. Darwin, Esq., J. D. H. Falkland Islands, most abundant.

The two plants here united under one specific name are decidedly mere varieties. The β . *empetrifolia* is by far the most abundant, and its prostrate stems sometimes attain the length of two feet. Var. *minor*, in its smallest

state, appears, at first sight, sufficiently distinct; but it often runs out to a considerable length, when the leaves become much more laxly imbricated.

Both pink and white berries are found on this species; also cones, similar to those described under *P. mucro*nata, and diseased ramuli, densely eovered with minute, crect, linear leaves.

Dr. Gillies' Arbutus vaccinioides, from the Andes of Chili, which appears identical with Poeppig's A. leucocarpa (Pernettya, DC.), is most likely another form of this plant, the length of the pedicels affording no character either in the flower or fruit.

2. GAULTHERIA, Kalm.

1. GAULTHERIA microphylla, Hook.fil.; pnmila, ramosa, ramis gracilibus setosis, foliis late ovatis v. oblongis obtusis marginibus incrassatis obscure serratis, pedicellis axillaribus brevibus fasciculatis unifloris recurvis, fructibus globosis vel turbinatis. Pernettya serpyllifolia, *DC. Prodr.* vol. vii. p. 587. Arbutus serpyllifolia, *Lam. Encycl.* vol. i. p. 228. A. microphylla, *Forst. Comm. Goett.* vol. ix. p. 32. (TAB. CXII. sub nom. G. Antarcticæ).

HAB. Strait of Magalhaens, Commerson; Port Famine, Capt. King. Good Success Bay, Banks and Solander; Staten Land, Webster; Hermite Island and East Falkland Island. J. D. H.

Suffruticulus 3-4-uneialis, vage ramosus, ramis gracilibus subfiliformibus rufo-brunneis parce setosis. *Folia* sparsa, brevissime petiolata, coriacea, glaberrima, 2-3 lin. longa, læte viridia, nitida. *Flores* parvi; eorolla globosa, alha. *Bacca* pallide rosea, foliis æquilonga.

When figuring this species, I gave it the trivial appellation of *Antarctica*, not being aware of its identity with *Pernettya serpyllifolia*, DC., and *Arbutus serpyllifolia*, Lam., all which names must yield to that of *G. microphylla*, the plant being undoubtedly the little-known *Arbutus microphylla* of Forster.

The genera *Gaultheria* and *Pernettya* are the representatives, in the high southern latitudes, of the *Arbuti*, of the family of *Ericeæ* in the northern and Arctic regions.

PLATE CXVI. Fig. 1, apex of flowering branch; fig. 2, flower; fig. 3, the same laid open; fig. 4, germen, hypogynous glands and stamen; fig. 5, stamen; fig. 6, longitudinal section of germen; fig. 7, transverse section of the same; fig. 8, ripe fruit; fig. 9, longitudinal section of the same; fig. 10, seed; fig. 11, longitudinal section of the same; fig. 12, seed with outer testa removed; fig. 13, longitudinal section of the same; fig. 14, embryo:— all magnified.

XXX. EPACRIDEÆ, Br.

1. LEBETANTHUS, Endl.

1. LEBETANTHUS Americanus, Endl. MSS. in Enchirid. Bot. Allodape Americana, Endl. Gen. Plant. p. 749. Walpers Repert. Bot. Syst. vol. ii. p. 733. Prionotes Americana, Hook. Ic. Plant. t. 30. DC. Prodr. vol. vii. p. 766. Azalea bullata, Forst. MSS. in Mus. Banks. cum icone. Jacquinotia prostrata, Homb. et Jacq. Foy. au Pole Sud, Bot. Dicot. t. 22. R.

HAB. Strait of Magalhacns, Port Famine, *Capt. King*; and thence south throughout the wooded portion of Fuegia and Staten Land, *Forster, C. Darwin, Esq., Mr. Webster, Sc.*

It is certainly very remarkable that the sole American representative hitherto noticed of the order *Epacrideæ*, is also among the very few that so deviate from one of the most important diagnostic characters of that order, as to present a distinctly two-celled anther. Labillardière rightly described the stamens of the Tasmanian *Prionotes*

cerinthoides, Br., as having this structure, and it is a singular circumstance that these two plants, which, through their bilocular anthers and hypogynous filaments, completely unite the *Ericeæ* of the northern hemisphere with their southern representatives in Australia, the *Epacrideæ*, are both natives of very humid climates and densely wooded regions, and not of such localities as the majority of either Order (but especially the *Epacrideæ*) affect.

The subscandent habit of *L. Americanus* is very peculiar; it grows on the trunks of trees, and often creeps up them for some feet. This is also the case with some other distichous-leaved Antarctic plants, as *Callixene*, and *Luzuriaga*, and with the *Prionotes* and *Decaspora* of Tasmania.

XXXI. GENTIANEÆ, Juss.

1. GENTIANA, L.

1. GENTIANA Magellanica, Gaud. in Ann. Sc. Nat. vol. v. p. 89, et in Freyc. Voy. Bot. p. 134. D'Urville, in Mém. Soc. Linn. Paris, vol. iv. p. 607. Grisebach, Gen. et Sp. Gent. p. 237, et in DC. Prodr. vol. ix. p. 99.

HAB. Strait of Magalhaens; Port Famine, Capt. King; south part of Fuegia, C. Darwin, Esq. Falkland Islands, Gaudichaud, D'Urville, Mr. Wright, J. D. H.

2. GENTIANA Patagonica, Grisebach, Gen. et Sp. Gent. p. 237, et in DC. Prodr. vol. ix. p. 99. (TAB. CXV. sub. nomine G. Magellanica).

Var. B, Darwinii, Griseb. l. c.

HAB. Strait of Magalhaens; Elizabeth Island, C. Darwin, Esq.

I can hardly consider Mr. Darwin's specimens to be even a variety of the plant collected by Capt. King at Cape Fairweather (not Port Jamaica, vid. Griseb.), on the coast of Patagonia.

Except the rather broader and more obtuse segments of the less deeply divided calyx, there is nothing to distinguish this from the Tasmanian and New Zealand G. montana, Forst.

PLATE CXV. (under the name of G. Magellanica). Fig. 1, flower; fig. 2, stamen; fig. 3, germen; fig. 4, ripe fruit; fig. 5, seed; fig. 6, the same with the testa removed :---all magnified.

3. GENTIANA prostrata, Haenk. in Jacq. Coll. vol. ii. p. 66. t. 17. f. 2. Griseb. Gen. et Sp. Gent. p. 271, et in DC. Prodr. vol. ix. p. 106.

HAB. Strait of Magalhaens; Cape Negro, C. Darwin, Esq.

For the widely extended gcographical distribution of this little species, see Part 1. p. 56. of the present work.

XXXII. CONVOLVULACEÆ, Juss.

1. CALYSTEGIA, Br.

1. CALYSTEGIA sepium, Br., Prodr. p. 483. Engl. Bot. t. 313. Choisy in DC. Prodr. vol. ix. p. 433. HAB. Chonos Archipelago, C. Darwin, Esq.

This plant, the common English Bind-weed, is universally diffused throughout the temperate regions, both of the northern and southern hemispheres. In the latter it inhabits New Holland, New Zealand, and the Island of Java, according to M. Choisy, in DC. Prodr. l. c.

XXXIII. BORAGINEÆ, Juss.

1. MYOSOTIS, L.

1. Myosotis albiflora, Banks et Sol. MSS.; caulibus e rhizomate valido plurimis prostratis gracilibus foliisque parce appresse pilosis, foliis radicalibus spathulatis petiolatis caulinis obovato-oblongis, floribus paucis axillaribus breviter pedicellatis calycibusque campanulatis appresse pilosis, corollæ tubo calycem superante limbi lobis late oblongis breviusculis. M. albiflora, *Banks et Sol. MSS. in Bibl. Banks. cum icone*.

HAB. Fuegia, Good Success Bay, Banks and Solander. South part of Tierra del Fuego, C. Darwin, Esq.

Rhizoma crassum, $\frac{1}{3}$ une. longum, fibras plurimas atras emittens, apiceque caules 5–8 gerens. Caules 2-unciales, prostrati, apice ascendentes, parce foliosi. Folia $\frac{1}{3}-\frac{1}{2}$ -uncialia, 3–4 lin. lata, apice obtusa, utrinque sed super præcipue pilis albidis appressis sparsa. Flores axillares, non racemosi, inconspicui, pedicellati; pedicello calyce æquilongo, sub $\frac{3}{4}$ lin. longo. Calyx 5-fidus, laciniis ovato-lanceolatis, acutis, corollæ tubo $\frac{1}{3}$ brevioribus. Corollæ tubus teres, fauce glandulis fornicatis superne medio emarginatis fere clausa. Stamina inclusa. Stylus stigmate clavato terminatus.

In size and habit this little species closely resembles the *M. Antarctica* (Part 1. p. 57. t. 38), but it is a much slenderer, less rigid, and comparatively glabrous plant, with larger, though still very inconspicuous, and white flowers. It evidently belongs, by its prostrate stems and axillary flowers, to the New Zealand group of the genus, which, under the species alluded to, I have noticed as very different from that including the majority of the genus.

XXXIV. SOLANEÆ. Juss.

1. SOLANUM, L.

1. SOLANUM tuberosum, Linn. Sp. Pl. 282. Dunal, Monogr. p. 135.

HAB. Chonos Archipelago, C. Darwin, Esq.

The true Potato plant reaches the boundary to which the Antarctic Flora of South America is confined, and is described as particularly abundant in the localities whence Mr. Darwin's specimens were brought. The nature of the present work forbids my dwelling on some of the peculiarities which mark the history and habitat of this plant; and I leave the subject with the less reluctance, because Mr. Darwin's own history of its discovery in an indisputably native state is already published in one of the most interesting 'Journals of a Naturalist' that has ever been written. The following remarks apply wholly to the botanical affinities of the individual species now universally cultivated in all temperate civilized countries.

There are in Sonth America several *Solana*, so closely allied to the true Potato, that it is exceedingly difficult to distinguish them specifically. Though differing materially in the shape of their calycine lobes, they display such variation in these organs, that no specific value can be attached to them alone. The fruit may afford better characters, but that of many is at present unknown. The following is an enumeration of those South American *Solana*, allied to, or varieties of, the true *S. tuberosum*, which exist in the Hookerian Herbarium. I shall commence with the specimens most similar to the common cultivated form.

STIRPS I. S. tuberosum, L.

Var. 1, *vulgare*, planta pubescens, caule robusto, foliis amplis, calycis majusculi lobis e basi late ovata in acumen subelongatum productis.

| Fuegia, the

HAB. Chonos Archipelago, C. Darwin, Esq. Specimens very luxuriant, altogether resembling wellgrown cultivated plants. Hills about Lima, J. Mac Lean, Esq.; several sub-varieties, marked as "yellow, mottled, white, or purple Potato"; but none are so luxuriant as Mr. Darwin's specimens. Juan Fernandez, Bertero; no flower;—Mr. Bertero remarks that it is possibly wild, the roots being bitter.

Var. 2, macranthum, foliolis multi-4-8-jugis ovato-lanceolatis glabratis, corymbis glabriusculis, laciniis calycinis subulatis, corollis amplis $1\frac{1}{2}$ unc. diametro.

HAB. Serras of Amancaes, Peru, Mathews, n. 847.

Var. 3, *puberulum*, foliolo terminali maximo, lateralibus parvis multoties minoribus, corymbis glabratis, calycibus minoribus glabriusculis, corollis amplis.

HAB. Puruchuca, Peru; Mathews, n. 772.

Var. 4, *multijugum*, totum ut in precedente, sed glabratum, foliolis æqualibus, lateralibus multijugis lanceolatis basi cordatis petiolulatis.

HAB. cum priore, Mathews, n. 771.

Var. 5, *polemoniifolium*, foliis incano-pubescentibus, foliolis plurimis parvis, calycis paulo minoris glabrati lobis brevioribus acutis.

HAB. Andes of Chili and Mendoza, Dr. Gillies.

This and the four preceding are all large-flowered states probably of the true *Solanum tuberosum*, npon the pubescence, or form and number of the leaflets of which no reliance is to be placed. If so, its range is from an elevation near Lima in Peru, to the level of the sea at Chonos Archipelago, and inland to the Andes of Mendoza in Chili.

STIRPS II. S. Commersonii, Poir.

Var. 1, glabriusculum, foliolo terminali lateralibus paucijugis majore, floribus majusculis, calycibus pubescentibus.

HAB. Buenos Ayres, Tweedie; Valparaiso, Bridges, n. 401.

Apparently the plant figured in Hort. Soc. Trans. vol. v., p. 249. t. 9, 10, 11, from Commerson's own specimens.

Var. 2, *pilosiusculum*, foliis amplis, foliolis multijugis æqualibus, floribus majusculis, calycibus pubescentibus.

HAB. Mountains of Mendoza, Dr. Gillies; "cult. ad Buenos Ayres sub nom. S. tuberosi," Herb. Hook.

Hardly different from the former variety. Apparently the S. tuberosum of Hort. Soc. Trans., the experiments upon which are there detailed.

Var. 3, glanduloso-pubescens, foliolis parvis ovatis basi cordatis petiolulatis.

HAB. Foot of the mountains of Mendoza, Dr. Gillies.

A smaller plant than either of the foregoing.

Var. 4, glabratum, foliolis paucijugis terminali majore, corymbo paucifloro, floribus minoribus :---an sp. distincta?

Buenos Ayres, in hedges, Dr. Gillies.

FLORA ANTARCTICA.

I think there can hardly be a doubt that the largest-flowered plant, whose varieties I have included under the Stirps I., is the true cultivated Potato, a species, in its wild state, confined to the west of the Andes. Whether the *S. Commersonii*, which chiefly differs in the size of the flowers, be really distinct or not, is another question. Ranging as as it does from one side of the continent to the other, it may perhaps have some claims to be considered the type of the Potato, of which the large-flowered variety, now commonly cultivated with us, is confined, as just observed, to the Pacific side of South America.

That both produce tubers, called "Papas" and "Maglia," is evident, for the specimen from which the tubers were reared to the size of ordinary Potatos, in the Horticultural Society's Gardens, is certainly referable to the small-flowered Valparaiso plant, also collected by Bridges; and the large-flowered species of Lima presents the ordinary varieties of the well-known vegetable, as does the Chonos Archipelago and Mendoza one.

To show how little evidence is to be derived from the mere fact of the species producing tuberous roots, I may mention that there is a third plant, allied to both the former, and found over a great part of extra-tropical South America, bearing tubers, altogether similar to those of the two foregoing *Solana*. This I refrain from naming, though unable to ascertain that it is previously described, but it may readily be recognized by its great general resemblance to *S. Commersonii*, from which it differs in the small fruit, and in the short cupuliform or hemispherical calyx, whose lobes are short, broad, and rounded; while in other respects, as regards pubescence and size and form of the leaflets, it is as variable as the two former. I have seen specimens from Antuco (*Reynolds*), Valdivia (*Bridges*, 719), Valparaiso (*Cuming*, 555), and Uraguay (*Tweedie*), to the last of which the collector has added on the ticket, "This bears a considerable quantity of nasty soft watery Potatos at its root, called Papas Amargas, in consequence of their bitter taste."

With regard to Mr. Darwin's specimens, in producing an abundance of tubers they only follow the habit of *Cardamine hirsuta* and many other plants, when inhabiting such a soil as a shingly beach. In the absence of a bitter principle, evident in the wild tubers of the "Maglia" of the drier parts of South America, the Chonos Archipelago Potato may be compared with the Celery mentioned at p. 287, whose insipidity I attribute partly to the dampness of the elimate, and still more to the absence of the direct rays of the sun.

Professor Henslow, who has investigated the subject of the native Potato with his usual care and skill, agrees with me in cousidering this of Mr. Darwin's to be quite identical with the common cultivated Potato; and he further remarks the differences between it and the "Maglia" of Chili, without, however, pronouncing them specifically distinct.

Thus, from the information I have been able to obtain, it appears very possible that the plant experimented upon in the Horticultural Society's Gardens, is even specifically distinct from the common cultivated Potato, for it is certainly the small-flowered "Maglia" of Chili, and not the large-blossomed "Aquinas" of Chiloe and the Chonos Archipelago. It would be very interesting to introduce the tubers of Mr. Darwin's *S. tuberosum*, and the *S. Commersonii* (the latter both from the east and west coasts), into our gardens; along with, if possible, the short-calyxed species, which is also stated by Mr. Tweedie to produce tubers.

Though I have spoken of these three Solana as all tuberous-rooted, it is more than probable that they are not always or necessarily so, and that the absence of those hybernacula does not indicate specific distinction. Mr. Crnikshanks, who has studied one of the above species in its cultivated state in Chili (probably the S. Commersonii), says of it, that the "Papas Amarillas," or Yellow Potato of Peru, which was grown in the Horticultural Society's Gardens, is a variety of the Solanum tuberosum, differing from all other known varieties of that species in its partiality for a particular climate. Mr. Cruikshanks also remarks, "that it will not produce bulbs near the coast in Peru, nor at Valparaiso, but only on the higher parts and in a very few spots; but that further south in Chili, as near Valdivia, it is very productive." The explanation seems to be, that this yellow Potato, whether a species or variety, is dependent upon a moist and cool climate for the formation of tubers, or, as the inhabitants of Peru express it, on the "tiemperamento de la Sierra". Hence, too, may arise their absence on Dr. Lindley's S. etuberosum, which is intermediate between two of Mr. Mathews' Peruvian states of S. tuberosum, having the foliage and colour of the flowers of his No. 847, which I have made the second variety of S. tuberosum, and the smooth paniele and small calyx of Mathews' No. 771, or my fourth variety of the same species.

Genus SOLANEIS relatum.

1. DESFONTAINEA spinosa, Ruiz et Pavon, Fl. Per. vol. ii. p. 47. t. 186. Don, in Ed. Journ. of Sc. 1831. p. 275. Hook. Ic. Plant. t. 33. D. splendens, H. B. K. Plant. Equinoct. vol. i. p. 157. t. 45.

HAB. Staten Land; Mr. Webster.

Much has been written regarding the affinities of this curious genus; for several reasons, I retain it near Solaneæ, to which Order it was doubtfully referred by the authors of the "Plantes Equinoctiales," and more recently by M. Endlicher. M. Kunth afterwards suggested its relationship with *Theophrasteæ*, which Mr. Don had also suspected. The last-mentioned author has more recently arranged it in *Gentianeæ*, and is followed by Dr. Lindley, in 'The Vegetable Kingdom', who had previously placed it in Aquifoliaceæ (*Nat. Syst. of Bot.*). My own impression is that its proper place is nearer to the order *Ericeæ*, an hypothesis strengthened by the observations of my friend M. Planchon, who has studied this plant most attentively, and who pointed out its affinity with the anomalous genus *Galax*, and particularly with the Arctic European and American *Diopensia Lapponica*, in the position of the anther and some other points.

Capt. King's collection contains a very curious plant from Port Famine, which, from the nature of the fruit and testa of the unripe seeds, I presume, approaches *Ericeæ*, though wholly differing in habit and in some other points which ally it to *Diapensia*. Unfortunately all the specimens are out of flower, which I exceedingly regret, for it may afford characters which will throw a light upon these and other obscure genera of *Monopetaleæ*. Its seeds are enclosed in a double testa, a structure which occurs, though rarely, in several orders of monopetalous Dicotyledons.

The geographical distribution of this curious genus is, like that of *Gunnera*, very extended, from the Andes under the equator, alt. 12,000 feet, to the level of the sea at Staten Island, in lat. 53° south.

XXXV. SCROPHULARINEÆ, Juss.

1. CALCEOLARIA, L.

1. CALCEOLARIA Fothergillii, Sol. in Ait. Hort. Kew. vol. i. p. 30. t. 1. Car. Ic. vol. v. t. 442. f. 1. Bot. Mag. t. 348. Benth. in DC. Prodr. vol. x. p. 208. C. Neeana, Spreng. Syst. Veget. vol. i. p. 44. (TAB. CXVII., left-hand figure).

HAB. Strait of Magalhaens; Port Famine, Capt. King. Falkland Islands, very abundant.

Though very inferior in stature and beauty to most of its congeners, this is among the pretticst of the wild flowers of the Falkland Islands, and the attention of the voyager who is familiar with the genus *Calceolaria* only in the conservatories of Britain, must be attracted by its appearance on the exposed shores of these inhospitable Islands. I have already mentioned several decidedly English plants, which are natives of this portion of the opposite hemisphere; interesting in themselves, they become still more so when contrasted with such foreign-looking associates as the present, or the nodding bells of the *Sisyrinchium*, which sometimes whiten the plains, or the deeporange blossoms of the Falkland Island violet, invariably seen growing with this *Calceolaria*.

PLATE CXVII., left hand figure. Fig. 1, ripc capsules; fig. 2, transverse section of the same; fig. 3, seed; fig. 4, longitudinal section of the same :----all magnified.

2. CALCEOLARIA nana, Sm.; herbacea, glabra v. tenuissime viscoso-tomentella, caule brevissimo, foliis

FLORA ANTARCTICA.

petiolatis ovatis obtusis basi longe angustatis integerrimis crenulatisve, pedunculis scapiformibus unifloris, calycis minute viscoso-tomentelli laciniis late ovatis obtusis, corollæ labio superiore calyce parum breviore inferiore dependente obovato basi longe contracto ultra medium aperto. *Benth. in DC. Prodr.* vol. x. p. 208. *Smith, Icon. ined.* vol. i. p. 1. t. 1. C. uniflora, *Lam. Illust. Gen.* t. 15. f. 3.

HAB. Strait of Magalhaens, Commerson. Port Gregory, Capt. King.

Caules $\frac{1}{2}$ -1-pollicares, apice pedicellos 1-2-subtripollicares ferunt. Staminum filamenta quam in affinibus longiora. Benth. l. c.

The foliage alone is insufficient to distinguish this species from a small state of *C. Fothergillii*, but they are very dissimilar in the calyx and size of the corolla, the sepals of the former being very broad and almost eucullate, covered externally with a viscid yellow tomentum, while in *C. nana*, they are smaller, narrower, and simply pubescent. The corolla of *C. nana* almost equals that of *C. Darwinii*, to which, in every respect, it is nearly allied.

This species has been also found at Cape Fairweather by Capt. King.

3. CALCEOLARIA Darwinii, Benth.; glabra, caule brevi, foliis late oblongis integerrimis vel remote paucidentatis in petiolum longe angustatis, pedunculis scapiformibus 1-3-floris, calycis minute puberuli laciniis late ovatis obtusis, corollæ labio superiore calycem subæquante inferiore dependente maximo late obovato basi longe contracto ultra medium aperto, antherarum loculis ovatis. *Bentham, in DC. Prodr.* vol.x. p. 207. (TAB. CXVII., *right-hand figure*).

HAB. Strait of Magalhaens; Elizabeth Island, C. Darwin, Esq.

Habitus C. polyrhize, corollæ is C. Fothergillii forma similes sed majores, speciosæ, maculatæ. Benth. l. c.

Much the handsomest species of the small section "Scaposæ," to which all the Antarctic Calceolariæ belong. Though very distinct at first sight from the former, the individual parts are so hable to vary that it becomes almost impossible to draw up an absolute distinctive character. For instance, the leaves in one specimen have the same form as in a Cape Fairweather individual of *C. nana*, and, though more glabrous than in most states of the latter plant, they are not universally so; the corollæ are quite alike in the two, and the difference in the length of the filaments is hardly appreciable. The calyx of *C. Darwinii* varies exceedingly in the size and form of its segments, they are sometimes large, broad, and obtuse, as in *C. Fothergillii*, or small and narrow like those of *C. nana*.

PLATE CXVII., right hand figure. C. Darwinii, the natural size.

4. CALCEOLARIA polyrhiza, Cav. Ic. Rar, vol. v. p. 25. t. 441. Benth. in DC. Prodr. vol. x. p. 207.

HAB. Falkland Islands; Neé.

Of this plant I have seen no Falkland Island specimens. It is also a native of Port Desire on the coast of Patagonia, where Mr. Darwin gathered it.

5. CALCEOLARIA plantaginea, Smith, Icon. ined. vol. i. p. 2. t. 2. Hook. in Bot. Mag. t. 2805. Lodd. Bot. Cat. t. 1402. Benth. in DC. Prodr. vol. x. p. 208. C. biflora, Lam. Encycl. vol. i. p. 556. Bæa plantaginea, Persoon, Synops. vol. i. p. 15.

HAB. Strait of Magalhaens, Commerson; Elizabeth Island, C. Darwin, Esq.

Very abundant between the latitude of Valparaiso and the Strait of Magalhaens, though confined to a narrow belt, which runs obliquely across the continent of South America, from lat. 33°, to lat. 53°. In the northern half of its range, between the parallels of Valparaiso and Chiloe, it is chiefly confined to the west of the Andes; in the southern half, between the latter locality and the Strait of Magalhaens, it crosses to the east side of South America; thus avoiding equally the wet, cold, and stormy latitudes of South-west Chili and Fuegia, and the arid plains of Patagonia.

2. LIMOSELLA, Linn.

1. LIMOSELLA aquatica, Linn. Sp. Pl. p. 881. Engl. Bot. t. 357. Benth. in DC. Prodr. vol. x. p. 427.

Var. β, tenuifolia. L. tenuifolia, Nutt. Gen. N. Am. vol. ii. p. 43. Gaudichaud, in Ann. Sc. Nat. vol. v. p. 102, et in Freyc. Voy. Bot. p. 133. D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 607. Benth. in DC. Prodr. vol. x. p. 427. L. australis, Brown, Prodr. p. 443.

HAB. Falkland Islands, Gaudichaud, J. D. H. Kerguelen's Land, J. D. H.

I am convinced there is no specific distinction between the *Limosella aquatica*, L., and *L. tenuifolia*, Nutt., and have consequently united them. In the specimens from the southern hemisphere which I have examined, the leaves do not attain the breadth which those of the northern temperate regions generally present; though, on the other hand, both European, Asiatic, and North American plants of the *L. aquatica* have the foliage narrow as that of *L. tenuifolia*, to which variety some Arctic individuals of *L. aquatica* are quite similar.

The range of this species is nearly identical with that of *Callitriche aquatica* and *Montia fontana*, and there is also a considerable resemblance in the mode and extent of their variation between these three plants. This is not remarkable with regard to *Callitriche* and *Montia*, which are very frequently seen associated together, invariably so in Kerguelen's Land, in the Falkland Islands, in Lord Auckland's Group and Campbell's Island, and thus are influenced in common by every fluctuation of climate and temperature, and by the depth or rapidity of the current, when growing in the water; but the *Limosella* does not occur mixed with these two genera, even though inhabiting the same islands.

In Kerguelen's Land the *Limosella* is found in the muddy bottom of a lake, and probably flowers all the year round. I gathered it in the month of July (mid-winter), beneath two feet of water, covered with two inches of ice; even then it had fully-formed flowers, whose closely imbricating petals retained a bubble of air, the anthers were full of pollen and the ovules apparently impregnated. The climate of Kerguelen's Land being such, that this lake is perhaps never dried, it follows that the plant has here the power of impregnation when cut off from a free communieation with the atmosphere, and supplied with a very small portion of atmospheric air generated by itself. My Falkland Island specimens are in a very poor state. Gaudichaud, who first detected it in that Island, considers it identical with the European plant.

3. VERONICA, L.

1. VERONICA elliptica, Forst.; Fl. Ant. part 1. p. 58. V. deeussata, Ait. ct auctor.

HAB. Strait of Magalhaens to Cape Horn in Fuegia, *Commerson, Banks and Solander*, and all succeeding voyagers. West Falkland Island, chiefly on the southern and western coasts.

2. VERONICA serpyllifolia, Linn. Sp. Pl. p. 15. Engl. Bot. t. 1075. Gaud. in Ann. Sc. Nat. vol. v. p. 102, et in Freye. Voy. Bot. p. 133. D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 607.

HAB. Falkland Islands, abundant near the colonized parts of the Islands; D'Urville, &c.

This species, in affecting principally the vicinity of the settlements and ground much frequented by cattle, was probably introduced originally from Europe into the Falkland Islands. It is found no where else in the southern hemisphere, except the neighbourhood of Quito, where Mr. Kunth doubts its being indigenous, or in equally equivocal situations.

4. OURISIA, Comm.

1. OURISIA Magellanica, Juss.; caule repente, foliis subradicalibus longe petiolatis cordato-ovatis

orbiculatisve obtusis erenatis floralibus orbiculatis semiamplexicaulibus, pedunculis dissitis, calyeis laciniis ovatis obtusiusculis ciliatis subbilabiatim connatis. *Benth. in DC. Prodr.* vol.x. p. 492. *Gærtner, fil. de Fruct.* vol. iii. p. 44, non Poepp. et Endl. Chelone ruelloides, *Linn. fil. Suppl.* p. 271.

HAB. Strait of Magalhaens, Commerson; Good Success Bay, Banks and Solander; Staten Land, Mr. Webster.

Caules breves, erassiusculi uti petioli nervi foliorum et calycis margo pilis nonnullis patentibus ciliati; planta cæterum glabra. *Petioli* 3-4 poll. longi. *Folia* erassiuscula, majora $2-2\frac{1}{2}$ pollicaria, crenis inæqualibus; floralia semi-pollicem lata. *Racemus* fere a basi scapi florifer. *Pedicelli* fructiferi ultra pollicares. *Calycis* laciniæ $2-2\frac{1}{2}$ lin. latæ. *Corolla* 7-8 lin, longa, tubo amplo incurvo, limbi laciniis retusis. *Capsula* late orbiculata, compressiuscula. *Benth. l. c.*

This, of which I have seen but a single specimen, must be one of the handsomest Fuegian plants; it is apparently very searce, for it does not exist in the collections of Capt. King or Mr. Darwin, nor have I myself gathered it.

2. OURISIA breviflora, Benth.; humilis, pilosa, caule ascendente foliato 2–4-floro, foliis petiolatis ovatoorbiculatis basi truncato-subcordatis floralibus sessilibus ovatis, calveis segmentis lineari-oblongis tubo corollæ longioribus. *Benth. in DC. Prodr.* vol. x. p. 493. (TAB. CXVIII. sub nomine O. Antarcticæ).

HAB. Strait of Magalhaens, Port Famiue, *Capt. King*; South part of Tierra del Fuego, *C. Darwin, Esq.* Hermite Island, in elefts of rocks on the mountains, *J. D. II*.

Caules basi ramosi, 2-3-pollicares. Folia vix semi-pollicaria. Calycis segmenta fere 3 lin. longa, angusta, obtusa. Corollæ limbus valde obliquus, laciniis emarginatis, infima quam tubus paulo longior.

A pretty little species, probably not rare in Fuegia, though readily overlooked from its very diminutive size. When the accompanying plate was prepared and the name *O. Antarctica* applied to it, I was not aware of Mr. Bentham having named the plant in the then unpublished volume of De Candolle's Prodromus.

The genus *Ourisia* is highly interesting, from being among those peculiar to the Antarctic or higher latitudes of the southern regions, which have no analogue in the northern, but which, though most abundant in Antarctic America, have representative species in the temperate portions of Terra Australis (*O. integrifolia*, Sm.), and in New Zealand or temperate Polynesia (*O. macrophylla*, Hook.).

PLATE CXVIII. (under the name of O. Antarctica). Fig. 1, flower; fig. 2, corolla laid open; fig. 3, ovarium; fig. 4, transverse section of the same; fig. 5, ripe fruit; fig. 6, transverse section of the same; fig. 7, seed; fig. 8, longitudinal section of the same :—all magnified.

5. EUPHRASIA, L.

EUPHRASIA Antarctica, Benth.; minima, subsimplex, pubescens, foliis euncato-trifidis lobis obtusis brevibus, corollæ tubo exserto limbi lobis brevibus subintegris, capsula ovata obtusa. Benth. in DC. Prodr. vol. v. p. 555.

IIAB. Strait of Magalhaens; Cape Negro, C. Darwin, Esq.

Herba perpusilla, vix pollicaris, glanduloso-puberula. *Caulis* crectus, simplex v. divisus, foliosus. *Folia* 1-3 lin. longa, cuneata, in lacinias 3 lineares obtusas ad medium fissa. *Flores* inter folia summa sessiles, pro planta magnæ. *Calyx* tubuloso-campanulatus, glabriusculus, breviter 5-fidus, lobis obtusis, apice puberulis, marginibus siccitate atratis. *Corollæ* tubus ealycem superans, lobis oblongis oblique emarginato-truncatis, galea vix sub lobis concava. *Stamina* corollam subæquantia, antheris basi bi-aristatis.

A very minute species; also found at Coquimbo in Chili, by M. Gay. It is the southern representative of

its European ally, *E. officinalis*, L., and is still more nearly allied to a Himalayan plant, detected by my friend Mr. Edgeworth, whose researches in the Indian Alps have been rewarded with the discovery of some well-marked types of an American Flora, occurring together where they might have been least expected.

XXXVI. LABIATÆ, Juss.

1. SCUTELLARIA, L.

1. SCUTELLARIA *nummularia folia*, Hook. fil.; parvula, glanduloso-puberula, caulibus gracilibus basi prostratis ascendentibus, foliis breviter petiolatis late elliptico-oblongis rotundatisve obtusis integerrimis subenerviis floralibus conformibus, floribus sparsis axillaribus breviter pedicellatis.

HAB. East coast of Tierra del Fuego, C. Darwin, Esq.

Caules graciles, diametro pennæ passerinæ, basi ramosi; ramis diffusis, simplicinsculis, elongatis, 2-4-uncialibus tenuiter puberulis. Folia $\frac{1}{3}-\frac{1}{2}$ unc. longa, subcoriacea, utrinque subglanduloso-puberula, apice rotundata, basi in petiolum brevem $1-1\frac{1}{2}$ lin. longum angustata. Flores pauci, majusculi, foliis longiores, breviter pedicellati, pedicello calyce puberulo æquilongo. Corolla calyce ter longior, e basi sensim ampliata, rosea (?), pubescens, v. glabrata, lobis superioribus lateralibusque liheris brevibus obtusis, inferiore subpendulo, fance piloso. Achænia immatura lævia.

Allied to the North American *S. antirhinoides*, Benth., but much smaller, and very different in the size of the flowers. A variety, also gathered by Mr. Darwin at Port St. Julian on the Patagonian coast, is more stunted, densely publicent, with shorter leaves, and the lower lip of the corolla bearded internally. The discoverer of this species remarks that the climate and productions of the particular locality which it inhabits, are intermediate in character between those of Patagonia and Fuegia.

2. STACHYS, L.

1. STACHYS *Chonotica*, Hook. fil.; herbacea, erecta, hispido-pilosa, foliis petiolatis oblongo-lanceolatis ovato-oblongisve obtusis acutisve basi cordatis obtuse crenato-serratis floralibus bracteæformibus inferioribus calyce longioribus, verticillastris 4–8-floris remotis, calycis hispidi campanulati dentibus ovatis aristatis, corollæ glabriusculæ tubo calyce longiore.

IIAB. Chonos Archipelago; C. Darwin, Esq.

Species *S. sylvaticæ* simillina, sed folia angustiora, brevius petiolata et obtusiora creuisque obtusioribus; labium inferius corollæ minus profunde secta. *S. Macræi*, Benth., (planta admodum variabili) quoque approximat habitu formaque foliorum, sed tubo corollæ elongato exserto lobisque latioribus labii inferioris sat differt.

A plant, so very closely resembling the *S. sylvatica*, L., of Great Britain, that I long hesitated on the propriety of erecting it into a new species, but do so in concurrence with the opinion of Mr. Bentham. Mr. Watson, also, upon whose thorough knowledge of British plants, in all their exotic forms to which he has had access, the greatest reliance may be placed, has, with his usual kindness, given much attention to the present plant, and sums up the differences between it and European *S. sylvatica*, in the leaves of the latter not being so obtuse nor so obtusely serrate, and in the lateral lobes of the lower lip of the corollæ being more deeply divided. The leaves of the European *S. palustris*, L., however, he adds, vary from very acutely to quite as obtusely serrate.

Not being versed in the whole genus *Stachys*, which contains upwards of one hundred species, I was inclined to regard this plant as possibly intermediate between the *S.sylvatica* of Europe, and *S. Macrai* of Chili. Mr. Bentham, however, entirely dissents from such an opinion after a most careful review of its characters, and, I need hardly add, that on his knowledge and experience we may rely for the validity of the species.

XXXVII. PRIMULACEÆ, Juss.

1. PRIMULA, L.

1. PRIMULA farinosa, Linn. Sp. Pl. p. 205. Engl. Bot. t. 6. Duby in DC. Prodr. vol. x. p. 44. Gaud. in Ann. Sc. Nat. vol. iv. p. 102, ct in Freyc. Voy. Bot. p. 133. D'Urville in Mém. Soc. Linn. Paris, vol. iv. p. 606.

Var. β, Magellanica. P. Magellanica, Lehm. Monogr. Prim. p. 62. t. 6. Duby, in DC. Prodr. vol. x. p. 45. P. decipiens, Duby, in DC. l. c. (TAB. CXX.)

HAB. Strait of Magalhaens to Cape Horn, Commerson, Capt. King, C. Darwin, Esq., J. D. H. Falkland Islands, most abundant, Gaudichaud, Sc.

The excellent plate, executed for this work by Mr. Fitch, enables the British botanist to form a just idea of the Antarctic state or variety of *P. farinosa*, L.; which, it will be seen, differs from the majority of those of Britain in the short peduacles of the white flower, in the position of the stamens, in the tube of the corolla, and in the colour of the flowers. The first of these characters is constant in all the Falkland Island and Magellanic specimens of this species, but is also seen in an individual of *P. farinosa*, gathered near Settle in Yorkshire by Mr. Tatham, for which I am indebted to my friend Mr. Watson, who not content with examining this plant with me, had the kindness to collate a suite of Antarctic specimens with many hundreds of British growth. The result of this examination has been, that except, perhaps, the colour of the flower, there is no constant character to distinguish the races of the opposite hemispheres, neither the length of the pedicels, of the calycine segments, of the tube of the corolla, nor the position of the stamens in the latter. If, again, we grant (with M. Duby) that the *P. Sectica*, Hook., is not even a variety of *P. farinosa*, the length of the pedicel is of still less value, for the North Scottish individuals are undistinguishable, except by the colour of the Strait of Magalhaens, by Mr. Darwin, these localities being the northern and southern extremes of its range in the Southern Hemisphere.

Lastly, on comparing var. β with foreign examples of *P. farinosa*, their identity is still more evident; for the latter attain the same great size in Austria that the var. β often does in the Falkland Islands, whilst Arctic American specimens of the two are entirely alike.

One argument which militates against the common origin of the individuals from the opposite hemispheres, must not be overlooked; it is the absence of the plant, and, indeed, of the whole genus, in any part of the Andes south of 39° north lat.; a circumstance which makes it very difficult to account for its appearance in the two opposite temperate zones, if all the individuals of both hemispheres are supposed to have sprung from one parent.

PLATE CXX. Fig. 1, flower; fig. 2, the same; fig. 3, the same laid open; fig. 4, ripe capsule; fig. 5, seed; fig. 6, longitudinal section of the same :---all magnified.

2. ANAGALLIS, Tourn.

1. ANAGALLIS alternifolia, Cav. Icones, vol. vi. p. 3. t. 506. f. 2. Duby in DC. Prodr. vol. x. p. 71.

Var. densifolia, Lysimachia repens, D'Urville, in Mém. Soc. Linn. Paris, vol. iv. p. 606. Gaud. in Freyc. Voy. Bot. p. 133. Hook. Ic. Plant. t. 536.

HAB. Strait of Magalhacns; Port Famine, Capt. King; Wollaston Island, C. Durwin, Esq.; Falkland Islands, D'Urville, Mr. Wright, J. D. H.

It appears to me that two very distinct species of this genus have been confounded, partly together, and partly with the *A. tenella*, L., of Europe. The first is confined to the damp western portions of middle and southern Chili, Fuegia, and the Falkland Islands, and there are two or perhaps three varieties of it; 1 take it to be the *A. alternifolia* of Cavanilles, a variable plant, with the peduncles of the same length as, or not much exceeding, the leaves, and the capsule shorter than the calyx. The figure of that author is very inaccurate and at variance with his description; for the plant is represented erect, instead of creeping, and the leaves scattered, though said to be, approximate. Supposing Cavanilles' plant to form one variety of *A. alternifolia*, a second is larger and also creeping, with prostrate branches, S-10 inches long, bearing broader, rounded and more acute leaves; it has been collected in Valparaiso by Mr. Cuming and Mr. Bridges. A third, intermediate between this and the Fuegian form, has the leaves more crowded, ovate-oblong, and smaller; it is possibly the state figured by Cavanilles, and has been gathered at Concepcion by Capt. King, at Valdivia by Mr. Bridges, and on the Andes of Mendoza by Dr. Gillies (*Ruellia caspitosa*, Gill. MSS.; and *Anagallis herpestoides*, Gill. MSS.). The fourth variety is what I have called *densifolia*; its leaves and stem are much smaller and crowded, and the whole plant is sneculent.

Another extra-tropical South American *Anagallis* is the *A. filiformis*, Link, (*A. tenella*, β . *filiformis*, St. Hil.), which approaches *A. tenella* so very closely, that M. St. Hilaire has united them specifically. It differs from *A. alternifolia* in the leaves being opposite, the stem slender, the peduncles longer, the calycine pieces narrower and twice as long as the capsule, and the whole plant not so succedent; from the European *A. tenella* in the leaves never heing so broad, in the longer peduncles and rigid stems.

The variety *densifolia* has a large capsule, always equalling the calyx in length, thus differing from the plant figured by Cavanilles. The capsule, though described by D'Urville as having the dehiscence of a *Lysimachia*, evidently opens transversely in the specimens I have examined, though it is sometimes, from pressure, split at the top also. Its habit resembles the Abyssinian *A. serpens*, Hochst.

3. SAMOLUS, L.

1. SAMOLUS littoralis, Brown, Prodr. p. 428. Duby in DC. Prodr. vol. x. p. 73. Scheffieldia repens, Forst. Nov. Gen. p. 18. t. 9.

HAB. Chonos Archipelago and Cape Tres Montes, C. Darwin, Esq.

A plant common to New Holland, New Zealand, and South Chili, and very variable in the size of its parts in all these countries. I have not seen Chilian specimens from a lower latitude than Valdivia, between which and Cape Tres Montes it seems limited.

2. SAMOLUS spathulatus, Duby, in DC. Prodr. vol. x. p. 74. Androsæa spathulata, Cavanilles Icones, vol. v. p. 56. t. 484. f. 1.

HAB. Strait of Magalhaens; Port Gregory, Capt. King. Elizabeth Island, C. Darwin, Esq.

The raceme, in most of Capt. King's specimens, is so much abbreviated that the flowers are almost capitate. The range of the species, between Port Desire and the Strait of Magalhaens, is remarkably limited.

XXXVIII. LENTIBULARIEÆ, Rich.

1. PINGUICULA, Linn.

1. PINGUICULA Antarctica, Vahl, Enum. p. 192. Alph. DC. Prodr. vol. x. p. 31. P. obtusa, Banks et Sol. MSS. in Bibl. Banks. (TAB. CXIX.)

HAB. Strait of Magalhaens; Port Famine, Capt. King; Good Success Bay, Banks and Solander; south part of Fuegia, C. Darwin, Esq.; Hermite Island, Cape Horn, J. D. H.

A very pretty little plant, the representative of the British *Pinguicula Lusitanica*, L., from which it differs in the narrower segments of the corolla and shorter spur. It is not uncommon on moist rocks in Fuegia.

PLATE CXIX. Fig. 1, lateral, and fig. 2, front view of tlower; fig. 3, calyx, germen, and stamens; fig. 4, stamen; fig. 5, germen; fig. 6, transverse section of the ovarium; fig. 7, ripe fruit; fig. 8, seed; fig. 9, embryo: ---all magnified.

XXXIX. PLUMBAGINEÆ, Juss.

1. STATICE, Tourn.

STATICE Armeria, Linn., Sp. Pl. p.394. Engl. Bot. t. 226. S. cæspitosa, Poiret, Encycl. p. 235. Gaud. in Ann. Sc. Nat. vol. v. p. 102. D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 606.

Var. B, alpina; Ed. Cat. p. 2. Hook. Brit. Fl. p. 270.

HAB. Var. a, Strait of Magalhaens, Commerson; Port Famine, Capt. King; Falkland Islands, most abundant near the sea; Gaudichaud, Sc. Var. β, on the mountains of Fuegia, C. Darwin, Esq., J. D. H.

There can, I think, be no question as to this being identical with the *S. Armeria* of the northern hemisphere; if any specific or other distinction exists, it has eluded Mr. Watson's and my examination. Both as an alpine and especially as a sea-side plant, its habits are those of the common Sca-Pink.

XL. PLANTAGINEÆ, Venten.

1. PLANTAGO, Linn.

1. PLANTAGO maritima, Linn., Sp. Pl. p. 165. Engl. Bot. t. 175. P. juncoides, Lam. Illust. Gen. n. 1683.

HAB. Strait of Magalhaeus; Port Famine and Port Gregory, Capt. King.

I am not aware of any South American stations for this plant except those mentioned above; it is also a native of the Cape of Good Hope, but not of Australia or New Zealand.

2. PLANTAGO *barbata*, Forst.; laxe cæspitosa simplex v. ramosa, foliis erectis stellatim patentibus recurvisve lineari-lanceolatis anguste lineari-elongatisve subacutis carnosis remote dentatis basi scariosis barbatis glabratisve, pedunculis folio subæquantibus, spicis 1-3-floris, capsulæ late obovatæ mcdio circumscissæ parte inferiore calycem vix excedente. P. barbata, *Forst. Comm. Goett.* vol. ix. t. 4. P. pauciflora, *Lam. Illust. Gen.* n. 1684. P. pauciflora, β , parva, *Barneoud Monogr. Plantag.* p. 17. P. polymorpha, *Banks et Sol. MSS. in Bibl. Banks. cum icone.*

Var. a, barbata; foliis stellatim patentibus spathulato-lanceolatis dentatis basi barbatis.

Var. β , *elongata*; caule simpliciusculo, foliis erectis anguste et longissime lineari-spathulatis obtusis remote sinuato-dentatis basi barbatis.

Var. γ , *imberbis*; caule ramoso, foliis patulis lanceolatis obtusis remote dentatis basi sub-barbatis. P. imberbis, *Hook. fil. MSS. in Part* 1. p. 66.

HAB. Var. a, Strait of Magalhaens, Commerson; Tierra del Fuego, Banks and Solander, Forster. Var. β, Port Gregory, Capt. King. Var. γ, Port Famine, Capt. King.

A highly variable plant; always, however, in all the specimens which I have examined, retaining the characters of a short capsule dehiscing across the middle, the broad lower half of which is as long as, or very little longer than the calyx, and of a different form from the narrow obconical elongated analogous organ of P. monanthos,

D'Urv. The nearest allies of this plant are its Australian and New Zealand representative, the *P. carnosa*, Br. (vid. Flor. Antarct. Pt. 1. p. 65), and the *P. Andicola*, Gill. MSS.; the former of these differs in its shorter eapsule, differently shaped seeds, erowded, more fleshy foliage; the latter in its eurious root and broader leaves, which are, however, very unimportant characters.

The Port Gregory specimens, of which I have made variety β , are perhaps drawn up, for the leaves are six inches long, which is at least four times the length of those of the ordinary state of the species.

3. PLANTAGO monanthos, D'Urv.; caulibus ramosis dense cæspitosis, foliis perplurimis basi arcte vaginantibus erectis subsquarrosis stellatim patentibusve anguste lineari-elongatis obtusis obscure dentatis marginibus cartilagineis pedunculis multoties longioribus basi glaberrimis, spiculis 1–4-floris, capsula anguste clavata infra medium circumscissa parte inferiore calyce bis terve longiore. P. monanthos, D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 606. Gaud. in Freye. Voy. Bot. p. 133. Barneoud, Monogr. Plantag. p. 17. exclud. syn. P. carnosæ, Br. (TAB. CXXI.)

Var. a, foliis erectis subsquarrosisve lineari-elongatis flaccidis.

Var. β , *albreviata*; caulibus cæspitosis, foliis brevioribus substellatim patenti-recurvis.

Var. γ , *muscoides*; caulibus densissime cæspitosis, foliis brevibus arcte imbricatis marginibus cartilagineis albis.

HAB. Var. α, Falkland Islands, D'Urville, J. D. H.; Hermite Island, Cape Horn, in moist places, J. D. H. Var. β, Hermite Island, amongst rocks; var. γ, the same locality, in clefts of exposed rocks, J. D. H.

A very different plant from the *P. carnosa*, Br., under which M. Barueoud has included it, especially in the habit, stems, foliage, comparative length of the peduncle, shape of the capsule and form of the seeds.

PLATE CXXI. Fig. 1, flower and braeteæ; fig. 2, germen; fig. 3, ripe fruit; fig. 4, transverse section of ditto; fig. 5, upper half of ditto, with dissepiment and seeds; fig. 6 and 7, dissepiment and seeds; fig. 8 and 9, front and back view of seeds; fig. 10, longitudinal section of seed :---all magnified.

PLANTAGO hirtella, H. B. K., Nov. Gen. et Sp. vol. x. p. 187. t. 127. Barneoud, Monogr. Plantag. p. 18. HAB. South Chili; Cape Tres Montes, C. Darwin, Esq.

Not at all an uncommon Chilian and Buenos Ayrean plant, exceedingly variable in the breadth and pubescence of the leaves, and also in the size of the flowers, which in these specimens are larger than in the figure quoted. It is also a Brazilian species, and occurs on the mountains of Peru and Columbia, and is very nearly allied to the *Pl. Virginica*, L. The character of the segments of the eorolla being patent or conniving is searcely tenable in this plant and its allies.

XLI. POLYGONEÆ, Juss.

1. POLYGONUM, L.

1. POLYGONUM maritimum, Linn. Sp. Pl. p. 519. Engl. Bot. Suppl. t. 2804. Meisner, Monogr. Polyg. p. 89.

HAB. South Chili; Cape Tres Montes, C. Darwin, Esq.; Strait of Magalhaens; Port Famine, Capt. King.

There appears no difference between these specimens and those of British growth. In the southern hemisphere the species occurs only at the Cape of Good Hope and in South Chili.

2. RUMEX, L.

1. RUMEX cuneifolius, Campd., Monogr. des Rum. p. 95. Fl. Antarct. pt. 1. p. 67.

HAB. South Chili; Chonos Archipelago, C. Darwin, Esq.

2. RUMEX crispus, Linn., Sp. Pl. p. 476. Engl. Bot. t. 1998. R. Patientia (?), Gaudichaud in Ann. Sc. Nat. vol. vi. p. 101. D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 605.

HAB. Falkland Islands; Berkeley Sound, undoubtedly introduced.

My specimens, though imperfect, are, I think, referable to this species.

3. RUMEX Acetosella, Linn., Sp. Pl. p. 481. Engl. Bot. t. 1674. Gaudichaud and D'Urville, l.c.

HAB. Falkland Islands; abundant near the settlements and on the mountains; Gaudichaud, Sc.

This, and the *R. Acetosa*, L., included in Gaudichaud's list, I consider undoubtedly as introduced plants, of which the seeds, being caten by the birds, are by their agency transported to otherwise inaccessible cliffs.

XLII. CHENOPODIACEÆ, Juss.

1. CHENOPODIUM, L.

1. CHENOPODIUM glaucum, Linn., Sp. Pl. p. 320. Engl. Bot. t. 1454.

Var. B, divaricatum; prostratum, ramosum, ramis gracilibus divaricatis.

HAB. Var. B, Chonos Archipelago; C. Darwin, Esq.

Evidently the *C. glaucum* of Great Britain, though the stem is more diffusely branched than in most English individuals. A precisely similar variety inhabits British North America, but I have seen no specimens from any part of the New World between that country and South Chili.

2. CHENOPODIUM macrospermum, Hook. fil.; glaberrinum, non glaucescens, caulibus validis succulentis basi divaricatim ramosis, foliis petiolatis deltoideo-oblongis obtusis sinuatis carnosis, racemis compositis densifloris aphyllis bracteatis, seminibus majusculis erectis subtilissime reticulatis.

HAB. Falkland Islands; Berkeley Sound and St. Salvador Bay, near the sea; C. Darwin, Esq., J. D. H.

Caules e radice descendente fusiformi solitarii v. plurimi, prostrati, 3-5-unciales, canaliculati v. angulati, crassi, diametro pennæ anserinæ. Folia longe petiolata, petiolo $\frac{1}{2} - \frac{3}{4}$ unc. longo, lamina æquilonga carnosa, utrinque opaca, siccitate flavo-virescentia. Flores fructusque multoties majores quam in affinibus.

This very distinct species has been used as a pot-herb by the colonists of the Falkland Islands, and was described to me as excellent. The great size of the seed at once distinguishes it from its nearest European allies, *C. rubrum*, L., and *C. polyspermum*, L. I have not included these two species under the genus *Blitum* because the seeds of *C. glaucum* are more frequently horizontal than erect, and neither of them possesses a calyx which is materially thickened after flowering.

XLIII. PROTEACEÆ, Juss.

1. EMBOTHRIUM, Forst.

1. EMBOTHRIUM coccincum, Forst., Gen. Plant. t. S. Comm. Soc. Reg. Goett. vol. ix. p. 24. Lamarck, Encycl. vol. ii. p. 351. Illust. Gen. n. 1284. t. 55. f. 2. Brown, in Linn. Soc. Trans. vol. x. p. 196.

4 G

HAB. Strait of Magalhaens, Commerson; Port Famine, Cupt. King; Fuegia, Banks and Solander, Forster, Sc.

This very handsome plant scems confined to the extreme sonthern part of South America, without, however, reaching Cape Horn itself; it is very nearly allied to the Chilian *E. lanceolatum*, R. and P., but differs in the nervation of the leaves.

2. LOMATIA, Brown.

1. LOMATIA ferruginea, Brown, in Linn. Soc. Trans. vol. x. p. 200. Embothrium ferrugineum, Cavanilles, Icones, vol. iv. p. 59. t. 385.

HAB. Chonos Archipelago, C. Darwin, Esq.

Like the former, this species has a very confined range, inhabiting the country between Valdivia and the Chonos Archipelago, including Chiloe, on the west side of the Andes only; and, according to Cavanilles, it is limited to places occasionally overflowed by the sea. Mr. Bridges states that the native name is "Romarilla".

XLIV. SANTALACEÆ, Br.

1. NANODEA, Gærtner, fil.

1. NANODEA muscosa, Gærtner, de Fruét. vol. iii. p. 251. t. 225. Gaud. in Ann. Sc. Nat. vol. v. p. 101. t. 2. f. 3, et in Freyc. Voy. Bot. p. 442. D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 605. Banks et Sol. MSS. in Bibl. Banks. cum icone. Balexerda muscosa, Commerson, MSS.

HAB. Strait of Magalhaens, Commerson; Port Famine, Capt. King; Fuegia, Good Success Bay, Banks and Solander; Hermite Island, J. D. H.; Falkland Islands, very common; Gaudichaud, &c.

2. ARJOONA, Car.

1. ARJOONA Patagonica, Homb. et Jacq.; stricta, erecta, ramosa, ramis simplicibus glaberrimis, foliis sparsis patulis breviter subulatis rigidis glaberrimis nervosis, inflorescentia sericeo-tomentosa capitata, bracteis concavis acutis tubo perianthii $\frac{1}{2}$ brevioribus. A. Patagonica, Homb. et Jacq. in Voy. au Pole Sud, Bot. Dicot. t. 15. A. sine descript.

HAB. Strait of Magalhaens; Port Peckett, Messrs. Hombron and Jacquinot.

This plant, of which I have examined specimens gathered by Capt. King on the Patagonian coast, is very probably only a variety of the A. tuberosa, Cav., of the same country, which varies in the size of the leaves, and in their being smooth, public, or tomentose. In Capt. King's specimens they vary from 1-4 lines long.

2. ARJOONA *pusilla*, Hook. fil.; caule erecto gracili simplici v. diviso, foliis flaccidis sæpius recurvis elongato-linearibus acuminatis marginibus glaberrimis subenerviis, floribus paucis, bractca exteriore majuscula cymbiformi obtusa glabrata, corolla extus sericco-tomentosa fance ampliata inter stamina fasciculis inconspicuis pilorum articulatorum ancta, stigmatibus 3 brevibus.

HAB. Strait of Magalhaens; Port Gregory, Capt. King; Cape Negro, C. Darwin, Esq.

Herba bi-tri-pollicaris. Caulis gracilis, erectus, simplex v. basi bis terve divisus. Folia flaccida, suberecta, $\frac{1}{2}-\frac{3}{4}$ unc. longa, sub 1 lin. lata, medio uninervia, apicibus acuminatis marginibus plerumque recurvis. Bractea $2\frac{1}{2}$ lin. longa, dorso glabrata, marginibus ciliatis, bracteolis interioribus in tubum apice inæqualiter 3-4-fidum ovario

FLORA ANTARCTICA.

subadhærentem obseure coalitis, extus pilosis. Perianthium $\frac{1}{2} - \frac{3}{4}$ unc. longum, extus pilis fulvis tomentosum, tubo gracili superne ampliato, laciniis ovato-oblongis, fance inter stamina barbata, pilis brevibus flaccidis articulatis creberrimeque transversim striatis. Stamina filamentis breviusculis, antherarum apicibus exsertis. Stigmata 3 parva, inter tubum corollæ retracta.

A very distinct species from the former, in the foliage especially. Hitherto it has been found in the Straits of Magalhaens only.

XLV. THYMELEÆ, Juss.

1. DRAPETES, Lam.

1. DRAPETES muscosa, Lamarck, Journ. d'Hist. Nat. vol. i. p. 186. t. 10. f. 1. Gærtner, de Fruct. vol. iii. p. 199. t. 215. Juss. in Annales du Mus. vol. vii. p. 479. Poiret, Encycl. Suppl. vol. ii. p. 523. t. 915. f. 1. D'Ure. in Mém. Soc. Linn. Paris, vol. iv. p. 605. Banks et Sol. MSS. in Bibl. Banks. cum icone.

HAB. Strait of Magalhaens, Commerson; and throughout Fuegia, on the mountains, Banks and Solander, Capt. King, Sc. Falkland Islands, D'Urville, J. D. H.

A curious little plant, confided in its geographical range to the mountains of Antarctic America, and represented in New Zealand by a very similar one, forming its only congener, the *D. Dieffenbachii*, Hook. (Lond. Journ. of Bot. vol. ii. p. 497. t. 17). However similar the two plants are in habit and in their more important structural characters, differences exist which some botanists may deem of generic value; these arc, the cylindrical continuous base of the perigonium, thickened faux and capitate glandular stigma of the New Zealand species, contrasted with the jointed angulated tube of the perigonium in the Antarctie American plant, which has an eglandulose faux and plunose stigma. The thickening of the throat of the perianth in *D. Dieffenbachii*, which almost causes the faux to be closed with scales, is effected by the three nerves of each segment being there joined by anastomosing venules, whilst in *D. muscosa* they run free to the apex of the segment.

XLVI. URTICEÆ, Juss.

1. URTICA, L.

1. URTICA *Darwinii*, Hook. fil.; caule gracili erecto sparsissime piloso v. glaberrimo, foliis membranaceis oppositis petiolatis ovatis acuminatis grosse æqualiter crenato-serratis basi rotundatis 3-nerviis utrinque subtilissime punctatis tenuiter puberulis, petiolo gracili, stipulis lineari-oblongis subacutis, floribus glomeratis glomerulis setosis in spicas graciles interruptas petiolo longiores dispositis.

HAB. Chonos Archipelago, C. Darwin, Esq.

Caulis penna corvina tenuior, flaccida, glaberrima, v. pilis raris albidis valde inconspicuis sparsa, internodüs $1\frac{1}{2}$ uncialibus. Stipulæ 3 lin. longæ, subacutæ. Petioli $\frac{1}{2}-\frac{3}{4}$ unc. longi, graciles parce puberuli. Folia 2-3 unc. longa, $1\frac{1}{4}-1\frac{1}{2}$ lata, grosse crenato-serrata, segmentis sinubusque latis acutis. Pedicelli axillares, subquaterni, patuli, penduli, petiolo $\frac{1}{2}$ v. bis longiores. Flores in glomerulos sparsos congesti; glomerulis setosis, paucis inferioribus masculis ceteris fœmineis.

In appearance this very closely resembles the *Pilea pumila* of North America, though it is more nearly related to the *Urtica gracilis* of the United States. Both this latter plant and the *U. Darwinii* differ from *U. dioica*, L., in the much larger flowers and achænia.

2. URTICA Magellanica, Poir.; caule valido crecto hispido-setoso, foliis subcoriaceis rugosis oppositis petiolatis ovatis ovato-lanceolatisve acuminatis basi cordatis argute serrato-dentatis, utrinque setosis subter leviter puberulis, stipulis lineari-oblongis acutis, floribus glomeratis, glomerulis setosis in spicas interruptas petiolo breviores v. clongatas dispositis. U. Magellanica, *Poiret, Encycl. Suppl.* vol. iv. p. 323.

HAB. Strait of Magalhaens, Commerson; Port Famine, Capt. King.

Caulis 2-pedalis crectus, validus, setis plurimis patentibus obtectus. Petioli $\frac{1}{4}$ -I une. longi. Folia $2\frac{1}{2}-3\frac{1}{2}$ uncialia, latitudine varia, basi plus minusve cordata rarius rotundata. Racemi seu spice penduli, monoici v. dioici, petiolo longiores rarius abbreviati. Flores majusculi, fœminei compressi, orbiculares, achænio conformes.

Not an uncommon species from Valparaiso to the Strait of Magalhaens, differing from the preceding in its robust habit, different texture of the leaves, and setose stem and foliage, all, I fear, very unimportant characters in this genus, but whose validity in the present species I have not sufficient materials for ascertaining. The characters drawn from the length of the racemes is a variable one, those bearing male flowers especially being the shortest, and sometimes, as described by Poiret, shorter than the petioles. The present appears very closely allied indeed to a South African species, and it may even be considered doubtful whether both are not states of *U. dioica*, with unusually large tlowers.

One of Anson's vessels, when detached from his squadron, put into a Bay near the western entrance of the Strait of Magalhaens, and recruited her crew, who were paralyzed by scurvy, by means of Nettle tops, most probably the produce of this or the former species.

The Urtica lanrifolia, Poiret, stated to have been brought from the Strait of Magalhaens by Commerson, does not appear to belong to this genus. I am wholly unacquainted with the U. gigantea, of the same author, also from the Strait of Magalhaens.

2. PILEA, Lindl.

1. PILEA *clliptica*, Hook. fil.; suberecta, caule debili herbaceo parce ramoso, foliis longe et graciliter petiolatis membranaceis ellipticis utrinque subobtusis grosse crenato-serralis trinerviis super subterque pilis appressis minimis conspersis, floribus masculis in umbellam capitatam longe pedicellatam congestis, fœmineis ad basin pedunculi sessilibus glomeratis, achænio orbiculari compresso apice oblique emarginato.

HAB. Chonos Archipelago; C. Darwin, Esq.

Caules uni-bipedales, crassitie pennæ corvinæ, rufescentes, punctis albidis elongatis notati. Petioli longitudine varii folio longiores v. breviores. Stipulæ membranaceæ, late ovatæ. Folia læte viridia, membranacea, exacte elliptica, imo basi obscure cordata, magnitudine varia, $\frac{1}{2}$ -2 unc. longa, grosse sed equaliter crenato-serrata; parenchyma corpusculis fusiformibus e epidermide translucida oculo nudo manifestis pilos appressos simulantibus farctum. Pedunculi petiolo æquilongi v. longiores, apice umbellulam simplicem florum masculorum gerentes, basi glomerulo florum fœmineorum aucti. Fl. MASC. Perianthium 4-partitum, laciniis late ovatis acuminatis inflexis. Fl. FœM, Perianthium valde compressum, 3-partitum, lacinia postica cucullata lateralibus oblongis multotics longiore. Achænium planum.

A very distinct species, confined to the S.W. portions of Chili between Valdivia and the Chonos Archipelago, a tract which may be considered as partaking of the Chilotean botany, the latter itself being a division of the Chilian Flora, only separable by the amount of specific difference from the other extra-tropical regions of western South America.

The appearance of the so-called pubescence of this species and many other *Urticeæ* is curious, and caused by the presence of numerous white fusiform raphides attenuated at both ends, which are scattered abundantly throughout

the parenelyma of the leaves and immediately beneath the surface of the stem; from the tenuity of the epidermis, and transparency of the leaves when dried, they form prominences on the euticle of a white colour, closely simulating the laterally attached hairs of Crucifera.

XLVII. EMPETRACEÆ, Nutt.

1. EMPETRUM, L.

1. EMPETRUM rubrum, Vahl, MS. et Willd. Sp. Pl. vol. iv. p. 713, ex Banks et Sol. MS. in Bibl. Banks. cum icone. Gaud. in Ann. Sc. Nat. vol. v. p. 103, et in Freyc. Voy. Bot. p. 134. D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 608. "Bruyère à fleurs d'un vert blanchatre," Pernetty, Voy. vol. ii. p. 64.

HAB. South Chili, Fuegia, and the Falkland Islands, most abundant, Commerson, Banks and Solander, and all future voyagers.

I am unable to detect any characters to separate the *Empetrum rubrum* from *E. nigrum*, beyond what is afforded by the colour of the berries. Though many of the northern specimens of *E. nigrum* are perfectly similar to Fuegian specimens of *E. rubrum* in every other respect, yet almost all the Falkland individuals, and many of those of Cape Horn, are more tomentose than any specimens of the Northern species that I have examined, Under these circumstances, the plants from the opposite hemispheres may be regarded as representative species, or varieties of the same; but, since all the specimens from the southern hemisphere present one constant character, distinguishing them from those of the northern, and since neither is known to occur in any part of the New World between the parallels of 45° N. and 33° S., I feel myself obliged to attach specific importance to the otherwise very trifling differences in the colour of the fruit.

The Empetrum rubrum is a very abundant western extra-tropical South American plant, from the latitude of Concepcion on the Pacific coast, and Mendoza on the Andes, to Cape Horn. In the latter country, as in the Falkland Islands, this species altogether simulates *E. nigrum* in the localities it affects, in its habit and mode of growth, stature, in the forms its varieties assume, and in the economy of nature, affording food to wild-geese, and, in Fuegia, to a bird allied to the grouse. The stems and leafy branches are much used for fuel in the Falklands, where the plant is called "Diddle-dee", they are especially employed in kindling fire, for even when sodden with rain, they speedily ignite, and burn with a bright and hot flame.

The affinities of this genus, or rather order, are yet undefined. I am inclined to adopt the opinion of Jussieu in allying it to *Ericeæ*, from the habit, foliage, the bracteæ, ealyx, and texture of the eorolla and anthers and some other characters.

XLVIII. CUPULIFERÆ, Rich.

1. FAGUS, L.

1. FAGUS Antarctica, Forst., ex Banks et Sol. MS. in Mus. Banks. cum icone. Hook. Bot. Journ. vol. ii. p. 15. t. VI. Calueechinus Antarctica, Homb. et Jacq. in Foy. au Pole Sud, Bot. Dicot. t. 14. Z. et Bot. Monocot. Phan. t. 6. O. C. Montagni, Homb. et Jacq. l. c. Bot. Dicot. t. 8. H. (TAB. CXXIII.)

HAB. South Chili and throughout Fuegia, very abundant, Commerson, Banks and Solander, and all succeeding voyagers.

This species and the following, form together so predominant a feature in the Fuegian landscape, that, though accurately described by several voyagers, especially Cook, King, and Fitzroy, and in the graphic narrative of my

friend Mr. Darwin, it is advisable to sum up the principal facts connected with their history, adding some little from personal observation. These remarks will be the more appropriately introduced here, from the two principal species having been first imported into England by the Antaretic Expedition, and now promising to become useful and ornamental additions to our forests; as, also, from their geographical range having been used as an indication of the limits of the Antaretic Flora.

The Fague Antarctica has always been recognized as a true Beech, from the very marked resemblance its deciduous foliage bears to that of the European F. sylvatica. The other common Fuegian species, F. betuloides, with coriaceous leaves of a deep green hue very similar to those of the Scottish Birch, was, up to the discovery of its flowers, considered to be a Betula. The habit of both species, however, is essentially that of the Beech, and so are the form of trunk, smooth bark, and especially the spreading ramification and horizontal divarienting ramuli; whilst their flowers and fruit resemble so closely in all but size, those of the European Fague sylcatica, that I consider them as undoubted eongeners of that plant.

I have elsewhere (vid. ante p. 277) alluded to the very common error of holding the locality in which a certain species particularly abounds, to be the principal habitat of the order or genus to which it belongs; this often arises from attaching a greater importance to the spread of the species than that of the genus. Naturalists unaequainted with the range of the Beeches, will be surprised to hear that they are more characteristic of the temperate and cold latitudes of the southern, than of the northern hemisphere, even in the proportion of five to one. Thus, one species alone is European, and one American; two are found on the mountains of Java; one is characteristic of the Alps of Tasmania, where the only Antaretic representatives of the Australian Flora are found; four inhabit the high mountains of the northern or lower levels in the middle and sonthern Islands of New Zealand; and, lastly, as many as seven * have been described from Chili and Fuegia.

Of the seven so called Chilian and Fuegian species, three are well marked, and afford instructive examples of the succession of species in proceeding northward from Cape Horn; they are the *F. Antarctica, betaloides*, and *obliqua*; the others, which may be varieties of the above, though from the want of copious suites of good specimens I advance this opinion with much hesitation, are *F. Pumilio*, Poepp. and Endl., *F. procera*, P. and E., *F. Dombeyi*, Mirb., and *F. alpina*, P. and E.

The Fagus Antarctica, justly so named, ascends even at Cape Horn much higher than F. betaloides, and nearly to the summits of the mountains, which are perhaps 1000 feet Lelow the assumed level of perpetual snow in that latitude, while at the sea it forms much the larger tree of the two. Supposing the continent of America to have been produced indefinitely to the southward, in a free ocean, the F. Antarctica would be found extending to as high a parallel as 62° S., whilst the F. betaloides would cease at the 60th degree: assuming that both species followed the same ratio of ascent that very many other Cordillera plants do, which ascend from the level of the sea in Fuegia to a considerable elevation in a lower latitude.

Fagus betaloides, though by far the most prevalent species in Hermite Island, and, indeed, throughout Fuegia, has its principal parallel about the Strait of Magalhaens, where it becomes a very large tree. It forms the prevailing feature in the seenery of Tierra del Fuego, especially in winter time, from having persistent, evergreen leaves, and from its upper limit being sharply defined and contrasting with the dazzling snow that covers the matted but naked branches of the *F. Antarctica*, which immediately succeeds it. Its upper limit at Cape Ilorn (lat. 56°) is about 800 feet; in the northern parts of Tierra del Fuego it reaches 1,400 feet; and, if the *F. alpina*, P. and E., be a state of the same species in its most northern locality, its level in lat. 36° is between 5,000 and 8,000 feet.

The following notice of the dimension the Evergreen Beech attains in the Strait of Magalhaens, is extracted from Capt. King's excellent 'Voyage of the Adventure and Beagle' (p. 576). "At Port Famine and in the neigh-

* An eighth, *F. glutinosa*, Poepp., is no *Fagus* at all, but, as my friend Mr. Miers assures me (and he has examined authentic specimens in M. Delessert's Herbarium), a species of *Eucryphia*.

FLORA ANTARCTICA.

bourhood, the Evergreen Beech grows in the greatest abundance and reaches a very large size. Trees of this species three feet in diameter, are abundant; of four feet there are many; and there is one tree (perhaps the very same noticed by Commodore Byron), which measures seven feet in diameter for seventeen feet above the roots, and then divides into three large branches, each of which is three feet through. This venerable tree seemed to be sound, but from our experience of several others that were cut down, might be expected to prove rotten in the centre. This tendency to decaying in the heart may be attributed to the coldness of the schistose sub-soil upon which the trees are rooted, as well as the perpetual moisture of the elimate."

The wood of these trees Capt. King describes as being heavy and far too brittle for masts, or even boat-hook staves; hut it euts up into tolerable planks, which, if seasoned, might serve for ship-building. During our sojourn in Hermite Island, Capt. Ross caused several thousand small trees, of both species, to be felled and barked; these we transported to the Falkland Islands, in which tree-less country they were highly prized for roofing houses, &e. The decidnous species appeared to afford the better wood of the two.

A more striking contrast between two so very closely allied plants, cannot well be imagined, than between *F. Autorctica* and *F. betaloides*, arising from the evergreen foliage of the latter being of a totally different texture and aspect from that of the former. Surely so strongly marked a difference between otherwise very nearly allied species, growing side by side under perfectly similar conditions, is a strong argument in favour of their being originally separate creations. We see, too, how the adaptation of particular forms of vegetation to certain elimates, even in this remote quarter of the globe, is exemplified in these trees; though both do grow together abundantly, they still have their preferences, the evergreen glossy foliage prevailing on the western coast, where the climate is damp and equable, whilst the deciduons-leaved plant seeks the heights more exposed to the vieissitudes of the weather, or the drier eastern parts of Fuegia, where the *F. betaloides* will not succeed. So it is with us in Great Britain; our glossy-leaved evergreens, whether native or introduced, thrive best in the climate of the west coast, where the summers are colder, the winters warmer, and all the seasons more humid than they are on the east.

The third species of *Fagus*, the *F. obliqua*, replaces *F. Antarctica* in South Chili, occupying the flanks of the Andes, between the altitudes of 1,000 and 5,000 feet, where it is the prevailing forest-tree. It appears to inhabit the level of the sea in the parallel of the Strait of Magalhaens, and is probably the third species of Beech alluded to by Capt. King (l. e. p. 576), for that voyager does not seem to have distinguished the *F. Pumilio* as a species.

The accompanying cut will explain better than words, the order of succession in latitude and in elevation that South American Fagi follow. Their southern ranges may be ascertained with tolerable precision, the exact altitude they attain in the two northern positions is more doubtful. The positions taken are, commencing from the southward, 1st, Hermite Island, lat. 56° ; 2nd, Strait of Magalhaens, lat. 54° ; 3rd, Antuco in Chili, lat. $36^{\circ}40'$. The upper eurve indicates the lower level of perpetual snow; the others, the upper limit of the tree whose name is found immediately under.

From the want of a suite of specimens I eannot speak with much confidence of the Chilian species, F. alpinaand F. Pumilio, the first appears, from the plate and description, a variety of F. betaloides, and, from occupying the position that F. betaloides would hold relatively to the others in South Chili, I have introduced it as such into the cut. The F. Pumilio is even more probably a variety of F. Antarclica. Both are said to occupy great heights in South Chili, the latter indeed only existing there as a stunted tree. There are still two other South Chilian species, F. procera, P. and E., and F. Dombeyi,* Mirb.; they inhabit the level of the sea in the parallel of 41°. The first I am inclined to regard as a variety of F. obliqua, or rather a large-leaved state of that plant descending to the eoast; the second is a similar form of F. betuloides. If my supposition prove correct, both species afford examples

* This is one kind of "Roble" of Capt. King's Narrative (p. 280); in speaking of the woods of Chiloe, he says, "Roble, (*Fagus obliqua*, Mirb.), is a large tree, and, from the durable quality of its timber, considered the

348

of mountain plants, having the upper limit in elevation which they attain sharply defined, throughout several degrees of latitude, but which descend and assume other aspects in a warmer climate. This, also, I have attempted to express on paper by dotted lines drawn down to the sea-level, from the Chilian positions of F. betaloides and F. obliqua. The abrupt termination of all the Beeches at about lat. 35° , occurs where the equally sudden ehange in the climate of northern and southern Chili takes place. These trees, like all extra-tropical plants, require a certain degree of cold, and in pursuing their range towards the warmer parallels, they ascend the mountains. They are, however, even more dependent upon humidity and an equable elimate than on temperature; and being further impatient of vicissitudes and dryness, they will not pass beyond the influence of those S.W. winds which drench all parts of western South America, alpine and lowland, south of the parallel of 37° .



One of the few attractions of spring in Antarctic America, is the bursting of the leaf and flower buds of the deciduous-leaved Beech from their resinous gummy scales; when a delightfully fragrant odour pervades the woods. The unfolding of the plaited foliage was watched with great interest, for we had not witnessed for years any process so closely resembling that of an English spring. It recalled Linnæus' enthusiastic description of the first burst of the birch leaf in Lapland.

best in the island, for ground-frames of houses, planks for vessels, and beams. The piraguas are built chiefly of this wood. There are two sorts, one an evergreen, and the other a deciduous-leaved tree. It is evidently a Beech, and the same that grows in all parts of the Strait of Magalhaens; the smooth-leaved sort is *F. obliqua*, Mirb." Capt. King attaches the name of "Roble" to his specimen of *F. Dombeyi*.

PLATE CXXII. Fig. 1, male flower; fig. 2, involucre and female flower; fig. 3, transverse section of ditto, more advanced; fig. 4, ripe achænium; fig. 5, the same; fig. 6, involucre after the achænia have fallen away:— all magnified.

2. FAGUS obliqua, Mirb., Mém. Mus. Hist. Nat. vol. xiv. p. 465. t. 4. Hook. Bot. Journ. vol. ii. p. 153.

HAB. Strait of Magalhaens; Port Famine, Capt. King.

This I take to be the third kind of Beech alluded to by Capt. King as a native of Port Famine, in his collections, however, no specimen of the present species occurs. It is distinguishable from the former chiefly by the larger, narrower, rhomboidal, more acute leaves.

3. FAGUS Pumilio, Poepp. et Endlicher, Nov. Gen. et Sp. Plant. Per. et Chili, vol. ii. p. 68. t. 195. Hook. in Journ. Bot. vol. ii. p. 154. Calusparassus Pumilio (?), Homb. et Jacq. in Voy. au Pole Sud, Bot. Dicot. t. 8. 4.

HAB. Strait of Magalhaens; Port Famine (?), Capt. King.

I have alluded to this Beech (under *F. Antarctica*) as perhaps only a state of that plant, differing in the leaves being publication both surfaces and more closely and deeply serrated. The figure of Poeppig and Endlicher is excellent; that of MM. Hombron and Jacquinot, in the 'Voy. au Pole Sud', represents a narrower and smallerleaved, perhaps, alpine state; or more probably a different species, those authors having included it in their not yet described genus *Calusparassus*. Judging from their figures of other Antarctic *Fagi*, also called *Calusparassi*, the genus appears to include only those evergreen species of which the leaves are not plicate in vernation, which those of the *F. Pumilio* decidedly are, both in our specimens and those described and figured by Poeppig.

The latter author states this to be a short prostrate tree, eight and twelve feet long, with a mode of growth not unlike that of *Pinus Pumilio*. It marks (in Chili) the transition zone, from the erect trees, whose superior limit is indicated by the *F. alpina*, to the frigid region, where snow lies for eight months of the year, and where the shrubby *Compositæ*, and the Violets that grow in dense capitate tufts, and other handsome plants, abound.

I have marked the habitat assigned to Capt. King's specimen with a query, the label attached to it bearing "Cape Fairweather", where it is exceedingly improbable that any *Fagus* should exist.

3. FAGUS betuloides, Mirb., Mém. du Mus. vol. xiv. p. 465. t. 4. Hook. Journ. Bot. vol. ii. p. 153. F. dubia, Mirb. et Hook. l. c. F. Forsteri, Hook. l. c. p. 156. t. viii. Calusparassus Forsteri, Homb. et Jacq. in Voy. au Pole Sud, Bot. Monocot. Phan. t. 6. z. C. betuloides, Homb. et Jacq. l. c. Bot. Dicot. t. 7. f. r. Betula Antarctica, Forst. Comm. Goett. vol. ix. p. 45. Willd. Sp. Pl. vol. iv. p. 466. Banks et Sol. in Bibl. Banks. cum icone. (TAB. CXXIV.)

HAB. South Chili to Cape Horn, very abundant; Commerson, Banks and Solander, Forster, and all succeeding voyagers.

The synonyms above ennmerated certainly all belong to one species, the common Evergreen Beech of Fuegia, and I incline to add the *F. alpina*, Poepp. and Endlicher, as stated at p. 347.

PLATE CXXIV. Fig. 1, male flower; fig. 2, involucre with female flowers; fig. 3 and 4, female flowers removed from ditto; fig. 5 and 6, longitudinal sections of the same, showing the ovules; fig. 7, involucre, after the achænia have fallen away :----all magnified.

XLIX. CONIFERÆ, Juss.

1. THUJA, Tourn.

1. THUJA tetragona, Hook., in Lond. Journ. of Bot. vol. iii. p. 144. t. 4.

HAB. South Chili and Strait of Magalhaens; Port Famine, Capt. King.

This species has been described, on the authority of Mr. Bridges, as the true "Alerse" of Chili. Upon showing my specimens, however, to M. Claude Gay, the celebrated Chilian traveller, he assured me that the "Alerse" was a totally different plant, and not a Thuja at all; a statement the more probable, from Capt. King's description of the Alerse leaves, which, he says, resemble those of a Pine in colour, but are only half an inch long; though the difference may arise from the young and old states of this, as of other Coniferæ, often bearing leaves of a very different appearance. When enumerating the woods in use in the Island of Chiloe, Capt. King mentions in one place (p. 281) the "Alerse" and "Cypress", which are thus usually considered as different plants, and says that the "Cypress" is brought to that Island in "tablones" (or planks), seven or eight feet long, two inches thick, and nine or twelve inches wide, as is also the "Alerse"; but the latter, from the facility in which it splits, is brought in boards also. The same voyager observes (p. 183) that, though the "Cypress" is thought to be a different tree from the "Alerse" he considers it only a variety, the wood being white, whilst that of the "Alerse" is deep red. Naturalists who are aware how uncertain are the limits of the acknowledged species and varieties of European Conifera, will readily appreciate the difficulty that attends the determination of those of an opposite hemisphere, only known to us through insufficient specimens, vague reports, and incorrect information. Capt. King evidently believes the "Cypress" and "Alerse" to be identical, for he affirms that the former grows commonly in the Strait of Magalhaens, in all parts west of Cape Forward, but that there, from the poverty of the soil, the wood is of very stunted growth (p. 283); and this description tallies with the specimens of *Thuja tetragona* in his Herbarinm.

Thuja tetragona is apparently a rare Magellanic plant. Capt. King says it is found on the north shore of the Strait between Cape Forward and Port Gallant, but not to the eastward, except on the sides of Mount Tarn, where it only reaches the height of three or four feet (King's Voy. p. 131). The same author elsewhere states that the natives make their spears of its wood (p. 568). In Hermite Island where the *Thuja* does not exist Drimys Winteri is used for that purpose.

L. ORCHIDEÆ, Juss.

1. CHLORÆA, Lindl.

1. CHLORÆA Gaudichaudii, Brongn., in Duperrey Voy. Bot. p. 189. t. 44. A. Lindl. Gen. et Sp. Orchid. p. 405. Arethusa lutea, Gaud. in Ann. Sc. Nat. vol. v. p. 101, et in Freye. Voy. Bot. t. 133. D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 604. "Satyrion," Pernetty, Voy. vol. ii. p. 54. t. 8. f. 5.

HAB. Falkland Islands, Gaudichaud, D'Urville, Mr. Wright, J. D. H.

Not uncommon in moist pastures of the Falkland Islands, varying a good deal in size and in the breadth of its leaves. It differs from the *C. alpina*, Poepp., of South Chili, by the flowers being very much smaller, and the sepals, petals, and labellum differently formed. Both the figure and description of Brongniart, are very good.

2. CHLORÆA Magellanica, Hook. fil.; labello ovato-cordato obsolete trilobo breviter unguiculato marginibus inflexis glandulis grossis elongatis stipitatis cristato axi sub-lamellato, lobis lateralibus sub-laceris

intermedio producto apice subdilatato obtuso incrassato nudiusculo, sepalis lateralibus linearibus ultra medinm incrassatis apice obtusis carnosis marginibus inflexis, petalis ovatis obtusis sepalis $\frac{1}{3}$ brevioribus, spica triflora, scapo foliato.

HAB. Strait of Magalhaens; Elizabeth Island, C. Darwin, Esq.

 $Planta 1-1\frac{1}{2}$ pedalis. Folia basi longe vaginantia; lamina ovato-lanccolata, sub-recurva. Bracteæ ovato-lanceolatæ, acuminatæ, membranaceæ, concavæ. Flores crecti, majusculi, speciosi. Sepala oblongo-lanceolata, omnino nuda, superiore obtuso, lateralibus linearibus, supra medium siccitate nigrescentibus, omnia petalaque venosa et transversim venulosa. Labellum coriaceum, recurvum, marginibus involutis, petalis æquilongum. Columna petalis paulo brevior, arcuata.

A perfectly distinct and very handsome species, confined in its habitat to that eastern portion of the Straits of Magalhacus, where, as Mr. Darwin remarks, the Floras of Fuegia and Patagonia are blended.

2. ASARCA, Poepp.

1. ASARCA Commersonii, Lindley, Gen. et Sp. Orchid. p. 405, sub Chloræa.

HAB. Strait of Magalhaens, Commerson; Falkland Islands (Western Island?); Mr. Wright, Mr. Chartres.

Brongniart's description and figure are very characteristic of the Falkland Island specimens of this plant, which is quite distinct from the following. I have seen no Magellanic or Fuegian individuals, Capt. King's Port Famine *Chloræa* or *Asarca* being a totally different species. Mr. Wright and Mr. Chartres having gathered it in the Western of the two Falkland Islands, and no other collectors having met with it in the Eastern, I am inclined to consider this plant as one of the Fnegian species which has not spread to the eastern parts of the group, as is the case with the *Veronica elliptica*.

2. ASARCA odoratissima, Poepp., Nov. Gen. et Sp. Plant. Per. et Chil. vol. ii. p. 13. t. 118. Lindley, Gen. et Sp. Orchid. p. 407.

HAB. Falkland Islands (Western Island?), Mr. Wright.

Mr. Wright's specimen of a spike of this plant, preserved in spirits, entirely accords with the figure of Poeppig.

3. ASARCA (?) *Kingii*, Hook. fil.; labello breviter unguiculato oblongo obtuso indiviso integerrimo nudo membranaceo nervis mediis vix incrassatis, sepalis lateralibus lanceolatis acuminatis apicibus simplicibus. petalis oblongo-obovatis obtusis sepalis labelloque paulo brevioribus, spica 6–8-flora.

HAB. Strait of Magalhaens; woods of Port Famine, Capt. King.

Herba pedalis. Folia radicalia 6-uncialia, lanceolata, acuminata. Scapus foliatus. Spica 2–4 unc. longa. Bracteæ ovato-lanceolatæ, aeuminatæ, membranaceæ, concavæ. Flores pro genere parvi, flavi. Sepala vix $\frac{1}{2}$ unc. longa, membranacea, venosa, lanceolata, lateralibus basi angustioribus. Petala sepalis paulo breviora. Labellum sepalis æquilongum, omnino indivisum. Columna brevissima.

The short column has induced me to refer this very distinct plant to the genus *Asarca*, for in a dried state it is almost impossible to determine whether the petals are patent or conniving.

3. CODONORCHIS, Lindl.

1. CODONORCHIS Lessonii, Lindl., Gen. et Sp. Orchid. p. 411. C. Poeppigii, Lindl. l. e. Calopogon Lessonii, Brongn. in Duperrey Voy. Bot. p. 188. t. 37. f. 1. Pogonia tetraphylla, Poepp. et Endl. Nov. Gen. §c. vol.ii. p. 16. t. 122. Epipactis Lessonii, D'Urv. in Mém. Soc. Linn. Paris, vol.iv. p. 605. (TAB. CXXV.) HAB. Strait of Magalhaens, and throughout Fuegia, Commerson, Banks and Solander, and all succeeding voyagers. Falkland Islands, D'Urville, &c.

The leaves of this plant vary from two to four, three being the prevailing number. The flowers, also, are very much larger in some specimens than others, and dissimilar in colour and spotting. Poeppig's *Pogonia tetraphylla*, from South Chili, is decidedly only a state of *Codonorchis Lessonii*, the glands on the labellum affording no more constant character in this plant than in the beautiful *Chiloglottis* of Tasmania.

PLATE CXXV. Fig. 1, ovarium, column, and labellum; fig. 2, labellum; fig. 3, column; fig. 4, anther-case; fig. 5, pollen-masses :---all magnified.

LI. IRIDEÆ, Juss.

1. SISYRINCHIUM, Tourn.

1. SISYRINCHIUM *filifolium*, Gaud.; caule simplici tereti striato basi folioso, foliis radicalibus filiformibus scapum æquantibus brevioribusve, scapo ultra bracteas in spatham elongatam producto, fasciculis florum sessilibus rarius pedunculatis solitariis v. rarissime geminis bibracteatis 2–S-floris, perianthii segmentis subæqualibus albis purpureo-venosis. S. filifolium, *Gaud. in Ann. Sc. Nat.* vol. v. p. 101, et in Freye. Voy. Bot. p. 133. D'Urv. in Mém.Soc. Linn. Paris, vol.iv. p. 604. S. Gaudichaudii, Dietrich. Sp. Pl. vol.ii. p. 505. (TAB. CXXVI.)

HAB. Strait of Magalhaens; Cape Gregory, Capt. King; Falkland Islands, Gaudichaud, and all succeeding voyagers.

Herba elegans, 4-unc. ad bipedalem. Radir e fibris plurimis horizontalibus earnosis. Caulis basi reliquiis fibrosis foliorum emortuorum obteetus. Folia pauca, pleraque radicalia, filiformia, scapo breviora v. elongata. Scapus gracilis, teres. Spatha 2-5-unc. longa, basi vaginans, superne in folium filiformem desinens. Pedunculi floriferi plerumque solitarii, rarius bini, brevissimi v. raro elongati, apice braeteas duas lanccolatas æquilongas unciales gerentes. Pedicelli filiformes, exserti, stricti v. flexuosi. Flores magnitudine varii, Galanthi nivalis æquantes v. dimidio terve minores, late campanulati, albi. Perianthii segmenta subæqualia, obovata, apiculata, membranacea, venis sæpius flexuosis purpureis ornata. Stamina fere omnino libera, antheris versatilibus brevibus. Stytus apice incrassatus, trifidus, ramis divaricatis. Capsula membranacea-coriacea. Semina obovata, lævia; testa reticulata, brunnea.

One of the most abundant and elegant plants in the Falkland Islands, where the grassy plains are, in the spring month of November, almost whitened by the profusion of its pendulous snowy bells.

A very similar species, if not the same, seems to be common in Chili, from Valparaiso to Concepcion; but its flowers are smaller than in the majority of the Falkland Island specimens.

PLATE CXXVI. Fig. 1, segment of the perianth; fig. 2, ovarium, stamens, style, and stigmata; fig. 3, transverse section of ovarium; fig. 4, ovule; fig. 5, ripe fruit; fig. 6, transverse section of the same; fig. 7, seed; fig. 8, the same, cut longitudinally:—all magnified.

2. SISYRINCHIUM laxum, Link., in Hook. in Bot. Mag. t. 2312.

Var. *major*; caule bifido foliisque latioribus, spatha bracteisque apices versus scaberulis, perianthii segmentis latioribus.

Var. *minor*; caule simplici foliisque angustioribus, spatha bracteisque glaberrimis, perianthii segmentis angustioribus.

HAB. Var. major, Chonos Archipelago, C. Darwin, Esq. Var. B, Strait of Magalhaens; Port Famine. Capt. King; Cape Negro, C. Darwin, Esq.

A species which has been erroneously included by Sprengel in the terete stemmed group, and even considered by Lindley and Dietrich to be synonymous with *S. iridifolium*, Kunth, (*Marica iridifolia*, Bot. Reg. t. 646). Such may be the case, but I have seen no specimens decidedly connecting these two species, and therefore hesitate before adopting a conclusion which would give this plant a geographical range from the equator to the Strait of Magalhaens. Still, the var. *major* is so decidedly scaberulous, so much larger, and so much more resembling the *S. iridifolium* than the var. *minor*, that there is nothing improbable in the supposition that both are varieties of one tropical species. *S. laxum* is also a native of Valparaiso.

3. SISYRINCHIUM (?) sp.

HAB. Strait of Magalhaens; Capt. King.

A curious plant, unfortunately too imperfect for description, but with very much the habit, foliage, and fruit of a *Sisyrinchium*. Root consisting of clongated fleshy fibres, 3-4 inches long. Rhizoma very short, fibrous, giving off at its apex a terete scape and short leafy stem. *Leaves* about three, with scariose sheaths at the base, filiform, terete, six or seven inches long. *Scape* shorter than the leaves, furnished at the middle with two leaf-like opposite bracteæ sheathing at the base. *Peduncle* solitary, one-flowered, shorter than the bracteæ, erect. *Fruit* immature, globose, trigonous (?), the size of a small pea, three-locular (?), each cell containing several seeds on parietal (?) placentæ.

The above diagnosis may serve to distinguish this curious plant, which differs chiefly from *Sisyrinehium* in the scape not springing from between the uppermost leaves, but from the base of the outer one, and in there being no spatha to any of the specimens; though the scape in one instance bears the scar of a fallen leaf, half-way between the insertion of the bracteæ and the rhizoma.

2. SYMPHYOSTEMON, Miers.

1. SYMPHYOSTEMON narcissoides, Miers, in Linn. Soc. Trans. v. xix. p. 97. Sisyrinchium narcissoides, Cav., Diss. vol. vi. p. 347. t. 191. f. 3. S. odoratissimum, Lindl. Bot. Reg. t. 1283. Galaxia narcissoides, Willd. Sp. Pl. vol. iii. p. 583. Gladiolus biflorus, Thunberg, Diss. Glad. n. 5.

HAB. Strait of Magalhaens, Commerson; Port Famine, Capt. King; Elizabeth Island, C. Darwin, Esq.

I quite agree with Mr. Miers in removing this plant from *Sisyrinchium*. Thunberg's habitat of the Strait of Magalhaens, from whence he originally described this plant as *Gladiolus biflorus*, has been replaced by that of the Cape in most succeeding authors, except Vahl (En. Plant. vol. ii. p. 97), and Willdenow (Sp. Pl. vol. i. p. 209).

3. TAPEINIA, Juss.

Perigonium corollinum, superum, hexaphyllo-partitum; laciniis basi connatis, subcarnosis, patentibus, apiculatis, 3 exterioribus majoribus, Stamina.3, imo perigonii inserta; filamentis in tubum trigonum connatis, supra medium liberis; antheris extrorsis, lineari-ovatis, basi profunde emarginatis. Ocarium lineari-obovatum, 3-loculare. Ovula plurima, basi anguli centralis loculi affixa. Stylus validus, supra medium in stigmata 3 erecta subulata apice dilatata papillosa fissus. Capsula coriacea, globosa, triloba, trilocularis, apice loculicido-trivalvis. Semina plurima, obovata, teretia; testa subcoriacea, grosse cellulosa; rhaphe indistincta; chalaza atra; embryo parvus, elongatoobconicus, basi albuminis duri immersus.—Tapeinia, Juss. Gen. p. 59, e schedis Commersonii.

I. TAPEINIA Magellanica, Juss., l. c. Witsenia pumila, Fahl, Enum. vol. ii. p. 48. Ræm. et Sch. Syst. Veg. vol. i. p. 371. Spreng. Syst. Veg. vol. i. p. 147. Dietrich. Sp. Pl. vol. ii. p. 559. W. Magel-4 K lanica, Pers. Synops. vol. i. p. 42. Ixia pumila, Forst. Comm. Goett. vol. ix. p. 20. t. 8. I. Magellanica, Lam. Ill. vol. i. p. 109. Moræa Magellanica, Willd. Sp. Pl. vol. i. p. 241. Galaxia obscura, Cav. Diss. vol. vi. p. 341. t. 189. f. 4. (Sisyrinchium pumilum, TAB. CXXIX.)

HAB. Strait of Magalhaens and throughout Fuegia, on the mountains, Commerson, Banks and Solander, and all succeeding voyagers.

The accompanying plate and analysis of this curious little plant represent all its characters, and especially those which have induced me to retain the genus which the illustrious Jussieu formed, but which has not been adopted by any succeeding author. It is to be distinguished from *Sisyrinchium* by its very remarkable habit, coriaceous perianth, and, more especially, by the capsule dehiscing at the apex, and the ovules and seeds occupying only the lower half of each placentiferous dissepiment. To the southward of the Strait of Magalhaens, where *Sisyrinchia* do not extend, this little plant represents that genus, and is also the analogue of the *Libertiæ* of New Zealand.

The curions and beautiful distichous arrangement of the foliage, is characteristic of this and of some other especially alpine Antarctic plants, belonging to several natural orders, amongst the majority of the species of which such a foliation is foreign or very rare. Thus, in *Cyperaceæ* it is seen in *Oreobolus pectinatus* (pt. 1. t. 49); amongst *Restiaceæ*, in *Gaimardia pallida* (p. 86); amongst *Alismaceæ*, in *Tetroncium Magellanicum* (t. 128); and amongst *Junceæ*, in the Peruvian *Distichya muscoides*, Nees and Meyen (Nov. Act. Acad. Cæs. vol. xix. Suppl. p. 77), which is probably the *Goudotia Tolimensis*, Decaisne (Ann. Sc. Nat. ser. 3. vol. iv. p. 83. t. 4). This tendency to assume a certain habit, which these otherwise wholly dissimilar plants present, is perfectly analogous to what occurs even more conspienously in the vegetation of the Cape of Good Hope and Australia; and one of the most singular phenomena of the vegetable kingdom.

PLATE CXXIX. Fig. 1, bracteæ and flower; fig. 2, expanded flower; fig. 3, stamens, styles and stigmata; fig. 4, ripe capsule; fig. 5, the same burst open; fig. 6, one value of the same, showing the insertion of the seeds; fig. 7, a seed removed; fig. 8, vertical section of the same; fig. 9, embryo (the figures 8 and 9 are inadvertently transposed):—magnified.

LII. SMILACEÆ, Br.

1. CALLIXENE, Comm.

1. CALLIXENE marginata, Commerson, ex Juss. Gen. n. 41. Lam. Illust. Gen. t. 248. Gaud. in Ann. Sc. Nat. vol. v. p. 101. t.2. f.2, et in Freyc. Voy. Bot. p. 133. D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 604. Enargea marginata, Banks et Sol. MSS. in Bibl. Banks. cum icone, et in Gærtner de Fruct. vol. i. p. 283. t. 59. f. 3.

HAB. Strait of Magalhaens and throughout Fuegia, Commerson, Banks and Solander, Sc. Falkland Islands, most abundant, Gaudichaud, and all succeeding voyagers.

A very elegant little plant, remarkable, especially in the Falkland Islands, for its very sweet-scented flowers.

The extrorse anthers of this genus have been hitherto overlooked, from the versatile nature of their attachment. The embryo, described as amphitropal, at first is nearly atropal; but apparently during the maturation of the ovarium the seed becomes partially inverted, so as to be placed at right angles with the funiculus, and the embryo is consequently heterotropal.

Callizene is an Antarctic-American, and New Zealand genus. From the latter country Mr. Colenso has sent the C. pareiflora, Hook. fil. (Hook. Ic. Plant. t. 632), which grows at the foot of large Beech trees, lying prostrate
against their trunks in the mountain forests, as the *C. marginata* does at Cape Horn. Their Australian representative is the *Drymophila cyanocarpa*, Br., a subalpine Tasmanian plant, very similar to them in habit.

2. CALLIXENE polyphylla, Hook. Ic. Plant. t. 674.

HAB. Cape Tres Montes, C. Darwin, Esq.

The *C. marginata* does not attain a lower latitude in South America than the Strait of Magalhaens, but is replaced in South Chili by the present species, which is much handsomer; this, again, is represented in Peru by the genus *Luzuriaga* of Ruiz and Pavon.

2. PHILESIA, Comm.

Flores hermaphroditi. Perigonium corollinum, campanulatum, sexpartitum, laciniæ exteriores interioribus multoties breviores. Stamina 6, imo perigonii inserta; filamenta filiformia, infra medium in tubum connata; antheræ inclusæ, lineares, extrorsæ. Ovarium parvum, nniloeulare. Ovula plurima, sub-biserialia, orthotropa, funiculis brevibus, plaeentis parietalibus elongatis adnexa. Stylus elongatus, simplex. Stigma exsertum, capitatum, plumosum, obscure 3-lobum. Bacca uniloeularis, polysperma. Semina pulpo glutinoso nidulantia, ascendentia, ovoidea, rugosa; testa tenuis, flavida; albumen corneum; embryo cavitate axili albuminis lente arcuatus, extremitate cotyledonari hilo oppositus.—Suffrutex Chilensis suberectus. Rami teretes, stricti v. flexuosi. Folia alterna, coriacea. Pedunculi ramis terminales. Flores magni, speciosi, basi bracteati. Philesia, Commerson, ex Juss. Gen. p. 41.

1. PHILESIA buxifolia, Lam., Illust. Gen. t. 248. Poiret, Encycl. vol. v. p. 269. Ram. et Sch. vol. vii. p. 314. Lindl. Veg. Kingd. p. 217.

HAB. Strait of Magalhaens, Commerson; Port Famine, Capt. King; Good Success Bay, Banks and Solander.

Except by the parietal placentation, the genera *Philesia* and *Lapageria* (themselves very closely allied), differ in no important points from *Callixene* and *Luzuriaga*, and since placentation does not afford characters of the importance amongst Monocotyledonous that it does in Dicotyledonous Orders, I see no objection whatever to arranging these two genera under *Smilaceæ* proper and next to *Callixene*.

In Asteliaceæ, as I have mentioned elsewhere, the placentæ are axile, parietal or pendulous; in Junceæ, parietal or basal; in Amaryllideæ, axile or parietal; in Liliaceæ, the same; and other orders equally display a very considerable amount of variation in the consolidation of the carpels, and consequent disposition of the placentæ, nnaecompanied, however, with any other characters of more than generic value.

In all other respects, *Philesia* is even generically very nearly related indeed to *Callixene*, through *Lucuriaga*, which has the three inner segments of the perianth still larger in proportion than in *Callixene*; and on the other hand, through *Lapageria*, in which they are all equal in size. The habit, texture, distiehous insertion of the leaves, which are all on the same plane with the ramuli; the texture, nervation, margination, and even form of the leaves, which are glaucous beneath, are alike in *Callixene* and *Philesia*; so are the terminal, large, solitary, bracteate flowers, the texture of the perianth, extrorse anthers, baccate fruit, the numerous ovules in two series on three rows of placentæ, the many ovoid seeds, delieate testa, dense albumen, and axile embryo which is of similar form in the two. The only difference in the ovules is, that those of the *Callixene* are heterotropal, those of *Philesia* nearly straight or atropal, charaeters rather indicating close affinity than the contrary.

With regard to the genus Lapageria, R. and P., it is so elosely allied to *Philesia* that I doubt its validity, the ehief differences being the nearly equally divided perianth of Lapageria, its more distinctly three-lobed stigma, oblong

FLORA ANTARCTICA.

berry, twining branches, and differently nerved leaves, in all which respects it is more evidently a genus of *Smilaceæ*, than either *Callixene* or *Philesia*. There is no reason for supposing Dombey's *Capia* to be other than *Lapageria rosea*.

It appears to me to be through these Antarctic and extra-tropical American genera, together with the *Callixene* of New Zealand and *Drymophila* of Tasmania, that the *Smilaceæ*, Lindl., are inseparably connected with the Tribe *Asparageæ*, Lindl., of *Liliaceæ*; groups which Dr. Lindley has placed in separate natural classes, on the ground chiefly of anatomical differences in their stems: and it further appears that all modifications of a stem typical of Endogens and one equally characteristic of Dictyogens may be traced amongst these plants.

My own observations on the wood of *Philesia* do not exactly lead to the conclusions that the learned author of the 'Vegetable Kingdom' has formed; what appears to be bark is at no period separable from the subjacent wood, and the pith is of undefined form. There is a resemblance between the bark of *Philesia* and that of an exogenous stem, but it is apparent and not real: the stem consists of one mass of cellular tissue, through which bundles of vascular tissue descend, between the axis and the cuticle; abundantly towards the latter, where they all coalesce, though always at a little distance within the circumference; more sparingly towards the axis, where a space is often left wholly unoccupied with woody fibres. A transverse section of such a stem thus presents, 1st, a cuticle; 2nd, a zone of cellular tissue, often formed of thick walled cells; 3rd, a zone of wood, dense and defined externally, gradually laxer towards the axis and separating into bundles which irregularly surround a central column of pith. The only difference, in short, between this and any other Endogenous stem, consists in the first-formed or outer bundles being disposed more symmetrically, and being combined into one zone.

If a branch of *Luzuriaga radicans* be examined, the same peculiarity will be perceived, with only this difference, that the zone of wood is narrower and the pith broader. In *Callivene polyphylla*, the woody zone, though still continuous, is narrower still. In *C. parviflora* both its edges (both inner and outer circumference) are elearly defined; and in *C. marginata* it is sometimes interrupted.

The Callizene marginata thus shows this disposition of the outer vascular bundles to unite in the lowest degree of these South American Smilaceæ, but in Lapageria the same tendency will be found in its highest, for the stem of that plant is almost wholly composed of woody matter, concentrated externally into a well-defined zone, rather looser towards the centre, and enclosing large tracheæ with very little cellular tissue intermixed. Externally to the wood is a very narrow layer of condensed parenchyma. In the first year's twig of this plant, the cellular tissue is proportionably abundant, with separate vascular bundles scattered through it, but is absorbed or obliterated afterwards. Nor is it in the genera of South America alone that these woody bundles are thus arranged, it is so in the Geitonoplesium (Luzuriaga cymosa, Br.) of New Holland, and in Drymophila, Br.; and even nearer home in Convallaria and probably in many Convallarieæ. To the last mentioned group the above named genera most assuredly belong; whether the venation be parallel as in Callizene, parallel and retose between the costæ as in Lapageria, or wholly retose as that of Philesia appears to be, from the two lateral of the three parallel costæ forming the thickened margin of the leaf.

On the other hand, if we turn to the *Smilaceæ* proper, as limited by Dr. Lindley, even they display no more deviation from the common Endogenous structure than do the *Convallarieæ*. A young shoot of *Rhipogonum* shows the same disposition of the woody and cellular tissue as *Callixene polyphylla*, with rather a broader zone of cellular tissue surrounding the wood; but in an older stem of the same, the wood so predominates over the parenchyma, that the zone of cellular tissue is only distinguished with difficulty. In the *Smilax excelsa*, L., of Europe, the woody zone of the young branch is neither so continuous nor regular, but it becomes so in the older state of the plant. *Tamus communis* presents the same arrangement. In the young stem of *Testudinaria elephantipes* I do not find the medullary plates described by Dr. Lindley; there appears to me to be a broad and perfectly continuous zone of wood, sending six or eight prolongations towards the axis, where there are further a few irregularly disposed bundles. I shall conclude this long digression by instancing the genus *Juncus* as of the furthest removed from

Dictyogens in every point of view, except that it possesses an equally continuous and defined zone of woody tissue, within the cuticle, separated from the latter by a zone of parenchyma, and enclosing a mass of pure pith.

The *Philesia buxifolia* is among the handsomest plants of the Antarctic American Flora; it occurs along the coast from the Strait of Magalhaens to Valdivia; to the northward of which, between Valdivia and Concepcion, it is replaced by the *Lapageria rosea*.

LIII. ASTELIEÆ, Brongniart.

1. ASTELIA, Banks et Sol.

1. ASTELIA pumila, Brown, Prodr. p. 291. Gaud. in Ann. Sc. Nat. vol. v. p. 100. et in Freyc. Voy. Bot. p. 132. D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 603. Fl. Antarct. vol. i. p. 76. Melanthium pumilum, Forst. Comm. Goett. vol. ix. p. 30. t. 6. Banks et Sol. MS. in Mus. Banks. cum icone. Funkia Magellanica, Willd. Mag. Naturf. Fr. vol. ii. p. 19. (TAB. CXXVII).

HAB. South Chili, from the Chonos Archipelago to Cape Horn, very abundant on the hills and in exposed places, *Commerson*, and all future voyagers ; Falkland Islands, *Gaudichaud*, Sc.

Under the description of A. linearis, in the first part of this volume, I mentioned that the placentation varies in the different species of this genus. In the majority, the ovules are numerous and arranged in two lines upon parietal placente; in one the seeds are numerous and pendent from the summit of a one-celled berry, whose dissepiments have probably been absorbed; a third form presents a three-celled ovarium, with several ovules pendulous from the summit of each cell; a fourth has a three- to six-celled subcapsular fruit, with a few pendulous seeds in each cell; while the present plant offers a fifth modification, for its placentation is decidedly axile, and the ovules are arranged in two rows along the inner angle of each of the three cells. This arises from the perfect consolidation of the carpels in a young state, when the edges of each carpellary leaf are so inflected as to meet in the axis of the pistil, where a triangular longitudinal cavity is often left (see fig. 5 of Plate CXXVII.). At an early period the cavity of each capsule is not apparent, the ovules being imbedded in a cellular mass, which in this species retires from between and around the ripering seeds, leaving a distinct cavity as the fruit advances to maturity, but in some others remains, partly attached to the placentæ and seeds, as a mucilaginous or gummy mass. At no time is the fruit of this plant truly even sub-capsular, its walls are always fleshy, and no trace of dehiscence can be seen along the furrows of each carpel, from which the seeds escape by the decay of the pericarp.

I have followed M. Brongniart in placing this genus by itself in a natural group, whose nearest affinities I have indicated in the first part of this work.

The Astelia pumila is a most abundant Fuegian and Falkland Island plant, forming, with the Caltha appendiculata especially, a large proportion of the peat in those countries. Its flowers are inconspicuous, and have a faintly sweet smell.

PLATE CXXVII. Fig. 1, three-flowered peduncle, bract and flower; fig. 2, flower removed; fig. 3, pollen; fig. 4, ovarium; fig. 5, transverse section of the same; fig. 6 and 7, ovules; fig. 8, ripe fruit; fig. 9, transverse section of ditto; fig. 10, ripe seed; fig. 11, the same with the outer osseous integument removed; fig. 12, the same, cut longitudinally; fig. 13, embryo:—all magnified.

LIV. JUNCEÆ, DC.

1. ROSTKOVIA, Desv.

1. ROSTKOVIA grandiflora, Hook. fil.; in Fl. Antarct. vol. i. p. 82. Marsippospermum calyculatum, Desv. Bot. Journ. vol. i. p. 330. M. grandiflorum, Hook. Ic. Plant. t. 533. Juncus grandiflorus, Linn. fil.

4 L

Suppl. p. 209. Forst. Comm. Goett. vol. ix. p. 27. t. 3. Gaud. in Ann. Sc. Nat. vol. v. p. 100, et in Freye. Voy. Bot. p. 132. D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 603.

HAB. Strait of Magalhaens and throughout Fuegia, Commerson, Banks and Solander, Sc. Falkland Islands, very abundant, Gaudichaud, and all succeeding voyagers.

The miserable natives of Fuegia weave the stems of this rush into baskets, and in doing so seem to exhaust their cunning, for such baskets appeared to us to be the only article they possessed, exhibiting any attempt at such handy-craft as demands the slightest ingenuity, except, perhaps, the moveable heads of their sealing spears.

2. ROSTKOVIA Magellanica, Hook. fil. l. c.

HAB. Strait of Magalhaens, Commerson; Hermite Island, Cape Horn, J. D. H.; Falkland Islands, very abundant, Gaudichaud, Sc.

I am not aware of this species having been gathered in Fuegia since Commerson's time, except by myself; and though abundant in Hermite Island, it is probably scarce and alpine to the north of that locality, as it is also in Campbell's Island.

2. JUNCUS, L.

1. JUNCUS scheuchzerioides, Gand.; Fl. Antarct. p. 79.

HAB. Strait of Magalhaeus; Port Famine, Capt. King; Hermite Island, Cape Horn, J. D. H.; Falkland Islands, very abundant, Gaudichaud, &c.; Kerguelen's Land, J. D. H.

Decidedly the most Antarctic *Juncus*, and exceedingly abundant at Cape Horn, the Falkland Islands, and Kerguelen's Land, where no other species of the genus exists. It is also a native of Campbell's Island and Lord Auckland's group.

2. JUNCUS planifolius, Brown, Prodr. p. 259. E. Meyer, Junci, n. 36, ct in Linnæa, vol. iii. p. 370. La Harpe, in Mém. Soe. Nat. Hist. Paris, vol. ii. p. 55. Kunth, En. Plant. vol. iii. p. 344.

HAB. Chonos Archipelago, C. Darwin, Esq.

These, and other specimens gathered at Valdivia by Mr. Bridges, are the only extra-Australian individuals of this species that I have seen. Meyer remarks (Herb. Hook.), that there is no specific difference between the specimens of the New and Old World.

3. JUNCUS graminifolius, E. Meyer, in Rel. Hank. vol. ii. p. 144. Cephaloxys graminifolia, Nees et Meyer, in Nov. Act. Acad. Cas. vol. xix. Suppl. p. 128. J. rivularis, Poeppig, fid. Meyer in Herb. Hook.

HAB. Chonos Archipelago, C. Darwin, Esq.

The present species, like the former, can scarcely be considered truly Antarctic, merely entering the northern limits which I have assigned to the Fnegian Flora. It ranges on the coast from Valparaiso to the latitude of Chonos Archipelago and is also found on the Cordillera of Peru.

Meyer (Hook. Herb.) remarks that this hardly belongs to the genus *Cephaloxys*, on account of the structure of its capsule.

3. LUZULA, DC.

1. LUZULA Alopecurus, Desv. Bot. Journ. vol. i. p. 159. E. Meyer, in Reliq. Hank. vol. ii. p. 145. Syn. Luzul. n. 5. La Harpe, in Mém. Soc. Hist. Nat. Paris, vol. ii. p. 177.

HAB. Strait of Magalhaens, Commerson, Capt. King; Falkland Islands, very abundant, Gaudichaud, Sc.

I have seen but an imperfect specimen of the *L. Peruviana*, Desv., to which the present is manifestly very closely related. E. Meyer (Herb. Hook.) observes, that though so much alike in the young state, when older they are very distinct species. The present is the most Antarctic of the genus, except the following, and is the South American representative of the *L. crinita* (Tab. XLVIII.) of Lord Auekland's group.

2. LUZULA sp.?

HAB. Hermite Island, Cape Horn, on the tops of the mountains, alt. 1,600 feet, J. D. H.

My specimens are only sufficient to prove this plant to be a Luzula; they are searcely two inches high, with a slender stem, and nodding small paniele; the whole somewhat resembling the L arcuata of Aretie Europe, whose Antaretie representative it probably is.

LV. ALISMACEÆ, Br.

1. TETRONCIUM, Willd.

Flores dioiei. FL. MAS. Perigonium obliquum, tetraphyllum, coloratum, foliolis coneavis inæqualibus, late ovatis, superioribus altius insertis, supremo majore. Stamina 4, foliorum perigonii basi inserta; filamenta brevissima; antheræ extrorsæ, late didymæ, basi fixæ. Ovarii rudimentum nullum. FL. FEM. Perigonium ut in mase., sed foliolis angustioribus. Stamina 0. Carpella 4, subulata, basi in ovarinm incomplete 4-loculare coalita, supra medium libera; styli subulati, divergentes, imo apiec inconspieue stigmatiferi; ovula quovis loculo solitaria, erecta, anatropa, foramine late aperto. Fructus indehiscens, 4-locularis, monospermus. Semen crectum, lineari-oblongum, compressum; testa tennissima; albumen farinaceum; embryo axilis, trigonus, longitudine albuminis, extremitate radiculari attenuata.—Herba Fuegiana et Falklandiea cæspitosa, perennis. Caulis basi radicans, squamis nitidis tectus, divisus. Folia plana, disticha, equitantia, lineari-ensiformia. Seapns terminalis, erectus. Flores spicati. Perigonium flaxescens, rufo-fusco maculatum. Stamina antheris magnis. Fructus deflexus, abortu monospermus, 4-cornutus. Tetroncium, Willd.

1. TETRONCIUM Magellanicum, Willd., in Berl. Mag. vol. ii. p. 17. Hook. Ic. Plant. t. 534. Kunth. En. Plant. vol. iii. p. 142. Triglochin reflexum, Vahl, ined. (ftd. Willd.). T. Magellanicum, Vahl, in Herb. Mus. Paris. Cathanthes, Rich. in Mém. Mus. vol. i. p. 365.

HAB. Strait of Magalhaens, Commerson; Port Famine, Capt. King. Good Success Bay, Banks and Solander, Forster; Hermite Island, Cape Horn, J. D. II.; Falkland Island, Mr. Wright, J. D. II.

The arrangement of all parts of the flower are quaternary in the specimens of this eurious plant that I have examined; in which respect it differs from the majority of, and in the albuminous seeds from all the order, *Alismaceæ*; without, however, shewing any further affinity with the *Naiadaceæ*, in which order Dr. Lindley has placed it.

The habit of *Tetroncium* is precisely that of *Narthecium*, but in most other points its alliance to *Triglochin* is evident, particularly in the spicate inflorescence, concave segments of the perianth, which are obliquely placed, the upper being larger and inserted above the rest; in the extrorse, nearly sessile anthers; the solitary, basal, anatropal ovules; and the erect seed, which, being albuminous, indicates an affinity with *Junceæ*.

PLATE CXXVIII. Fig. 1, male flower; fig. 2, segment of perianth and stamen; fig. 3, female flower; fig. 4, earpel eut open; fig. 5, ovnle; fig. 6, the same, with the primine partly removed; fig. 7, ripe fruit; fig. 8, transverse section of the same; fig. 9, the same, longitudinally divided; fig. 10, seed; fig. 11, embryo:—all magnified.

FLORA ANTARCTICA.

2. TRIGLOCHIN, Linn.

1. TRIGLOCHIN Monte-Vidense, Spreng., Syst. Veg. vol. ii. p. 145. Roem. et Sch. Syst. vol. vii. p. 1586. Kunth, En. Plant. vol. iii. p. 144. T. capense, Thunb. Prodr. p. 67. T. maritimum, Drege, in Herb. Hook. T. striatum, Cham. et Schlecht. fid. Kunth, l. c.

HAB. Cape Tres Montes, C. Darwin, Esq.

Variat magnitudine, scapoque foliis nunc longiore nunc multoties breviore.

Probably a very widely diffused, and certainly in size a variable plant, common to both coasts of extra-tropical South America, and to the Cape of Good Hope. To this may also belong the *T. Chilense*, of Meyer, of which a wholly insufficient character is given in a foot-note to that traveller's journey (Reise un die Erde. vol. i. p. 354). Its nearest ally is the *T. decipiens*, Br., of Australia, of which *T. filifolium*, Sieb. (inaccurately described as wanting the abortive carpels), is a synonym; indeed, the Australian differs from the South American plant only in the larger fruit, so far as my only specimen enables me to judge.

LVI. RESTIACEÆ, Br.

1. GAIMARDIA, Gaud.

1. GAIMARDIA australis, Gaud., in Ann. Sc. Nat. vol. v. p. 100, et in Freye. Voy. Bot. p. 419. t. 3. Kunth, En. Plant. vol. iii. p. 491.

HAB. Fuegia; Hermite Island, Cape Horn, J. D. H.; Falkland Islands, very abundant, Gaudichaud, D'Urville, J. D. H.

A particularly abundant plant on the hills of the Falkland Islands, forming, in boggy places, hard, extensive green patches, often several yards across, and contributing materially to the formation of peat-bog. It has representatives on Lord Auekland's Group and probably likewise in Tasmania.

1

LVII. CYPERACEÆ, DC.

1. OREOBOLUS, Br.

1. OREOBOLUS obtusangulus, Gaud., in Ann. Sc. Nat. vol. v. p. 99. t. 2. f. 1, et in Freye. Voy. Bot. p. 417. Kunth, En. Plant. vol. ii. p. 367.

HAB. Fuegia; Hermite Island, Cape Horn, J. D. H.; Falkland Islands, abundant, Gaudichaud, D'Urville, J. D. H.

It is difficult to suppose that a plant, so abundant in the Falkland Islands, should be rare on the mountains of the adjacent continent, where, however, it has only been gathered near Cape Horn, unless a species collected by M. Gondot full 4,000 miles further north, on the peak of Tolima in Colombia, should prove to be the same plant, as, judging from a barren specimen, it very likely may.

2. ELEOCHARIS, Br.

1. ELEOCHARIS palustris, Br., Prodr. p. 244. Engl. Bot. t. 131. Scirpus melanostachys, D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 603. Fimbristylis melanostachys, Brong. in Duperrey, Voy. Bot. p. 181.

HAB. Falkland Islands, D'Urville, J. D. H.

Evidently the European *E. palustris*, which is also a native of Patagonia, and very widely diffused throughout the temperate regions of both the northern and southern hemispheres. Hypogynous setæ are generally present, though those of my specimens vary in size; Kunth says, "setæ plane abortientes in *Scirpo melanostachyo*," and D'Urville and Brongniart have, from their occasional absence, included this species in *Fimbristylis*.

3. ISOLEPIS, Br.

1. ISOLEPIS pygmæu, Kunth, En. Plant. vol. ii. p. 191.

Var. brevis. Isolepis brevis, Brong. in Duperrey, Voy. Bot. p. 180. I. Magellanica, Gaud. in Duperrey, Voy. Bot. p. 414. I. Meyeniana, Nees, in Nov. Act. Acad. Cas. vol. xix. Suppl. p. 87.

Var. clongata. I. pygmæa, var. B, Kunth, l. c. I. trigona, Kunze, in Poeppig, Coll. n. 1. p. 27 (?).

HAB. Var. brevis, Falkland Islands, D'Urville, J. D. H. Var. elongatu, Cape Tres Montes, C. Darwin, Esq.

The variations in the size, form, and markings on the surface of the achænia of the otherwise almost identical forms of *Isolepis* seem really endless. Falkland Island specimens are short, with small spikes, and small fuscous achænia, which are broader than long and punctulate, but the puncta not in parallel lincs. Mr. Darwin's plant is much longer, and has rather longer spikes, with elliptical ovate larger achænia, which are longer than broad and similarly punctulate, its culms are often ten inches long. The *I. lepida*, Nees (in Linnæa, vol. iv. p. 291), judging by Cuming's Valparaiso specimens (in Herb. Hook.), resembles the plant of Mr. Darwin, its achænia are precisely similar to those of the Falkland Island variety in form, colour and surface, but scarcely half as large.

The Cape of Good Hope *I. pygmæa*, so called by Kunth, has the achænium of *I. lepida*, but pale coloured and smaller still; while the Auckland Island *I. Aucklandica* (p. 88, t. L) has larger fruit than any.

All of the above differ from the European *I. Savii*, Seb. and Manr., in the achænia not being so deeply punctate or striate. In size and form the pericarp of *I. Savii* resembles that of the Falkland Island plant.

4. CHÆTOSPORA, Br.

1. CH.ETOSPORA Antarctica, Hook. fil.; culmis dense cæspitosis teretibus basi foliosis, foliis culmum vix æquantibus anguste lineari-elongatis rigidis semiteretibus super canaliculatis glaberrimis, spiculis sub 6 in paniculam brevem coarctatam involucro 5-phyllo breviorcm aggregatis 1-floris, squamis distichis carinatis imberbibus, setis hypogynis 6 capillaribus nucem superantibus. (TAB CXLVII.)

HAB. Cape Tres Montes; Patch Cove, alt. 2,000 feet, C. Darwin, Esq.

Radix e fibris crassis descendentibus. Rhizoma breve, inclinatum. Culmi dense cæspitosi, rigidi, erecti. Folia 6-pollicaria, basi in vaginam castaneam chartaceam 1 unc. longam dilatata; lamina vix $\frac{1}{2}$ lin. lata, apice acuminata. Panicula sub 1 unc. longa, coarctata, involucro basi vaginante $\frac{1}{2}$ brevior. Spiculæ erectæ, pedicellatæ, inferiores involucratæ, $\frac{1}{2}$ unc. longæ, lineari-oblongæ, uniflores. Squamæ sub 5, pallide flavo-fuseæ, nitidæ, linearioblongæ, acuminatæ, dorso carinatæ, inferiores supremaque vacuæ. Stamina 3. Setæ hypogynæ 6, squamis breviores, graciles, scaberulæ. Nux elliptico-oblonga, angulis costatis, glaberrina, polita, pallide fusea. Stylus gracilis, elongatus, apice stigmataque filiformia exserta.

PLATE CXLVII. Fig. 1, spikelet; fig. 2, flower with the anthers fallen away :-- both magnified.

2. CHÆTOSPORA *laxa*, Hook. fil.; culmis dense cæspitosis teretibus basi foliosis, foliis culmo brevioribus anguste lineari-elongatis rigidis semiteretibus super canaliculatis glaberrimis, spiculis plurimis in paniculam

laxam subeffusam involucratam involucris breviorem dispositis 2-floris, squamis distichis carinatis exterioribus dorso scaberulis, setis hypogynis 4–6 rigidis scabridis nuce $\frac{1}{2}$ longioribus. (TAB. CXLVI.)

HAB. South Chili; Cape Tres Montes, C. Darwin, Esq.

Culmi pedales. Folia ut in priore sed duplo longiora. Panicula 2-3 une. longa, paree ramosa. Involucri foliola 2 exteris longiora, panieulam superantia. Spiculæ $\frac{1}{4}$ unc. longæ, ovato-oblongæ, compressæ, bifloræ, inferiores longius et graciliter pedieellatæ. Squamæ sub 6, atro-eastaneæ, coneavæ, ovatæ, aeutæ v. sub-acuminatæ, nitidæ, inferiores vacuæ. Stamina 3. Setæ hypogynæ 4-6, rigidæ, seabridæ. Nux breviter stipitata, late elliptiea, angulis costatis, stylo elongato stigmatibus 3 capillaribus exsertis terminato.

One of these two species of *Chætospora* may be considered the Antarctic representative of the *Schænus nigricans* of Europe. Neither of them appears to inhabit a high south latitude, though the *C. Antarctica*, ascending to an elevation of 2,000 feet in South Chili, might have been expected to grow at the level of the sea in Fuegia.

PLATE CXLVI. Fig. 1, spikelet; fig. 2, the same with the lower scales removed; fig. 3, achænium :---all magnified.

5. CARPHA, Banks et Sol.

1. CARPHA schanoides, Banks et Sol. MS.; culmis cæspitosis teretibus lævibus, foliis breviusculis culmo $\frac{1}{2}$ brevioribus semiteretibus, spiculis sub 2-floris in paniculam paucifloram involucratam dispositis, setis hypogynis 6 ad apicem plumosis. C. schænoides, *Banks et Sol. MS. in Bibl. Banks. cum icone.* (TAB. CXLVIII.)

HAB. Southern parts of Tierra del Fuego; Good Success Bay, Banks and Solander; Hermite Island, Cape Horn, J. D. H.

Culmi 6-8-unciales, exspitosi, basi foliati, radices plurimas fibrosas erassas demittentes. Folia plurima, basi vaginantia, lævia, vaginis pallidis, lamina lineari-subulata, acuta, semiterete, super anguste canaliculata. Panicula involucro $\frac{1}{3}$ brevior. Spiculæ sub 3, pedicellatæ, pedicello compresso infra squamas ancipiti. Squamæ sub 5, $\frac{1}{3}$ unc. longæ, lineari-oblongæ, acuminatæ, 2 inferiores vacuæ, dorso carinatæ, carina obseure scaberula, superiores dorso convexæ, floriferæ, supremo minore vacuo. Setæ hypogynæ 6, planæ, lineares, utrinque ciliato-plumosæ, longitudine squamas æquantes, basi in tubum brevem cyathiformeni connatæ. Stamina 3, fauce tubi perigonii insertæ. Nux obovato-oblonga, stipitata, 3-costata angulis incrassatis, stylo coronata. Stylus persistens, trigonus, angulis scrratis, inferne attennatus, apice acuminatus, validus, rigidus. Semen solitarium, erectum, nuci conforme ; raphe et chalaza prominentes ; embryo parvus, oetohædrns, basi albumine inclusus, extremitate cotyledonari attenuata.

The Carpha schemoides of the hills of Fuegia, and the C. alpina, Br., of the loftier mountains of Tasmania, are two closely-allied representative species, both apparently very rare and local plants. C. alpina is replaced further north, in Anstralia, by the C. densta, Br., a native of the colony of Port Jackson, but hitherto no South American species except the one here described has been noticed. These three form together a very distinct group, as Mr. Brown has indicated (Prodr. p. 230).

PLATE CXLVIII. Fig. 1, spikelet; fig. 2, floriferous and empty scale; fig. 3, achænium, filaments, and setæ; fig. 4, base of setæ and filaments; fig. 5 and 6, achænium; fig. 7, seed; fig. 8, same, cut open; fig. 9, embryo:—all magnified.

6. CAREX,* L.

1. CAREX ovalis, Good. in Linn. Trans. vol. ii. p. 148. Engl. Bot. t. 306.

Var. B, minor, Brongn. in Duperrey, Voy. Bot. p. 149. C. Macloviana, D'Urv. in Mém. Soc. Linn. Paris, vol. v. p. 599.

* The species of this genus, and of Uncinia, have been determined and described by my kind friend Dr Boott.

362

HAB. Falkland Islands; D'Urville.

I have seen no Falkland Island specimens of this plant; can it be the C. festiva?

2. CAREX festiva, Dewey; spica composita e spiculis pluribus androgynis basi masculis in capitulum ovato-suborbiculatum arcte congestis, stigmatibus 2, perigyniis ovatis acuminato-rostratis bifidis ore antice oblique fisso nervosis marginatis denticulato-serratis squamam lanceolatam acutam æquantibus vel ea longioribus. Boott. C. festiva, Dewey, in Sill. Journ. vol. xxix p. 446. C. propinqua (?), Nees et Meyen.

HAB. Strait of Magalhaens; Port Gregory and Port Famine, Capt. King.

Culnus subpedalis, strictus, inferne glaber, foliis vaginantibus, rudimentisque foliorum pallide castaneis tectus, superne nudus, acutangulus, serrato-scaber. Folia 2 lin. lata, culmum æquantia vel breviora, margine scabra. Spica 8-9 lin. longa, 7-8 lin lata, nuda, vel braetea brevi subfoliacea basi suffulta. Spiculæ 8-12, vel plures, subrotundæ, arctissime congestæ, ferrugincæ, concolores. Squamæ lanceolatæ, aeutæ, apice membranaceo-pallidæ. Antheræ hispido-apiculatæ. Stylus exsertus. Stigmata 2, longa. Perigynium 2 lin. longum, lineam latum, utrinque nervosum, ferrugineum, marginibus alatis, e medio sursum denticulato-serratis. Achænium 8-9 lin. long., 5-9 lin. latum, oblongum, compressum, ferrugineum, basi styli abrupte apiculatum. Boott.

A C. ovali, Good., solum, spiculis pluribus, subrotundis, in capitulum arete congestis, perigyniis paululum brevioribus, differt. Boott.

Dr. Boott has kindly favoured me with the range of this species, which is so wide in the northern hemisphere that we should quite expect that it will hereafter be found along the chain of the Cordillera. Commencing in Greenland on the cast, it crosses to Unalaschka on the west by Cumberland House on Bear Lake, and thence runs south along the Rocky Monutains. In Europe it has hitherto been found in Lapland only.

3. CAREX curta, Good., in Linn. Trans. vol. ii. p. 145. Engl. Bot. t. 386, C. spicata, Banks et Sol. MSS. in Mus. Banks. cum icone. C. similis, D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 599. Kunth, En. Plant. vol. ii. p. 403.

HAB. Strait of Magalhaens; Port Famine, Capt. King; Good Success Bay, Banks and Solander; Falkland Islands, abundant, D'Urville, J. D. H.

The present, Dr. Boott remarks, is decidedly the European C. curta, one Falkland Island specimen alone, out of very many, differing from the others in having ten spiculæ, the average number being six to cight.

The geographical distribution of this species is very wide, for it inhabits all Europe from the latitude of Lapland, where, according to Wahlenberg, it is excessively common, to the Mediterranean region, which it does not enter. In Arctic America again it is abundant, extending in the United States as far south as New York.

4. CAREX acaulis, D'Urv., in Mém. Soc. Linn. Paris, vol. iv. p. 599. Brong. in Duperrey, Voy. Bot. p. 153. t. 28 A.

HAB. Falkland Islands, D'Urville.

A species wholly unknown to me, except through the figure and description of M. Brongniart.

5. CAREX decidua, Boott; spicis 4-7 atro-purpureis erectis, suprema mascula vel androgyna basi vel apice et basi mascula, reliquis fœmineis, superioribus sessilibus contiguis oblongis, inferioribus cylindraceis bracteatis evaginatis rarius geminatis, infima brevi pedunculata subremota, stigmatibus 2, perigyniis oblongoovatis rostellatis ore integro utrinque nervosis stipitatis pallidis deciduis squama oblonga obtusa atropurpurea nervo pallido decidua longioribus latioribusque. *Boott.* C. cæspitosa, *Banks et Sol. MSS. in Mus. Banks. cum icone.*

FLORA ANTARCTICA.

[Fuegia, the

HAB. Tierra del Fuego; Good Success Bay, Banks and Solander; Falkland Islands, J. D. H.

Radix stolonifera. Culmus $1-l\frac{1}{2}$ pedalis, triqueter, glaber, pars spicas gerens 2-3 poll. longa. Folia $l-l\frac{1}{2}$ lin. lata, culmo longiora, flaccida, marginibus scabridis. Bracteæ foliaccæ, evaginatæ, inferiores culmum superantes ; auriculis 2 subrotundis, ferrugineis. Spica terminalis, subpollicaris, sæpius androgyna, basi, vel apice et basi mascula; sterilis $l-l\frac{1}{2}$ lin lata; androgyna 3 lin. lata; spicæ fæmineæ 8-15 lin. longæ, 3-4 lin. latæ, inferiores interdum geminatæ, vel basi spicula minori anctæ; infima rarius l-2 pollices remota. Squamæ omnes obtusæ, atro-purpureæ, nervo pallido infra apicem evanescente; fæmineæ perigynio breviores. Pedunculus infimus 3-6 lin. longus. Perigynium (cum stipite) $l\frac{4}{9}-l\frac{7}{9}$ lin. longum, $\frac{7}{9}$ lin. latum, rarins ad margines superne denticulato-serratum, plus minus nervosnm. Achænium $\frac{8}{9}$ lin. longum, $\frac{7}{9}$ lin. latum, orbiculatum, lenticulare, fuscum, impresso-punctulatum, basi styli æquali apiculatum.

Habitus C. Goodenovii, Gay, et forsau ejus nil nisi forma aberrans. Differt spica terminali sæpius androgyna; fæmineis rarius geminatis vel compositis, perigyniisque margine scabris, culmo glabro.

6. CAREX Andersoni, Boott; spicis 7–9 atro-purpurcis erectis, terminali mascula, fæmineis 6–8 oblongis cylindraceisque superioribus geminatis ternatisque sessilibus inferioribus simplicibus pedunculatis omnibus interdum apice masculis, stigmatibus 2, perigyniis ellipticis brevi-rostratis ore integro valide nervosis stipitatis pallidis squama ovata obtusa vel lanceolata acuta atro-purpurea nervo pallido brevioribus. *Boott*.

HAB. Strait of Magalhaens; Port Famine, Capt. King.

Culmus sesquipedalis, firmus, superne acutangulus, scaber, basi vaginis foliorum tectus, pars spicas gerens 3-5 poll. longa. Folia 2-3 lin. lata, margine scabra, culmo longiora; vagina intus albo-membranacea, longa, cylindracea; ligula ad foliam alligata, ferruginea, obtusa. Bracteæ evaginatæ, suprema setacea, reliquæ latæ, foliaceæ, culmum longe superantes: auricula indivisa, amplectente, ferruginea. Spica mascula solitaria, 12-14 lin. longa, 2 lin. lata, vel exemplare unico spicâ alterâ minimâ (3 lin. longa) ad basin aucta. Squamæ latæ, obtusæ, atro-purpuræ, nervo viridi in inferioribus infra apicem evanescente; spicæ fœmineæ 6-8, contiguæ, 6-17 lin. longæ, 2-3 lin. latæ, inferiores longiores, simplices, cylindraceæ, brevi-pedunculatæ: superiores geminatæ vel ternatæ, inæquales, sessiles: omnes fœmineæ vel apice masculæ. Squamæ atro-purpurææ, ovatæ, vel inferiores lanceolatæ, muticæ, nervo pallido. Stylus inclusus. Stigmata 2, longa. Pedunculus infimus 2-8 lin. longus. Perigynium (floriferum) 1 $\frac{1}{9}$ lin. longun, lineam latun, stipitatum, breve cylindracco-rostratum, utrinque crebre et valide nervosum, pallidum, papillosum, superne marginibus parce serrato-scabriusculum, ore integro. Achænium suborbiculatum, compressum. Boott.

A *C. decidua* differt culmo validiori, firmo, acutangulo, scabro; foliis bracteisque lationibus; auricula indivisa, amplectente; spicis longioribus; terminali mascula, fœmineis sæpe apice masculis, mediis geminatis ternatisque; perigyniis paululum latioribus, squama interdum lanceolata acuta brevioribus. *Boott.*

7. CAREX *Darwinii*, Boott; spicis 8–12 ferrugineis cylindraceis longe pedunculatis nutantibus 2 terminalibus masculis, fœmineis 6–10 remotis geminatis ternatisque foliaceo-bracteatis evaginatis basi laxifloris rarius infima simplici, stigmatibus 2, perigyniis ellipticis brevi-rostratis ore integro nervosis stipitatis papillosis squamâ lanceolatâ acuminatâ hispido-cuspidata ferruginea latioribus brevioribusque. *Boott.* (TAB. CXLV.)

HAB. Chonos Archipelago, C. Darwin, Esq.

Culmus tripedalis, validus, triqueter, glaber, sulcatus, basi foliatus, apice gracillimus, pars spicas gerens 12 poll. longa. Folia bipedalia et ultra, 3-4 lin. lata, nervosa : margine carina apiceque serrato-scabra, supra nervis 2 prominentibus notata, infra ad interstitia nervorum squamato-punctata. Bracteæ emarginatæ, foliaceæ, inferiores culmum longe superantes, auricula oblonga, ferruginea. Pedunculi triquetri, scabri, inæqnales, $\frac{1}{2}$ -3 poll. longi. Spicæ fœmineæ $\frac{1}{2}$ -3 poll. longæ, 3 lin. latæ, cylindraceæ, basi laxifloræ, intervallis 2-4-pollicaribus remotæ, inferiores geminatæ, superiores ternatæ (spica interdum unica abbreviata sessili), exemplare solitario spica infima simplici,

nonnullisque apice masculis. Squamæ ferrugineæ, nervo pallido, inferiores hispido-cuspidatæ; (spicarum terminalium basis solum unius inferioris mascula sessilis adest, ceteræ disruptæ.) Perigynium $1\frac{2}{3}$ lin. longum, $\frac{8}{9}$ lin. latum, ellipticum, breve acuminato-rostratum, ore integro, utrinque 4–5-nervatum, papilloso-asperatum, maculis ferrugincis notatum, stramineo-pallidum. Achænium 7–9 lin. longum, $\frac{5}{9}$ lin. latum, orbiculato-obovatum, lenticulare, castaneum, basi styli æquali terminatum. Boott.

A C. decidua et C. Andersono spicis ferrugineis, elongatis, longo pedunculatis, nutantibus, remotis; squamis acutis, hispido-cuspidatis; perigyniis glabris, squamâ duplo brevioribus, satis distincta! Boott.

PLATE CXLV. Fig. 1, scale and stamens of male spike; fig. 2, scale and female flower; fig. 3, perigynium; fig. 4, the same cut open, showing the achaenium :-- all magnified.

8. CAREX Magellanica, Lamk.; spicis 3-4 androgynis basi masculis atro-purpureis concoloribus oblongis pedunculatis nutantibus bracteatis approximatis vel infima subradicali vaginata, stigmatibus 3, perigyniis suborbiculatis stipitatis rostellatis ore integro papillosis squama lanccolata apice acuminata involuta vix duplo brevioribus. Boott. C. Magellanica, Lam. Encycl. vol. iii. p. 385. Schkuhr, Caric. vol. i. p. 52. vol. ii p. 42. t. N. f. 51. Kunth, En. Plant. vol. ii. p. 435. C. atrata, β. Magellanica, Vahl, Act. Hafn. 1803. (TAB. CXLIII.)

HAB. Strait of Magalhaens, Commerson; Port Famine, Capt. King; Good Success Bay, Banks and Solander.

Radix e fibris lutescente-lanatis. Culmus 6-9-poll., acute triqueter, gracilis, firmus, apice scabriusculus, filiformis, pars spicas gerens plerumque $2-2\frac{1}{2}$ poll. longa. Folia $1-1\frac{1}{2}$ lin. lata, culmo breviora vel æquantia, margine carinaque scabra. Bractea infima foliacea, culmum æquans, reliquæ angustæ, spicis suis breviores, demum setaceæ, basi ligula ferruginea amplectente. Spicæ 6-9 lin. longæ, 5 lin. latæ, basi flosculis masculis paucis instructæ. Squamæ lanceolatæ, acuminatæ, apice involutæ, atro-purpureæ, concolores vel ad margines rufæ, eximie papillosæ, dorso trinerves. Pedunculi 8-14 lin. longi, capillares, apice infra spicam clavati, scabriusculi. Perigynium $1\frac{4}{9}$ lin. long., 1 lin. latum, suborbiculatum, basi productum vel latiuscule stipitatum, minime rostellatum, ore integro, obsolete vel utrinque leviter 4-5-nervatum, stramineo-pallidum, superne atro-purpureo tinctum, papillosum; stylo persistente porrecto. Achænium lineam longum, $\frac{5}{9}$ lin. latum, oblongo-triquetrum (uno specimine rarius 4-angulum).

Affinis C. limosæ, L., et congeneribus, præsertim C. irriguæ, Sm.

PLATE CXLIII. Fig. 1, scale and stamen of male spike; fig. 2, scale and female flower; fig. 3, perigynium; fig. 4, achænium:-all magnified.

9. CAREX Banksii, Boott; spicis 3-5 atro-purpureis oblongis basi cuneatis crassis exserte pedunculatis nutantibus terminali androgyna basi mascula reliquis fœmineis infima remota, stigmatibus 3, perigyniis hyalino-tenuissimis albidis compressis nervosis late ovatis cylindraceo-rostratis ore obliquo bifido glabris squama atro-purpurea oblongo-spathulata emarginata aristata brevioribus latioribusque. Boott. (TAB.CXLII.)

HAB. Tierra del Fuego; Good Success Bay, Banks and Solander, C. Darwin, Esq.

Culmus sesquipedalis, acute triqueter, glaber, basi foliatus, pars spicas gerens 4-7 poll. longa. Folia 2-3 lin. lata, carinata, culmo breviora, apice triquetro-acuminata, nervo marginibusque tuberculato-scabra. Bracteæ foliaceæ, vaginantes, infima culmum subæquans, supremæ squamæformes. Ligula elongata. Vaginæ 7 lin.-1 $\frac{1}{2}$ poll. longæ. Spicæ 3-5, omnes pedunculatæ, 7-14 lin. longæ, 4-5 lin. latæ, superiores approximatæ, infima intervallo $1\frac{1}{2}$ -4 poll. longo remota. Squamæ omnes atro-purpureæ, nervo pallidiori, laxiuscule imbricatæ. Perigynium brevissime stipitatum, $3\frac{4}{9} - \frac{7}{9}$ lin. longum, $1\frac{5}{9} - \frac{8}{9}$ lin. latum, album, tenuissimum, nervis tenuibus, rostro angusto, cylindraceo, fusco, oblique bifdo. Achænium longe stipitatum, $\frac{2}{3}$ lin. longum (cum stipite $1\frac{2}{9}$ lin. longum), $\frac{1}{2}$ lin. latum, castaneum, acute triquetrum, lateribus concavis. Boott.

4 N

FLORA ANTARCTICA.

In Herb. Banksiano sunt specimina plura (cel. Banks et Solander in Tierra del Fuego lecta) sub nominibus *C. atratæ* et *C. Magellanicæ*, quarum omnia spicam terminalem androgynam basi masculam habent, sed spicæ cylindraceæ evadunt, et hinc ad *C. germanam* tendunt. *Boott.*

Affinis C. Mertensio, Prescott.

PLATE CXLII. Fig. 1, scale and stamens of male flower; fig. 2, female flower; fig. 3, perigynium; fig. 4, ovarium, style, and stigmata; fig. 5, ovule; fig. 6, ripe perigynium; fig. 7, ripe achaenium: fig. 8, seed; fig. 9, longitudinal section of the same :---all magnified.

10. CAREX germana, Boott; spicis 4-6 fusco-ferrugineis 1 v. 2 terminalibus masculis, fæmineis 3-5 crassis cylindraceis densifloris superioribus sessilibus contiguis erectis inferioribus subnutantibus exserte pedunculatis infima interdum remota, stigmatibus 3, perigyniis ellipticis hyalino-tenuissimis albidis compressis brevi cylindraceo-rostratis ore obliquo bidentato nervosis squamam oblongam fusco-ferrugineam emarginatam aristatam subæquautibus. *Boott*.

HAB. Cape Tres Montes, C. Darwin, Esq.

Culmus 12-15-pollicaris, obtusangulus, glaber, foliis 2-3 vaginantibus instructus, apicem culmi attingentibus, basi foliatus, pars spicas gerens 4-10 pollices longa. Folia 2-3 lin. lata, glanco-viridia, apice triquetro-acuminata, nervis carina marginibusque tuberculato-scabra. Bracteæ vaginautes, superiores setaceæ, inferiores foliaceæ, culmum superantes. Vaginæ 3 lin.-2½ poll. longæ, basi purpureo-tinctæ. Spica mascula (uno specimine spicula altera minori basi aucta), 8-12 lin. longa, 2 lin. lata, squamis serrato-mucronatis. Spicæ fæminæ 12-17 lin. longæ, 4 lin. latæ, cylindraceæ, densifloræ, obtusæ, superiores sessiles vel brevi exserte pedunculatæ (uno specimine), infima remota pedunculo tres pollices extra vaginam bipollicarem exserto instructa. Squamæ arete imbricatæ, fuscoferruginæ, oblongo-spathulatæ, emarginatæ, obtusæ, nervo pallido serrato aristatæ. Perigynium 2½-3 lin. longum, 1¼ lin. latum, album, nervis subnovenis pallide ferrugineis teneribus notatum, ellipticum, apice acuminatum, rostro brevi cylindraceo fusco-purpureo, ore obliquo bidentato. Achænium longe stipitatum, $\frac{2}{3}$ lin. longum (cum stipite 12 lin. longum), $\frac{4}{3}$ lin. latum, pallide castaneum, acute triquetrum, lateribus concavis. Boott.

Affinis C. Banksio, et quoad fructum non distinguenda. Differt spica terminali mascula, fœmineis cyliudraceis densifloris, superioribus sessilibus nec basi cuneatis; squamis fusco-ferrugineis, arcte imbricatis; pedunculis validioribus; culmo obtusangulo foliisque glauco-viridibus. Boott.

I append the description of two new species of extra-tropical South American Carices; which, with those enumerated in the body of this work, include all that I know to exist in western Chili and Fuegia *.

1. CAREX acutala, Boott; spicis 5-6 erectis cylindraceis fuscis masculis 1-2 sessilibus reliquis 4 formineis sæpe apice subulato-acutatis masculis densifloris sessilibus vel pedunculatis longe foliaceo-bracteatis alternatim contiguis, stigmatibus 3, perigyniis elliptico-lanceolatis subinflatis nervosis glabris uitidis squama purpureo-ferruginea concolori vel apice hydina ciliata hispido-aristata longioribus. *Boott.* C. physocarpa, *Nees, in Herb. Hooker (non Presl.*)

HAB. In America merid. Ins. Chiloe, *Cuming*, n. 43. In Mont. Pilzhum, Columbia, ad alt. 12,000 ped., Jameson, (Herb. Hooker.)

Culmus acutangulus, validus, scabriusculus, pars spicas gerens 3-6 poll. longa. Folia 4 lin. lata, culmo longiora. Bracteæ foliaceæ, infima 3 lin. lata, culmum longe superans, nunc brevissime vaginata. Spicæ masculæ sessiles, $1-1\frac{1}{2}$ poll. longæ, $1\frac{1}{2}$ lin. latæ. Squamæ ferrugineæ, concolores, vel apice hyalino-albidæ, ciliatæ, brevi hispidomucronatæ. Spicæ fæmineæ 4, (pars suprema plerumque tertia omnium sæpius subulato-acutata, mascula vel sterilis) $1\frac{3}{4}$ poll. longæ, 4 lin. latæ, densifloræ, superiores sessiles, approximatæ, inferiores plus minus longe pedunculatæ, intervallis $1\frac{1}{2}-2$ poll. longis, remotæ, tamen omnes ob longitudinem pedunculorum contiguæ. Squamæ purpuræ,

11. CAREX indecora, Kunth; spicis 3-5 oblongis erectis terminali mascula clavata subsessili reliquis fœmineis bracteatis sessilibus contiguis vel infima subremota exserte pedunculata, stigmatibus 3, perigyniis oblongo-ovatis acuminato-rostratis bidentatis demum ore integro utrinque leviter nervosis stramineis lucidis squama æquilata purpurea trinervi obtusa vel emarginata hispido-cuspidata longioribus. *Boott*.

Var. β , humilis. C. fuscula, D'Urville, in Mém. Soc. Linn. Paris, vol. iv. p. 599. Brongn. in Duperrey, Voy. Bot. p. 154. t. 28 b.

HAB. Falkland Islands, D'Urville, J. D. H.

Culnus 4-12 poll., obtusangulus, lævis, basi foliatus, versus medium folio vaginante instructus, pars spicas gerens $\frac{1}{2}$ -5 poll. longa. Folia $1\frac{1}{2}$ -2 lin. lata, plana, culmo breviora, rigidiuscula, flavescente-viridia. Bracteæ erectæ, infimæ culmum superantes, vaginatæ, superiores angustæ, vaginæ 2-7 lin. longæ. Spicæ 3-5, omnes interdum congestæ, sessiles; spica mascula 3-6 lin. longa, lineam lata, clavata, sessilis vel brevi-pedunculata. Squamæ uninerves, obtusæ, cuspidatæ. Spicæ fæminæ 4-7 lin. lougæ, 2-3 lin. latæ, contiguæ, vel infima intervallo 1-5 poll. longo remota, exserte (vel binæ inferiores plus minus longe exserte) pedunculata. Squamæ purpuræ, trinerves, obtusæ vel emarginatæ, valide hispido-cuspidatæ. Pedunculi 6-12 lin. longi, glabri, nunc vix exserti. Stylus inclusus. Perigynium $1\frac{7}{9}$ lin. long., $\frac{7}{9}$ lin. latum, oblongo-ovatum, sensim acuminato-rostratum, bidentatum, demum ore integro, stramineum, punctis ferrugineis notatum, lucidum, punctulatum, glabrum, vel rarius superne ad margines serrato-scabrum, leviter (luci subjectum) utrinque nervosum. Achænium $\frac{7}{9}$ lin. long., $\frac{5}{9}$ lin. latum, pallidum, subrotundo-triquetrum, punctulatum, basi styli æquali apiculatum. Boott.

concolores vel apice hyalino-albidæ, ciliatæ, nervo lato viridi in aristam latam hispidam producto. Pedunculi validi, erecti, infimus $\frac{1}{2}-2$ poll. longus, evaginatus vel e vagina 4 lin. longa exsertus. Perigynium $1\frac{8}{9}$ lin. longum, $\frac{5}{9}$ lin. latum, nitidum, erebre nervosum, pallide viride, basi purpureo tinctum, pellucido-punctatum. Achænium (vix maturum) $\frac{8}{9}$ lin. longum, oblongo-triquetrum, pallide stramineum, basi styli incrassato terminatum. Boott.

Affinis C. paludosæ, Good.

2. CAREX paleata, Boott; spicis 7–10 cylindraceis masculis 2–4 sessilibus contiguis extremis longioribus infima longe bracteata fœmincis 3–7 remotis exserte ligulato-pedunculatis longissime bracteatis densifloris basi attenuatis inferioribus nutantibus, stigmatibus 2–3, perigyniis obovatis rostellatis bifidis nervatis nervisque 2 marginalibus pallidis scabris cinctis olivaceis purpureo-maculatis squama ovata paleacea obtusa vel acuta trinervi late hispidocuspidata brevioribus longioribusque. *Boott*.

HAB. In Ins. Juan Fernandez, Dr. Scouler. (Herb. Hooker et Fielding.) Cuming, n. 1341. (Herb. Boott.)

Culmi pars superior solum adest, triquetra, lævis, inter spicas scabriuscula, pars spicas gerens 10 poll. ad 2 ped. louga. Folia desunt. Bracteæ omnes culmum superantes, infima 2 lin. lata, superiores sensim angustiores. Spicæ masculæ 2-4, sessiles, contiguæ, 7-20 lin. longæ, $1-1\frac{1}{2}$ lin. latæ, castaneæ, extremæ longiores, infima longe bracteata. Spicæ fæminæ 5-7, intervallis $2\frac{1}{2}-3\frac{1}{2}$ poll. remotæ, $1\frac{1}{2}-2\frac{1}{2}$ -poll. longæ, 2 lin. latæ, cylindraceæ, densifloræ, basi attenuatæ, duæ superiores nunc apice masculæ, suprema interdum inclusa, pedunculata. Pedunculi ligulato-compressi, inferiores $2-2\frac{1}{2}$ poll. longi, glabri, supremus interdum abbreviatus, infimusque versus apicem squamas alternas steriles longe cuspidatas ferens. Vaginæ 3 lin.-2 poll. longæ, glabræ. Perigynium $1\frac{3}{9}$ bin. longun, $\frac{7}{9}$ bin. latum, obovatum, rostellatum, bifidum, laciniis serratis, nervatum, nervisque 2 marginalibus prominentibus pallidis superne scabris ciuctuu, pallide olivaceum, purpureo-maculatum, plano-convexum vel triquetrum, coriaceum. Achænium $\frac{6}{9}$ bin. longun, $\frac{5}{9}$ lin. latum, obovatum, plano-triquetrum, atro-olivaceum, cavitatem perigynii implens. Squamæ omnes ovatæ, acutæ vel obtusæ, trinerves, late hispido-cuspidatæ; masculæ castaneæ; fœmineæ membranaceo-pallidæ. Boott.

Affinis C. lucidæ, Boott.

Affinis C. extensæ, Good., quæ perigyniis costato-nervosis, glaucis, squamis masculis muticis, foliis, bracteisque patentibus vel recurvis, sæpe involutis, differt. Boott.

12. CAREX trifida, Cav., vid. Fl. Antarct. Pt. 1. p. 89.

HAB. Cape Tres Montes, C. Darwin, Esq.; Falkland Islands, abundant, D'Urville, Capt. Sulivan, J. D. H.

A very noble species, abundant in the Falkland Islands, growing with, and emulating in size, young Tussock grass. Mr. Darwin alone has gathered it on the American continent, and he only at Cape Tres Montes. Its confined range is very singular, for it can scarcely have been overlooked in Fuegia or the Strait of Magalhaens, had it existed there; and it is also probably the only plant common to New Zealand and the Falkland Islands, not found abundantly in Tierra del Fuego.

Carex trifida affords a remarkable instance of apparent caprice in its choice of habitat; for though common in the Falklands, along with the Dactylis caspitosa (Tussock grass), and though there these grow in company, and under precisely the same conditions, yet the Tussock grass in America only appears in the southern extreme of Fuegia, where it is unaccompanied by Carex trifida; whilst the latter is confined to a latitude eight hundred miles north of Cape Horn. There is nothing whatever in the climate or soil of any part of western South Chili, or Fuegia, that can be pronounced unfavourable to the growth of this *Carex*, whose absence there naturally leads to the question, how is its presence in Cape Tres Montes and the Falkland Islands to be accounted for? did it originate in each of these two isolated localities? was the seed transported over the intervening land, by an agent whose operations were limited to the eastern, and western extremes only of Antarctic America? or, have the individuals that once tenanted the intervening land, been destroyed? Any one of these hypotheses is at first sight plausible, and the first, perhaps, the most so, New Zealand being a third, and far more remote, habitat for this same species, which may thus be supposed to have had three separate origins. Such a question should not be discussed with reference to a single species, but as one which concerns all organized nature, whose phenomena are amenable to general laws. Hypotheses, adopted to account for exceptional cases, if not viewed in reference to the general rule from which these exceptions deviate, are generally fallacious; and however much so, they still are apt to be magnified into laws. If we knew only such plants as are sporadic (the term given to species which inhabit unconnected and remote localities) we might, perhaps, be justified in assuming it as an axiom, that individuals of a species have sprung, at isolated localities, from as many similar parents : the cases which appear to demand this solution are, however, exceptions in Botanical Geography.

The study of the distribution of any one species or genus, or of the Flora of any one country, does not afford scope enough for investigating satisfactorily such a subject as the origin of the individuals of plants. If species, genera, and small natural orders were sporadic, recurring wherever climate and soil presented similar conditions, several points of origin for the same species might be assumed. But it is not so: species, genera, and orders are distributed within geographical limits, according to their extent : the great mass of individual plants in the one case, and of forms in the other, appear to have sprung from single centres, in the former case from a common parent, and to have radiated from one point to greater or less distances around it, in proportion to the facilities for migration and absence of checks to diffusion. The explanation of exceptions to this prevailing rule must then be sought in some natural cause, capable of counteracting the general law, and not what, if adopted for the case of one species, must be conceded with respect to all, and consequently force us to conclude that two classes of agents are required to effect one object, namely, the dispersion of vegetables.

7. UNCINIA, Pers.

1. UNCINIA tenuis, Poepp., Synops. Plant. Am. Austr. vol. iii. n. 240. Kunze, Synops. der Reidgr. t. 21. Kunth, En. Plant. vol. ii. p. 525.

HAB. Strait of Magalhaens; Port Famine, Capt. King; Hermite Island; Cape Horn, J. D. H.

A species entirely confined to South Chili, between Concepcion and Cape Horn.

The four species enumerated in this work, together with *U. erinacea*, Pers. (a native of Valdivia and Chili) and two new ones *, diagnoses of which Dr. Boott has kindly given, include all the extra-tropical American *Unciniæ* known to me.

2. UNCINIA phleoides, Persoon, Synops. vol. ii. p. 534. Brongn. in Duperrey, Voy. Bot. p. 158 (excl. syn. U. Maelovianæ). Hook. et Arn. in Bot. Voy. Beechey, p. 50. Carex phleoides, Cav. Icon. vol. v. p. 40. t. 464. f. 1.

HAB. Chonos Archipelago; C. Darwin, Esq.

On several occasions I have alluded to the change which occurs in the vegetation of the western coast of South America, at, or about, the latitude of the Chonos Archipelago. This arises from many species extending to (but not crossing) that limit, both from much lower and higher latitudes, of which the present plant affords an example. U. phleoides inhabits the plain of Quito, under the equator, at an elevation of 8,000 feet; it grows also at

1. UNCINIA *multifaria*, Nees; spica crassa densifiora basi attenuata apice conico mascula nuda, stigmatibus 3, perigyniis (arista divaricata vix duplo brevioribus) linearibus ore truncato striato-nervosis scabris margine ciliatis squama oblonga obtusa pallida apice albo-membranacea ciliolata angustioribus longioribusque. *Boott.*

HAB. Chiloe, Cuming (n. 44. Herb. Hooker.)

Culmus subbipedalis, triqueter, firmus, lævis, inferne foliatus. Folia 3-4 lin. lata, culmo longiora vel æquantia, glaucescentia, margine versus apicem facicque scabra, supremum angustum. Spica $2\frac{1}{2}$ poll. longa, superne 6 lin. vel aristis divaricatis mensurata 10 lin. lata, basi attenuata, (1 lin. lata), nuda; apice conico, (4 lin. longo), mascula. Squamæ oblongæ, obtusæ, pallidæ, demum fuseæ, apice ciliolatæ, albo-membranaceæ, infra apicem ferrugineo-zonatæ, nervo dorsali vix prominente; masculæ breviores. Perigynium $3\frac{2}{9} - \frac{3}{9}$ lin. longun, $\frac{1}{2}$ lin. latum, biconvexum, superne præcipue scabrum, margine ciliatum, pilis sursum longioribus demum fasciculatis, ore truncato ciliolato, arista 2 lin. extra os exserta, 5 lin. longa, divaricata, imo basi torta. Achænium 2 bin. longum. $\frac{4}{9}$ lin. latum, triquetrum, utrinque sursum convexum, fuscum, impresso-punctulatum, apice et basi attenuatum. Stylus basi subincrassatus. Stigmata 3, non plumosa. Boott.

Ab U. erinacea, Pers., perigymis linearibus diversa.

2. U. Douglasii, Boott; spica clongata lineari nuda apice mascula conformi, stigmatibus 3, perigyniis (arista $\frac{1}{3}$ brevioribus) lanceolatis convexo-concaviusculis basi obconico attenuatis ore truncato plurinerviis margine scabris superne pilis appressis utrinque exasperatis pallidis squama amplectente ovata acuminata obtusa flavescenti-viridi angustioribus sublongioribusque. Boott.

HAB. Ins. Juan Fernandez. David Douglas. (Herb. Hooker.)

Culmus bipedalis, gracilis, lævis, nudus, basi foliatus. Folia $1-1\frac{1}{2}$ lin. lata, culmo longiora, utrinque marginibusque scabra. Spica $5\frac{1}{2}-6$ poll. longa, lincam lata, pars suprema mascula, subpollicaris, conformis. Squamæ ovatæ, acuminatæ, obtusæ, amplectentes, flavescenti-virides, striatæ, margine pallide-ferrugineæ, perigynio vix longiores, omnes conformes. Perigynium $2\frac{1}{2}-3$ lin. longun, $\frac{1}{2}$ lin. latum, lineare, hine convexum, inde concaviusculum, basi obconico-attenuatum, dorso plurinervium, marginibus e basi scabrum, pilis sursum longioribus, superne pilis brevioribus appressis utrinque exasperatum, pallidum, lineolis ferrugineis maculatum, ore truncatum. Achænium $1\frac{5}{9}$ lin. longum, $\frac{3}{9}$ lin. latum, lineare, convexo-concaviusculum, facie dorsali linea centrali (angulo) notatum, castaneum, impresso-punctulatum. Arista $3-\frac{3}{9}$ lin. longa, pallida, filiformis, apice ferruginea, perigynio $\frac{1}{3}$ longior. Stylus inclusus. Stigmatibus 3. Boott.

4 0

Valparaiso, and Concepcion, again at Valdivia, finally disappearing at the Chonos Archipelago. Though we are now fairly acquainted with the botany of America south of lat. 33°, a more complete collection from the coast and mountains between the southern extreme of Chiloe and Cape Tres Montes is wanting; the proportion of new species would probably be small, but the investigation would exhibit the range of many Valdivian and Fuegian plants, not contained in the invaluable Herbarium of Mr. Darwin, the only naturalist whose good fortune it has been to visit and explore that unfrequented line of coast.

3. UNCINIA Macloviana, Gaud., in Ann. Sc. Nat. vol. v. p. 99, et in Freye. Voy. Bot. p. 412. Kunth, En. Plant. vol. ii. p. 526.

HAB. Falkland Islands; Gaudichaud.

When botanizing in the Falkland Islands early in the winter of 1841, I found what I considered to be this plant, growing amongst grass in wet spongy bogs; it was, however, in a very bad state, and the specimens, unfortunately, lost.

Brongniart unites this with U. phleoides, Pers.; but M. Kunth has kept it distinct.

4. UNCINIA *Kingii*, Boott; spica capitata fusca nuda apice mascula, stigmatibus 3, perigyniis (arista ¹/₃ brevioribus) lanceolatis superne angustiori cylindraceis ore truncato oblique fisso ferrugineis glabris squama lanceolata fusco-ferruginea nervo pallido angustioribus longioribusque. *Boott*. (TAB. CXLV.)

HAB. Strait of Magalhaens; Port Famine, Capt. King.

Cæspitosa. Radix repens, fibroso-lanatus. Culmus 2-4 poll., lævissimus, sudcatus, basi vaginis foliornm castaneis laceratis tectus. Folia angusta, involuta, hine filiformia, culmo breviora, apice margineque scabra. Spica 5-7 lin. longa, 3-6 lin. lata, congesto-capitata, apice flosculis masculis paucis inconspicuis, basi fæmineis 9-16 instructa. Squamæ fæmineæ lanceolatæ, infima mucronulata. Perigynium (cum arista, stipiteque) $4\frac{1}{2}$ -5 lin. longum, $\frac{5}{9}$ (ad basin) latum, superne cylindraceo-attenuatum, ore oblique fisso, fusco-ferrugineum, basi pallidum. Achænium 1 lin. long., $\frac{1}{2}$ lin. latum, oblongo-triquetrum, pallidum, basi styli incrassato apiculatum. Arista $4-4\frac{1}{2}$ lin. longa, canaliculata, pallida, superne ferruginea, imo apice dilatata. Stylus inclusus. Stigmata 3, brevia. Boott.

PLATE CXLV. Fig. 1, scale and male flower; fig. 2, scale and female flower; fig. 3, female flower, removed from the perigynium :--all magnified.

L. GRAMINEÆ,

1. ALOPECURUS, L.

1. ALOPECURUS alpinus, Smith, Engl. Bot. t. 1126. Kunth, En. Plant. vol. i. p. 25.

Var. β, aristatus. A. alpinus, Trinius, Ic. Gram. vol. i. t. 38. A. pratensis, Banks et Sol. in Mus. Banks. A. pratensis, var. spica ovata; Ledebour, in Herb. Hook. A. Baicalensis, Turz. in Herb. Hook. A. Antarcticus, Vahl, Symb. vol. ii. p. 18. Brongn. in Duperrey, Voy. Bot. p. 16. Kunth, Agrost. p. 25. A. Magellanicus, Lamk. Illust. Gen. vol. i. p. 168. Gaudichaud, in Ann. Sc. Nat. vol. v. p. 100, et in Freye. Voy. Bot. p. 131. D'Urville, in Mém. Soc. Linn. Paris, vol. iv. p. 600. (TAB. CXXX.)

Var. y, gracilior; spica angustiore.

HAB. From the Strait of Magalhacus to Cape Horn, and throughout Fuegia and the Falkland Islands, abundant, Commerson, Banks and Solander, and all succeeding voyagers. Var. β , Port Gregory, Capt. King.

This plant I believe to be specifically the same with the North-European and American *A. alpinus*, of which Smith considered it to be a variety. What appeared specific differences, were pointed out by Mr. Brown (*in Appendix*

to Parry's 1st Voyage, p. 184.). Since the publication of the last mentioned work it has been universally looked as an Antaretie species alone, and its close affinity with the *A. pratensis*, of the Northern Hemisphere was never alluded to. The ordinary states of the latter plant have a longer and less hairy spike; but amongst the varieties of it which occur in North Western Asia, and N. Eastern America, there is one wholly undistinguishable from Antaretie individuals; and how far these may be constantly distinct appears very doubtful to me. Mr. Brown, in drawing up the characters of *A. alpinus*, alludes to his having gathered Scotch specimens with an arista twice as long as the glumes, such is the case with all the Antaretie ones, and in Trinius's figure of A. alpinus; but is at variance with Smith's specific character, (founded on Mr. Brown's specimens) and with the ordinary state of the Scotch plant. Mr. Watson, however, has gathered the same aristate variety of *A. alpinus* in Scotland, and has cultivated both forms in his garden. His garden specimens of both states are now before me, the long awned one retaining its characters, and the awns of the common form decidedly elongating under cultivation. The comparative length of the lamina and vagina of the uppermost leaf, is also very variable, even in *A. alpinus*, these being sometimes of equal length, while in the Antarctic plant the lamina is sometimes considerably the shorter; and, again, I have examined an European specimen of A. pratensis, in which the lamina is even longer than the vagina. The other characters of A. pratensis, used by Mr. Brown, are those of the glumes being acute, and villous only at the sides: this is the case with the British examples that I have studied, but not with the Siberian, which certainly present intermediate forms between this species, and its Fuegian congener. The Antaretic specimens vary exceedingly in size, from four inches, to two and even three feet high; the culms are generally tumid above the upper leaf and contract gradually towards the paniele; or they are slender, cylindical and terete: the lamina of the upper leaf is occasionally far shorter than at other times, equal in length to, or much longer than its vagina. Spikes nearly eylindrical, 2-3 to $1\frac{1}{2}$ inches long, generally rather more than twice as long as broad, but now and then much narrower. Glumes always more or less villous all over.

Admitting the foliage to afford no specific character between *A. alpinus, A. pratensis*, and *A. Antarcticus*, and the length of the arista to be very variable in the first of these, there remains no constant character to distinguish these three; for between *A. Antarcticus* and *A. pratensis* the only apparent distinctions lie in the villosity of the glumes, and the form of the spike, differences which do not hold in Siberian specimens of the latter. I have added a plate of the common Falkland Island state of this species.

PLATE CXXX. Fig. 1, glumes and floret; fig. 2, floret removed from the glumes; fig. 3, pistil :---all magnified-

2. PHLEUM, L.

1. PHLEUM alpinum, Linn. Sp. Pl. p. 88. Banks et Sol. in Bibl. Banks. Engl. Bot. t.519. P. Hænkeanum, Presl, Rel. Hænk. vol. i. p. 245. Nees, in Nov. Act. Acad. vol. xix. Suppl. p. 140.

HAB. Strait of Magalhaens; Port Famine and Port Gregory, Capt. King. Good Success Bay, Banks and Solander.

This species, which is associated in the mountains of Scotland with *Alopecurus alpinus*, also accompanies that plant in the southern regions. It has been gathered by Mr. Bridges, on the east side of the Andes of Chili, at an elevation of 6-7,000 feet; and also on the Cordillera of Mexico by Lindeu, and by Galcotti on the Peak of Orizaba, at an elevation of between 10 and 12,000 feet.

3. MÜHLENBERGIA, Schreb.

1. MÜHLENBERGIA rariflora, Hook. fil.; rigida, glaberrima, panieula effusa pauci- sub IO-flora, glumis subæqualibus enervibus flosculo paulo brevioribus, palea inferiore lanceolata coriacea basi glaberrima in aristam longissimam rigidam scaberulam desinente superiorem breviorem amplectante, culmo foliato, foliis rigidis sctaceis marginibus involutis. (TAB. CXXXI.)

HAB. Cape Tres Montes; Patch Cove, 2,000 feet, C. Darwin, Esq.

372

FLORA ANTARCTICA.

Gramen rigidum, cæspitosum, 4-6 pollicare. Culmi basi ascendentes, phries divisi, vaginis coriaceis nitidis striatis foliorum vetustorum obtecti, parte superiore usque ad paniculam vaginati. Foliorum vagina 1-2 unc. longa, teres, glaberrima, profunde striata; ligula brevis; lamina vagina brevior v. superans, erecta, culmo brevior, rigida, anguste setacea, apice pungens, folii superioris paniculam fere superans. Panicula $1\frac{1}{2}$ unc. longa, pedunculo pedicellisque flexuosis, elongatis, lævissimis. Spiculæ purpureæ, nitidæ, vix 2 lin. longæ. Glumæ membranaceæ, lanceolatæ, flosculo paulo breviores, inferiore paulo majore. Flosculus brevissime pedicellatus, pedicello barbato. Palea inferior in aristam desinens; arista $1-1\frac{1}{2}$ unc. longa, siccitate curvata, madore recta, rigida, sub lente scaberula, apice gradatim attenuata, basi obscure articulata, haud v. vix torta, angulata. Squamulæ 2, lineari-oblongæ, obtusæ. Stamina 3. Ovarium stipitatum, supra medium constrictum.

Allied to *M. capillaris* of North America, in the form of the locustæ; but a very different species, and, I think, decidedly of the genus *Mühlenbergia*. The rigidity of the arista is quite like that of *Stipa*, as is the harsh foliage, while in other respects the plant has more affinity with the *Agrostideæ*.

PLATE CXXXI. Fig. 1, locusta; fig. 2, floret with portion of the awn removed; fig. 3, squamula; fig. 4, ovarium:---all magnified.

4. AGROSTIS, L.

1. AGROSTIS tenuifolia, Bieb., Flor. Taur. Cauc. vol. i. p. 56. Trinius, Ic. vol. iii. t. 35. Kunth, En. Plant. vol. i. p. 220.

Var. Fretensis; locustis paulo majoribus.

HAB. Var. Fretensis, Strait of Magalhaens; Port Famine, Capt. King.

I have compared this grass most carefully with authentic specimens of A. tenuifolia from Persia and the Caucasus, without being able to detect any further difference than in the size of the locustæ, which in the Antarctic plant are $\frac{1}{8}$ of an inch long, the Caucasian scarcely $\frac{1}{10}$. Intermediate between them is a common Rocky Mountain species, collected by Douglas, and described as A. exarata, β ., in the 'Flora Boreali-Americana' (vol. 2. p. 239). There are, however, two forms of A. exarata β , one from the east side of the Rocky Mountains, which has the scabrid broader leaves of the true A. exarata, and a distinct upper palea (this is the A. Drummondi, Torrey MS.), the other (or Douglas's), from the west side of the dividing ridge, is smaller, more slender, with small locustæ, and no upper palea; it agrees closely with the Magellanic plant in size and foliage, and bears the name of A. tenuifolia? Bieb., appended to it by Dr. Torrey.

The culms of \mathcal{A} . tenuifolia β . are 15 inches to $2\frac{1}{2}$ feet long, smooth, erect, and very slender. Leaves subsetaceous, obscurely scabrid. Lower palea truncate, 4-toothed and 4-nerved, with or without a short dorsal awn. Upper palea none, or when present extremely short.

2. AGROSTIS alba, Linn., Sp. Pl. p. 93. Engl. Bot. t. 1189. A. cæspitosa, Gaud. in Ann. Sc. Nat. vol. v. p. 100, et in Freyc. Foy. Bot. p. 131. Kunth, Agrost. p. 219.

Var. β , stolonifera. A. stolonifera, Linn. Sc.

HAB. Falkland Islands, both varieties abundant, but possibly introduced; Gaudichaud, Mr. Wright, J. D. H.

The lower palea in my specimens is obscurely 4-nerved, or, in var. β ., 5-nerved, with occasionally a very short awn, never projecting beyond the glumes. The upper palea is one third shorter than the lower. This grass forms a very good pasturage, both in the upland and lowland districts about Port Louis, but is not very abundant, which it may become if it be an introduced plant. The var. *stolonifera* is the famous 'Fiorin grass,' or 'Squitch' of Dr. Richardson and the Irish agriculturists.

3. AGROSTIS *prostrata*, Hook. fil.; eulmo longe procumbente geniculato stolonifero, panicula erecta contraeta lineari-oblonga densiflora, glumis latiusculis aeuminatis carina scabrida flosculum basi glaberrimum superantibus, palea inferiore truncata apice erosa enervi superiore bis longiore, arista nulla, foliis breviusculis planis glaberrimis v. obscure scaberulis.

HAB. Falkland Islands; boggy ground on Hog Island, Berkeley Sound, rare, J. D. H.

Gramen humile, glaberrimum, repens. Culmi prostrati, 3–4 unc. longi, parte ascendente bi- tri-pollicare, nodosi, stoloniferi, foliosi, glaberrimi, internodo terminali solummodo erecto, unifoliato. Foliorum vagina elongata, profunde striata, glaberrima; ligula breviuscula, late ovata, membranacea; lamina vagina brevior, patens, sub 1 unc. longa, plana, striata, e basi latiuscula ad apicem acuminatum gradatim angustata. Panicula uncialis, $\frac{1}{3}$ unc. lata, interrupta, densiflora, ramis ramulisque brevibus, fastigiatis. Locustæ glaberrimæ, $1\frac{1}{2}$ lin. longæ, mitidæ.

To all appearance a very distinct species, allied to *A. alba*, var. *stolonifera*, but differing in the much smaller size, coarctate panicle, smaller locustæ and florets, absence of an arista, &c.

4. AGROSTIS *Falklandica*, Hook. fil.; dense cæspitosa, glaberrima, panicula gracili ramis erectis, glumis æqualibus ovato-laneeolatis acuminatis glabratis carina scaberula flosculis basi nudis $\frac{1}{2}$ longioribus, palea inferiore apiec erosa truncata obscure 5-nervi nervo medio infra medium evanido, arista glumis breviore v. nulla, palea superiore nulla, foliis radicalibus filiformibus culmo graeili erecto longioribus.

Var. a, culmo folia bis terve superante.

Var. β , culmo folia vix superante.

HAB. Falkland Islands; var. a and β in marshy places, on rocks near the sea, and on the hills, abundant.

Gramen dense cæspitosum, gracile, 3 unc. ad pedalem. Culmi e basi crecti, foliis interdum duplo triplove longiores, basi foliati, superne exemplaribus elatioribus longe nudi, læves, obscure striati, glaberrimi, Foliorum inferiorum vagina breviuscula, gracilis, striata, 1 unc. longa, superiornm elongata profundius striata, 2–3 unc. longa; ligula membranacea, truncata; lamina angustissima, filiformis, erecta, herbacea, glaberrima, 3–5 unc. longa, apice gradatim angustata. Panicula $\frac{3}{4}-1\frac{1}{2}$ unc. longa, ramis erectis elongatis paucifloris, in var. β brevioribus. Locustæ sub $1\frac{1}{2}$ lin. longæ, juniores puberulæ, demum glabratæ.

Apparently a variable plant, its very narrow filiform leaves are characteristic of this species amongst its Antarctic allies.

5. AGROSTIS Magellanica, Lamk. (?); glaberrima, cæspitosa, panicula elongata laxiflora nutante v. inclinata, ramis pedicellisque scabridis, glumis majusculis æquilongis glaberrimis nitidis carina scabridis flosculo basi barbato fere triplo longioribus, palea inferiore apice truncata 4-dentata obscure 5-nervi, nervo intermedio ad medium in aristam recurvam glumas superantem desineute, palea superiore inferiore $\frac{1}{2}$ breviore, foliis planis longe lineari-lanceolatis gradatim angustatis, culmis cæspitosis vaginatis. A. Ma gellanica, Lamk. Illust. Gen. n. 807. Poiret, Encycl. Méth. Suppl. vol.i. p.207. Kunth, Agrost. p.221.

HAB. Strait of Magalhaens; Port Famine, Capt. King.

Culmi erecti, basi cæspitosi, $1-1\frac{1}{2}$ pedales, glaberrimi, nitidi, herbacei, foliis vaginati, stricti, infra paniculam scaberuli. Folia panca; vagina elongata, 3-5 unc. longa, profunde striata, glaberrima; ligula membranacea, oblonga, obtusa; lamina suberecta v. patens, plana, linearis, striata, gradatim acuminata, herbacea, vaginæ suæ subequilonga. Panicula subcontracta, elongata, 3-5-pollicaris, nutans v. inclinata; ramis verticillatis, erectis, divisis pedicellisque scaberulis. Locustæ sub 2 lin. longæ, micantes. Glumarum valvæ subæquales, compressæ, dorso scaberulæ, acuminatæ, flosculo fere ter longiores. Flosculi basi barbati. Palea inferior membranacea,

4 P

nervis obscuris, dorso arista basi recurva deinde incurva instructa. Arista glumas superans, gracilis, scaberula. Palea superior latiuscula, membranacea, obscure bifida,

I have presumed this to be the *A. Magellanica* of Lamarck, for it agrees with his insufficient description, and also with the longer one given by Poiret, except that the awn is not terminal, though so described (possibly through inadvertence) by that author. As a species it is very nearly allied to the following, but may be distinguished by the larger glumes, greater size, and conspicuous upper palea.

6. AGROSTIS Antarctica, Hook. fil.; erecta, cæspitosa, panicula elongata nutante v. inclinata subdensiflora, ramis subverticillatis pedicellisque scabridis, glumis æqualibus pilosiusculis glabratisve carina scabridis flosculum basi glaberrimum bis longioribus, palea inferiore apice truncata 4-cuspidata 5-nervi, nervo intermedio ad medium in aristam glumas superantem desinente, superiore parva, squamulis oblongoacinaciformibus subacutis. A. Magellanica, *Gaud. in Ann. Sc. Nat.* vol. v. p. 100, *et in Freye. Voy. Bot.* p. 131 (?). (TAB. CXXXII.)

HAB. South Chili and Fuegia, from the Chonos Archipelago to Cape Horn, the Falkland Islands and Kerguelen's Land, very abundant.

Statura variabilis. *Culmi* 2 unc. ad bipedalem, graciles, superne nudi v. vaginis foliorum tecti. *Folia* et *inflorescentia A. Antarctica*, sed foliorum vaginæ plerumque latiores, panicula densior, locustæ minores, arista paulo longior, paleaque superior multoties minor.

Agrostis Antarctica is one of the most abundant of grasses in the regions it inhabits, especially in swampy grounds, which seem particularly favourable to its growth. It is also a very elegant plant, from its graceful habit and the form of its nodding panicle. It may be the true *A. Magellanica* of Lamarck, and judging from its abundance, appears natural to suppose so; but the very short upper palea is not alluded to in that author's description, and Poiret's observation that the upper is the longest, would imply that there is no remarkable difference in their length. Considering the invalid nature of the characters afforded by the comparative length of the palea in this genus, it is probable that this and the preceding are but varieties of one and the same plant.

Kerguelen's Land specimens are frequently monstrous; the lower glume being then provided with two parallel distinct nerves, and in other cases I have seen three distant valves, two outer and one inner. The lower palea again has the arista sometimes placed on one side of its base.

PLATE CXXXII. Fig. 1, locusta; fig. 2, floret; fig. 3, squamulæ and pistil; fig. 4, squamula :---all magnified.

5. POLYPOGON, Desf.

1. POLYPOGON *Chonoticus*, Hook. fil.; panicula ampla oblonga subeffusa lobata densiflora, ramis glabriusculis pedicellisque scaberulis, glumis pubescentibus apice oblique truncatis aristis valvis bis longioribus, palea inferiore superne 5-nervi truncata 5-aristata aristis 2 lateralibus subelongatis intermedio palea triplo longiore, culmo vaginato, foliis planis scaberulis striatis vaginis brevioribus.

HAB. Chouos Archipelago and Cape Tres Montes, C. Darwin, Esq.

Gramen pulchrum, bipedale. Culmi validi, erecti, per totam longitudinem vaginati. Folia radicalia breve vaginantia, superiorum vagina internodos fere æquans, glaberrima, lævis, profunde striata; ligula breviuscula; lamina 5-pollicaris, lanceolato-subulata, e basi latiuscula gradatim angustata, super præcipue scaberula. Panicula 4-5 une. longa, $1-1\frac{1}{2}$ lata, lobata, sericea, ramis e copia locustarum velatis. Glumæ $1\frac{1}{2}$ lin. longæ, pubescentiscaberulæ, carina scabrida, apice oblique truncata, vix acuta, in aristam pallidam v. purpuream desinentes, flosculos longiores. Palea inferior membranacea, basi enervis, superne 5-nervis, nervis 2 lateralibus in aristas paleæ

æquilongas productis, nervo intermedio in aristam terminalem tenuissimam aristis glumarum breviorem producto; palea superior brevior, apice bidentata.

The four-aristate lower palea of this species distinguishes it at once from any of its congeners. Mr. Darwin's, and one gathered in Chiloe by Capt. King, are the only specimens I have seen.

6. ARUNDO, L.

1. ARUNDO pilosa, D'Urville, in Mém. Soc. Linn. Paris, vol. iv. p. 600. Kunth, Agrost. p. 247. Ampelodesmos anstralis, Brongniart, in Duperrey Voy. Bot. p. 31. t. 6.

HAB. Falkland Islands, abundant; D'Urville, Mr. Wright. Capt. Sulivan, J. D. H.

A fine species and first pointed out to me by my friend Governor Moodie, as forming, next to the Tussock, the most useful grass in the Falkland Islands, for fodder. It abounds both in wet and dry places, in the upland and low grounds, affording excellent pasturage, and even when cut and dried it is eaten with avidity by horses, sheep, and cattle. A very similar congener inhabits the lofty peak of Tolima, in New Grenada, north of the Equator.

7. HIEROCHLOE, Gmel.

1. HIEROCHLOE Magellanica, Hook. fil. Torresia Magellanica, Pal. Beauv. Agrost. p. 63. Roem. et Schultes, Syst. Veg. vol. ii. p. 516. H. Antarctica, var. redolens, Brongn. in Duperrey, Voy. Bot. p. 144. t. 23. optime. Avena redolens, D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 601.

HAB. Strait of Magalhaens and throughout Fuegia and the Falkland Islands, very abundant, *Banks* and Solander, and all succeeding voyagers.

Under *H. redolens*, in the first part of this work, I have pointed out the very slight distinctions that separate this plant both from it and from the Tasmanian *H. Antarctica*, Br.: I consider them scarcely valid, though constant in specimens from the three widely separated localities they inhabit. In the Falkland Islands this grass is particularly abundant, forming large tufts and often beds, especially near running water and on wet rocks close to the sea, and is much frequented by sea-birds, as a building place. The scent is very strong, and retained in the dried specimens. Living plants introduced, by means of Ward's cases, into the Kew Gardens, have flourished luxuriantly, hitherto without flowering.

8. AIRA, L.

1. AIRA flexuosa, Linn., Sp. Pl. p. 96. Engl. Bot. t. 1519. Gaud. in Ann. Sc. Nat. vol. v. p. 100. et in Freyc. Voy. Bot. p. 100. D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 600.

HAB. Strait of Magalhaens; Port Famine and Port Gregory, Capt. King; Falkland Islands, Gaudichaud, and all succeeding voyagers.

An exceedingly abundant Falkland Island grass, and a great ornament to the black peat bogs, which are frequently clothed with its elegant purple panicles. The foliage is too scanty and of too rigid a texture to afford good pasturage.

I do not detect any difference between Falkland Island aud European specimens. Mr. Watson remarks that this is, perhaps, the *A. uliginosa*, Weihe; a plant I do not know, but quoted by Kunth as synonymous with *A. flexuosa*.

2. AIRA caryophyllea, Linn., Sp. Pl. 97. Engl. Bot. t. 812.

FLORA ANTARCTICA.

HAB. Falkland Islands, Mr. Wright. J. D. H.; probably introduced.

There are specimens of this species in the Hookerian Herbarium, marked as collected in the Falkland Islands by Admiral D'Urville, and others sent from Valdivia by Mr. Bridges. The plant is unquestionably the common European "Silver Hair Grass", and accidentally introduced, for, as Mr. Curtis remarks, "so insignificant an annual can hardly be worth cultivating."

3. AIRA *Kingii*, Hook. fil.; glaberrima, elata, panicula elongata effusa, ramis gracilibus subverticillatis, glumis lanceolatis acuminatis albidis nitidis vix puberulis flosculis pedicellatis ter longioribus, palea inferiore basi longe et dense sericeo-barbata apice truncata vix bifida irregulariter 4-dentata puberula obscure 3-nervi, nervo intermedio supra medium in aristam rectam palea paulo longiorem desinente superiore bifido, foliis lineari-elongatis culmo brevioribus vaginis profunde striatis duplo brevioribus. (TAB. CXXXV.)

HAB. Strait of Magalhaens; Port Famine, Capt. King; South part of Tierra del Fuego, C. Darwin. Esq.

Gramen elatum, 2–4-pedale, perenne, glaberrimum, nitens. Culmi cæspitosi, erecti, simplices, 2–3-nodosi, crassitie pennæ anatinæ, obscure striati, internodiis 4 unc. ad spithamæum et ultra. Foliorum radicalium vagina 4–5 unc. longa, lamina brevior, caulinorum internodiis brevior, profunde striata, fere ad basin hians; ligula oblonga, scariosa, alba; lamina angusta, herbacea v. subcoriacea, linearis, glaberrima, striata, marginibus siccitate involutis. Panicula 6–10 unc. longa, inclinata, effusa, ramis fasciculatis verticillatisve, gracillimis, divisis, inferioribus $\frac{3}{4}$ paniculæ æquantibus, glaberrimis, superioribus pedicellisque scaberulis. Spiculæ lineari-oblongæ, fere $\frac{1}{3}$ unc. longæ, albidæ, basi purpurascentes, scariosæ, nitidæ. Glumæ 1–nerves, angustæ, acuminatæ. Flosculi parvi, sub-longæ pedicellati, inclusi, glumis ter breviores, pedicello ciliato. Paleæ puberulæ, albidæ, micantes, scariosæ, subæquilongæ. Stamina sub-inclusa, antheris breviusculis. Squamulæ oblique lanccolato-ovatæ, acuminatæ. Ovarium compressum, obovato-oblongum, stylis basi discretis.

A very handsome grass, somewhat resembling the British A. caspilosa, but with very different locusta and florets.

PLATE CXXXV. Fig. 1, locustæ; fig. 2, floret; fig. 3, stamens and pistil; fig. 4, squamula:-all magnified.

4. AIRA *Magellanica*, Hook. fil.; puberula, panicula effusa pauciflora rachi ramisque elongatis gracilibus pubescenti-scaberulis, glumis ovato-lanccolatis acuminatis subæqualibus pubescentibus dorso scabridis flosculis stipitatis longioribus superiore basi 3-nervi, palca inferior late ovata basi sericco-barbata puberula 5-nervi, nervo intermedio infra apicem irregulariter 4-dentatum in aristam strictam glumis inclusam desinente, foliis planis latiusculis super pubescentibus. (TAB. CXXXIV.)

HAB. Strait of Magalhaens; Port Famine, Capt. King.

Species parvula, erecta, 6-8-uncialis. *Culmi* erecti, basi cæspitosi, simplices, foliati. *Foliorum vagina* teres, striata, hians, glabrata v. glaberrima; *ligula* brevis, ovata, obtusa; *lamina* vagina plerumque brevior, $1-l\frac{1}{2}$ unc. longa, flaccida, lineari-lanceolata, acuminata, plana, striata, super pilis sparsis puberula, subter glaberrima. *Panicula* exemplaribus verosimiliter immaturis basi vaginæ inclusa, gracilis, inclinata, effusa; ramis filiformibus, fasciculatis verticillatisve, divisis, flexuosis. *Glumæ* sub $\frac{1}{5}$ unc. longæ, subæquales, compressæ, ovato-lanceolatæ, acuminatæ, puberulæ, herbaceæ, virides et purpurascentes, opacæ. *Flosculi* glumis ter breviores, cum arista omnino inclusi. *Paleæ* subæquilongæ, inferior latiuscula. *Stamina* inclusa, antheris latiusculis. *Squamulæ* lineares, acuminatæ, *Ovarium* breviter stipitatum.

Capt. King's specimens of this pretty *Aira* are scarcely mature, though sufficiently developed for examination; the species ranks very near a Rocky Mountain one, chiefly differing in its smaller size, and in the pubescent upper surface of its leaves and glumes.

PLATE CXXXIV. Fig. 1, locusta; fig. 2, floret; fig. 3, stamens and pistil; fig. 4, squamula:-all magnified.

5. AIRA Antarctica, Hook.; panicula effusissima ramis fasciculatis capillaribus pedicellisque elongatis, spiculis lanceolatis 1-2-floris setulaque flosculi secundi tertiive auctis, flosculis puberulis pedicellatis basi sericeis, palea inferiore profunde bifida basi aristata, arista glumas superante, culmo brevi, foliis subulatis longe vaginantibus. A. Antarctica, *Hook. Ic. Plant.* t. 150. (TAB. CXXXIII.)

HAB. Hermite Island, Cape Horn, the Falkland Islands, and Kerguelen's Land, abundantly, J.D.H.; New South Shetlands, Dr. Eights.

Planta phænogamica ante omnia Antarctica. Culmi dense cæspitosi, breves, 1-3-unciales, erecti procumbentesve, foliosi. Folia glaberrima, herbacea, longe vaginantia; vagina $\frac{1}{2}-1\frac{1}{2}$ unc. longa, teres, striata; ligula linearis, $\frac{1}{4}$ unc. longa; lamina anguste lineari-subulata, marginibus involutis, vaginæ æquilonga v. longior. Panicula pro planta maxima, 4-6 unc. longa, effusa, 3-6 unc. lata, v. ob ramos appressos angustior; ramis 1-5 unc. longis pedicellisque scaberulis, capillaribus. Spiculæ angustæ, fere $\frac{1}{4}$ unc. longæ, 1-2 flores, uniflores semper biflores sæpissime pedicello ciliato floris alterius auctæ. Glumæ lanceolatæ, carina scabrida marginibus sub lente ciliatis, apicibus acutis, flosculis pedicellatis inclusis bis longiores. Palea inferior ciliata, oblongo-lanceolata, scarioso-membranacea, bifda v. fere bicuspidata, sinu quadrato bidentato, dorso basi aristata; arista reeta, scaberula, paulo ultra glumas exserta. Stamina exserta, antheris brevibus. Squamulæ oblique ovatæ acuminatæ. Ocarium breviter stipitatum, stylis basi discretis divaricatis.

This elegant grass, appropriately named A. Antarctica, attains a higher southern latitude than any other flowering plant, being the only phænogamic species that inhabits the South Shetland Islands. Kerguelen's Land in latitude 48° is its northern limit; but that Island being situated in a longitude where the rigour of the Antarctic climate extends further north than in any other, this grass is even there more typical of the frigid zone than the latitude would indicate, and always seeks the most sheltered places. In the Falkland Islands again, the most temperate region it inhabits, it invariably avoids shelter, being found chiefly in open marshy places near the sea, fully exposed to the violence of the winds.

PLATE CXXXIII. Fig. 1, two locustæ and portion of panicle; fig. 2, a floret from the same; fig. 3, squamula; fig. 4, single-flowered locusta; fig. 5, floret from the same :---all magnified.

6. AIRA *parvula*, Hook. fil.; cæspitosa, puberula v. glabrata, panicula erecta contracta subsimplici pauciflora ramis brevibus locustisque erectis, glumis lanceolatis acuminatis flosculis pedicellatis triplo longioribus, palea inferiore late ovata basi barbata apice bifida inter segmentos acutos bidentata dorso supra basin aristata, arista geniculata glumas vix excedente, foliis setaceis culmo brevioribus.

HAB. Hermite Island, Cape Horn; rocks near the mountain tops, J. D. H.

Gramen 3-5-unciale, foliosum, dense cæspitosum, rigidiusculum. Culmi erecti, basi fibrosi et pluries divisi, foliis perplurimis vaginati. Folia 2 unc. longa, stricta, erecta, anguste subulata, glabrata v. pilis patulis puberula, subcoriacea, marginibus involutis; ragina latiuscula, membranacea; ligula valde elongata, scariosa, linearis, acuminata. Panicula $1-1\frac{1}{2}$ -uncialis, stricta, erecta; ramis paucis, brevibus, 1-floris, paniculæ appressis. Locustæ $\frac{1}{3}$ unc. longæ. Glumæ æquales, glabriusculæ. Flosculorum pedicelli sericeo-barbati; flosculi superioris palea superior setula aucta. Squamulæ ovatæ, acuminatæ. Ovarium obtusum, stylis discretis, lateralibus.

A remarkably distinct little species, most nearly allied to *A. Antarctica*, but distinct in the foliage, the very different panicle, and shorter florets.

9. TRISETUM, Kunth.

1. TRISETUM subspicatum, Beauv., Agrost. p. 88. Fl. Antarct. Pt. 1. p. 97. T. andinum, Benth. Plant. Hartweg. p. 261. n. 1449.

4 q

FLORA ANTARCTICA.

HAB. Strait of Magalhaens; Port Famine, Capt. King; Hermite Island, on sandy beaches near the sea, J. D. H. Falkland Islands, most abundant, D'Urville, J. D. H.

In the first part of this work I have given the geographical range of the *Trisetum subspicatum*, when noticing it as a native of Campbell's Island; at which time I was not aware of any other South American station for it than the Andes of Peru. Since then I have seen several specimens collected both in the Cordillera of Columbia and in Mexico, whence it is evident that this plant, like many common to the opposite temperate zones, has availed itself of the direct communication afforded by the Andes of the American continent for migrating from the Northern to the Southern Hemisphere. Its great abundance in the New World and especially in the extreme South of America, coupled with its rarity in the southern regions of the Old World, where it is only known on the tops of the mountains of Campbell's Island, seem to indicate its having been transmitted from east to west, or against the course of the prevailing winds in the Antarctic regions.

10. AVENA, L.

1. Avena leptostachys, Hook. fil.; glaberrima, nitida, panicula gracillima flexuosa nutante ramis breviusculis subverticillatis capillaribus paucifloris, glumis inæqualibus inferiore flosculo $\frac{1}{2}$ breviore, palea inferiore lanceolata basi barbata bicuspidata inter segmenta aristata, arista gracili reflexa locusta bis longiore, culmis gracilibus, foliis planis elongatis.

HAB. Strait of Magalhaens; Port Famine, Capt. King.

Culmus exemplare incompleto pedalis, gracilis, erectus, debilis, foliis vaginatus, nitens. Folia caulina longe vaginantia; vagina teres, striata, 5-unc. longa; ligula membranacea, ovata, fimbriata; lamina 6-S-pollicaris, $\frac{1}{3}$ unc. lata, flaccida, membranacea, striata. Panicula 6 unc. longa; ramis capillaribus, $\frac{1}{2}$ -1-uncialibus, glaberrimis. Locusta $\frac{1}{4}$ unc. longæ, biflores; flosculis pedicellatis; superiore longius pedicellato, setula ciliata aucto. Glumæ ovatolanceolatæ, acuminatæ, glaberrimæ; inferiore $\frac{1}{3}$ -minore, 1-nervi; superiore 3-nervi. Palea inferior lanceolata, puberula, 1-nervis, nervo dorso scaberulo; superior brevior, apice bicuspidata. Squamulæ 2, oblongæ, laceræ. Ovarium obovatum, breviter stipitatum, apice barbatum; stybis lateralibus, basi paulo discretis.

A very elegant species, of which I regret having seen but one colm, which wants the rooting portion. It is nearly allied to the United States *Avena palustris*, Mich.; from which it may readily be distinguished by the smaller locustæ, more exserted florets, and unequal glumes.

11. POA, L.

1. PoA scaberula, Hook. fil.; erecta, gracilis, scabrida, panicula subsecunda coarctata densiflora, glumis 3-floris subæqualibus 1-nerviis puberulis dorso scabridis, flosculis pubescentibus basi lanatis breviter pedicellatis, palea inferiore subcarinata 3-nervi, nervis lateralibus tenuissimis inconspicuis, carina dorso superne scabrida inferne ciliato-plumosa, superiore $\frac{1}{3}$ breviore apice 2-dentata, foliis lineari-setaceis scaberulis eulmo gracili erecto scabrido multotics brevioribus.

HAB. Strait of Magalhaens; Port Famine, Capt. King.

Radix fibrosa. Culmi erecti v. basi ascendentes, pedales et ultra, graciles, striati, scaberuli. Folia pauca, longe vaginantia; vagina scaberula, profunde striata; lamina 3-5-uncialis, setacea, involuta; ligula ovata, obtusa, membranacea. Panicula 2-3-pollicaris, coaretata, basi interrupta, unilateraliter secunda, $\frac{1}{2} - \frac{3}{4}$ unc. lata. Locustæ parvæ, $\frac{1}{8}$ unc. longæ, puberulæ, purpurco-pictæ, late ovatæ, sub 3-flores. Glumæ virescentes, compressæ, locusta $\frac{1}{3}$ breviores, acutæ. Flosculi basi longe arachnoidco-lanati. Palea superior acuta, membranaceo-marginata. Squamulæ parvæ, ovatæ, acuminatæ.

I know of no species with which the present can be confounded. The scabridity, coarctate paniele, dense locustæ, and other characters at once distinguish it from its congeners.

2. PoA nemoralis, Linn., Sp. Pl. 102. Engl. Bot. t. 1265.

HAB. Strait of Magalhaens; Port Gregory, Capt. King.

Most distinctly the *P. nemoralis* of Ben-Lawers, whose flowers are slightly webbed at the base. It is also a Rocky-Mountain plant, but has not hitherto been found on the Cordillera of South America.

3. Pos pratensis, Linn., Sp. Pl. 99. Banks et Sol. in Mus. Banks. Engl. Bot. t. 1073. P. compressa, var. viresceus, D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 600. P. alpina, Brongn. in Duperrey, Voy. Bot. p. 44, non Linn.

Var. 1, ligula folii superioris oblonga, panicula laxa, glumis 4-floris paleisque angustioribus.

Var. 2, panicula subcoarctata, glumis latioribus brevioribusque 2-3-floris.

Var. 3, panicula effusa, glumis 3-5-floris.

Var. 4, panicula coarctata, glumis sub 4-floris paleisque angustioribus, flosculis basi fere nudis.

Var. 5, 6-uncialis, locustis minoribus 2-3-floris.

HAB. Strait of Magalhaens; Port Famine, Capt. King (vars. 1, 3, and 4), Good Success Bay, Banks and Solander; Falkland Islands, abundant (vars. 2 and 5).

I cannot ascertain the identity of this species with the European *P. pratensis*, so satisfactorily as that of the former with *P. nemoralis*; nevertheless, the more the present grass is studied, the more difficult it appears to detect specific characters. The five varieties enumerated, appear all to belong to one plant; except, perhaps, the *var.* 1, in which the ligula of the upper leaf is oblong as in the European *P. alpina*, to a North American state of which I should have referred that variety, had its florets not been webbed, a character, which, though of trifling importance, (perhaps even less than the form of the ligula) does not exist in any of the numerous individuals of *P. alpina* that I have examined.

In British, and, indeed, in European examples of P. alpina, we are accustomed to see a small panicle of short and broad flowers, with a different aspect to that of P. pratensis; but North American individuals are subject to great variations in the size and outline of the panicle, so great that without connecting forms it would be very difficult to recognize them. Mr. Watson is equally persuaded with myself of the close affinity between this Antarctic Poa and P. alpina, though neither of us can adduce a tangible character beyond the webbed florets to separate the plants of Arctic and Antarctic America. I have not seen any of these species from the intervening Cordillera, a circumstance of little importance, the Gramineæ of these regions having been very much neglected by all collectors since the period of the travels of the illustrious Humboldt.

12. TRIODIA, Br.

1. TRIODIA Kerguelensis, Hook. fil.; parvula, dense cæspitosa, panicula simplici pauciflora scaberula, glumis inæqualibus acutis 3-nerviis 2-floris, flosculis breviter pedicellatis glumis inclusis basi nudis, palcis æquilongis inferiore dorso convexa obscure 5-nervi, superiore æquilonga bifida, foliis setaceis, culmis brevibus basi foliosis. (TAB. CXXXVIII. sub nomine Poæ).

HAB. Kerguelen's Land; rocky places, at an elevation of 300-1200 feet.

Gramen parvulum, dense cæspitosum, 2-4-unciale. Folia glaberrima, brevia, recta sed vix rigida, setacea, marginibus involutis; lamina vix pollicaris vagina tumida longior; ligula ovata, subacuta. Panicula seu racemus simplex, 3-5 lin. longus, erectus; rachi flexnosa, scaberula. Locustæ pedunculatæ, $1\frac{1}{2}$ lin. longæ, ovato-oblongæ, virides. Glumæ margine dorso nervisque scaberulæ, concavæ, coriaceo-chartaceæ; superior longior, paulo angustior; inferior oblique acuminata. Flosculi glumas vix superantes, inferior subsessilis, superior breviter pedieellatus. Paleæ æquilongæ; inferior 3-nervis, dorso basi serieeo, acuto v. obscure et oblique truncato; superior bicarinata, apice bifida. Antheræ parvæ, late oblongæ. Caryopsis ovoidea, glaberrima.

I was long doubtful whether to refer this curious little species to *Poa*, *Festuca*, or *Triodia*, to all which genera (like some other grasses) it has nearly equal affinity; to *Poa* in the form of the locustæ and florets, and to *Festuca* in the acute glumes and palea; hut eertainly most to *Triodia*, in habit, form of panicle, included florets and obscurely 3-dentate lower palea.

PLATE CXXXVIII. Fig. 1, portion of culm with vagina, base of lamina of leaf, and ligula; fig. 2, locusta; fig. 3, glume; fig. 4, floret; fig. 5, do with ripe caryopsis; fig. 6, caryopsis: ---all magnified.

2. TRIODIA Antarctica, Hook. fil.; parvula, dense eæspitosa, glaberrima, panicula subsimplici coaretata, locustis breviter pedunculatis, glumis subæqualibus lanceolatis 3-floris, flosculis basi nudis paleis subæquantibus inferiore 5-nervi acuta obscure 3-dentata folis basi longe membranaceis vaginantibus culmum fere æquantibus, lamina setacea. Festuca pusilla, Banks et Sol. in Mus. Banks.

HAB. Tierra del Fuego; C. Darwin, Esq. Rocks near the tops of the mountains of Hermite Island. J. D. H.

Culmi dense fastigiati, basi inclinati, valde foliosi, 4-pollicares. Folia plurima, erecta, substriata sed non rigida; vagina elongata, striata; lamina pollicaris, setacea, marginibus involutis; ligula ovata, aeuminata. Panicula fere uneialis, simplex v. basi ramosa, erecta. Locustæ parvæ, 3-flores, glabriusculæ. Glumæ subæquales, flosculis breviores, lanceolatæ, concavæ, 3-nerves. Flosculi 2 superiores pedicellati, basi omnino nudi. Palea inferior late ovata, concava, apice breviter truneata et tridentata, dente intermedio paululum elongato, 5-nervis; nervis dorso obscure scaberulis; superior æquilonga, biearinata, apice bifida. Antheræ parvæ, late oblongæ.

A peculiar species, allied to the last and to no other with which I am acquainted. The obliquely truncated apex of the lower palea in the *T. Kerguelensis*, is here, as it were, exaggerated by that organ becoming decidedly though minutely trifid at the apex, as in the European *Triodia decumbens*, a genus to which both species ought from this circumstance be referred, and from their peculiar panicle and locustæ.

In habit the similarity between this plant and the former is very great, and apparent in the size, foliage, and locality they both affect, in their respective Islands; the differences in the details of the florets, are, on the other hand, sufficiently wide.

13. FESTUCA, L.

1. FESTUCA Fuegiana, Hook. fil.; erecta, elata, culmis basi praccipue foliosis scaberulis glaberrimisve, panicula effusa v. subcoarctata, glumis ovato-lanceolatis acuminatis subcarinatis, flosculis breviter pedicellatis basi araneosis, superioribus viviparis, palea inferiore acuminata puberula 5-nervi nervis dorso sericeis superiore æquilonga bifida, foliis breviusculis subacutis marginibus involutis, ligula oblonga. (TAB. CXLI.)

Var. a, panicula contracta, culmo superne præcipue scabrido. Aira cæspitosa. Banks et Sol. in Mus. Banks. (in part).

Var. β. panicula effusa, magis vivipara, culmis glaberrimis. Poa alpina, var. vivipara, Banks et Sol. in Mus. Banks.

HAB. Strait of Magalhaens, Port Famine and Port Gregory, Capt. King. South part of Fuegia, C. Darwin, Esq.

Gramen erectum, 1- ad 2-pedale. Culmi dense fastigiati, basi valde foliosi, superne glaberrimi v. scabridi. Folia breviuscula, 3-4-uncialia, glaberrima, substricta sed non rigida, late linearia, acuta, marginibus involutis,

381

vaginis striatis breviora; *ligula* late elongata, oblonga, apice fimbriata. *Panicula* 3-5 nnc. longa, contracta v. effusa, ramis scaberulis. *Locustæ* 4-5 lin. longæ, viviparæ, pollieares et ultra. *Glumæ* chartaceæ, æquales, acuminatæ, superiore 3-nervi, subcarinata, carina scaberula. *Flosculi* sub 5, basi appresse araneosi, lana albida. *Palea* inferior flosculorum snperiorum sæpissime in folium apice uncinatum lignla et vagina 5-nervi instructum desinens. *Antheræ* lineares. *Ovarium* late obovatum, supra basin contractum, basi squamulis acinaciformibus instructum. *Styli* breviusculi, ad basin plumosi.

A very handsome grass, which, perhaps, properly belongs to *Poa*, though the paleæ are so decidedly acuminate that I prefer retaining it under *Festuca*. The two varieties enumerated are not always constant to the characters assigned to them.

In general appearance this species resembles the British *Aira caspitosa*, which is frequently similarly viviparous on the mountains, and the modifications the paleæ consequently undergo both in these and some other grasses, is a subject well worthy of study. When the inflorescence becomes foliaceous, the palea itself, which is distinctly 5-nerved is represented by the (equally 5-nerved) vagina of the leaf; the ligula of the latter holds the position of the membranous and often divided apex of such a palea as that of Aira, whilst the lamina answers to a dorsal awn; or rather, in the case of Festuca Fuegiana, to five awns (such as those of Polypogon Chonoticus, p. 374), united by parenchyma. That the arista of the lower palea in grasses is the produced mid-rib of a modified leaf, is perhaps generally admitted, but the exact relation of the apex of the palea to a ligula is not so evident in all aristate florets, as it is in those where the middle nerve is not percurrent but separates from the palea in the form of an awn. One apparent objection to this view may be adduced in the distinctly articulate awn of Corynephorus and Stipa, which may further be supposed favourable to M. Raspail's theory, that the mid-rib of the palca is an axis of development in cohesion with the bracts; such articulations are, however, exceptional, and their position 1 am inclined to consider as indicating the point of union of the leaf with the vagina, where an angle is always observable. Viviparous grasses, too, would be expected to produce constantly additional organs from the portion of the transformed palea beyond the ligula, if M. Raspail's view were correct, but, this, on the contrary, is seldom the case. There is a similarity between the palea of a viviparous grass and the upper bract of each spikelet in some Marisci: for in them the dilated lower portion of the braet, or the true continuation of the rachis, somewhat resembles, without however being strictly analogous to, the lower palea of a locusta, and the uppermost flower is borne in a position, similar to the axle of the ligula on the leaf of a grass.

PLATE CXLI. Fig. 1, locusta; fig. 2, floret; fig. 3, ovary; fig. 4, squamula; fig. 5 and 6, viviparous portions of a spikelet; fig. 7, palea transformed into a leaf:—all magnified.

2. FESTUCA Arundo, Hook. fil. F. Alopecurus, D'Urville in Mém. Soc. Linn. Paris, vol. iv. p. 604. Brong. in Duperrey Voy. Bot. p. 32. Poa (?) Alopecurus, Kunth, En. Plant. vol. i. p. 256. Arundo Alopecurus, Gaud. in Ann. Sc. Nat. vol. v. p. 100., et in Freyc. Voy. Bot. p. 409.

Var. β . minor, foliis angustioribus culmo brevioribus.

Var. γ . pedalis, glumis et paleis latioribus brevioribusque.

Var. 8. culmo graciliore, panicula sub-nutante, flosculis-sæpius basi parce lanatis.

HAB. Falkland Islands, all the varieties forming very large tufts; on the sea-sand abundant; D'Urville, J.D.H. Var. γ . Strait of Magalhaens; Port Gregory, Capt. King.

Next to the Tussock, the present is the largest grass in the Falkland Islands, though, like that plant, it is very variable in size. The largest specimens are three or even four feet high, the smaller scarcely one. Though a conspicuous object, its varieties are not always easily recognizable; for the most prominent characters of the typical state, which are the great size of the locustæ, and the narrow paleæ and glumes with slender attenuated apices, are quite fallacious. All my large specimens of var. a have either a minute turbinate ovarium or a small

caryopsis, and are never staminiferous; thus it is very possible that some of the varieties enumerated may be the males of this, the largest form.

M. Brongniart has suggested the propriety of erecting the present plaut, together with the *F. Antarctica*, into a new genns, and they certainly are more nearly allied to one another than to any of their congeners; still I doubt the possibility of finding any character of generic value common to them both. They also resemble some South Brazilian and Patagonian grasses, as the *Poa lanuginosa*, Nees, and other undescribed species.

If I had seen only single specimens of the different varieties, I should certainly have considered three of them to be as many species; but a very large collection of individuals, from various parts of the Island, has convinced me, that neither the comparative length, breadth, or attenuation of the apices of the glumes and paleæ, nor the woolliness of the base of the florets, or length of the leaves, afford any grounds for a further subdivision; at least I have been unable to effect such, either when examining the fresh specimens, or, more lately, when comparing the dried ones. Dissimilar as the following plant appears, I am not at all positive of its claims to the rank of a separate species; for some of its characters may be due to the different locality it generally affects; and specimens of the var. δ . approaching the *F. Arundo* far too nearly.

Though a large and very handsome grass, the *Festuca Arundo* is so harsh and rigid as to be quite unpalateable to cattle; this is the more obvious from its often growing side by side with the untritious Tussock, out of the same sand-heap.

3. FESTUCA Antarctica, Kunth, Gram. vol. i. p. 132. En. Plant. vol. 1. p. 408. Arundo Antarctica, D'Urv. in Mém. Soc. Linn. Paris. vol. iv. p. 602.

Var. a, culmo pedali, foliis strictis rigidis, panicula erecta, flosculis basi fasciculis pilorum instructis. Arundo Antarctica, Brong. l. c.

Var. β . culmo pedali et ultra, foliis elongatis flexuosis, panicula nutante, fasciculis pilorum rarissimis.

Var. γ . omnia varietatis β ., sed flosculis omnino nudis.

Var. δ . habitu varietatis α . flosculisque varietatis γ .

HAB. Falkland Islands, most abundant; vars. a. and δ . on sandy shores; vars. β . and γ . in rocky places, both near the sea and upon the hills, sometimes also on the sandy shores.

Few botanists would, I think, venture to separate any of the varieties enumerated above from *F. Antarctica*, and very many others would unite all with the preceding species, and perhaps correctly. M. Kunth describes a specimen of this grass (received from D'Urville) as having the flosculi naked at the base, exactly as in my varieties γ . and δ . (*En. Plant. Suppl.* p. 340). The locustæ vary in my specimens, being from two- to four-flowered.

4. FESTUCA archaria, Lamk., Encycl. vol. i. p. 191. D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 602. Brongniart, in Duperrey Voy. Bot. p. 35. Kunth, En. Plant. vol. i. p. 408.

HAB. Strait of Magalhaens, Commerson; Port Famine, Capt. King; Falkland Islands, marshy and sandy places, Gaudichaud, Sc.

The lower palete of this species are frequently notched on each side, below the apex, as in a genuine *Dactylis*, and in the following plant. Fuegian specimens are often viviparous.

5. FESTUCA *Cookii*, Hook. fil.; panicula elougata erecta contracta fastigiatim v. verticillatim ramosa, glumis subæqualibus ovato-lanceolatis acuminatis glaberrimis 4-floris superiore 3-nervi, flosculis basi uudiusculis palea inferiore puberula 5-nervi dorso basi sericeo-barbata, apice acuminata integra v. 3-dentata, culmo diviso folioso basi radicante, foliis distichis culmum superantibus. (TAB. CXXXIX.)

HAB. Kerguelen's Land, abundant; Anderson (in Cook's Voyage), J. D. H.

Gramen foliosum, 3 unc. ad bipedale. Culmi robusti, basi pluries divisi, prostrati v. repentes; pars repens validus, sæpe pedalis; pars erectus per totam longitudinem foliosus, compressus. Folia plurima, distiche inserta, elongata, coriacea sed non rigida; vagina aperta, folio brevior, compressa, striata; ligula brevis, fimbriata; lamina culmum paniculamque superans, plana v. dorso subcarinata, utrinque lævis, 2-4-lin. lata, gradatim in apicem acuminatam angustata. Panicula 2 ad 8 unc. longa, erecta v. paulo inclinata, $\frac{3}{4}$ unc. lata, subcontinna v. verticillatim interrupta. Locustæ late ovato-oblongæ, 3-4-flores, sub 8-lin. longæ. Gluma superior 3-nervis, inferiore 1-nervi longior, flosculis breviter pedicellatis brevior. Palea inferior ovato-lanceolata, acuminata, concava, vix carinata; nervo medio dorso superne ciliato, inferne barbato; flosculo superiore infimoque apice acuminato, integerrimo, duobus intermediis apiccm versus utrinque uni-dentatis; palea superior inferiore $\frac{1}{3}$ brevior, bifida. Antheræ lineares. Caryopsis oblonga, cylindracea.

The commonest grass in Kerguelen's Land and a very valuable one, affording a rich and abundant fodder. The tendency in the palea to become toothed on each side towards the apex, and the distichous, long, and particularly rich foliage, show its affinity with the Tussock and with the *Festuca foliosa* of Lord Auckland's group, which chiefly differs from this in its larger panicle. These three grasses are certainly representatives of one another, and all typical of moist Insular climates; their northern analogues are evidently the *F. Donax*, Lowe, of Madeira, and *F. albida*, Lowe, of the same island.

PLATE CXXXIX. Fig. 1, locusta; fig. 2, floret; fig. 3, squamula; fig. 4, caryopsis :---all magnified.

§ 2. Flosculis arista terminatis.

6. FESTUCA *purpurascens*, Banks et Sol. MSS.; elata, panicula laxa ramis elongatis apicibus paucifloris, locustis oblongis multi- 8-floris, glumis trinerviis lanceolatis superiore ter majore, flosculis glabriusculis, paleis 5-nerviis apice 3-dentatis dente intermedio in aristam producto, foliis planis culmo brevioribus. (TAB. CXL.)

HAB. Strait of Magalhaens; Port Famine, Capt. King; Fuegia; Good Success Bay, Banks and Solander, C. Darwin, Esq.

Radix stolonifera. Culmi 2-4-pedales, graciles, erecti, glaberrimi, striati, politi, remote nodosi. Folia pauca, culmum vaginantia, patentia, flexuosa; vaginæ pedales, teretes, superne hiantes; ligula brevis, transversa; lamina plana, utrinque lævis, vagina brevior longiorve, gradatim supra medium acuminata. Panicula 6 unc. longa, laxa, inclinata; ramis paucis, elongatis, filiformibus, versus apices divisis. Locuslæ fere $\frac{1}{2}$ -unc. longæ. Glumæ flosculis pedicellatis basi nudis breviores. Patea inferior dorso convexa, superne præcipue pilosiuscula, superiorem bifidam paulo breviorem amplectans, nitida, purpureo-picta, 5-nervis, nervis scaberulis. Squamulæ 2, profunde bifidæ. Ovarium obovatum, superne pilosum; styli basi discreti.

A tall and handsome grass, allied to the South Brazilian *F. fimbriata*, Nees, in which the paleæ are not awned, and the leaves are convolute. In general habit it resembles somewhat the European *F. elatior*.

PLATE CXL. Fig. 1, locusta; fig. 2, floret; fig. 3, pistil and squamulæ; fig. 4, squamulæ:-all magnified.

7. FESTUCA duriuscula, Linn., Sp. Pl. 108. Engl. Bot. t. 470.

HAB. Strait of Magalhaens; Port Famine, Capt. King.

Capt. King's specimens are nearly two feet high, in which respect only they differ from ordinary forms of those of British growth. The leaves are erect and involute.

S. FESTUCA gracillima, Hook. fil.; elata, glaberrima, panicula simplici elongata pauciflora inclinata, locustis majusculis pedunculis compressis longioribus multi-7–9-floris, glumis inæqualibus lineari-oblongis

late scarioso-marginatis superiore latiore 3-nervi, flosculis basi remotiusculis, palea inferiore obscure puberula in aristam brevem producta, culmis gracillimis folium lineari-filiforme involutum superantibus.

HAB. Strait of Magalhaens; Port Famine, Capt. King.

Radix fibrosa, nunc repens? Culmi 3-pedales, gracillimi, erecti, læves, nitidi. Folia $1-1\frac{1}{2}$ -pedalia. Panicula sub 5-unc. longa, 6-8-fiora. Locustæ $\frac{1}{2}$ ad $\frac{3}{4}$ pollicares. Glumæ latiusculæ, concavæ, non carinatæ, inferior 1-nervis, superior latior, 3-nervis. Paleæ lineares, inferior sursum puberula, arista breviuscula recta auctæ; superior æquilonga, bifida.—Species elongata, gracillima, priori simillima, sed elatior, foliis longioribus, locustis majoribus, glumis plurifloris latioribusque differt.

A very elegant species, allied to the former; but, judging from my specimens, distinct, especially in the form of its glumes; although in British examples of F. rubra that organ varies much in breadth and the locustæ in sizc.

9. FESTUCA bromoides, Linn., Sp. Pl. 109. Engl. Bot. t. 1412. D'Urville, in Mém. Soc. Linn. Paris, vol. iv. p. 601.

HAB. Falkland Islands, found only near the settlement, D'Urville, J. D. H.

Apparently identical with the European plant, and most probably introduced.

10. FESTUCA Magellanica, Lamk., Illust. vol. i. p. 119. Encycl. vol. ii. p. 461. D'Urville, in Mém. Soc. Linn. Paris, vol. iv. p. 601. Brong. in Duperrey, Voy. Bot. p. 38. Kunth, En. Plant. vol. i. p. 396.

Var. β . culmo elongato, foliis glaberrimis.

HAB. Strait of Magalhaens, *Commerson*; Falkland Islands, on rocks near the sea, D'Urville, J. D. H.Var. β . Port Famine, *Capt. King*.

The var. β ., from Port Famine, is almost identical with Austrian specimens of *F. pallens*, Host., and it comes very near some British states of *F. duriuscula*, apparently differing chiefly by the membranous margins of the sheaths of the leaves. Falkland Island specimens vary considerably in stature and in the size of their locustæ; the foliage is very rigid in all, though more or less publicent in different specimens.

11. FESTUCA erecta, D'Urville, in Mém. Soc. Linn. Paris, vol. iv. p. 601. Brongniart, in Duperrey Voy. Bot. p. 37. t. 7. Kunth, En. Plant. vol. i. p. 398.

HAB. Tierra del Fuego; Hermite Island, J. D. H.; Falkland Islands, D'Urville, J. D. H.; Kerguelen's Land, R. M'Cormick, Esq.

Variable in the comparative length of the leaves and stem, as also in size, but otherwise a well-marked species.

14. DACTYLIS, L.

1. DACTYLIS cæspitosa, Forst., in Comm. Goctt. vol. is. p. 22. Willd. Sp. Pl. vol. i. p. 407. Hook. fil. in Lond. Journ. of Bot. vol. ii. p. 298. t. 9 and 10. Festuca cæspitosa, Roem. et Sch. Syst. Veg. vol. ii. p. 732. Kunth, En. Plant. vol. i. p. 408. F. flabellata, Lamk. Encycl. vol. ii. p. 462. Gaud. in Ann. Sc. Nat. vol. v. p. 100, et in Freyc. Voy. Bot. p. 409. D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 603. Brongniart in Duperrey Voy. Bot. p. 36. "Glayeux," Pernetty, Voy. vol. i. p. 343. (TAB. CXXXVI.—CXXXVII.)

HAB. Strait of Magalhaens, *Commerson*; and throughout Fuegia; Staten Land, *Forster*; Hermite Island, Cape Horn, J. D. H.; Falkland Islands, most abundant, *Gaudichaud*, and all subsequent voyagers.

Though much has lately been written in the 'Journal of Botany' upon this plant, the famous Tussoek Grass of the Falkland Islands, it appears advisable to sum up here the principal facts connected with its history.

Commerson was doubtless the discoverer of it in the Strait of Magalhaens, in 1767, and it has been gathered in Fuegia by several suecceding yoyagers; but as it nowhere forms so conspicuous a feature as in the Falkland Islands, it is most appropriately considered in reference to them alone.

A French colony was established on the Falklands, by Admiral Bougainville, in 1766, when cattle and horses were landed, which, no doubt, soon manifested a predilection for this noble grass. Pernetty, the historiographer of the Voyage, in describing the remarkable plants of those Islands, alludes particularly to it under the name of "Glayeul"; but it was not until the recent colonization of the Falklands by the British that attention was particularly directed to the Tussock, in consequence of accounts forwarded to the Colonial Office by Governor Moody, and to the Admiralty by the Antarctic Expedition.

The peculiar mode of growth of *Dactylis caspitosa* enables it to thrive in pure sand, and near the sea, where it has the benefit of an atmosphere loaded with moisture, of soil enriched by decaying sca-weeds, of manure, which is composed in the Falkland Islands of an abundant supply of animal matter in the form of Guano, and of the excrements of various birds, who deposit their eggs, rear their young, and find a habitation amongst the groves of Tussoek. Its general locality is on the edges of those peat-bogs which approach the shore, when it contributes eonsiderably to the formation of peat. Though not universal along the coast of these Islands, the quantity is still prodigious, for it is always a gregarious grass, extending in patches sometimes for nearly a mile, but seldom seen except within the influence of the sea air. This predilection for the ocean does not arise from an ineapaeity to grow and thrive except close to the salt water, but because other plants, not suited to the sea-shore, already cover the ground in more inland localities, and prevail over it : I have seen the Tussoek on inaccessible cliffs in the interior, having been brought there by the birds and afterwards manured by them; and, when cultivated, it thrives both in the Falklands and in England, far from the sea.

I know of no grass likely to yield nearly so great an amount of nourishment as the Tussock, when thoroughly established; in proof of which I quote my friend Governor Moody's printed report, for the truth of which I can vouch, both from my own experience and from his having kindly given me ample means for judging of the correctness of his interesting and useful observations, when drawing up the report from which the following extract is made.

"During several long rides into the country I have always found the Tussock flourishing most vigorously in spots exposed to the sea, and on soil unfit for any other plant, viz. the rankest peat-bog, black or red. It is wonderful to observe the beaten foot-paths of the wild cattle and horses, marked like a foot-track aeross fields in England, extending for miles over barren moor-land, but always terminating in some point or peninsula covered with this favourite fodder; amid which, one is almost certain to meet with solitary old bulls, or perhaps a herd of cattle; very likely, a troop of wild horses, just trotting off as they scent the coming stranger from afar. To cultivate the Tussock grass I should recommend that its seeds be sown in patches, just below the surface of the earth and at distances of about two feet apart; it must afterwards be weeded out, for it grows very luxuriantly, frequently attaining a height of six or seven feet. It should not be grazed, but cut or reaped in bundles. If cut, it quickly shoots again; but is much injured by grazing; for all animals, especially pigs, tear it up to get at the sweet nutty-flavoured roots. I have not tried how it would be relished if made into hay, but eattle will eat the dry thatch off the roof of a house in winter ; their preference to Tussock grass being so great that they seent it a considerable distance and use every effort to get at it. Some bundles, which had been staeked in the yard at the back of Government House, were quickly detected, and the eattle in the village made, every night, repeated attempts to reach them, which occasioned great trouble to the sentry on duty."

Since the above was written, the Tussoek has been used abundantly when made into hay, being preferred by cattle even to the green state of any of the other excellent grasses in the Falklands. Governor Moody informs me that in his garden it grows rapidly and improves by eutting.

385

There is, however, one draw-back to the value of the Tussock; it is a perennial grass, of slow growth, and some disappointment has already been experienced in England from this cause. Each Tussock consists of many hundreds of culms, springing together from a mass of roots, which have required a long series of years to attain their great and productive size. Our cultivated specimens in the Royal Gardens of Kew, now nearly three years old, are in a fair way of becoming good Tussocks; for the quantity of stems from each root, the produce of one seed, is incalculably more than any other grass throws np, and these are already forming a ball of root-fibres which in time will form a mound; but this ball, now scarcely six inches across and not two in height, must have grown to six or eight feet high, with a diameter of three or four feet; instead of forty culms there must be four hundred; and the leaves, now three feet long, must attain seven; ere the Tussock of England can compete with its parent in the Falklands. Though, however, the stoles (if 1 may so call the matted roots of this grass) in the most vigorons native specimens attain a height of seven feet, it is certain that they are very productive before they have reached two or three. By the time the leaves have gained their great size, the bases of the culms are nearly as broad as the thumb, and when pulled out young, they yield an inch or two of a soft, white, and sweet substance, of the flavour of a nut, and so nutritions, that two American sealers, who deserted a vessel in an unfrequented part of the Falklands, subsisted on little else for fourteen months.

Again, the Tussock-grass field, when fully established, must not be grazed indiscriminately by cattle. These creatures and the pigs have already diminished its abundance in the Falklands; for, after devouring the foliage, they cat down the stumps of the cuhns, greedily following them into the heart of the muss of roots from which they spring, for the sake of the white core just described; the rain-water lodges in the cavity thus formed, and decay so surely follows, that I have seen nearly half a mile of Tussock-grass plants entirely destroyed by no other means.

Although in the Falklands this plant will grow on pure sand near the sea, and there reach as great a size as on any other soil, it is not likely to do so in the drier climate of Britain, where the absence of an equally humid atmosphere must be artificially remedied. A wet, light, peaty soil has in England been found to favour its growth; sca-weed manure might probably be added with advantage, and certainly guano. Slow its progress assuredly is, but it may be hastened by such stimulants. In the mean time the cultivator has no just cause for complaint; the plant is already increasing unusually at the base, and thence sending up many more culms than other grasses, though, springing from one small base, they do not make such a show, but form a compact mass of living roots which in the case of other Gramineæ would spread over ten times the area that this occupies, and they annually increase in vigour and productiveness. And, lastly, it must be borne in mind that the farmer here obtains an enormous crop from a very small surface. Each great Tussock is the produce of one seed and is an isolated individual plant, which, though standing upon perhaps only two square yards of ground, yields annually a produce equal to that of a much greater surface of land, if cropped with hay or clover. The number of seeds required to stock an acre in Tussock and one in grass is in the proportion of tens to thousands ; and we may be well content to know that the number of months required to ensure a profitable return is not in the same ratio.

There are few plants which from perfect obscurity have become objects of such interest as this grass. The Tussock in its native state scens of almost no service in the animal economy. A little insect, and only one that I observed, depends on it for sustenance; and a bird, no bigger than the sparrow, robs it of its seeds; a few seafowl build amongst the shelter of its leaves : penguins and petrel seek hiding-places amongst the roots, because they are soft and easily penetrated, and Sea-lions cower beneath its huxuriant foliage : still, except the insect, I know no animal or plant whose extinction could follow the absence of this, the largest vegetable production in the Falklands, which does not even support a parasitical fungus. These same sea-birds breed and burrow where no Tussock grows; rocks elsewhere suit the Sea-lion's habits equally well; and the sparrow, which subsists on other food eleven months of the year, could surely make shift without this for a twelfth. Certain it is, that the Tussock might yet be unknown and unprized amongst plants, if eattle had not been introduced to its locality by man;

who thus became, first the injurer, and then the protector and propagator of the existence of this noble grass; for the herbivorous quadrupeds which he carried to the Falklands and left there, were surely extirpating the Tussock, when man returned, and, by protecting, perpetuating, and transporting it to other countries, he has widely dispersed it. It appears singular that so striking a grass should abound where there is no native herbivorous animal to profit by its luxuriance; but it is no less certain that had not civilization interfered, the Tussock might have waved its green leaves undisturbed over the waters of the stormy Antarctic Ocean, for ever perhaps, or until some fish, fowl, or seal, should be so far tempted by the luxuriance of the foliage as to transgress the laws of nature, and to adapt its organs to the digestion and enjoyment of this long-neglected gift of a bounteous Providence.

It must appear strange to all who know grasses only in the pastures of England, that the patches of Tussock resemble nothing so much as groves of small low Palm-trees! This similarity arises from the matted roots of the individual plants springing in cylindrical masses, always separated down to the very base, and throwing out a waving head of foliage from each summit. Bogs and damp woods in Britain very frequently produce a Sedge (*Carex paniculata*), whose mode of growth is, on a small scale, identical with that of the Tussock-Grass, and to which the name of Tussock is applied. I have seen them two to three feet above the ground, in South Wales; and if they were higher, larger, and placed closer together, the general resemblance would be complete. The effect in walking through a large Tussock grove is, very singular, from the uniformity in height of these masses, and the narrow spaces left between them, which form an effectual labyrinth; leaves and sky are all that can be seen overhead, and their curious boles of roots and decayed vegetable matter on both sides, before and behind; except now and then, where a penguin peeps forth from his hole, or the traveller stumbles over a huge Sea-lion, stretched along the ground, blocking up his path.

PLATE CXXXVI.—CXXXVII. Fig. 1, locusta; fig. 2, floret; fig. 3, squamulæ, stamens, and pistil; fig. 4, squamula; fig. 5, pollen; fig. 6, caryopsis:—all magnified.

15. CATABROSA, Beauv.

1. CATABROSA *Magellanica*, Hook. fil.; glaberrima, panicula elongata laxe ramosa, ramis apice floriferis elongatis, glumis inæqualibus apice erosis 4–6-floris superiore majore 3-nervi, palea inferiore ovato-oblonga obtusa 5-nervi glaberrima vix costata, culmo erecto foliorum vaginis tecto, foliorum lamina involuta vagina breviore.

HAB. Strait of Magalhaens; Port Famine, Capt. King.

Gramen pedale, erectum, glaberrimum. Culmi basi prostrati, divisi. Foliorum vagina latiuscula, 3-5 unc. longa, striata, hians; ligula ovata, acuta; lamina 2-3-uncialis, anguste lineari-subulata, superne scaberula, marginibus involutis. Panicula 5-7 unc. longa, erecta; ramis gracilibus verticillatis v. fastigiatis, inferioribus 4 unc. longis, filiformibus, glaberrimis, flexuosis, apices versus divisis et floriferis. Locustæ $\frac{1}{3}$ unc. longæ, sub 4-floræ. Gluma inferior lanceolata, acuta v. truncato-erosa; superiore oblongo-lanccolata 3-nervi obtusa erosa $\frac{1}{2}$ breviore. Flosculi basi dissiti, glaberrimi, cylindracei. Palea inferior oblongo-ovata, obtusa, sub-erosa, obscure 5-nervis, ecostata, marginibus subciliatis, superior brevior, apice bidentata. Antheræ parvæ, late oblongæ.

Quite a distinct species, and differing from the typical plants of the genus in having many florets contained in each locusta.

16. BROMUS, L.

1. BROMUS pictus, Hook. fil.; strictus, ercetus, simplex, puberulus, panicula simplici, locustis sub 4 magnis pedunculis longioribus, glumis lineari-oblongis subacutis medio nervosis flosculisque purpureo-pictis

FLORA ANTARCTICA.

sub 5-floris, palea inferiore lineari-ovata obtusa infra apicem arista brevi capillacea instructa 7-nervi inferne sericco-puberula.

HAB. Strait of Magalhaens; Port Gregory, Capt. King.

Gramen pedale. Culmus simplex, basi geniculatus, erectus, gracilis, strictus, puberulus. Folia pauca, culmo breviora; vagina striata; ligula ovata, lacera; lamina vagina brevior, pilosa, involuta. Panicula 2 une. longa. Locustæ fere pollicares, $\frac{1}{3}$ une. latæ. Flosculi nervosi, superne glaberrimi, nitidi, inferne dorso sericei; arista $\frac{1}{3}$ flosenli æquans, scaberula. Palea superior inferiori æquilonga, sed $\frac{1}{2}$ angustior.

A very distinct little species, only found in the eastern parts of the Strait of Magalhaens, and more characteristic of the grassy plains of Patagonia than of an Antaretic vegetation.

17. ELYMUS, L.

1. ELYMUS Antarcticus, Hook.fil.; erectus, glaberrimus, panicula spicæformi lineari-oblonga, spiculis binis collateralibus 2-floris, glumis subæqualibus lanceolatis aristato-acuminatis integris v. bifidis nervosis, flosculis brevissime pedicellatis, palea inferiore lanceolata in aristam gluma breviorem desinente 5-nervi superne puberula, superiore brevissime bidentata, foliis planis vaginis brevioribus.

HAB. Strait of Magalhaens; Port Famine, Capt. King.

Culmi erecti, bipedales, subvalidi, foliosi, glaberrimi. Folia eoriacea sed non rigida, culmo breviora; ragina teres, striata, 5 unc. longa; ligula brevissima; lamina lincari-subulata, utrinque lævis, basi plana, superne marginibus involutis, gradatim acuminata, 3–4 unc. longa. Panicula 3–4-uncialis, strieta, creeta, continua, $\frac{1}{2}$ unc. lata. Locustæ creetæ, imbricatæ, appressæ, scaberulæ. Glumæ liberæ, ad basin eujusvis articulationis quaternæ, quarum exterior lateris unici sæpe ad medium fissa evadit, fere $\frac{1}{2}$ unc. longæ, aristatæ. Flosculi cum aristis $\frac{2}{3}$ unc. longi, inferior vix, superior longius, pedicellatus; arista paleis paulo brevior.

This a good deal resembles a Chilian species, which possesses longer aristæ to the glumes and palea, and is otherwise different. One glume ont of the four at each articulation is often bifid, this is always the outer, and all such are on one side of the panicle only. The general appearance is not dissimilar to that of *Elymus Europæus*, from which the present species may be distinguished at once by the glumes being free to the base.

18. HORDEUM, L.

1. HORDEUM jubatum, Linn., Sp. Pl. 126. Kunth, En. Plant. vol. i. p. 457.

HAB. Strait of Magallaens; Port Famine, Capt. King.

North American specimens do not appear to differ from the Magellanic, or from others gathered at Cape Fairweather. 1 find the sheaths of the (old) leaves sometimes pilose, whence it seems very probable that the Chilian *II. carnosum*, Presl, is only a state of this plant which varies a good deal in size, in the stoutness of the culm, and length of its paniele. *II. jubatum* had been considered as confined to North America, where it ranges from the Missouri to the Saskatehewan, and from Boston to the Colombia river.

2. HORDEUM *pubiflorum*, Hook. fil.; spicis oblongis, glumis æquilongis omnibus setaceis basi pubescentibus superne scabridis, flosculis lateralibus neutris, intermedio basi setula aucto, palea inferiore scabridopalois lanceolata arista glumis æquilonga terminata, foliis caulinis longe vaginantibus radicalibus subsetaceis.

HAB. Strait of Magalhaens; Port Famine, Capt. King.

388

Radix fibrosa, subrepens. Culmi 8-10 unc. alti, basi inclinati, glaberrimi. Folia radicalia, panea; vagina uncialis glaberrima v. obscure pilosa; lamina subulata, involuta, 2-pollicaris; caulinorum vaginæ elongatæ, tumidæ, striatæ; lamina brevissima, subulata. Spica $1\frac{1}{2}$ -uncialis, fusco-purpurea. Glumæ $\frac{3}{4}$ -unciales, recurvæ, basi floseulique pube scabrida, pilis brevibus intermixtis vestita.

Very distinct from the former (than which it is a good deal smaller) and from any other species, and may be recognized by the pubescence of the florets; its general appearance resembles the European *H. maritimum*.

19. TRITICUM, L.

1. TRITICUM repens, Linu., Sp. Pl. 128. Engl. Bot. t. 909.

Var. palea superiore semper pubescente floribusque plerumque majoribus. T. repens, *var.* pungens, *Brongniart in Duperrey, Voy. Bot.* p. 57. T. glaucum, *Lamk.* (?) *D'Urville in Mém. Soc. Linn. Paris,* vol. iv. p.601.

Sub-var. 1, spiculis majoribus $\frac{2}{3}$ -uncialibus, glumis paleisque angustioribus, palea inferiore breviter aristata apice trifida v. integra.

Sub-var. 2, spiculis majoribus, glumis paleisque latioribus, palea inferiore apice mucronata subaristata integra v. obscure trifida.

Sub-var. 3, spiculis apiee majoribus, palea inferiore acuminata pungente trifida.

Sub-var. 4, spiculis minoribus, glumis angustioribus, palea inferiore apiee 3-dentata breviter aristata v. mutica.

Sub-var. 5, spiculis minoribus, glumis latioribus, palca inferiore apice 3-dentata nunquam aristata.

HAB. Strait of Magalhaens; Port Famine, *Capt. King* (sub-vars. 4 and 5); South Fuegia, Good Success Bay, and Hermite Island, *Banks and Solander, C. Darwin, Esq., J. D. H.* (sub-vars. 2 and 3); Falkland Islands, abundant, *D'Urville, &e.* (sub-vars. 1, 2, and 3).

A very variable plant, and all the more perplexing from some of the larger varieties differing more in appearance than they do in reality from the common European *T. repens*. The lower palea is generally, but not constantly tridentate at the apex, with the middle tooth sometimes produced into a short awn; it is, however, always hairy, as may be seen in Siberian, Arctic American, and Rocky Mountain plants, which latter, indeed, are sometimes villous. The North American specimens are generally larger than the European. This species is also a native of Cape Fairweather on the Patagonian coast, and exceedingly abundant throughout Fuegia and the Falkland Islands.

20. LOLIUM, L.

1. LOLIUM perenne, Linn., Sp. Pl. 122. D'Urville, in Mém. Soc. Linn. Paris, vol. iv. p. 601. Brongn. in Duperrey, Voy. Bot. p. 57.

HAB. Falkland Islands, on the sandy sea-shores, frequent; D'Urville.

The habitat assigned by D'Urville to this plant renders its being indigenous very probable.

LI. FILICES, Juss.

1. HYMENOPHYLLUM, Sw.

1. HYMENOPHYLLUM cruentum, Cav., Præl. 1801, n. 684. Swartz, Syn. Fil. p. 145. Hook. Sp. Fil. vol. i. p. 87. t. xxxi. A.

4 т

HAB. Chonos Archipelago; C. Darwin, Esq.

Even amongst the Ferns we have much peculiarity attending the Flora of S.W. Chili and Fuegia. I have elsewhere alluded to the botanical division of that line of coast into a northern and southern portion, differing specifically in their productions, but not generically to any very great amount. These two divisions are, 1st, the Valdivian or Chilotian, which stretches from Concepcion to Cape Tres Montes; and, 2nd, the Magellanic or Fuegian, commencing at Cape Tres Montes and terminating at Cape Horn. From the lower latitude and consequently higher temperature of the northern of these two countries, and from its greater surface, containing also mountains that reach the limits of perpetual snow, its Flora is by very far the richest, including a larger proportion of the Fuegian plants than Fuegia does of the Chilotian. There are also many species, which, though conspicuous in the southern Flora, are either unknown even on the Alps of the northern, or appear there only under very different aspects.

Many more species common to both these divisions, Fuegian plants especially, prevail through the whole line of coast, than its great extent would lead us to expect. This proceeds from a mutual interchange of individuals between two countries whose Floras may be supposed to have been originally quite distinct. The inosculation of the Floras is most conspicuous at Cape Tres Montes and the Chonos Archipelago, and is not accompanied by any tendency in those species, which there come iuto juxta-position, to change, each into that which represents it in the other. The union or mingling is complete, but there is no blending of two species, no obliteration of specific characters, nothing to indicate either that the peculiar plants of one country have originated from what pre-existed in the other under a different form; or, still less, that all have sprung from one common source, lower in the scale of organization.

2. HYMENOPHYLLUM pectinatum, Cav., Pred. 1801, n. 687. Swartz, Syn. Fil. p. 146. Willd. Sp. Pl. vol. v. p. 425. Hook. Sp. Fil. vol. i. p. 96. t. 34. D.

HAB. Chonos Archipelago, C. Darwin, Esq.

3. HYMENOPHYLLUM Wilsoni, Hook., Brit. Flor. ed. 5. p.446. Wilson, in Engl. Bot. Suppl. t. 2686.

Var. y. Hook. Sp. Fil. vol. i. p. 96.

HAB. South part of Tierra del Fuego, C. Darwin, Esq.; Hermite Island, Cape Horn, J. D. II. Falkland Islands, quartz rocks on the hills, J. D. II.

Found in all the four quarters of the globe, also in Australia and New Zealand.

4. HYMENOPHYLLUM Chiloense, Hook., Sp. Fil. vol. i. p. 90. t. 32. B.

HAB. Chonos Archipelago; C. Darwin, Esq.

The specimen in Mr. Darwin's herbarium is very small, but I think referable to this species; certainly to no other published one.

5. HYMENOPHYLLUM caudiculatum, Martius, Pl. Crypt. Bras. p. 102. t. 67.

Var. *β. Hook. Sp. Fil.* vol. i. p. 102.

HAB. Chonos Archipelago; C. Darwin, Esq.

6. HYMENOPHYLLUM tortuosum, Banks et Sol., MSS. Hook. et Grev. Ic. Fil. t. 129. Hook. Sp. Fil. vol. i. p. 99.

HAB. Chonos Archipelago, C. Darwin, Esq.; and throughout Fuegia, Banks and Solander, Sc.

One of the most common Antaretic American ferns, from the latitude of Valdivia to Cape Horn.
7. HYMENOPHYLLUM secundum, Hook. et Grev., Ic. Fil. t. 133. Hook. Sp. Fil. vol. i. p. 100.

HAB. Staten Land, Menzies; Hermite Island, Cape Horn, J. D. H.

Decidedly the most Antarctic of Ferns, occurring only at the very extremity of the American continent, where it is tolerably abundant in the woods.

8. HYMENOPHYLLUM rarum, Brown, Prodr. p. 159. Fl. Antaret. p. 105. H. semibivalve, Hook. et Grev. Ic. Fil. t. 83.

Var. B. Hook. Sp. Fil. l. c. H. imbricatum, Colenso, in Tasm. Phil. Journ. vol. ii. p. 187.

HAB. Var. β. South part of Tierra del Fuego, C. Darwin, Esq.; Hermite Island, Cape Horn, J. D. H.

A species exhibiting a singular predilection for those insular and peninsular localities, which terminate the continents in the Southern Ocean. Thus it occurs only at the very southern extremity of America and Africa; at Ceylon, which is nearly the southernmost land of the vast Indian empire; in Tasmania, which is an analogous position in Australasia; and in New Zealand and Lord Auckland's group, which bear the same geographical relation to Polynesia. As it also inhabits Bourbon and the Mauritius, it appears to exist all round the world, resting on the highest southern lands of each longitude.

2. TRICHOMANES, Sm.

1. TRICHOMANES flabellatum, Bory, in Duperrey Voy. Bot. Crypt. p. 281. Hook. Sp. Fil. vol. i. p. 119. T. flabellula, D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 597.

HAB. Falkland Islands; Gaudichaud, D'Urville.

A plant which I have never seen, though I diligently sought for it in the Falkland Islands. It is remarkable that both the French Naturalists who met with it, should have failed to notice the *Hymenophyllum Wilsoni*, which is sufficiently abundant, and generally accompanies the following species.

2. TRICHOMANES caspitosum, Hook., Sp. Fil. vol. i. p. 132. t. 40 B. Hymenophyllum cæspitosum, Gaud. in Ann. Sc. Nat. vol. v. p. 908, et in Freyc. Voy. Bot. p. 374. t. 5. f. 2. D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 597.

HAB. Southern parts of Fuegia; Staten Land, *Menzies*; Hermite Island, Cape Horn, abundant on trunks of trees, J. D. II.; Falkland Islands, clothing the quartz rocks on the hills; *Gaudichaud*, D'Urville, J.D. II.

This singular little species was discovered by the lamented Menzies, in Staten Land. Cape Horn is its southern limit, and Chiloe its northern. It has been probably overlooked in the intervening latitudes.

3. CISTOPTERIS, Bernh.

1. CISTOPTERIS fragilis, Bernh., Neu Journ. Bot. vol. n. p. 27. Engl. Bot. t. 1587. Hook. Sp. Fil. vol. i. p. 197.

HAB. Strait of Magalhaens; Port Famine, Capt. King; Falkland Islands (West Island?), Capt. Sulican.

One of the most extensively diffused of all vegetables, or even Ferns, though avoiding such hot and equable elimates as the low lands of the Tropics. In America, it ranges along the Cordillera, from the Arctic Sea and Greenland to the Strait of Magalhaens; in Enrope, from Iceland and Lapland to the Mediterranean; in Asia, between Kamtschatka and the Himalaya Mountains; but in Africa it is confined to the Canary Islands and the Cape of Good Hope. Its aversion to damp or uniform heat is conspicuously displayed in its not being a native of New Zealand or Fuegia proper on the one hand, or of India or tropical Africa on the other.

4. ASPIDIUM, L.

1. ASPIDIUM (Polystichum) Mohrioides, Bory, in Mém. Soc. Linn. Paris, vol. iv. p. 597, et in Duperrey, Voy. Bot. Crypt. p. 267. t. 35. f. 1. (TAB. CXLIX.)

HAB. Strait of Magalhaens; Port Famine, Capt. King; Falkland Islands, D'Urville, Sc.

The Magellanic specimens are larger, and have longer and more laxly imbricating pinnæ, than those from the Falkland Islands; which are characteristic of a climate less favourable to Ferns.

PLATE CXLIX. Fig. 1, fertile pinna; fig. 2, sterile ditto; fig. 3, sorus and involucre :- magnified.

2. ASPIDIUM (Polystichum) coriaceum, Swartz, Syn. Fil. p. 57.

HAB. Chonos Archipelago; C. Darwin, Esq.

A species apparently impatient of cold, for though inhabiting the damp west coast of Chili, as far south as the Chonos Archipelago and the dry climate of Patagonia, reaching there the latitude of Port St. Elena, it neither enters the Strait of Magalhaens, nor occurs in the Falkland Islands or Fuegia. It is almost universally diffused throughout the Tropics, and the temperate regions of the southern hemisphere.

3. ASPIDIUM (Polystichum) vestitum, Swartz, Syn. Fil. p. 53. Polypodium, Forster, Prodr. n. 445.

Var. pinnulis profundius sectis apicibus acutis.

HAB. Var. Tierra del Fuego, sonth part, C. Darwin, Esq.

The only specimen which I have seen is imperfect, but appears merely a variety of the A. restitum, with rather narrower and more deeply cut pinules, which are acute, but not pungent or spinulose; the segments of the pinules also are narrower, and the whole frond smaller. In other respects, and particularly in the clothing of the stipes, rachis, &c., it exhibits all the characters of the species I have referred it to, which is a native of Juan Fernandez and Chiloe. I am not prepared to say how far all may be distinct from the British A. aculeatum, the incisions of the broader mucronate pinules in the European plant are closer, and all aculeate, which is not the case with the typical states of A. restitum; and the clothing, too, is different.

This species is represented by the A. renustum, Homb. and Jacq., in Lord Anckland's group, and by A. proliferum, Br., in Tasmania.

5. ASPLENIUM, L.

1. ASPLENIUM Magellanicum, Kaulf. En. Fil. p. 175. Hook. et Grev. Ic. Fil. t. 180.

HAB. Strait of Magalhaens, Commerson; Port Famine, Capt. King; Hermite Island, Cape Horn, J. D. H.

A very pretty and distinct little species; probably not uncommon between the latitudes of Concepcion and Cape Horn, on the west coast of South America. It has a very nearly allied representative in New Zealand; and another, the *Asplenium laxum*, Br., in Tasmania.

6. LOMARIA, Willd.

1. LOMARIA alpina; Stegania, Brown, Prodr. p. 152. S. alpina, β. latinseula, Bory, fid. D'Urville, in Mém. Soc. Linn. Paris, vol. iv. p. 597. Lomaria polypodioides, Gaud. in Ann. Sc. Nat. vol. v. p. 908.

et in Freyc, Foy. Bot. p. 374. L. australis, Kunze, Coll. Plant. Poeppig, p. 57 (fid. sp. in Herb. Hook.). L. decurrens, Kunze, MSS. L. blechnoides, Bory (?), in Duperrey, Voy. Bot. Crypt. p. 273. L. Sellowiana, Presl, in Herb. Reg. Berol. p. 100 (fid. sp. in Herb. Hook.). L. Antarctica, Carmichael, in Linn. Soc. Trans. vol. xii. p. 512. L. linearis, Colenso, in Tasman. Phil. Journ. vol. ii. p. 176. Polypodium Pinna-marina, Poiret, Encycl. (TAB. CL.)

HAB. South Chili, Fuegia, and the Falkland Islands, most abundant; Kerguelen's Land, very scarce, J. D. II.

One of the commonest Ferns between the latitudes of Concepcion and Cape Horn on the west coast of South America, and also in the Falkland Islands, often covering the ground for many yards. It has also been collected in South Brazil by Sellow, and in Tristan d'Acunha, Kerguelen's Land, New Zealand, and in Tasmania; throughout all which countries it retains its characters very markedly, and is altogether a most distinct species. The *Blechnum loreale* is evidently its representative in the northern hemisphere, and is very similar in size, form, and habit, though abundantly distinct in the nature of its involuce.

PLATE CL. Fig. 1, portion of sterile pinna; fig. 2, fertile pinna; fig. 3, transverse section of the same.

2. LOMARIA Magellanica, Desvaux, in Mag. Nat. Berlin, 1811, p. 330, ct in Mém. Soc. Linn. Paris, vol. vi. p. 289. L. Magellanica, β. angustiseta, Bory, in Mém. Soc. Linn. Paris, vol. iv. p. 597. L. setigera, Gaud. in Ann. Sc. Nat. vol. v. p. 98, et in Freye. Voy. Bot. p. 130. L. robusta, Carm. in Trans. Linn. Soc. vol. xii. p. 512. L. zamioides, Gardner, MSS. in Herb. Hook. Pteris palmæformis, Petit Thouars, Flore de Tristan d'Acunha, p. 30. "Cetcrach," Pernetty, Voy. vol. ii. p. 56.

HAB. South Chili, Fuegia, and the Falkland Islands, very abundant.

This species is more confined in longitude but has a much wider range in latitude than *L. alpina*. I have examined what appears to be the same from British Guiana (possibly a distinct species), from Brazil, and La Plata, on the east coast of South America; and from Peru, Juan Fernandez, and Chili, on the west; it also inhabits Tristan d'Acunha. Its New Holland representative is the *L. procera*, Br.

Mr. Gardner's name of *zamioides* is peculiarly applicable both to his Brazilian and my Falkland Island specimens, they singularly resemble a *Zamia* in habit and general appearance.

7. GRAMMITIS, Sw.

1. GRAMMITIS australis, Brown, Prodr. p. 146. Fl. Ant. p. 111.

HAB. Strait of Magalhaens; Port Famine, *Capt. King.* Hermite Island, Cape Horn, abundant in the woods and on the rocks upon hills, J. D. H.

This, again, is a Fern of the Southern Ocean, being found in Tasmania, New Zealand, Lord Auckland's group and Campbell's Island, the west coast of South America, from Cape Horn probably all the way north to Lima, aud on Tristau d'Acunha. I have seen no American specimens but Capt. King's, my own, and Cuming's (n. 1052). Its tropical representative is the beautiful little *G. marginella*.

8. GLEICHENIA, Sm.

1. GLEICHENIA acutifolia, Hook., Sp. Fil. vol. i. p. 7. t. 7. A.

HAB. Strait of Magalhaens; Port Famine, Capt. King.

Like all the pedate, erect, southern species of *Gleichenia*, this has a very narrow range, and is probably confined to the coast between the Strait of Magalhaens and Chiloe, whence the specimens quoted as Patagonian in Species

4 U

Filicum were brought by Capt. King. It is represented in New Zealand by the G. Cunninghamii, Hook.; in Tasmania by G. tenera, Brown, and G. flabellata, Br.; and in Chili, north of Valdivia, it appears to be in a great measure replaced by the G. pedalis, Kaulf.

2. GLEICHENIA eryptocarpa, Hook., Sp. Fil. vol. i. p. 7. t. 6 A.

HAB. Falkland Islands (West Island?); Lieut. Robinson, Capt. Sulivan, Mr. Chartres.

Very nearly allied to the *G. acutifolia*, but readily distinguishable by the revolute margins of the pinnæ covering the sori; this character, together with the stout, rigid, and very coriaceous habit, appear to indicate its being a native of drier places than the former. It has hitherto been found only in the Falklands, in plains of Valdivia, and on the Island of Chiloe.

9. SCHIZÆA, Sm.

1. SCHIZÆA australis, Gaud., in Ann. Sc. Nat. vol. v. p. 98. Fl. Antarct. p. 111.

HAB. Falkland Islands; Gaudichaud.

I have seen no Falkland Island specimens except those collected by M. Gaudichaud, which are identical with others gathered by myself in Lord Auckland's group and Campbell's Island.

LII. LYCOPODIACEÆ, Swartz.

1. LYCOPODIUM, Br.

1. LYCOPODIUM Selago, Linu., Sp. Pl. p. 1565. Engl. Bot. t. 233.

Var. Saururus. L. Saururus, Lam. Encycl. Bot. vol. iii. p. 625. Bory, Voy. aux quatre Iles, Sc. vol. i.
p. 344. t. 16. f. i. L. crassum, H. B. K. Nov. Gen. vol. i. p. 33. Hook. et Grev. Ic. Fil. t. cexxiv.
L. insulare, Carm. in Trans. Linn. Soc. vol. xii. p. 509. L. elongatum, Swartz, Syn. Fil. p. 175. L. carinatum, Desv. Monogr. n. 5. Poiret, Encycl. Bot. vol. iii. p. 555. Selago etc. Dillenius, Hist. Musc. t. 84. f. 3.

HAB. Falkland Islands, Gaudichaud, Sc. Var. Saururus, Kerguelen's Land, J. D. H.

The Falkland Island specimens are perfectly identical with others of British growth, but the var. Saururus is so different from any aspect of *L. Selago* found in Europe, that it requires the most perfect suite of specimens, showing the gradual passage of the one into the other, to prove their common origin. Such, however, exist, especially in volcanic islands, which seem peculiarly favourable to the production of this variety.

In its largest and handsomest form, the var. *Saururus* inhabits the Andes of Pern, the Island of Bourbon, and Kerguelen's Land. A more sleuder state, but not slenderer than what occurs amongst Andes specimens, is found in Tristan d'Acunha, St. Helena, and in some of the West Indian Islands.

The narrow form of var. Saururus was brought from St. Helena as early as 1702, and given to Dillenius, who figured it. On the Andes, and there alone, the Selago division of Lycopodia assume a deep brick-red colour, which, however, affords no specific character, for the Saururus is as often wholly green as red, and at other times is only tinged with the latter colour at the apices of the leaves; and this is the case with the Kerguelen's Land specimens. It is difficult to imagine the cause for this tint of plants. On the gloomy Island of Desolation, it cannot be due to the tropical sun, nor to colouring matter contained in the soil, for it also occurs in two species which I believe are always parasitical.

2. LYCOPODIUM clavatum, Linn., Sp. Pl. p. 1564. Engl. Bot. t. 224.

Var. Magellanicum, foliis apice muticis. Fl. Ant. p. 133.

HAB. Var. Magellanicum, Strait of Magalhaens and throughout Fuegia, the Falkland Islands, very abundant; Kerguelen's Land, J. D. H. Var. fustigiatum, Port Famine, Capt. King.

I have in the former part of this work given my reasons at length for assigning these varieties to *L. clavatum*. The var. *fastigiatum* is a plant of a warmer climate than the var. *Magellanicum*, which inhabits not only the low-lands of Fuegia, the Falklands, and Kerguelen's Land, but also the lofty heights of the Cordillera of Peru and Colombia, and the mountains of New Zealand, Tasmania, and Lord Auckland's group.

LIII. MARSILEACEÆ, Br.

1. AZOLLA, Lam.

1. AZOLLA Magellanica, Willd., Sp. Pl. vol. v. p. 541. A. filiculoides, Lam. Encycl. vol. i. p. 340.

HAB. Strait of Magalhaens, Commerson; Falkland Islands, Gaudichaud.

I am quite nnacquainted with this species, either as a Falkland Island or Magellauic plant.

LIV. CHARACEÆ,

1. CHARA, L.

1. CHARA flexilis, Linn., Sp. Pl. 1624. Smith, Engl. Bot. t. 1070.

HAB. Kerguelen's Land, in the fresh-water lake above Christmas Harbour, abundant, J. D. II.

After a careful comparison of this plant with English specimens of *C. flexilis*, I cousider them to be the same species, and am confirmed in this opinion by my friend Mr. Wilson, who has studied the British species of this difficult genus very carefully; he says, that the points at the apices of the branches are, perhaps, longer than common in the Antarctic specimens. It is probably not an unfrequent plant in the southern temperate zone.

LV. MUSCI, L.

By W. Wilson, Esq., and J. D. Hooker.

1. ANDREÆA,* Ehrh.

1. ANDREÆA alpina, Linn.; caule ramoso elongato, foliis undique imbricatis patentibus apice incurvis obovatis acumiuatis concavis infra medium contractis siccitate appressis. A. alpina, *Dill. Hist. Musc.* t. 73. f. 39. *Hook. et Tayl. Musc. Brit.* ed. 2. p. 2. t. 8.

Var. 1. foliis inferioribus squarrosis subrectis.

Var. 2. caulibus gracilioribus, foliis confertis.

HAB. Var. 1 and 2, Hermite Island, Cape Horn; Kerguelen's Land, var. 1, on alpine rocks.

This species has in Europe frequently been confounded with *A. rupestris*, and we cannot assent to the remark in the 'Muscologia Britannica, that Dr. Mohr was the first to distinguish it accurately; since neither the description of Weber and Mohr (*Bot. Tasch.* p. 383), nor their citation of Dillenius (*Hist. Musc.* t. 73. f. 40), as a synonym for *A. rupestris* instead of *A. Rothii*, tends to prove that these authors understood the species. The illustrative figures

^{*} For the generic characters and remarks on this and other genera, see the 1st Part of this work.

FLORA ANTARCTICA.

and observations show that *A. alpina* of Weber and Mohr (*Bot. Tasch.* t. 11. f. 3, 5), is only a state of *A. rupestris*, with leaves spreading in all directions. This is confirmed by the fact that *A. alpina* of Mougeot and Nestler (*Stirp. Crypt. Voges.* no. 115), is that very form of *A. rupestris* to which we allude. The figures of Dillenius doubtless refer to *A. alpina*; but the absence of a separate figure of *A. rupestris*, and the remark, "in rupibus surchi e fusco rufescunt" (p. 507), prove that this author considered both these as varieties of one species. The true *A. alpina* may always be known from *A. rupestris* by its obovate dark glossy leaves, and its longer and more robust stems.

2. ANDREÆA marginata, Hook.fil. et Wils.; caulibus laxe cæspitosis subramosis, foliis erecto-patentibus incurvis ovatis longe acuminatis infra medium contractis enerviis marginibus inferne pallidis caulinis majoribus superioribus confertis, perichætialibus longioribus elongatis late lanceolatis convolutis, theca exserta. (TAB. CLI. Fig. I.)

HAB. Hermite Island, Cape Horn, frequent on rocks ou the mountains; J. D. H.

Caules unciales. Folia e basi ovata longe acuminata, concava, superiora in caule fertili sensim majora, laxiora et magis erecta, omnia puniceo-atra, nitentia, areolis minutis, oblongis, marginalibus inferne minoribus, pallidioribus.

With some hesitation we venture to separate this from A. *alpina*, on account of the more gradual acumination of its leaves, which are full twice as long. It differs from A. *acutifolia* in having the leaves considerably larger, the lower part more suddenly dilated, and the base not gibbous; also in their dark colour and glossiness, and in the more evident perichaetium. It is easily recognized by its habit.

PLATE CLI. Fig. I.-1, a specimen :-natural size; fig. 2 and 3, leaves; fig. 4, theca:-magnified.

3. ANDREÆA acutifolia, Hook. fil. et Wils., vid. Part 1. p. 118.

Var. β . rufescens, ramis fastigiatis. (TAB. CLI. Fig. II. 2.)

Var. γ . foliis latioribus.

Var. 8. foliis superioribus subsecundis. (TAB. CLI. Fig. II. 1.)

Var. c. foliis superioribus erectis elongatis.

Var. ¿. foliis erectis obtusiusculis atro-sanguineis.

HAB. Var. α , β , and ϵ , Hermite Island, Cape Horn. Var. δ and ζ , Falkland Islands. Var. γ , Kerguelen's Land.

All these varieties differ somewhat, though slightly, from that gathered in Campbell's Island.

PLATE CL1. Fig. II.—1, specimen of var. δ , and 2, specimen of var. β :—natural size; fig. 3, perichaetium and theca; fig. 4, perichaetial leaves; fig. 5, cauline leaf of var. δ ; fig. 6, leaf, and fig. 7, theca of var. β :—magnified

4. ANDREÆA rupestris, Linn.; caule humili subramoso, foliis e basi vaginante patentibus (interdum secundis) ovato-lanceolatis vel ovatis superne attenuatis acutiusculis enervibus superioribus siccitate appressis, perichætialibus longioribus ovato-lanceolatis convolutis, theca exserta. A. rupestris, *Hook. et Tayl. Musc. Brit.* p. 2. t. viii.

ILAB. Hermite Island, Cape Horn, frequent on maritime rocks.

Slightly differing from British specimens in its wider, less evidently papillose leaves, which are more suddenly dilated near the middle. Bridel (Bryol. Univ.) cites with doubt, Dillenius (Musc. p. 507. t. 73. f. 40) as a synonym for this species, but the description of Dillenius distinctly mentions the nerved leaves which are characteristic of A. Rothii, to which species even the description of Linnæus (as Bridel properly remarks) seems to refer. The secund foliage, generally ascribed to A. rapestris, is not a constant character, even in specimens gathered in the same locality.

5. ANDREÆA mutabilis, Hook. fil. et Wils., vid. Part 1. p. 119. pl. lvii. f. ii.

Var. y, subsecunda; foliis laxioribus inferioribus secundis.

Var. 8. uncinata; foliis dissitis falcato-secundis.

HAB. Falkland Islands; both varieties, abundant.

The var. β . of this species is a Lord Auckland's group and Campbell Island plant.

6. ANDREÆA *laxifolia*, Hook. fil. et Wils.; caulibus laxe cæspitosis parce ramosis, foliis lanceolatosubulatis obtusiusculis concavis enerviis ramulinis falcato-secundis caulinis erectis subsecundis laxe imbrieatis, perichætialibus elongatis ovato-lanccolatis convolutis, theca exserta ovato-oblonga. (TAB. CL1. fig. IV.)

Var. β . *minor*; theca subexserta.

HAB. Hermite Island, Cape Horn; not rare, on moist rocks in the higher parts of the Island. Var. β . on rocks near the sea.

Caulis vix uncialis. Folia ramulorum conferta, falcato-secunda, luteo-viridia; caulina dissita, majora, vix secunda, subamplexicaulia. Florescentia monoica: flos masculus primo terminalis, folia perigonialia rotundo-ovata, acutiuscula, concava. Antheridia 6. Paraphyses numerosæ, duplo longiores. Theca siccitate turbinata, basi pallida.

PLATE CLI. Fig. IV.-1, a tuft of the natural size; 2, 3, 4, and 5, leaves; 6, theca:-magnified.

7. ANDREEA subulata, Harvey; vid. Part 1. p. 119. pl. lvii. f. i.

Var. β . rigida; foliis minus falcatis crassioribus luridis.

IIAB. Hermite Island, Cape Horn, and the Falkland Islands; not uncommon.

The Auckland and Campbell Island variety differs slightly from the above.

Subgen. ACROSCHISMA, Hook. fil. et Wils. (Theca cylindracea, e basi ad medium et ultra indehiscens, apicem rersus tantum in valvulis 4 vel 8 fissa ;—an genus proprium?)

8. ANDREÆA (Acroschisma) *Wilsoni*, Hook. fil.; caule laxe cæspitoso elongato ramoso, foliis ramulorum undique pateutibus squarrosis spathulato-lanceolatis obtusiusculis caulinis erectis laxe imbricatis ovatolanceolatis basi angustatis amplexicaulibus omnibus enervibus concavis marginibus inflexis, perichætialibus elongatis elliptico-oblongis convolutis, theca exserta cylindracea apicem versus fissa. (TAB. CLI. fig. III.)

HAB. Hermite Island, Cape Horn; on maritime rocks near the spray of rivulets, rare.

Caules sesquiunciales, graciles, erecti, per intervallos breves innovantes, subdichotomi, steriles vage ramosi, rami patentes. Folia ramulorum lanceolata, basi subamplexicaulia, erecta, deinde patula, squarrosa, apice subincurva, obtusiuscula, lateribus inflexis, luteo-viridia, caulina majora, erecta, infra medium repente augustata, flavescentia, enervia, areolis majusculis elongatis. Florescentia monoica: antheridia circiter 11, paraphysibus numerosis longissimis. Theca elongata, maxima, inferne integra, badia, apice in valvulis 4 vel 8 fissa.

Allied to *Andreæa laxifolia*, but differing in the remarkable capsule and also in the form of the leaves, which do not taper gradually from the base, but are widened near the middle.

PLATE CLI. Fig. III.—1, a plant of the natural size; 2, branches; 3, young theca, &c.; 4, lcaf; 5, mature theca; 6, perichætial leaf:—all magnified.

2. SPHAGNUM, L.

In addition to the former remarks on the structure of *Sphagnum*, we would observe, that the spirally lined cellules of the leaves do not constitute the proper parenchyma. The cellules, which contain the chlorophyll, are

4 x

those which are interposed between the larger utricles, and which form the network of the leaf. This is well explained by the cauline leaves of *Sphagnum fimbriatum* (Wils. MSS.), a very curious British and Antarctic species, long confounded with *S. aculifolium*; in them the spirally lined cells are altogether absent. Spirally lined cells communicate with each other by pores, as we have ourselves witnessed the passage of animalcules (vibrio) from one cell into another.

1. SPHAGNUM cymbifolium, Dill.; caule elongato, ramis crassis, foliis imbricatis patentibus ovatis obtusis concavis superne deuticulatis cellulis ramulorum spiraliter lineatis. S. cymbifolium, Nees et Hornsch. Bryol. Germ. vol. i. p. 6. t. 1. f. 1. S. obtusifolium, Hook. et Tayl. Musc. Brit. ed. 2. p. 13. t. 4 (ex parte).

Var. 2. condensatum, Hook. fil. et Wils.; caule humili, ramulis brevissimis undique dense confertis. S. condensatum, Brid. Bryol. Univ. vol. i. p. 18 (?).

HAB. Falkland Islands; common in streams, bogs, and peat-ponds. Var. 2. Strait of Magalhaens, Port Famine, Capt. King.

In general aspect exceedingly like the more compact form of *S. compactum*, Bridel, but preserving the true character of the species to which we refer it, in the shape of the leaves, and in the markings of the ramuline cellules. Our specimens are all fertile, the stems not two inches in length. An example occurs where two capsules are produced upon the same pseudopodium.

Mr. Valentine was the first to point out (in the 'Muscologia Nottinghamensis') the structure of the cells of the ramuli, which, from oft-repeated observation, we consider a valid specific character, distinguishing this species from all others. On the other hand, the characters derived from the length of the peduncle and the disposition of the branches appear to be fallacions.

2. SPHAGNUM *fimbriatum*, Wils. MSS.; caule longiusculo gracili subramoso, foliis dimorphis, caulinis obovato-subrotundis obtusissimis fimbriatis, rameis ovato-lanceolatis acuminatis concavis acutis, perichætialibus obovatis obtusis valde concavis, theca brevi-peduuculata.

HAB. Hermite Island, Cape Horn, and the Falkland Islands.

Caulis plerumque gracilis. Rami 3-nati subinde 4–5-nati, longiusculi, apice attenuati. Folia caulina erecta, subrotunda, obtusissima, fimbriata!, cellulis propriis (chlorophyllo farctis) reticulum formantibus, interstitiis (e defectu utriculorum linea spirali notatorum) vacuis, folia ramorum conferta, creeto-patentia, apice subrecurva, concava, acuta, perichætialia subcucullata, subretusa, obtusissima, concava, thecam immaturam arete amplectentia. Theca matura globosa, pedicello breviusculo exserto.

From *Sphagnum acutifolium*, Ehrh., this species may be readily known by its more slender habit, and is essentially distinguished by the peculiar cauline leaves, which consist of an open net-work of parenchymatous cells without any intermediate ones lined with spiral filaments; the perichaetial leaves are also very different in shape, and those of the hranches are more acute, their reticulation also is, especially at the summit, considerably smaller.

The specimens here described are not so slender as others gathered in Britain; but possess all their essential characters; the *S. acutifolia* of Montagne (Voy. au Pole Sud, Bot. Crypt. p. 282) is probably the same plant.

3. SPHAGNUM cuspidatum, Ehrh.; ramulis attenuatis laxis, foliis lanceolato-subulatis laxis patulis siccitate undulatis marginibus reflexis perichætialibus acutis. S. cuspidatum, Nees et Hornsch. Bryol. Germ. vol. i. p. 13. t. 4. f. 9. Hook. et Tuyl. Musc. Brit. p. 15. t. iv.

HAB. Hermite Island, Cape Horn, and the Falkland Islands; common.

Neither this, nor any of the other southern *Sphagna*, are so universal in the Antarctic bogs as they are in the European and Arctic.

FLORA ANTARCTICA.

Falklands, etc.]

3. SCHISTIDIUM, Bridel.

Stoma nudum. Calyptra mitræformis s. campanulato-conica, in lacinias plures subæquales basi fissa, rarius integra. Theca æqualis, exapophysata.

1. SCHISTIDIUM marginatum, Hook. fil. et Wils.; caule erecto, foliis erecto-patentibus lanceolatis subapiculatis marginatis solidi-nerviis subdenticulatis, theca immersa subrotunda, operculo conico-rostrato erecto. (TAB. CLI. fig. VI.).

HAB. Kerguelen's Laud; not uncommon on moist banks, alt. 500 feet, forming large patches on wet rocks, and on the debris of precipices.

Caules crecti, vix ramosi, 4 lin. longi, cæspitosi, pallide rubri, succulenti. Folia imbricata, erecto-patentia, siccitate paulo tortilia, lanceolato-oblonga, subundulata, inferiora spathulato-lingulata, nervo valido rufo subexcurrente instructa, rufo-viridia, areolis subquadratis, minutis; perichætialia paulo longiora, angustiora, erecta. Seta theca brevior, erecta, fusco-lutea. Theca ovato-cyathiformis, badia, basi rotundata, ore aperto. Annulus persistens, inconspicuus. Operculum conico-subulatum, capsula paulo longius. Calyptra campanulata, apice rufo-brunnea, basi pallida, membranacea, demum lacera. Sporæ minimæ, lutescentes. Species dioica?

Very distinct from all other described species.

PLATE CLI. Fig. VI.-1, a tuft of the natural size; 2 and 3, leaves; 4, theca; 5, calyptra:-magnified.

4. GYMNOSTOMUM, Hedw.

Stoma nudum. Calyptra cuculliformis vel ventricoso-subulata, latere fissa. Theca æqualis, exannulata.

 GYMNOSTOMUM Heimii, Hedwig; foliis patentibus oblongo-lanceolatis subconcavis apice denticulatis nervo subcontinuo, theca truncato-obovata et oblonga, operculo oblique rostellato columellæ insidente.
 G. Heimii, Hedwig, Stirp. Crypt. vol. i. p. 84. t. 30. Hook. et Tayl. Muse. Brit. p. 22. t. vii. Pottia Heimii, Bruch et Schimper, Bryol. Europæa.*

Var. 1, foliis angustis apice vix serrulatis.

Var. 2, foliis subovato-acuminatis margine pellucidioribus, theca turbinata brevi.

Var. 3, foliis latioribus obovatis concavis.

HAB. Falkland Islands, var. 1 and var. 3 (imperfect) perhaps a distinct species, found near the sea, in sandy places. Hermite Island, Cape Horn; var. 1, on maritime rocks; var. 2, on sandy banks.

Subgenus Physcomitrium, Bridel.

2. GYMNOSTOMUM (Physcomitrium) *laxum*, Hook. fil et Wils.; cæspitosum, caule simpliciusculo, foliis erecto-patentibus laxe imbricatis elliptico-lanceolatis acutis concaviusculis integerrimis reticulatis siccitate vix crispatis, nervo sub apice evanido. (TAB. CLI. fig. V.)

^{*} For the sake of brevity, we omit in this work the generic distinctions proposed by Bruch and Schimper in the 'Bryologia Europæa', without, however, intending to question their validity in a natural system. *Pottia* comprises those terrestrial species of *Gymnostomum*, exclusive of *Physcomitrium*, which are of bi-triennial duration and have monoicous inflorescence.

FLORA ANTARCTICA.

HAB. Kerguelen's Land, not uncommon on moist sandy banks (barren). Closely allied to the British *Physcomitrium pyriforme*, of which it may be a variety. PLATE CLI. *Fig.* V.—1, a tuft of the natural size; 2 and 3, leaves; 4, apex of ditto:—*magnified*.

5. LEPTOSTOMUM, Br.

1. LEPTOSTOMUM *Menziesii*, Brown; caule subsimplici, foliis oblongo-ovatis apice denticulatis nervosis piliferis, theca oblongo-clavata subrecurva horizontaliter inclinata, operculo conico, rostro brevi obliquo. Gymnostomum Menziesii, *Hook. Musc. Exot.* t. 6.

HAB. Hermite Island, Cape Horn; very common from the sea coast to the top of the hills, alt. 1700 ft., on trees, rocks and banks, forming large noble tufts. Strait of Magalhaens, D'Urville.

6. SPLACHNUM, L.

1. SPLACHNUM Magellanieum, Brid.; caule crecto subdiviso, foliis oblongo-lanccolatis acuminatis serratis evanidinerviis, pedunculis aggregatis, thecæ oblongæ ovatæ apophysi obconica, operculo convexo. S. Magellanicum, Schwaegr. Suppl. I. pt. 4. p. 47. t. 14. Eremodon Magellanicus, Brid. Bryol. Univ. vol. i. p. 236.

HAB. Hermite Island, Cape Horn, on the horizontal limbs and dead stumps of old trees in the woods, growing in large tufts.

Calyptra conico-mitræformis, basi appendiculata, dcmum lacera, pallida, apice flavescens.

Our specimens, though not in the best state for the examination of the capsules, all possess apparently a peristome of eight teeth; nevertheless we do not consider it necessary to remove this moss from *Splachnum*. Bridel's genus *Eremodon* is not adopted by Bruch and Schimper, and is indeed untenable. Our moss has characters which induce us to doubt whether it should be referred to *Tayloria* or to *Splachnum* of Bruch and Schimper. Its affinity with the European *Tayloria serrata* and *Rudolphiana* is too striking to be overlooked; but the pale apophysis, though not wider than the capsule, is evidently that of a true *Splachnum*; while the peristome connects it with *Dissodon*, Br. and Sch.

7. GRIMMIA, Ehrh.

Peristomium simplex. Dentes sedecim, pyramidati, pertusi, rarius imperforati, reflexiles. Calyptra mitræformis. Theca æqualis.

1. GRIMMIA *tortuosa*, Hook. fil. et Wils.; caule pulvinato, foliis erecto-patentibus lineari-lanccolatis acuminatis piliferis siccitate tortuosis, theca immersa subsessili urceolata, operculo convexo apiculato. (TAB. CLI. fig. VII.)

IIAB. Falkland Islands; dry quartz rocks on Mount Vernet, alt. 1,000 feet, very scarce.

Caules 3-4-linearcs, pulvinati, subramosi. Folia conferta, erecto-patentia, subflexuosa, lineari-lanceolata, acuminata, pilifera, carinata, margine paulo incrassata, subplana, nervo valido, dorso prominente, excurrente, siccitate tortilia, subcrispata, opaca, atroviridia, areolis minutissimis, punctatis, basi majoribus, reticulatis, pellucidis; perichætialia similia. Seta brevissima, vix ulla. Theca immersa, subrotunda, erecta, fusca, ore patulo. Annulus nullus. Peristomii dentes conniventes, siccitate erecti, subreflexi, pyramidati, integri, rubri. Operculum planiusculum, subrostellatum. Sporæ minimæ, ferrugineæ. Calyptra brevis, basi lacera, fusca, apice brunnea.

Apparently a distinct species, somewhat allied to *G. apocarpa*, as to the fruit; but in the foliage, more nearly to *G. trickophylla*.

2. GRIMMLA *fulcata*, Hook. fil. et Wils.; caulibus laxe cæspitosis pendulis, foliis falcato-secundis lanceolato-subulatis crassinerviis canaliculatis integerrimis, theca immersa subsessili turbinata, operculo rostellato. (TAB. CLI. fig. VIII.)

HAB. Kerguelen's Land; on rocks and stones near a small waterfall.

Caules laxe cæspitosi, 1-3-unciales, penduli, flexuosi, ramosi, rami subincurvi. Folia imbricata, falcato-secunda, lanceolato-subulata, carnosa, integerrima, canaliculata, lateribus inflexis, nervo lato crasso excurrente, lurido-viridia, inferiora sæpe aqua destructa, nervo solo residuo, areolis minutis subquadratis. Perichætialia ovato-lanceolata, acuminata, thecam superantia. Vaginula conica. Theca subsessilis, subrotunda, ore patulo, brunnea. Annulus nullus? Peristomii dentes magni, apice subperforati, incurvi, dorso trabeculati, rubri, siccitate recurvi. Operculum hemisphæricum, rostellatum, capsula brevius. Calyptra fusca, mitræformis, brevis. Flos masculus in axillis ramulorum ad basin ramuli fructiferi. Antheridia plurima, eparaphysata.

Allied to *Grimmia apocarpa*, var. *rivularis*, but very distinct in its falcate leaves, and broad thick nerve. As in that species, the columella generally falls away with the operculum. The short fertile branches are often clustered two or three together.

PLATE CLI. Fig. VIII.—1 and 2, plants of the natural size; 3, apex of branch; 4, leaf; 5, perichætial ditto; 6, theca; 7, operculum:—magnified.

3. GRIMMIA maritima, Turner, Musc. Hib. p. 23. t. 3. f. 2. Hook. et Tayl. Musc. Brit. p. 66. t. xiii. Schistidium maritimum, Bruch et Schimper, Bryol. Europ. fasc. 25-28. p. 10.

HAB. Hermite Island, Cape Horn; on granite rocks near the sea at St. Joachim's Bay.

Ab exemplis Britannicis his notis differt : *caulibus* longioribus pluries ramosis ; *foliis* magis patulis, subrecurvis, perichetialibus apice diaphanis, nervo angustiore ; *capsula* majore.

This moss affects the same localities in the Antarctic regions that it does in England.

4. GRIMMIA apocarpa, Linn.; Hedw. Musc. Frond. vol. i. p. 104. t. 39. Hook. et Tayl. Musc. Brit. p. 65. t. xiii. Schistidium apocarpum, Bruch et Schimper, l. c. p. 7.

Var. 1, foliis subpiliferis subcrectis.

Var. 2, foliis obtusiusculis subpatentibus, perichætialibus obtusis.

Var. 3, foliis caulinis angustioribus, perichætialibus prægrandis obtusis.

Var. 4, foliis lineari-lanceolatis longioribus.

HAB. Falkland Islands; var. 1, slate rocks near the sea. Var. 2 and 3, Kerguclen's Land, on rocks, alt. 500 feet. Hermite Island, Cape Horn; on trap rocks near the sea.

The most striking feature of the three last named varieties is the large obtuse perichaetial leaves. The habit and place of growth are similar to what this moss inhabits in Britain.

8. DRYPTODON, Brid.

This Bridelian genus appears to have been properly reduced to a section of *Racomitrium*, in the 'Bryologia Europæa' of Bruch and Schimper.

4 Y

FLORA ANTARCTICA.

1. DRYPTODON *rupestris*, Hook. fil. et Wils.; caule gracili fastigiato-ramoso, foliis patentibus ovatolanceolatis carinatis margine recurvo nervo subcontinuo, seta brevi, theca elliptico-oblonga erecta, operculo rostrato. (TAB. CLII.fig. I.)

IIAB. Hermite Island, Cape Horn; moist rocks on the northern slope of Mount Foster, alt. 600 feet.

Caules 1–2-unciales, dense cæspitosi, gracilescentes, fastigiato-ramosi. Folia dense imbricata, patentia, ovatolanceolata vel ovato-acuminata, inferiora subsquarrosa, superiora recurva, acutiuscula, carinata, margine subrecurva, nervo rubello, siccitate appresso-incurva, plus minusve spiraliter contorta, lurido-viridia. Perichætialia latiora, elliptico-oblonga, obtusa. Seta brevis, $1\frac{1}{2}$ lin. longa, recta, siccitate tortilis. Theca erecta, elliptico-oblonga, subpyriformis, parvula, subcoriacea, rufo-brunnea, ore contracto. Peristomii dentes subsimplices, rubri, conniventes, siccitate patentes. Operculum rostro aciculari, theca paulo brevius. Calyptra non visa.

A more robust species than D. crispulus (nobis), with the leaves shorter, not piliferous, and the thece larger.

2. DRYPTODON crispulus, Hook. fil. et Wils.; vid. Part 1. p. 124. pl. lvii. f. ix.

Var. β . foliis siccitate patulis rigidioribus.

HAB. Kerguelen's Land, in gravelly beds of rivulets.

Also a native of Campbell's Island, but not seen at Cape Horn or the Falkland Islands.

9. RACOMITRIUM, Brid.

I. RACOMITRIUM protensum, Al. Braun; Bruch et Schimper, Bryol. Europ. fasc. 25-28. tab. Drypt. 2. R. aquaticum, Brid. Bryol. Univ. vol. i. p. 222. R. cataractarum, Braun, Brid. l. c. Suppl. p. 776.

Var. 1, subaquaticum, foliis acutis subsecundis.

Var. 2, caule humili, thecis minoribus.

Var. 3, caule robustiore, foliis longioribus secundis.

HAB. Hermite Island, Cape Horn; all the varieties. Falkland Islands; var. 2 (barren) and var. 3. Kerguelen's Land, also barren).

The last variety much resembles *R. aciculare*, Dill., but has acute leaves and the teeth of the peristome arc longer and more slender. It is perhaps still more nearly allied to *Trichostomum subsecundum*, Hook. and Grev., (Hook. le. Pl. t. 17. f. 5), chiefly differing in its more robust habit and shorter setae.

2. RACOMITRIUM fasciculare, Dill., Bridel, Br. Univ. vol. i. p. 218. Bruch et Schimp. Bryol. Europ. Trichostomum fasciculare, Schwaegr. Suppl. I. pt. 1. p. 155. t. 38.

Var. 2, caule gracili, ramulis brevissimis, foliis subsecundis luteo-viridibus, calyptra pallida.

Var. 3, caule gracili subsimplici, foliis acutiusculis siccitate subappressis, seta breviore, calyptra pallida.

HAB. Hermite Island, Cape Horn; all the varieties.

The last variety has some resemblance to *R. heterostichum*, var. γ . gracilescens, (Bruch and Schimper), but differs in having the leaves more acute and the teeth of the peristome larger and more regularly formed.

3. RACOMITRIUM heterostichum, Brid.; Bryol. Univ. vol. i. p. 214. Bruch et Schimper, Bryol. Europ. fasc. 25–28. p. 9. t. 1. Trichostomum heterostichum, Hedw. Musc. Frond. vol. ii. t. 25.

HAB. Falkland Islands; on moist rocks on the hills, barren and stunted.

4. RACOMITRIUM lanuginosum, Brid.; Bryol. Univ. vol. i. p. 215. Flor. Antarct. pt. 1. p. 124. Bruch et Schimper, Bryol. Europ. l. c. p. 11. t. 6. Trichostomum lanuginosum, Hedw. Musc. Frond. vol. jii. t. 2.

HAB. Falkland Islands; common on the hills, barren. Hermite Island, Cape Horn; also barren. Strait of Magalhaens; D'Urville.

This moss is very common throughout the Antarctic regions. The specimens from Hermite Island have the leaves more obscurely toothed than British examples, and the branches very short. It may be R. Borbonicum, Brid. (Br. Univ. vol. i. p. 218).

10. ORTHOTRICHUM, Hedw.

1. ORTHOTRICHUM crassifolium, Hook. fil. et Wils.; see Part 1. p. 125. pl. lvn. f. viii.

Var. γ , foliis superioribus ovato-lanceolatis acutiusculis.

Var. 8, foliis subsecundis.

HAB. Hermite Island, Cape Horn; vars. 1 and 3, very common on maritime rocks of granite and trap. Falkland Islands; var. 1, also on clay-slate. Kerguelen's Land; vars. 1 and 3 common.

Both these varieties differ from the Campbell's Island state of the moss.

2. ORTHOTRICHUM *luteolum*, Hook. fil. et Wils.; caule erecto ramoso, foliis erecto-patentibus siccitate crispulis anguste lineari-lanceolatis basi dilatatis margine planis, theca exserta ovali-oblonga siccitate sulcata, calyptra pilosa. Orthotrichum coarctatum, *Schwaegr. Suppl.* I. 2. p. 26. t. 52 (excl. syn. Belvisian.?). *Hook. et Grev. ! in Brewst. Journ.* vol. i. p. 125 (nec Orth. coarctatum, *Br. et Schimp. Bryol. Europ.*) (TAB. CLII. fig. II.)

Var. β . calyptra glabra.

HAB. Hermite Island, Cape Horn; var. a. and β . on stems of shrubs, especially of *Berberis ilicifolia*, from the sea to alt. 1,000 feet, abundant, forming round soft tufts, rare on rocks. South part of Tierra del Fuego, *C. Darwin, Esq.*, n. 140.

Caules laxe pulvinati, luteoli, subunciales, ramosi. Folia conferta, erecto-patentia, curvula, anguste linearilanceolati, basi dilatata, ovata, utrinque laxe et pellucide reticulata, margine plana, nervo rubello, areolis laxe punctatis, luteola, siecitate parum erispula. Faginula subpilosa, ovata, minuta. Seta longitudine varia folia perichætialia plerumque æquans vel superans, siecitate striata, in collum eapsulare sensim dilatata. Theca subpyriformis, parva, pallide luteo-fusca, 8-striata, sieca et vacua cylindracea, vix sulcata, ore haud constricto. Peristomii externi dentes 8, bigeminati, siecitate reflexi; interni cilia. Calyptra campanulata, pilosa, straminea, in var. β . glabra, brunnea. Florescentia monoica.

It will be seen how closely the description corresponds with that of *O. coarctatum*, Br. and Schimp.; but on comparing authentic specimens, we find the leaves in the latter much wider and carinate, less dilated at the base, more crisped when dry, the vaginula twice as long, capsule larger, and the habit considerably different. Our moss, unlike that, has very little resemblance to *O. crispum*, and is remarkable for its pale yellowish colour; it varies in the length of the seta. An original specimen of *O. coarctatum* (from P. de Beauvois in Professor Arnott's

Herbarium), as also the description (in Ætheog. p. 80), prove it to be identical with O. Ludwigii, Schwacgr., which therefore ought to have been named O. coarctatum.

PLATE CLII. Fig. II.—1, plant of the natural size; 2 and 3, leaves; 4, seta, theca, &c.; 5, calyptra; 6, theca; 7, teeth of peristome:—magnified.

3. ORTHOTRICHUM erispum, Hedw.; Musc. Frond. vol. ii. t. 35. Hook. et Tayl. Musc. Brit. p. 133. t. xxi. Bruch et Schimper, Bryol. Europ. fasc. 2-3. p. 23. t. 12.

HAB. Hermite Island, Cape Horn; on rocks and branches of trees near the sea, always barren.

This species often bears, at Hermite Island, jointed conferva-like gemmæ among the young leaves.

4. ORTHOTRICHUM Magellanicum, Mont., in Voy. au Pole Sud, Bot. Crypt. p. 290. t. 20. f. 2.

HAB. Strait of Magalhaens; M. Jacquinot.

11. MACROMITRIUM, Brid.

1. MACROMITRIUM longipes, Schwaegr.; Suppl. II. 2. p. 131.? Orthotrichum longipes, Hook. Musc. Exot. t. 24.

Var. ramis gracilioribus elongatis, foliis luridis crectis lineari-oblongis plicato-carinatis nervo excurrente. HAB. Hermite Island, Cape Horn, Mr. Davis; barren.

Perhaps a distinct species; but though different in aspect, obvious characters are wanting to distinguish it from the Hookerian specimens.

12. WEISSIA, Hedw.

1. WEISSIA crispula, Ludw.; vid. Part 1. p. 127. t. lviii. f. ii. Dicranum interruptum, Brid. Bryol. Univ. vol. i. p. 438. Bryum pilosum interruptum, Dill. Musc. p. 376. t. 47. f. 38.

HAB. Hermite Island, Cape Horn; rocks on Kater's Peak, alt. 1,000–1700 feet, growing in tufts. Apparently identical with the European plant, and also found in Campbell's Island.

2. WEISSIA contecta, Hook. fil. et Wils.; vid. Part. 1. p. 127. t. lxiii. f. iii.

IIAB. Kerguelen's Land; barren, on rocks.

Also a native of Campbell's Island.

3. WEISSIA acuta, Hedw.; Musc. Frond. vol. iii. t. 35. Hook. et Tayl. Musc. Brit. p. 87. t. 14.

Var. β . theca subrotunda, seta breviore arcuata, peristomii dentibus latioribus cribroso-pertusis.

HAB. Falkland Islands; at Port Louis, barren. Hermite Island, Cape Horn; var. β . on wet rocks in and near water-courses, in St. Martin's Cove; wet sandy banks on Mount Foster.

We have no specimen of Weissia acuta β ., Wahlenb. (Fl. Lapp.), which seems to differ, according to the description very little from our moss. The capsule of ours is turbinate when dry, with a very wide mouth. The seta is sometimes equally short in British specimens.

4. WEISSIA *stricta*, Hook. fil. et Wils.; caule ramoso, foliis subfalcatis lanceolato-setaceis rigidis canaliculatis integerrimis crassinerviis nervo longe excurrente, theca subrotunda, operculo rostrato. (TAB, CLHI. fig. IV.)

FLORA ANTARCTICA.

HAB. Kerguelen's Land; on rocks near the sea, not uncommon.

Caules unciales, laxe cæspitosi, ramosi. Folia suberecta, conferta, rigida, siccitate vix crispata, nervo lato crasso longe excurrente instructa, luteo-viridia; perichætialia longiora, basi latiora, crecta, convoluta. Seta 3-linearis, erecta vix tortilis, pallide rufa. Theca suberecta, subrotunda, ore contracta, rufo-brunnea, demum atrorubens, vernicosa. Peristomii dentes 16, parvuli, conniveutes, pyramidati, obtusiusculi, linea media notati. Operculum conico-rostratum, capsulæ longitudine, rostro curvato acuto. Calyptra dimidiata, subventricosa, fusco-lutea. Florescentia monoica.

We know of no described species with which this can be confounded. It is somewhat allied to the European W. acuta, but the capsules are larger, of a firm texture, retaining their shape when dry. In Dr. Lyall's specimens the leaves are more falcate and the seta shorter.

PLATE CLII. Fig. IV.—1, tufts of the natural size; 2, leaf; 3, young scta; 4, capsule; 5, the same before the fall of the calyptra :—all magnified.

5. WEISSIA *tortifolia*, Hock. fil. et Wils.; caule ramoso, foliis patentibus flexuosis siecitate crispatis lineari-subulatis canaliculatis integerrimis nervo excurrente, perichætialibus brevioribus convolutis, seta brevi, theca subrotunda, operculo rostrato. (TAB. CLII. fig. V.)

HAB. Kerguelen's Land, common on gravelly banks, from the sea to 1,000 feet.

Caules subunciales, dense cæspitosi, raunosi. Folia patentia, varie flexuosa, siccitate crispata vel tortuosa, nervo gracili excurrente instructa, lutescenti-viridia, inferiora fuscescentia; perichætialia breviora, ovata, acuminata, convoluta. Sela vix 2-linearis, erassiuscula, fusco-brunnea. Theca erecta, subrotunda, rufo-brunnea, vernicosa, demum indurata. Peristomii dentes 16, pyramidati, conniventes. Operculum conico-rostratum, capsulæ longitudine, rostro obliquo. Calyptra cucullata, capsulam æquans illamque obtegens, brunnea. Florescentia monoica.

Very closely allied to *Weissia stricta* (nobis), but differing in the crisped widely spreading leaves, which arc only half as long as in that species.

PLATE CLII. Fig. V.-1, tuft of the natural size; 2 and 3, leaves; 4 and 5, capsules :- all magnified.

13. DICRANUM, Hedw.

1. DICRANUM *aciphyllum*, Hook. fil. et Wils.; caule ramoso, foliis erecto-patentibus strictis rigidis lineari-lanceolatis integerrimis canaliculatis, nervo latissimo continuo, theca subcylindracea erecta, peristomii dentibus angustis subintegris, operculo longirostro. (TAB. CLII. fig. III.)

Var. 2. foliis secundis.

Var. 3. caule graciliore, foliis brevioribus siccitate subflexnosis, nervo tenuiore.

HAB. Staten-Land, A. Menzies, Esq., (1787). Hermite Island, Cape Horn, on rocks and on branches of trees on the hills, alt. 700 feet. Falkland Islands, on rocky ground among the hills, rare in fruit. Patch Cove, Cape Tres Montes, C. Darwin, Esq. Vars. 2 and 3, Hermite Island; var. 3 forming small tufts from the sea-side to the tops of the hills, alt. 1,740 feet.

Caules biunciales, parce ramosi, cæspitosi, siceitate parum fragiles. Folia erecto-patentia, vix secunda, siccitate erecta, lateribus inflexis, canaliculata, integerrima, luteo-viridia, nervo latissimo ultra laminam in acumen longum rigidum subulatum producto; perichætialia ovato-lanccolata, vaginantia, caulinis breviora. Seta uncialis, tortilis, fusco-lutea, superne pallida. Theca subcylindracea, erecta, subæqualis, basi attenuata, fusca, demum siccitate substriata. Peristomii dentes 16, breves, angusti, trabeculati, perforati, vix apice fissi, siccitate erecti, rubri. Sporæ minutæ, virides. Operculum conico-rostratum, thecæ longitudine, rostro obliquo. Calyptra luteola. Nearly allied to *D. longisetum*, Hook. (Musc. Exot. t. 139), but differs in the more robust habit, leaves longer, wider, more rigid, less setaceons above, without serratures, the nerve broader and thicker, capsule longer, peristome smaller, the teeth not divided to the base as in that moss.

PLATE CLII. Fig. III.—1, tuft of the natural size; 2, leaf; 3, perichaetial ditto; 4, theca; 5, calyptra; 6, theca and peristome; 7, teeth :—all magnified.

2 DICRANUM? imponens, Mont.; in Voy. au Pole Sud, Bot. Crypt. p. 298.

HAB. Strait of Magalhaens, MM. Hombron et Jaequinot. Hermite Island, Cape Horn; on the hills (barren).

Not having seen original specimens, it is necessary to mention that in the plant we refer to this species, the leaves are subsecund and entire; while in other respects they agree with the description quoted. *D. penicillatum*, Hornsch., to which Dr. Montague compares his moss, belongs to the genus *Campylopus*, Brid.

3. DICRANUM robustum, Hook.fil.et Wils.; caule elongato subramoso, foliis falcato-secundis longissimis lineari-lanceolatis setaceo-attenuatis convolutis spinuloso-serrulatis, nervo latiusculo excurrente, perichætialibus intimis obtusis enerviis, theca cylindraeea inelinata curvula strumulosa, operculo longirostro. D. scoparium, β. reflectens, Mont. in Voy. au Pole Sud, Bot. Crypt. p. 297? (TAB. CLII. fig. VIII.)

HAB. Hermite Island, Cape Horn; in woods, on the trunks and roots of trees, and in the open country, growing in large tufts, very abundant. Kerguelen's Land, Dr. Lyall.

Caules 4-unciales et ultra, robusti, parce ramosi, subinde incurvi, siccitate parum fragiles. Folia conferta, 6–7 lineas longa, apice setacea, inferne convoluta, superne carinata; margine dorsoque spinuloso-serrulata, viridia, inferiora squalida, perichætialia exteriora squarrosa, basi rotundato-ovata, acuminata, interiora erecta, convoluta, 3-linearia, elliptico-oblonga, obtusa cum mucrone lineari longiusculo, enervia. *Faginula* linearis, elongata. Seta 7–8 lin. longa, crassiuscula, siccitate vix tortilis, rubella. *Theca* cylindracea, inclinata, curvula, rufo-brunnea, strumulosa. *Peristomii* dentes bifidi, rubri. *Operculum* theca paulo longius. *Calyptra* straminea, apice fusca.

A larger and more robust moss than D. pungens, nobis, which it much resembles, differing in its longer leaves, which are more decidedly serrated, less convolute, nerve broader and excurrent, the perichætial ones much shorter, and wholly different in shape, capsule longer and strumose, peristome larger. In general aspect it is not unlike the British D. majus, Turn., but is distinguished by the very long and attenuated leaves.

PLATE CLII. Fig. VIII.-1, moss, natural size; 2 and 3, leaves; 4 and 5, thece:--all magnified.

4. DICRANUM pungens, Hook. fil. et Wils.; vid. Pt. 1. p. 129. t. 59. f. 1.

Var. 2. foliis vix secundis.

Var. 3. foliis minoribus, vix secundis.

Var. 4. lucidum; foliis aureo-nitentibus falcato-secundis.

HAB. Hermite Island, Cape Horn. Vars. 1 and 4 (barren) on rocks, trunks of trees, and the ground, very abundant. Kerguelen's Land; Vars. 2 and 4, Cumberland Bay, R. M'Cormick, Esq.

The var. 4 is a very elegant moss, differing in aspect from the other varieties, but we are unable to detect sufficient characters to establish it as a species.

5. DICRANUM Boryanum, Schwaegr., Suppl. II. vol.i. p. 71. t. 121. Cecalyphum dichotomum, P. Beauv. Prodr. p. 41. Oncophorus dichotomus, Brid. Br. Univ. vol. i. p. 401.

HAB. Kerguelen's Land, common on the hills in large dense tufts, barren.

Our specimens agree precisely with an original one from P. de Beauvois himself. The leaves are more falcate and wider at the base than those of *D. Billardieri*, to which this moss is very closely allied.

6. DICRANUM Billardieri, Schwaegr., Suppl. II. vol. i. p. 170. t. 121. Fl. Antarct. p. 119.

Var. caulibus apice ramosis, ramis brevibus confertis flagelliformibus.

HAB. Falkland Islands, amongst stones on Mount Vernet, not common, and always barren.

This peculiar condition of the moss is probably the result of its exposure to a dry atmosphere. Similar appearances occur in such British species as *Campylopus flexuosus*.

7. DICRANUM Starkii, Web. et Mohr, Bot. Tasch. p. 189, 471. Hook. et Tayl. Muse. Brit. t.xvii. p. 97. Var. 2. foliis vix secundis.

HAB. Hermite Island, Cape Horn, both varieties, the first in clefts of rocks, common, but rare in fruit. Thece smaller and more creet than in European specimens. Intermediate between the usual form and D. Sphagni, Wahl.

S. DICRANUM *tenuifolium*, Hook. fil. et Wils.; caule ramoso, foliis circinatim falcatis basi lanccolatis longissime capillaceis integerrimis nervo lato percursis, theca turbinata, operculo longirostro. (TAB. CLII. fig. VII.)

HAB. Hermite Island; moist shelving rocks on Mount Foster, above Deep-water Bay, scarce.

Caules unciales, parce ramosi, cæspitosi, curvuli, luridi. Folia longissima, angustissima, conferta, eleganter eircinato-falcata, lurido-viridia, inferiora atrata; perichætialia basi latiora, vaginantia. Seta 3-4 lineas metiens, crassinseula, pallide lutescens. Theca parva, subcrecta, turbinata, ore patulo, senior fusca. Perislomii dentes rubri. Operculum oblique longirostrum. Calyptra dimidiata, scariosa, fuseo-lutea.

Our specimens are not in a very good state, being too far advanced. Allied to the European *D. falcatum*, from which it differs in having narrower and longer leaves, and a very small nearly crect capsule destitute of a struma.

PLATE CLII. Fig. VII.-1, moss, natural size; 2 and 3, leaves; 4-6, thece; 7, peristome :--all magnified.

9. DICRANUM vaginatum, Hook., Musc. Exot. t. 141.

HAB. Hermite Island, on moist banks of sea-sand in St. Joachim's Bay, rare.

Our moss differs from original specimens gathered by Humboldt on the Andes of New Grenada, in the following particulars: teeth of the peristome broader; eapsule erect, short and turbinate; seta shorter and thicker; leaves entire, acute.

14. CAMPYLOPUS, Bridel.

1. CAMPYLOPUS introflexus, Bridel; Br. Univ. vol. i. p. 472. Fl. Antarct. pt. 1. p. 130. Dicranum introflexum, Hcdw. Sp. Musc. p. 147. t. 29.

HAB. Falkland Islands, common on the ground, especially in peaty situations. Also a native of Lord Auckland's group and Campbell's Island.

2. CAMPYLOPUS *flexuosus*, Bridel; *Br. Univ.* vol. i. p. 469. *Fl. Antaret.* l. c. Var. foliis piliferis. Dicranum clavatum, *Schwaegr. Suppl.* t. 255?

HAB. Amsterdam Island, South Indian Ocean, *Lieut. Smith*, R.N. Found also in Campbell's Island.

15. DIDYMODON, Hedw.

1. DIDYMODON capillaceus, Web. et Mohr, Bot. Tasch. p. 155. Hook. et Tayl. Musc. Brit. p. 119. t. 20. Swartzia capillacea, Hedw. Musc. Frond. vol. ii. p. 26.

HAB. Hermite Island, Cape Horn; on sandy banks near the sca, St. Joachim's Bay.

2. DIDYMODON longifolius. Trichostomum longifolium, Brid. Br. Univ. vol. i. p. 496. Trichostomum pallidum, β. strictum, Schwaegr. Suppl. II. vol. i. p. 77. t. 123?

Var. 2. tenuifolius; foliis basi magis dilatatis membranaceis, nervo duplo latiore.

Var. 3. *penicillatus*; eanle fastigiato-ramoso, foliis longioribus crectis subquadrifariam imbricatis, nervo erassiore.

Var. 4. curvifolius; foliis subfalcatis sceundis, e basi latiore sensim angustatis.

HAB. Staten Land, A. Menzies, Esq. (1787). Hermite Island, Cape Horn, vars. 1 and 4, abundant. South part of Fuegia, C. Darwin, Esq. Falkland Islands, var. 2, on the ground, rare in fruit; var. 3, on stones in streams, (barren).

From *Trichostomum pallidum* our moss differs essentially in the inflorescence, in the dilated base of the leaf, stronger nerve, cylindrical capsule, and also in the structure of the peristome. The var. 2 has at times an elliptical theca.

3. DIDYMODON? glacialis, Hook. fil. et Wils.; caule ramoso fastigiato-cæspitoso, foliis erecto-patentibus apice incurvis fragilibus ovato-lanceolatis lineari-acuminatis solidinerviis. (TAB. CLII. fig. VI.)

HAB. Cockburn Island, lat. 64° S. 57° W. (barren).

Caules 4 lin. longi, ramis erectis. *Folia* e basi concaviuscula, repente in acumen lineare producta, vel nervo crasso longe excurrente, margine haud reflexa, arcolis minimis, subrotundis, inferioribus majoribus pellucidis, e fusco lurido-viridia.

One of the only three mosses which have hitherto been detected in a higher latitude than that of Cape Horn.

PLATE CLII. Fig. VI.-1, tuft of the natural size; 2, branch; 3, 4 and 5, leaves :-- all magnified.

16. CERATODON, Brid.

1. CERATODON purpureus, Brid., Br. Univ. vol. i. p.480. Fl. Antarct. pt.1. p.131. Didymodon, Hook. et Tayl. Musc. Bot. p. 113. t. 20.

HAB. Strait of Magalhaens, *M. Jacquinot* (in D'Urville's Voyage). Falkland Islands, common on clay soil and on the sand-hills about Port Louis. Not seen on Hermite Island.

A very abundant Antarctic plant in many situations.

17. TORTULA, Hedw.

We retain this name, instead of *Barbula*, for the following reasons: Schreber is the first authority for the union of the two Hedwigian genera *Tortula* and *Barbula* in the year 1791 (Gen. Plant.). He adopted the name *Tortula* which stands first in Hedwig's arrangement for both. This fact is overlooked by Bruch and Schimper, who in their history

of this genus (Bryol. Europ.), rely chiefly on the authority of Bridel in support of *Barbula*. But Bridel's authority is in favour of *Tortula*, for he adopted it in his earlier work. It was discontinued by him in 1819 (*Mantissa*) under the erroneous impression that the name had been legitimately given to a phenogamous genus; whereas *Barbula* had been applied nine years before by Loureiro to designate a Chinese shrub. Hence Schrader, Sibthorpe, Swartz, Roth, and all British writers ou Mosses, retained *Tortula*, a name which would be at present unoccupied if not employed in conformity with the views of Hedwig and Schreber.

1. TORTULA *densifolia*, Hook. fil. et Wils.; caule humili subdiviso, foliis patentibus confertis lanccolatoacuminatis acutis marginatis apice serratis solidinerviis, theca oblonga, peristomii dentibus contortis, membrana basilari breviuscula, operculo subulato. (TAB. CLIII. fig. I.)

HAB. Falkland Islands, on clayey rocks near the sea at Port Louis, scarce.

Dioica? Caules 4-6 lin. longi, laxe cæspitosi, subdivisi. Folia dense conferta, basi erecta, dein patentia, stricta, lanccolato-subulata, acuta, subcarinata, rigidiuscula, margine cartilaginea vix incrassata, apice dentato-serrata, nervo valido rubello, lutescenti-viridia, areolis opacis minimis, basi majoribus subdiaphanis; perichætialia minora crecta. Seta 6-8 lin. longa, tortilis, fusca. Theca oblonga, erecta, fusca. Peristomii dentes basi membrana latius-cula conjuncti, rubelli. Operculum subulatum, capsulæ longitudine. Calyptra dimidiata, fusca.

The only described species with which this can be compared is *Barbula marginata*, Bruch and Schimp. (Bryol. Europ.); but that is a smaller moss, having leaves not at all acuminated, and an excurrent nerve.

PLATE CLIII. Fig. I.-1, tuft of the natural size; 2, leaf; 3, apex of ditto; 4, thece:-magnified.

2. TORTULA *robusta*, Hook. et Grev.; caule clongato subramoso, foliis patulo-recurvis lanceolatis subcarinatis acutis apice serratis solidinerviis, theca cylindracea curvula, peristomii dentibus contortis tubo ad tertiam partem producto, operculo subulato. T. robusta, *Hook. et Grev. in Brewst. Ed. Journ.* vol. i. p. 299. t. 12. (TAB. CLIII. fig. II.)

Var. β . foliis laxioribus viridibus.

HAB. Hermite Island, on sandy ground amongst grass at the head of St. Joachim's Bay, forming large patches. Var. β . Falklaud Islands, common in moist saudy places (barren).

Dioica, cæspitosa. Caules sesquiunciales, robusti, ramosinseuli. Folia lanceolata, vix acuminata, patentia, recurva, subcarinata, acuta, apice serrata, margine recurva, flavescentia, nervo tenui saturatius colorato percursa, siccitate erecta, incurva, subtortilia, areolis subrotundis, basi majoribus diaphanis, perichætialia similia erecta. Seta uncialis, sinistrorsum tortilis, rubella. Theca cybindracea, suberecta, curvula, rufo-fusca, ore rubello. Annulus persistens, albidus. Peristomii tubus basilaris dentium tertiam longitudinis partem æquans, albidus, dentes contorti, puchre rubelli. Operculum subulatum capsula dimidio longius, flavescens. Calyptra dimidiata, castanea.

From all the European Syntrichiæ this species is distinguished by the servated leaves. In size and general aspect it is not unlike Barbula Mulleri, Br. and Schimp.

After careful examination of the original specimens of *Tortula robusta* and *T. serrulata*, we have reason to believe that one, if not both, of these mosses is the same species as our Antarctic plants, differing only in the narrow leaves; but they are in too imperfect a state to determine very satisfactorily. The figure of *T. robusta* represents the leaves much too widely spreading, and they are also serrulate at the apex, just as in *T. serrulata*.

PLATE CLIII. Fig. I.—1, plant of the natural size; 2, leaf; 3, theca; 4, apex of theca and peristome :--all magnified.

3. TORTULA Mulleri. Barbula Mulleri, Bruch et Schimp. Bryol. Europ. fasc. 13-15. p. 44. t. 28.

HAB. Falkland Islands, on sand-hills near the Lagoon at Uranie Bay.

In our specimens the inflorescence is variable, even on the same stem; in some the antheridia and paraphyses are abundant, in others entirely wanting.

4. TORTULA lævipila, (Barbula), Bruch et Schimp. l. c. p. 40. t. 25.

Var. 1. foliis erecto-patentibus dorso margineque papillosis, florescentia monoica (interdum hermaphrodita).

Var. 2. foliis ovalibus suberectis dorso lævibus.

Var. 3. caule gracili ramoso, foliis brevioribus subcrectis elliptico-oblongis apice pilo brevi instructis margine subincurvis.

Var. 4. foliis obtusis areolis majusculis subrotundis.

HAB. Falkland Islands, vars. 1 and 2, on sandy soil near the sea; vars. 3 and 4, Cockburn Island, lat. 64° S., long. 57° W., both barren.

5. TORTULA gracilis. Barbula gracilis, Bruch et Schimp. l. c. p. 22. t. 8. (TAB. CLIII. fig. III.) HAB. Cockburn Island, (barren).

Our specimens differ from authentic examples in having the leaves more crowded, more pellucid at the base and less acuminated; the nerve also is wider.

PLATE CLIII. Fig. III.-1, tuft of the natural size; 2 and 3, leaves; 4, areolæ:-magnified.

6. TORTULA hyperborea, Mont. in Voy. au Pole Sud, Bot. Crypt. p. 302. t. 20. f. 4. Syntrichia hyperborea, Brid. Bryol. Univ. vol. i. p. 583. S. mucronifolia, Br. in Parry's 1st Voy. App. p. 198. excl. synonym.

HAB. Strait of Magalhaens; M. Jacquinot.

18. POLYTRICHUM, Linn.

1. POLYTRICHUM compressum, Hook. fil. et Wils.; caule subsimplici, foliis suberectis subulatis concavis subserratis, theca inclinata ovata compressa microstoma, operculo conico-rostrato, calyptra apice subpilosa. (TAB. CLIII. fig. IV.)

HAB. Hermite Island, in various situations, chiefly on wet rocks, from the sea to the tops of the hills, rare in fruit.

Dioieum. Caules plus minus dense cæspitosi, vix ramosi, nunc luridi, nunc rufo-ferruginei, subinde luteovirides. Folia erecto-patentia, imbricata, lanceolato-subulata; in caule masculo ovato-lanceolata, breviora; apice incurva, obtusiuscula, concava, mollia, subcarnosa, obscure serrata, nervo angusto haud lamellato instructa, areolis minutis subrotundis, opacis; perichætialia longiora, erecta. Seta uncialis, crassa. Theca inclinata, subinde horizontalis, ovata, obliqua, compressa, microstoma, hurido-fusca. Columella compressa. Peristomii dentes circiter 32, irregulares, albidi. Sporæ minutæ, ferrugineæ. Operculum conico-rostratum, capsula duplo brevius. Calyptra apice subpilosa, latere fissa, parvula, fusca.

Allied to the Icelandic P. lævigatum, Hook., but abundantly distinct in its compressed theca and narrower leaves.

PLATE CLIII. Fig. IV.—1, plant of the natural size; 2 and 3, leaves; 4 and 5, thece; 6, calyptra:—all magnified.

2. POLYTRICHUM juniperinum, Hedw., Sp. Musc. p. 89. t. 28. Hook. et Tayl. Musc. Brit. p. 45. t. 10. Var. foliis confertis subcrectis strictis.

HAB. Falkland Islands, on the moors, (barren). Hermite Island, Cape Horn, (barren). Evidently the British species, and a very widely dispersed one.

3. POLYTRICHUM alpestre, Hoppe; Bridel, Br. Univ. vol. ii. p. 140. P. juniperinum, var. Hook. et Tayl. l. c. p. 45.

HAB. Strait of Magalhaens, Port Famine, Capt. King.

This, which is also a British species, has been collected by Capt. King only.

4. POLYTRICHUM piliferum, Schreb.; Schwaegr. Suppl. I. vol. ii. p. 313. t. 153. Hook. et Tayl. l. c. p. 44. t. 10.

HAB. Falkland Islands, not uncommon on the moors, rare in fruit.

Affecting the same locality and habit in the Falklands that it does in England.

5. POLYTRICHUM deudroides, Schwaegr. Suppl. II. vol. ii. p. 2. t. 151.

HAB. Strait of Magalhaens, D'Urville. Hermite Island, in the woods, on steep banks by rivulcts, not uncommon, but rare in fruit.

In fertile specimens the branches are arranged at intervals around a common axis, the lower ones usually subdivided. The peristome has some analogy to that of *Lyellia*, the teeth being very small and the connecting base remarkably thick and prominent. The calyptra is quite glabrous. Columella apparently winged. In habit this moss belongs to *Pogonatum* of Bruch and Schimper, but its other characters do not correspond.

6. POLYTRICHUM squamosum, Hook. fil. ct Wils.; caule elongato squamoso fastigiato-ramoso, ramis patentibus brevibus densis curvatis, foliis dense imbricatis crecto-patentibus lanccolato-subulatis strictis serratis. (TAB. CLIII. fig. VIII.)

HAB. Hermite Island, on the hills at an altitude of 1,000-1,500 feet, scarce and scattered, always barren.

Caulis subspithameus, aseendens, firmus, subflexuosus, atro-purpureus, maxima ex parte humo sepultus, triqueter, superne squamis appressis luteo-fuscis scariosis nitidis in folia caulina gradatim abeuntibus vestitus, apice dendroideoramosus; rami vix semiunciales, subsimplices, patentes, fastigiati. *Folia* dense conferta, subcreeta, ad apices ramorum subsecunda, siecitate appressa, e basi membranacea pellucida semiamplexicauli lanceolato-subulata, vix trilinearia, superne lamellata, dorso carinaque scabra.

Although this moss is very different in aspect from P. dendroides, the characters seem searcely sufficient to distinguish it. The short straight leaves are only half as long as in that species.

PLATE CLIH. Fig. VIII.-1, plant of the natural size; 2 and 3, leaves :- magnified.

7. POLYTRICHUM Magellanicum, Hedw., Sp. Musc. p. 101. t. 20. Fl. Antarct. pt. 1. p. 132. t. bx.

HAB. Strait of Magalhaens; *D'Urville, Hombron*. Hermite Island, on fallen trunks of old trees, &c., in the woods; also on alpine rocks, in clefts. Falkland Islands, not uncommon on the ground and in clefts of quartz rocks on the hills, (always barren).

19. CONOSTOMUM, Sw.

1. CONOSTOMUM australe, Swartz, Schwacgr. Suppl. II. vol. i. p. 108. t. 130. Fl. Antarct. pt. 1. p. 182.

HAB. Hermite Island, on open rocky ground on the hills, occupying the same situations as *C. boreale* in Europe. Falkland Islands, in similar localities, bearing fruit in November.

Entircly the representative of the British and Arctic C. boreale. Also found in Lord Auckland's group.

FLORA ANTARCTICA.

20. BARTRAMIA, Hedw.

1. BARTRAMIA patens, Schwaegr., Suppl. I. vol. i. p. 55. t. 62. Fl. Antarct. pt. 1. p. 133.

Var. B. intermedia; caule minore.

HAB. Hermite Island, common in the woods, from the sea to the tops of the hills, in crevices of rocks, growing in dense soft tufts. Falkland Islands, common on wet clay-slate rocks near the sea, not found on the hills, abundant in fruit. Var. β . on clayey ground and rocks near the sea.

The variety β . differs only in its smaller size and in the general aspect, which resembles that of the British *B. ithyphylla*.

2. BARTRAMIA pendula, Hook.; Musc. Exot. t. 21. Fl. Antarct. pt. 1. p. 133.

Var. 1. foliis e basi crecta patulo-squarrosis, margine evidentius recurvis parcius denticulatis.

Var. 2. caule longiore robustiore vix tomentoso, foliis latioribus subsecundis.

ILAB. Hermite Island, in rocky places near the tops of the hills, scarce, abundant in fruit near the spray of a waterfall on the south side of St. Martin's Cove, forming large patches. Var. 2, always barren.

In some respects our moss resembles B. tomentosa, Hook. (Musc. Exot. t. 19), which we scarcely consider to be a distinct species, but the capsule is oblong and pendulous.

The var. 2 may be distinct. It resembles *Hypnum elongatum*, nobis. We have seen only the male inflorescence, which is truly that of *Bartramia*.

3. BARTRAMIA pomiformis, Hedw.; var. crispa. B. crispa, Swartz. Mont. in Voy. au Pole Sud, Bot. Crypt. p. 307.

HAB. Strait of Magalhaens; Admiral D'Urville et M. Jacquinot.

21. BRACHYMENIUM, Hook.

1. BRACHYMENIUM? *ovatum*, Hook. fil. et Wils.; caule humili, foliis laxe imbricatis erecto-patentibus quinquefariis ovato-oblongis nervo valido apiculatis. (TAB. CLIII. fig. IV.)

HAB. Falkland Islands, amongst dry quartz rocks on the hills, (barren).

Caules cæspitosi, inferne dense radiculosi, spongiosi, 1–2-unciales, parce ramosi. Folia erecto-patentia, 5-faria, ovato-oblonga, planiuscula, integerrima, luteo-viridia, siecitate subincurva, nervo valido excurrente apiculata, areolis minimis, rotundis. In axillis foliorum fasciculi corporum fascorum confervæ instar evadunt.

This moss resembles *Tetraphis pellucida*, but differs in the excurrent nerve and in the disposition and texture of the leaves.

PLATE CLIII. Fig. IV.-1, tuft of the natural size; 2 and 3, leaves :- magnified.

22. ORTHODONTIUM, Schwaegr.

1. ORTHODONTIUM *australe*, Hook. fil. et Wils.; caulc ramoso fastigiato humili, foliis erecto-patentibus subrecurvis anguste linearibus subflexuosis, nervo subcontinuo, theca suberceta oblonga brevicolla, operculo brevirostro. (TAB. CLIH. fig. V.)

HAB. Falkland Islands, alt. 900 feet, upon the fibrous roots of the Tussae grass; observed in one spot only. Hermite Island, in clefts of rocks on the hills and on wet banks, not uncommon.

FLORA ANTARCTICA.

Caules cæspitosi, 2-3 lin. longi, ramis brevibus apice coma incrassatis. Folia densa, e basi lineari longissime attennata, subflexuosa, carinata, integerrima, viridia, nervo conspicuo sub apicem evanido, cellulis elongatis: perichætialia longiora, similia, antheridiis in axillis eorum positis, paraphysibus paulo longioribus immixtis. Seta 4-5 lin. longa, gracilis, rubra, siccitate tortilis. Vaginula angusta, oblonga. Theca lanceolato-oblonga, inclinata, sub-

inde erecta, orc angustato, demum rufescens, siccitate substriata. Sporangium internum paulo brevius. Annulus obscurus, operculo adhærens. Operculum basi conicum, rostello brevi obliquo, interdum conicum, acuminatum, rectum. Peristomium breve; dentes externi hyalini, transverse trabeculati, siccitate inflexi; interni processus 16 longiores, carinati, linca media notati, membrana basilari connexi, siccitate erecti, subincurvi. Sporæ minimæ, luteæ. Calyptra latere fissa, pallida, apice brunnea.

This differs from Orthodontium lineare, Schwaegr. (Suppl. t. 188), in the oblong subcrect capsule, and in the absence of terminal male flowers, thus deviating from the generic character proposed by Schwaegrichen. In the inflorescence it agrees with the British Orthodontium gracile, Bruch and Schimper, but differs in the form of the capsule and the stronger nerve of the leaf.

PLATE CLIII. Fig. V .--- 1, plant of the natural size; 2, leaf; 3 and 4, thece; 5, peristome :-- all magnified.

23. BRYUM, Dill.

1. BRYUM nutans, Schreb.; Hook. et Tayl. Musc. Brit. p. 203. t. 29. Fl. Antarct. pt. 1. p. 134.

HAB. Falkland Islands; from the sea to the hill-tops, varying in size and habit. Hermite Island, Cape Horn; in clefts of rocks in the woods, and on hard soil by streams.

A common Antarctic moss; also a native of Lord Auckland's group.

2. BRYUM lacustre, Brid.; Bruch et Schimp. Bryol. Europ. Monogr. p. 16. t. 2.

HAB. Hermite Island; on sandy banks close to the brushwood a little above high water mark, St. Joachim's Bay.

3. BRYUM bimum, Schreb.; Bruch et Schimp. l. c. p. 50. t. 21.

HAB. Kergnelen's Land (barren).

4. BRYUM Billardieri, Schwaegr.; Suppl. I.vol.ii. p. 115. t. 76 (non Bruch et Schimp. l. c. p. 58. t. 26.)

HAB. Falkland Islands; on elay-slate rocks at Port Louis, rare and barren.

The European specimens, described by Bruch and Schimper, belong to B. Canariense, Schwaegr. (Suppl. t.214 b); we do not however contend for that moss being a really distinct species.

5. BRYUM argenteum, Linn.; Hook. et Tayl. Musc. Brit. p. 199. t. 29. Bruch et Schimp. l.c. p. 78. t.41.

Var. foliis arcte imbricatis augustioribus acuminatis.

HAB. Falkland Islands; in sandy places near the sea, common. Cockburn Island, (barren).

This variety is connected with the ordinary states of the species by intermediate forms which Mathews gathered at Casapi (Peru). The Hookerian Herbarium contains a *Bryum*, collected by Humboldt in South America, with muticous convolute leaves, allied to this, but probably a distinct species.

BRYUM cæspititium, Linn.; Hook.et Tayl. Musc. Brit. p.201.t. 29. Bruch et Schimp. l.c. p.70. t. 34.
 Var. β. gracilescens, Bruch et Schimp.

HAB. Strait of Magalhaens, *M. Jacquinot*. Falkland Islands, with unripe fruit. Var. *B*. Falkland Islands (barren).

5в

BRYUM pallescens, Schwaegr.; Suppl. I. vol. ii. p. 67. t. 74. Bruch et Schimp. l. e. p. 51. t. 22.
 HAB. Falkland Islands, on sand-hills at Uranie Bay. Hermite Island, Cape Horn.

8. BRYUM Antarcticum, Hook. fil. et Wils.; monoicum, caulibus ramosis cæspitosis, foliis confertis imbricatis erecto-patentibus ovatis acuminatis concavis reticulatis evanidinerviis margine planis. (TAB,CLIII. fig. VI.)

HAB. Cockburn Island, lat. 64° S., long. 57° W., with young setæ.

Caules 2 lin. longi, rubelli, inferne radiculosi. Folia late ovata, acumine brevi, apice diaphana, subdenticulata, rufescentia, nervo latiusculo rubello sub apice evanido, areolis subquadratis; perichætialia majora, acumine longiore. Seta vix $1\frac{1}{2}$ lin. longa, crassiuscula, rubra. Calyptra rubra. Flos masculas in ramulis brevibus per innovationes lateralis floribus fæmineis alternaus; antheridia cum paraphysibus longioribus subclavatis; archegonia paraphysibus brevioribus filiformibus immixta.

Nearly allied to the British *B. Zierii*, Dicks., but differing essentially in the inflorescence. The leaves are more crowded and have smaller areolæ.

PLATE CLIII. Fig. VI .-- 1, tuft of the natural size; 2, stem and 3, leaf, both magnified.

9. BRYUM Wahlenbergii, Schwaegr.; Bruch et Schimper, l. c. p. 44. t. 17., Fl. Ant. pt. 1. p. 134.

Var. 1. caule rubro, foliis ovatis rubellis.

Var. 2. foliis ovato-lanceolatis laxis viridibus.

Var. 3. caule elongato 2-3-unciali ramoso, foliis rubescentibus ovato-lanceolatis secundis.

Var. 4. foliis ovatis secundis minoribus nigro-viridibus.

HAB. Hermite Island, Cape Horn; var. 1, sand near the sea. Falkland Islands; var. 3, slate rocks near the sea (barren); Kerguclen's Land; var. 1, 2, and 4, all abundant.

The var. 1 is very similar to British specimens, differing in the colour of the leaves and in their being less acute. This moss also inhabits Lord Auckland's group.

10. BRYUM vagans, Hook. fil. et Wils.; caule vage ramoso, foliis patentibus secundis ovato-lanceolatis apiculatis submarginatis apice serratis, nervo subcontinuo. (TAB. CLIV. fig. 1.)

HAB. Hermite Island; marshy places in the woods, especially on slopes, frequent, (barren.)

Caulis basi procumbens, fere repens, biuncialis, apice ascendens, sæpe incurvus, vage ramosus, rami crecti, apice curvati. Folia laxe imbricata, patentia, secunda, mollia, ovato-lanceolata, apiculata, concaviuscula, submarginata; nempe areolis marginalibus angustioribus, confertis, cæteris majusculis, subrhomboideis, apice serrulata, basi haud decurrentia, nervo tenui subcontinuo instructa, juniora pallide viridia, vetustiora luteo-viridia. Flos masculus discoideus, terminalis; folia perigonialia late ovata, basi erecta, concava, superne patula, serrulata; antheridia numerosa, cylindracea, paraphysibus filiformibus immixta.

Allied to *B. Wahlenbergii*, but larger, the leaves twice as long, less succulent, not decurrent at the base, margined, apiculate, the nerve extending higher. During the winter months, owing to shifting of the watercourses, the banks on which this moss grows become inundated, and the varieties that ensue are very puzzling. The original plant is gradually covered by a carpet of young branches of a bright green colour, the whole forming a soft spongy and treacherous covering to the bogs.

PLATE CLIV. Fig. 1.-1, plant of the natural size; 2, leaf; 3, areolæ of ditto,-magnified.

414

11. BRYUM *Lævigatum*, Hook. fil. et Wils; caule ramoso, foliis erecto-patentibus imbricatis ovatis concavis integerrimis siccitate erectis, nervo subcontinuo, theca pendula ovato-oblonga, operculo conico obtuso. (TAB. CLIV. fig. III.).

Var. β . foliis angustioribus minus concavis.

HAB. Hermite Island; wet rocks on Mount Foster, (barren); Falkland Islands; common in bogs, (always barren.) Var. β. Kerguelen's Land (barren).

Dioicum? Caules unciales-triunciales, steriles longiores, inferne radiculis tomentosis nigricantibus intertexti, ramosi. Folia imbricata, subpatentia, ovata vel elliptico-oblonga, subcallosa, concava, integerrima, nervo valido vix sub apice evanido instructa, læte viridia, nitentia, inferiora e fusco-purpurascentia, siccitate subappressa. Seta 7-8 lin. longa, rufa, nitida, haud tortilis. Theca pendula, subpyriformis, oblonga, ore subpatulo, rufo, nitido. Peristomii externi dentes ferruginei, siccitate erecti; interni cilia perforata ciliolis singulis interjectis. Operculum breve, conicum, obtusum.

Our description and figure are drawn up in part from specimens gathered in Van Diemen's Land, by Mr. Lawrence. A very distinct species, characterised by its concave shining subcoriaceous leaves, retaining their shape when dry, intermixed with dark purple radicles.

PLATE CLIV. Fig. III.—1 and 3, stems of two states, from Hermite Island; 2, a third state, from Van Diemen's Land, of the natural size : 4 and 5, leaves; 6, thece;—magnified.

12. BRYUM truncorum, Bridel, Bryol. Univ. vol. 1. p. 699.

HAB. Falkland Islands; with fruit rare, Dr. Lyall; and a taller barren state in marshy places, not uncommon.

24. MNIUM, Bruch et Schimper.

1. MNIUM rostratum, Bruch et Schimper; Bryol. Europ. Monogr. p. 27. t. 7. Bryum rostratum, Hook. et Tayl. Musc. Brit. p. 208. t. xxx.

HAB. Strait of Magalhaens, Port Famine; Capt. King, (barren).

25. FUNARIA, Schreb.

1. FUNARIA hygrometrica, Hcdw.; Fl. Antarct. pt. 1. p. 135., Hook. et Tayl. Musc. Brit. p. 171. t. xx. HAB. Falkland Islands; common at Port Louis, on burnt ground. Not hitherto found in Fuegia, but a native of Campbell's Island.

26. ANŒCTANGIUM, Brid.

1. ANGECTANGIUM Humboldti, Brid. Hedwigia Humboldti, Hook. Musc. Exot. t. 137; Fl. Antarct. pt. 1. p. 135.

Var. β . australe.

HAB. Hermite Island; on a moist sloping rock exposed to the north, on Mount Foster, alt. 1000 ft.; in large barren patches resembling a discoloured mass of *Sphagnum*.

These specimens are intermediate between the typical form and that of Lord Auckland's and Campbell Island.

FLORA ANTARCTICA.

27. LEUCODON, Schwaegr.

1. LEUCODON Lagurus, Hook; Musc. Exot. t. 126. Fl. Antarct. pt. 1. p. 136.

HAB. Strait of Magalhaens; Port Famine; D'Urville et Jacquinot. Hermite Island; on trees in the forest and on rocks from the sea to an altitude of 1200 feet in large tufts.

Larger than the specimen figured in the 'Musci Exotici'; the capsules inclined, substrumose, the teeth of the peristome united regularly in pairs by transverse bars, pale yellow, leaves nerved half-way.

28. LESKIA, Hedwig.

1. LESKIA *nitida*, Hook. fil. et Wils.; caule vage ramoso, ramis longiusculis subsimplicibus teretibus, foliis imbricatis subcrectis ovato-oblongis acuminatis concavis integerrimis basi binerviis, seta lævi, theca eylindracea subcrecta curvula, operculo brevirostri. (TAB. CLIV. fig. VI).

HAB. Staten Land; A. Menzies, Esq. (1787). Hermite Island; Cape Horn; barren.

Caules sesquiunciales, steriles longiores, molles, virides; rami teretes, filiformes, apice e foliis convolutis cuspidati. Folia dense imbricata, erecto-patentia, ovato-oblonga, acuminata, acumine vix tertiam partem folii æquante, concava, subconvoluta, integerrima, nervis basilaribus duobus instructa, lutescenti-viridia, sericeo-nitentia, tenuissime elongato-areolata; perichætialia longiora, subsquarrosa. Seta uncialis, tortilis, gracilis, rubra. Theca cylindracea, subcrecta, curvula, interdum subcernua, basi attenuata, brunnea. Operculum basi conicum, rostello obliquo, capsula dimidio brevius, badium. Annulus operculo adherens. Peristomii externi dentes lutei, linea media notati, acuminati; interni processus breviores, angusti, carinati, ciliolis nullis.

This moss has cousiderable resemblance to *Hypnum stramineum*, but differs in having the leaves almost piliferous, in the rostrate operculum and in the structure of the peristome. It is also allied to *Hypnum crinitum*, nobis, from Van Diemen's Land.

29. HYPNUM, Dill.

a. Foliis distichis.

1. HYPNUM *politum*, Hook. fil et Wils.; caule ramoso compresso, foliis distichis patentibus oblongis compresso-carinatis subpiliferis integerrimis enerviis, seta lævi, theca suberecta oblonga. (TAB.CLIV. fig. II.)

HAB. Hermite Island; common in woods near the sea. Kerguelen's Land; in rocky places (barren).

Caules unciales et ultra, cœspitosi, subramosi, complanati, distiche ramosi; rami compressi. Folia arcte imbricata, patentia, disticha, elliptico-oblonga, scaphæformia vel compresso-carinata, apice cucullata, subpilifera, integerrima, enervia, læte viridia, sericco-nitentia, tenuissime arcolata; perichætialia ovata, longe acuminata, erecta, integerrima, caulinis duplo breviora. Seta vix uncialis, læv¹s, rufo-fusca. Theca oblonga, subcrecta, sub-apophysata, ore patulo. Peristomii externi dentes lutci, incurvi, linea media notati; interni cilia ciliolis interpositis. Calyptra dimidiata, straminea. Operculum non visum.

A beautiful species, unlike any hitherto described.

PLATE CLIV. Fig. II.-Specimen of the natural size : 2 and 3, leaves ; 4, thecæ; 5, peristome ; all magnified.

2. HYPNUM denticulatum, Dill. Linn.; Hedw. Musc. Frond. vol. 4. t. 31. Hook. et Tayl. Musc. Brit. p. 153. t. xxiv.

HAB. Hermite Island; on moist banks, wet rocks, &c., not uncommon, (barren).

3. HYPNUM reticulatum, Hook. fil. et Wils.; caule erecto simpliciusculo, foliis distichis patentibus ovato-lanceolatis acuminatis vix piliferis submarginatis reticulatis apice serrulatis, nervo tenui subexcurrente. (TAB. CLIV. Fig. V.)

HAB. Hermite Island; on the ground in damp woods, rare (barren).

Caules laxe cæspitosi, unciales, subelongati, erecti, plerumque simplices, complanati, molles. *Folia* disticha, patentia, subobliqua, acutissima, fere pilifera, areolis marginalibus angustioribus confertis, cæteris majusculis sub-rhomboideis, recentiora læte viridia, nitentia.

The many points of correspondence between this moss and *Bryum vagans*, nobis, have not escaped our notice. The specimens being few and barren, we are unable to pronounce with confidence on the validity of the species. It differs from *H. subbasilare* in the acuminated distichous leaves and almost excurrent nerve. In habit it much resembles *H. denticulatum*.

PLATE CLIV. Fig. V.-1, plant of the natural size; 2, leaf; 3, apex of ditto :- magnified.

4. HYPNUM riparium, Dill. Liun.; Hedw. Musc. Frond. vol. 4. t. 3. Hook. et Tayl. Musc. Brit. p. 152. t. xxiv.

Var. 2. caule elongato rigido, foliis dissitis minoribus rigidulis.

HAB. Kerguelen's Land; both varieties, in the lake near Christmas Harbour (barren).

With the habit of *Fontinalis*; probably a distinct species, but the specimens are not in a state to be determined satisfactorily.

b. Foliis imbricatis, seta radicali.

5. HYPNUM mnioides, Hook.; Musc. Exot. t. 77. Mont. in Voy. au Pole Sud, Bot. Crypt. p. 329. H. subbasilare (ex errore), Schwaegr. Suppl. t. 256.

HAB. Strait of Magalhaens, D'Urville. Hermite Island; abundant everywhere in the woods. Closely allied to *H. spiniforme*, from which it chiefly differs in its broader leaves.

6. HYPNUM subbasilare, Hook.; Musc. Exot. t. 10. H. mnioides (ex errore), Schwaegr. Suppl. t. 257.

HAB. Hermite Island; in moist woods, at the roots of trees, very common, growing in tufts.

This species so much resembles *H. mnioides*, as to be scarcely distinguishable from it until gathered,

In our specimens the leaves are by no means bifarious, as stated in the *Musci Exotici*, and are compressed only when dry. The perichætial leaves are erect, almost piliferous, and nerved to the apex; young calyptra coriaceous, slightly ventricose, not subulate, at length dimidiate. The operculum is absent from all our specimens.

c. Foliis imbricatis ruptinerviis, seta laterali.

7. HYPNUM rutabulum, Dill. Linn.; Hedw. Musc. Frond. vol. 4. t. 12. Hook. et Tayl. Musc. Brit. p. 176. t. xxvi. Fl. Antarct. pt. 1. p. 138.

Var. 1. foliis apice attenuatis, perichatialibus erectis.

Var. 2. foliis majoribus læte viridibus.

Var. 3. caulibus 2-3-pollicaribus, foliis angustioribus luteo-viridibus nitentibus inferioribus fuscis.

Var. 4. caule clongato graciliore.

Var. 5. caule elongato, foliis subcirrhosis brevinerviis.

HAR. Hermite Island, Cape Horn; var. 1. wet rocks in the woods; var. 2, roots of trees (barren). Falkland Islands; var. 5. springy places, forming large green masses. Kerguelen's Land; var. 3. wet places on the hills (barren); var. 4. wet bogs (barren).

The second of these varieties resembles very closely the European plant.

8. HYPNUM *subpilosum*, Hook. fil. et Wils.; caule fastigiato-ramoso, foliis cordato-ovatis imbricatis suberectis acuminatis subpiliferis concavis striatis serrulatis ruptinerviis. (TAB. CLIV. Fig. IV.)

IIAB. Hermite Island, Cape Horn; in moist earth, near the tops of the hills, altitude 1500 feet.

Monoicum. *Caules* laxe cæspitosi, sesquiunciales, vage ramosi, subfastigiati; rami patentes, subrecurvi. *Folia* arete imbricata, suberecta, cordato-ovata, repente acuminata, acumine fere piliformi, reflexiuscula, concava, subplicata, serrulata, nervo crasso medio exarata, læte viridia, iuferiora squalida, areolis angustis. *Seta* inferne lævis, superne scabriuscula. (Cætera desunt).

Closely allied to *H. rutabulum*, but smaller and more rigid, the leaves more closely imbricated and almost piliferous.

PLATE CLIV. Fig. IV.-1, plant of the natural size; 2, leaf :- magnified.

9. HYPNUM albicans, Dill. Neck.; Hedw. Musc. Frond. vol. 4. t. 5. Hook. et Tayl. Musc. Brit. p. 167. t. xxv.

Var. caule elongato, foliis luteo-viridibus.

HAB. Hermite Island; in moist places and streams, not uncommon (barren).

This resembles Var. 4. of *II. rutabulum*, but has more distinctly striated leaves.

10. HYPNUM serpens, Dill. Linn.; Hedw. Musc. Frond. vol. iv. t. 18. Hook et Tayl. Musc. Brit. p. 153. t. xxiv.

Var. 1. foliis subsecundis subsolidinerviis.

Var. 2. foliis ovatis brevioribus latioribus.

HAB. Kerguclen's Land; both varieties on the rhizomata of the "Cabbage", Pringlea, (barren).

d. Foliis imbricatis subencrviis, seta laterali.

11. HYPNUM chlamydophyllum, Hook. fil. et Wils.; Fl. Antarct. pt. 1. p. 139. t. lix. fig. i.

HAB. Hermite Island; moist rocks on the hill-tops, altitude 1400 ft., rare.

Also a native of Tasmania and Campbell's Island.

12. HYPNUM auriculatum; Montagne in Voy. au Pole Sud, Bot. Crypt. p. 331. t. 20. f. 3. HAB. Strait of Magalhaens, M. Jacquinot.

13. HYPNUM *lucidulum*, Hook. fil. et Wils.; caule ramoso humili, foliis erecto-patentibus ovato-acuminatis apice attenuatis integerrimis margine reflexis basi 1-2-nerviis, seta lævi, theca cernua ovato-oblonga. (TAB. CLV. Fig. I.) HAB. Hermite Island; on banks and moist rocks in the evergreen beech-woods, abundant; also in crevices of rocks on the hills.

Caules seminneiales et ultra, cæspitosi, ramosi, molles, fragiles. Folia conferta, imbricata, erecto-patentia vix secunda, ovato-acuminata, concavinscula, apice attennata, subpilifera, tenera, margine reflexa, basi 1-2-nervia, pallide viridia, nitida, tennissime arcolata; perichætialia ovata, breviter acuminata, erecta, integerrima. Seta 7-8 lin. longa, kevis, apice incurvata, rubra. Theca ovato-oblonga, basi attenuata, enrvula, e setæ apice curvata cernna, brunnea. Peristomium externum luteum, internum albescens, ciliolis binis.

This moss bears some resemblance to *H. Silesianum*, Schwaegr., but is quite distinct in character, and allied also to *H. adnatum*, Hedw., from which it differs in the flaceid leaves, recurved at the margin.

PLATE CLV. Fig. I.-1, plant of the natural size; 2 and 3, leaves; 4 and 5, thece :- magnified.

e. Foliis patulis squarrosis.

14. HYPNUM aciculare, Brid.; Schwaegr. Suppl. I. vol. ii. p. 280. t. 92. Fl. Antarct. pt. 1. p. 140.

HAB. Staten Land, A. Menzies, Esq. Hermite Island; in woods near the sea, not uncommon.

A very abundant plant in the South temperate and colder regions.

f. Foliis secundis nervosis.

15. HYPNUM conspissatum, Hook. fil. et Wils.; caule elongato ramoso, foliis patentibus secundis ovatolanccolatis acuminatis integerrimis margine incrassatis solidinerviis. (TAB. CLV. Fig. III.)

Var. 2. caule longiore, foliis latioribus.

Var. 3. foliis longioribus magis acuminatis.

HAB. Kerguelen's Land; var. 1, boggy places, common (barren). Falkland Islands; var. 2, (barren), Dr. Lyall; var. 3, growing in waters (barren).

Caules fluitantes, biunciales ad semipedales, ramosi, fastigiati ; *rami* simplices, ascendentes. *Folia* laxe imbricata, plus minus falcato-seconda, rigidula, crassiuscula, opaca, margine valde incrassata, nervo continuo exarata, juniora intense viridia, cætera lurido-viridia, interdum fuscescentia, inferiora nisi nervus margoque incrassata plerumque tabescentia. (Cætera desunt.)

In the leaves this moss has a close affinity with Cinclidotus, but the aspect is that of Hypnum ruscifolium.

PLATE CLV. Fig. III.-1, plant of the natural size; 2 and 3, leaves; 4, apex of ditto :--magnified.

16. HYPNUM filicinum, Dill. Linn.; Hedw. Sp. Musc. p. 258. t. 76. Hook. et Tayl. Musc. Brit. p. 183. t. xxvi. Fl. Antarct. pt. 1. p. 141.

Var. 2. robustius, foliis elliptico-lanceolatis angustioribus.

Var. 3. omnia var. 2, sed foliis vix secundis.

Var. 4. foliis vix secundis latioribus erectis acuminatis.

HAB. Falkland Islands; var. 1, rocky fresh-water streams, and wet sandy places; var. 3, *Dr. Lyall.* Hermite Island; Cape Horn; var. 4, wet rocks and sandy places. Kerguelen's Land; var. 2, boggy places. (barren).

These varieties are all very similar to the plant mentioned in the first part of this work.

17. HYPNUM *paradoxum*, Hook. fil et Wils.; caule repente subpinnato, foliis falcato-secundis ovatolanceolatis acuminatis striatis serrulatis ruptinerviis, seta scabra, theca cernua obovato-oblonga. (TAB. CLV. Fig. II.) Var. β . foliis laxe imbricatis substriatis, theca ovata, operculo conico.

HAB. Hermite Island, Cape Horn; var. β . on moist rocks and at the roots of trees; scarce.

Caules 2-3-unciales, rami ascendentes. Folia falcato-secunda, plicato-striata, serrulata, nervo ultra medium producto, luteo-fusca, subspadicea, in var. β . læte viridia. Seta semiuncialis, brunnea. Theca horizontalis, turgida, sub ore contracta, rufa. Peristomium externum ferrugineum, internum flavum.

This moss resembles H. aduncum in everything but the scabrous seta, and the serrulate leaves; the var. β again approaches very nearly to some varieties of H. velutinum.

PLATE CLV. Fig. II.-1, plant of the natural size; 2 and 3, leaves; 4 and 5, thece; 6, peristome :- magnified.

18. HYPNUM fluitans, Linn.; Hedw. Musc. Frond. vol. iv. p. 36. Fl. Antarct. pt. 1. p. 141.

HAB. Hermite Island, Cape Horn, in bogs, very common by the margins of mountain lakes. In fruit amongst wet stones.

Also found, but barren, in Campbell's Island.

19. HYPNUM aduncum, Dill. Linn.; Hedw. Musc. Frond. vol. iv. t. 24. Hook. et Tayl. Musc. Brit. p. 186. t. 26.

Var. c. revolvens, Bridel, Bryol. Univ. Hook. et Tayl. l. c.

HAB. Hermite Island, Cape Horn, (barren).

20. HYPNUM falcatum, Bridel, Bryol. Univ. vol. ii. p. 526. Schwaegr. Suppl. II. vol. i. p. 162. t. 145.

HAB. Falkland Islands, common in watery places, rare in fruit.

Apparently the moss mentioned by Gaudichaud under the name of *H. aduncum*, to which indeed it is nearly allied. It differs from *H. fluitans*, in its very strong, often percurrent nerve. The leaves vary in length.

21. HYPNUM uncinatum, IIall.; Hedw. Musc. Frond. vol. iv. t. 25. Hook. et Tayl. Musc. Brit. p. 187. t. xxvi.

HAB. Kerguelen's Land, in bogs, not uncommon (barren). Hermite Island, Cape Horn, also in wet places.

g. Fotiis secundis enerviis.

22. HYPNUM lithophilum, Hornschuch; ramis elongatis, foliis erecto-patentibus secundis siccitate erectis ellipticis acutiusculis (rameis elliptico-oblongis obtusiusculis) concavis margine reflexis tenuissime lineari-areolatis, perichætialibus erectis. H. lithophilum, Hornschuch, in Endlich. et Mart. Flora Brasil. p. 84, in part.

HAB. Hermite Island, Cape Horn; with H. amanum (barren).

In the Hookerian Herbarium we find a Brazilian specimen, apparently authentic, of *H. lithophilum*, labelled "supra lapides rivulorum in novo Friburgo", but consisting of two different species mixed together; and as the description in Fl. Bras. appears to have been drawn up from both, we append a diagnosis of the other species.*

* HYPNUM succedaneum, nobis; foliis laxe imbricatis sicco ac humido pariter patulis subsquarrosis late ovatis concaviusculis margine subreflexis, areolis paralellogrammis, perichætialibus longioribus apice subsquarrosis.

HAB. Brazil, Province of Rio, New Friburg, Martius.

FLORA ANTARCTICA.

Our specimens agree with H. lithophilum, except that their branches are more elongated and the leaves faintly two-nerved at the base. A Brazilian specimen from Raddi corresponds with the H. succedaneum, which is allied to H. molle.

23. HYPNUM micans, Wils.; in Hook. Brit. Flora, v. 2. p. 83. Engl. Bot. Suppl.

Var. laxum, foliis secundis patentibus ellipticis concavis serrulatis enerviis.

HAB. Hermite Island; in moist places in the woods, and on the hills.

Caules semiunciales, prostrati, parce ramosi; rami graciles.

This variety is allied to *H. gracile*, nobis, but differs in the reticulation of the leaves, and in their margins being recurved at the base.

24. HYPNUM amænum, Hedw., Sp. Musc. p. 292. t. 77. Isothecium amænum, Brid. Bryol. Univ. vol. ii. p. 382.

HAB. Hermite Islaud; on wet rocks by streams of water, generally in woods, growing in patches, rare in fruit.

Allied on the one hand to *H. tenuirostre*, Hook. (Musc. Brit.), and on the other to *H. leptorhynchum*, Schwaegr. From the first of these it differs in the longer more attenuated circinate leaves; from the latter in its larger size and in the elliptical shape of the lower part of the leaf, which is not reflexed at the margin. The operculum, as in those species, has a long slender beak and is somewhat longer than the capsule.

25. HYPNUM leptorhynchum, Brid.; Schaegr. Suppl. I. v. 2. p. 295. t. 93. Fl. Antaret. pt. 1. p. 140.

HAB. Hermite Island; very common on the rocks and banks, and on trunks of trees, taking the place of *H. cupressiforme*, which, strange to say, has not hitherto been found in any part of Fuegia or the Falkland Islands.

30. HOOKERIA, Sw.

a. Foliis marginatis enerviis.

1. HOOKERIA apiculata, Hook. fil. et Wils.; caule compresso subramoso, foliis distiche imbricatis rotundatis apiculatis marginatis enerviis siccitate undulatis, seta scabriuscula, capsula cernua, calyptra pilosa. (TAB. CLV. fig. VI.)

HAB. Hermite Island; ou moist shady rocks near the sea (barren), forming green tufts.

Caules steriles uneiales, erecti, densius cæspitosi, parce ramosi, inferne radiculis nigris obsiti, ramis erectis compressis; fertiles procumbentes, humiles, vix semiuneiales. Folia laxe imbrieata, lateralia patentia, eætera appressa, rotundato-ovata, apiculata, rigidiuseula, marginata, euervia, siccitate paulo undulata, apice subinde denticulata, areolis majusculis hexagonis; perichætialia erecta, minora, ovato-lanceolata, acuta. Seta seabriuscula, 2–3 lin. longa, flexuosa. Capsula eernua vel horizontalis, ovata, subapophysata. Operculum basi hemisphærico-conicum, rostratum, capsula paulo brevius, rostro reeto. Calyptra parva, pilosa, albida. Florescentia dioica.

Allied to *Hookeria asplenioides*, Schwaegr., but smaller, and having the margin of the leaves thickened and undulated when dry. Described from fertile specimens, gathered on the bark of trees, in Tasmania, by Mr. Gunn.

PLATE CLV. Fig. VI.-1, tuft of the natural size ; 2 and 3, leaves :- magnified.

b. Foliis marginatis evanidinerviis.

2. HOOKERIA Dicksoni, Hook. in Brewst. Edinb. Journ. of Science, vol. 2. p. 226.

HAB. Falkland Islands; on shady clay-banks near the sea, at Port Louis (barren). Hermite Island; common on mossy banks and on the trunks of old trees in the woods of evergreen beech, abundant in fruit.

Very closely allied to *Hookeria pulchella*, nobis (part 1. p. 142. t. lxii); but the leaves are more erect, less crowded, acuminated, with larger reticulations, thece larger and decidedly cernuous. The calyptra in both these species is fringed at the base.

3. HOOKERIA *flaccida*, Hook. fil. et Wils.; caule debili elongato erecto subramoso, foliis imbricatis erecto-patentibus ellipticis concavis obtusis subapiculatis integerrimis anguste marginatis evanidinerviis, seta clongata lævi, theca erecta obovato-oblonga, operculo rostrato, calyptra basi fimbriata. (TAB. CLV. fig. V.)

HAB. Hermite Island; in wet bogs on the hills, amongst other mosses and grass, very rare in fruit.

Caules unciales ad triunciales, graciles, debiles, parce subpinnatim ramosi, rufo-fusci, ramis compressiusculis. Folia laxe imbricata, erecto-patentia, flaccida, elliptico-oblonga, concava, obtusa, brevissime apiculata, inferiora subobovata, omnia integerrima, margine tenui cartilagineo nervoque tenuissimo sub apice evanido instructa, sordide ac pallide viridia, siccitate crispata, areolis parvulis rotundatis; perichætialia triplo minora, ovata, enervia. Seta uncialis, vix tortilis, rubra. Theca erecta, obovato-oblonga, brunnea, subapophysata, ore subpatulo. Peristomii externi dentes lutei, incurvi, trabeculati, linea media notati, interni processus albidi. Sporæ minimæ, luteo-virides. Opercutum conico-acuminatum, theca paulo brevius. Calyptra elongato-conica, acuminata, basi fimbriata, fusca, capsulæ dimidiam partem obtegens.

 Λ remarkably soft and delicate species, bearing much the same analogy to its congeners that *Hypnum strami*neum does to other *Hypna*.

4. HOOKERIA Magellanica, P. Beauv.; caule ramoso erecto, foliis ovato-oblongis acuminatis marginatis evanidinerviis, calyptra basi fimbriata.

HYPNUM Magellanicum, P. Beauv. Æthcog. p. 66.

HAB. Strait of Magalhaens.

An authentic specimen in Professor Arnott's Herbarium is closely allied to *Hookeria flaccida*, nobis. It differs in having narrower acuminated leaves, which do not fully recover their shape after long immersion in water.

c. Foliis emarginatis.

5. HOOKERIA deuticulata, nobis; vid. Pt. I. l. c. 145. tab. lxii. f. 2.

HAB. Falkland Islands; in tufts of *Riccia* and *Jungermannia*, on rocks near the sea, frequent (barren). Hermite Island, Cape Horn; on the wet ground in woods, not uncommon (also barren).

6. HOOKERIA cristata, Hedw.; Sp. Musc. p. 211. t. 49. Schwaegr. Suppl. t. 278. A.B.

HAB. Hermite Island, Cape Horn.

A solitary barren stem of this occurs in the collection of Hermite Island plants.

31. HYPOPTERYGIUM, Bridel.

Our reasons for not having previously admitted this genus will be found in the former portion of this work. We have seen since, that the male flowers are occasionally, though rarely, inserted beneath the accessory leaves, and

therefore we retain this name for a genus which certainly claims to be separated as well from Leskia as from Hookeria.

1. HYPOPTERYGIUM laricinum, Bridel; Bryol. Univ. v. 2. p. 714. Hypnum laricinum, Hook. Musc. Exot. t. 35. Hypnum tamariscinum, Swartz !

HAB. Hermite Island; in wet places on the ground, very common in the woods, forming large green patches (always barren).

Under Leskia tamariscina two species have been confounded by Hedwig (Sp. Musc. p. 212). The name ought to be applied to the present moss, if the inconvenience of changing names generally received did not forbid.

2. HYPOPTERYGIUM Thouini, Schwaegr.; Suppl. t. 289 (sub nom. Hypnum). Hypnum Arbuscula, P. Beauv. Ætheog. p. 61! Hypopterygium Thouini, Montagne in Ann. Sc. Nat., Aug. 1845, p. 86.

HAB. Strait of Magalhaens; Port Famine, Capt. King.

Our specimens are not so large as those described by P. de Beauvois, though evidently belonging to the same species. Dr. Montagne has properly remarked that this species differs from *H. laricinum* in the flabelliform, not pinnate, disposition of its branches, which all spring from one central point and take a horizontal direction. Fertile specimens from Colchagua, in Chili, have also a more pendulous oblong capsule and shorter operculum.

ORD. LIII. HEPATICÆ, Juss.

(By Dr. THOMAS TAYLOR and J. D. HOOKER.)

1. JUNGERMANNIA, L.

(1. GYMNOMITRION, Nees.)

1. JUNGERMANNIA *physocaula*, Hook. fil. et Tayl.; caule gracili disperso suberecto ramoso cellulosotumente, ramis apice curvatis incrassatis, foliis laxe cellulosis imbricatis distichis concavis oblique erectis late ovatis quadrato-rotundatisve ad medium bifidis segmentis late subulatis integerrimis. Nobis in *Lond*. *Journ. Bot.* v. 3. p. 455. (TAB. CLVI. Fig. I.)

HAB. Hermite Island, Cape Horn; creeping through tufts of J. densifolia, Hook.

Caules 1-2 unc. longi, graciles, vage parce ramosi; rami solitarii v. bi-terni, pallide olivacei v. albidi, nunc rufobrunnei, apice curvati. Folia tumida, arcte imbricata, cauli appressa; segmentis formæ subvariis, integerrimis. Stipulæ nullæ.

Allied to the Scottish *J. concinnata*, Lightf.; but readily distinguishable by the stems not being tufted, the shoots slender and flexile, the larger more cellular leaves, which are far more deeply divided, and have lanceolate segments, and by the cellular stem.

PLATE CLVI. Fig. I.-1, plant of the natural size; 2, portion of stem : 3, leaf :- magnified.

2. JUNGERMANNIA atrocapilla, Hook. fil. et Tayl.; caule tenuissimo procumbente implexo parce ramoso flexuoso basi longe nudo, foliis remotis erectis cauli appressis concavis late ovato-quadratis integris eroso-

FLORA ANTARCTICA.

Fuegia, the

emarginatis, perichætialibus majoribus imbricatis in capitulum clavatum congestis. Nobis in Lond. Journ. of Bot. vol. 5. p. 258.

HAB. Foul Haven, Kerguelen's Land; on clay banks, at an elevation of 600 ft.

Cæspites extensi, 1-2 une. lati, valde inconspicui. *Caules* atri, diametro setæ equinæ, crassiusculi, subnudi, basi hie illic cicatricati, superne foliis parvis tumidis appressis remotis quasi nodosi, fertiles apices versus foliosi, e foliis perichætialibus gradatim majoribus arcteque imbricatis clavati. *Folia* late quadrata, supra medium apicibusque erosis pallida.

A remarkably distinct little species, forming very obscure black patches on the ground. Stems wiry when dry, and loosely tufted; those of the perichætium paler and olive-brown, having their apiees twice as broad as any other parts of the shoot. Perichætial leaves more imbricated, rounder, broader, and more concave than the cauline, enclosing a pair of minute whitish connivent seariose scales, but without any trace of ealyptra or barren pistilla.

(2. GOTTSCHEA, Nees.)

3. JUNGERMANNIA lamellata, Hook.; Muse. Exot. t. 49. Gottsche, Lind. et Nees; Syn. Hep. p. 30. HAB. Staten Land, Menzies. Hermite Island, Cape Horn, in dense woods abundant.

This beautiful species is apparently peculiar to the southern extreme of the American continent.

4. JUNGERMANNIA leucophylla, Lehm. MS. Gottsche, Lind. et Nees; Syn. Hep. p. 17.

HAB. Strait of Magalhaens; Commerson (in Hb. Reg. Berol.).

5. JUNGERMANNIA splachnophylla, Hook. fil. et Tayl.; caule crasso subdisperso procumbente simplici recurvo e foliis complicatis densissimeque imbricatis squamoso dorso fibrillis squamisque densissime obsito, foliis erecto-patentibus undulato-complicatis carnosis marginibus sub-erosis, lobo ventrali oblongo-ovato, dorsali subæquali semi-ovato, ala lineari undulata. Nobis *in Lond. Journ. Bot.* vol. iii. p. 455. (TAB. CLVI. Fig. II.)

HAB. Hermite Island, Cape Horn; on the ground amongst underwood, alt. 800-1000 ft., and thence to the hill-tops.

Caules 2-unc. longi, erassi, terræ appressi, sub $\frac{1}{4}$ -une. lati, dorso densissime filamentosi, substuposi. Folia sordide alba, densissime imbricata et complicata, carnosa et aquosa, fragilissima, marginibus hine sæpissime erosis, paulo incurvis, basi sese arcte auplectentia.

A very singular plant, differing in its earnose texture from all the previously described species. The leaves are so thick, brittle, and watery as to be erushed to pieces readily between the finger and thumb, whence the analysis of the dried specimeus is extremely difficult.

TAB. CLVI. Fig. II:---1, plant of the natural size : 2, front, and 3, back view of leaf; 4, leaf from lower portion of stem :---magnified.

6. JUNGERMANNIA pachyla, Hook. fil. et Tayl.; caule cæspitoso erecto subramoso ramisque apice incurvis anguste linearibus, foliis inflatis dense imbricatis crecto-patentibus, lobis ovato-oblongis acuminatis apicibus incurvis, dorsali integerrimo dorso convexo, ventrali undulato horizontali margine anteriore basi dentato, ala anguste lineari, stipulis majoribus late ovato-quadratis bifidis segmentis lanceolatis apice incisis. Nobis in Lond. Journ. Bot. vol. iii. p. 456. (TAB. CLVI. Fig. III.)

HAB. Hermite Island, Cape Horn; on the bare ground in wet places.

424

Cæspites extensi, laxi, luride rufo-brunnei. *Caulis* creetus, parce ramosus, dorso radiculis fibrillosis purpureis per totam longitudinem instructus, apicibus subcurvatis. *Folia* latiuscula, concava, patentia, marginibus undulatis, erosis. *Stipularum* laciniæ sæpius inæquales.

One of the more slender species of the genus, with the leaves short and concave. Colour a dark reddish brown. The leaves are closely imbrieated, the lobes uniting by one-fourth of their length. The smaller lobe is folded at the margin, and receives in the sinus thus formed a similar fold of the leaf above; its inner rounded margin is sharply inciso-dentate. Specifically this is remarkably distinct from any of its congeners, and like the former, is rather an abnormal form.

PLATE. CLVI. Fig. III:---1, plant of the natural size: 2 and 3, back and front views of leaf and stipule; 4, leaf with the stipule removed, showing the form of the smaller lobe; 5, stipule; ---magnified.

7. JUNGERMANNIA laminigera, Hook. fil. et Wils.; caule cæspitoso subcrecto ramoso planiusculo, foliis imbricatis patentibus eroso-ciliatis subter lamellatis marginibus lamellisque undulatis ciliato-dentatis, lobo ventrali lanccolato basi bilobo, dorsali semi-cordato, stipulis majoribus late rotundato-quadratis 4–5fidis ciliatis, calyce terminali oblongo compresso spinuloso ore laciniato ciliato obscure bilobo. Nobis *in* Lond. Journ. Bot. v. 3. p. 456. (TAB. CLVI. Fig. IV.)

HAB. Hermite Island, Cape Horn; on the ground in the woods, abundant.

Cæspites laxi, superne pallide flavo-virescentes, inferne sordide brunnei. Caules 1–3 unc. longi ; ramis erectis, subfastigiatis. Folia patentia, dorso carinis lamellisve plurimis eristatis infra apicem evanidis ornata, versus apices setosa, marginibus ereberrime spinuloso-dentatis. Stipulæ majuseulæ, dorso basi obscure lamellatæ, segmentis linearibus subobtusis, marginibus recurvis ciliato-dentatis. Calyx oblongus, $\frac{1}{2}$ -exsertus, pallidus, extus spinulosus, vix lamellatus. Seta uncialis. Capsula eylindracea.

This a good deal resembles the *J. lamellata* (v. supra), but is more robust though smaller, the lobes of the leaves are more united throughout their whole length and the stipules are 4–5-fid. It varies much in size, some of our specimens being hardly an inch long.

PLATE CLVI. Fig. IV.—1, plant of the natural size; 2, upper, and 3, under surface of the leaf; 4, stipule; 5, calyx :—magnified.

(3. PLAGIOCHILA, Nees et Mont.)

S. JUNGERMANNIA ansata, Hook. fil. et Tayl.; eaule gracili laxe eæspitoso elongato subramoso, foliis laxe imbricatis plauis erectis appressis secundis oblique rotundatis basi decurrentibus integerrimis fuscis. Nobis *in Lond. Journ. Bot.* v. 3. p. 457. (TAB. CLVI. Fig. VI.)

HAB. Falkland Islands; amongst moss on the hills, abuudant.

Cæspites laxi, inter muscos implexi, pallide brunnei. *Caules* flexuosi, graeiles, tenues, vix ramosi, *Folia* submembranacea, integerrima, rotundata, erecta, cauli appressa, hine homomalla, margine anteriore obscure recurvo, posteriore decurrente.

In habit resembling the *J. colorata*, Hook. (v. infra), but the stems are more clongated, and the leaves quite entire and free, not united into opposite pairs at their bases. The same characters distinguish it from the *Pl. Braunii* of Java, which further has heteromallous leaves. The trivial name alludes to the produced lower margin of the rounded leaf forming a handle.

PLATE CLVI. Fig. V1.-1, plant of the natural size; 2 and 3, front and back view of a leaf :--magnified.

9. JUNGERMANNIA unciformis, Hook. fil. ct Wils.; caulibus cæspitosis subsimplicibus erectis apice

5 E

plerumque curvatis, foliis crassiusculis laxe imbricatis erectis secundis appressis oblique ovato-rotundatis, margine inferiore gibboso obscure sinuato-dentato superiore incurvo basi celluloso. Nobis *in Lond. Journ. Bot.* v. 3. p. 457. (TAB. CLVI. Fig. V.)

HAB. Hermite Island, Cape Horn; on the trunks of trees near the ground.

Cæspites 2-4 une. lati, rufo-brunnei. Caulis $\frac{1}{2}$ une. longus, apice strictus v. sæpius curvatus, nune hamatus. Folia e basi caulis gradatim majora, imbricata, homomalla, compressa, margine superiore incurvo, plica tumida elongata, inferiore tumido obscure et obtuse sinuato; cellulæ minimæ, densæ, nisi ad basin foliorum ubi majores pallidioresque evadunt, maculam latam efficientes.

A species allied to the *J. biserialis*, L. and L., of Tasmania, but less than half the size, with more minute denticulations to the leaves, which are widely ovate, not round or decurrent at the anterior margin, nor bispinous at the apex.

PLATE CLVI. Fig. V.—1, plant of the natural size; 2, front, and 3, back view of leaf and portion of stem; 4 and 5, similar views of leaves removed from the stem;—magnified.

10. JUNGERMANNIA Magellanica, Lindb.; Sp. Hep. p. 164. Gottsche. Lindb. et Nees, Syn. Hep. p. 53. Mont. in Foy. au Pole Sud, Bot. Crypt. p. 271.

HAB. Strait of Magalhaens, St. Nicholas Bay; M.M. D'Urville et Jacquinot.

We owe our acquaintance with this plant to the liberality of our learned friend M. Montagne, who most generously has communicated to us his own examples of such Antarctic species as we desired for comparison or examination, unhesitatingly confiding his unique specimens to the care of the post-office, that we might profit to the fullest by his labours, and avoid unnecessary errors. It differs from our *J. unciformis* in the larger and narrower leaves, which are acute and servato-dentate along the inner margin.

11. JUNGERMANNIA *duricaulis*, Hook. fil. et Tayl.; caulibus cæspitosis duris robustis flexuosis erectis ramosis, foliis amplis subimbricatis patentibus oblique ovato-cordatis basi decurrentibus argute denticulatis, basi postica porrecta verticali, margine inferiore lente recurvo. Nobis *in Lond. Journ. Bot.* v. 3. p. 458. (TAB. CLVI. Fig. IX.)

HAB. Hermite Island, Cape Horn; abundant, in the woods.

Species insignis. Cæspites laxi, majusculi; externe pallide sed luride olivacei. Caules 4 unc. longi, irregulariter ramosi, ramis compressis. Folia $\frac{1}{8}$ unc. longa, arete laxiusve imbricata, marginibus dorsalibus parium suboppositorum rotundatis postice porrectis appressis carinamque cauli quasi efficientibus; margine superiore paulo incurvo, marginibus omnibus minute sed creberrime et regulariter denticulatis. Perigonia in spicam brevem terminalem disposita.

Most nearly related to the *Pl. flaccida*, Lindb., of St. Vincent, which has a very similarly hard and woody stem but the present may be known by the greater breadth of its branches and foliage, by its more compound ramification and the minute denticulation of its leaves.

PLATE CLVI. Fig. IX .--- 1, plant of the natural size ; 2, stem and opposite pair of leaves ; 3, leaf :-- magnified.

12. JUNGERMANNIA asplenioides, Linn.; Sp. Pl. p. 1597. Mont. Foy. au Pole Sud, Bot. Crypt. p. 268.

HAB. Strait of Magalhaens; D'Urville.

We have seen no Fuegian specimens of this species. Those M. Montagne has examined, are in a very unsatisfactory state.
13. JUNGERMANNIA sphalera, Hook. fil. et Tayl.; caule laxe cœspitoso erecto basi ramoso apice incurvo, foliis vix imbricatis subhorizontaliter patentibus secundis siccitate suberectis late oblique ovato-rotundatis acutis, apice inæqualiter bifido v. bidentato, margine superiore incurvo integerrimo, inferiore planiusculo dentato laxe celluloso. Nobis *in Lond. Journ. Bot.* vol 3. p. 458. (TAB. CLVI. Fig. VIII.)

HAB. Hermite Island, Cape Horn; growing amongst mosses in the woods.

Cæspites laxi, pallide olivacei. Caules 1-2 unc. longi, raro in ramos 2-3 erectos divisi. Rami compressi, recti v. curvati, apicibus rotundatis. Folia remotiuscula, madore patentia, sed secunda, apicibus subrecurvis, basi contracta, margine inferiore grosse irregulariter serrato.

Nearly allied to *J. uncialis*, but taller, with the leaves more remote, less imbricated and secund, more loosely cellular and not so strongly dentate. When moistened the differences are more apparent, the leaves in particular of *J. sphalera* being distinctly narrowed at the base,

PLATE CLVI. Fig. VIII.--1, plant of the natural size; 2 and 3, front view of leaf, and portion of stem; 4, the same detached from the stem:-magnified.

14. JUNGERMANNIA *uncialis*, Hook. fil. et Tayl.; caule breviusculo cæspitoso suberecto v. prostrato et ascendente ramoso, foliis imbricatis erecto-patentibus concavis late ovatis acutis argute irregulariter ciliatodentatis sublaxe cellulosis, margine inferiore subrecurvo, calyce majusculo terminali compresso late obovato, ore oblique subrotundato dentato-ciliato. Nobis *in Lond. Journ. Bot.* vol. 3. p. 459. (TAB. CLVI. Fig. VII).

HAB. Hermite Island, Cape Horn; on damp rocks and the trunks of trees.

Cæspites late extensi, pallide flavo-virescentes. Caules vix 1 unc. longi, subprostrati, rarius erecti, vage ramosi, ramis fructiferis subfastigiatis. Folia vix decurrentia, perichætialia calyce $\frac{1}{2}$ breviora. Calyx obovato-cuneatus, compressus, ore obtuse rotundato, oblique fisso, serrulato. Capsula oblongo-sphærica, vix exserta. Perigonia in spicas breves secus ramos disposita.

In habit the present approaches the African *P. sarmentosa*, Lindb., but in character it is more nearly allied to our Tasmanian *J. aculeata*. The former, whose fructification is unknown, has larger and more rounded leaves. The *J. aculeata* is a much larger plant; its leaves have a narrower base, and their superior margin is recurved with a broader fold, the denticulation is coarser, and calyx shorter, being scarcely exserted beyond the perichaetial leaves; above all, the cellulation of the *J. ancialis* is much coarser though belonging to a smaller plant.

PLATE CLVI. Fig. VII.—1, plant of the natural sizc; 2, stem, perigonium and leaves; 3, front, and 4, back view of portion of stem and leaf; 5, calyx and capsule :—magnified.

15. JUNGERMANNIA Jacquinotii, Mont., in Voy. au Pole Sud, Bot. Crypt. p. 273.

HAB. Strait of Magalhaeus; D'Urville.

A very different plant from any collected by the Antarctic Expedition.

16. JUNGERMANNIA minutulu, Hook. fil. et Tayl.; cæspitosa, caule brevissimo erecto parce ramoso, foliis imbricatis erectis appressis obovato-rotundatis convexiusculis, margine anteriore subdecurrente posteriore recurvo, supremis majoribus denticulatis. Nobis *in Lond. Journ. Bot.* vol. 3. p. 459. (TAB. CLVII. Fig. I).

HAB. Kerguelen's Land; on the ground and on moist rocks.

Cæspites late extensi, atro-virides. *Caules* crecti, crassiusculi. *Rami* primarii vix $\frac{1}{8}$ unc. longi. *Folia* inferiora minuta, subintegerrima, caule vix latiora, gradatim majora, superiora arctius imbricata, in capitulum compressum dilatatum congesta, superiora crenato-denticulata, omnia crassa, obscure cellulosa; cellulis parvis, opacis, margina-libus conspicuis.

Each branch is short and bears but few pairs of leaves :---these are narrow at the base, gradually widening upwards to the top of the branches, where they are collected into a flattened head three or four times wider than the inferior part of the shoot. This resembles in general appearance the *P. pusilla*, Mont. (of Tasmania), but is more minute, has not curved stems, there are fewer leaves on the shoots, and the anterior margin of the leaf is decurrent.

PLATE CLVII. Fig. 1.—1, plant of the natural size; 2 and 3, front and back views of leaf and portion of stem: magnified.

17. JUNGERMANNIA heterodonta, Hook. fil. et Tayl.; cæspitosa, caule erecto v. prostato ramoso, ramis ascendentibus subfastigiatis, foliis crecto-patentibus late ovatis obovatisve grosse inæqualiter eroso-dentatis margine superiore deeurrente, inferiore apice obscure bifido, calyce terminali foliis periehætialibus breviore angustioreque obovato-rotundato, ore contracto æquali truncato ciliato-dentato. Nobis in Lond. Journ. Bot. vol. 3. p. 460. (TAB. CLVII. Fig. II).

HAB. Kerguelen's Land; on moist rocks near the sea.

Cæspites late extensi, rupibus appressi, læte olivaeco-virides. *Caules* sub 2 une. longi, irregulariter vage ramosi. *Folia* subarete imbrieata, oblique rotundata, apiee latiuscula, dentibus marginalibus, nunc manifeste nunc obseure bifida. *Calyx* perichætio brevior, obseure bilabiatus ; labiis rotundatis, crenatis et minute eiliatis.

Allied to the *P. sciophila* of Nepaul, which has emarginato-dentate leaves, but from which the present may be distinguished by its smaller size, erect growth, and closely imbricated foliage, which is more toothed. The perigonia, with which the Antaretie species is supplied, are in the form of a narrow spike, whose leaves are minute, erect, imbricated, and bidentate with somewhat squarrose apiecs.

PLATE CLVII. Fig. II.—1, plant of the natural size; 2, leaf and portion of stem; fig. 3, ditto removed from stem; 4, ealyx:—magnified.

18. JUNGERMANNIA Chonotica, Tayl.; cæspitosa, surculis erectis subramosis complanatis basi nudiusculis, foliis imbricatis erecto-patentibus oblongo-rotundatis basi angustatis convexis, marginibus recurvis spinosodentatis. Tayl. in Lond. Journ. Bot. vol. 5. p. 260.

HAB. Chonos Archipelago; C. Darwin, Esq.

Cæspites pallide fulvi. *Caules* 2-3 une. longi, dendroidei, v. nudi basique simplieiuseuli. *Perigonia* parva, brevia, in spieam linearem arete imbricatam disposita, foliolis parvis tumidis dentieulatis.

Resembles the *P. fasciculata*, Lindb., of New Holland and Lord Auckland's group; the shoots however are much narrower, leaves shorter, more distinctly and minutely toothed, the branches fascicled and the cells of the leaves much more minute.

19. JUNGERMANNIA distinctifolia, Tayl. l. e.; Lindb. Sp. Hep. p. 17. t. 3. Gottsche, Lindb. et Nees, Syn. Hep. p. 30.

HAB. Staten Island; Menzies in Herb. Hook.

We have not seen specimens of this from the Antarctic Expedition. It is also a native of Jamaica and the Brazils.

(4. JUNGERMANNIA, L. et auct. recent.)

20. JUNGERMANNIA colorata, Lehm. in Linn. vol. 4. p. 366. Gottsche, Lindb. et Nees, Syn. Hep. p. 86. Fl. Antarct. Pt. 1. p. 149.

HAB. Hermite Island, Cape Horn; from the sea to the mountain-tops. Falkland Islands; on the hills. Kerguelen's Land; particularly abundant on the ground, on the hills.

Also found in New Holland, Tasmania, and New Zealand, Lord Auckland's group, the Cape of Good Hope, and Juan Fernandez. In the Falkland Islands it forms large black patches on the alpine rocks, resembling an Andreæa.

21. JUNGERMANNIA byssacea, Roth; Cat. Bot. vol. 2. p. 158. Engl. Bot. t. 2463.

HAB. Falkland Islands; amongst mosses on the hills.

22. JUNGERMANNIA bicuspidata, Linn.; Hook. Brit. Jung. t. 11. Engl. Bot. t. 2239.

HAB. Falkland Islands; on moist rocks.

One of the, comparatively speaking, few *Hepaticæ*, which, according to the modern limitation of species, is acknowledged to be a cosmopolite.

23. JUNGERMANNIA *rigens*, Hook. fil et Tayl.; minima, laxe cellulosa, cæspitosa, caule prostrato subpinnatim ramoso, ramis erectis, foliis laxe imbricatis suberectis concavis late oblongis bifidis, segmentis incurvis late subulatis integerrimis, stipulis ovato-rotundatis concavis bifidis segmentis late subulatis integerrimis. Nobis *in Lond. Journ. Bot.*, vol. 3, p. 461. (TAB. CLVII. Fig. III).

HAB. Falkland Islands; on moist maritime rocks.

Cæspites parvi, pallide olivaceo-flavescentes. *Caules* 2–3 lin. longi, carnosiusculi, simplices v. ramosi. *Folia* sursum gradatim minora, imbricata, tumida, hine caulis submoniliformis. *Stipulæ* pro planta amplæ, foliis consimiles, ad medium v. supra medium bifidæ.

Allied to the British J. Francisci, Hook., but more minute, the leaves more concave, and the stipules of a very different form.

PLATE CLVII. Fig. III.---1, plant of the natural size; 2, portion of stem, leaf, and stipule; 3, stipule, removed :-- magnified.

24. JUNGERMANNIA *tubulata*, Hook. fil. et Tayl.; parvula, caule laxe cæspitoso procumbente ramoso, foliis laxis suberectis oblongis bifidis segmentis acutis acuminatisve, calyce terminali anguste lineari-elongata tubulata, basi oblongo, ore plicato minutissime denticulato, foliis perichætialibus segmentis lanceolatis integerrimis. Nobis *in Lond. Journ. Bot.* vol. 3. p. 463. (TAB. CLVII. Fig. VI).

HAB. Falkland Islands; on moist rocks near the sea.

Caules graciles, laxe cæspitosi, simpliciusculi v. ramosi, vix $\frac{1}{2}$ unc. longi, sæpissime ramulis flagelliformibus nudis aucti, (ut in *J. bicuspidata*, L.), pallide virescentes. *Folia* crecta, pallida, pellucida, basi concava, in segmentis duobus lanceolatis apice subulatis divisa, sinu angusto acuto. *Calyces* conspicui, albidi, clongati, superne subinflati et plicati, ore minutissime denticulato. *Folia* perichætialia erecta; segmentis angustis, integerrimis. *Capsula* oblonga. *Sporæ* numerosissimæ, luteo-brunneæ, subangulatæ. *Elateres* e helice duplici constantes.

So very near the European *J. bicuspidata*, as to be hardly distinguishable from it specifically: the capsules are however shorter, the perichætial leaves entire, the calyx longer, and the areolæ of the foliage smaller. The calyces are always terminal, whereas in *J. bicuspidata* they are more frequently lateral.

5 f

25. JUNGERMANNIA vasculosa, Hook. fil. et Tayl.; flaccida, tenerrima, cæspitosa, caule procumbente subramoso, foliis imbricatis secundis erectis rotundato-quadratis, basi lato decurrente, margine integerrimo undulato, stipulis majoribus ovatis concavis bifidis, segmentis lanceolatis integerrimis v. basi utrinque unidentatis. Nobis in Lond. Journ. Bot. vol. 3. p. 461. (TAB. CLVII. Fig. IV).

HAB. Falkland Islands; on wet rocks near the sea, abundant.

Cæspites late extensi, 2-4 unc. lati, atro-virides. *Caulis* 2 unc. longus, parce ramosus. *Folia* laxiuscule imbricata, integerrima; marginibus subinflexis undulatis, superiore subgibboso, inferiore longe decurrente; substantia tenerrima, flaccida, cellulis majusculis. *Stipulæ* conspicuæ, caulem amplectentes; segmentis late lanceolatis, erectis, sinn obtusiusculo.

This has a good deal of resemblance to the *J. cordifolia*, Hook., of Britain, but the presence of stipules will at once distinguish the Antarctic plant. In many respects it has an equal claim to be considered a *Lophocolea* as a *Jungermannia*; on the whole, however, we incline to retain it in the latter genus.

PLATE CLVII. Fig. IV.—1, plant of the natural size; 2, portion of branch, with leaf and stipule; 3, stipule: both magnified.

26. JUNGERMANNIA erinacea, Hook. fil. et Tayl.; tenerrima, cæspitosa, caule suberecto ramoso, ramis erecto-patentibus, foliis imbricatis patentibus flaceidis ciliato-dentatis apice obtusis plus minusve profunde emarginatis, lobo inferiore ovato-rotundato superiore libero ovato adpresso stipulis majoribus late rotundatis irregulariter ciliato-dentatis integris bifidisve. Nobis *in Loud. Journ. Bot.* vol. 3. p. 462. (TAB. CLXI. Fig. IV).

HAB. Falkland Islands; on moist rocks near the sea.

Pallide olivaceo-flava. Caspites laxi, extensi. Caulis $1\frac{1}{2}-2$ unc. longus, flaccidus, ramosus, ramis subfastigiatis. Folia laxe imbricata, tenerrima et flaccida sed areolis minutis, apice plerumque bifida, sinu lato rotundato, ciliis marginalibus basi latiusculis sæpe repente in apicem articulatam desinentibus. Stipulæ rotundatæ, ambitu ciliatæ, lobulo minore folii majores, bifidæ.

A very beautiful species, allied to our *J. diplophylla* (Pt. 1. p. 152. t. 64. f. iv); where fructification is unknown, they together appear to form as natural a genus as any which has been proposed out of *Jungermannia*, and differ from *Scapania* in the presence of stipules. The present is readily distinguishable from *J. diplophylla* by the separation of the two lobes of the leaf, by their emarginate tips, by the larger and closer ciliation of their margins and by the less deeply but more frequently divided and broader stipules.

27. JUNGERMANNIA *humectata*, Hook. fil. et Tayl.; laxe cæspitosa, flaccida, caule erecto parce ramoso, foliis remotis tenuibus erecto-patentibus undulatis basi amplexicaulibus ovato-rotundatis emarginatis bifidisve segmentis obtusiusculis subdivaricatis iutegerrimis v. utrinque dentatis, stipulis foliis consimilibus sed minoribus. Nobis *in Lond. Journ. Bot.* vol. 3. p. 462. (TAB. CLVII. Fig. V).

HAB. Falkland Islands; on wet sand by the sides of mountain-streams.

Cæspites laxi, extensi, inferne atro-brunnei; ramis paucis, erectis, pallide fusco-olivaceis. *Folia* alterna, remotiuscula, basi caulem totam fere amplectentia, late obovato-oblonga v. rotundata, bifida, sinu acuto v. obtuso.

Possibly from its rather anomalous locality, an altered state of some other species, though we cannot say of what. In the wet place of growth, erect habit and general outline of the leaf, it resembles the British J. Lyoni,

Tayl. (J. socia. var., Gottsche, Lind. et Nees), differing in the paler green colour of the young shoots, in the more delicate foliage, smaller areolæ, deeper emargination and clasping leaves.

PLATE CLVII. Fig. V.-1, plants of the natural size; 2, stipule; 3, leaf :- both magnified.

28. JUNGERMANNIA austrigena, Hook. fil. et Tayl.; laxe cæspitosa, caule elongato ascendente subramoso, surculis incurvis, foliis imbricatis subsecundis erecto-patentibus rotundatis convexis integerrimis marginibus recurvis perichætialibus rotundatis, stipulis majoribus rotundatis, marginibus reflexis integerrimis bidentatisve, calyce terminali oblongo compresso ore subintegro trigono. J. austrigena et J. cavispina. Nobis in Lond. Journ. Bot. vol. 3. p. 463 et 466. (TAB. CLVII. Fig. VII. and TAB. CLVIII. Fig. V).

HAB. Hermite Island, Cape Horn; moist banks in woods (fruit). Falkland Islands, along with J. humeetata.

Cæspites laxi, lati, pallide flavidi, virides v. atro-brunnei. Caules 2-3 unc. longi, vage ramosi, flexuosi, crassiusculi. Folia arete imbricata, subopposita, autice decurrentia, siccitate plerumque crispata, recurva; madore suberecta, appressa, marginibus plus minusve recurvis. Stipulæ rotundatæ, basi utrinque decurrentes, integerrimæ v. apice bidentatæ, marginibus valde deflexis porrectis, intra margines posticos foliorum verticaliter compressæ. Calyæ majusculus, oblougus, latiusculus, trigonus.

A very curious species, and unlike any with which we are acquainted. The habit of the Falkland Island specimens when dried, is, owing to their having grown in water, so peculiar, that we regarded them at first as a different species, which we described as *J. cavispina*, from the reflexed margins of the closely imbricating stipules, giving a grooved appearance to the back of the stem. The calyx is that of a *Lophocolea*, from which group the presence of stipules and the entire leaves remove it.

PLATE CLVII. Fig. VII. Falkland Island state, (sub nom. J. cavispinæ).—1, plant of the natural size; 2, front, and 3, back view of stem and leaf; 4, stem and stipule; 5 and 6, stipules :—magnified. PLATE CLVIII. Fig. V. (Hermite Island state).—1, plant of the natural size; 2, lateral, 3, back, and 4, front view of stem, leaf, and stipule; 5, stem and stipule; 6 and 7, stipules :—magnified.

29. JUNGERMANNIA palustris, Hook. fil. et Tayl.; caule elongato disperso flaccido ramoso, foliis laxe imbricatis erecto-patentibus tenuissime membranaceis rotundatis valde concavis marginibus incurvis medio longitudinaliter undulatis integerrimis, stipulis majoribus ovalibus cymbiformibus integerrimis. Nobis in Lond. Journ. Bot. vol. 3. p. 464. (TAB. CLVII. Fig. VIII).

IIAB. Hermite Island, Cape Horn; on the borders of an alpine lake, growing in the water.

Caules inter *Muscos* aliasque *Hepaticas* demersas ascendentes, sparsi, 3 unc. longi ; ramis erectis, flaccidissimis. *Folia* tumida, varie incurva, medio plerunque plica longitudinali notata. *Stipulæ* valde concavæ, subimbricatæ.

Allied to the British J. Doniana, and to the J. involutifolia, Mont. (v. infra), but very distinct from both. The leaves resemble those of Hypnum cochlearifolium, Schwaegr.

PLATE CLVII. Fig. VIII.-1, plant of the natural size; 2, front view of stem, leaves, and stipules; 3, back view of ditto:-magnified.

30. JUNGERMANNIA involutifolia, Mont. in Gottsche, Nees et Lindb. Syn. Hep. p. 81. Voy. an Pole Sud, Bot. Crypt. p. 260.

HAB. Strait of Magalhaens; on tufts of Hypnum fluitans: M. Hombron.

The nearest ally to this plant is J. notophylla, nobis.

[Fuegia, the

31. JUNGERMANNIA *fulvella*, Hook. fil. et Tayl.; parvula, caule implexo procumbente ramoso, ramis cylindraceis, foliis arcte imbricatis patentibus concavis marginibus incurvis remote dentatis carnosiusculis pellucidis laxe cellulosis caulinis oblique rotundatis, rameis minoribus arctius imbricatis rotundatis, stipulis erectis concavis late rotundatis integerrimis v. irregulariter dentatis, calyce laterali obovato cylindraceo foliis perichætialibus integerrimis duplo longiore. Nobis *in Lond. Journ. Bot.* vol. 3. p. 464. (TAB. CLVIII. Fig. I.)

HAB. Hermite Island, Cape Horn; in the woods on dead timber, trunks and twigs of trees; abundant. South part of Tierra del Fuego; C. Darwin, Esq.

Cæspites 3 unc. lati, pallide fulvi. Caulis 1 unc. longus, pluries vage ramosus, ramis tenuibus. Folia caulina rameis laxius imbricata, basi latiora planiora, dentibus valde irregularibus. Stipulæ rameæ caulinis breviores, sed latiores, rariusque dentatæ. Folia perichætialia parva, oblonga, concava, calyce ter breviora. Calyx lineari-oblongus, trigonus, subtumidus, ore angustato. Seta $\frac{1}{3}$ unc. longa. Capsula late oblonga.

With much the appearance of a *Herpetium*, but having no flagelliform shoots, and the leaves are irregularly dentate. It is a most distinct species.

PLATE CLVIII. Fig. I.—1, plant of the natural size; 2, part of stem, leaf, and stipule; 3, leaf; 4, upper leaf; 5, stipule; 6, portion of branch with leaves, calyx, seta, and capsule; 7, calyx and perichaetium; 8, corolla:—*magnified*.

32. JUNGERMANNIA obvoluta, Hook. fil. et Tayl.; cæspitosa, caule ascendente vage ramoso, ramis cylindraccis elongatis flaccidis suberectis, foliis imbricatis patentibus membranaceis laxe cellulosis late quadratis margine incurvo undulatis bifidis sinu angusto hic illic grosse dentatis subdecurrentibus, stipulis majoribus rotundato-ovatis concavis emarginatis utrinque uni-dentatis. Nobis *in Lond. Journ. Bot.* vol. iv. p. 80. (TAB. CLXI. Fig. I.).

HAB. Hermite Island, Cape Horn, and the Falkland Islands; on the trunks of trees, and on the ground.

Cæspites pallide olivacei, straminei. Caules $\frac{1}{2}-\frac{3}{4}$ unc. longi. Folia secus partem caulis superiorem involuta. Stipulæ cmarginatæ, sinu latiusculo.

Allied to J. oligophylla, nob., but quite different specifically from that, and from any other species with which we are acquainted.

PLATE CLXI. Fig. I.-1, plant of the natural size; 2, portion of stem, leaves, and stipules; 3, leaf; 4, stipule: *magnified*.

33. JUNGERMANNIA madida, Hook. fil. et Tayl.; cæspitosa, caule elongato planiusculo erecto ramoso, ramis erectis fastigiatis, foliis laxe imbricatis patentibus semiamplexicaulibus concavis ovato-quadratis bifidis, segmentis acutis incurvis integerrimis v. apices versus. 2–3-dentatis, stipulis majoribus foliis paulo brevioribus concavis late ovatis bifidis subintegerrimis, calyce terminali cylindraceo recto apice obscure dentato, capsuke valvis lineari-elongatis. Nobis *in Lond. Journ. Bot.* vol. 3. p. 465. (TAB. CLVIII. Fig. II).

HAB. Hermite Island, Cape Horn; on moist banks, and in bogs on the mountains, forming dense tufts.

Cæspites densi, 2-3 unc. lati, locis humidioribus obscure virescentes, siccis rufescentes. *Caules* 2 unc. longi. *Folia* laxe imbricata, ad $\frac{1}{3}$ longitudinis bifida, minute cellulosa, integerrima v. apices versus 1-2-dentata. *Stipulæ* foliis subæquales, late oblongæ, bifidæ, segmentis obscure dentatis. *Calyx* elongatus, superne attenuatus, apice truncatus. *Capsula* elongata, cylindracea.

Very closely allied to the *J. serrulata*, Sw. (*Musc. Exot.* t. SS), of the West Indies, but the leaves are not so densely imbricated, are scarcely serrulate, their areolæ are more minute, and the stipules are different. When growing in moist places the plant is greener and larger, and the leaves more generally serrulate than when found in drier situations.

PLATE CLVIII. Fig. II.—1, plant of the natural size; 2, portion of stem, leaf, and stipule; 3, leaf; 4, stipule; 5, perichætial leaf; 6, calyx, seta, and capsule; 7, corolla; 8, capsule:—magnified.

34. JUNGERMANNIA æquata, Hook. fil. et Tayl.; caule brevi implexo procumbente ascendente ramoso flexuoso, ramis compressis curvatis, foliis imbricatis secundis appressis suboppositis rotundatis margine incurvis integerrimis crassis opacis ima basi inter se et cum stipula parva ovata bifida v. integra connatis. Nobis in Lond. Journ. Bot. vol. iii. p. 465. (TAB. CLVIII. Fig. III.)

HAB. Hermite Island, Cape Horn; on the trunks of trees in the woods.

Caspites parvi, inter Muscos aliasque Hepaticas nidulantes, rufo-brunnei. Caules unciales, vage sed parce ramosi, basi nudi, sursum curvati. Folia arcte imbricata, oblongo-rotundata, madore e marginibus incurvis tumida, opposita, basi antice connata, postice cum stipula adnata. Stipula ovata, bifida v. varic secta, segmentis subulatis.

The form of the leaves, their opposite arrangement and connexion in front, are similar to J. Brankiana, Necs, but that species is destitute of stipules.

PLATE CLVIII. Fig. III.—1, plant of the natural size; 2, portion of branch; 3, ditto with front view of leaf and stipule, 4 and 5, stipules:—magnified.

35. JUNGERMANNIA otophylla, Hook. fil. et Tayl.; caule debili flavido elongato subramoso, foliis oppositis secundis erecto-patentibus imbricatis flavidis et membranaceis late reniformi-rotundatis basi latissime cauli adnatis integerrimis, margine superiore basi tumido recurvo, stipulis majoribus concavis late rotundatis emarginatis integerrimis obscure sinuatisve. Nobis *in Lond. Journ. Bot.* vol. iii. p. 466. (TAB. CLVIII. Fig. IV.)

HAB. Hermite Island, Cape Horn ; in alpine bogs.

Cæspiles laxi, luride olivacei v. albescentes. Caulis gracilis, 3 unc. longus, parce ramosus; ramis erectis. Folia tenuissime membranacea, latissime oblonga v. rotundata, basi ad marginem anteriorem quasi auriculata. Stipulæ amplæ, subimbricatæ, marginibus incurvis, apicibus emarginatis, sinu lato, nunc apice sinuato.

In habit and general appearance this approaches our J. palustris, which inhabits similar localities, but they are in many respects widely different plants.

PLATE CLVIII. Fig. IV.—1, plant of the natural size; 2, front, and 3, back view of portion of stem and leaf; 4, stipule:—magnified.

36. JUNGERMANNIA densifolia, Hook., Musc. Exot. t. 36. Scapania? densifolia, Gottsche, Lindb. et Nees, Syn. Hep. p. 72.

HAB. Hermite Island, Cape Horn; in wet bogs, &c.

A very abundant species in Hermite Island.

37. JUNGERMANNIA chloroleuca, Hook. fil. et Tayl.; caule erecto cæspitoso parce ramoso flavido, foliis subapproximatis imbricatis patentibus ovatis v. ovato-oblongis inferne tumidis semi-amplexicaulibus bipartitis ciliato-dentatis, segmentis linearibus ligulatisve summo apice bifidis. Nobis *in Lond. Journ. Bot.* vol. iii. p. 467. (TAB. CLXI. Fig. V.)

HAB. Hermite Island, Cape Horn; on moist banks near the sea.

Cæspites subereeti, flavo-vireseentes. *Folia* disticha, vix imbrieata, segmentis plerumque recurvis, ramis hinc (siceitate præcipne) squarrosis.

So nearly allied to the previous species that a particular description is hardly required; it differs conspicuously in the colour. These species were never seen passing into one another, and both are remarkably constant to their characters. They belong, with the *J. vertebralis*, Gottsche (Pt. 1. p. 153), of Lord Auckland's group and Tasmania, also a very closely allied plant, to a distinct section of the genus. The *J. chloroleuca* differs from *J. vertebralis* in colour, size, and different texture, also in the form of the leaves and ciliation; and from *J. densifolia*, besides the colour, mentioned above, in the denser tissue of the remote leaves, which are much broader at the base, and whose areolæ are blended together, in their long ciliation and bifid apiees.

PLATE CLXI. Fig. V.-1, plant of the natural size; 2, portion of stem and leaf :- magnified.

38. JUNGERMANNIA clandestina, Mont., in Voy. au Pole Sud, Bot. Crypt. p. 264. t. 16. f. 4. Gottsche, Lindb. et Nees, Syn. Hep. p. 73.

HAB. Strait of Magalhaens; Port Famine and Port Gallant, M. Hombron.

39. JUNGERMANNIA schismoides, Mont., vid. Pt. 1. p. 150. (TAB. CLXI. Fig. IX.)

HAB. Hermite Island, Cape Horn; creeping through tufts of mosses in the woods.

The leaves of these specimens are slightly serrulate along the margins, in which respect alone the plant differs from that found in Lord Auekland's group.

PLATE CLX1. Fig. IX.-1, plant of the natural size; 2, 3, and 4, leaves :- magnified.

40. JUNGERMANNIA crebrifolia, Hook. fil. et Tayl.; caule cæspitoso erecto ramoso, ramis suberectis, foliis carnosulis arcte imbricatis erecto-patentibus secundis coneavis late ovato-rotundatis bilobis, lobis ovatis subaeutis integerrimis inferiore minore basi dentato v. integerrino, calyce minimo laterali obovato plicato, ore scarioso laeiniato, laeiniis lanceolatis. Nobis *in Journ. Lond. Bot.* vol. iii. p. 467. (TAB.CLVII. Fig.IX.)

HAB. Hermite Island, Cape Horn.

Dense eæspitosa, rufo-bruunea. *Caules* fere 2 une. longi, siceitate fragiles, irregulariter repetitim ramosi, rarius superne paulo incrassati. *Folia* arcte imbricata, valde coneava, marginibus apieibusque madore ercetis, lobo superiore majore, inferiore basi supra eaulem producto, integerrimo v. uni-dentato. *Calyces* minuti, valde inconspieui, ore albido scarioso.

Closely allied to the *J. cryptodon*, Wils. MS., of the Andes of Colombia, which has a similarly toothed lower lobe of the leaf, equally produced at the base aeross the stem. The present is a larger plant, with more imbricated and erect leaves, their lower lobe smaller, and the produced portion larger in proportion.

PLATE CLVII. Fig. IX .---- 1, plant of the natural size ; 2 and 3, leaves :--- magnified .

41. JUNGERMANNIA *humilis*, Hook. fil. et Tayl.; parvula, caule implexo procumbente radicante ramoso, foliis subimbricatis erecto-patentibus secundis rotundatis concavis integerrimis crassiusculis, stipulis minutis ovatis integris v. bifidis segmentis unidentatis v. irregulariter sectis. Nobis *in Lond. Journ. Bot.* vol. iii. p. 468. (TAB. CLVIII. Fig. VI.)

HAB. Kerguelen's Land; on tufts of Azorella Selago.

Cæspites lati, pallide flavo-olivacei. Caulis vix uncialis, irregulariter ramosus. Folia laxe imbricata, basi late eaule adnata sed non decurrentia, patentia, homomalla. Stipulæ eauli æquilatæ, varie sectæ, emarginatæ, bifidæ v. irregulariter sinuato-dentatæ.

Allied to *J. turgescens*, nobis (Pt. 1. p. 150, t. lxiv. f. 2.), of Lord Auekland's group; but the present may be readily distinguished by its smaller size, more olive colour, its toothed stipules, more patent and differently shaped leaves, whose attachment is also different, and which are not decurrent; and by their larger areolæ.

PLATE CLVIII. Fig. VI.—plant of the natural size. 2, stem, back view of leaf and stipude; 3, front view of leaf; 4, stipule:—magnified.

42. JUNGERMANNIA minuta, Crantz, vid. Fl. Antaret. Pt. 1. p. 152.

HAB. Kerguelen's Land; on tufts of mosses, &c., on the hills.

Also found in Lord Auekland's group, but hitherto not elsewhere in the Southern Hemisphere; nor out of Europe in the Northern.

43. JUNGERMANNIA quadripartita, Hook., Muse. Exot. t. 117. Gottsche, Lindb. et Nees; Syn. Hep. p. 146.

HAB. Staten Land, Menzies (in Herb. Hook.); Hermite Island, Cape Horn; on moist banks, Mr. Davis.

(5. GYMNANTHE, Tayl.)

44. JUNGERMANNIA Urvilleana, Mont., vid. Fl. Antarct. Pt. 1. p. 153.

HAB. Hermite Island, Cape Horn; in the woods.

Also a native of Lord Auckland's group and Tasmania.

(6. LOPHOCOLEA, Nees.)

45. JUNGERMANNIA *textilis*, Hook. fil. et Tayl.; caule laxe implexo prostrato parce vage ramoso plano, foliis distichis horizontaliter patentibus complanatis approximatis late ovato-quadratis apice bifidis planis laxe cellulosis segmentis subulatis acutis integerrimis, stipulis ovatis bipartitis segmentis linearibus divaricatis. Nobis *in Lond. Journ. Bot.* vol. iii. p. 468. (TAB. CLVIII. Fig. IX.)

HAB. Hermite Island, Cape Horn; in woods. Falkland Islands; on wet rocks near the sea, very common.

Cæspites late extensi, pallide sed læte flavo-virides. *Caules* 2 unc. longi, laxe intertexti, terræ appressi. *Folia* disticha, omnia horizontaliter patentia, basi lata, eauli adnata sed non decurrentia, margine superiore subrotundata inferiore reeta; substantia tenera, laxe arcolata. *Stipulæ* parvæ, eauli æquilatæ, bipartitæ, segmentis subulatis acuminatis. *Perigonia* nune secus ramos obvia, plerumque apices versus; foliis arete appressis erectis, basi tumidis.

In some particulars resembling our J. *planiuscula* (Pt. 1. p. 156. t. lxv. f. 2), of Lord Auckland's group, which is a larger plant, with leaves rounded and otherwise of a very different form. The whole stratum is very flat and appressed, wide, of a fine shining green colour, and soft texture.

PLATE CLVIII. Fig. IX.-1, plant of the natural size; 2, stem and leaves; 3, stipule:-magnified.

46. JUNGERMANNIA *leptantha*, Hook. fil. et Tayl.; caule flaceido implexo procumbente ramoso, foliis distichis planis subimbricatis patentibus ovato-oblongis integerrimis margine superiore subrecurvo apice emarginato-bidentatis dentibus elongatis sinu rotundato, stipulis porrectis minutis bipartitis segmentis subulatis extus unidentatis 4-partitisve, calyce terminali lineari-oblongo trigono ore trifido segmentis emarginatobidentatis serratis rarius subintegris. Nobis *in Lond. Journ. Bot.* vol. iii. p. 471. Lophocolea coadunata Nees, fid. Montagne in Voy. au Pole Sud, Bot. Crypt. p. 256 (non Jung. coadunata, Swartz.) (TAB. CLIX. Fig. VI.)

HAB. Hermite Island, Cape Horn; moist places in the woods.

Cæspites late extensi, plani, pallide fusco-olivacei. Caules 1-2 unc. longi. Folia vix imbricata, patentia, ovata, apicem versus dilatata et in segmenta 2 subulata subcaudata fissa, laxe cellulosa. Stipulæ parvæ, cauli subæquilatæ. Folia perichætialia erecta, lateralibus emarginatis dentatisque, intermedio seu stipulari bifido integerrimo. Seta uncialis. Capsula ovalis.

This is one of the many southern forms of *Lophocolea* nearly related to one another, and to *J. bidentata* of Enrope; it differs from *J. secundifolia*, in the leaves being horizontally patent and not secund; from *J. diademata*, nob., of New Zealand, in the calyx and less spreading foliage; and from *J. physantha*, nob., of the same country, also by the totally different calyx, from all three by the divisions of the mouth of the latter organ being dentate. We are indebted to our friend M. Montagne for a specimen of his *J. coadunata*, so named by Nees, but which we do not consider to be the plant of Swartz.

PLATE CLIX. Fig. VI.-1, plant of the natural size; 2 and 3, leaves; 4 and 5, stipules; 6, calyx, seta, and capsule :-- all magnified.

47. JUNGERMANNIA *humifusa*, Hook. fil. et Tayl.; caule flaccido procumbente implexe ramoso, foliis approximatis subimbricatis horizontaliter patentibus planis oblongis antice gibbosis emarginato-dentatis integerrimis, stipulis bipartitis segmentis subulato-setaceis extus unidentatis quadripartitisve. Nobis *in Lond. Journ. Bot.* vol. iii. p. 472. (TAB. CLIX. Fig. V.)

HAB. Kerguelen's Land; on the rhizomata of Pringlea.

Cæspites lati, explanati, appressi, pallide virides. Caulis uncialis, irregulariter ramosus, ad stipulas radicans. Folia approximata, vix imbricata, basi latiora, decurrentia, sinu apice formæ irregulari. Calyx ovato-oblongus, trigonus; angulo unico alato, subdentato. Perigonia in spicas ovato-lanceolatas secus ramos obvias disposita, foliolis imbricatis, ventricosis.

Very nearly allied to *J. leptantha* and perhaps not distinct from it; though we have preferred separating species from such widely-severed localities as these affect, when, as in this case, they present tangible characters. Those of this species will be found in the more erect leaves, with lax areolæ, wide and decurrent bases, and more setaceous stipules.

PLATE CLIX. Fig. V .-- 1, plant of the natural size ; 2 and 3, leaves ; 4, stipule :-- magnified.

48. JUNGERMANNIA alternifolia, Hook. fil. et Tayl.; caule gracili laxe implexo procumbente parce ramoso, foliis flaccidis laxe reticulatis alternis patentibus planis triangulari-ovatis emarginatis decurrentibus segmentis spinoso-acuminatis integerrimis, stipulis minutis quadripartitis segmentis setaceis, calyce terminali triangulari-cylindraceo ore trilabiato ciliato. Nobis *in Lond. Journ. Bot.* vol. iv. p. 83. (TAB. CLXI. Fig. II.)

HAB. Falkland Islands; on moist banks near the sea.

Caspites huride virides, ramis substrictis. Folia basi decurrentia, laxe reticulata; perichatialia $\frac{1}{2}$ longitudine calycis, erecta, concava, subciliata. Capsula oblongo-rotundata.

Related to *J. humifusa*, but distinguishable by the deep division of the apex of the leaf, the longer segments, the more decurrent bases and the wider segments of the stipules. This species was erroneously described (Lond. Journ. Bot. l. c.) as a native of New Zealand; from whence we have never seen specimeus.

PLATE CLXI. Fig. II.-1, plant of the natural size; 2 and 3, branch and leaf; 4, lcaf; 5, stipule:-all magnified.

49. JUNGERMANNIA divaricata, Hook. fil. et Tayl.; caule implexo procumbente ramoso, foliis approximatis suberectis secundis e basi angusta oblongis convexis bifidis segmentis lanceolatis acuminatis divaricatis subflexuosis, stipulis bifidis segmentis subulatis extus unidentatis. Nobis *in Lond. Journ. Bot.* vol. v. p. 367. (TAB. CLXI. Fig. VIII.)

HAB. Hermite Island, Cape Horn; in tufts of mosses, &c.

Cæspites pallide flavo-virescentes. *Caules* 1 unc. longi, basin versus præcipne ramosi. *Folia* laxe imbricata, grosse reticulata, ad medium in segmenta dua acuminata divaricata fissa, margine dorsali decurrente et recurvo. *Perigonia* in spicas terminales disposita; foliolis arcte appressis, basi tumidis, antheriferis, segmentis foliis caulinis brevioribus.

Allied to J. leptantha, but a smaller plant, with leaves of a different shape, being narrower at the base and deeply divided beyond the middle. The stipules are bipartite.

PLATE CLXI. Fig. VIII.-1, plant of the natural size; 2 and 3, brauch and leaf; 4, leaf; 5, stipule :-- all magnified.

50. JUNGERMANNIA sabuletorum, Hook. fil. et Tayl.; minima, caule cæspitoso crassiusculo prostrato ramoso, ramis ascendentibus apice recurvis, foliis approximatis subremotisve erecto-patentibus secundis subquadratis angulis obtusis integerrimis apice retusis laxe cellulosis, stipulis minutis ovatis lanceolatisve bipartitis, segmentis subulatis incurvis. Nobis *in Lond. Journ. Bot.* vol. iii. p.469. (TAB. CLVIII. Fig. VIII.)

HAB. Falkland Islands; on wet sand and clay-slate.

Cæspiles sub 2 unc. lati, pallide flavo-virides. *Caules* breves vix $\frac{1}{4}$ unc. longi. *Rami* e caule prostrato erecti, curvati, demum horizontales. *Folia* versus apices ramorum laxe imbricata, parva, madore homomalla, apice plerumque retusa v. emarginata, rarius rotundata, basi late adnata, laxc cellulosa, paria ultima sæpissima appressa, apicibus ramulorum hine compressis. *Stipulæ* caule subangustiores.

Perhaps the most minute of the Lophocole α , from all the species of which its habit and the form of the leaves amply distinguish it.

PLATE CLVIII. Fig. VIII.—I, plant of the natural size; 2, portion of branch, leaves, and stipules; 3, leaf; 4, stipule:—magnified.

51. JUNGERMANNIA *rivalis*, Hook. fil. et Tayl.; flaccida, căule cæspitoso ascendenti v. erecto ramoso gracili, foliis distichis laxe imbricatis approximatisve inferioribus remotis teneribus flaccidis oblongoquadratis angulis obtusis integerrimis basi late adnatis decurrentibus apice retuso, stipulis ovatis bifidis segmentis integerrimis v. extus unidentatis. Nobis *in Lond. Journ. Bot.* vol. iii. p. 469. (TAB. CLVIII. Fig. VII.)

HAB. Falkland Islands; on wet rocks, &c., near the sea, abundant.

Cæspites laxi, basi sæpe submersi, atro-fusci, superne pallide olivacei. *Caulis* 1-2 unc. longus, ramosns, ramis erectis gracilibus, laxe foliosis. *Folia* tenerrima, membranacca et flaccida, minute arcolata, basi lata decurrente, oblongo-rotundata v. subquadrata, apice plerumque obscure retusa. *Stipulæ* caule vix latiores, basi subrotundatæ, v. late ovatæ, bifidæ; segmentis integerrimis uni-dentatisve.

Allied to *J. planiuscula* (Pt. 1. t. 65. f. 2), which is a larger plant, with differently shaped stipules. Also near the following, which, again, is smaller than either, with leaves of another form.

PLATE CLVIII. Fig. VII.-1, plant of the natural size; 2 and 3, leaves; 4 and 5, stipules :- magnified.

5 н

52. JUNGERMANNIA grisea, Nobis in Fl. Antarct. Pt. 1. p. 154. t. lxiv. f. 8.

Var. β , *laxa*; caule ramoso flexuoso, foliis laxius insertis subremotis alternantibus. (TAB. CLX. Fig. IV.)

HAB. Falkland Islands; on clay banks near the sea. Var. β , in similar situations.

Also a native of Lord Auckland's group.

PLATE CLX. Fig. IV.—1, plaut of the uatural size; 2, stem and leaves; 3 and 4, leaves; 5 and 6, stipules: magnified.

53. JUNGERMANNIA *reclinans*, Hook. fil. et Tayl.; caule prostrato implexo ramoso, foliis imbricatis patentibus siccitate explanatis madore secundis e basi lata ovato-rotundatis integerrimis apice rotundatis v. obscure retusis, stipulis 2–4-partitis, segmentis setaceis intermediis elongatis. Nobis *in Lond. Journ. Bot.* vol. iii. p. 470. (TAB. CLIX. Fig. I).

HAB. Falkland Islands; on wet rocks near the sea.

Cæspites laxe intertexti, pallide flavo-olivacei, inter Muscos Hepaticasque alias repentes. Caules supini, vix $\frac{1}{2}$ unc. longi, parce ramosi. Folia margine superiore sursum producto plerumque trilobo, rarius bilobo, lobis latis obtusis, folia nunc integra. Stipulæ basi augustatæ, quadratæ, bipartitæ, segmentis plerumque uui-dentatis omnibus setaceis articulatis incurvis.

With much affinity to the *J. multipenna* of Lord Aucklaud's group; but the upper margin of the leaf is not so gibbous or produced upwards, the stipules have the inner segments straight or recurved, but not reflexed, and much more slender.

54. JUNGERMANNIA secundifolia, Hook. fil. et Tayl.; parvula, caule subcæspitoso procumbente subramoso, foliis imbricatis erectis secundis oblongis emarginato-bifidis, segmentis lanceolatis integerrimis, stipulis bipartitis, segmentis inæqualiter bifidis laciniis subulato-sectaceis, calyce terminali oblongo trigono, ore trilaciniato laciniis dentatis. Nobis in Lond. Journ. Bot. vol. iii. p. 471. (TAB. CLIX. Fig. II.)

HAB. Falkland Islands; on tufts of mosses.

Cæspites parvi, inter *Muscos* intricati, pallide olivacei. *Caules* vix $\frac{1}{2}$ unc. longi, prostrati, demum ascendentes, apicibus supinis radicantibus. *Folia* imbricata, madore erecta et secunda. *Stipulæ* amplæ, segmentis setaceis incurvis. *Calyx* majusculus, trigonus, latere inferiore latiore, ore ciliato-dentato.

This in some respects approaches the British *J. bidentata*, but is even more like *J. helerophylla*, from which it may eventually prove not distinct; its claims rest on the closely imbricated and secund leaves, and more entire segments of the stipules.

PLATE CLIX. Fig. II.—1, plant of the natural size; 2, apex of stem, perichætium, and calyx; 3, leaf; 4, stipule:—magnified.

55. JUNGERMANNIA *subviridis*, Hook. fil. et Tayl.; parvula, caule cæspitoso prostrato ramoso, foliis laxe imbricatis secundis erecto-patentibus erectisve oblique obovatis quadratisve emarginato-bifidis segmentis obtusiusculis, margine anteriore gibboso, inferiore decurrente, stipulis ovatis bifidis utrinque uni-dentatis. Nobis *in Lond. Journ. Bot.* vol. iii. p. 473. (TAB. CLIX. Fig. IV.)

HAB. Hermite Island, Cape Horn; on the ground.

Cæspiles lati, 2 une. diametro, pallide virides, intricati. Caules vix $\frac{1}{2}$ une. longi, decumbentes v. prostrati, apicibus ascendentibus. Folia remotiuscula, margine superiore sursum gibboso; substantia dense celludosa.

Closely allied to the *J. discedens*, Nees, of the East Indies; but the leaves are shorter, wider, have a deeper sinus and more acute segments; and the stipules are not so slender.

PLATE CLIX. Fig. IV.—1, plant of the natural size; 2, branch, with leaves and stipules; 3 and 4, leaves; 5 and 6, stipules :—all magnified.

56. JUNGERMANNIA trachyopa, Hook. fil. et Tayl.; parvula, caule implexe ramoso procumbente flaccido, foliis tenerrimis laxe cellulosis imbricatis erectis subsquarrosis latissime rotundato-quadratis profunde bitrifidis grosse inæqualiter spinuloso-dentatis, stipulis late ovatis bifidis segmentis acuminatis grosse dentatis, calyce terminali oblongo trigono, ore laciniato-dentato. Nobis *in Lond. Journ. Bot.* vol. iii. p. 471. (TAB. CLIX. Fig. III.)

HAB. Hermite Island, Cape Horn; growing in the woods on Anthoceros punctatus, L.

Caules vix $\frac{1}{2}$ unc. longi, irregulariter ramosi, pallide virides. Folia arete imbricata, in lacinias duas v. plerumque plures lanceolato-subulatas divisa. Calyx pro planta majusculus. Capsula ovalis. Sela cauli æquilonga.

A very distinct species from any of the forcegoing, and a beautiful object under the microscope from the delicacy and reticulation of its leaves.

PLATE CLIX. Fig. III.—1, plant of the natural size; 2, stem and leaf; 3-6, stipules; 7, perichætium, calyx, seta, and capsule :--all magnified.

57. JUNGERMANNIA *triacantha*, Hook. fil. et Tayl.; caule implexo procumbente vage ramoso, foliis planis approximatis patentibus oblongo-ovatis trifidis segmentis subulato-lanceolatis, stipulis subquadratis bifidis segmentis bilaciniatis laciniis subulatis. Nobis *in Lond. Journ. Bot.* vol. v. p. 368.

HAB. Falkland Islands; on the ground.

Cæspiles densi, luride olivacei. *Caulis* uncialis, ramis paucis patentibus. *Folia* basi vix imbricata, patentia, divaricata, oblonga, apices versus sinubus duobus excisis aueta; laciniis 3 subulatis, porrectis, subparallelis. *Stipulæ* liberæ, crecto-patentes, quadrifidæ, segmentis subulatis.

Most nearly allied to *L. chlorophylla*, nobis, of New Zealand, which is, however, a smaller plant and has shorter wider subsecund leaves, and rounded dentate stipules,

(7. CHILOSCYPHUS, Nees.)

58. JUNGERMANNIA *pallido-virens*, Hook. fil. et Tayl.; majuscula, caule implexe subramoso procumbente, foliis patentibus imbricatis late ovato-oblongis apice retusis integerrimis margine anteriore recurvo, stipulis minutis recurvis oblongis bifidis segmentis subulatis extus unidentatis quadrifidisve, calyce in ramo abbreviato terminali oblongo tri-alato compresso apice laciniato-ciliato. Nobis *in Lond. Journ. Bot.* vol. iii. p. 473. (TAB. CLIX. Fig. IX.)

HAB. Hermite Island, Cape Horn; on the ground near the sea.

Cæspites late extensi, pallide flavo-virescentes, demum fuscescentes, interdum (status minor) omnino fuscati. Caules 2 une. longi, parce ramosi, ramis $\frac{1}{8}$ une. latis. Folia dense reticulata, arcolis minutis, laxe imbricata, erectopatentia, ope stipulæ basi connexa, apice rotundata seu truncata, unidentata v. emarginata. Stipulæ minimæ, caulis $\frac{1}{2}$ latitudine, concavæ, recurvæ, oblongæ; segmentis setaceis extus unidentatis. Perichælium ramum abbreviatum terminans, e paribus 2-3 foliolorum erectorum appressorum constans, foliolo interiore 4-5-partito. Calyx oblongocampanulatus, latere unico profunde fissus. This handsome plant resembles the *C. Endlicherianus*, Nees, of Norfolk Island, more than any other species; differing, however, materially in its great size, the less rounded tops of the leaves, their more convex figure, their perfectly entire margins, and by the less laciniated stipules.

PLATE CLIX. Fig. IX.—1 and 2, plants of the natural size; 3, stem and leaf; 4, back view of ditto and stipules; 5, stipule; 6, perichaetial leaf; 7, calyx:—magnified.

59. JUNGERMANNIA grandifolia, Hook. fil. et Tayl.; caule procumbente subsimpliei laxe implexo, surculis planis, foliis arcte imbricatis patentibus quadrato-rotundatis antice basi gibbosis margineque recurvis integerrimis, stipulis minutis sub 4-laciniatis. Nobis in Lond. Journ. Bot. vol. iii. p. 474. (TAB. CLIX. Fig. VIII.)

HAB. Hermite Island, Cape Horn; in the woods.

Cæspites 3-4 unc. longi, $\frac{1}{6}$ unc. lati, superne pallide virescentes, inferne rufo-brunnei. Folia ampla, margine superiore basi præcipne recurvo, inferiore basi simplici non decurrente, flaccida, crassiuscula, pellucida, areolis parvis, rarius cum stipulis imo basi connexa, plerumque libera.

The largest and handsomest species of *Chiloscyphus*, in which the disproportion between the leaves and stipules is very remarkable. The broader and shorter leaves, their larger areolæ and more laciniated stipules, are alone sufficient to distinguish it from the former.

PLATE CLIX. Fig. VIII.--1, plant of the natural size; 2, back of stem, stipules, and leaves; 3, stem and leaf; 4, stipule:-magnified.

60. JUNGERMANNIA *fusco-virens*, Hook. fil. et Tayl.; caule implexo procumbente subramoso, surculis ascendentibus, foliis imbricatis verticalibus patentibus secundis rotundatis integerrimis, stipulis bi-quadripartitis, segmentis radiantibus, calyce in ramo brevi terminali oblongo-campanulato triplicato, ore truncato integro. Nobis *in Lond. Journ. Bot.* vol. iii. p. 474. (TAB. CLIX. Fig. VII.)

HAB. Hermite Island, Cape Horn; forming dense tufts on the tops of the mountains, alt. 1,700 feet.

Cæspites parvi, densi, rufo-brunnei, ramis junioribus virescentibus. *Caules* 2 unc. longi, secus totam longitudinem radiculas dense fasciculatas demittentes. *Folia* planiuscula, paria opposita basi valde approximata, opaca, crassinscula, cellulis parvis. *Stipulæ* basi breves, in segmenta dua v. plura subulata setaceave fissa. *Calyces* bini v. plures, folüs pallidiores. *Seta* fere uncialis. *Capsula* oblongo-rotundata.

The narrow segments of the differently shaped stipules and calyx afford the best means of distinguishing between this, and *J. australis*, nob., of Campbell's Island. The latter is also a smaller plant, of a darker colour. The size, large arcoke of the leaves, and their not being connate at the base with the stipules, at once remove the present from the following species.

PLATE CLIX. Fig. VII.—1, plant of the natural size; 2, stem and leaf; 3, back view of ditto and stipules; 4, stipule; 5, perichætium, calyx, seta, and capsule:—magnified.

61. JUNGERMANNIA surrepens, Hook. fil. et Tayl.; caule disperso simplici repente, foliis imbricatis patentibus rotundatis integerrimis stipula ovata subquadrifida connatis. Nobis in Lond. Journ. Bot. vol. iii. p. 475. (TAB. CLX. Fig. I.)

HAB. Hermite Island, Cape llorn; on J. Magellanica.

Caules plerumque subsolitarii, supiui, pallide brunnei v. albidi. *Fotia* opposita, dorso ope stipulæ basi connexa. *Stipula* caule vix latior, ovata, bifida, segmentis subulatis extus dente majuscula auetis.

Near the last, but a very different plant in size, and in the disposition of its leaves.

PLATE CLX. Fig. I.—I, plant of the natural size; 2, back view of stem, leaves and stipules; 3, stem and leaf; 4, stipule:—magnified.

62. JUNGERMANNIA retusata, Hook. fil. et Tayl.; caule implexo procumbente subsimplici rectiusculo, foliis patentibus planis late oblongis obtusis retusisque integerrimis hinc stipulæ minutæ setacco-bipartitæ connexis. Nobis in Lond. Journ. Bot. vol. iv. p. 84. (TAB. CLXI. Fig. III. sub. nom. J. reclinatæ.)

HAB. Falkland Islands; on the ground.

Cæspites laxi, pallide fusco-olivacei. Caules unciales. Folia remotiuscula, late oblonga, obscure emarginata, cum stipula caule æquilata connexa.

Allied to the *Ch. integrifolius*, Gottsche, of Chili, but the leaves are more distant, shorter, and wider; the stipule more divided and the whole plant of a darker colour.

PLATE CLXI. Fig. III. (under the name of J. reclinata).-1, plant of the natural size; 2, 3, and 4, stem and leaves; 5, stipule: -magnified.

63. JUNGERMANNIA horizontalis, Hook., Musc. Exot. t. 96. Gottsche, Lindb. et Nees, Syn. Hep. p. 178. HAB. Staten Land; Menzies (in Herb. Hook.).

Not in the collections of the Antarctic Expedition.

64. JUNGERMANNIA amphibolia, Nees, in Martius, Flor. Bras. vol. i. p. 334. Gottsche, Lindb. et Nees, Syn. Hep. p. 178.

HAB. Hermite Island, Cape Horn; mixed with J. uncialis.

Also a native of the Brazils.

(8. LEPIDOZEA, Nees.)

65. JUNGERMANNIA tetradactyla, Hook. fil. et Tayl.; in Fl. Antarct. Pt. 1. p. 158. Gottsche, Lindb. et Nees, Syn. Hep. p. 213.

HAB. Hermite Island, Cape Horn; moist places near the sea.

Likewise found both in Lord Auckland's group and New Zealand.

66. JUNGERMANNIA plumulosa, Lehm. et Lindb., Pugill. p. 30. Gottsche, Lindb. et Nees, Syn. Hep. p. 211.

HAB. Staten Land, Menzies. Strait of Magalhaens, D'Urville. Hermite Island, Cape Horn; on moist banks.

67. JUNGERMANNIA lævifolia, Hook. fil. et Tayl.; in Fl. Antarct. Pt. 1. p. 157. Gottsche, Lindb. et Nees, Syn. Hep. p. 208.

HAB. Falkland Islands; on moist rocks near the sea.

First described from Auckland Island specimens; also found in New Zealand and Tasmania.

68. JUNGERMANNIA oligophylla, Lehm. et Lindh., Pugill. vi. p. 26. Gottsche, Lindh. et Nees, Syn. Hep. p. 201.

IIAB. Staten Land, Menzies. Hermite Island, Cape Horn; in moist places.

69. JUNGERMANNIA tridactylis, Lehm. et Lindb.? fid. Montagne, in Voy. au Pole Sud, Bot. Crypt. p. 243. HAB. Strait of Magalhaens; M.M. Hombron et Jacquinot.

Fuegia, the

70. JUNGERMANNIA filamentosa, Lehm. et Lindb., Pugill. vi. p. 29. Montagne in Voy. an Pole Sud, Bot. Crypt. p. 246.

HAB. Strait of Magalhaens; M. Hombron.

A plant we do not recognize amongst the numerous forms, from Fuegia and Lord Auckland's Island, of this most difficult, and perhaps too extended group.

71. JUNGERMANNIA Javanica, Mont., in Foy. au Pole Sud, Bot. Crypt. p. 246.

HAB. Strait of Magalhaens; Port Famine, M. Jacquinot.

This may be one of the above enumerated species, though we have failed in identifying it. It is also a native of Java.

72. JUNGERMANNIA chordulifera, Tayl., in Lond. Journ. Bot. vol. v. p. 371. (TAB. CLXI. Fig. VI.)

HAB. Chonos Archipelago, C. Darwin, Esq.

A very handsome species, allied to the J. pendulina of New Zealand.

PLATE CLXI. Fig. VI.-1, plant of the natural size; 2, stem, leaves, and stipules; 4, stipule :- magnified.

(9. MASTIGOPHORA, Nees.)

73. JUNGERMANNIA hirsuta, Nees; Fl. Antarct. Pt. 1. p. 160. Sendtnera ochrolenca, Nees, in Gottsche, Lindb. et Nees, Syn. Hep. p. 240.

HAB. Hermite Island, Cape Horn; on roots of stunted trees, &c., alt. 1,000 feet. Falkland Islands; rocks on the hill tops, rare.

A widely distributed plant, being found in Mexico and Java, at the Cape of Good Hope, and Lord Auckland's group.

(10. RADULA, Nees.)

74. JUNGERMANNIA *Helix*, Hook. fil. et Tayl.; parvula, caule repente implexo subflexuoso pinnatim ramoso, foliis remotis alternis oblongis alte concavis integerrimis basi gibbosis, lobo superiori ovato-oblongo obtuso, inferioris ovati tumidi involuti apice subacuto superiori appresso. Nobis *in Lond. Journ. Bot.* vol. iv. p. 475. Gottsche, Lindb. et Nees, *Syn. Hep.* p. 260. (TAB. CLX. Fig. II.)

HAB. Hermite Island, Cape Horn; growing with J. colorata.

Caspites vix 1/4 unc. lati, pallide straminei. Folia pauca, alterna, nisi apicem caulis versus remota.

A very distinct little species, found growing on large masses of J. colorata, with the purple colour of which its pale stems contrast conspicuously. It is smaller and has more tunid leaves than any of its congeners; the latter resemble in form the shell of *Helix putris*, whence the trivial name.

PLATE CLX. Fig. II.--1, plant of the natural size; 2, stem and leaves; 3 and 4, leaves :--magnified.

75. JUNGERMANNIA physoloba, Mont.; Fl. Antarct. Pt. 1. p. 161. J. flavifolia, nobis in Lond. Journ. Bot. vol. iv. p. 476. Gottsche, Lindb. et Nees, Syn. Hep. p. 259. J. complanata, β, Hook. (TAB. CLX. Fig. III.)

HAB. Hermite Island, Cape Horn; on trunks of trees.

An abundant boreal plant, inhabiting Europe from Switzerland to Iceland. In the southern hemisphere it has hitherto been seen only in Lord Auckland's group and at Cape Horn. It is rather a variable species, and we have

PLATE CLX. Fig. III.—1, plant of the natural size; 2, branch and leaf; 3, leaf; 4, perichætium and calyx; 5, ealyx and capsule :—magnified.

(11. POLYOTUS, Gottsche.)

76. JUNGERMANNIA Magellanica, Lamk. Fl. Antarct. Pt. 1. p. 162.

HAB. Hermite Island, Cape Horn; on trunks of trees, abundant. Strait of Magalhaens, Commerson. Staten Land, Menzies.

Also found in Campbell's Island, Tasmania, and New Holland.

77. JUNGERMANNIA Menzicsii, Hook., Musc. Exot. t. 118.

HAB. Hermite Island, Cape Horn; mossy trunks of trees, wet rocks, &c.; also on the summits of the mountains. Staten Land, *Menzies*.

JUNGERMANNIA palpebrifolia, Hook., Musc. Exot. t.71. Gottsche, Lindb. et Nees, Syn. Hep. p. 246.
HAB. Strait of Magalhaens, D'Urville.

(12. FRUILANIA, Raddi.)

79. JUNGERMANNIA cyperoides, Schwaeg., Prodr. Hep. 14. Gottsche, Lindb. et Nees, Syn. Hep. p. 420. HAB. Strait of Magalhaens; (fid. Schwaegrichen).

80. JUNGERMANNIA lobulata, Hook., Musc. Exot. t. 119. Gottsche, Lindb. et Nees, Syn. Hep. p. 445. HAB. Hermite Island, Cape Horn; in woods. Staten Land, Menzies. Falkland Islands; on rocks near the hill tops.

S1. JUNGERMANNIA Magellanica, Spreng., in Annal. des Wetter. Ges. vol.i. p. 25. t.4. f. 10, (fid. Gottsche, Lindb. et Nees, Syn. Hep. p. 447.)

HAB. Strait of Magalhaens; on Drimys Winteri and Berberis ilicifolia; Forster (fid. Gottsche).

(13. LEJEUNIA, Spreng.)

S2. JUNGERMANNIA subintegra, Hook. fil. et Tayl.; caule breviusculo cæspitoso procumbente elongato subsimpliei, foliis subimbricatis erectiusculis integerrimis, lobo superiorc oblongo-rotundato, inferiore $\frac{1}{3}$ breviore tumido involuto angulo superiore acuminato, stipulis caule paulo latioribus ovatis acutis integerrimis v. summo apice fissis. Nobis in Lond. Journ. Bot. vol. iv. p. 477. Gottsche, Lindb. et Nees, Syn. Hep. p. 377. (TAB. CLX. Fig. V.)

HAB. Falkland Islands; in wet places near the sea.

Cæspites late extensi, pallide olivacei. *Caules* vix $\frac{1}{2}$ unc. longi, plerumque simplices. *Folia* amplectantia, suberecta, concava, diametro caulis duplo latiora, laxe cellulosa, areolis majusculis. *Stipulæ* majusculæ, integerrimæ, v. imo apice solum fissæ, segmentis approximatis.

The great size of the stipules comparatively to the leaves and their very obscure division, afford sufficiently distinctive characters of this species.

PLATE CLX. Fig. V.-1, plant of the natural size; 2, stcm; 3, lcaf; 4 and 5, stipules :-magnified.

83. JUNGERMANNIA parasitiea, Hook. fil. et Tayl.; caule subimplexo procumbente pinnatim ramoso, foliis subapproximatis patentibus valde concavis integerrimis v. obscure dentatis, lobo superiore triangulariovato acuto v. acuminato apice subrecurvo, inferiore oblongo acuminato, stipulis parvis obovato-quadratis bilobis lobis rotundatis integerrimis. Nobis *in Lond. Journ. Bot.* vol. iii. p. 477. J. marginalis, *nobis*, *l. c.* vol. iv. p. 91. (TAB. CLX. Fig. VI.)

HAB. Hermite Island, Cape Horn; parasitical on J. uneialis, and amongst lichens on trunks of trees.

Caules gracillimi, minimi, oculo nudo invisibiles, pallide virides, inter Lichenes aliasque Hepaticas dispersi, $\frac{1}{10}$ une. longi, subpinnatim ramosi, ramis suberectis. Folia subremota, patentia, basi latiusenla, apicibus acuminatis, incurvis v. recurvis, lobo inferiore margine involuto subundulato.

This has precisely the habit and appearance of the Irish J. orata, Tayl. MSS., an equally minute parasite, with stipules of the same form. The present differs from that in the shape of the more distant leaves, which are shorter, wider at the base, and more acute above. The whole plant is of a paler colour, and from the smaller size of the cells of the leaves their tissue is more dense.

PLATE CLX. Fig. VI .--- 1, plant of the natural size ; 2, stem, leaves, and stipules ; 3, leaf ; 4, stipule :-- magnified.

84. JUNGERMANNIA rufescens, Lindb., in Gottsche, Lindb. et Nees, Syn. Hep. p. 366.

HAB. Staten Land, Menzies (in Herb. Hook.).

(14. DIPLOLÆNA, Nees.)

85. JUNGERMANNIA pisicolor, Hook. fil. et Tayl.; fronde laxe eæspitosa ereeta dichotome divisa basi in stipitem teretem gradatim attenuata, lobis linearibus obtusis emarginatis uninerviis integerrimis. Nobis in Lond. Journ. Bot. vol. iii. p. 478. (TAB. CLX. Fig. VII.)

HAB. Hermite Island, Cape Horn; at the bottom of an alpine lake.

Frondes e radice repente elongato crecti, sub 3 une. lati, pisicolores, demum flavescentes, parte inferiore atra, bis terve dichotome divisi, lobis plerumque madore concavis, e margine proliferi ; juniores basi rotundati, primum liberi?, demum radices emittentes. *Nervus* validus, percurrens, siecitate tenuis, albescens, madore dilatatus, fronde concolor, axillis acutis.

A very distinct and curious plant, allied to the *J. tenuinervis*, nob., of New Zealand, from which it may be known by its yellow-green colour, greater size, and taller habit, narrower and more elongated lobes, and, above all, by its being creet, differing remarkably in that respect from its northern allies, *J. Hibernica*, Hook., and *J. Lyellii*, Hook.

PLATE CLX. Fig. VII.-1, plant of the natural size; 2, apex of frond :--magnified.

(15. ANEURA, Nees.)

86. JUNGERMANNIA multifida, Linn.; Fl. Antaret. Pt. 1. p. 166.

Var. *β. submersa*; fronde anguste lineari-elongata pellueida parce ramosa plana, ramis brevibus, perigoniis marginalibus alternis brevissimis, cellulis densis.

Var. γ . nana; parvula, ramosa, cellulis latioribus brevioribusque.

HAB. Hermite Island, Cape Horn; var. a and β , very abundant; var. γ , in a fresh-water lake amongst the mountains. Falkland Islands; abundant.

An extremely abundant plant in the southern extra-tropical regions.

87. JUNGERMANNIA pinguis, L. Hook. Brit. Jung. t. 46.

HAB. Hermite Island, Cape Horn; wet places in the woods.

88. JUNGERMANNIA alcicornis, Hook. fil. et Tayl.; fronde cæspitosa erecta alterne bipinnatim ramosa, caule ramisque linearibus laciniis lobisve brevibus subtruncatis, calyptris lateralibus linearibus albidis scabridis apice laceris. Nobis in Lond. Journ. Bot. vol. iii. p. 479. (TAB. CLX. Fig. VIII.)

HAB. Hermite Island, Cape Horn; mossy places in the woods.

Frondes cæspitosi, planiusculi, unciales, circumscriptione oblongi, juniores læte virides, demum fuscescentes et siccitate nigrescentes. *Rami* scu laciniæ basi subdichotomi, superne subpinnatim divisi; lobudis brevibus, obtusis. *Calyptræ* parte frondis inferiore laterales, valde elongatæ, cylindraceæ, basi curvatæ, carnosæ, papillosæ. *Seta* sub $\frac{1}{2}$ unc. longa. *Capsula* cylindracea.

A very beautiful little species, allied to the *J. palmata*, Hoffm., of Europe; but much more divided, with the divisions pinnate and not palmate; the lobules also are much narrower, and the calyptra is borne higher up in the frond.

PLATE CLX. Fig. VIII.-1, plant of the natural size; 2, ditto; 3, branch and calyptra:-magnified.

(16. METZGERIA, Nees.)

89. JUNGERMANNIA furcata, L. Flor. Antarct. Pt. 1. p. 167.

Var. B. pubescens, J. pubescens, Brit. Jung.

HAB. Hermite Island, Cape Horn; in woods, &c.; both varieties abundant.

90. JUNGERMANNIA *prehensilis*, Hook. fil. et Tayl.; fronde laxe cæspitosa, ramis erectis incurvis alatis, lobis secundis alternis pinnatis, pinnulis linearibus planis crassinerviis, calyptra e basi anguste elongatoobovata basi squamosa, perigoniis clavatis. Nobis *in Lond. Journ. Bot.* vol. iii. p. 480. (TAB. CLX. Fig. IX.)

HAB. Hermite Island, Cape Horn; on moist banks near the sea.

Frondes laxe cæspitosi, apicibus latiusculis hamatis. *Caulis* planus, brunneus, pubescens, pinnulis glabratis pallide olivaceis. *Calyptræ* brunneæ, apices versus frondis laciniarum basi superficie inferiore sitæ, binc occlusæ, interdum binæ. *Perigonia* plantis aliis obvia, clavata, c ramo pinnato frondis constantia, lobulis pinnisve incurvis singulis antheram majnsculam sphericam pedicellatam foventibus.

A remarkably distinct and fine species, most resembling the *J. eriocaula*, Hook., of New Zealand; though the frond is of a darker colour aud tripinnate, the pinnules much narrower, and the whole plant more elongated and divided. The perigonia and calyptræ occupy similar positions on different plants.

PLATE CLX. Fig. IX.—1, plant of the natural size; 2, upper, and 3, under surface of branch with calyptra; 4, calyptra :—magnified.

(17. NOTEROCLADA, Tayl.)

Involucrum apicem versus frondis concavi tumidi inflatum, ore libero sub-bilobo. Capsula quadrivalvis, seu irregulariter rumpens, pedicellata. Elateres spirales seminibus immixtæ. Antheræ fronde immersæ. Frons pinnatim lobata, v. fohis basi latissimis longe decurrentibus subspiraliter dispositis ornata.—Stirps inter Jungermannias foliosas frondosasque quasi media, his tamen accedens.—Androcryphia, Gottsche.

91. JUNGERMANNIA confluens, Tayl. in Lond. Journ. Bot. vol. iii. p. 478. (TAB. CLXI. Fig. VII. in part.)

HAB. Hermite Island, Cape Horn; on the bare ground in woods. Falkland Islands and Christmas Harbour, Kerguelen's Land; on moist banks.

Laxe cæspitosa. Frondes flaccidissimi, 2 unc. longi, erecti. Caulis simpliciusculus, foliis imbricatis omnino occlusus, subrufescens, gracilis. Folia alterna, tenerrima, madore carnosiuscula, siccitate membranacea, ægre resuscitentia, et inter se quasi confluentia (hinc frons prima visu continua et lobata), basi latissima, bis latiora quam longa, longe decurrentia, fere amplexicaulia, semi-orbicularia, apice rotundata v. retusa, areolis majusculis. Involucrum terminale, sessile, erectum, cylindraceum, compressum, ore eroso-dentato. Seta uncialis, gracilis. Capsula ovato-globosa, irregulariter v. regulariter rumpens. Elateres brevissimi. Calyptra irregulariter rupta, parte superiore stylo persistente terminata. Anlheræ ovato-oblongæ, biseriales, substantia frondis immersæ, liquido oleaginoso scatentes.

A genus allied to *Fossombronia*, but the structure of the involuce, apparently formed very much out of the frond itself, is quite dissimilar. The involuce is terminal in this species, but lateral in a Brazilian congener, which was long regarded as identical, and smooth; when terminal, winged from the adhesion to its surface of the upper abbreviated leaves: it is either truncated or obscurely two-lipped. The young spores are united by fours in a transparent membrane. The drawing of the fruit is taken from Brazilian specimens of an allied species, or perhaps variety, collected by Mr. Gardner; the leaves of the Antarctic plant having become so firmly united under pressure, that no maceration would separate them satisfactorily.

In the 'Synopsis Hepaticarum' of Nces, Lindenberg and Gottsche, the generic name has been changed to *Androcryphia*, with the following explanation. "Noteroclada nomen Græce cum sonet neque Græci esse possit originis, (seil. $\nu\omega\tau\sigma\sigma$ tergum non dat $\nu\omega\tau\epsilon\rho\sigma\nu$, neque $\chi\lambda\dot{a}\delta\sigma\nu$ sive rami character hoc loco succurrit); substituere aliud nomen ægre id quidem mecum sustinui." (*l. c.* p. 470.) The derivation of the name being, however, $\nu\sigma\tau\eta\rho\dot{o}s$ "madidus," sufficiently vindicates the adoption of Noteroclada.

PLATE CLXI. Fig. VII. (in part.)—1, Brazilian, and 2, Falkland Island specimen, of the natural size; 3, branch, leaves, &c., of the Brazilian specimen; 4, leaf of ditto; 5, corolla of ditto:—magnified.

(18. FOSSOMBRONIA, Nees.)

92. JUNGERMANNIA pusilla, L.

HAB. Kerguelen's Land; on banks amongst moss, &c.

Also a native of New Zealand, and probably not an uncommon plant in the temperate parts of the Southern as it is of the Northern hemisphere.

2. MARCHANTIA, March.

1. MARCHANTIA polymorpha, L. Flor. Antarct. Pt. 1. p. 168.

HAB. Fuegia, the Falkland Islands, and Kerguelen's Land; very abundant.

This is perhaps the most widely dispersed of *Hepaticæ*, ranging from the Arctic circle to the 57th degree of south latitude.

3. ANTHOCEROS, Michel.

1. ANTHOCEROS punctatus, L.

HAB. Hermite Island, Cape Horn, and the Falkland Islands; very common.

LIV. FUNGI, L.

(By the Rev. M. J. BERKELEY.)

1. AGARICUS, L.

1. AGARICUS *longinquus*, Berk.; pileo obliquo suborbiculari albo demum pallide fusco glabro nitente, strato superiore gelatinoso, stipite curto quandoque brevissimo pallide flavo-fusco basi albo-floccoso, lamellis albis subfurcatis, interstitiis lævibus. (TAB. CLXIII. Fig. V.)

HAB. Hermite Island, Cape Horn; on dead wood near the sea.

Pileus $\frac{3}{4}$ unc. et ultra latus, tenuis, oblique ovatus vel semiorbicularis, primum subtiliter pruinosus, mox autem glaberrimus, nitens; margine striato; stratum epidermale gelatinosum. *Stipes* vix liueam exsuperans quandoque obsoletus, æqualis, primum ceutralis, deinde excentricus aut omnino lateralis, subtiliter pruinosus, demum glaber, pallide flavo-fuscus, ad matricem basi tomentosa affixus. *Lamellæ* albæ, distantiusculæ, subfurcatæ, prope marginem præsertim subventricosæ, decurrentes, interstitiis lævibus, non reticulatis.

The nearest ally of this species is perhaps Ag. mitis, Pers., but the stem is not dilated upwards and the pileus is seldom perfectly lateral. Its colour, too, is different, and it has not the same opake appearance when dry. The upper stratum is gelatinous, though the pileus is dry externally; a character common to several closely allied species. It resembles also some smaller forms of Ag. algidus, Fr., but that is at first resupinate, whereas in the present species the stem is normal, though, as in most of the smaller excentric Agarics, occasionally obsolete. I do not know any other species with which it is necessary to compare it.

PLATE CLXIII. Fig. V.—1, Agaricus longinquus, Berk., of the natural size, from rather young specimens; 2, another :---magnified.

2. AGARICUS *exquisitus*, Berk.; minimus, tenerrimus, pileo ferrugineo subtiliter pulverulento, stipite brevi filiformi sursum incrassato pulverulento pileo concolori, basi dilatata irregulariter floccoso-membranacea, lamellis paucissimis ventricosis subliberis albo-marginatis interstitiisque lævibus ferrugineis.

HAB. Port Louis, Falkland Islands; on stems of Chiliotrichum amelloides.

Pileus membranaceus, 1 liu. latus, orbicularis, subtiliter pudverulentus, ferrugineus. *Stipes* 2 lin. altus, filifornis, sursum incrassatus, flexuosus, pulverulentus, pileo concolor, basi disco irregulari flexuoso-membranaceo affixus. *Lamellæ* paucæ (6), subliberæ, ferrugiueo-fuscæ, interstitüs latis lævibus.

This elegant little Agaric differs from Ag. horizontalis, Bull., an imperfectly known species, which I have received from Dr. Léveillé and have myself gathered on the trunks of trees in the park at Burleigh, near Stamford, in being altogether more delicate, in the membranaccous pileus, and especially in the absence of smaller gills between the larger. The pileus of Ag. horizontalis also is much darker when dry.

3. AGARICUS *Glebarum*, Berk.; pileo carnoso late umbonato glabro pallide fusco, stipite brevi glabro sursum subinerassato solido, basi mycelio floccoso affixa, lamellis latiusculis adnatis horizontalibus fulvis. (TAB. CLXII. Fig. III.)

HAB. Berkeley Sound, Falkland Islands; growing out of tufts of *Bolax* on the hills, where it endures a great degree of wet and cold.

Pileus $\frac{3}{4}$ uuc. latus, glaberrimus, subhemisphericus, umbonatus, pro ratione valde carnosus, senectute rugosus ; nequaquam viscidus. *Stipes* $\frac{3}{4}$ unc. altus, $1\frac{1}{2}$ lin. crassus, solidus, subtiliter fibrillosus, sursum incrassatus, basi ad surculos foliaque marcida mycelio floccoso affixus. Lamellæ latiusculæ, horizontales, dente obscuro adnatæ, subdistantes, quandoque furcatæ. Sporæ oblique ellipticæ, sub lente aureo-fuscæ, nucleo parvo globoso. Margo pilei interdum reflexus discusque exinde depressus, tunc ctiam lamellæ, ni caute perscrutentur, liberæ habcantur.

This species is allied to Ag. innocuus, Tasch, and Ag. cerodes, Fr. From the latter it differs in its solid stem; from the former, in its decidedly carnose unbonate pilcus.

PLATE CLXII. Fig. III.—1, Ag. Glebarum, Berk., of the natural size, on tufts of Bolax; 2, vertical section of the same; 3, spores :—highly magnified.

4. AGARICUS fascicularis, Huds., Fl. Angl. p. 615. Fries, Ep. p. 222.

HAB. Hermite Island, Cape Horn; on the trunk of a dead tree, Mr. Davis.

A single specimen only was found.

5. AGARICUS papilionaceus, Bull., t. 561. f. 2. Pers. in Freyc. Voy. p. 168. Ag. fimetarius, Gaud. in Ann. des Sc. Nat. vol. v. p. 97.

HAB. Falkland Islands; Gaudichand.

This species was not met with during the visit of the Erebus and Terror. Ag. Glebarum could not have been considered as belonging to the subdivision of Coprini. The pileus is said by Persoon to be broader, and the stem shorter than in Bulliard's figure.

2. COPRINUS, Pers.

1. COPRINUS *Flosculus*, Berk.; minimus, tenerrimus, pileo glabro ovato demum expanso hemispherico fisso sulcato vertice depressiusculo, stipite brevi, lamellis liberis paucis linearibus remotis. (TAB. CLXII. Fig. II.)

HAB. Berkeley Sound, Falkland Islands; on dung.

 $Pileus 1\frac{1}{2}$ lin. altus, 1 lin. latus, ovatus, profunde ex ipso vertice sulcatus, interstitüs striatis, glaber, subgriseus, margine crenulato, demum expansus, hemisphericus, fissus. $Stipes \frac{1}{2}$ lin. altus, filiformis, primum leviter ad basin turgidus, demum æqualis. Lamellæ primariæ subdecem, liberæ, remotæ, lineares. Sporæ ovatæ, atro-purpureæ.

A minute Coprinus, belonging to the same section with Coprinus Hemerobius, but differing from it and from the other species of the section in various characters. It resembles in habit C. Hendersonii, Berk., but wants the ring which is characteristic of that species.

PLATE CLXII. Fig. II.—1, Coprinus Flosculus, of the uatural size; 2, ditto :—magnified; 3, hymenium, viewed vertically with the spores on the sporophores; 4, spores :—highly magnified.

3. POLYPORUS, Fries.

1. POLYPORUS versicolor, Fr., Ep. p. 473. Berk. in Ann. Nat. Hist. vol. iv. p. 292.

HAB. Falkland Islands; on the underside of timber, C. Darwin, Esq.

This can scarcely be considered indigenous. The mycelium in all probability existed on the timber when imported.

4. CORTICIUM, Fries.

1. CORTICIUM tremellinum, Berk.; confluenti-effusum, gelatinosum, pellucidum, candidum, quandoque opacum, subtiliter pruinosum, siccum non rimosum decoloratum.

HAB. Hermite Island, Cape Horn; on bark of the Deciduous Beech in damp woods.

Primum maculas orbiculares exhibens, quæ demum confluendo areolas tenues longe effusas omnes matricis inæqualitates observantes efficiunt; album, ut plurimum pellucidum, quandoque opacum, tenue gelatinosum, subtiliter pruinosum atque exinde nitidulum, inodorum, insipidum; exsiccatum sordide umbrinum. Margo tenuis, nequaquam fimbriatus, hic illic exsiccatione liber. *Sporæ* ellipticæ, majores.

Nearly allied to *Corticium viscosum*, but not in the least cracked when dry. I have found the same species, apparently, in Sherwood Forest, which I had referred to *C. viscosum*; but the characters given by Fries, in his 'Epicrisis,' indicate a distinct species.*

5. TREMELLA, L.

1. TREMELLA mesenterica, Retz, in Vetensk Ac. Handl. 1769, p. 249. Engl. Bot. t. 709.

HAB. Hermite Island, Cape Horn; on a dead trunk of Deciduous Beech, almost covered with former winters' snow, 1,200 feet above the sea, in an exposed place.

The only specimen seen.

6. EXIDIA, Fries.

1. EXIDIA Auricula Judæ, Fries, Ep. p. 590.

HAB. Port Famine; on Beech, C. Darwin, Esq. Hermite Island, Cape Horn; J. D. H.

The specimens collected in the latter locality are small and less tomentose than the more usual state of the species.

7. CRUCIBULUM, Tul.

1. CRUCIBULUM vulgare, Tul. Ann. Sc. Nat. Ser. 3. vol. i. p. 90. Cyathus Crucibulum, Pers. Syn. p. 238. Grev. Scot. Crypt. Fl. t. 34.

HAB. Hermite Island, Cape Horn; on moss near the sea, always solitary.

The specimens differ from the ordinary form, which occurs in the southern as well as in the northern hemisphere, in their solitary habit, more conical peridia, which are of a semi-transparent dirty orange-yellow, and in the more irregular sporangia. In structure I find no difference.

S. LYCOPERDON, Tourn.

1. LYCOPERDON calatum, Bull. Champ. vol. i. p. 156. t. 430.

HAB. Falkland Islands; on a tuft of Bolar.

One specimen only was met with.

It is not possible to speak very positively of a single old specimen and which had been evidently much exposed to the weather. It is, however, certainly neither *L. gemmatum*, nor *L. pyriforme*, and appears to me to be a state of *L. cælatum*. *L. arenarium*, Pers., will be found under the genus *Bulgaria*.

9. LEPTOTHYRIUM, Kze.

1. LEPTOTHYRIUM *decipiens*, Berk.; suborbiculare, atrum, nitidum, sporis tenerrimis irregulari-subfusiformibus quandoque curvatis. (TAB. CLXIII. Fig. III.)

* An authentic specimen, however, received from Mons. Lindblad, since the above was printed, is not more cracked than the Antarctic plant. *Corticium tremellinum* must be considered, therefore, mcrely a highly developed form of *C. viscosum*.

HAB. Falkland Islands; on dead stems of Rostkovia grandiflora.

Puncta irregularia suborbicularia picea nitida in culmos exsiceatos efformans. *Perithecia* valde depressa, demum basi squamæ instar dehiscentia. *Sporæ* irregulares, fusiformes, quandoque curvatæ, tenerrimæ, albæ, pellucidæ; endochromium varie partitum, non autem septatum.

A species which, examined superficially, may be passed over as *Leptostroma junceum*, differing merely in its more shining perithecium. The spores are, however, of a very different form, and many times larger. In that species, as published in 'British Fungi' (No. 197), and by Madame Libert (No. 260), they are extremely minute and obtuse at either extremity; the perithecium also is more closely cellular. In the specimens published by Klotzsch and Fries (in my copy at least), there is no fructification. It resembles also, externally, *Leptostroma vulgare*, but there is as decided a difference as in the former case between the spores.

PLALE CLXIII. Fig. III.—Leptothyrium decipiens, Berk., of the natural size; 2, portion of stem of Rostkovia grandiflora, with base of peridium adhering to it :—magnified; 3, spores :— highly magnified.

10. SPHÆRONEMA, Fries.

1. SPHÆRONEMA sticticum, Berk.; minutissimum, punctiforme, innatum, atrum, nitidum, demum collapsum, sporis minutissimis ellipticis. (TAB. CLXIII. Fig. I.)

HAB. Hermite Island, Cape Horn; on dead leaves of the Deciduous Beech (Fagus Antarctica.)

Minutissimum, punctiforme, atrum, nitidum, demum collapsum, præcipue venis foliorum innatum, unde dispositionem reticulatam exhibit. *Sporæ* minutissimæ, sporophoris brevibus filiformibus affixæ.

Not to be confounded with Spharia punctiformis, Pers., (Fr. Sc. Suec. No. 56), which has true asei, assuming the production published by Fries, which exactly accords with specimens gathered in Northamptonshire, to be the type of the species. Both Desmazière's (No. 984), and Mougeot's, and Nestler's (No. 662) plants appear to me quite different. Unfortunately in neither have I been able to detect fructification. In Mougeot's plant the perithecia arc strongly collapsed, which is by no means the case with that of Fries; and that of Desmazière approaches Sp. maculaformis.

The genus Sphæronema is here considered as comprising such species of the genus Sphæria as have simple spores, never included in asci, such as Sp. acuta, &c.

PLATE CLXIII. Fig. I. — 1, Sphæronema sticticum, Berk., upon leaves of Fagus, of the natural size; 2, portion of leaf and fungus; 3, spores on their sporophores; 4, spores :— all highly magnified.

11. SPORIDESMIUM, Lk.

1. SPORIDESMIUM adscendens, Berk., in Ann. Nat. Hist. vol. iv. p. 292. t. S. f. 1. 1840.

HAB. Falkland Islands; on the underside of Polyporus versicolor, C. Darwin, Esq.

The species is nearly allied to Sp. vagum, Necs, from which it differs merely in having constantly a single globose nucleus in each articulation, presuming that Corda's figure, published in the same year with that in the Annals of Natural History, is the plant of Necs.

12. ÆCIDIUM, Gmel.

1. ÆCIDIUM Magellanicum, Berk.; hypophyllum, totam faciem inferiorem occupans inque petiolos sparsum, rarissime epiphyllum, maculis rubellis, peridiis urceolatis elougatis, sporis pallidis irregulariter orbicularibus. (TAB. CLXIII. Fig. II.)

HAB. Strait of Magalhaens; Port Famine; on Berberis ilicifolia, Capt. King.

Maculæ rubellæ; peridia plus minus elongata, urceolata, sursum leviter constricta, vel omnino eylindraeca, fragilia, totam superfieiem inferiorem investientia, plus minus in petiolos descendentia. *Sporæ* pallidæ (saltem in exemplaribus exsiceatis) irregulariter subglobosæ, angulatæ. Rarissime pauca peridia epiphylla sunt.

Resembling much in external appearance *Ec. sambucinum*, Schwein. It is at once distinguished from *Ecidium Berberidis* by its very different habit.

PLATE CLXIII. Fig. II.-1, leaves of Berberis and *Ecidium Mageltanicum*, of the natural size; 2, portion of leaf and fungus; 3, spores :- highly magnified.

13. UREDO, Pers.

1. UREDO candida, Pers., Syn. p. 223.

HAB. Falkland Islands; on Arabis Macloviana, Capt. Sulivan.

The mycelium is very visible in these specimens. There is no difference in the spores.

14. MORCHELLA, Dill.

1. MORCHELLA scmilibera, Dec., Fl. Fr. vol. ii. p. 212.

HAB. Falkland Islands; on the ground.

I have seen a single imperfect specimen only, which is scarcely more than sufficient to determine the genus. I believe it, however, to be the species of De Candolle, above cited.

15. PEZIZA, Dill.

1. PEZIZA Kerguelensis, Berk.; media, cupula plana adnata coccinea extus setis brevioribus obsita. (TAB. CLXIV. Fig. III.)

HAB. Hermite Island, Cape Horn; on dead branches amongst the snow, alt. 1,000 feet. Christmas Harbour, Kerguelen's Land; May and June; on bare boggy earth near the sea, growing amongst *Conferva*.

 $Cupula \frac{1}{2} - \frac{3}{4}$ unc. lata, plana, adnata, margine tantum ut plurimum libero, coceinea, externe setis brevioribus subflaceidis plus minus contextis primum pallidis deinde saturate rubris vestita, dissepimentis demum absorptis. Asci lineares, obtusi; sporidia late elliptica, glabra, nucleo unico globoso; paraphyses apice clavulatæ.

Allied to *P. scutellata* and *P. umbrosa*, but larger than either. The bristles are short and somewhat flaceid, in which it differs strikingly from the former species, as also in its broader sporidia. From the latter it differs principally in its larger size and less conspicuous hairs. I am not able, in the absence of authentic specimens, to compare the sporidia; but if that species be the same with *P. trechispora*, Berk., and Broome, which is not impossible, the difference is considerable.

PLATE CLXIV. Fig. III.—1, Kerguelen's Land, aud 2, Cape Horn specimens; of the natural size; 3, setæ: --magnified; 4, asci, sporidia and paraphyses; 5, sporidia:--very highly magnified.

2. PEZIZA stercorea, Pers. Obs. vol. ii. p. 89. (TAB. CLXIII. Fig. IV.)

• HAB. Port Louis, Falkland Islands; on cow-dung.

Not distinguishable from European specimens. I caunot detect Ascobolus furfuraceus, which is so generally its eompanion in Europe.

PLATE CLXIII. Fig. IV.—1, bristles from cup; 2, asci and paraphyses, in the broken ascus the inner membrane is visible, projecting below; 3, sporidia :—all highly magnified.

16. BULGARIA, Fries.

1. BULGARIA arenaria, Lév., Ann. Sc. Nat. Ser. 3. vol. v. p. 253. Lycoperdon arenarium, Pers. in Freye. Voy. p. 179. t. 1. f. 2. Gaud. l. c.

HAB. Falkland Islands; "très-commun en Mars et Avril, au sommet des dunes de sable qui bordent le contour de la baie Francaise au Camp de l'Uranie."

This species unfortunately was not found during the visit of the Erebus and Terror. M. Léveillé has had an opportunity of inspecting an original specimen, and finds its slender asci to contain simple sporidia.

17. CYTTARIA, Berk.

1. CYTTARIA *Hookeri*, Berk.; parva, turbinato-obovata, obtuse papillata, pallide fusca, cupulis paucis. (TAB. CLXII. Fig. I.)

HAB. Hermite Island, Cape Horn; on living branches of the Deciduous Beech.

Receptacula communia obovata, e disco oblongo corticali enata, $\frac{3}{4}$ -1 une. alta, $\frac{1}{2}-\frac{3}{4}$ une. crassa, basi attenuata, apiee obtuse papillæformi, pallide fusea, glabra; contextu ut in aliis speciebus gelatinoso-carnoso, e fibris anastomo-santibus; eupulis paucis, primum materic gummosa repletis, demum vacuis; ascis linearibus truncatis, paraphysibus linearibus quandoque furcatis immixtis. *Sporidia* ignota.

The genus *Cyttaria* is peculiar to the Southern hemisphere, and unless Commerson's habitat, to be mentioned presently, should prove correct, to the more temperate latitudes. All the species known at present grow on living beech; *Cyttaria Berteroi* on *Fagus obliqua*, the Fuegian species on *Fagus betuloides*, that of Tasmania on *Fagus Cumninghami*, and *Cyttaria Hookeri* on *Fagus Antarctica*. The species, on which *Cyttaria disciformis*, Lév., grows, has not been ascertained. It is probable that the genus occurs also in New Zealand, where there is a species of beech closely allied to *Fagus Cunninghami*. There exists, indeed, in Monsicur B. Delessert's Herbarium, a species purporting to have been collected in the Isle of Bourbon, by Commerson, but though the locality* is very precisely indicated, it is probable, both on account of the difference of climate and the absence of the genus *Fagus* in that island, that there is some mistake about the specimen.

All the species seem to grow from a distinct disc, which doubtless, as in *Podisoma*, produces a fresh erop every season. The disc bursts through the cuticle, and is formed either entirely of the lower portion of the bark, or of that and the upper stratum of the wood, which are split longitudinally or in the direction of the medullary rays, the fissures being traversed by loose threads of mycelium. Sometimes, also, there are traces of mycelium in portions of bark where no disc has been protruded. The structure of the bark is often much deranged, and sometimes quite disorganized. The base of the receptacles is attenuated, and penetrates generally to the dotted vessels. In *Cyttaria Gramii*, which seems more truly cortical, there appears always to be a fasciele of such vessels in connexion with the base penetrating through the cortical stratum. I do not find this to be the case in *Cyttaria Hookeri*. The structure of the substance of the receptacles is so different in the plant when dry, from that in the same species when

^{*} The label attached to the specimen is literally as follows :---

[&]quot;ELVELA CLATHRUS: sessilis seutellæ instar concava, brunnea subterius murina Commerson. Vel acaulis seutelliformis in concavitate fusca subterius e murino cinerascens. Comm. Entre la Rivière du Rampart et Langevin St. Vincendan, à Bourbon.

Envoié à M. Linné sous le No. 1 (inconnu à M. Linné) An. 1779." (Such appears to be the date, but Commerson died at the Isle of France, in 1773, and the elder Linnæus in 1778.

preserved in spirits, as to be scarcely recognizable. I had, at first, on examining dried specimens of *Cyttaria Gumnii*, fancied that I had made some mistake in the analysis given in my paper in the 19th volume of the Linnean Transactions. The fact is, that when a very thin slice of the dried plant is placed on the field of the microscope, the gelatinous coat of the threads of which it is composed becomes visible; while in the plant preserved in spirits, the jelly seems to form one common mass in which the central tube alone is exhibited, and when the plexus of filaments is drawn out with the point of a lancet, they appear far less curled than they do *in situ*. Perfect sporidia have not at present been observed in any species.*

PLATE CLXII. Fig. 1.—1, Cyttaria Hookeri, Berk., of the natural size, on a living twig of Fagus Antarctica; 2, vertical, and 3, transverse sections of a single plant, of the natural size; 4, asci and paraphyses; 5, curious state of asci; 6, part of the tissue from the darker part of a specimen preserved in alcohol; 7, ditto from lighter part artificially extended; 8, portion of intercellular tissue of Cyttaria Gunnii as seen in a dry specimen; (the same structure is found in dry specimens of Cyttaria Hookeri, and in Cyttaria Darwinii, after it has been preserved in alcohol and dried for the Herbarium); 9, horizontal slice from portion of bark nearest to the wood, in a part of a twig not externally attacked by Cyttaria, to show the mycelium penetrating the cells; 10, slice of fructifying disc, showing two sorts of tissue of the bark, interrupted by a cavity which is traversed by mycelium; 11, slice of bark infested with mycelium; 12, slice from the outer surface of the wood; 13, section through a fructifying disc, showing fissures radiating from wood through the spongy portion of the bark, which is greatly increased in volume, and also a cavity traversed by mycelium parallel to the cuticle. The lower portion of the fungus penetrates in this case to the wood; occasionally, however, it does not penetrate quite so far :—all the above figures, with the exception of the first two, are more or less magnified.

18. ASTERINA, Lév.

1. ASTERINA *pelliculosa*, Berk.; effusa, tenuissima, peritheciis punctiformibus depressis atro-fuscis in mycelio fusco a matrice solubili sparsis. (TAB. CLXIV. Fig. I.)

HAB. Chonos Archipelago; on leaves of an Eugenia; C. Darwin, Esq.

Amphigena, atro-fusca ; maculæ irregulares, varie effusæ punetiformesque, e fibrillis radiantibus intertextis ramis patentissimis formatæ, demum e matrice solubiles ; interstitiis sæpe strato celluloso tenuissimo repletis.

I have not detected fructification; but the species certainly belongs to the genus *Asterina*, which is very properly separated from *Dothidea* by Léveillé. The cells of which the perithecium is composed are elongated, but very irregular, and I find similar cells often filling up the interstices left by the crossing of the radiating threads. Sometimes the mycelium is very obscure and the species then assumes quite a different appearance, the fructifying cells predominating and the patches presenting merely a brown stain studded with darker specks.

PLATE CLXIV. Fig. I.—1, leaves of Eugenia, with Asterina pelliculosa, Berk., of the natural size; 2, part of perithecium seen from the under side; 3, filaments of mycelium :—highly magnified.

2. ASTERINA *stictica*, Berk.; minutissima, omnino punctiformis, mycelio obscuro, peritheciis depressis atro-fuscis margine membranaceo pellucido. (TAB. CLX1V. Fig. IV.)

* The Tasmanian species, of which I have seen dried specimens only, differs from *Cyttaria Darwinii* in the total absence of the granulations at the base of the receptacle. It may be characterized,—

CYTTARIA Gunnii, Berk.; receptaculo subgloboso demum cavo, basi non primum distincte stipitiformi nec scabra, cupulis parvis.

HAB. Tasmania; on Fagus Cunninghamii, R. C. Gunn, Esq.

The specimens are hollow when dry. I cannot say whether such is also the case in Cyttaria Darwinii.

5 м

HAB. Hermite Island, Cape Horn; on leaves of Viola tridentata.

Amphigena, punctiformis. Mycelium valde obscurum, e filamentis paucis brevibus parce ramosis. *Perithecia* depressa, atro-fusea, e cellulis radiantibus elongatis subregularibus formata; margine tenui lacerato membranaceo pellucido.

Neither have I been able to detect fructification in this plant, but the genus is I believe certain.

PLATE CLXIV. Fig. IV.—1, Viola attacked with Fungus of the natural size; 2, leaf of ditto and Fungus; 3, perithecium; 4, portion of edge of ditto :---highly magnified.

3. ASTERINA *Darwinii*, Berk.; epiphylla, maculis parvis orbieularibus e fibrillis radiantibus artieulatis inartieulatisque, peritheciis irregularibus demum depressis centralibus margine laciniato. (TAB. CLXIV. Fig. II.)

HAB. Cape Tres Montes; on Azura lanceolata; C. Darwin, Esq.

Maeulæ epiphyllæ, orbiculares, $\frac{1}{2}$ -1 lin. latæ, e fibrillis radiantibus reticulatisque, partim e margine perithecia, partim e superficie inferiore enatæ, breviter articulatæ, aut omnino simplices. *Perithecia* primum irregularia, subelevata, demum depressa, margine laciniato laciniis denticulatis. *Asci* ut in reliquis speciebus globosi; *sporidia* oblonga, biloculata.

Apparently different from A. Azaræ, Lév., in its perithecia, which are not depressed in the centre, as in that species. Unfortunately I have no opportunity of comparing them. The perithecinm, both here and in Asterina microscopica, splits from the centre in a radiating manner when slightly pressed.

PLATE CLXIV. Fig. II.—1, Asterina Darwinii, Berk., on leaves of Azara lanceolata, of the natural size; 2, perithecia and mycelium; a, cuticle of matrix; b, incipient perithecium; c, curious processes given off from threads of mycelium; 3, fibres of mycelium; 4, portion of border of perithecium; 5, processes on threads of mycelium; 6, asei; 7, sporidia:—all very highly magnified.

19. EUROTIUM, Lk.

1. EUROTIUM herbariorum, Lk., Obs. vol. i. p. 29. f. 44.

HAB. On biseuit on board the 'Erebus', Jan. 3rd, 1841.

The sporangia in the specimens before me, which are very scanty, are almost destitute of floeci, but accompanied by an abundant tawny mycelium, thus confirming the opinion of Fries and Corda, that *Eurotium epixylon* is not really a distinct species. I cannot, however, think with Corda that it has the slightest affinity with *Physarum*. The morphosis has not at present been traced, and till this is done it appears better to let it remain where Fries has placed it, in the neighbourhood of *Mucor*.

. The peridium is lined with a stratum of gelatinous cells, which vanish in a great measure as the plant approaches maturity. The flocci in Kze. and Schm., n. 83, are rough and dark, but I find great variation both of surface and colour.

LV. ALGÆ, L.

1. D'URVILLÆA, Bory.

1. D'URVILLÆA utilis, Bory, in Duperrey Voy. Bot. p. 65. t. 1 et 2. f. 2. Fl. Antarct. Pt. 1. p. 167. Laminaria capæstipes, Montagne in Voy. D'Orbigny, Bot. Crypt. p. 11. t. 2.

HAB. Tierra del Fuego, the Falkland Islands, and Kerguelen's Land; very abundant at half-tide mark and below it; also in the open ocean, between lat. 45° and 55° S., reaching the 65th degree of south latitude in the meridian of New Zealand.

This, the Lessonia, and Macrocystis are the three most remarkable Alga of the Antarctic regions, especially on account of their size; the present exceeding any sea-weed, except the Lessonia and the Ecklonia buccinalis of the Cape of Good Hope, in bulk; while the Macrocystis, to which we shall afterwards allude, is the longest vegetable production known.

The nearest affinity of D'Urvillæa was considered, in the 'London Journal of Botany' (vol. ii. p. 325), to be with Himanthalia of the Northern and Arctie seas, an opinion to which one of us was led by observing how, in habit and locality, these species represented each other in the opposite Polar oceans. Wahlenberg, Bory de St. Vincent and Greville, all regard the curious pezizæform organ of Himanthalia as the frond, and the deciduous strap-shaped laciniæ as receptacles, which view is also maintained in the 'Phycologia Britannica' (t. lxxviii.) Lyngbye (the founder of the species) and Agardh, on the other hand, pronounce the frond to be swollen at the base into a bladdery stipes, furnished with strap-shaped laciniæ, over whose surface the conceptacles are scattered as in D'Urvillæa; and in Xiphophora, a genus (as pointed out by Montagne) nearly allied to the present, and which represents it in a lower latitude of the Southern Ocean. In the 'London Journal of Botany' the true analogy to the bladder of Himanthalia was sought in the trumpet-shaped stipes of Ecklonia buccinalis, but in that plant the growth of stipes and frond proceeds from the earliest stage, pari-passu, whilst the bladder of Himanthalia is fully developed before the straps appear.

We have nowhere seen a good representation of the beautiful cellular tissue of D'Urvillaa utilis, which, in its fresh state, is so regular and large as to resemble perfectly in size and structure one of the two layers of cells found in honey-comb. Most of the specimens brought to Europe are injured by pressure, which can however hardly have caused the total obliteration of structure which M. Bory's plate represents; the most accurate figure we know is given in the beautiful plate accompanying M. Decaisne's 'Essay on the fructification of Algæ'.

The spores of this and the following species are divided into four, and we cannot doubt but that this division is followed by the complete breaking up of the organ into four sporules, whose future germination resembles that described by MM. Decaisne and Thurct in *Fucus servatus* ('Annales des Sc. Nat.' Ser. 3. vol. iii, p. 10. t. 2). The conceptacles contain probably both antheridia and spores, so far as we can judge from drawings taken from the living plant, though at the time these bodies were not recognized as belonging to two differents classes of organs.

The northern limit of D'Urvillæa will probably be found to be the latitude of Valparaiso, or 33° S., on the West coast of South America, and 50° S., on the opposite shores of the same continent. In New Zealand it attains the parallel of 40° , but whether it inhabits any of the shores of Tasmania, or is there represented by the *Fucus* potatorum, is a question we caunot answer. Though carried by the currents along the ocean to the south of the Cape of Good Hope, (for it was collected in that meridiau in the 51st degree, floating in the open ocean,) it does not appear to inhabit or be cast upon the southern extremity of Africa; and in the Indian Ocean, again, its range is not likely to be north of the Islets of Prince Edward's, the Crozet group and Kerguelen's Land. On the other hand, the south latitude it attains is probably regulated by the position of the Paek Ice, to within a few miles of which it was traced by the Antarctic Expedition, on one oceasion, south of New Zealand to the 65th degree, which is probably its "ultima Thule" in any longitude; for it was there the last trace of vegetation. It grows invariably accompanied by the Macroeystis pyrifera.

Bory de St. Vincent states, on the excellent authority of D'Urville, that the poorer classes of West Chili nse this plant for food, and that when made into soup it is very palateable, being sweet and mucilaginous. In Kerguelen's Land its enormous and weighty fronds, sometimes ten feet long, and almost too heavy for a man to lift, form the only shelter for the shells and soft animals, which there find a refuge from the flocks of aquatic birds that cover the shores and follow the receding tide.

2. D'URVILLÆA *Harveyi*, Hook. fil.; radice e fibris crassis demum anastomosantibus constante, stipite perbrevi valido compresso in laminam subsolidam coriaceam apice laciniatam gradatim dilatato. Nobis *in Lond. Journ. Bot.* vol. iv. p. 249. Himanthalia D'Urvillæi, *Bory? in Duperrey Voy. Bot.* p. 135. (TAB. CLXV, CLXVI.)

HAB. Hermite Island, Cape Horn, and the Falkland Islands; abundant.

Radix fibrosus, fibris crassis, inter se intricatis, denum anastomosantibus, discum callumve pertusum 2-4 une. diametro efficientibus. Stipes 3-4-uneialis, $\frac{1}{2}-\frac{3}{4}$ une. diametro, valde compressus, in laminam forma variam gradatim dilatatus. Lamina 4-8-pedalis, supra medium 1-2 ped. lata, plerumque late lanceolata, basi augustata, apieem versus in lacinias plures lineari-elongatas ligulatas abbreviatasve acutas truneatasve fissa, sieeitate atro-fusea v. subpieea, opaea, dura, subfragilis, lineis superficialibus striata, v. subretieudata, e conceptaculis prominulis mamillosa; madore olivaceo-brunnea, coriaeea v. flaccida, plana, lævis, intus solida, 1-4 lin. crassa. Conceptacula sphærica, per totam frondem sparsa, poro ineonspicuo pertusa, filis articulatis sporisque basifixis repleta. Sporæ ut in D.utili, varie quaternatim divisæ, limbo hyalino cinctæ.

Always considerably smaller than the *D. utilis*, of a much thinner texture, and readily distinguishable by its fibrous root. I have never observed the frond of even the largest state of this species to be filled with those elongated transverse cells which distinguish the former.

The structure of the fronds is seen to consist, on a transverse section, of a dense narrow layer of cortical substauce, which gradually becomes more open inwards, and there breaks up into parallel lamellæ projecting towards the centre of the frond. These are less densely packed inwards, and are united at right angles by similar very short plates, together forming a loose cellular tissue, whose walls are thickened at the angles; which, again, at the very centre of the frond, are gradually resolved into a mass of slender, short, waved filaments, free or anastomosing and floating in a gelatine.

The affinity of the Laminaria potatorum is probably with this genus; it is described, by M. Kützing, under the generic name of Sarcophycus (Phycologia, p. 392). I have examined a very small fragment of the plant, and find the spores to be contained in cysts, altogether like those of D' Urvillæa and Xiphophora.

PLATES CLXV, CLXVI.--1, transverse slice of frond; 2, vertical section of ditto; 3, spores and antheridia; 4, spores :--highly magnified.

2. SCYTOTHALIA, Grev.

1. SCYTOTHALIA Jacquinotii, Mont., in Voy. au Pole Sud, Bot. Crypt. p. 86. t. 5.

HAB. Graham's Land; lat. 63° S., floating in the ocean, Dr. Lyall. Deception Island, New South Shetlands; Mr. Webster.

An accurate description of this noble sea-weed is given by its discoverer, Mr. Webster, R.N., in the Appendix to the Narrative of Capt. Foster's Voyage; though nothing was known of the species, botanically, until specimens were received by Dr. Montagne, from the Herbarium of the French South Polar Expedition, collected within a very few unles of the spot where it was again seen by the Antaretie Expedition, and obtained by our indefatigable friend, Dr. Lyall.

The existence of this sea-weed on the Icy shores of an Antaretie land, in the longitude of Cape Horn, is a most singular and anomalous fact; for I believe it to be the only species of the tribe *Cystoseireæ*, which inhabits the colder or Antaretie seas of South America; though many abound in similar temperate latitudes of New Zealand, Lord Auekland's group, New Holland, and Tasmania. We have thus, under the most rigorous skies, the representative of a group, the total absence of whose other species in warmer seas of the same longitude, was supposed to be owing to a low degree of temperature being destructive to its life. The said group of *Cystoseireæ* is not here represented by a species in any way indicative of its habitat being far removed from its eongeners, or of its locality being uncongenial,

We are accustomed to regard the ocean as so ever-active and powerful an agent in facilitating migration, and its uniform temperature is so conducive to the general diffusion of species, that it seems almost wonderful that Algaeshould have limits to their distribution, especially in waters which gird the globe on the same parallel of latitude, and whose unchecked swells and currents literally extend over every degree of longitude. The remarkable increase in temperature of the tropical over the polar seas of the Atlantic may, and probably alone does, check the progress of the *Macrocystis* in its course from Cape Horn to the Equator in that ocean, for, as I shall afterwards show, the same sea-weed can float with the colder currents of the Pacific from the same Cape to Behring's Straits; but no such obstacle prevents the fullest interchange of *Cystoseireæ* between New Zealand and the temperate seas of South America. It, however, is the fact, that whilst this group literally abounds in certain latitudes and longitudes, which are those of New Holland and the West Pacific, they are nearly absent from analogous positions in the longitude of South America.

Throughout all latitudes the two tribes *Fucoideæ* and *Cysloseireæ* form that prevailing marine vegetation to which the name *sea-weed* is commonly applied; and the different genera so far arrange themselves within geographical limits as to present, with such few exceptions as the *Scytothalia Jacquinotii*, a most harmonious assemblage. Thus, in the opposite colder and frigid zones the waters are inhabited by certain genera of *Fucoideæ* which are in a great measure representatives of one another; as, in

the north cool zone
$$\left\{ \begin{array}{c} Fucus \text{ proper, and} \\ Himanthalia, \end{array} \right\}$$
 are represented in analogous $D'Urvillea$, and $Sarcophycus$, Kütz.

None of these genera approach the tropics, for the *Fucoideæ* abound towards the poles, and there attain their greatest bulk, diminishing rapidly towards the Equator, and ceasing some degrees from the Line itself. The representatives of the *Cystoseireæ* in the higher latitudes of the opposite hemisphere, are equally appropriate with those of *Fucoideæ*, for we have in

the north cool zone
$$\left\{ \begin{array}{c} Cystoseira, \text{ and} \\ Halidrys, \end{array} \right\}$$
 represented in the south cool zone, by $\left\{ \begin{array}{c} Blossevillea, \text{ and} \\ Scytothalia; \end{array} \right\}$

whilst the immense genus Sargassum finds its maximum in lower latitudes, and under the Equator itself.

Such are the salient features of the distribution of these tribes, which are not influenced by the minor divisions, chiefly local assemblages of small genera, affecting exclusively certain coasts or bays.

3. LESSONIA, Bory.

1. LESSONIA fuscescens, Bory, in Duperrey Voy. Bot. Crypt. p. 75. t. 2. f. 2. et t. 3. Post. et Ruppr. Illust. Alg. p. 2. t. 3 et p. 4. t. 39. f. 14-18. L. flavicans, D'Urville, in Mém. Soc. Linn. Paris, vol. iv. p. 594. (TAB. CLXVIII., CLXVIII. A., and TAB. CLXXI. D.)

HAB. Hermite Island, Cape Horn, and Falkland Islands; most abundant, always far beyond low-water mark. Christmas Harbour, Kerguelen's Land; rare

The fructification of the species of *Lessonia* occurs, as in *Macrocystis*, upon the surface of the fronds, and there forms large patches. In the present species the sori are situated beyond the middle of the leaf, they are oblong and nearly as broad as the lamina, of which they carry away the upper part when decaying, causing their broad apices to be two-horned. In none of our specimens is the point perfect, all the spores we have seen being situated on the edges of the sorus, which has itself fallen away from the frond. The air-cells are less numerous, and the spores are smaller, shorter, more densely packed than in the following species, and covered

with a very thin cutieular layer of the frond. The presence of this cuticle is owing to the peculiar manner in which the superficial or sporiferous cells of the frond dehisce transversely, allowing the dispersion of the spores (shown in the dissection of *Macrocystis*, given at PLATE CLXIX., CLXX. *Fig.* 2).

This and the following arc truly wonderful *Alyæ*, whether seen in the water or on the beach; for they are arborescent, dichotomously branched trees, with the branches pendulous and again divided into sprays, from which hang linear leaves 1-3 feet long. The trunks usually are about 5-10 feet long, as thick as the human thigh, rather contracted at the very base, and again diminishing upwards. The individual plants are attached in groups or solitary, but gregarions, like the pinc or oak, extending over a considerable surface, so as to form a miniature forest, which is entirely submerged during high-water or even half-tide, but whose topmost branches project above the surface at the ebb. To sail in a boat over these groves on a calm day affords the naturalist a delightful recreation; for he may there witness, in the Antarctic regions, and below the surface of the ocean, as busy a scene as is presented by the coral reefs of the tropics. The leaves of the Lessonia are crowded with Sertularia and Mollusca, or encrusted with Flustræ; on the trunks parasitie Algæ abound, together with Chitons, Patellæ, and other shells; at the bases, and amongst the tangled roots swarm thousands of Crustacea and Radiata, whilst fish of several species dart amongst the leaves and branches. But it is on the sunken rocks of the outer coasts that this genus chiefly prevails, and from thence thousands of these trees are flung ashore by the waves, and with the Macrocyslis, and D'Urvillea, form along the beach continued masses of vegetable rejectamenta, miles in extent, some yards broad, and three feet in depth; the upper edge of this belt of putrefying matter is well in-shore, whilst the outer or seaward edge dips into the water, and receives the accumulating wreck from the sub-marine forests throughout its whole length. Amongst these masses the best Alga of the Falklands are found, though if the weather be mild, the stench, which resembles putrid cabbage, is so strong as to be almost insufferable. The ignorant observer at onee takes the trunks of Lessonia thus washed up for pieces of drift-wood, and on one occasion, no persuasion could prevent the captain of a brig from employing his hoat and boat's crew, during two bitterly cold days, in collecting this incombustible weed for fuel !

The trunks, which contract to one-fourth of their original dimensions when dry, and become deeply furrowed, are perfectly smooth and cartilaginous when fresh. On being cut across, the curious appearance of concentric elliptical rings, in many respects similar to, though very different from, those of an Exogenous trunk, is very evident. These rings surround a lance-shaped pale line, which occupies the broad axis of the compressed stem, without reaching across it, and appears to afford some rude indication of the age of the plant, though of this we could by no means satisfy ourselves. It is singular that this, the most arborescent of the *Alga*, and the beautiful *Usnea melaxantha*, the most tree-like in form of the *Lichens*, are nearly the only plants of the Orders to which they respectively belong, conspicuously presenting even a semblance, if it be no more, to a growth that indicates an increase by periodical accessions to the circumference.

The substance of the trunk of the *Lessonic* is very usefully employed by the Gauchoes, for knife-handles*; the haft of the instrument is plunged into a radely-shaped piece of this weed, which contracts into a substance harder than horn. The range of the present species is from the Falkland Islands to Cape Horn, and thence north along the coasts of South America probably to Valparaiso.

PLATE CLXVII.—CLXVIII. \mathcal{A} .—apex of a branch and fronds, of the natural size; \mathcal{A} . 1, portion of stem showing layers of cellular tissue and air-cell :—*magnified*.

2. LESSONIA nigrescens, Bory, in Duperrey Voy. Bot. Crypt. p. 80. t. 5. Post. et Ruppr. Illust. Alg. pp. 2 et 4. t. 4 et 39. f. 11 et 13. (TAB. CLXVII.—CLXVIII. C.)

HAB. Hermite Island, Cape Horn, and the Falkland Islands; very abundant, with the former species. A species very similar to the preceding in general appearance, but of a different consistence and colour.

^{*} The stipes of *Laminaria digitata* is used by the Orkneymen for similar purposes, as is noticed by our excellent friend Dr. Neill, in his interesting account of the Orkney Islands.

Fructification forming a large oblong or linear sorus between the base and middle of the frond, of a rich red-brown colour when held between the eye and light, imbedded in the thickened substance of the frond, which decays with it. On a transverse section the soriferous lamina is seen to be hollow in the centre; or rather the sorus is formed of two parallel plates, each covered externally with densely aggregated spores, which occupy what are the superficial cells of other parts of the frond. Below the superficial series of cells, and especially in fertile specimens beneath the spores, are several, 4-6 or many more, air-cavities, reposing on, and separated from each other by a loose cellular tissue, which is hexagonal, transparent, the cells becoming transversely elongated and finally towards the centre of the frond breaking up into a layer of matted filaments, which surrounds the cavity, a structure resembling very closely that of *Fucus confluens* as given by Turner. When dry, the surface of the plant is covered with white efflorescence, similar to that of Laminaria saccharina, it has been analysed by my friend Mr. Stenhouse of Glasgow, who finds it to contain excellent Manna, and who further informs me that this and the other larger Antarctie Algæ are peculiarly rich in Iodine.

The Lessonia quercifolia of Bory, is described and figured as having the frond eovered with cavities containing spores, whence it would appear to belong to Fucoideæ, and to be more allied to D'Urvillea than to this genus.

Lessonia ciliata of Postel and Ruppreeht, is certainly only the young state of Macrocystis purifera.

PLATE CLXVIII.-CLXVIII.-C. transverse section of frond in fructification :- highly magnified.

3. LESSONIA ovata, Hook. fil. et Harv.; stipite brevi vage dichotome ramoso, ramis brevibus divaricatis, frondis laciniis breviter petiolatis, petiolo in laminam ovatam lineari-ovatamve olivaceo-fuscescentem submembranaccam dilatato. (TAB. CLXVII.—CLXVIII. B; et TAB. CLXXI. C.)

HAB. Hermite Island, Cape Horn, and the Falkland Islands; very abundant.

Radix e fibris perplurimis crassis intricatis massam 1-2 ped. latam efficientibus. Stipites e radice plurimi (ut in Macrocysti) 4-6 unc. longi, torti v. flexuosi, erassitie pollicis humanæ, dichotome fissi, demum solitarii, incrassati, subarborescentes. Laminæ pedales, colore et substantia L. fuscescentis, juniores basi obscure sinuato-dentatæ; adultæ integerrimæ.

Certainly very near L. fuscescens; but as far as could be judged on examining the plant, both on the shores it inhabits and in the herbarium, it has good claims to be considered a distinct species, especially in the many short stipites, short branches and broad leaves. Never having seen the fruit, however, it may prove the young of L. fuscescens, for we can well suppose only one out of the many stems of that plant to attain any great dimensions, and the lamina of the young state to be broader than that of the adult.

The ramification of all the species of Lessonia is dichotomous; each plant in a young state consists of a few rooting and clasping fibres, giving off a single stem (or petiole) and frond. This frond splits at the base, and as the growth proceeds, the fissure extends vertically upwards, till the original frond is bisected; each of the two parts is now a complete frond, altogether similar to the primary one, and provided with a petiole of its own : these again divide, and the process is repeated. Hence the rapid growth of this genus, and hence the origin of the flattened form of ramuli and elliptic core which is placed in the long axis of these ramuli and across the axis of the terete stem. It was not observed whether any relation existed between the number of branches on the whole frond and of concentric rings in the trunk. The latter are probably the indices of the number of times that a subdivision of the laminæ has occurred, supposing that all split at about the same epoch, rather than a register of the years the vegetable has existed; as the following account of the anatomy of this species will show.

A branched portion of the plant, terminated by four laminæ, necessarily presents subdivisions of three periods of growth: 1st, the petioles of the four laminæ; 2nd, the two ramuli from which the four are given off; and 3rd, the one branch which gives off the two latter: these were successively examined.

1. The base of the lamina or petiole is exceedingly compressed, and composed of a mass of cellular tissue of

459

different textures, all, however, very gelatinous, and modifications of the three layers forming the leaf, there are 1st, the superficial tissue (or cortex) consisting of small cells, closely packed and full of chromule, gradually opening out into, 2nd, an intermediate tissue of much larger cells more loosely placed, with little or no contained chromule, separated by much gelatine; and 3rd, an elliptical core placed in the long axis of the petiole, composed of still smaller cells, separated by broader masses of gelatine, which latter is permeated by canals, full, as are the small cells, of chromule.

2. Each ramulus, from which proceed the two periodes, whose structure we have just described, presents no very important difference from them; the core no longer stretches across it, however, but the whole petiole within the superficial portion is augmented by a newly developed though indistinct zone of cellular tissue, thus deposited between the superficial (or cortical) and intermediate tissue. At this period the cortex is somewhat broader, and the intermediate tissue has become, through the absorption of the gelatine, much more conspicuous; the cells being larger and the spaces between them narrower; little or no change is perceptible in the core itself.

3. The branch is very materially different from either of the above, for what was hitherto the petiole is now enclosed (all but its cortex) in a very broad zone of cellular tissue, whose cells are large and thin towards the old tissue, elongated and of a different shape, so as to show the line of separation between the two periods of growth (see B 1, of the plate Lessoniæ).

From this time forward the normal mode of growth followed by the stem exhibits an additional layer or zone of cellular tissue for every subdivison of the frond, (shown at \mathcal{A} 1, where six are interposed between the cortex and core). It is not probable, however, that this numerical relation can be always evident, or that the number of subdivisions of the frond will indicate the rings of growth in a large stem. This uncertainty arises from the branches being frequently broken off; added to which, the growth of the sea-weed is very rapid, and there being no period of rest, irregular zones may be expected, or their absence from those branches of the plant whose leaves are injured.

In their anatomy the stems of *L. fuscescens* and *L. nigrescens* do not differ much from that of this species, except that the air-cells are copious in the stems of the former, and much rarer in the latter; in which also the cortical substance is much broader.

In the elegant *Lessonia Sinclairi*, Harv. MSS., from California, the stipes (which bears but a solitary lincar frond) is terete, and in the specimen we examined, contains a central core, reaching half-way across the diameter. There are apparently two rings of tissue beneath the cortex, separated by a zone of very large cells (air-cells?); whence it is difficult to account for the stem being terete, for the frond is plane, and the core three times longer than broad. Nor is it easy to explain the origin of the two zones surrounding the core; if they really be successively deposited, it is possible that the frond is two years old; if not, that the large cells are air-cells, and do not indicate a line of separation between two successive deposits.

I have stated the growth of the Lessonia to be very rapid; this is proved by the zones of a five-ringed stem being progressively broader towards the circumference. The probability, too, of one being added for every time the laminæ divide, and the fact that the process of subdivision is continued in geometrical progression, all favour the opinion that these Algæ attain their enormous bulk in a very few months. The vast masses washed up on the outer eastern shores of the East Falkland Island, and the rapidity with which they decay, are additional proofs of a singularly rapid development.

The analogy between the mode of growth exhibited by this genus and an Exogenous tree, is, though incomplete, very obvious; both increase by layers deposited outside one another, within a cortical substance, and both contain an axis of tissue different from that forming the greater part of the trunk : here, however, there are no traces of medullary rays. We conclude this subject with the observation, that the periodical increment of the trunk being dependent on, or coincident with, the formation of the laminæ, these appear to perform the office of the leaves in the higher order of plants; and that the *Lessonia* is also in this respect analogous to an Exogenous plant,

deprived of its woody tissue, for it is a stem composed of layer upon layer of cellular tissue, deposited round an axis, which, like the pith, when once formed, is afterwards but slightly modified.

PLATE CLXVII.—CLXVIII. B. apex of branch and frond of the natural size; B 1, transverse section of young stem :—magnified.

4. MACROCYSTIS, Ag.

1. MACROCYSTIS pyrifera, Agardh, Sp. Alg. vol. i. p. 47. Nov. Act. Nat. Cur. vol. xix. p. 297. t. 26 f. 2. Post. et Ruppr. Illust. Alg. p. 9. t. 6; et p. 4. t. 39. f. 22, 23. Fl. Antarct. Pt. 1. p. 178. M. communis, Bory, in Dict. Class. v. x. p. 8. M. planicaulis, Agardh in Nov. Act. Nat. Cur. l. c. Lessonia eiliata, Post. et Ruppr. l. c. (young state).

Var. B. integrifrons; foliis fere integerrimis planis rugoso-plicatisve. M. integrifrons, Bory, l. c. t. 6.

Var. γ . angustifrons; vesiculorum parietibus tenuibus, foliis ut in *M. pyrifera*. M. angustifrons, *Bory*, *l. c.* t. S. *Agardh*, *l. c.* t. 26. f. 4 and 5. *Post. et Ruppr.* t. 5.

Var. 8. zosteræfolia; foliis anguste lineari-elongatis planis. M. zosteræfolia, Bory, Sc.

Var. e. luxurians; foliis 3-S-pedalibus S unc. latis basi cordatis membranaccis plicatis margine longe eiliato-dentatis, vesiculis late obovatis parietibus tenuibus, caule gracili. (TAB. CLXIX.-CLXX.)

Var. ¿. membranacea; foliis ut in M. pyrifera sed tenuissime membranaceis planis, vesieulis parvis elliptico-ovatis utrinque subacutis.

Var. n. Humboldtii; foliis lineari-elongatis planiusculis, vesiculis globosis tenuibus. M. Humboldtii, auct. M. pomifera, Bory.

HAB. Throughout the Antarctic scas, between the parallels of 40° and 64°, both attached, and floating over the whole ocean.

After a very attentive examination of many hundreds of specimens, we have arrived at the conclusion that all the described species of this genus which have come under our notice may safely be referred to *Macrocystis pyrifera*. Nor can these variations excite surprize, when it is considered that this gigantic weed is subject to every vicissitude of elimate, of temperature, and exposure; that it literally ranges from the Antarctic to the Arctie eirele, through 120 degrees of latitude; that it lives and flourishes, whether floating or attached, growing in bays, harbours, or the open sea when most distant from laud; and, lastly, that it equally adapts itself to the ealmest or most tempestuous situations, to waters of uniform depths or those which rise and sink with the tide, to dead water or to strong currents. One thing alone it requires, and that is, a mean depth of six or more fathoms; for, like the *Laminariæ* of our own shores, it, and others of the same tribe in the sonth, invariably form the outer belt of marine vegetation.

A few remarks upon the above varieties may be interesting; showing how much their characters depend upon natural eauses, and how much more upon mutilations of the specimens, or changes during the operation of drying.

Variety β . integrifrons. This we have received from various parts of the west coast of South America; its characters rest almost entirely on the want of ciliation at the margin of the frond, which is much dependent upon the portion of the plant from which the specimen is taken, the lower leaves being always nearly entire; also on the state of the waters, those plants which grow in quiet bays having very much developed ciliæ, whilst those from the main ocean or stormy coasts are generally more entire.

Variety δ . zosteræfolia, is a plane and narrow-leaved state of M. pyrifera; we have traced all the changes in one specimen of M. pyrifera, from very rugose to perfectly plane. Young specimens and terminal leaves are generally

plane, and it often depends on the smoothness of the water how long they may remain so. This variety is abundant everywhere in the Antaretic seas.

Variety γ . angustifrons. The character, drawn from the tenuity of the vesicles, is utterly unsatisfactory, being attributable to the drying of the specimen, and the locality of the live plant. Besides the Antarctic habitats of this variety, it has been found in Chili, New Zealand, and the Indian Ocean.

Varieties ϵ . luxurians, and ζ . membranacea. If any form of this genus deserves specific distinction it is surely the noble one we have designated ϵ . luxurians; and yet permanent characters, distinguishing it from pyrifera, were vainly sought in plants gathered on the shores of Berkeley Sound. Both there and at Cape Horn these two states inhabited deep and still waters, where, as might be expected, the *Macroeystis* would acquire its greatest development, where its substance would be most membranous, its stems most slender, and the vesicles broad with thin walls, and the base of the frond broadest. We have seen no specimens of these varieties except what were brought home by the Antarctic Expedition.

Variety η . Humboldlii, at first sight appears different, and the specimens found on the outer shores of the Falklands we once thought might belong to a distinct species. The rounded form of the vesicles, however, which affords the main character, is not constant on specimens collected in the Coral Islands by Captain Beechey. It has been gathered at various places along the west coast of South America, from Cape Horn to the Equator, and far westward in the Pacific amongst the Coral Islands.

With regard to other states, which we have not seen, the most remarkable is the *M. Orbignyana* of Montagne (Sert. Patagon. p. 12. t. 1.), which has the vesicles remarkably lengthened and the leaf attenuated at the base above the vesicle into a distinct petiole. The *M. latifolia*, Bory, is intermediate between our ϵ . *luxurians* and *pyrifera*. *M. tenuifolia*, Post. and Ruppr., is apparently between *M. pyrifera* and *M. zosteræfolia*. The character of *M. planicanlis* is founded on the compression of the stem, produced by drying, and we have therefore quoted it as a synonym.

In thus bringing together under one, the ten species which have been described by five authors, of whom hardly one has ever seen even the genus in a living state, we are only taking advantage of opportunities which a long residence in the Southern Hemisphere has afforded. Without studying these plants on the coasts they inhabit, it is impossible to judge of the influence of local causes on their plastic forms. We venture to say that few botanists in Europe have seen even tolerable specimens from one single plant of this Alga, such, we mean, as give a fair idea of the differences between the leaves and bladders, along, perhaps, 300 feet of stem, with the submerged fructifying fronds from the root. Out of some thirty specimens brought home by ten different collectors and preserved in the Hookerian Herbarium previous to our visit to the seas which M. pyrifera inhabits, not one conveys any notion of the variations which even a solitary individual can assume.

The fructification of this plant appears to be produced only on the young newly-formed submerged leaves, where it forms large irregular brown patches or sori, causing the frond to separate into two laminæ, as in Lessonia. The spores are fusiform, first divided into four, each afterwards breaking up into as many sporidia. Under a high power the surface of the fertile frond is seen generally to be covered with anastomosing raised lines of a dark colour, on which the spores are placed; the spaces between are pale and transparent. We have not noticed spores, like what are figured by Agardh (l. c. t. 28. f. 11), but plenty of the kind he represents at f. 10^b of the same plate, though not contained in sporangia. These, magnified as highly as his f. 10^b, are evidently divided, as in $D^{i}Urvillæa$. The granules also, which occur abundantly with the spores, are surrounded by a hyaline border, and divided into two to four sporidia; we suppose them to be merely small spores.

It is seldom that the history of an *Alga* is likely to afford interest or annusement to the general reader, nnless it be a positively valuable plant in an economic point of view. Like the Sargasso-weed of the Tropics, however, the *Macrocystis* is so conspicuous, and from its wandering habits, often occurs so unexpectedly, that the attention of our carliest voyagers has been directed to it, and we are consequently led back by our enquiries into its first
in the stormy seas of the south. "Nihil vilior Alga", is a saying more trite than true, and one which a seaman can never use; for these weeds often prove his unerring guide towards land, as they surely are to the direction of the currents; or become of more importance still in the case of the present plant; for it is, where growing, not only the infallible sign of sunken rocks, but every rock that can prove dangerous to a ship is conspicuously buoyed by its slender stem and green fronds, and we may safely affirm that without its presence many channels would be impractieable, and numerous harbours in the south closed to our adventurous mariners.

The first notice of the *Macrocystis*, with which we are acquainted, is of so early a date as the middle of the 16th century, and occurs in a copy of sailing directions for mariners, with the title "A Rutticr from the River of Plate to the Streight of Magelana", and forms part of "A special note concerning the currents of the sea between the Cape of Buena Esperanza and the coast of Brazilia, given by a French pilot before Sir John Yorke, Knt., before Sebastian Cabote, which pilot had frequented the shores of Brazilia eighteen voyages." (Hakluyt, ed. 2. vol. iv. p. 219). In describing the above-mentioned route, after passing Cape Sta. Martha, the trusty pilot's direction to the mariner is to "goe S.W. by W. until he be in 40 degrees, where he shall find great store of weedes which come from the coast"; and again, in pursuing the voyage after entering the Straits, "if you see beds of weede, take heed of them and keep off from them." Now, both the position assigned to the great masses of floating weed and the value of those which are attached in denoting hidden dangers, are conclusive as applying to the Macrocystis. These directions bear no date; but the discovery of the Strait of Magalhaens was in 1520, and the death of Sebastian Cabote took place in 1556, so that we have sufficient proof that this plant attracted the attention of the earliest Antarctic voyagers in the longitudes of Cape Horn; though it may have been noticed previously on the southern extreme of Africa or the China seas. Nor can we wonder that the attention of our forefathers should have been so early called to it, when even now it is of the first importance that the look-out man should use his utmost vigilance to detect, and promptitude to report, this weed, on approaching any of the straits and bays of the shores of Tierra del Fuego and similar latitudes. In the latest voyages that have been published, those of Capts. Foster, King, and Fitz-Roy, we find a constant watch for the "kelp" to have been kept, and caution used to avoid the "moored" pieces, together with instructions how to distinguish them from those which are floating.

The earliest scientific notice which we find of it, consists of a rude figure and description in Bauhin's "Historia Plantarum," published in 1651, where it is designated "Fneus marinus crinitus." In the year 1764, the Freuch Navigator, Bougainville, fell in with the Macrocystis, Lat. 42°S., Long. 57°W., and a long description appeared in 1771, by Don Pernetty, the historiographer of that voyage, together with a sketch, when it was published as a native also of the Falkland Islands. (Pernetty Voy., vol. ii. p. 67. t. ix. f. 5.) About the same period (1771) specimens were collected by Emmanuel Kœnig (of Bale) on his voyage to India, and transmitted to Linnæus, with this observation, "Habitat in oceano Æthiopico e profundissimo mari sæpe enatans insulasque quasi formans," (MSS.); it was then published in 1771, as Fucus pyriferus (Linn. Mantiss. p. 311,) with the additional remark of "maximus forte omnium Fucorum." We are not sure of the precise habitat of Konig's specimens; but by "oceanus Æthiopicus" he probably alluded to the seas of the Cape of Good Hope, which he doubled on his way to India, and where this weed abounds.

So remarkable a plant was not likely to escape the notice of Cook, and especially of the illustrious companion of that navigator's first voyage, and we accordingly find in his narrative repeated allusions to it. It engaged the attention of Banks when entering the Straits of Le Maire in 1769, and frequently afterwards in the cooler latitudes of the southern ocean. To him we owe the first account of its gigantic dimensions. Captain Cook says, on the authority of Banks and Solander, who called it Fucus giganteus, that the stems attain a length of 120 feet. That these dimensions are considerably under the mark there is little doubt; though the report that specimens have been measured upwards of 1000 feet is perhaps as much of an exaggeration. Still it must be remembered that, provided the water be smooth and of sufficient extent, there are no impediments to the almost indefinite

elongation of the upper part of a plant which never branches, and whose growth is independent of all below it, even of the root. Specimens measuring between 100 and 200 feet are common in the open ocean, and these are always broken off at the lower end, either from the division of the frond by sea-amimals, through whose agency the plant increases and the floating island it forms dilates, or from the impossibility of securing the whole mass from the motion of the vessel or the swell of the sea, in latitudes where no boat can be lowered. Again, D'Urville, upon whose observations in natural history the utmost reliance may be placed, states it to grow in eight, ten and even fifteen brasses of water, from which depth it ascends obliquely and floats along the surface nearly as far : this gives a length of 200 feet. In the Falkland Islands, Cape Horn and Kerguelen's Land, where all the harbours are so belted with its masses that a boat can hardly be forced through, it generally rises from eight to twelve fathom water, and the fronds extend upwards of one hundred feet upon the surface. We seldom, however, had opportunities of measuring the largest specimens, though washed up entire on the shore; for on the outer coasts of the Falkland Islands, where the beach is lined for miles with entangled eables of Macrocystis, much thicker than the human body, and twined of innumerable strands of stems coiled together by the rolling action of the surf, no one succeeded in unravelling from the mass any one piece upwards of seventy or eighty feet long; as well might we attempt to ascertain the length of hemp fibre by unlaying a cable. In Kerguelen's Land, the length of some pieces, which grew in the middle of Christmas Harbour, was estimated at more than three hundred feet; but by far the largest seen during the Antarctic Expedition, were amongst the first of any extraordinary length which the ships encountered, and they were not particularly noticed, from the belief that the report of upwards of 1000 feet length was true; or, at any rate, that better opportunities of testing its truth would arise in the course of a three years' voyage, than the first week of our explorations could afford. These occurred in a strait between two of the Crozet Islands, where, very far from either shore, in what is believed to be forty fathoms water, somewhat isolated stems of Macrocystis rose at an angle of 45° from the bottom, and streamed along the surface for a distance certainly equal to several times the length of the 'Erebus';-data, which if correct, (and we believe them so) give the total length of the stems as about 700 feet.

That isolated patches of weed should rise through such a volume of water is not incompatible with the statements we have elsewhere made, that eight or ten fathoms is the utmost depth at which, judging by our experience, submerged sea-weed vegetates in the Southern temperate and Antaretic Ocean. These exceptional eases are probably due to the parent plant having attained such a size in its birth-place near shore, as to weigh its stony moorings and deposit itself in deeper water, where an increase of the roots would unite the original base to other rocks, and thus gain a footing that defies the power of the elements.

We have stated that the elongation of the *Macrocystis* may be indefinite; but this is only true partially and in the case of detached patches: for the stem of the attached plant does not gain bulk or tenaeity, after a certain period; whilst the growing dimensions of the floating portion are increasing the difference between the specific gravity of the vegetable and the element it inhabits, and consequently augmenting the strain upon the slender stem by which it is attached. At some period or other, the resistance is overcome and the floating part detached from the submerged : though at what epoch this may take place, or whether it be coincident with other phases in the life of the plant, is beyond our conjecture.

The fact that fructification is produced only on the submerged young bladderless and small frond, within a few inches of the very root, is highly remarkable. What then is the function of the floating mass of the plant? to one of whose thousand leaves, each four to six feet long, the fructifying part bears an inconceivably small proportion. Were this a phænogamie plant, we should recognize, in such foliaceous expansions, organs which fulfil a respiratory and digestive office and are subservient and necessary to the development of the more important parts of the vegetable; but in this case such a mutual dependency is not so easily traced. As in *Lessonia* the multiplication of the leaves is intimately connected with the development in diameter of the stem, so in *Macrocystis* the development of fructifying fronds may take place only at the root of the barren ones, on whose previous existence they may be dependent for their origin. These are, however, questions which propose themselves to us in the closet only,

when the prospect of solving them is gone by; and when they but add to the thousand regrets over lost opportunities, the remembrance of which weighs so heavily on the mind of every naturalist, that the brightest prospects of discovery in the fair future can never obliterate them.

So many interesting points are connected with the *Macrocystis*, that a book might be instructively filled with its history, anatomy, physiology and distribution; whilst its economy, its relation to other vegetables and to the myriads of living creatures which depend on it for food, attachment, shelter and means of transport, constitute so extensive a field of research that the mind of a philosopher might shrink from the task of describing them. We conclude with an outline of its dispersion over the surface of the globe, which is wider than that of any of the large *Algæ*.

As already mentioned, the Macrocystis girds the globe in the Southern temperate zone, but not in the Tropics or Northern Hemisphere, and this is a most curious trait in its history. We may first, however, trace the southern edge of the belt which it forms, and we are the better enabled to do so, because the limits of its existence, as a floating plant, were observed in six different longitudes in the passage of the Antarctie Expedition as often between the Southern Sea and the Southern Lee, within which there is no vegetation. The southern boundary of the "Macrocystis sea" is very much determined by the position of the ice, and the northern by the currents and temperature of the water. Thus, in the longitude of New Zealand, where open sea extends to the 65th degree, this plant is found as far as 64°, the specimens having probably been drifted originally from Kerguelen's Land or the Crozets, which are the great nurseries for it in the Eastern Hemisphere, and from whence all those drifting islets have been wafted which occur between their longitude and Cape Horn. In the longitude of Cape Horn, 58° or 60° is the highest parallel it attains, for it has not been found amongst the South Shetlands; further east, in the South Atlantic, its parallel is probably still lower; till in the meridian of the Cape of Good Hope it is 40 degrees removed from the Pole, being scen no further south than 50° 30'. There the Atlantic Ocean specimens are derived from the southern extreme of America and the neighbouring islands. Its northern range on the other hand is dependent, 1st, on the temperature of the ocean ;- for it neither enters the Tropics of the Atlantic, nor passes up the shores of Africa or into the Indian Ocean ; whilst it does inhabit the whole surface of the Pacific Ocean and the west coast of both Americas : 2ndly, on the currents, for when north of the influence of the uniform westerly movement of the waters in the Antarctic Ocean, it is deflected with their courses and carried, while temperature allows, to whatever seas receive those waters. Thus, the South Polar current divides at Cape Horn, one portion following the west coast of South America to Cape Blanco and the Galapagos Islands under the Equator, carrying the Macrocystis with it, which then enters the cold waters which flow from the Arctie Islands of the Pacific, and over whose entire surface it is spread, reaching Kamtschatka, New California, and the Alentian Islands : so that in the longitude of Western America the Macrocystis ranges from the Arctie to the Antarctic circle. The eastern branch of the Cape Horn current passes between the Falkland Islands and Fuegia, conveying vast masses of this sea-weed 200 miles north of the Falklands, as low as the 44th degree, and some even reaching the Plate river in 35°, its northern limits in the Western Atlantic. Further west in the Antarctic ocean its distribution is less known; but since it does not occur far north of the Cape of Good Hope in that meridian, we may conclude that it ceases about the 34th degree. With regard to the South African habitat, it is difficult to account for so vast a quantity as the Agulhas Bank exhibits, for these waters, 130 miles in breadth, flowing with a rapid stream from the N.E. or Indian Occan, literally swarm with Macrocystis, which possibly is taken up from the northern edge of the westerly Polar current (which flows along the parallel of 45° S.) by the Indian (or N. E.) current in question.

Its northern limit in the Indian Ocean is not ascertained, but it lies probably south of a line drawn northeast from the Cape of Good Hope to Australia, upon whose western shores the plant is found, as also in New Zealand, and on the coast of China to the north, to which sea it perhaps migrates from the North Pacific Ocean, Kamtschatka, &c.

PLATE CLXIX., CLXX. Frond of M. pyrifera, var. luxurians, of the natural size; 1, thin slice of fructifying

frond; 2, portion of ditto showing the two modes of escape of the spores, either free or contained in the original cells:--very highly *magnified*. (The quaternary division of the spores was not seen in this dissection.)

5. LAMINARIA, Lamourx.

1. LAMINARIA fascia, Ag., Syst. p. 273. Harv. Phyc. Brit. t. 45.

HAB. Hermite Island, Cape Horn, and the Falkland Islands; not common.

These specimens do not appear to differ from British ones in any particular. The *L. fascia* is a northern, but not Arctic species, found along the shores of England, Ireland, and the German Ocean; and under the name of *L. debilis*, it also inhabits the Mediterranean Sea.

6. CAPEA, Montagn.

1. CAPEA biruncinata, Montagn. in Flor. Canar. Crypt. p. 140. t. 7. Laminaria biruncinata, Bory in Duperrey Voy. Bot. Crypt. p. 101. t. 10. L. radiata, β . exasperata, Turner Hist. Fue. vol. ii. p. 16.

HAB. Hermite Island, Cape Horn; J. E. Davis, Esq.

Our solitary specimen is barren, and does not appear different from the plant of the Canary Islands, New Holland, and New Zealand.

We quite agree with Dr. Montagne as to the propriety of separating this genus from Laminaria and Ecklonia, which latter is its Cape of Good Hope representative, and is reported to be a native of the Falklands, probably erroneously; as is the station assigned to the Macrocyslis of the Canary Islands. On the other hand, there is no reason why the Macrocyslis should not accompany the Capea, whose principal parallel is certainly in the Southern Hemisphere.

The fructification of *Ecklonia* is scarcely known; we have seen what appear to be young sori in the form of opaque thickened spots on the frond. A transverse section shows the cortical layer to be thickened and formed of parallel tubes full of granules, analogous to what such fructification as Montagne's beautiful analysis of *Capea* represents would be in an immature state, but we are extremely doubtful of our analysis.

7. DESMARESTIA, Lamourx.

1. DESMARESTIA media, Grev., Synops. p. 40. Sporochnus medius, Agardh, Ic. Alg. p. 259. t. 16. D. anceps, Montagne in Foy. au Pole Sud, Bot. Crypt. p. 51?

HAB. Cockburn Island, (lat. 64° 12' S., long. 57° W.); floating in the sea, abundant.

One of the most Antarctic of plants and probably common in many latitudes; for, under other names, it has been noticed as a native of Peru, of various parts of the Pacific Ocean, and even north to the Arctic circle. It is singular that a plant of this small genus, and from which the present is perhaps not specifically distinct, the *D. aculeata*, should be among the highest Arctic *Algæ*, inhabiting Spitzbergen in $S0^{\circ}$ N. lat. Montagne's *D. anceps*, confessedly described from imperfect specimens, is very probably this plant, having been gathered in nearly the same locality : or else it is the *D. ligulata*, a Cape Horn species.

2. DESMARESTIA viridis, Lamourx.-Flor. Antarct. Pt. 1. p. 178.

Var. B. distans; ramis remotioribus.

HAB. Hermite Island, Cape Horn, Falklaud Islands, and Kerguelen's Land; very abundant. Var. β , Kerguelen's Land.

The range of *D. viridis* in the Northern Hemisphere is not extensive, and almost confined to the shores of England and of the German Ocean. It is, however, found at Unalaschka, according to Postel and Rupprecht.

3. DESMARESTIA ligulata, Lamourx. Grev. Alg. Brit. p. 37. t. 5. Turner Hist. Fuc. t. 99.

HAB. Hermite Island, Cape Horn; rare.

Our specimens in no way differ from European ones. It is probably a common Antarctic species, for we have received it from the east coast of Patagonia.

4. DESMARESTIA herbacea, Lamourx. Montagne in Voy. au Polc Sud, Bot. Crypt. p. 50.

HAB. Port Famine, Strait of Magalhaens; M. Hombron.

We have seen no Antarctic specimens of this plant, which is also a native of the Cape of Good Hope, Concepcion on the west coast of South America, and of North-west America.

5. DESMARESTIA chordalis, Hook. fil. et Harv.; fronde eoriaceo-cartilaginea compressa anguste lineari tri-quadripinnata, pinnis pinnulisque longissimis oppositis distantibus apice longe nudis, pinnulis sæpe alternis elongatis inermibus chordiformibus. Nobis *in Lond. Journ. Bot.* vol. iv. p. 249.

HAB. Christmas Harbour, Kergueleu's Land; very abundant, in two to five fathom water.

Alga socialis, rupicola, cæspitem gramineam submarinam late extensam efficiens. Frondes e radice anguste scutata valde elongatæ, 3-5-pedales. Stipes $1-1\frac{1}{2}$ lin. diametro. Pinnæ paulo angustiores, pinnulis $\frac{1}{2}$ lin. latis.

A very noble species, recognizable at once by the long whip-like naked apices of its pinnæ. In the great abundance of this *Alga* consists one of the peculiarities in the submarine vegetation of Kerguelen's Land.

6. DESMARESTIA *Rossii*, Hook. fil. et Harv.; fronde eoriaeeo-cartilaginea compressa lineari bi-tripinnata circumscriptione anguste lanceolata, pinnis pinnulisque omnibus oppositis basi apiceque attenuatis aeutis erectis v. ultimis appressis margine integerrimis. Nobis *in Lond. Journ. Bot.* vol. iv. p. 249. (TAB. CLXXIII., CLXXIII.)

HAB. Staten Land, A. Menzies, Esq. Hermite Island, Cape Horn, and the Falkland Islands; very abundant.

Frondes 4-8 ped. longæ, anguste lanceolatæ, nunc apices versus dilatatæ, inferne bipinnatæ, supra medium tripinnatæ, valde coriaceæ. Caules stricti, 2-3 lin. lati. Pinnæ 1 lin. latæ pinnnulæque oppositæ, subercetæ, axillis acutis, basi apiceque attenuatæ.

Even a nobler species than the last, from its great size and opposite ramification. In general form it resembles the *D. ligulata*, but is of a totally different consistence, being much more rigid, coarser, and thicker. The fronds are sometimes curiously dilated towards the apex, when they are generally linear below the expanded portion. This appearance is produced by the upper pinnæ elongating and becoming thrice pinnated; they are then also of a paler colour than the lower ones.

PLATE CLXXII., CLXXIII. Fig. 1, transverse section of stem; fig. 2, horizontal, and fig. 3, vertical slice of the same :--magnified.

8. DICTYOSIPHON, Grev.

1. DICTYOSIPHON fasciculatus, Hook. fil. et Harv., in Fl. Antarct. Pt. 1. p. 178. t. 49. f. 1.

HAB. Berkeley Sound, Falklaud Islands, and Christmas Harbour, Kerguelen's Land.

We have before noticed this plant, which is the Southern representative of the Northern and Arctic D. faniculaceus.

9. STEREOCLADON, Hook. fil. et Harv.

Frons solida, olivacea, filiformis, ramosissima, e eellulis endochromate repletis longitudinaliter seriatis formata. Sporidia solitaria, sparsa, in frondis peripheria immersa, nigro-olivacea, elliptica.—Genus dubiæ affinitatis, vix in tribu Dictyotearum includendum.

1. STEREOCLADON Lyallii. (TAB. CLXXIV.)

HAB. Cape Horn, and the Falkland Islands; thrown up on the beach, rare.

Frons 5-6 une. longa, setacea, decomposito-ramosissima, ramificatione valde irregulari. Caulis percurrens v. parce divisus, vix dichotome ramosus. Rami alterni, patentes, flexuosi, decompositi; ramuli omnes patentes, flexuosi v. squarrosi, multifidi, apices acuti. Substantia rigidula, chartæ laxe adhærens. Color olivaceus. Sporidia numerosissima, per frondis partem superiorem dense sparsa, immersa.

This remarkable plant resembles, to the naked eye, *Dictyosiphon faniculaceus*; but its stem and branches are solid throughout, and the seeds are immersed endwise, in the substance of the branch.

PLATE CLXXIV. Fig. 1 and 2, portions of branch and ramuli; fig. 3, segment of ramulus; fig. 4, transverse section of fructifying stem :---magnified.

10. CHORDA, Stackh.

1. CHORDA lomentaria, Grev., Alg. Brit. p. 50. t. 9. Fl. Antarct. Pt. 1. p. 179.

HAB. Berkeley Sound, Falkland Islands, and Christmas Harbour, Kerguelen's Land; abundant.

Very abundant on the shores of Europe, from the Mediterranean Sea to the German Ocean. Also found in Lord Auckland's Group, but not, that we are aware, within the Tropics.

11. ASPEROCOCCUS, Lamours.

1. ASPEROCOCCUS sinuosus, Bory, Morea, p. 326. Encelium sinuosum, Ag. Sp. Alg. vol. i. p. 136.

HAB. Falkland Islands; abundant. Hermite Island, Cape Horn.

The distribution of this species is very wide, continuing through the Tropics from the latitude of Spain to the Falkland Islands. We have specimens from the collections of Humboldt; also from Vogel, gathered in Tropical Africa, and from the Red Sea and Persian Gulf. It neither inhabits Northern Europe nor is found on any shores south of the Falklands.

12. ADENOCYSTIS, Hook. fil. et Harv.

1. ADENOCYSTIS Lessoni, Hook. fil. et Harv., Fl. Antarct. Pt. 1. p. 179. t. 69. f. 2.

HAB. Hermite Island, Cape Horn; Falkland Islands; Kerguelen's Land, and Cockburn Island, lat. 64° 12' S., long. 57° W.; very abundant.

Apparently quite an Antarctic species, though much resembling some of the *Dumontiæ* figured in Postel and Rupprecht's great work on the *Algæ* of the Arctic and Paeific Occans.

2. ADENOCYSTIS D'Urvilleai, Hook. fil. et Harv. Asperococcus D'Urvillai, Bory in Duperrey Foy. Bot. p. 200. t. 11. f. 3.

HAB. Berkeley Sound, Falkland Islands; with the former.

We are inclined to regard this as a slender state of the *A. Lessonii*, which is exceedingly variable when young.

13. SPHACELARIA, Lyngb.

1. SPHACELARIA oborata, Hook. fil. et Harv.; parvula, gracilis, pallide viridis, stupa nulla, fronde circumscriptione obovata, caule gracili articulato basi longe nudo supra medium ramis plurimis tenuibus clongatis crecto-patentibus laxe distiche pinnatis ornato apicibus sphacelatis. Nobis *in Lond. Journ. of Bot.* vol. iv. p. 251.

HAB. St. Martin's Cove, Hermite Island, Cape Horn, in about eight fathom water; very scarce.

Cæspitosa, superne fastigiatim ramosa. Frondes $1-1\frac{1}{2}$ unc. longæ, eaule ramisque gracilibus, per totam longitudinem articulatis.

We have seen no specimens of this, but what were dredged up from a considerable depth; and, if fully grown, the outline of the frond alone is sufficient to distinguish it from its congeners.

2. SPHACELARIA funicularis, Mont. Fl. Antarct. Pt. 1. p. 180.

HAB. Cape Tres Montes, South Chili; C. Darwin, Esq.

The representative of the European S. scoparia. We have a note, unaccompanied, however, by any specimen, purporting that this species was also found in the Falkland Islands.

14. CLADOSTEPHUS, Ag.

1. CLADOSTEPHUS spongiosus, Agardh, Sp. Alg. vol. ii. p. 15. Engl. Bot. t. 2427. f. 1.

HAB. Hermite Island, Cape Horn, and the Falkland Islands; abundant.

This plant varies considerably at several periods of the year, becoming bare of ramuli in the winter. It is abundant in the German Ocean and on the Atlantic shores of Europe, and extra tropical North America, in the Mediterranean Sea and Canary Islands, but has not been hitherto found within the Tropies. We regard these specimens as specifically identical with others of British growth.

15. ECTOCARPUS, Lyngb.

1. ECTOCARPUS tomentosus, Lyngb. Grev. Crypt. Flor. t. 316.

HAB. St. Martin's Cove, Hermite Island, Cape Horn; rare.

The European shores of the Atlantic Ocean and the German Sea are the only previously recorded habitats for this species.

2. ECTOCARPUS siliculosus, Lyngb. Engl. Bot. t. 2319.

HAB. Hermite Island, Cape Horn, and Berkcley Sound, Falkland Islands ; abundant.

Apparently a much more widely distributed species than the former, ranging from the Baltic Sea and German Ocean to the Mediterranean and Canary Islands, also along the shores of the United States. In the Southern hemisphere it has been found at the Cape of Good Hope and New Zealand.

3. ECTOCARPUS geminatus, Hook. fil. et Harv.; cæspite basi intricato ramoso olivacco v. vircscentc, filis majusculis tenuibus ramosissimis apice liberis plumosis, ramis ramulisque patentibus oppositis quaternisve ultimis brevibus, utriculis sessilibus oppositis conicis basi sæpe ramulo brevi bractcæformi suffultis. Nobis in Lond. Journ. Bot. vol. iv. p. 251.

HAB. Hermite Island, Cape Horn, and the Falkland Islands; abundant.

Cæspites 4-5 unc. longi, basi e ramulis perplurimis implexis intricati. Rami primarii circumscriptione lineariobovati, plerumque quaterni, secundarii ramulique ultimi oppositi, patentes. Utriculi semper oppositi, ramulo bracteæformi duplo longiores.

In habit and general appearance resembling the European *E. granulosus*; but abundantly different in the constantly opposite sessile conical capsules or utricles, which are generally subtended by a minute ramulus half their own length.

16. MESOGLOIA, Ag.

1. MESOGLOIA *linearis*, Hook. fil. et Harv.; virescens, fronde circumscriptione lineari, caule gracili indiviso v. rarius diviso ramis brevibus ornato, ramis alternis crebris abbreviatis flexuosis erecto-patentibus, ramulis subsecundis. Nobis *in Lond. Journ. Bot.* vol. iv. p. 251.

HAB. Hermite Island, Cape Horn; rare.

Frondes 4-6 unc. longi, vix $\frac{1}{2}$ lin. diametro, pallide flavo-virescentes. Caulis gracilis, indivisus v. basi ter quaterve divisus, rarius medium versus in ramos 2 primarios fissus. Rami perplurimi, $\frac{1}{4}-\frac{1}{2}$ unc. longi, flexuosi, interdum basin versus caulis nudi v. ramulis paucis aucti. Ramuli secundarii plerumque e margine inferiore seu exteriore ramorum orti. Peripheriæ filamenta moniliformia, e substantia gelatinosa vix exserta.

Probably the representative of the European M. vermicularis, from which it may be at once distinguished by the ramification.

17. DELESSERIA, Lamourx.

1. DELESSERIA sanguinea, Lamourx. Engl. Bot. t. 1041.

Var. B. lancifolia; fronde clongata anguste lineari-lanceolata ligulatave utrinque angustata.

HAB. Hermite Island, Cape Horn; on rocks, abundant near the shore, also dredged up in five or six fathom water.

Hitherto known only as an inhabitant of the seas of the Northern hemisphere; where its range is not wide. In the southern it appears to be confined to the extreme south of America, flourishing in the deep bays which indent the coasts of Fuegia. Some specimens are altogether similar to those of European growth; others, of which we have constituted the var. *lancifolia* are larger, longer, sometimes almost a foot long, much narrower, and more attenuated at both ends. It is a very handsome variety.

2. DELESSERIA *Davisii*, Hook. fil. et Harv.; caule cartilagineo alato, lamina profunde pinnatifida v. pinnata, laciniis pinnisve cultrato-lanceolatis obliquis costatis penninerviis, nervis alternis, demum inter nervos alterne v. secunde lacerato-laciniatis, lacinulis erecto-patentibus costatis. Nobis *in Lond. Journ. Bot.* vol. iv. p. 52. (TAB. CLXXV.)

HAB. Hermite Island, Cape Horn; abundant. Falkland Islands, Berkeley Sound, Dr. Lyall.

Frons 5-7-uncialis, alata v. latiuscule marginata, basi in caulem brevem abrupte attenuata, circumscriptione late ovato-rotundata, in lacinias perplurimas simplices v. partitas distichas costatas divisa, rosea, membrauacea. Laciniæ plerumque sceus marginem exteriorem oblique ad costam fissæ, $\frac{1}{4}-\frac{1}{2}$ unc. latæ, obtusæ v. subacutæ.—Inter D. alatam et D. sanguineam quasi media, sed utraque sat diversa.

This is perhaps most closely allied to *D. sanguinea*, from which it may be distinguished by the alternate nervation and dividing of the frond. The *D. alata*, which it is also near, differs in the texture and colour of its frond. No fruiting specimens were found.

FLORA ANTARCTICA.

PLATE CLXXV. The absence of fructification precludes the necessity of dissections.

3. DELESSERIA platycarpa, Lamourx., in Ann. du Mus. vol. xx. p. 124. Agardh, Sp. Alg. vol. i. p. 188. Turner, Hist. Fuc. t. 144.

HAB. Falkland Islands; abundant in Berkeley Sound and Port William.

Originally discovered at the Cape of Good Hope, where it is very abundant. More recently it has been gathered on the coast of California, whence its existence in the Falkland Islands might have been considered probable, even before its recent discovery in that part of the Southern Ocean. It is very plentiful on the long shelving beaches of the islands in question, but was not seen on the deeper and more rocky shores of Fuegia.

4. DELESSERIA crassinervia, Mont. Fl. Antarct. p. 184.

Var. β . costa angustiore.

HAB. Hermite Island, Cape Horn, and the Falkland Islands; both varieties abundant. Kerguelen's Land; Christmas Harbour, var. a only.

Were the var. β . found upon the coast of Europe, it would be undonbtedly referred to *D. hypoglossum*, and it may well be considered very doubtful whether the *D. crassinervia* of the Southern Hemisphere be the representative of its northern congener, into the likeness of which it varies; or whether, as is perhaps more probable, both are not varieties of the same species. The true *D. ruscifolia* is a native of the Cape of Good Hope, of Tasmania, and, according to Gaudichaud, of the Falkland Islands also.

5. DELESSERIA quercifolia, Bory in Duperrey Voy. Bot. p. 186. t. 18. f. 1.

HAB. Hermite Island, Cape Horn, and the Falkland Islands; most abundant both on the outer coasts and in the harbours.

A very handsome species, of which M. Bory gives a sufficiently characteristic but discoloured figure. The original colour of the plant is a rosy or vinous red. In every respect, except the position of the granules, which are here scattered over the surface of the frond, this is very nearly allied to D. sinuosa, Ag.

6. DELESSERIA Lyallii, Hook. fil. et Harv.; fronde lineari-oblonga obtusa costata penninervi argute serrato-dentata, nervis oppositis, margine incrassato folia consimilia petiolata emittente, dentibus subulatis simplicibus v. latere inferiore plerumque erosis, coccidiis frondis pagina sparsis, granulis in soros inter nervos laciniarum sitos dispositis. Nobis *in Lond. Journ. Bot.* vol. iv. p. 252. (TAB. CLXXVI.)

HAB. Falkland Islands; on the outer coast only, probably washed ashore from the exposed rocks. Kerguclen's Land; Christmas Harbour, washed up on the shores.

Frons primaria exemplaribus Kerguelensibus 9 unc., Falklandicis 4-5 unc. longa, $1-1\frac{1}{2}$ unc. lata, in petiolum cylindraceum gradatim angustata, oblonga v. linearis, apice rotundata, costa valida percursa, venosa, venis oppositis, erecto-patentibus, margine argute serrato-dentata, sed non sinuata, incrassata, foliola seu frondes secundarias emittens? *Frondes secundariæ* primariis consimiles, sed plerumque minores, omnes evidenter petiolatæ, c margine incrassato frondis primariæ ortæ, venis ejus oppositæ v. alteruæ. *Color* luride sanguineus.

D. sinuosæ habitu formaque frondis colore et substantia simillima : differt margine incrassato, dentato, non sinuato, et præsertim frondibus secundariis evidenter petiolatis, nunquam e laciniis frondis primariæ ortis.

A very noble species, of which fine specimens were collected, thrown up on the stony shores of Kergueleu's Land and the outer coasts of the Falkland Islands; and which, as it was never seen attached, either in the shallow or deep bays of any of the coasts visited by the Expedition, finds, we conclude, a congenial home amongst the wild breakers that fringe many parts of these iron-bound coasts. Specifically it is allied to the European *D. sinuosa*,

from which it differs remarkably in the thickened margin of the frond not being sinnated, but proliferous; in the leaves all being petiolate and arising from the margin, and not from lacinize of the frond; and in the position of the fructification.

We have, in figuring the nobler species of this and some other genera, endeavoured to commemorate the services rendered to the botany of the Antarctic regions by those officers of the Antarctic Expedition who particularly devoted themselves to increasing the botanical collections. Their names appear to be more properly associated with the Algae, than with any other tribe of plants; comprising, as these do, the greater part of the vegetation of that element which these gentlemen have adopted for their home, and being natives of the regions they have so successfully explored.

PLATE CLXXV. Fig 1, apex of frond aud sori; fig. 2, portion of ditto showing the spherospores :- highly magnified.

18. NITOPHYLLUM, Grev.

1. NITOPHYLLUM *lividum*, Hook. fil. et Harv.; fronde e stipite brevi filiformi cartilagineo late expansa tenerrima basi vix venosa furcata v. dichotoma margine undulata livido-purpurea, laeiniis patentibus oblongis obtusis, soris minutissimis punctiformibus coecidiisque perplurimis per totam frondem sparsis. Nobis *in Lond. Journ. Bot.* vol. iv. p. 253. (TAB. CLXXIX.)

HAB. Falkland Islands; Berkeley Sound and Port William, not uncommon.

Stipes cartilagineus, filiformis, $\frac{1}{2}-1$ unc. longus, ad basin frondis evanidus. Frons 4 unc. longa, 6 v. plures lata, in lacinias paucas latinsculas furcatas apice obtusas divaricatas divisa, avenia, nisi ad imam basin, ubi stipes in venas breves evanidas abiit. Substantia tencrrima. Color livide purpureus, ut in Porphyra, sed vix nitens.

The colour affords a very distinctive character for this species, in which particular it resembles only one of its congeners, the *N. Gunnianum*, Harv., of Tasmania. But that plant, is of a much thicker texture and less lubricous. A single imperfect specimen from Cape Horn probably belongs to the *N. lividum*. Of the mass of radiating spores contained in the capsules of the species, only those at the base of the cavity are fertile.

2. NITOPHYLLUM *fusco-rubrum*, Hook. fil. et Harv.; stipite filiformi elongato nunc dichotome ramoso nudo, ramis frondiferis, frondibus flabelliformibus lobatis v. longitudinaliter fissis crasso-membranaceis fuscorubris, basi cuneatis in stipitem gradatim angustatis tenuiter venosis, margine plano subintegerrimo, apicibus (exemplaribus nostris) laceris, soris minutissimis punctiformibus coccidiisque numerosissimis per totam frondem sparsis. Nobis *in Lond. Journ. Bot.* vol. iv. p. 254.

HAB. Kerguelen's Land; parasitical on larger sea-weeds in Christmas Harbour.

Stipes 1-8 unc. longus, simplex v. irregulariter ramosus, ramis in frondes cuneatas elongatas exeuntibus. Frondes 3-5 unc. longæ, latitudine variæ, ima basi obseure venosæ, irregulariter profunde fissæ, laciniis euneatis linearibusve. Sori minimi, inconspicui. Splærosporæ plerumque solitariæ, per totam paginam frondis creberrime sparsæ. Coccidia frondibus distinctis numerosa. Substantia firma, basi subcartilaginea. Color luride fusco-ruber.— Stirps N. ulvoideo, Hook. similis, sed abunde differt colore, sphærosporis sparsis, stipiteque ramoso elongato.

Apparently a native of Kerguelen's Land only, where it was found sparingly, adhering to the stems of larger Alga. The colour, texture, and branching stem at once distinguish this from its congeners.

3. NITOPHYLLUM Crozieri, Hook. fil. et Harv.; fronde basi longe cuneata in stipitem angustata linearilanceolata v. ovata v. late ovato-lanceolata integerrima v. in lacinias plurimas longitudinaliter fissa enervi

tenerrima rosea, soris majusculis oblongis coccidiisque per frondem sparsis. Nobis in Lond. Journ. Bot. vol. iv. p. 254. (TAB. CLXXVII.)

HAB. Hermite Island, Cape Horn; abundant in deep water.

Radix discus cartilagineus. Stipes gracilis, $\frac{1}{2} - \frac{3}{4}$ unc. longus, cartilagineus, superne alatus, basin in frondem cuncatam abeuns, deinde gradatim evanescens. Frons 8–12 unc. longa v. longior, latitudine varia, lineari-oblonga v. late ovato-rotundata, integerrima v. in lacinias fissa, adultior punctis crebriformibus pulcherrime terebrata, apice exemplaribus normalibus attenuata, margine integerrima, undulata, plus minusve in lacinias fissa, subavenia v. basi solum nervis indistinctis e apice stipitis ortis notata. Sori numerosissimi, per totam frondis paginam sparsi. Substantia tenerrima. Color pulcherrime roseus.

One of the most beautiful of the genus, and probably the southern representative of the European *N. punctatum*, chiefly distinguishable from it by the long euncate base of the frond passing into a filiform stem and by the absence of dichotomous divisions with wide axils. The traces of the stem become gradually more faint at a short distance from the base of the frond, but do not break up into numerous veins. The normal form of the frond is broadly lanceolate, tapering to an acute point; with waved but entire margins, which are, however, often split and torn into numerous linear ribbon-like segments, caused by injury and not the natural divisions of the frond.

PLATE CLXXVII.-Fig. 1, portion of frond and sorus :- magnified.

4. NITOPHYLLUM *multinerve*, Hook. fil. et Harv.; fronde breviter stipitata elliptica v. ovata subintegerrima v. lobata, nervis pluribus parallelis distinctis dichotomis apicem versus frondis evanescentibus, soris? Nobis *in Lond. Journ. Bot.* vol. iv. p. 255.

HAB. Hermite Island, Cape Horn, and Falkland Islands; Berkeley Sound, on rocks, not common.

Stipes plerumque $\frac{1}{8} - \frac{1}{4}$ unc. longus, nunc pollicaris, simplex. Frons in lacinias plurimas obtusas cito fissa, 2-4 unc. longa, basi rotundata v. cuncata; nervis plurimis, validis, ad apicem frondis evanidis. Substantia membranacea. Color roscus.

Of this plant we have not very satisfactory specimens, or which may not belong to the *Delesseria dichotoma*: except that in the present species the nerves are much fainter, less distinct from the lamina, and vanishing further from the apex of the frond, which evinces no tendency to form distinct leaves.

5. NITOPHYLLUM *Smithii*, Hook. fil. et Harv.; fronde stipitata flabelliformi lobata basi cuneata superne divisa et lacera, apicibus laciniarum obtusis, marginibus planis, colore rubro subfuscescente, nervosa, nervis gradatim evanescentibus basilari centrali crasso lateralibus radiantibus tenuibus nunc evanidis, soris minutis rotundatis margines versus laciniarum frondis densissime sparsis. Nobis *in Lond. Journ. Bot.* vol. iv. p. 256. (TAB. CLXXVIII.)

HAB. Falkland Islands; in the bays and along the outer sea coast, growing on the roots of larger Alga.

Frons 4-7 une. longa, angusta v. latiuseula, in stipitem simplicem v. ramosam $\frac{1}{2}$ unc. longam et ultra desinens, forma varia; nunc elongatæ, laceræ; juniores latiores, lobatæ: laciniis latis, obtusis, emarginatis retusisve. Nervus centralis elongatus, basi latus, frondibus senioribus ultra medium extensus, junioribus eito evanidus; laterales ad basin frondis flabellatim expansi, oblique areuati. Substantia crassiuscula. Color ruber, demum fuscescens.

This is a very distinct species ; but, like its congeners, so variable in form that little dependence can be placed on the characters drawn from its ontline, or from the length of the stipes. Our figure gives a very faithful representation of the colour and normal form of the old and young states, both veined and nearly veinless.

PLATE CLXXVIII .- Fig. 1, portion of frond and tetraspores; fig. 2, ditto with coccidium :- both magnified.

5 r

6. NITOPHYLLUM Bonnemaisoni, Grev.; Alg. Brit. p. 81.

Var. laciniatum, fronde flabelliformi profunde digitatim lohata v. subdichotome pinnatifida, laciniis inciso-dentatis. N. laciniatum, nobis in Lond. Journ. Bot. vol. iv. p. 256.

HAB. var. laciniatum. Hermite Island, Cape Horn; on rocks in deep water. Falkland Islands; not common.

Though not exactly identical with our European N. Bonnemaisoni, we cannot consider this as more than a variety of that plant, which is occasionally found in Britain nearly as much laciniated as the specimens before us are. Stipes, in the Antarctic specimens, $\frac{1}{4}-\frac{1}{2}$ inch long, terminating in the thickened cuncate base of the frond, which is 4-5 inches long, and deeply cleft into 5-9 segments, either radiating from the centre in a digitate manner, or springing like pinnules from a lengthened rachis. Colour, a pale brownish-red at the base, becoming rosy upwards.

The N. Bonnemaisoni in the Northern Hemisphere inhabits the Orkney Islands, the west coasts of Scotland and Ireland, and the south-western shores of England.

19. PLOCAMIUM, Lyngb.

1. PLOCAMIUM coccineum, Lyngb. ; Harv. Phyc. Brit., t. 44. Fl. Antarct. p. 186.

HAB. Hermite Island, Cape Horn, and the Falkland Islands; very abundant.

The present is one of the most widely diffused of the *Florideæ*, both in the Northern and Southern Hemisphere. In Europe it ranges from the shores of the Mediterranean Sea to the North Cape. In Africa it occurs at the Cape of Good Hope, in North America on the coast of California, and it is abundant, on the Atlantic shores of the United States. It inhabits both coasts of South America, but particularly the western; Tasmania, New Zealand, and the Anekland Islands. The Antarctic specimens are equally luxuriant with those of the Northern Hemisphere.

2. PLOCAMIUM Hookeri, Harv.; fronde cartilaginea anguste lineari compressa plana distiche decomposite ramosa, ramis primariis subdichotomis patentibus, secundariis alternis flexuosis folia ramulosque alterne gerentibus, foliis planis aveniis oblique obovato-lanceolatis obtusis basi angustatis cultratis integerrimis v. margine exteriore crenatis, ramulis linearibus alterne et secunde pectinato-multifidis, stichidiis brevibus lateralibus dense fasciculatis digitatis laciniatis simplicibus obtusis, coccidiis lateralibus sessilibus sparsis. *Harv. in Lond. Journ. Bot.* vol. iv. p. 257.

HAB. Kerguelen's Land; Christmas Harbour; thrown upon the beach, rare.

Frons 8-10 une. longa, vix lin. diametro, plano-compressa, cartilaginea, nunc subdichotome nunc pinnatim ramosa. Rami primarii patentes; secundarii eircumscriptione lineares, alterni, flexuosi, ramulos decompositos foliaque gerentes: foliis $\frac{1}{2}$ une. longis, $1\frac{1}{2}$ -3 lin. latis, anguste obovatis lanceolatisve, obtusis, aveniis, integerrimis v. rarius secus marginem exteriorem crenatis. Color luride roscus.

One of the most singular species of the genus, from the curious leaf-like appendages on the secondary and lesser branches. It is a very rare plant in Christmas Harbour, and cannot be confounded with any of its congeners.

3. PLOCAMIUM? Magellanicum, Hook. fil. et Harv. Thamnophora Magellanica, Montagne in Voy. au Pole Sud, Bot. Crypt. p. 142. t. S. f. 2.

HAB. Hermite Island, Cape Horn; the Falkland Islands, and Kerguelen's Land; very abundant.

FLORA ANTARCTICA.

475

Of this plant we have very copious suites of specimens, gathered in the localities above enumerated and varying in length from 2 or 3, to 4 or 8 inches; most of them are covered with coccidia, though none presents us with a single stichidium; which is the more remarkable, because, in other species of this genus the latter description of fruit is the most general.

Under the *P. coccineum*, in the first portion of this work, the reasons for abandoning the genns *Thamnophora* are detailed; whether or not the present plant belongs to *Plocamium* even, must remain uncertain until the nature of the stichidia is known.

20. RHODYMENIA, Grev.

1. RHODYMENIA palmata, Grev.; Alg. Brit. p. 93. Fucus palmatus, Engl. Bot. t. 1306.

HAB. Berkeley Sound, Falkland Islands; abundant. Hermite Islands, Cape Horu; rare.

The Dulse, so commonly eaten on the coasts of Scotland, is not an unfrequent sea-weed on the shores of the Falkland Islands, where it was quickly recognized by some of the north-country seamen of the 'Erebus' and 'Terror.' In Europe its distribution is from the Canary Islands and Mediterranean Sea, to the coasts of Norway and Ireland. Dr. Greville mentions that it is a native of the shores of Brazil.

2. RHODYMENIA sobolifera, Grev.; Alg. Brit. p. 95. Fucus soboliferus, Eng. Bot. t. 2133.

HAB. Falkland Islands; in Berkeley Sound, and on the exposed outer sea-coast; abundant.

Apparently identical with a sea-weed which inhabits the western shores of Ireland, Scotland, and the Orkney Isles, and has also been gathered on the west coasts of France. It is scarcely more than a variety of R. palmata, whether occurring in the north or south temperate oceans.

3. RHODYMENIA corallina, Bory; in Duperrey Voy. Bot. Crypt. p. 175. t. 16.

HAB. Christmas Harbour, Kerguelen's Land; rare.

We have referred our single specimen, without fruit, to this species; with which it appears entirely to agree. The species is not uncommon along the Pacific shores of South America, between the latitude of Concepcion and the Equator.

4. RHODYMENIA Palmetta, Grev.; Alg. Brit. p. 88. t. 12.

HAB. Straits of Magalhaens, D'Urville; Falkland Islands, Gaudichaud.

Of this species we have seen no southern examples.

5. RHODYMENIA fimbriata, Grev.; Synops. p. 48. Sphærococcus fimbriatus, Agardh, Spec. Alg. vol. i. p. 299.

HAB. Falkland Islands, Gaudichaud.

This again is a plant which we do not recognize amongst the collections brought from the Southern Hemisphere.

6. RHODYMENIA variegata, Montague; in D'Orbigny Voy. p. 22. and 116 in Obs. Halymenia variegata, Bory in Duperrey Voy. Bot. Crypt. p. 179. t. 14. R. Hookeri, Harv. in Lond. Journ. of Bot. vol. iv. p. 258. R. glaphyra, Suhr, in Flora, 1839, vol. i. p. 69. t. 2. f. 43.

Var. a. flabellata; fronde stipitata rosea v. sanguinea flabellata fere ad basin partita, laciniis manifeste flabelliformibus basi cuneatis repetite di-tri- vel palmatim dichotomis, laciniis linearibus $\frac{1}{4} - \frac{1}{2}$ unc. latis, margine lacinulis brevissimis truncatis quadratis alternis ornato, axillis rotundatis :—R. Lambertæ forma similis, sed substantia differt.

Var. β . *atro-sanguinea*; fronde substipitata atro-sanguinea palmato-fissa, laciniis obtusis crectis subdichotome v. alterne divisis margine proliferis, axillis rotundatis. Color luridior quam in var. a.

Var. γ . *latissima*; fronde 10 unc. longa ad pedalem, laciniis parum divisis apice truncatis 1-4 unc. latis. Varietas a cæteris valde diversa, scd certe nil nisi forma gigantea.

Var. 8. lacerata ; inter varietates a et β media :--frons subsessilis divisa.

Var. ϵ . prolifera ; fronde $1\frac{1}{2}-2$ unc. longa subdichotoma, laciniarum marginibus proliferis lacinulas numerosas angustissimas furcatas v. irregulariter ramulosas acutas emittentibus.

Var. 4. pulcherrima; laciniis angustis decomposito-ramosis, pinnulis ultimis elongatis emarginatis.

HAB. Hermite Island, Cape Horn; var. a. Falkland Islands; var. a. β . (on the outer sea-coast) and ζ . (in Berkeley Sound) Kerguclen's Land; vars. a. γ . δ . and ϵ . All very abundant in Christmas Harbour.

A more variable species can scarcely be imagined: so dissimilar are its forms that the more distinct of them were unhesitatingly pronounced to be different species, before the whole suites of specimens were collated. In Kerguelen's Land it is one of the most common of Alga; and the varieties, collected there and noted as belonging to the one species, are connected by various links with one another, and with the forms of Cape Horn, the Falkland Islands, and of the American coast. The dark coloured variety, β . *atro-sanguinea*, is evidently sea-beaten, and though generally destitute of the marginal tooth-like laciniæ, so conspicuous in *a*, there are specimens possessing them, which connect the two forms. The var. γ . *latissima* is the best marked, more, perhaps, by its great size than by its presenting any positive characters: it was gathered along with *a* and δ , and referred when fresh to the same species. In the Falkland Islands the var. ζ . is conspicuous for having few, and but sparingly divided principal segments, about $\frac{1}{4}$ inch wide, suddenly passing into narrow much-divided minor segments from $\frac{1}{2}-1$ line broad. Though at first sight abundantly distinct from β . or γ ., it is immediately connected with them both through var. *a*.

We follow Endlicher in quoting Suhr's figure of R. glaphyra as a synonym of this species: the representation is, however, anything but characteristic of an Alga.

7. RHODYMENIA variolosa, Hook. fil. et Harv.; fronde carnoso-membranacea sanguinea in lacinias plures late lineares v. cuneatas elongatas furcatas dichotomasve ad basin fere divisa, laciniis basi angustatis erectis apice obtusis emarginatisve, coccidiis? superficialibus densissime conspersis sessilibus pedicellatisve deciduis. Nobis *in Lond. Journ. Bot.* vol. iv. p. 259. (TAB. CLXXX.)

HAB. Christmas Harbour; Kerguelen's Land.

Frons 2-3 v. 7-8 unc. longa, ad basin fere in lacinias fissa. Laciniæ $\frac{1}{2}$ -1 unc. latæ, e basi angustata sensim latiores, furcatæ, bis terve dichotome divisæ, sinubus latis obtusis, margine integerrimo v. parce prolifero, super sub-terque corporibus granuliferis (coccidia?) sparsæ. Coccidia? (in genere abnormalia) superficialia, subglobosa, basi angusta frondis affixa, cito decidua.

Allied to the R. variegata, especially in form, but of a firmer texture and brighter colour; and very different in the nature of the fructification, which is easily detached, leaving only a small puncture on the surface of the laciniæ: this constitutes the peculiar character of the species. In the structure and form of the frond there is some affinity with the *Gracilaria polycarpa* of the south of England and California; but the fruit of that plant is quite different.

PLATE CLXXX.—Fig. 1, portion of frond and coccidia; fig. 2, portion of ditto vertically sliced; fig. 3, spores from the same; fig. 4, immature ditto :—all highly magnified.

21. ACANTHOCOCCUS, Hook. fil. et Harv.

Frons linearis, compressa, distiche ramosa, cartilagineo-carnosa, rosea. Axis solidus, densus, e cellulis minutissimis formatus, tubulis magnis pluriseriatis extus sensim minoribus circumdatus. Peripheria celluli parvis reticulata. Coccidia globosa, in apicibus ramulorum immersa, sporis numerosissimis repleta.

1. ACANTHOCOCCUS Antarcticus, nobis, in Lond. Journ. Bot. vol. iv. p. 261. (TAB. CLXXXI.)

HAB. Cape Horn and the Falklaud Islands; not uncommon, and parasitic.

Frons 4-8 unc. longa, compressa, anguste linearis, basi semilineam vix lineam latitudine, sursum sensim angustata, distiche ramosissima. Rami patentes vel divaricati, nunc flabellatim multifidi, nunc pinnati et bipinnati; secundarii nunc breves subsimplices, nunc longissimi, ramosissimi. Ramuli per totam frondem sparsi, apicem versus erebriores, erecti et erecto-patentes, subulati, 1-3 lineas longi, alterni vel sæpius secundi, simplices vel parum divisi. Coccidia solitaria, globosa, spinis 4-6 magnis subulatis armata, in apicibus ramulorum immersa, sporis numerosissimis miuutis repleta. Tetrasporæ ignotæ. Color intense ruber, siecitate obscurior. Substantia firma, cartilagineocarnosa :---chartæ adhæret.

We cannot satisfactorily include this plant under any established genus. It belongs, unquestionably, to the *Sphærococceæ* and will stand near *Hypnea*, from which it differs in the structure of the frond, as well as in the fructification. The densely cellular axis, surrounded by large empty cellules or tubes, is seen in *Hypnea musci-formis*, and also in *Gracilaria purpurascens*. Outwardly there is a close resemblance between our plant and *Heringia rostrata*, J. Ag., (*Gelidium ? rostratum*, Griff.; *Fucus alatus*, and *angustissimus*, Turn.); but, besides the dissimilar fructification, the structure of that plant is uniformly dense, without a trace of large cellules, or tubes. Again, the present plant may be compared with *Microcladia*, which it approaches in habit; and to a certain extent, the spinous *coccidia* may be deemed analogous to the involucrated *favellæ* of that genus; but, in *Microcladia*, the axis, far from being the most dense part of the frond, is tubular.

PLATE CLXXXI.—Fig. 1, ramulus; fig. 2, apex of ditto with coccidium; fig. 3, transverse section of ramulus; fig. 4, longitudinal section of coccidium:—all magnified.

22. GRACILARIA, Grev.

1. GRACILARIA (?) nigrescens, Hook. fil. et Harv.; radice fibrosa, frondibus purpurascentibus cæspitosis e basi irregulariter dichotoma et intricata ramosissimis gracilibus subcylindraceis obscure compressis flexuosis flaccidis carnoso-membranaceis, axillis obtusis sæpissime latis, ramis decompositis sensim angustatis, ramulis filiformibus v. subulatis acutis, ultimis sæpe secundis. G. obtusangula, nobis in Lond. Journ. Bot. vol. iv. p. 260. Sphærococcus subulatus, β. nigrescens, Agardh, Sp. Alg. p. 329.

Var. B. tenuior, ramis strictioribus divaricatis, axillis patentibus.

HAB. Hermite Island, Cape Horn, and the Falkland Islands; not uncommon. Var. β . Falkland Islands.

Frons basi repens et fibrosa, filiformis, 4-6 unc. longa, $\frac{1}{4}$ lin. lata, subcompressa. Color luride purpureus, ut in G. purpurascente.

In the absence of fructification we refer this doubtfully to the genus *Gracilaria*, chiefly from its close resemblance to the *G. purpurascens* in the essential characters of the frond. We have never seen original or any other specimens of the *Spharococcus subulatus*, var. *nigrescens*, and owe the identification of our specimens with that plant to the kindness of our friend Dr. Montagne.

2. GRACILARIA (?) aggregata, Hook. fil. et Harv.; cæspitosa, nigrescens, frondibus fastigiatis e basi communi late scutata carnosa ortis plurimis filiformibus, primariis cylindraceis cartilagineis vage subdichotome ramosis, axillis augustis, ramis erectis simplicibus furcatisve ommibus compresso-cylindraceis filiformibus superne subfastigiatis apicibus obtusis, fructu ——? Nobis *in Lond. Journ. Bot.* vol. iv. p. 261.

HAB. Falkland Islands; on rocks in Berkeley Sound, Dr. Lyall.

Frondes 3-4 une. altæ, vix $\frac{1}{2}$ lin. diametro, e basi scutata, $\frac{1}{2}$ unc. latæ, dense fastigiatæ, siecitate rigidæ, madore cartilagineæ, irregulariter superne præcipue ramosæ, basi cylindraceæ, interdum simplices, nunc e basi regulariter dichotome ramosæ. Rami omnes erecti, axillis angustis, sinubus obtusiusculis, ultimi paulo latiores. Color nigro-purpurascens, siecitate ater ;—chartæ non adhæret. Habitu Polyidis rotundi.

Of this, again, we have seen no fruit; but the structure of the frond is exactly that of the genus *Gracilaria*, to which we consequently refer it.

23. POLYSIPHONIA, Grev.

1. POLYSIPHONIA atro-rubescens, Grev. Harv. in Brit. Fl. vol. ii. p. 331.

HAB. Berkeley Sound, Falkland Islands; scarce.

Only a few specimens of this species, and in an immature state, were procured: they very closely approach British ones in structure, but are, perhaps, more branched; though we can detect no specific differences between them. They differ from the following species in substance and in wanting the fibrillous apices of the ramuli; the latter, however, is an unimportant character.

The species ranges from the Mediterranean Sca to the coasts of Scotland.

2. POLYSIPHONIA *fusco-rubens*, Hook. fil. et Harv.; atro-rubescens, obscure articulata, rigidiuscula, multistriata, frondibus irregulariter dichotomis, caule angulatim flexuoso gracili sensim attenuato, ramis majoribus dichotomis, minoribus alternis strictis elongatis, ramulis paucis subulatis erecto-patentibus, axillis primariis patentibus secundariis acutis, articulis e tubulis octo radiantibus coloratis formatis, ramorum diametro duplo-triplo-quadruplove, ramulorum sesquilongioribus. (TAB. CLXXXII. Fig. I.)

HAB. Falkland Islands; Mrs. Capt. Sulivan.

Caules primarii ramosissimi, fasciculati, 6-8-unciales, graciles, filiformes, basi uudi, superne sub-augulatim flexuosi, ad angulos ramos 2 unc. longos emittentes, rigiduli. Rami ramulique crecto-patentes, laxe pluries divisi, fasciculati, ultimi longitudine varii, $\frac{1}{4}$ -2 lin. longi, elongati v. subulati, omnes rigidiusculi. Color atro-rubeseens :--- ehartæ laxe adhæret.

Very similar indeed to the *P. atro-rubescens*, but differing in the flexuose stem; also allied to *P. anisogona*, nob., but of a totally different texture and consistence.

PLATE CLXXXII. Fig. 1.-1, ramuli; 2, apex of ditto; 3, transverse section of ditto :--all magnified.

3. POLYSIPHONIA anisogona, Hook. fil. et Harv.; atro-rubescens, flaccida, madore fragillima, frondibus cæspitosis irregulariter ramosissimis equalibus setaceis articulatis vix attenuatis, ramis ramulisque alternis subdichotomisve crectis v. appressis, axillis angustissimis, articulis variis inferioribus diametro sextuplo, superioribus duplo triplove longioribus, ultimis sesquilougioribus v. quadratis, omuibus striis sex notatis e tubulis duodecim tenuibus radiantibus endochromate repletis formatis, ceramidiis ——? Nobis in Lond. Journ. Bot. vol. iv. p. 268. (TAB. CLXXXII. Fig. II.)

HAB. Hermite Islaud, Cape Horn, and the Falkland Islands; not common.

Dense cæspitosa. Cæspites 4-5 une. longi, intricati. Articuli longitudine varii, inferiores valde clongati, supremi brevissimi, omnes striis sex rectis spiralibusve notati, e tubulis duodecim tenuibus coloratis eirea cavitatem centralem dispositis conflata.

A fine species, and evidently quite distinct from any previously described; but unfortunately so tender that it eannot be removed from the paper without breaking; for which reason our description of the ramification is not so perfect as is desirable. It differs, in the substance especially, from the British *Polysiphonia atro-rubescens*, being more fragile and tender.

PLATE CLXXXII. Fig. II.—.A. and B. different states of P. anisogona; 1, ramulus; 2, apex of ditto; 3, transverse section of ditto:—all magnified.

4. POLYSIPHONIA *tenuistriata*, Hook. fil. et Harv.; rubescens, articulata, multistriata, frondibus gracillimis capillaceis flaceidis elongatis circumscriptione ovato-lanceolatis, caule primario subsimplici flexuoso alterne irregulariter dichotome ramoso, ramis remotis circumscriptione ovatis ramulisque erecto-patentibus sensim attenuatis apice fibrillosis, axillis acutis, articulis ramorum diametro multiplo, ramulorum duplo triplove longioribus sex-striatis e tubulis duodecim tenuissimis radiantibus coloratis formatis, geniculis incrassatis. Nobis *in Lond. Journ. Bot.* vol. iv. p. 266. (TAB. CLXXXII. Fig. III.)

HAB. Hermite Island, Cape Horn; dredged up in about six fathom water.

Basis frondis deest. *Caulis* primarius solitarius? 4–6 unc. longus, capillaris, flexuosus, alterne ramosus. *Rami* 2–3-unciales, ramulique tenuissimi.

A very elegant species, allied to *P. anisogona*, but much more slender, very differently branched, and not fragile when moistened after being once dried.

PLATE CLXXXIII. Fig. III.—1, plant of the natural size; 2, branch aud ramulus; 3, ramulus; 4, transverse section of ditto:—magnified.

5. POLYSIPHONIA *Sulivanæ*, Hook. fil. et Harv.; pusilla, badia, articulata, flaccida, multistriata, frondibus flabellatim ramosis, ramis alterne decompositis fastigiatis, ramulis sparsis alternis subulatis simplicibus subquadrifariis, articulis ramorum diametro subduplo, ramulorum sesquilongioribus omnibus e tubulis duodecim angustis radiantibus formatis. (TAB. CLXXXII. Fig. IV.)

HAB. Falkland Islands; Mrs. Capt. Sulivan.

Radix? Frondes cæspitosæ, 1 une. longæ, e basi valde fastigiatim ramosæ, flabellatim expansæ. Caulis primarius brevissimus, ramos plurimos capillares repetitim divisos fasciculatos emittens, rami penultimi subpectinati, ultimi curvati patentes, axillis obtusiuseulis, supremi ramos ramulosque terminantes arete incurvi sese invicem ampleetentes, apices frondinm hine nodosi v. incrassati apparent. Color badius. Substantia flaceida, tenax :— chartæ adhæret.

A pretty but small species, allied to the Anekland Islands *P. ceratoclada*, Mont., but slenderer; with the stem formed of a greater number of tubes, more flaceid, &c. The ultimate ramuli are longer in proportion than those they spring from, they are slender, patent, rather uniform in length, and eurve upwards, hence giving a somewhat pectinated appearance to the ultimate divisions of the frond. The hue, which is pale brown in this specimen, may be somewhat faded, and here and there shows indications of the plant having been originally rose coloured.

PLATE CLXXXII. Fig. IV.-1, branch and ramuli ; 2, apex of branch :- both magnified.

6. POLYSIPHONIA microcarpa, Hook. fil. et Harv.; atro-rubescens, cæspitosa, frondibus tenuissimis capillaribus membranaceis flaccidis tenacibus oligosiphoniis equalibus vix attenuatis irregulariter repetitim dichotomis, ramis ramulisque erecto-patentibus crebre divisis, articulis bistriatis e tubulis quatuor formatis, ramorum majorum diametro multiplo, minorum triplo quadruplove, ramulorum sesqui duplove longioribus, ceramidiis pusillis ovatis breve pedicellatis. Nobis in Lond. Journ. Bot. vol. iv. p. 265. (TAB. CLXXXII. Fig III.)

HAB. Hermite Island, Cape Horn; very rare.

Fil. 3-4 une. longa, capillaria, flaccida, tenacia sed non fragilia, deuse cæspitosa, basi irregulariter dichotome ramosa, ramis omnibus diametro æqualibus. *Ceramidia* minima, lateralia, elliptico-urceolata. *Color* luride ruber.

PLATE CLXXXII. Fig. III.-1, portion of branch; 2, ditto with ceramidium :- both highly magnified.

7. POLYSIPHONIA abscissa, Hook. fil. et Harv.; coccinea, frondibus circumscriptione ovatis tenuibus membranaceo-gelatinosis flaccidis tenacibus oligosiphoniis, caule primario parce diviso flexuoso ramos secundarios alternos multifidos circumscriptione obovatos emittente, ramis filiformibus minoribus alternis subdichotome divisis, ramulis fastigiatis (quasi abscissis) fibrilliferis, articulis ramorum diametro quadruplo v. sextuplo, ramulorum duplo triplove longioribus bistriatis, ceramidüs pusillis ovatis breviter pedicellatis. Nobis in Lond. Journ. Bot. vol. iv. p. 266. (TAB. CLXXXIII. Fig. II.)

HAB. Hermite Island, Cape Horn ; dredged up in about six fathom water.

Frons seu ramus primarius 3-4 unc. longus, filiformis v. capillaceus, flexuosus, alterne ramosus, ramis gradatim brevioribus, hine circumscriptio frondis totius ovata evadit. Rami inferne nudiusculi, superne fastigiatim ramulosi; ramulis ultimis tenuissimis, confertis. Caulis tubuli sub quatuor. Color roseus v. cocciueus.

A beautiful species, of which the only specimens were procured with the dredge in St. Martin's Cove. The fastigiate branching of the ramuli and the colour arc, of themselves, sufficient to distinguish this from the *P. microcarpa*, to which it is most nearly related. The principal stem is very conspicuous though slender, and the branching regular and tolerably uniform.

PLATE CLXXXIII. Fig. II.—1, plant of the natural size; 2, branch and ramuli; 3, portion of ramulus; 4, portion of ramus; 5, ditto, with ceramidium :—all highly magnified.

8. POLYSIPHONIA *flabelliformis*, Hook. fil et Harv.; pusilla, setacea, badia, rigidula, fronde brevi basi simplici stipitiformi apice flabellatim ramosa, ramis irregulariter dichotomis multifidis apice subfastigiatis, ramulis ultimis erectis longe nudis, axillis angustis, articulis multistriatis inferioribus diametro multiplo superioribus sesquilongioribus. Nobis *in Lond. Journ. Bot.* vol. iv. p. 266. (TAB. CLXXXIII. Fig. I.)

HAB. Crozet Islands; on Macrocystis pyrifera.

Frons uncialis, solitaria, rigida, crassitudine setæ porcinæ, inferne simplex, superne distiche flabellatim ramosa, crassiuscula, circumscriptione orbiculari. Rami multifidi, irregulariter dichotomi, fastigiati, ramulis erectis. Articuli rami valde clongati, strüs numerosis notati. Color badius, vix rufescens :---chartæ vix adhæret.

Only one specimen of this very distinct species was procured, from a piece of *Macrocystis* floating off the Crozet's Islands, of which group the present, the *Callithamnion Ptilota*, nob., and *Ballia Brunonis*, are the only known vegetable productions.

PLATE CLXXXIII. Fig. III.—1, plant of the natural size; 2, branch and ramuli; 3, portion of ramulus, and 4, of stcm :—highly magnified.

9. POLYSIPHONIA (Heterosiphonia) Berkeleyi; Het. Berkeleyi, et Pol. punicea, Mont. Voy. au Pole Sud, Bot. Crypt. p. 128. t. 5. f. 3. Fl. Antarct. Pt. 1. p. 182.

Var. β. Davisii; robustior, caule primario regulariter ramoso, ramis erecto-patentibus sub-bipinnatim ramulosis, ultimis erectioribus densioribus parciusque divisis. P. Davisii, nobis in Lond. Journ. Bot. vol. i. p. 267.

HAB. Hermite Island, Cape Horn; Falkland Islands, and Kerguelen's Land; abundant. Var. Davisii, Hermite Island; rare.

The somewhat different habit, more regular primary ramification, and more erect, denser, and less divided ramuli, had induced us to separate the var. β . from the original *P. Berkeleyi*: an opinion we have now abandoned, after a careful examination of very many specimens; amongst which, forms connecting the two may be found.

Though not included by Montagne under his genus *Heterosiphonia*, the structure of the tubes forming the frond of *P. punicea* is the same with that of *Heterosiphonia Berkeleyi*, of which we have examined an authentic specimen, communicated by our friend the Rev. M. J. Berkeley, and differing in no respect from *P. punicea*. We scarcely, however, think that the varying diameter of the tubes in the genus *Polysiphonia* authorizes a division of the genus; for, in some species, as the present, the increased size of two of the tubes, though conspicuous under favourable circumstances, affords but an obscure character; and in some species the difference of diameter is trifling.

24. RHODOMELA, Ag.

1. RHODOMELA *patula*, Hook. fil. et Harv.; fronde cylindracea brunnea cellulis irregularibus notata vage bipinnatim ramosa, ramis alternis elongatis horizontalibus suberecto-patentibusve minoribus elongatis patentibus subsimplicibus alternatis nudis. Nobis *in Lond. Journ. Bot.* vol. iv. p. 264. (TAB. CLXXXIII. Fig. IV.)

HAB. Falkland Islands; Port William and Berkeley Sound, rare.

Frons 4-6 unc. longa, basi diametro $\frac{1}{2}$ lin., vage et patentim ramosa. Caulis primarius subsimplex, ramos alternos, patentes, elongatos emittens. Rami laxe ramulosi, ultimi breviusculi, e tubulis 4 magnis circa cavitatem centralem dispositis extus strato cellulornm confertorum circumdatis conflati. Substantia membranacea. Color luride brunneus v. fuliginosus:---chartæ adheret.

Similar to the following, and, perhaps, not specifically distinct : it differs in the ramification ; and in the absence of the very numerous short ultimate ramuli so copiously scattered over the branches of *R. Gaimardi*.

PLATE CLXXXIII.—Fig. IV.—1, portion of plant of the natural size; 2, portion of stem and ramulus :—magnified.

2. RHODOMELA Gaimardi, Ag.; fronde cylindracea flabellatim ramosissima, stipite simplici filiformi, ramis primariis divaricatis, secundariis patentibus bipinnatim multifidis segmentis alternis, ramulis brevibus setaceis simplicibus furcatis quadrifidisve sæpe secundis per totam frondem sparsis. Nobis in Lond. Journ. Bot. vol. iv. p. 264. Agardh, Spec. Alg. vol. iv. p. 380. (non Mont. in Voy. au Pole Sud). (TAB. CLXXXIV.)

HAB. Hermite Island, Cape Horn, and in Berkeley Sound, Falkland Islands; not uncommon.

Frons 4-6 unc. longa, crassitudine setæ porcinæ, basi simplex, superne in ramos 3-4 primarios flabellatim divisa. Rami primarii subdichotomi v. irregulares, divaricati, repctitim bifarie ramulosi; rami secundarii tertiariique elongati, simplicinsculi, filiformes, ramulis brevibus ornati. Ramuli 2-3 lin. longi, sæpissime secundi, tenuissimi. Structura ut in R. patula. Color huridus.

This, which we doubtfully referred in the London Jonrnal of Botany to the R. Gaimardi, Ag., appears to us decidedly the plant of Agardh; and our friend, Dr. Montagne, has kindly furnished us with a specimen of the Auckland Island species, to which he had applied this name, and which belongs to another plant. The R. Gaimardi of Dr. Montagne is assuredly our *Polysiphonia botryocarpa*, (Pt. 1. p. 181.) and has very much the appearance of a *Rhodomela*. The specimens, from which the above description is taken, were gathered in the same locality as that from whence the typical plant of Agardh was brought by Gaudichaud; and they agree with

5т

the description of that author in every particular : except that the compression attributed to the frond is certainly not a character of our specimen, and most probably originated in that of Gaudichaud from bad drying.

PLATE CLXXXIV.—Two states of R. Gaimardi, of the natural size. Fig. 1, ramuli and stichidia; fig. 2, tetraspores; fig. 3, section of stem:—magnified.

3. RHODOMELA? comosa, Hook. fil. et Harv.; ramosissima, atro-rubescens, caule cylindraceo frondem percurrente ramis crebris alternis ornato, ramis cylindraceis elongatis pluries alterne divisis erecto-patentibus sensim utrinque attenuatis, ramulis ultimis setaceis acutis abbreviatis vagis, capsulis ovatis breve pedicellatis. Nobis *in Lond. Journ. Bot.* vol. iv. p. 263. Harv. Ner. Aust. t. xi. (TAB. CLXXXV.)

Var. B. fibrillifera; fronde tenuiori laxius ramosa, apicibus fibrilliferis.

HAB. Berkeley Sound, Falkland Islands; both varieties abundant.

Caulis cylindraceus, 6-9 unc. longus, $1-1\frac{1}{2}$ lin. diametro (in var. β gracilis) indivisus v. e basi in ramos primarios 3-4-divisus. Rami primarii secundariis perplurimis aucti, secundarii ramulis brevibus setaceis ornati, ultimis in var. β . fibrilliferis : omnes e tubulis septem circa axin centralem articulatam dispositis ct strato externo cellulorum densorum circumdatis conflati. Ceramidia numerosa, secus ramulos ultimos tertiariosque disposita, parva, ovata, breviter pedicellata. Substantia flaccida, opaca, primo visu inarticulata, sed vere articulata. Color huride rufobrunneus :---chartæ arcte adhæret.

A very much branched species, variable in size and in the density of the ramification. In old specimens the stem becomes considerably incrassated and constricted at irregular intervals. Being unacquainted with the secondary fructification, we doubtfully refer this plant to *Rhodomela*: it may belong to *Dasya*.

PLATE CLXXXV.—Two vars. of *R. ? comosa*, of the natural size. *Fig.* 1 *a*, branch and ramuli of var. *a*; *fig.* 2 *a*, section of ditto; *fig.* 3 *a*, tissue of ditto; *fig.* 1 *b*, portion of branch and ramulus of var. β . with ceramidia; *fig.* 2 *b*, fibrilliferous apex of ditto:—highly *magnified*.

25. MELOBESIA, Lamx.

1. MELOBESIA verrucata, Lamx. Polyp. flexibles, p. 315. Decaisne in Ann. Sc. Nat. Ser. ii. vol. xviii. p. 126.

Var. *Antarctica*; fronde circumscriptione orbiculari lobata medio adnata margine integerrima libera superficie lævi lineis concentricis undulata, ceramidiis depresso-hemisphæricis :—an species distincta?.

HAB. Var. B. Hermite Island, Cape Horn; the Falkland Islands, and Kerguelen's Land; encrusting shells, and the stems of Alga, particularly of Ballia Brunonis.

The *M. verrucata* is a native of the Atlantic Ocean and Mediterranean Sea; the var. β . of the Antarctic Ocean; of Lord Auckland's Group, New Zealand and Tasmania. The ceramidia are intermediate in size between those of *M. verrucata* and *M. pustulata*, Lamx. We have little doubt of this being a new species; but the materials for its determination are wanting.

26. DASYA, Ag.

1. DASYA *pectinata*, Hook. fil. et Harv.; setacea, rigida, purpurea, fronde basi nuda superne distiche decomposito-pinnata, ramis articulatis tri-striatis pectinato-pinnatis, ramulis (v. pinnulis) simplicibus alternis brevibus subulatis articulatis monosiphoniis, articulis diametro sesquilongioribus, ceramidiis urceolatis pedicellatis. Polysiphonia pectinata, *nobis in Lond. Journ. Bot.* v. iv. p. 267.

HAB. Hermite Island, Cape Horn; dredged up from about six fathom water, and on rocks at lowwater mark, very rare; Falkland Islands, Mrs. Capt. Sulivan.

Species pulcherrima, habitu Bonnemaisoniæ asparagoidis. Frons 2-4 unc. longa, circumscriptione late ovata, 2 unc. lata, rigida, distiche ramosa, v. ramosissima, ramis setaceis decomposito-pinnatis. Caulis primarius simpliciusculus, basi inarticulatus, superne articulatus, tri-striatus, compressus v. angulatus. Rami minores ramulis alternis ornati, omnes breves, subulati, e singulo serie cellularum formati, hine monosiphonii. Articuli omues breves, caulini e tubis quatuor inæqualibus (quorum 2 lateralibus latioribus,) circa cavitatem ceutralem dispositis conflati; articuli ramulorum Callithamnio forma et structura simillimi. Ceramidia secus ramulos disposita. Color pulchre purpureo-roseus.

A very beautiful and rare species : distinct from any of its European and exotic congeners that have been described. Mrs. Sulivan's specimens are much finer and more branched than those from Cape Horn.

27. STICTOSIPHONIA, Harv.

Frons purpurea, filiformis, cylindracea, ramosa, tubulosa, extus stictis quadratis notata, intus diaphragmatibus scptata. Peripheria e cellulis quadratis tubum centralem cavum radiatim cingentibus formata. Ceramidia ; Stichidia lanceolata, ramulos terminantia, tetrasporas pluriseriatas foventia.—Algæ pusillæ, cæspitosæ, e filis repentibus ortæ, rupes marinas Antillauas, Austro-Atlanticas, Antarcticasque vix demersas v. ad limitem pleni maris æstus sitas incolentes.— Genus Bostrychiæ, Mout. valde affine.

A very natural little group: composed of a few species, which occupy the same position with regard to the high-water mark in the Southern Ocean, that *Lichina* and *Catenella* do in the Northern. As a genus it differs from *Bostrychia*, Mont., only in the more simple internal structure of the frond, and broad, apparently septate, tubes, surrounded by only one row of cells occupying the centre of the frond : in habit and other respects they are so closely allied, that it is doubtful whether *Stictosiphonia* should not rather be regarded as a subgenus of *Bostrychia*. The structure of the frond is very similar to that of *Polysiphonia*, differing chiefly in the cellules of the periphery being very short; whilst those constituting the axis are lengthened.

1. STICTOSIPHONIA *Hookeri*, Harv.; caulibus indivisis curvatis apice involutis, ramis lateralibus abbreviatis alternis subquadrifariis crecto-patentibus, inferioribus subulatis simplicibus furcatisve, superioribus alterne multifidis, ramulis subulatis acutis erectis, axillis acutis, stictis subtriseriatis, stichidiis lanceolatis acutis ramulos minores terminantibus. Bostrychia Hookeri, Harvey *in Lond. Journ. Bot.* vol. iv. p. 269. (TAB. CLXXXVI. Fig. II.)

HAB. Hermite Island, Cape Horu; and the Falkland Islands: on rocks close to high-water mark; abundant.

Frons $1-l\frac{1}{2}$ unc. longa, dense cæspitosa, rigida, atro-purpurea. *Caulis* plerumque simplex, per totam longitudinem ramulis brevibus lateralibus ornatus. *Rami* nunc omnes 1 lin. longi et indivisi v. superiores clongati 2-4 lin. longi, repetitim ramosi. *Ramuli* ultimi subulati, erecti erecto-patentesve. *Rami ramulique* omnes apicibus plerumque arcte involutis :—chartæ laxe adhærct.

A beantiful little plant, marked all over, under the microscope, with three rows of dark purple dot-like cells.

PLATE CLXXXVI. Fig. II.—1, plant of the natural size; 2, stem, &c.; 3, ramulus and stichidium; 4, portion of stem; 5, longitudinal and 6, horizontal section of ditto; 7, tetraspores :--all magnified.

2. STICTOSIPHONIA fastigiata, Hook. fil. et Harv.; caulibus fastigiatis multifidis apicibus involutis,

ramis æquilongis curvatis, ramulis alternis subulatis furcatis v. alterne multifidis, axillis acutis, stictis 3-4- v. pluriseriatis. Bostrychia fastigiata, nobis in Lond. Journ. Bot. vol. iv. p. 269.

HAB. Hermite Island, Cape Horn; on stones near high-water mark.

Pusilla, dense fastigiata. Frondes $\frac{1}{2}$ unc. longæ, e basi in ramos plurimos primarios divisæ, rubro-purpureæ. Caulis brevissimus. Rami elongati, curvati, apicibus arete incurvis, ramulis simplicibus multifidisve ornati :—ehartæ laxe adhæret.

Possibly only a variety of the preceding; from which, however, it differs conspicuously in the very abbreviated stem, the consequently longer, more divided branches and the duller colour.

3. STICTOSIPHONIA vaga, Hook. fil. et Harv.; caulibus flexuosis vage dichotome ramosis, ramis paucis nudis simplicibus filiformibus subcapillaribus arcnatis medio incrassatis apicibus incurvis, ramulis nullis, axillis patentibus, stictis minutis multiseriatis, stichidiis longissime pedunculatis lanccolatis acutis. Bostrychia vaga, nobis in Lond. Journ. Bot. vol. iv. p. 270. (TAB. CLXXXVI. Fig. I.)

HAB. Christmas Harbour, Kcrguelen's Land; on rocks and stones above high-water mark, and in damp places at a considerable distance from the sea; abundant.

Dense cæspitosa, filis intertextis quasi crinita. Frondes $\frac{1}{2}-1$ unc. longæ, flexuosæ, irregulariter ramosæ, capillarcs. Stictæ parvæ, 6-8-seriatæ. Substantia rigida. Color luride purpureus :—chartæ laxe adhæret.

A remarkably distinct little species, of very simple structure. It is abundant in Kerguelen's Land, sometimes inhabiting places some hundreds of feet above the sea, but probably always within reach of the spray.

PLATE CLXXXVI. Fig. I.—Plant of the natural size; 2, rami of ditto; 3, portion of ditto; 4, incrassated ramulus; 5, ramulus and stichidium; 6. tetraspores :—all magnified.

28. LAUREN IA, Lamx.

1. LAURENCIA pinnatifida, Lamx. Var. y. angustata, Hook.; Fl. Antarct. Pt. 1. p. 184.

HAB. Berkeley Sound, Falkland Islands; abundant on the beach.

One of the most widely dispersed of the *Algæ*, inhabiting the shores of Europe from Norway to the Mediterranean; the Canary Islands; west coast of Africa, and Cape of Good Hope; the Peninsula of India; Australia and New Zealand; the Pacific Islands, and both coasts of North and South America. This very extended range has, however, its limits; the plant is neither found so far north as Iceland in the Arctic Sea, nor in the south is it kuown to inhabit Cape Horu or Kerguelen's Land.

29. DELISEA, Mont.

1. DELISEA pulchra, Mont. in Ann. Sc. Nat. Ser. iii. vol. i. p. 158. Bowiesia pulchra, Grev. Synops. Alg. p. 57. Bonnemaisonia elegans, Endl. Suppl. vol. iii. p. 44. Calocladia pulchra, Grev. Herb. Sphærococcus flaccidus, Suhr. (fid. Mont.)

HAB. Christmas Harbour, Kerguelen's Land; common.

Magnificent specimens of this noble *Alga* were collected by the Antarctic Expedition, though only in Kerguelen's Laud. The previously assigned habitat for the species is New Holland or Tasmania; but we have seen no other specimens than Mr. Fraser's original one, labelled as from that quarter of the world. It therefore appears to us probable, that the specimen sent by Mr. Fraser, may have been collected in Me'Quarrie's Island; whence other Antarctic plants were brought to that gentleman in Sydney, some of which have since found their way into our Herbaria as of Australian origin.

30. IRIDÆA, Bory.

1. IRIDÆA Radula, Bory; Fl. Antarct. Pt. 1. p. 188.

HAB. Hermite Island, Cape Horn; Falkland Islands and Kerguelen's Land, very abundant. Cockburn Island; at the limits of southern vegetation, on the beach, rare and bleached.

So abundant are the *Irideæ* in the South Polar Oceau, and so variable in their form and texture, that we can searcely hope to arrive at any accurate knowledge of the species until they shall have been studied in a living state; and then it is not improbable that the genus will be considerably reduced; and one or two of the more common species be found to assume forms as dissimilar as those of our *Laurencia pinnatifida*.

There exist in the Hookerian Herbarium, authentic specimens of the *Fucus bracteatus* of Gmelin, as figured in Turner's 'Historia,' collected both at the Cape of Good Hope and in North West America, by Mr. Menzies. These are (as is generally the case with the specimens of the larger *Fuci*, preserved in our Herbaria) smaller and of that lanceolate form which other *Irideæ* present in a young state. Their texture is very thick, densely cartilaginous, opaque; and covered with tubereles which fall away, leaving a cribriform frond both when immature and older. This great density is a very remarkable character, and observable in the plant here referred to that species, which, when full grown, becomes broadly ovate, or orbicular, and eordate or rounded, or narrowed at the base; with the lamina more or less and variously divided, sometimes three feet broad, or upwards. The largest specimens we have never seen attached, though they are abundant, washed up on the beach, and probably attain their great size on the outer rocks.

Since the publication of the first part of this work, we have, through Dr. Montagne's kindness, had the opportunity of inspecting the *I. laminarioides*, Bory, of Lord Auckland's Group : specimens of which are in our Herbarium from the same island; but which we had previously regarded as a more delicate state of *I. Radula*. Even what we consider the true *I. Radula* of Lord Auckland's Group and Kerguelen's Land, is not so dense in the frond as the specimens of the Falkland Islands and Cape of Good Hope are. Both this and the following species have the surface frequently covered with granules, tubercles or pedicellate pear-shaped organs; or in the young state with elongated fleshy bodies similar to those of the *I. stiriata*, Bory. The *I. stiriata*, according to the descriptions, may belong to a state of this, or the following, or many other forms of the genus: it is, however, a narrower, smaller species, with a much more dense frond than even *I. Radula*.

2. IRIDÆA cordata, Bory, in Duperrey Voy. Bot. p. 104; et I. micans, p. 110. t. 13 et 13 lis. Halymenia cordata, Agardh, Sp. Alg. p. 201. Fucus cordatus, Turner Hist. Fuc. t. 116.

Var. *β. ciliolata*; stipite brevi cartilagineo euneato ciliato-dentato mox in frondem simplicem ovatolanceolatam desinente, fronde latissima basi euneata v. eordata apice obtusa v. acuta v. emarginato-bifida membranacea rubra plana nitente lævi margine vix undulata. Nobis *in Lond. Journ. Bot.* vol. iv. p. 263.

Var. γ . dichotoma; stipite brevi mox cuneato furcato v. pluries dichotomo sensim in frondem late cuneatam obovatamve desinente, segmentis integris vel divisis margine dentatis lobatis proliferisve.

HAB. Hermite Island, Cape Horn, and the Falkland Islands; both varieties very abundant.

This species, when fresh, well deserves the brilliant description of its beautiful tints, given by M. Bory on the anthority of Admiral D'Urville and M. Gaudichaud. It is one of the most common Alga of the southern extremity of America and the Falklands. In its younger state, the fronds are obovate or spathulate, like those of *I. laminarioides*, figured by Bory, and soon expand into laminæ, variously modified, according to situation and exposure, with relation to the force of the sea, the nature of the bottom, the currents, depth, and protection afforded by other Alga; for no two fronds of a similar shape are usually to be found within a few yards. Indeed, I question whether *I. micans* be more than a membranous form of *I. Radula*: the former always preferring the quieter harbours, where its fronds are sometimes as thin as those of a *Delesseria*, quite unfitted to withstand the rough seas of the outer coasts, which wash the almost uninjured fronds of the *I. Radula* ashore in broad sheets, as large and as red as an ordinary pocket-handkerchief.

Though sometimes almost equally thin, the substance of the *I. micans* is never so membranous as that of a *Delesseria*. The colour, though not so bright a rose, or so delicate when the plant is dried, is, when seen in the living state, much more varied and more beautiful. The texture is such that the slightest motion of the water causes the frond to undulate throughout from the base upwards without falling into folds: each portion of the surface, when presented at a certain angle to the eye, reflecting back the most brilliant metallic tints of azure, steelblue, pink, and purple. A more beautiful object in the water is not to be found in the whole order of Alga than this, when seen from a boat in calm weather and sunshine; though it is seldom that such opportunities occur in the latitudes it inhabits. I have not been able to detect any striæ on the surface of the frond, which is formed of cells so densely packed that they coalesce into a homogeneous cartilaginous tissue.

We have no hesitation in pronouncing this as identical specifically with the *I. cordata* of the Banks of Newfoundland and the Cape of Good Hope; of which species there is an excellent figure in the 'Historia Fucorum', coinciding with that of Bory in Duperrey's Voyage. The descriptions, both of Agardh and Turner, particularly mention the iridescence of their specimens. The only differential characters noted by Bory, who justly indicates the close affinity of *I. micans* with *I. cordata*, are the slight discrepancy in the bluntness of the apices of the fronds and depth of the lobes at the cordate base. We are, however, well assured that such characters are all too slight; for we could not, either at the Cape of Good Hope or the Falkland Islands, distinguish between the forms of this *Iridæa* with a cordate and those with a cuneate base to the frond. We are, however, far from asserting that there may not be from the two last-named localities two species here confounded (one of which, the *I. micans* of Borv, is the same with the *F. cordatus* of Turner), though we strongly incline to the opposite opinion.

31. PHYLLOPHORA, Grev.

1. PHYLLOPHORA cuneifolia, Hook. fil. et Harv.; fronde stipitata basi ramosa lato-cuneata prolifera integra emarginata v. biloba e margine disco v. apice frondes consimiles emittente.

HAB. Port William and St. Salvador Bay, Falkland Islands; Christmas Harbour, Kerguelen's Land; rare.

Frondes omnes stipitatæ. Stipes compressus interdum subplanus, ima basi plerumque angustissima, sensim in lamiuam latam cuneatam deltoideamve dilatatus, basi divaricatim ramosus, bis, ter pluriesve divisus. Frondes primariæ 1-2 unc. longæ, $1-1\frac{1}{2}$ latæ; apice latiore late rotuudato, emarginato, retuso v. bilobo; segmentis rotundatis, rarius erosis; secundariæ primariis omnino similes sed colore pallidiores et basi simplices, sæpe frondes tertiarias emitteutes, hinc planta vetusta catenatim ramosa evadit. Fructus — ? Substantia tenuiter cartilaginea, subcornea, basi opaca. Color ut P. Brodiæi.—Chartæ vix adhæret.

Certainly distinct from *P. obtusa*, the only one of the genus hitherto described as a native of the southern temperate hemisphere, but perhaps not equally so from *P. Brodiæi*. Still our specimens are very different from the ordinary British form of that plant, in the much shorter stipes, and larger broader frond, which is much less lobed and the lobes are not so narrow or elongated, or separated by so deep a sinus.

2. PHYLLOPHORA obtusa, Grev. Fl. Antarct. Pt. 1. p. 187.

HAB. Hermite Island, Cape Horn; dredged up from five fathom water, very rare.

The specimens of this species are sufficiently characteristic, though few in number. It is also a native of the Cape of Good Hope and Lord Auckland's Group.

FLORA ANTARCTICA.

32. NOTHOGENIA, Mont.

1. NOTHOGENIA variolosa, Mont. Fl. Antarct. Pt. 1. p. 188.

HAB. Hermite Island, Cape Horn; the Falkland Islands; and Christmas Harbour, Kerguelen's Land; on rocks, very abundant.

An exceedingly variable plant in size and in the breadth of its fronds, simulating in the high southern latitudes the *Chondrus crispus*, as far as locality and abundance are concerned. The southern species representing our *Chondrus crispus* is the *C. tuberculatus* in Lord Auckland's Group, (where the *Nothogenia* also abounds,) and at the Cape of Good Hope the *C. dilatatus*.

33. DUMONTIA, Lamx.

1. DUMONTIA filiformis, Grev. Fl. Antarct. Pt. 1. p. 189.

HAB. Berkeley Sound, Falkland Islands; rare.

Apparently identical with the European plant, which ranges from the Mediterranean to the British coasts.

34. GIGARTINA, Lamx.

1. GIGARTINA plicata, Grev. Alg. Brit. p. 15. Fucus plicatus, Engl. Bot. t. 1089.

HAB. Cape Pembroke, Falkland Islauds; Christmas Harbour, Kerguelen's Land; abundant.

These examples so entirely accord with others of British growth, that it is unnecessary to separate them spefically. No specimens considered by any systematic botanist to belong to this *Gigartina* have been found between the latitudes of the south of Europe and Kerguelen's Land, except (according to Montagne) at Callao : yet the genus, under one or other of its Protean aspects, abounds throughout all tropical and temperate seas.

35. PTILOTA, Ag.

1. PTILOTA *Harveyi*, Hook. fil.; caule compresso cartilagineo inarticulato anguste lineari furcato inordinateve ramosissimo, ramis distichis pinnatim decomposito-ramosis majoribus minoribusque pectinatim pinnulatis costa articulata percursis, pinnulis creberrimis simplicibus articulatis monosiplioniis abbreviatis subulatis oppositis, pinnularum articulis quadratis, favellis in ramulos terminalibus ramellis pinnatis involucratis, tetrasporis ad apices pinnularum aggregatis nudis breve pedicellatis. Hook. fil. *in Lond. Journ. Bot.* vol. iv. p. 271. (TAB. CLXXXVII.)

Var. β . pinnulis subdistantibus.

HAB. Hermite Island, Cape Horn, and on the outer coasts of the Falkland Islands; abundant.

Species pulcherrima, prima visu *P. plumosæ* referenda, sed distinctissima. *Frons* 8 unc. ad pedalem, e ramis patulis ejusdem latitudinis. *Stipes* gracilis, $\frac{1}{2}$ lin. diametro, et per totam frondem æquilatus, irregulariter furcatim v. dichotome v. sub-pinnatim ramosissimus. *Rami* minores majoresque (juniores præcipue) ramulis creberrimis articulatis 1 lin. longis pulcherrime pectinati. *Ramuli* simplices, serie unica cellularum quadratarum endochromate roseo repletarum constantes, ramis *Callithamnio* subsimiles.

This lovely plant is the Cape Horn and Falkland Island representative of the Boreal and Arctic *P. sericea*, Harv. (*P. elegans*, Kutz., *Fucus sericeus*, Gmel.) and of the Auckland Island *P. formosissima*, (t. LXXVII.) From

the former of these it differs in being larger, more rigid, and having ramuli of much greater diameter, so that under the microscope it is impossible to confound them. Its Cape of Good Hope representative, and indeed, very near ally, is the *P. setigera*, Harv. (Nereis Australis.)

PLATE CLXXXVII.—Fig. 1, branch and ramuli; fig. 2, portion of a ramulus; fig. 3, another ramulus; fig. 4, favella; fig. 5, spores from ditto; fig. 6, tetraspores:—all magnified.

36. CERAMIUM, Adans.

1. CERAMIUM rubrum, Ag. Fl. Antarct. Pt. 1. p. 191.

HAB. Hermite Island, Cape Horn; Falkland Islands, and Christmas Harbour, Kerguelen's Land; very abundant.

These two *Ceramia* (*rubrum* and *diaphanum*) are very widely distributed throughout the temperate regions of both hemispheres : they are also found on the shores of Peru and Brazil.

2. CERAMIUM diaphanum, Ag. Fl. Antarct. Pt. 1. p. 191.

HAB. Hermite Island, Cape Horn; Falkland Islands; and Christmas Harbour, Kerguelen's Land; abundant.

37. GRIFFITHSIA, Ag.

1. GRIFFITHSIA Antarctica, Hook. fil. et Harv.; filis cæspitosis dichotome ramosis flaccidis, axillis inferioribus patentibus, superioribus acutis, ramis elongatis ramulisque nudis ad nodos constrictis, articulis cylindraceis superne paulo incrassatis, ramorum diametro sextuplo, ramulorum subtriplo longioribus;—fructificatio deest.

HAB. Hermite Island, Cape Horn, and the Falkland Islands; on rocks.

Fila sub 3 unc. longa, fastigiata, parce ramosa, ramis elongatis, distanter ramulosis, ramulis brevibus. Color roseus :----chartæ arcte adhæret.

Allied to the G. secundiflora, J. Ag., but smaller in all its parts.

2. GRIFFITHSIA corallina, Ag. Conf. corallina, Engl. Bot. t. 1815.

HAB. Hermite Island, Cape Horn; dredged up in about seven fathom water.

A solitary barren specimen, resembling the British G. corallina, which is also a native of the Mediterranean Sea, Canary Islands, and Cape of Good Hope.

3. GRIFFITHSIA equisetifolia, Ag. Conf. equisetifolia, Engl. Bot. t. 1479.

HAB. Falkland Islands; (Agardh.)

38. BALLIA, Harv.

1. BALLIA Brunonis, Harv. Fl. Antarct. Pt. 1. p. 190.

Var. B. Hombroniana, Fl. Antarct. l. c.

HAB. Hermite Island, Cape Horn; Falkland Islands; Christmas Harour, Kerguelen's Land, and the Crozet Islands; most abundant; always (?) parasitical.

One of the handsomest, and certainly the most common and widely distributed of the Antarctic *Floridea*: its northern limit in the New World is Patagonia, and the Bay of Islands in New Zealand in the Old. With regard to the adoption of the trivial appellation of "*Callitricha*," which Agardh proposed for this species (under *Sphacelaria*), it was waived in compliance with the wish of the first discover of the plant, whose name it now bears.

39. CALLITHAMNION, Lyngb.

1. CALLITHAMNION Plumula, Agardh, Sp. Alg. vol. ii. p. 159.

HAB. Hermite Island, Cape Horn; dredged up from about seven fathom water; very rare.

Decidedly the same as the European and North American plant.

2. CALLITHAMNION *simile*, Hook. fil. et Harv.; fronde subsolitaria rigidiuscula ramosissima, ramis alternis v. subdichotomis articulatis enerviis, ramulis brevissimis oppositis distichis crassis sursum pectinatis e quoque ramorum articulo horizontaliter porrectis, pinnulis robustis simplicibus ramosisve, articulis ramorum diametro sesqui-duplo longioribus, ramulorum diametrum subæquantibus.

HAB. Christmas Harbour, Kerguelen's Land; rare.

Frons 2-5 unc. longa, gracilis, rigidiuscula, repetitim distiche ramosa; ramis omnibus articulatis, æquilatis. Ramuli $\frac{1}{2}$ lin. longi, distiche oppositi, e medio articuli cujusvis per totam frondis longitudinem orti, robusti, subacuti, horizontaliter patentes, secus marginem superiorem dispositi, ramulis secundariis obsiti. Color fusco-ruber.

C. Plumulæ simillimum, sed rigidiusculum, ramis latioribus, ramulis robustioribus articulisque brevioribus.

This so closely resembles the *C. Plumula*, that it is difficult by mere words to discriminate them; yet, on comparing them under the microscope, they are obviously distinct. *C. simile* is a much coarser and more rigid plant, with the ramuli more robust in proportion to the diameter of the articulation they spring from, and the articulations themselves are shorter. Again, from the circumstance of the true *C. Plumula* occurring at Cape Horn, where this, (the only Kergueleu's Land species), does not appear, we incline to regard the present as a representative species rather than a variety.

3. CALLITHAMNION *Ptilota*, Hook. fil. et Harv.; parvum, rigidulum, setaceum, fronde pinnatim ramosissima, ramis vix distichis venoso-striatis subopacis, secundariis opposite pinnulatis, pinnulis simplicibus patentibus subulatis e quoque ramorum articulo ortis, articulis diametro duplo longioribus. Nobis *in Lond*. *Journ. Bot.* vol. iv. p. 272. (TAB. CLXXXIX. Fig. I.)

HAB. Crozet Islands; on a floating mass of *Macrocystis pyrifera*.

Frons 1-2 unc. longa. Caulis crassitudine setæ cquinæ, repetitim pinnatim ramosus; ramis suboppositis patentibus demum deflexis, inferioribus subuncialibus; secundariis breviusculis pinnatis, pinnis plurimis densis, oppositis, simplicibus, subulatis, e omni articulo rami ortis. Color fusco-ruber:—chartæ vix adhæret.

Only oue specimen of this very distinct little species was found : it grew on a piece of floating sea-weed, picked up at a considerable distance from the shore.

PLATE CLXXXIX. Fig. I.-1, plant of the natural size; 2, ramuli; 3, ditto with sphærospores :--all magnified.

4. CALLITHAMNION *ternifolium*, Hook. fil. et Harv.; perpusillum, vage dichotome ramosum, ramis pellucide articulatis, ramulis sæpissime ternis e omni ramorum articulo ortis erecto-patentibus brevibus gracilibus simplicibus, articulis ramorum diametro 4–5-plo ramulorum subduplo longioribus, favellis magnis bilobis ramos terminantibus. Nobis *in Lond. Journ. Bot.* vol. iv. p. 272. (TAB. CLXXXIX. Fig. II.)

HAB. Hermite Island, Cape Horn; dredged up from about eight fathom water; parasitic on other Algo.

Species perpusilla, parasitica, cæspitosa, sub $\frac{1}{2}$ unc. longa, vage ramosa, rosea, flaccida et membranacea. *Ramuli* plerumque e quoque articulo terni, raro bini quaternive, graciles, breves, simplices. *Articuli* caulis ramorumque elongati. *Favellæ* magnæ.

A very small plant, of which but few specimens were obtained, and by the dredge only : they are in a good state of fruit, and probably characteristic of the species.

PLATE CLXXXIX. Fig. II.—1, plant of the natural size; 2, portion of ditto; 3, ramuli; 4, ditto, with favella; 5, tetraspores :—magnified.

5. CALLITHAMNION *flaceidum*, Hook. fil. et Harv. ; gracillimum, flaceidum, membranaceum, fronde laxe et vage decomposite ramosa, ramis primariis et secundariis oppositis alternisve distichis elongatis patentibus, ramulis ultimis brevibus simplicibus patentibus oppositis secundisve apice incurvis, articulis ramorum primariorum diametro multoties secundariorum 6–10-plo ramulorum sesquilongioribns pellucide roseis enerviis. Nobis *in Lond. Journ. Bot.* vol. iv. p. 273. (TAB. CLXXXVIII. Fig. I.)

Var. B. alternifolium; ramis ramulisque alternis secundisve rarissime paucis oppositis.

HAB. Hermite Island, Cape Horn; dredged up from about seven fathom water; abundant.

Frons 2-4 unc. longa, laxe ramosa, membrauacea, flaccida, rosea. Caules ramique gracillimi, filiformes.

A very beautiful and delicate species, remarkable for the ramification being often truly opposite, always so in var. a, two branches springing from opposite points of the same articulation. On the other hand, when, as in var. β , they are alternate or secund, it is owing to the inner ramuli on the branches becoming abortive, those along the outer edge alone being developed. There is no other difference between the two varieties. Specifically the present is most closely allied to the *C. Turneri*, but it is much larger and more branching. The colour is a very bright rose, and from the delicacy of the filaments, the plant forms a beautiful object when properly displayed upon paper. The articuli of the stem are often singularly elongated.

PLATE CLXXXVIII. Fig. I.—1, plant of the natural size; 2, ramus and ramuli; 3, apex of ramulus; 4, favella:—magnified.

6. CALLITHAMMION scoparium, Hook. fil. et Harv.; caespitosum, caulibus fastigiatis, primario crasso inarticulato fibris intertextis flexuosis stuposis vestito flabellatim ramoso, ramis primariis cauli similibus, secundariis strictis gracilibus pellucidis creberrime pinnatis bipinnatisve quadrifariis e primariorum apicibus fasciculatim ortis fastigiatis, articulis diametro duplo triplo longioribus. Nobis in Lond. Journ. Bot. vol. iv. p. 173. (TAB. CLXXXIX. Fig. III.)

Var. B. ramulosum; pinnis apice ramulis secundis ornatis.

HAB. Var. a. Berkeley Sound, Falkland Islands; on rocks. Var. B. Hermite Island, Cape Horn; rare.

Frons 2-3 unc. longa, dense fastigiata. Caules robusti, inarticulati, basi integra, fibris stuposis dense vestiti, in discum latiusculum expansi. Rami cauli subsimiles, per totam longitudinem ramulis strictis dense fastigiatis quadrifariis vestiti (ut in Sphacelaria scoparia); secundarii articulati, vage ramosi, pinnati v. dichotomi secundive; omnes erecti, ramulique plerumque appressi; apicibus obtusis v. acutis simplicibus v. ramulis brevibus pectinatis ornati. Substantia rigida. Color luride purpurcus.

A densely tufted species, with the habit of *Sphacelaria scoparia*, resembling amongst its congeners the *C. tetricum* of Britain, but abundantly different under the microscope. It has also been found in Tasmania.

PLATE CLXXXIX. Fig. III.-1, plant of the natural size; 2, ramus and ramuli; 3, apex of the latter; 4, fibres at the base of the stem :--magnified.

7. CALLITHAMNION Montagnei, Hook. fil.; fronde fruticosa ramosissima, caulibus primariis decompositis sensim alternatis crassis quadrifariis inarticulatis opacis, ramis inarticulatis striatis ramulis quadri-

fariis plumosis densissime obsitis, ramulis (seu pinnulis) brevibns pinnatis bipinuatisve articulatis pellucidis roseis, pinnulis patentibus, inferioribus simplicibus elongatis subulatis superioribus furcatis v. iterum pinnulatis, articulis diametro subduplo longioribus. C. Gaudichaudii, Ag.? Nobis *in Lond. Journ. Bot.* vol. iv. p. 274. (TAB. CLXXXVIII. Fig. II.)

Var. β . caulibus elongatis laxius ramosis basi nudis, ramulis paucioribus gelatinosis.

HAB. Hermite Island, Cape Horn; and Berkeley Sound, Falkland Islands. Var. B. Falkland Islands.

Radix scutata. Frons 2-3 unc. (in var. β . 4-5 unc.) longa, fruticulosa, ramosissima. Caulis crassiusculus, $\frac{1}{2}$ lin. fere diametro, e basi ramosus v. nudus superne præcipue in ramos undique patentes divisus. Rami primarii pluries divisi; secundarii ramulis pinnulisve parvis $1-l\frac{1}{2}$ lin. longis undique vestiti. Favellæ magnæ, 2-3-lobatæ, lobis granulis plurimis farctis. Color siccitate atro-purpureus, madore sub lente roseo-purpureus. Substantia caulis ramo-rumque cartilaginea, ramulorum tener, chartæque adhærens.

In the London Journal of Botany we referred this plant with a mark of doubt, (and erroneously as it subsequently appears) to the *Cal. Gaudichaudii* of Agardh: a Falkland Island species, with which it seemed to agree in many particulars. Our kind friend, Dr. Montagne, has, with his usual liberality, supplied us with a portion of the original specimen of *C. Gaudichaudii*, which proves to be quite distinct. Its nearest northern allies are *C. Arbuscula* and *C. Brodiæi*, between which it appears almost intermediate, having the large size and robust habit of the former, with longer and more compound pinnules, and being much stouter than *C. Brodiæi*, having more opaque stems. The var. β . may be only an advanced state, having been gathered in the same locality with var. *a.*, but three months later in the season. It chiefly differs in its more tender and gelatinous substance, and in the branches being less densely clothed with ramuli, and nearly naked at the base. Its outward appearance is very much that of *C. tetragonum*, Ag.

PLATE CLXXXVIII. Fig. II.--1, plant of the natural size ; 2, branch and ramuli ; 3, ditto with favella :--magnified.

8. CALLITHAMNION Gaudichaudii, Ag. Sp. Alg. vol. ii. p. 173.

HAB. Falkland Islands; Gaudichaud.

9. CALLITHAMMION leptocladum, Montagne in Voy. au Pole Sud, Bot. Crypt. p. 91.

HAB. Strait of Magalhaeus ; D'Urville.

40. CODIUM, Stackh.

1. CODIUM tomentosum, Stackh.; Fucus tomentosus, Engl. Bot. t. 712.

HAB. Hermite Island, Cape Horn; and the Falklaud Islands; abundaut. Kerguelen's Laud?

This curious plant is equally widely diffused in the southern as in the northern and tropical zones; and the specimens from the different localities are very similar. What we believe to have been this species was collected in Kerguelen's Land, but no specimens appear to have been preserved.

41. CLADOTHELE, Hook. fil. et Harv.

Frons cylindracea, filiformis, viridis, solida, ramosa, extus papillosa. Axis cellulosa, densa, e cellulis magnis hyalinis vacuis cellulam centralem radiatim cingentibus formata. Peripheria cellulosa, cellulis coloratis (viridibus) pluriscriatis. Utriculi papillæformes, totam superficiem vestientes.—Alga marina Falklandica, irregulariter ramosa, sordide viridis, ecorticata. 1. CLADOTHELE Decaisnei, Hook. fil. et Harv.; in Lond. Journ. Bot. vol. iv. p. 293. (TAB. CXC.)

HAB. Berkeley Sound, Falkland Islands; in the sea.

Radix fibrosa? Frondes 4-6 unc. altæ, cæspitosæ, filiformes, seta porcina crassiores, cylindraceæ, flexnosæ, plus minusve ramosæ, ramificatione valde irregulari. Rami primarii elongati, sæpe simplices, ramulis longis simplicibus sæpissime secundis curvatis v. incurvis vix attenuatis laxe donati. Substantia tenax. Color sordide viridis, siecitate cinerascens :---chartæ laxe adhæret.

A very curious plant, certainly related to *Codium*, especially to *C. simpliciusculum*, by the structure of the papillæ that cover its surface, and from which we have derived the generic name. The axis is, however, of very different structure from that of *Codium* or of any other genus of *Siphoneæ*, and more closely resembles that of *Polysiphonia*. In the specific name we wish to pay a deserved compliment to our friend M. Decaisne, who has thrown much light on the affinities of the corallinoid Algæ, especially those related to *Siphoneæ*.

PLATE CXC.—Fig. 1, plant of the natural size; 2 ramus and ramuli; 3, longitudinal, and 4, vertical section of branch; 5, cellular tissue of ditto:—magnified.

42. BRYOPSIS, Lamx.

1. BRYOPSIS plumosa, Grev. Alg. Brit. p. 187.

Var. B. Arbuscula, J. Agardh, Alg. Medit. p. 21. B. Arbuscula, Ag. Sp. Alg. p. 451.

HAB. Hermite Island, Cape Horn, and the Falkland Islands; abundant, both varieties.

The branches of the frond are narrower and pinnated nearcr to the base, with the ramuli more uniform and shorter in some of the specimens than in others.

2. BRYOPSIS Rosa, Ag. Syst. Alg. p. 179. Bory in Duperrey Voy. Bot. p. 211. t. 24. fig. 1.

HAB. Hermite Island, Cape Horn, and the Falkland Islands.

None of our specimens equal those figured by Bory in size, though they coincide in all other respects with the descriptions published by that author and Agardh. Our opinion is, that the present plant is not distinct from the *B. plumosa*, but is a large state of that very sportive species, depending probably on the temperature of the ocean it inhabits for its development. Some other species of this highly Protean genus are equally difficult to define; and we cannot but expect that a copious suite of specimens from different shores and depths will considerably diminish it.

43. VAUCHERIA, DC.

1. VAUCHERIA Dillwynii, Ag.; Grev. Alg. Brit. p. 191. t. 19. Conferva frigida, Dillwyn, t. 19.

HAB. Christmas Harbour, Kerguelen's Land; on the ground amongst the Penguin Rookeries.

The patches are very extensive and rather more glaucous than others collected near Edinburgh, with which the Antarctic specimens appear otherwise entirely to agree. The capsules are not always globose, as figured by Mr. Hassall, (Brit. Fresh-water Algæ) but often, if not more generally, horizontally elongated and gibbons; as shown in Greville's 'Algæ Brit.' (l. c.) where there is an excellent figure of this species.

2. VAUCHERIA cæspitosa, Ag.; Grev. Alg. Brit. p. 194.

HAB. Berkeley Sound, Falkland Islands; on the moist borders of fresh-water lakes, and in pendent masses from dripping rocks.

These specimens are dried very badly, so that we have not much confidence in our identification of the species.

FLORA ANTARCTICA.

44. BATRACHOSPERMUM, Roth.

1. BATRACHOSPERMUM vagum, Ag.; Harvey, Manual, p. 119. Lyngb. Hydroph. Dan. t. 44.

HAB. Hermite Island, Cape Horn; in an alpine pool.

We cannot distinguish these from British specimens; an alpine locality is common to both, the English plant having been gathered on the summit of Snowdon.

45. DRAPARNALDIA, Bory.

1. DRAPARNALDIA *pusilla*, Hook. fil. et Harv.; filis perpusillis densissime cæspitosis gelatinosis parce vage ramosis flexuosis, ramulis perpaucis brevibus apice non setigeris simplicibus, articulis coloratis luteoviridibus diametro sub-duplo longioribus. Nobis *in Lond. Journ. Bot.* vol. iv. p. 296. (TAB. CXC. Fig. II.)

HAB. Berkeley Sound, Falkland Islands; growing on the roots of Crantzia lineata, in fresh-water.

Fila sub $\frac{1}{4}$ unc. longa, radices radiculasque submersas vestientia, diametro *D. tenuis*, cui verosimiliter species affinis; differt præcipue filis rigidioribus ramulisque non setigeris.

We have referred this and the following species to the genus *Draparnaldia* with little hesitation, from their affinity with *D. tenuis*, Ag. Those naturalists, however, who know the great difficulty of examining such things in a dried state, will best understand the uncertainty which attaches to all determinations of species belonging to these tribes which are not from the fruit, or from characters of higher importance than the filamentous branches, evidently referable to known forms.

PLATE CXC. Fig. II.—1, plant of the natural size, on roots of Crantzia lineata; 2 and 3, threads; — highly magnified.

2. DRAPARNALDIA sp.?

HAB. Hermite Island, Cape Horn; in stagnant water on the hills.

The filaments of this species are infinitely more slender than those of the last, but similarly gelatinous and of the same structure.

46. CONFERVA, Ag.

1. CONFERVA clavata, Ag.? Syst. Alg. p. 99.

Var. *Darwinii*; pro genere maxima, filis 2 uncialibus e basi gradatim incrassatis, articulis ad nodos constrictis diametro paulo longioribus, inferioribus longioribus, supremis ¹/₅ unc. latis. (TAB. CXCII. Fig. I.)

HAB. Cape Tres Montes, on Sphacelaria funicularis; C. Darwin, Esq.

Of this variety we have seen but one specimen and refer it doubtfully to the *C. clavata* of the Cape of Good Hope and New Zealand, to which it is certainly very closely allied.

PLATE CXCII. Fig. I.—Plants of C. clavata, var. Darwinii, of the natural size, parasitical on Sphacelaria funicularis, Mont.

2. CONFERVA Linum, Ag.; Harv. Man. Brit. Alg. p. 128.

HAB. Christmas Harbour, Kerguelen's Land; in the sea, on rocks near high-water mark.

A widely distributed species, found from the Canary Islands, Mediterranean and Black Sea, to the coasts of

5 Y

Scotland, also on the east coast of North America, the West Indies, and on the west coast of South America. It is singular that Kerguelen's Land should be its only hitherto recorded locality in the Southern Hemisphere.

3. CONFERVA Sandvicensis, Ag.; Syst. Alg. p. 92. (TAB. CXCII. Fig. II.)

HAB. Berkeley Sound, Falkland Islands; in pools of fresh water, and hanging from wet rocks.

Massa pedalis et ultra, mollissima, pallide sed læte viridis, e filis tenuissinis arachnoideis densissime fastigiatis constans. *Fila* hyalina, flaccidissima, simplicissima, vix intertexta, aciem oculorum fugientia. *Articuli* diametro duplo longiores, subvitrei; sacculo endochromatis interno medio constricto, læte virente, pellucido.

Our specimens entirely agree with Agardh's description of a plant brought from the Sandwich Islands by M. Gaudichaud.

PLATE CXCII. Fig. II.--1, plant of the natural size; 2, portion of a thread; 3 & 4, other portions of ditto: --both very highly magnified.

4. CONFERVA angulata, Hook. fil. et Harv.; fluitans reptansve, filis simplicibus tenuissimis brevibus strictiusculis hic illic incrassatis angulatisque angulis radiculo ramulove abnormali auctis, articulis diametro 3-5-plo longioribus coloratis, endochromate siccitate contracto. Nobis *in Lond. Journ. Bot.* vol. iv. p. 295. (TAB. CNCI. Fig. II.)

HAB. Christmas Harbour, Kerguelen's Land; common in streams, pools, and lakes of fresh water.

Fila intertexta, tenuissime capillacea, in massam pallide virescentem subnitentem conferta. Articuli siccitate contracti, æquales, nunc medio incrassati, nucleati, rarissime ramum emittentes, sæpissime radicula parva uniarticulata aucti, nunc geniculatim eurvati.

Allied to the British *C. bombycina*, but readily distinguishable by its greater rigidity, angular flexures, radicles and different incrassations, which do not appear to us of the same character as those of Mr. Hassall's genus '*Vesiculifera*.' The filaments are sometimes ramified, though very rarely, and perhaps only at the very base: the branch is always at right angles to the filament.

PLATE CXCI. Fig. II.—1, plant in mass, of the natural size; 2, thread of ditto; 3, portion of ditto with branch; 4, ditto with rootlets?; 5, ditto with swollen joint :--all very highly magnified.

5. CONFERVA ambigua, Hook. fil. et Harv.; filis basi intertextis adnatis? capillaribus rigidulis nigrovirescentibus longe fluctuantibus simplicibus hic illic spurie? ramosis radicantibus, nunc processubus lateralibus anastomosantibus auctis diametro 2–3-plo longioribus opacis sacculo endochromatis repletis. Nobis in Lond. Journ. Bot. vol. iv. p. 295. (TAB. CXCI. Fig. I.)

HAB. Christmas Harbour, Kerguelen's Land; in the sea.

Fila 4-5 unc. longa, basi in stratum densum intertexta, deinde libera, elongata, massam crinitam efficientia.

PLATE CXCI. Fig. 1.-1, plant of the natural size; 2, filament from ditto :- magnified.

6. CONFERVA quadratula, Hook. fil. et Harv.; pusilla, filis tenuibus pallide viridibus flexuosis intricatis cylindraceis, articulis quadratis siccitate endochromate collapso notatis. (TAB. CXCI. Fig. IV.)

IIAB. Christmas Harbour, Kerguelen's Land; in pools and streams of fresh water; very common.

Fila simplicissima, sub $\frac{1}{2}$ unc. longa, in strato dilute viridia, implicata, erispata, diametro C. floccosæ duplo triplove superantia. Articuli longitudine diametrum æquantes, cylindracei, ad nodos non constricti, pellucidi, endochromate plerumque in massam linearem viridem collapso medio notati.

PLATE CXCI. Fig. IV.---1, plant of the natural size; 2, thread from ditto; 3, portion of ditto:--both very highly magnified.

7. CONFERVA *podagraria*, Hook. fil. et Harv. ; filis simplicibus basi intertextis breviusculis fluctuantibus flexuosis flaccidis flavo-viridibus cylindraceis, articulis opacis elongato-quadratis diametro $\frac{1}{2}$ -2-plo longioribus sacculo endochromatis repletis integumento externo sæpissime incrassato nodoso. (TAB. CXCI. Fig. III.)

HAB. Christmas Harbour, Kerguelen's Land; in streams of fresh water, attached to stones or earth.

Massæ unciam latæ, nunc latius extensæ, pallide virescentes. *Fila* flexuosa, $\frac{3}{4}$ unc. longa, laxe intertexta, opaca, e basi simplicissima, cylindracea, subæquilonga. *Articuli* cylindracei, sacculo endochromatis repleti, sæpissime, ob tegumentum externum morbo affectum, incrassati et nodosi.

A remarkably distinct little species, forming patches in the water. The threads are densely tufted, curled, and ascending, rather stout in proportion to their length, but flaccid and somewhat soft in consistence; they are generally covered at some part of their length with a thickened opaque substance, of irregular form, extending over several of the joints at once, but more or less evidently protuberant on one side of the thread. This appearance seems due to a diseased condition of the outer membrane; for the sac of endochrome is often seen to be unchanged beneath this thickening, which sometimes increases the filament to twice its usual diameter.

46. CLADOPHORA, Kütz.

1. CLADOPHORA rupestris, Linn.; Dillw. Hist. Brit. Conf. t. 23.

HAB. Christmas Harbour, Kerguelen's Land; on rocks in the sea.

These specimens are very characteristic of the northern *C. rupestris*, which inhabits all latitudes between the Arctic Circle and Mediterranean Sea on the west coast of Europe.

2. CLADOPHORA flexuosa; Dillw. Hist. Brit. Conf. t. 10.

Berkeley Sound, Falkland Islands; in the sea.

Specimens not very satisfactory, but we think referable to this species. The ramuli are secund, and the other characters of *C. flexuosa* are tolerably evident.

3. CLADOPHORA arcta; Dillw. Brit. Conf. Suppl. t. E.

Var. centralis, Conferva centralis, Lyngb. et auct.

HAB. Hermite Island, Cape Horn, and in the Falkland Islands; very abundant, in the sea.

Decidedly the European plant of the name, which is a native of the German and North Atlantic Ocean.

4. CLADOPHORA riparia, Roth; Engl. Bot. t. 2100.

HAB. Christmas Harbour, Kerguelen's Land; on rocks near high-water mark.

A native also of the German Ocean, the North Sea, and West Indian Islands.

5. CLADOPHORA *Falklandica*, Hook. fil. et Harv.; filis densissime cæspitosis flaccidis flexuosis intricate ramosissimis læte virescentibus, ramis secundariis longissimis subsimplicibus undulatis flexuosis brevibus secundis, ramulis patentibus distantibus, articulis granuliferis diametro triplo-quintuplo longioribus. Nobis *in Lond. Journ. Bot.* vol. iv. p. 294. (TAB. CXCIII. Fig. I.)

HAB. Berkeley Sound, and St. Salvador Bay, Falkland Islands; on muddy rocks in the sea, abundant.

Cæspites 6-10 unc. longi, densissime fastigiati, e filis flexuosis intertextis gracillimis quasi crinitis formati. Rami flexuosi, elongati : secundarii valde elongati, simplices, ramulis brevibus longioribusve patentibus secundis ornati.

Apparently very distinct from any hitherto described species : its remarkable characters are its wavy habit, and the great length and simplicity of the upper branches, which are furnished with more or fewer, short, patent, secund ramuli.

PLATE CXCHI. Fig. I.--1, plant of the natural size; 2, branch and ramuli; 3, apex of branch; 4, portion of ditto :--very highly magnified.

6. CLADOPHORA *incompta*, Hook. fil. et Harv.; filis intricatis incomptis atro-viridibus opacis rigidis setaceis tortuosis vix ramosis, ramis longe nudis v. ramulis brevibus pectinatis circinato-inflexis ornatis, ramulis ultimis secundis v. alternis patentissimis obtusis approximatis remotisve, articulis diametro brevioribus quadratis v. $\frac{1}{2}$ longioribus. Nobis *in Lond. Journ. Bot.* vol. iv. p. 294. (TAB. CXCII. Fig. III.)

HAB. Hermite Island, Cape Horn; in the sea.

Cæspiles intertexti, horizontaliter extensi, atro-olivacei, rigidi. Fila intricata, irregulariter parce ramosa, latiuscula, C. simpliciusculæ diametro duplo excedentia. Rami flexuosi, sæpe undi, non raro ramulis involutis pectinatis obsiti, ut in C. flexuosa. Color huride ater v. virescens, opacus. Substantia siccitate rigida :---chartæ minime adhæret.

PLATE CXCII. Fig. III.—1, plant of the natural size; 2, portion of ditto, highly magnified, with abbreviated ramuli; 3, another portion of ditto and branch :—still more highly magnified.

7. CLADOPHORA simpliciuscula, Hook. fil. et Harv.; filis intricatis incomptis atro-viridibus opacis tlexuosis rigidiusculis capillaribus irregulariter subramosis, ramis valde remotis elongatis simplicibus, ramulis perpaucis patentissimis filiformibus sæpe secundis articulis diametro æqualibus v. $\frac{1}{2}$ -2 plo longioribus, sacculum endochromatis intus foventibus. Nobis in Lond. Journ. Bot. vol. iv. p. 295. (TAB. CXCII. Fig. IV.)

HAB. Hermite Island, Cape Horn, and the Falkland Islands; on sea-weeds, stones, and shells.

Caspites intertexti, opaci, luride virides, subhorizontaliter extensi. Fila 1-2 unc. longa, remote et irregulariter ramosa; rami ramulis perpaueis aucti :-- chartæ non adhæret.

Allied to *C. riparia*, but more robust, also near the *C. flagelliformis* of the Cape of Good Hope, but with a very different habit from that plant.

S. CLADOPHORA glaucescens, Griff.? Harv. Manual, p. 139.

HAB. Hermite Island, Cape Horn; rare.

We are not at all satisfied with the reference of the Antarctic plant to the British *C. glaucescens*, which has slenderer filaments. The specimens resemble that species more nearly than any other, and are not in a sufficiently good state for a proper comparison.

48. OSCILLATORIA, Vauch.

1. OSCILLATORIA *purpurea*, Hook. fil. et Harv.; strato gelatinoso tenaci siccitate translucente purpureo, filis violaceis omnium tenuissimis dense intertextis curvatis longe radiantibus, striis inconspicuis. Nobis *in* Lond. Journ. Bot. vol. iv. p. 297.

HAB. Kerguelen's Land; in alpine rivulets, alt. 300-700 feet.

Species admodum singularis, Lyngbyæ prolificæ, Grev. (Scot. Crypt. Flor. t. 303,) plerisque notis affinis, nec

non (suadente clariss. Berkeley), cum Bysso aquatico, D.C., (Geneva Trans. vol. ii. p. 29,), Oscillatoria rubescente, Bory et cum Conferva purpurea quoque conferenda. Fila muscos submersos strato gelatinoso translucente vestientia, dum maxime amplificata lineis transversis obscure notata.

2. OSCILLATORIA autumnalis, Agardh Syst. p. 62.; Harvey, Manual of Brit. Alg. p. 165.

HAB. Falkland Islands; on wet rocks; Cockburn Island, Graham's Land (Lat. 64° S. Long. 57°.W.) in moist places.

We have carefully compared this with Captain Carmichael's Appin specimens of *O. autumnatis*, and find them to be quite the same species, which is considered common in England, though Mr. Hassall quotes Captain Carmichael's habitat as the only one. The figure in the last named author's 'British Fresh-water Algæ,' is very unlike either Captain Carmichael's or the Antarctic specimens; in both of which the striæ are nearer to one another than the filament is broad. The diameter of the Cockburn Island filaments is $\frac{1}{2600}$ of an inch.

49. CALOTHRIX, Alg.

1. CALOTHRIX olivacea, Hook. fil. et Harv.; cæspite majnsculo intense olivaceo v. ærugescente erecto strictiuseulo, filis basi dichotome v. alterne divisis luteis flavidisve superne strictiusculis flexuosisve in funiculos crispatos tenaces cohærentibus per totam longitudinem connexisve apice liberis obtusiusculis, endochromate opaco obscure striato, articulis diametro longioribus brevioribusve. Nobis *in Lond. Journ. Bot.* vol. iv. p. 296. (TAB. CXC. Fig. III.)

HAB. Christmas Harbour, Kerguelen's Land; in alpine rivulets, adhering to stems and leaves of mosses, &c.

Cæspites fasciculati, interdum extensi, $\frac{1}{2}$ unc. longi, intense colorati, olivacci læte ærugescentesve, siccitate vix nitentes. *Fila C. distorta* multoties latiora, flexnosa sed non torta, sæpissime in fasciculos siccitate erispatos connexa, circa $\frac{1}{2000}$ unc. lata, basi flavescentia, ramosa v. divisa.

A very pretty species and quite distinct from any European one with which we are acquainted.

PLATE CXC. Fig. III. —1, plant of the natural size ; 2, filaments ; 3, upper, and 4, lower portion of ditto :— much magnified.

2. CALOTHRIX DISTORTA, Harvey, Manual of Brit. Alg. p. 158. Engl. Bot. t. 257.

HAB. Falkland Islands; in pools of fresh water.

The specimens, though in a very indifferent state, are clearly referable to the English C. distorta.

50. LINGBYA, Ag.

1. LYNGBYA muralis, Agardh; Conferva murahis, Dillwyn, Brit. Conferv. t. vii.

HAB. Falkland Islands; on the ground, abundant.

These we have carefully compared with original British specimens of *L. muralis*, and find them to differ only in having the filaments rather broader and more opaque.

2. LYNGBYA *fragilis*, Hook. fil. et Harv.; filis minntis tenuissimis fragilibus flavo-viridibus tortuosis implexis in stratum tenue lutescens cohærentibus, striis densissimis. Nobis *in Lond. Journ. Bot.* vol. iv. p. 296. (TAB. CXCIII. Fig. II.)

HAB. Berkeley Sound, Falkland Islands; on the fur of a dead rabbit.

Fila L. murali subsimillima, sed diametro $\frac{1}{2}$ angustiona, fragillimaque. *Stratum* tenue, lutescens v. flavo-virescens, vix nitens.

Fuegia, the

A plant so nearly related to the common L. muralis, of Britain, as to require no detailed description.

PLATE CXCIII. Fig. II.—1, plant of the natural size; 2, thread, highly magnified; 3, portion of ditto :--still more highly magnified.

3. LYNGBYA subarticulata, Hook. fil. et Harv.; filis tenuissimis laxe implexis vix tortis hic illic obscure subarticulatis, sporidiis disciformibus diametro variis filum vix $\frac{1}{2}$ æquantibus ad articulos spurios fili solutis.

HAB. Christmas Harbour, Kerguelen's Land; creeping amongst Ulvæ &c., on wet rocks near the sea.

Fila cylindracea, sub $\frac{1}{1000}$ unc. lata, pellucida, obscure articulata, articulis diametro ter longioribus, intus cum axi e sporidiorum disciformium composito ancti. *Sporidia* læte viridia, opaca, ter quaterve latiora quam longa.

A very different species from either of the two former. Each filament is a transparent tube, very obscurely incrassated, as if jointed here and there, and containing an axis of sporidia about half its own diameter. The sporidia are discoid, of various breadth and length, but always much broader than long, of a bright green colour, and interrupted opposite the spurious articulation of the filament.

51. MICROCOLEUS, Desmaz.

1. MICROCOLEUS repens, Harvey; Manual, p. 168. Oscillatoria repens, Agardh Syst. p. 61. O. chthonoplastes β, Harvey in Hook. Brit. Fl. vol. ii. p. 373.

HAB. Cockburn Island, Graham's Land; (Lat. 64° S. Long. 57° W.) on the ground.

Onr specimens are very poor, as might be expected from the native place being on the limits of vegetation in that quarter of the globe which the plant inhabits.

52. ULVA, L.

1. ULVA Lactuca, Linn; Grev. Scot. Crypt. Flor. t. 313.

HAB. Hermite Island, Cape Horn; sparingly.

2. ULVA latissima, Linn.; Engl. Bot. t. 1551.

HAB. Hermite Island, Cape Horn; Falkland Islands and Kerguelen's Land; very abundant.

There is probably no shore between that of Iceland and Cape Horn, that does not produce abundantly this species of Alga.

3. ULVA rigida, Agardh, ? Syst. Alg. p. 189.

HAB. Port William, Falkland Islands; common.

Our specimens appear to be only a young dark-coloured variety of the former. The species is also a native of Brazil, and the west coast of South America.

4. ULVA Linza. Linn.; Harv. Phyc. Brit. t. XXXIX.

HAB. Falkland Islands and Hermite Island, Cape Horn; abundant.

An abundant plant in the Atlantic and Mediterranean shores of Europe; it has also been collected in New Zealand.

5. ULVA crispa, Lightf.; Harvey, Manual, p. 171.
HAB. Berkeley Sound, Falkland Islands; on moist rocks; Cockburn Island, Graham's Land; very abundant.

A highly interesting species, because it is one of the very few terrestrial plants that have been gathered on the limits of vegetation both in the Northern and Southern Hemispheres. It was collected in Spitzbergen, (in 80° . N.) by the officers of Captain Parry's Expedition towards the North Pole, and is a native of many intervening latitudes. We have carefully compared these specimens with Agardh's original ones of U. crispa, from Norway, and find them to be identical. The Cockburn Island specimens are in fine fruit

6. ULVA cristata, Hook. fil et Harv.; pusilla, stratum continuum furfuraceum efficiens, frondibus crispatis lacunosis latioribus quam longis supra medium in laciniis perplurimis fissis, laciniis filiformibus fistulosis tortis pluries divisis processubus corniculatis simplicibus ramosisque undique obsitis, substantia tenerrima, sporis confertis irregulariter dispositis rarius quaternis.

HAB. Kerguelen's Land; in moist clefts of rocks overhanging Christmas Harbour, growing with Trypothallus anastomosans.

Frondes singulæ 2–6 lin longæ, latiores quam longæ, sessiles, basi contractæ, læte virescentes, fragiles, marginibus crispatis, superficie lacinioso v. profunde rugoso; laciniis perplurimis gracilibus compressis v. teretibus, fistulosis, processubus divaricatis undique ornatis.

A species so closely resembling the U. crispa, that we at first sight confounded it with that plant: it is, however, abundantly distinct, in the much smaller spores, and in the curious long and slender lacinize of the frond, which are tubular in the specimens we have examined, and, as well as the margins of the sessile frond, are studded with short simple or divided horn-like processes, or abbreviated ramuli.

53. MASTODIA, Hook. fil. et Harv.

Frons plana, membranacea v. subcarnosa, viridis, late expansa, inordinate areolata. Fructificatio duplex : 1°. Sporidia granulæformia, in areolis indefinita (ut in Ulva) fronde immersa. 2° Conceptacula mammæformia, fronde immersa, apice mamilla instructa, materie grumosa repleta, sporasque ellipticas foventia.—Genus Ulvæ proximum, et nisi presentia conceptaculorum nullo modo distinguendum.

1. MASTODIA tessellata, Hook. fil. et Harv. Ulva tessellata, nobis in Lond. Journ. Bot. vol. iv. p. 297. (TAB. CXCIV. Fig. II.)

Var. a. fronde tenuissima, laciniis longioribus.

Var. β . fronde carnosa siccitate rigida, lacimis rotundatis.

HAB. Kerguelen's Land; var. a. in streams of fresh-water. Var. β . on stones occasionally exposed in a fresh-water lake.

Frons foliacea, 1-2 unc. lata, luride viridis, subplicata, siccitate rigidiuscula, suberecta v. in var. β . horizontaliter expansa, sub lente granulis majușculis opacis in areolas quadratas compositas dispositis pulcherrime quasi tessellata, demum in lacinias plurimas undulato-crispatas rotundatas fissa; areolis quadratis, lineis hyalinis circumscriptis, granulis magnis quaternis. Conceptacula exemplaribus omnibus nobis visis perplurima, ad angulos areolarum majorum sita, elevata, mammæformia, apice pallidiore, crassa et carnosa, intus cava, materic grumosa sporisque lineari-ellipticis viridibus immixtis farcta.

Erroneously described as a marine species in the London Journal of Botany. Even when destitute of fruit it is specifically very distinct from any Ulva, especially in the great size of the granules, and their comparative remoteness from one another. The curious hemispherical bodies are abundant in all the specimeus, and resemble in some degree

the capsules of a *Nitophyllum*: whether they be an abnormal development, or organs of fructification rarely developed in the genus, may be a point of dispute. At first sight they were supposed to be caused by the puncture of an aquatic insect or other animal; but their appearing in specimens from different localities; their position, constantly at the angles of the greater areolæ, where four of these meet; their uniform size; the constant presence of the cavity filled with elliptic spores, taken especially along with the fact, that there is no sign of disease or lesion in the frond, would indicate these to be organs in a normal condition.

PLATE CXCIV. Fig. II.—1, Var. a.; 2, var. β .; both of the natural size; 3, apex of frond; 4, portion of ditto with conceptacles; 5, portion of ditto more highly magnified; 6, vertical section of conceptacle; 7, grumous contents from ditto; 8, spores from ditto:—all very highly magnified.

54. ENTEROMORPHA, Link.

- 1. ENTEROMORPHA compressa, Grev. Alg. Brit. p. 180. t. 18.
- HAB. Hermite Island, Cape Horn, Falkland Islands, and Kerguelen's Land; very abundant.
- 2. ENTEROMORPHA intestinalis, Link. Grev. Alg. Brit. p. 179.
- HAB. Hermite Island, Cape Horn; Falkland Islands, and Kerguelen's Land; with the former.

These two species enjoy equally wide ranges with the Ulva latissima. I have found it very difficult to distinguish between this and the former species, even when growing, and between E. compressa and Ulva Linza in a young state. In the Falkland Islands the U. latissima abounds in the land-locked Lagoons, and the U. Linza in the harbours where no heavy seas run; whilst the Enteromorpha compressa, and intestinalis, may be collected on the shores of the weather-beaten coasts. Hence it becomes difficult for the collector to regard these species, whose structure and organization are so similar, as anything more than states of one plant, which commences as a pyriform bladder wherever it germinates, but whose future outline is determined by the depth and tranquillity or the reverse of the clement it inhabits, and other natural causes. Such specimens as our Herbaria generally afford, are too often, if not fragmentary, immature; the full development of the species being arrested by the collector, who is content with one entire specimen in whatever stage of growth, and generally preserves it without any note of the conditions under which it was gathered. A few observations on the forms which the Alga assume during different stages of their growth, would be eminently useful: portions of a crop of such species as this, which often covers shells or pebbles, might readily be transported to other waters, whose state is very different from what the plant enjoyed before. It cannot be doubted that great changes in form would be the consequence ; and it is on outline alone that specific characters are chiefly founded.

55. PORPHYRA, Ag.

1. PORPHYRA vulgaris, Ag. Grev. Alg. Brit. p. 169.

HAB. Hermite Island, Cape Horn; the Falkland Islands, and Kergnelen's Land; very abundant.

This has as wide a range in latitude and longitude as Ulva latissima.

2. PORPHYRA laciniata, Ag. Ulva umbilicata, Engl. Bot. t. 2296.

IIAB. Hermite Island, Cape Horn; the Falkland Islands, and Kerguelen's Land; very abundant.

Obviously a variety, or rather state of *P. vulgaris*; of which the *P. Columbina*, Mont., is probably the young, and *P. Capensis*, Kütz. another variety.

56. TRYPOTHALLUS, Hook. fil. et Harv.

Frons subcartilaginco-carnosa, vix gelatinosa, undulato-crispata, lobata, e cellulis hyalinis in stratum conglobatis

efformata, primum continua, matura terebrata v. clathrata (cellulis in lineis anastomosantibus dispositis) demum in massam gelatinosam subgranulosam collabens. *Sporæ* (seu granulæ) plerumque binæ, anguste lineari-oblongæ. — *Genus* Palmellæ *affine*, *sed indole frondis diversissimum*.

1. TRYPOTHALLUS anastomosans; Hook. fil. et Harv.; Palmella? anastomosans, nobis in Lond. Journ. Bot. vol. iv. p. 298. (TAB. CXCIV. Fig. I.)

HAB. Christmas Harbour, Kergnelen's Land; in clefts of rocks, and in damp caves, near the sea.

Frons $\frac{1}{4} - \frac{1}{2}$ unc. longa, undulato-crispata, pallide viridis, prima facie Ulvam furfuraceam referens, e strato unico cellularum formata, translucida; junior continua, suberecta v. horizontaliter extensa, margine lobata; matura (e cellulis in lineas dispositis) pulcherrime clathrata, foraminibus diametro variis pertusa. Cellulæ hyalinæ, marginibus sub lente vix distinctis, dense aggregatæ, rotundatæ v. obtuse angulatæ. Sporæ axi cellularum immersæ, plerumque binæ, læte virides (sub lente), post marcescentiam frondis diametro auctæ, cellulasque fere implentes.

We referred this plant doubtfully to *Palmella*, in the London Journal of Botany; and now, unhesitatingly, we place it in a new genus, most distinct from any previously defined. Under the microscope it is a very beantiful object, the full grown specimens appearing as a transparent frond, firmer and more membranous than *Palmella*, and much thicker in proportion than any *Ulva*, beautifully clathrate or formed of anastomosing branches: the branches are composed generally of one, or more rarely, of two collateral lines of cells, each containing a pair of parallel minute spores, of a bright green colour, placed at right angles to the axis of the branch.

PLATE CXCIV. Fig. I.—1, plant in its foliaceous and reticulated condition; 2, the same at a later stage, forming a gelatinous mass; 3, the same with the spores disunited;—all of the natural size; 4, portion of foliaceous state; 5 and 6, portions with anastomosing structure; 7 and 8, spores:—all very highly magnified.

57. PROTOCOCCUS, Ag.

1. PROTOCOCCUS stercorarius, Berk.; strato aurantiaco demum subrimoso, globulis demum margine pellucidis nucleis subgranulatis conformibus.

HAB. Falkland Islands; on cow-dung; abundant.

P. nivali, Desm., (quæ eadem est ac P. pluvialis, Flotow) affinis, sed globulis minoribus saturatius coloratis, Hæmatococco Orsinii, Menegh., quoque referens.

This curious vegetable appears abundantly in places frequented by cattle, covering their droppings with a pale orange stratum in a very short space of time. For the identification and description of the species we are indebted to our learned friend Mr. Berkeley, who had previously observed the same plant in England.

58. NOSTOC, Vauch.

1. Nostoc commune, Vauch. Conferv. p. 223. t. 16. f. l.

HAB. Christmas Harbour, Kerguelen's Land; on wet rocks near the sea.

The Rev. M. J. Berkeley has favoured us by examining this, the following, and several other of the lower forms of Alga collected during the Antarctic Expedition : of the present he says that it scarcely differs from the N. commune, of England, which is common throughout Europe, and in Bolivia and the Canary Islands.

2. Nostoc microscopicum, Carm.?; Harv. Man. Brit. Alg. p. 184.

HAB. Christmas Harbour, Kerguelen's Land; on wet rocks near the sea.

Specimens rather larger than those of British growth, but not otherwise different. Mr. Hassall considers the N. microscopicum to be a variety of N. muscorum, Ag.

59. ANABAINA, Bory.

1. ANABAINA tenax, Hook. fil. et Harv.; strato globuloso definito lobato gelatinoso fluctuante æruginoso, filis densissime intertextis flexuosis moniliformibus inæqualibus hic illic interruptis, articulis plerumque globosis angulatisve nunc transverse elongatis, majoribus ellipticis oblongis limbo hyalino cinctis solitariis plurimisve. Sphærozyga tenax, *nobis in Lond. Journ. Bot.* vol. iv. p. 298. (TAB. CXCIII. Fig. III.)

HAB. Falkland Islands; in small pools of water on the hills.

Stratum 1-3 une. latum, e massis $\frac{1}{4} - \frac{1}{2}$ uneialibus conglobatis efformatum, gelatinosum, hyalinum, pulchre æruginosum, natans. Substantia gelatinosa, sub lente oculum fugiens. Fila perplurima, dense aggregata, diametro varia. Articuli sub lente glauco-virescentes, opacæ, majores translucidæ.

A very distinct and beautiful species, evidently congeneric with the *Sphærozyga Jacobi*, of which the Rev. M. J. Berkeley has published an excellent figure in the Supplement to English Botany, (t. 2826. fig. 2.) but which we do not consider generically distinct from *Anabaina*. The granular substance of the larger articuli is of a different nature from that filling the smaller one, being more transparent, and confined in a proper cyst, between which and the border of the articulation there is a transparent space. The stratum is as firm as that of *Nostoc cæruleum*, and the specimens preserved resemble a dried mass of *Oscillatoria*.

Specifically this differs from A. Jacobi in the form of the stratum, and from A. flos-aquæ in the straightness of the larger articulations.

PLATE CXCIII. Fig. III.-1, plant of the natural size; 2, threads; 3, portion of a thread with spores; 4, spores :-highly magnified.

60. CHROOLEPUS, Ag.

1. CHROOLEPUS aureus, Harv. in Hook. Brit. Flor. vol. ii. p. 380. Conferva aurea, Dillwyn, Hist. Conf. t. 35.

HAB. Hermite Island, Cape Horn; Kerguelen's Land, and the Falkland Islands; very abundant on the under surfaces of rocks near the sea, &c.

One of the commonest vegetable productions in the Antarctic Islands, growing under circnmstances where no *Lichen*, or other cryptogamic plant, flourishes. It was always found near the *Lecanora miniata*, and is very abundant in situations sheltered from the direct rays of the sun. When fresh, or rather during drying, it emits a very evident smell of violets.

2. CHROOLEPUS ebeneus, Ag. Syst. Alg. p. 36. Conferva ebenea, Dillwyn, t. 101. Byssus niger, Engl. Bot. t. 702.

HAB. Hermite Island, Cape Horn; in clefts of rocks in the woods.

Like the former, this species, invariably shuns the light in the south. It was found in damper places than *C. aureus.* Both are, very probably, abnormal states of some *Lichen*.

LVI. DIATOMACEÆ, Ag.

The Waters and the Ice of the South Polar Ocean were alike found to abound with microscopic vegetables belonging to this Order. Though much too small to be discernible by the naked eye, they occurred in such countless myriads, as to stain the Berg and the Pack-Ice, wherever they were washed by the swell of the sea; and when enclosed in the congealing surface of the water, they imparted to the Brash and Pancake-Ice a pale ochreous colour. In the open ocean, northward of the Frozen Zone, this Order, though no doubt almost universally present, generally eludes the search of the naturalist; except when its species arc congregated amongst that mucous scum which is sometimes seen floating on the waves, and of whose real nature we are ignorant; or when the coloured contents of the marine animals who feed on these Algæ arc examined. To the south, however, of the belt of ice which encircles the globe, between the parallels of 50° and 70° S., and in the waters comprised between that belt and the highest latitude ever attained by man, this vegetation is very conspicuous, from the contrast betweeu its colour and the white snow and ice in which it is imbedded. Insomuch, that, in the eightieth degree, all the surface-ice carried along by the currents, the sides of every berg, and the base of the great Victoria Barrier itself, within reach of the swells, were tinged brown, as if the Polar waters were charged with oxide of iron.

As the majority of these plants consist of very simple vegetable cells, enclosed in indestructible silex (as other *Algæ* are in carbonate of lime), it is obvious that the death and decomposition of such multitudes must form sedimentary deposits, proportionate in their extent to the length and exposure of the coast against which they are washed, in thickness to the power of such agents as the winds, currents and sea, which sweep them more energetically to certain positions, and in purity to the depth of the water and nature of the bottom. Hence we detected their remains along every ice-bound shore, in the depths of the adjacent ocean, between eighty and 400 fathoms. Off Victoria Barrier (a perpendicular wall of ice, between one and two hundred feet above the level of the sea), the bottom of the ocean was covered with a stratum of pure white or green mud, composed principally of the siliceous cells of *Diatomaceæ*. These, on being put into water, rendered it cloudy, like milk, and took many hours to subside. In the very deep water off Victoria and Graham's Laud, this mud was particularly pure and fine; but towards the shallower shores, there existed a greater or less admixture of disintegrated rocks and sand; so that the organic compounds of the bottom frequently bore but a small proportion to the inorganic.

Being indebted to the works of the illustrious Ehrenberg for all I knew of these organisms, previous to the sailing of the Antarctic Expedition, I had supposed the *Diatomaceæ* to belong to the Animal Kingdom *; and as they are unaccompanied in the Antarctic region by any evidence of a higher order of plants, I had always supposed vegetation to cease at a much lower latitude than these productions actually attain. The species were, however, collected on every available occasion, and transmitted, on my return to England, to Professor Ehrenberg, whose determination of the genera and species is here introduced, at the suggestion of the Rev. M. J. Berkeley and other eminent Cryptoganuic botanists.

* It is well known that the true uature of the *Diatomaceæ* has been long and unsuccessfully disputed, being claimed both by botamists and zoologists. No conclusive evidence on this subject had been adduced, till, within these very few days, it was the singular good fortune of my friend, Mr. Thwaites, of Bristol, a most acute observer and profound Cryptogamist, to detect several species of *Diatomaceæ* conjugating, in a manner perfectly analogous to that pursued by the *Zygnemata*: a fact which leaves no doubt of their vegetable origin in the minds of persons acquainted with his interesting observations. I am indebted to Mr. Thwaites for specimens of three British species of *Eunotia*, and *Gomphonema*, illustrating this important discovery, and mounted in fluid, after the beautiful plan invented by that gentleman for preserving vegetable tissues moist, and always ready for the microscope in the form of slides.

I must offer some apology for omitting a class of organisms which have been investigated, and considered of vegetable origin, by Prof. Ehrenberg, and which are almost equally abundant in the Antarctic Ocean with the *Diatomaceæ*, whether on the surface or at the bottom of the sea: these are the *Phytolitharia*, Ehrb. I am not aware of the precise limits of this Order, and of many of the genera composing it; but from casual allusions, I gather that the term *Phytolitharia* is a conventional one, employed to designate the siliceous and other inorganic particles, deposited in plants of a higher structure. Thus, *Lithodermatium* is a genus whose species are represented by modifications of the siliceous epidermis of one or many species of *Equisetum*; and the *Lithostylidia* are the siliceous cells of *Gramineæ**. It is not my object to discuss in this place the expediency of constituting such orders, genera, and species. The total absence of *Equiseta* from the Antarctic Flora, and of *Gramineæ* or other phænogamic plants from any position within 700 miles of Victoria Barrier where the *Phytolitharia* abound, renders it in the highest degree improbable that the latter should be of vegetable origin.[†]

A few remarks on the phases and situations under which these curious vegetables occurred, will not be misplaced here, especially as I have little to add to what is already known of their habits and organization.

Scattered on the surface of the ocean, the Antarctic *Diatomaceæ* were seen connected in filaments, or resolved into the simple frustules, of which they are composed. When entire, they shewed no signs of motion or irritability. The grumous or granular contents of the cells were yellow under the microscope; but in mass the same species assumed an orange-brown, or burnt Sienna colour; the intensity of which depended on the denseness with which they were packed together.

The various means employed for selecting the species varied according to circumstances, as the following enumeration of the processes pursued will show. 1. Sea-water was filtered through closely woven bibulous paper (filter-paper), which latter was folded, dried, and carefully put away. If a certain measure of water be always thus treated, an approximate knowledge of the abundance and scarcity of the various species and genera occurring at different positions, may be gained. 2. The scum of the ocean almost invariably contains many species entangled in its mass; it was preserved in small phials, well secured. 3. A tow-uet of fine muslin, used when the vessel's rate does not exceed two or three knots, secures many kinds, which may be washed off the muslin, and collected on filter paper. 4. The stomachs of Salpæ[‡] and other (especially of the naked) mollusca, invariably contain Diatomacee, sometimes several species. These Salpe were washed up in masses on the Pack ice, and in decay they left the snow covered with animal matter impregnated, as it were, with Diatomacea: the reliquia were preserved in spirits. 5. The dirt and soil of the Penguin Rookeries, and especially their Guano, abound in Diatomaceæ, perhaps originally swallowed by the Salpæ and Cuttle-fish, which themselves become the prev of the Penguins. 6. Ice encloses Diatomaceae: they are deposited on the already formed ice by the waves, or frozen into its substance during calm weather, when the npper stratum of water rapidly congeals. Ice, so formed, generally breaks up by the swell of the sea into thin angular masses, which become orbicular by attrition, whence the name Pancake-ice. The Pancake-ice was often seen a few hours after a calm, covering leagues of ocean, and uniformly stained brown from the abundance of these plants. It was taken in buckets, and when removed from the water appeared perfectly pure and colourless. On melting, however, it deposited a pale red clondy precipitate, excessively light, consisting wholly of Diatomacea. This precipitate was bottled on the spot, and proved

* See Ehrenberg, in Schrift. Berlin Akad., June, 1841.

† On the contrary, I caunot but suspect that some of these *Phytolitharia* are the remains of *Crustacea*, and especially the siliceous (?) particles, which occur in the tunics of naked *Mollusca*.

‡ I do not remember to have examined the contents of the stomach of any Salpa between the latitudes of the N. Tropic and the 80° S., which did not contain the remains of Diatomaceæ. Dictyocha aculeata was universally found in the stomachs of those I opeued when off Victoria Land.

more rich in species than any of the other collections. The specimens were also the best preserved; for Professor Ehrenberg observes, that some* thus obtained, appeared as if still alive, though collected three years previous to his examination, and subjected to many vicissitudes of climate. The snow sometimes falls on the surface of the still ocean-water, and does not freeze, but floats a honey-like substance, often called Brash-ice: treated in the same way as the Pancake-ice it yielded an abundant harvest. 7. The mud and other soundings from the bottom of the ocean, when brought up on the arming of the deep sea-lead, or the chlam or dredge, generally contain the siliceous skeletons or coatings of many species, with the markings on their surface retained.⁺ 8. The fresh and salt waters and muddy estuaries of the Falkland Islands, and similar localities, present us with species, occurring under circumstances, altogether similar to what accompany their allies in Europe.

The universal existence of such an invisible vegetation as that of the Antarctic Ocean, is a truly wonderful fact, and the more from its not being accompanied by plants of a high Order. During the years we spent there, I had been accustomed to regard the phenomena of life as differing totally from what obtains throughout all other latitudes; for everything hving appeared to be of animal origin. The ocean swarmed with Mollusca, and particularly entomostracous Crustacea, small whales and porpoises: the sea abounded with penguins and seals, and the air with birds: the animal kingdom was ever present, the larger creatures preving on the smaller, and these again on smaller still: all seemed carnivorous. The herbivorous were not recognized, because feeding on a microscopic herbage, of whose true nature I had formed an erroneous impression. It is, therefore with no little satisfaction that I now class the Diatomaceæ with plants, probably maintaining in the South Polar Ocean that balance between the animal and vegetable kingdoms, which prevails over the surface of our globe. Nor is the sustenance and nutrition of the animal kingdom the only function these minute productions may perform : they may also be the purifiers of the vitiated atmosphere, and thus execute, in the Antarctic latitudes, the office of our trees and grass-turf in the temperate regions, and the broad leaves of the palm, &c., in the Tropics. Though we possess incontestible proofs of the abundance of *silica*, contained in the ocean, from its being secreted so copiously by these plants, we are ignorant of the process by which it is assimilated, and the chemical state in which it is suspended in the sea-water. The end these plants serve in the great scheme of nature is apparent, on inspecting the stomachs of many sea-animals, as above stated. Owing to the indestructible nature of their shields, they tell their own tale.

I shall now notice the most remarkable feature in the distribution of these organisms. They possess more than ordinary interest, many of the species being distributed from Pole to Pole; while these, or others, are preserved in a fossil state, in strata of great antiquity. There is probably no latitude between that of Spitzbergen and Victoria Land, where some of the species of either country do not exist: Iceland, Britain, the Mediterranean Sea, North and South America, and the South Sea Islands, all possess Antarctic *Diatomaceæ*. The siliceous coats of species only known living in the waters of the South Polar Ocean, have, during past ages, contributed to the formation of rocks; and thus they outlive several successive creations of organized beings. The Phonolite stones of the Rhine, and the Tripoli stone, contain species identical with what are now contributing to form a sedimentary deposit (and perhaps at some future period a bed of rock), extending in one continuous stratum for 400 measured miles. I allude to the shores of the Victoria Barrier; along whose coast the soundings examined were invariably charged with Diatomaceous remains, constituting a bank which stretches 200 miles north from the base of Victoria Barrier, while the average depth of water above it is 300 fathoms, or 1,800 feet.[‡]

* Fragilaria pinnulata, and some Coscinodisci.

[‡] This great depth, reaching to within a quarter of a mile of the Barrier, whose height appeared nowhere to exceed 200 feet, proves that the latter does not rest on this bank. The accumulation, however, of snow on the

[†] The soundings were invariably in greenish mud, into which the lead sometimes sunk for two feet. At times, this mud seemed almost wholly composed of Diatomaceous remains.

Again, some of the Antarctic species have been detected floating in the atmosphere which overhangs the wide ocean between Africa and America. The knowledge of this marvellous fact we owe to Mr. Darwin, who, when he was at sea near the Cape de Verd Islands, collected an impalpable powder which fell on Captain Fitzroy's ships. He transmitted this dust to Ehrenberg, who ascertained it to consist of the siliceous coats, chiefly of American *Diatomaceæ*, which were being wafted through the upper regions of the air, when some meteorological phenomenon checked them in their course, and deposited them on the ship and surface of the ocean.

The existence of the remains of many species of this Order (and amongst them some Antarctic ones), in the volcanic ashes, pumice, and scorize of active and extinct volcanoes (those of the Mediterranean Sea and Ascension Island for instance), is a fact bearing immediately upon the present subject. Mount Erebus, a volcano 12,400 feet high, of the first class in dimensions and energetic action, rises at once from the ocean, in the 78th degree of south latitude, and abreast of the *Diatomaceæ* bank, which reposes in part on its base. Hence it may not appear preposterous to conclude, that, as Vesuvius receives the waters of the Mediterranean, with its fish, to eject them by its crater'; so the subterranean and subaqueous forces which maintain Mount Erebus in activity, may occasionally receive organic matter from this bank, and disgorge it, together with those volcanic products, ashes and pumice.

Along the shores of Graham's Land and the South Shetland Islands, we have a parallel combination of igneous and aqueous action, accompanied with an equally copious supply of *Diatomaceæ*. In the Gulf of Erebns and Terror, 15 degrees north of Victoria Land, and placed in the opposite side of the globe, the soundings were of a similar nature with those of Victoria Land and Barrier, and the sea and ice as full of *Diatomaceæ*. This was not only proved by the deep-sea lead, but by the examination of bergs, which, once stranded, had floated off and become reversed, exposing an accumulation of white friable mud, frozen to their bases, which abounded with these vegetable remains.

The following systematically arranged catalogue of the hitherto described Antarctic species is drawn up from various papers by Professor Ehrenberg, but principally from that which appeared in the 'Monatsberichten der Berliner Akad. der Wissenschaften '' for May, 1841, and which has been reprinted in Taylor's 'Annals of Natural History', and in the Appendix of Sir James Ross' 'Narrative of the Antarctic Expedition'. A few Falkland Island and Kerguelen's Land species have subsequently been examined by Mr. Thwaites, to whom, and to the Rev. Mr. Berkeley, I am much indebted for the assistance they have afforded me in this group. The arrangement of the genera followed is that of M. Kützing's great work on this order.

1. EUNOTIA, Ehrb.

1. EUNOTIA gibberula, Ehrb. Epithemia gibberula, Kütz. Kieselsch. Bacill. p. 35. t. 29. f. 54, c.

HAB. Open Ocean, in Pancake-iee, Lat. 75° S. Long. 170° W.

An inhabitant of the Baltic Sea. Found *fossil* at Newhaven, in Connecticut, in volcanic ashes from the Rhine and amongst an atmospheric dust which fell near the Cape de Verd Islands.

2. EUNOTIA amphioxys, Ehrb. Kütz. l. c. p. 44. t. 30. f. l.

HAB. Falkland Islands, Lesson. Cockburn Island, amongst the guano of a Penguin rookery.

surface of the barrier, in a climate where there is no thaw throughout the year, and where snow lies perennially, will result in the sinking of the barrier and its base becoming imbedded in this stratum of vegetable debris. Supposing the barrier, then, to have a progressive motion, such as smaller but similar glaciers exhibit, the result would be flexnres of the pasty stratum of mud upon whose edge it rests, and against whose walls it would in time abut, as the deposit thickens.

Found living in the German Ocean and Sandwich Islands; in the natural paper of Silesia, and dead in the guano of Peru. *Fossil* as floating in the air with the former species. In peat, Iceland; earth, Labrador, and in strata on the banks of the Euphrates and Oxus. In the volcanic tuff of the Rhine and in Phonolite.

3. EUNOTIA Faba, Ehrb. Epithemia Faba, Kütz. l. c. p. 36. t. 5. f. 21.

HAB. Falkland Islands; on marine Confervæ, Lesson.

Abundant both recent and fossil, in Germany, Sweden, Finmark, Newfoundland, Labrador, and the Oregon. Also found in the volcanic tuff of the Rhine.

4. EUNOTIA biceps, Ehrb. Kütz. l. c. p. 37. t. 29. f. 65. c.

HAB. Falkland Islands; on marine Confervæ, Lesson.

Found in earth at Labrador and the Oregon.

2. FRAGILARIA, Lyngb.

1. FRAGILARIA amphiceros, Ehrb. Schrift. Berl. Akad. Feb. 1844.

11AB. Victoria Barrier, in Pancake-ice and in mud from 190 fathoms. Graham's Land, in mud from 270 fathoms. In a floating scum, Lat. 64° S. Long. 160° W.

Occurs fossil in Virginia, U.S.

2. FRAGILARIA acuta, Ehrb. Kütz. Kieselsch. Bacill. p. 46. t. 16. f. 7. C.

HAB. In Pancake-ice off Victoria Barrier, and in Lat. 75° S. Long. 170° W. In the stomach of a Salpa, taken in the open occan, Lat. 64° S. Long. 157° W.

Previously only known as a fossil, occurring near Freiberg.

3. FRAGILARIA n. sp.? (indicated by Ehrb.)

HAB. Victoria Barrier; in mud from 190 fathoms.

4. FRAGILARIA pinnulata, n. sp. Ehrb. Schrift. Berl. Akad. May, 1844.

HAB. Pancake-ice, Lat. 75° S. 170° W., and near the continent of Victoria Land, 76° S., in Brashice. Graham's Land, in mud from 270 fathoms.

Onc of the most abundant Victoria Land Diatomaceæ.

5. FRAGILARIA rotundata, n. sp. Ehrb. l. c.

HAB. Pancake and brash-ice off Victoria Land and Barrier. In the stomachs of Salpa, taken in Lat. 66° S. and Long. 170° W. Graham's Land, in mud from 207 fathoms.

6. FRAGILARIA n. sp.? (indicated by Ehrb.)

HAB. Victoria Land; in Brash-ice.

7. FRAGILARIA granulata, n. sp. Ehrb. l. c.

HAB. In the stomachs of Salpæ, Lat. 66° S. Long. 157° W. In the open sea near Cape Horn (M. Schayer, fid. Ehrb.). In occanic scum, Lat. 64° S. Long. 160° W.

Lately indicated to exist as a fossil near the Araxes river.

S. FRAGILARIA constricta, Ehrb. Kütz. l. c. p. 46. t. 29. f. 25, c.

HAB. Falkland Islands; on marine Confervæ, Lesson.

Found in the volcanic tuff of the Rhine, and in Mexico.

9. FRAGILARIA rhabdosoma, Ehrb. F. capreina, Kütz. p. 45. t. 36. f. iii.

HAB. Falkland Islands; on marine Conferva, Lesson.

A frequent inhabitant of pools and ditches in England, and many other parts of the world, including Asia, Africa, America and the South Sea Islands. In the sand-hills of Patagonia, and in the volcanic tuff of the Rhine.

10. FRAGILARIA Trachea, n. sp. Ehrb. Schrift. Berl. Akad. l. c.

HAB. Falkland Islands; on marine Confervæ, Lesson.

11. FRAGILARIA Ventriculus, n. sp. Ehrb. l. c.

HAB. Falkland Islands; on marine Confervæ, Lesson.

3. MELOSEIRA, Ag.

1. MELOSEIRA n. sp.? filis moniliformibus tenuissime striatis pedunculo gelatinoso affixis, frustulis per paria coadunatis, junioribus sphæricis demum compressis, apicibus utrinque convexis. *Thwaites, MS*.

HAB. Kerguelen's Land; in the sea.

M. globiferæ, Harv. simillima, sed frustulis adultioribus semper compressis differt. Fila striata ut in M. globifera. Theorites, MS.

4. PYXIDICULA, Ehrb.

1. PYXIDICULA dentata, n. sp. Ehrb., Schrift. Berl. Akad. May, 1844.

HAB. Victoria Barrier; in Pancake-ice.

2. PYXIDICULA Hellenica, Ehrb. l. c.

HAB. Victoria Barrier; in Pancake-ice. Graham's Land; in mud from 270 fathoms (doubtful as to species).

This has been found fossil in Bermuda, the Ægean Sea, and Maryland, U.S.

3. PYXIDICULA n. sp.? Ehrb. l. c.

HAB. Victoria Barrier; in mud from 190 fathoms.

4. PYXIDICULA sp.?

HAB. In the stomachs of Salpa, Lat. 66° S. Long. 157° W.

5. HEMIZOSTER, N. G. Ehrb.

1. HEMIZOSTER tubulosus, Ehrb., Schrift. Berl. Akad. May, 1844. HAB. Victoria Barrier and Land; in Pancake-ice.

FLORA ANTARCTICA.

509

6. GALLIONELLA.

1. GALLIONELLA pileata, n. sp. Ehrb. l. c.

HAB. Victoria Barrier; in Pancake-ice.

2. GALLIONELLA sulcata, Ehrb., Schrift. Berl. Akad. April, 1837.

HAB. Victoria Land; in Pancake-ice (doubtful). Graham's Land; in mud from 270 fathoms.

An Arctic plant, having been observed at Melville Island. Also in the open ocean off Rio de Janeiro. It inhabits Peruvian and African guano; has been found in the sand-hills of Patagonia, fossil in Bermudas, Sicily, Algiers, Maryland and Virginia, U.S.; and in volcanic ashes from the Patagonian coast.

3. GALLIONELLA Sol, n. sp. Ehrb.; Schrift. Berl. Akad. May, 1844.

HAB. Victoria Barrier; in mud at 190 fathoms. Graham's Land; in mud from 207 and 270 fathoms.

4. GALLIONELLA tympanum, n. sp., Ehrb. l. c.

HAB. Graham's Land; in mnd from 207 fathoms.

5. GALLIONELLA Oculus, n. sp., Ehrb. l. c.

HAB. Graham's Land; in mud from 270 fathoms.

7. CAMPYLODISCUS, Ehrb.

1. CAMPYLODISCUS Clypeus, Ehrb. Kütz. Kieselsch. Bacill. p. 59. t. 2. f. v. 1-6.

HAB. Graham's Land; in mud from 270 fathoms.

Found fossil in Germany and Italy, in the Bermudas and in the *Mastodon* earth of the Plate river, in African guano, in the atmospheric dust near the Cape de Verd Islands, and in the volcanic tuff of the Rhine.

S. SURIRELLA, Ehrb.

1. SURIRELLA (?) australis, Ehrb.; Schrift. Berl. Akad. l. c. HAB. Falkland Islands; on marine Conferva, Lesson.

9. SYNEDRA, Ehrb.

1. SYNEDRA Ulna, Ehrb.; Infus. t. 17. f. 1. Kütz. l. c. p. 66. t. 30. Exilaria Ulna, Hassall, Brit. Fresh-water Alg. p. 433. t. 97. f. 2.

HAB. Victoria Barrier; in Pancake-ice (doubtful). Graham's Land; in mud 270 fathoms.

One of the most abundant and easily recognized of the *Diatomaceæ*, not only in Europe but throughout the globe. Mr. Hassall states it to be of very frequent occurrence in fresh-water ponds and ditches of England. It is also found in Icelandic peat, in marine mud from Spitzbergen, in the natural paper of Silesia, and in the Tropical Ocean off Rio. As a fossil or dead, it has occurred in Oran and Sicily, the United States, in alluvial deposits in Brazil, the Euphrates River, and in atmospheric dust off the Cape de Verd Islands. It is also found in the volcanic tuff of the Rhine and in Peruvian guano.

6 c

10. DICLADIA, N. G. Ehrb.

1. DICLADIA antennata, Ehrb.; Schrift. Berl. Akad. May, 1844.

HAB. Victoria Barrier; in Pancake-ice.

2. DICLADIA bulbosa, Ehrb. l. c.

HAB. Victoria Barrier; in Pancake-ice.

This and the preceding always occurred abundantly in the stomachs of the Victoria Land Salpa.

11. SCHIZONEMA, Ag.

1. SCHIZONEMA sp.? filis ramosis siccitate luride viridibus, frustulis ovato-oblongis siccis collapsis. Thwaites, MS.

HAB. Kerguelen's Land; abundant in the sea.

Very similar to, and possibly not distinct from, the British S. implicatum, Harv.

12. EX1LARIA, Grev.

1. EXILARIA, n. sp.? frustulis linearibus striatis e dorso visis leviter versus apices truncatos attenuatis, e latere visis utrinque obtuse apiculatis. *Thwaites, MS.*

HAB. Kerguelen's Land; on marine Confervæ.

G. truncatæ forma accedit, sed frustulis striatis ut in Synedra Ulna.

13. COCCONEIS, Ehrb.

1. COCCONEIS Placentula, Ehrb. Kütz. p. 73. t. 28. f. 13. c.

HAB. Falkland Islands; on marine Confervæ, Lesson.

Found living in the fresh waters of Europe; also in Iceland, Mexico and the Oregon river, Chili, the Sandwich Islands, and in African (?) guano.

2. COCCONEIS Scutellum, Ehrb. Kütz. l. c. t. 5. f. vi. 3-6.

HAB. Falkland Islands; ou marine Conferva, Lesson. Kerguelen's Land, also on marine Conferva.

Found living in the Atlantic, German, and Mediterranean Seas; on the coast of Iceland, in Peruvian guano and in volcanic tuff.

14. ACHNANTHES, Ay.

1. ACHNANTHES pachypus, Kütz. p. 76. t. 21. II. f. 3 and 29. f. 83.

HAB. Falkland Islands; on marine Conferva, Lesson.

Recorded by Montagne to be a native of Callao, in Peru.

2. ACHNANTHES longipes, Ag. Harv. Man. Alg. Brit. p. 200.

HAB. Kerguelen's Land; on Alga.

Specimen in a very bad state, but probably referable to this species. Thwaites, MS.

FLORA ANTARCTICA.

3. ACHNANTHES turgens, Ehrb. l. c.

HAB. Graham's Land; in mud from 270 fathoms.

15. LICHNOPHORA, Ag.

1. LICHNOPHORA abbreviata, Ag. Podosfenia abbreviata, Ehrb. Infus. p. 24. t. 18. f. 7.

HAB. Kerguelen's Land; in the sea.

Frustula omnino ut in exemplaribus ab Ehrenberg depietis, sed plura, stipiteque majore. *Thwaites*, *MS*. Oceurs as a parasite on *Ceramium rubrum*, in the Mediterranean Sea.

16. DIATOMA, Ag.

1. DIATOMA, n. sp.? frustulis angustissimis omnino aut fere lævissimis e dorso visis versus apices truncatos sub-dilatatis, e latere visis infra apices rotundatos constrictis.

HAB. Falkland Islands; on marine Confervæ.

D. etongata, Ag. simillima, differt frustulis nunquam aut vix striatis. Thwaites, MS.

17. COCCONEMA, Ehrb.

1. COCCONEMA Lunula, Ehrb. Cymbella maeulata, Kütz. p. 79. t. 29. f. 32. c.

HAB. Falkland Islands; on marine Confervæ, Lesson. In a white pigment used by the Fuegians, C. Darwin, Esq.

Found throughout Europe, in Mexico and Chili, the Ægean Sca, the Oregon River, and in earth from Labrador. It also occurs in the white pigment used by the natives of Fuegia, and in atmospheric dust at the Cape de Verd Islands.

18. GOMPHONEMA, Ag.

1. GOMPHONEMA clavatum, Ehrb.; Infus. t. 18. f. vi.

HAB. Falkland Islands; on marine Conferva, Lesson.

Throughout the European and North American shores, those of Iecland, the Marian and Sandwich Islands. Fossil in Virginia, U.S.

2. GOMPHONEMA minutissimum, Grev. in Hook. Brit. Fl. vol. ii. p. 209. G. curvatum, β. salinum, Kütz. p. 85. t. 8. f. l.

HAB. Falkland Islands; on marine Confervæ; Lesson.

A British and common Atlantic species; found also in the Oregon territory and fossil in Virginia, U.S.

19. PINNULARIA, Ehrb.

1. PINNULARIA borealis, Ehrb.

HAB. Cockburn Island; in the soil of a Penguin rookery. In a pigment used by the Fuegians; C. Darwin, Esq.

Found also in Peruvian guano, in the *Mastodon* earth of the Plate river, and in the Cape de Verd atmospheric dust. Also in voleanie ashes from Ascension Island and Patagonia; in voleanie tuff of the Rhine and Phonolite stone.

2. PINNULARIA peregrina (?), Ehrb. Navicula peregrina, Kütz.

HAB. Cockburn Island; in the dirt of a Penguin rookery. Falkland Islands, Lesson.

The true *P. peregrina* is a native of the open ocean near the Brazilian coast, and has been found fossil in Virginia, U. S., and living at St. Domingo, Cuba, and Labrador.

20. NAVICULA, Bory.

1. NAVICULA elliptica, n. sp. Ehrb. Schrift. Berl. Akad. May, 1844.

HAB. Graham's Land; in mud from 270 fathoms.

2. NAVICULA amphioxys, Ehrb. Kütz. p. 91. t. 28. f. 37.

HAB. Falkland Islands; on marine Confervæ, Lesson.

This has also been collected living in Chili, Cayenne and Cuba, in various alluvial deposits, as the Brazils, Iceland, and in the natural paper of Silesia.

3. NAVICULA Didyma (?), Kütz.

HAB. Falkland Islands; on marine Confervæ, Lesson.

The true N. Didyma is a native of a salt-water Lagoon in Germany.

4. NAVICULA Lyra, Ehrb. Kütz. p. 94. t. 28. f. 55, c.

HAB. Falkland Islands; on marine Confervæ, Lesson.

5. NAVICULA viridis, Kütz. p. 97. t. 4. f. 18. and t. 30. f. 12.

HAB. Falkland Islands; abundant on marine Conferva, Lesson, J. D. H.

One of the most widely dispersed of all *Diatomacea*, found alive in fresh waters of England, Scotland, and Ireland, in the natural paper of Silesia, in the Sandwich and Marian Islands, and West Tropical Africa, also in alluvial deposits of Iceland, Labrador and Peru.

21. STAUROPTERA, Ehrb.

1. STAUROPTERA aspera, Ehrb. Infus. Amer. p. 134. t. 1. Kütz. p. 106. t. 12, c.

HAB. Graham's Land; in mnd from 270 fathoms. Falkland Islands, Lesson.

This has been collected in Norway, Spitzbergen, Iceland and Labrador, Mexieo, Cuba, Peru; on the sand-hills of Patagonia, and in Peruvian guano.

2. STAUROPTERA capitata, n. sp. Ehrb. Schrift. Berl. Akad. May, 1844.

HAB. Cockburn Island; on the ground in a Penguin rookery.

22. AMPHORA, Ehrb.

1. AMPHORA Libyea, Ehrb. Kütz. p. 107. t. 29. f. 28, c.

HAB. Graham's Land; in mud from 270 fathoms.

Originally detected in the oasis of Sivah, and since found in various quarters of the globe, as leeland, Labrador, the Oregon River and United States; at the Euphrates River, in African guano, and in the volcanic tuff of the Rhine.

FLORA ANTARCTICA.

2. AMPHORA navicularis, Ehrb. ; l. c.

HAB. Falkland Islands; on marine Confervæ: Lesson.

23. ASTEROMPHALOS, N. G. Ehrb.

1. ASTEROMPHALOS Hookeri, Ehrb. Schrift. Berl. Acad. May, 1844. cum ic.

HAB. Victoria Barrier; in Pancake Ice and in mud from 190 fathoms. In a scum floating in the ocean, Lat. 64°S., Long. 160°W.

2. ASTEROMPHALOS Rossii, Ehrb. l. c. cum ic.

HAB. Victoria Barrier and Victoria Land; in Pancake Ice. In scum with the previous species.

3. ASTEROMPHALOS Buchii, Ehrb. l. c. cum ic.

HAB. Victoria Land and Barrier, with the *A. Rossii*, also in mud from 190 fathoms and in a floating scum with the two preceding species.

4. ASTEROMPHALOS Beaumontii, Ehrb. l. c. cum ic.

HAB. Victoria Barrier; in Pancake Icc.

5. ASTEROMPHALOS Humboldtii, Ehrb. l. c. cum ic.

HAB. Victoria Barrier; in Pancake Ice and in mud from 190 fathoms. In a floating scum with A. Hookeri.

6. ASTEROMPHALOS Cuvieri, Ehrb. l. c. cum ic.

HAB. Victoria Barrier; in Pancake Ice and in mud from 190 fathoms.

7. ASTEROMPHALOS Darwinii, Ehrb. l. c. cum ic.

HAB. In a scum floating in the ocean, in Lat. 64° S., Long. 160° W.

24. HALIOMYX, N. G. Ehrb.

1. HALIOMYX senarius, Ehrb. in Schrift. Berl. Akad. May, 1844. HAB. Victoria Barrier; in the Pancake Ice.

2. HALIOMYX duodenarius, Ehrb. l. c.

HAB. Victoria Barrier; in Pancake Ice. In the stomachs of Salpæ taken in Lat. 64°S., Long. 157°W.

25. HEMIAULUS, N. G. Ehrb.

1. HEMIAULUS Antarcticus, Ehrb. in Schrift. Berl. Akad. May, 1844.

HAB. Victoria Barrier and Victoria Land; in Pancake Ice and in mud from 190 fathoms. Open sea off Cape Horn, Mr. Schayer (Ehrb.).

2. HEMIAULUS? obtusus, Ehrb. 1. c.

HAB. In a floating scum, Lat. 64°S., Long. 160°W. Graham's Land ; in mud from 207 and 270 fathoms.

2 D

26. SYMBOLOPHORA, Ehrb.

1. SYMBOLOPHORA? Microtrias, n. sp. Ehrb.; Schrift. Berl. Akad. May, 1844.

HAB. Victoria Barrier; in Pancake Ice and in mud from 190 fathoms. Graham's Land; in mud from 270 fathoms.

2. SYMBOLOPHORA? Tetras, n. sp., Ehrb.; l. c.

HAB. Victoria Barrier; in Pancake Ice and in mud from 190 fathoms. Graham's Land; in mud from 270 fathoms.

3. SYMBOLOPHORA? Pentas, n. sp., Ehrb. l. c.

HAB. Victoria Barrier; in Pancake Ice and in mud from 190 fathoms. Graham's Land; in mud from 270 fathoms.

4. SYMBOLOPHORA? Hexas, n. sp., Ehrb. l. c.

HAB. Victoria Barrier; in Pancake Ice and in mud from 190 fathoms. Graham's Land; in mud from 270 fathoms.

27. CHÆTOCEROS, N. G. Ehrb.

1. CHÆTOCEROS Dichaeta, Ehrb.; Schrift. Berl. Akad. May, 1844.

HAB. In a scum floating on the surface of the ocean, in Lat. 64° S., Long. 160° W.

2. CHETOCEROS Tetrachæta, Ehrb. l. c.

HAB. In a floating scum with the former species.

Two other species of this new genus have recently been discovered in the Bermuda Islands.

28. ANAULUS, N. G., Ehrb.

1. ANAULUS scalaris, Ehrb.; Schrift. Berl. Akad. May, 1844. HAB. Graham's Land; in mud from 207 and 270 fathoms.

29. RHAPHONEIS, N. G. Ehrb.

1. RHAPHONEIS *fasciolata*, Ehrb.; *l.c.* HAB. Graham's Land, in mud from 207 and 270 fathoms.

2. RHAPHONEIS scutellum, Ehrb.; l. c.

HAB. Cockburn Island; in the dirt of a Penguin rookery.

30. PODOSPHENIA, Ehrb.

1. PODOSPHENIA cuneata, Ehrb.; Infus. t. 17. f. viii. Kutz. Kieselsch. Diat. p. 121. t. 9. f. 13. 1-4.

FLORA ANTARCTICA.

HAB. Graham's Land; in mud from 270 fathoms.

Also found in the Atlantic, German, and Mediterranean seas, and in Peruvian guano.

31. GRAMMATOPHORA, Ehrb.

1. GRAMMATOPHORA parallela, Ehrb.

HAB. Graham's Land; in mud from 207 and 270 fathoms.

Found fossil in Sicily, Oran and Virginia, U.S.

2. GRAMMATOPHORA Africana, Ehrb.; Kütz. p. 129.

HAB. Graham's Land; in mud from 270 fathoms.

Exists also in the Mediterranean Sea and German Ocean; and fossil in Sicily, Oran, and Virginia, U.S., and in Peruvian and African? guano.

3. GRAMMATOPHORA serpentina, n. sp., Ehrb.; Schrift, Berl. Akad. May, 1844. (non Kütz.)

HAB. Graham's Land; in mud from 270 fathoms.

4. GRAMMATOPHORA Oceanica, Ehrb. G. marina, Kütz. Diatoma marinum, Lyngb. Harv. Brit. Alg. p. 201.

HAB. Falkland Islands; on marine Conferra: Lesson.

Abundant on the Atlantic shores of both the Old and New World, and in the Mediterranean Sea. Found fossil in Virginia, U.S., and Peruvian guano; also in the atmospheric dust of the Cape de Verds.

5. GRAMMATOPHORA stricta, Ehrb.; Kütz. p. 129. t. 29. f. 76. c.

HAB. Falkland Islands; on marine Confervæ: Lesson.

This occurs in Peruvian guano, and in deposits at Vera Cruz, in Mexico, and North America.

32. COSCINODISCUS, Ehrb.

1. COSCINODISCUS? actinochilus, n. sp., Ehrb.; Schrift. Berl. Akad. May, 1844. HAB. Victoria Barrier; in Pancake Ice.

2. CosciNodiscus Apollinis, n. sp., Ehrb.; l. c.

HAB. Victoria Barrier; in Pancake Ice and in mud from 190 fathoms. Stomachs of Salpæ, in Lat. 64°S., Long. 157°W. Graham's Land; in mud from 207 and 270 fathoms.

3. Coscinodiscus cingulatus, n. sp., Ehrb.; l. c.

HAB. Victoria Barrier; in Pancake Ice. Graham's Land; iu stomachs of Salpæ with the preceding.

4. Coscinodiscus? gemmifer, n. sp., Ehrb.; l. c.

HAB. Victoria Land; in mud from 190 fathoms, and in Pancake Ice. Graham's Land; in the stomachs of Salpa with the preceding.

This species has also been found in the Bermuda Islands, and fossil at Oran and Sicily.

5. Coscinodiscus Luna, n. sp., Ehrb.; l. c.

HAB. Victoria Barrier and Land; in Pancake Ice and in mud from 190 fathoms. In stomachs of Salpa, taken in Lat. 66°S., Long. 157°W. Graham's Land; in mud from 207 fathoms.

6. COSCINODISCUS eccentricus, Ehrb.; Leb. Kr. p. 66. Kütz. Kiesel. Bacill. p. 131. t. 1. f. 9.

HAB. Victoria Barrier; in Pancake Ice.

Found on the European shores of the Atlantic, and at Vera Cruz; in deposits at Oran, Bermuda, Virginia, U.S., and in Pernvian and African? guano.

7. Coscinodiscus limbatus, Ehrb.; Schrift. Berl. Akad. 1840. Kütz. l. c. p. 131.

HAB. Victoria Barrier; in mud at 190 fathoms.

Also found in the Ægean sea.

8. Coscinodiscus lineatus, Kütz. p. 131. t. 1. f. 10.

HAB. Victoria Barrier and Land; in Paucake and Brash Ice, also in mud from 190 fathoms. Stomachs of *Salpæ* within the Antarctic circle. In a floating scum Lat. 64° S., Long. 160° W. Graham's Land; in mud from 270 fathoms.

A very widely dispersed species, inhabiting Mclville Island, Sicily, Virginia, Maryland, and Peruvian guano.

9. Coscinodiscus Oculus-Iridis, Ehrb.; Leb. Kr. l. c. Kütz. l. c. p. 132.

HAB. Victoria Land and Barrier; in Pancake Ice. Graham's Land; in mud from 270 fathoms.

Found in the Atlantic Ocean, Bermudas, Mcditerranean Sea, and in Peruvian guano.

10. Coscinodiscus radiolatus, Ehrb.; Kütz. p. 132. t. 1. f. 18.

HAB. Victoria Barrier and Land; in Pancake Ice, and in mud from 190 fathoms. Graham's Land; in mud from 207 fathoms.

This occurs in the Mediterranean Sea, as also in deposits at Oran, Sicily, the Bermudas, the United States, Peru and Cuba.

11. COSCINODISCUS subtilis, Ehrb.; Schrift. Berl. Akad. Feb. 1844. Kütz. l. c. p. 132. t. 1. f. 16.

HAB. Victoria Land and Barrier; in Pancake Ice. Stomachs of Salpæ and oceanic scums within the Antarctic circle. Graham's Land; in mnd from 270 fathoms.

Previously found in deposits only, as in Sicily, the Bermudas, the United States, the Mastodon earth of the Plate river, Vera Cruz, and Peruvian and African guano.

12. COSCINODISCUS velatus, Ehrb.; Schrift. Berl. Akad. Feb. 1844.

HAB. Victoria Barrier; in Pancake Ice. Graham's Land; in und from 207 fathoms.

Known previously only in the fossil deposits of Virginia and Maryland, U.S.

33. FLUSTRELLA, Ehrb.

1. FLUSTRELLA concentrica, Ehrb.; Schrift. Berl. Akad. Feb. 1844.

HAB. Victoria Barrier; in Pancake Ice. Graham's Land; in mud from 270 fathoms.

In a fossil state this species occurs in Sicily, Oran, the Ægeau Sea, Maryland, U.S, and in the Bermuda Islands.

34. ACTINOCYCLUS, Ehrb.

1. ACTINOCYCLUS senarius, Ehrb.; Schrift. Berl. Akad. June, 1844. HAB. Falkland Islands; on marine Confervæ : Lesson.

35. ACTINOPTYCHUS, Ehrb.

1. ACTINOPTYCHUS biternarius, Ehrb.; l. c.

HAB. Victoria Barrier; in Pancake Ice.

Also occurs in tertiary deposits in Virginia and Maryland, U.S., and in the Bermuda Islands.

36. DISCOPLEA, Ehrb.

1. DISCOPLEA Rota, n. sp., Ehrb.; Schrift. Berl. Akad. May, 1844. HAB. Graham's Land; in mud from 270 fathoms.

2. DISCOPLEA Rotula, n. sp., Ehrb.; l.c.

HAB. Graham's Land; in mud from 270 fathoms.

37. LITHOBOTRYS, Ehrb.

1. LITHOBOTRYS? denticulata, n. sp., Ehrb.; l. c.

HAB. Victoria Barrier; in Pancake Icc and in mud from 190 fathoms. In a floating scum, Lat. 64° S., Long. 160° W.

The only other species of this genus is a Virginian fossil.

38. LITHOCAMPE, Ehrb.

1. LITHOCAMPE Australis, n. sp., Ehrb.; Schrift. Berl. Akad. May, 1844.

HAB. Victoria Barrier; in Pancake Ice.

This genus was known only as a fossil, occurring in the United States and the Mediterranean Sea, previous to the detection of this and the following species.

LITHOCAMPE Antarctica, n. sp., Ehrb.; l. c.

HAB. Graham's Land; in mud from 270 fathoms.

39. TRIAULACIAS, N.G., Ehrb.

I. TRIAULACIAS triquetra, Ehrb.; Schrift. Berl. Akad. May, 1844. HAB. Victoria Barrier; in mud from 190 fathoms.

40. BIDDULPHIA, Gray.

1. BIDDULPHIA ursina. n. sp., Ehrb.; l. c.

HAB. Graham's Land; Gulf of Erebus and Terror, in mud 207 and 270 fathoms.

6 E

41. ZYGOCEROS, Ehrb.

1. ZYGOCEROS Australis, n. sp., Ehrb.; l. c.

HAB. Victoria Barrier; in Pancake Ice. Graham's Land; Gulf of Erebus and Terror, in mud at 207 fathoms.

42. DENTICELLA, Ehrb.

1. DENTICELLA lævis, n. sp., Ehrb.; l. c.

HAB. Graham's Land; Gulf of Erebus and Terror, in mud at 270 fathoms.

43. MESOCENA, Ehrb.

MESOCENA? Spongolithis, n. sp., Ehrb.; l. c.
 HAB. Victoria Barrier; in mud from 190 fathoms. Graham's Land; in mud at 270 fathoms.

44. ACTINISCUS, Ehrb.

1. ACTINISCUS lancearius, n. sp., Ehrb.; l. c.

HAB. Open ocean, Lat. 66° S., Long. 157° W., in the stomach of a Salpa.

45. DICTYOCHA, Ehrb.

1. DICTYOCHA aculeata, Ehrb.; Leb. Kr. p. 68. Kütz. Kiesel. Bacill. p. 140.

HAB. Victoria Land and Barrier; in Pancake Ice. In the stomachs of *Salpæ*, Lat. 66° S., Long. 157 W. Entangled in a surface scum, Lat. 64° S., Long. 160° W.

Found living in the North Sea, fossil in Sicily, the Ægean, North Africa, and Virginia, U.S.

2. DICTYOCHA binoculus, Ehrb.; Schrift. Akad. Berl. May, 1844.

HAB. Victoria Barrier; in Pancake Ice. Entangled in a floating scum, Lat. 64° S., Long. 160° W. Also occurs in tertiary deposits in the Ægean Sea.

3. Diстуосна *biternaria*, Ehrb.; *l. c.* Нав. Victoria Barrier; in Pancake Ice.

4. DICTYOCHA Epiodon, Ehrb.; Schrift. Berl. Akad. Feb. 1844.

HAB. Victoria Barrier; in Pancake Ice.

Originally described from specimens occurring in a tertiary deposit in Virginia, U.S., also found in Peruvian guano.

5. DICTYOCHA octonaria, Ehrb.; Schrift. Berl. Akad. May, 1844.

HAB. Victoria Barrier; in Pancake Ice.

FLORA ANTARCTICA.

6. DICTYOCHA Ornamentum, Ehrb.; l. c. Feb. 1844.

HAB. Victoria Barrier; in Pancake Ice. In a floating scum, Lat. 64° S., Long. 160° W. Tertiary deposits of Sicily contain this species.

7. DICTYOCHA septenaria, Ehrb.; l.c.

HAB. Victoria Barrier; in Pancake Ice, and in mud from 190 fathoms.

Previously found fossil in the tertiary deposits of Oran.

8. DICTYOCHA Speculum, Ehrb.; Kutz. Kiesel. Bacill. p. 140. t. 21. f. 22. c.

HAB. Victoria Barrier; in Pancake Ice, and in mud at 190 fathoms. In the stomachs of Salpee, Lat. 66° S., Long. 157° W. In a floating scum, Lat. 64° S., Long. 160° W.

A widely distributed species, found living in the North Sca and Atlantic, fossil at North Africa, Greece, and Sicily, and in Maryland, U.S.

46. RHIZOSOLENIA, Ehrb.

1. RHIZOSOLENIA Calyptra, n. sp., Ehrb.; Schrift. Berl. Akad. May, 1844.

HAB. Victoria Barrier; in Pancake Ice. Graham's Land; Gulf of Erebus and Terror, in mud from 270 fathoms.

2. RHIZOSOLENIA Ornithoglossa, n. sp., Ehrb.; l. c.

HAB. Victoria Barrier; in Pancake Ice, and in mud from 190 fathoms. Graham's Land; Gulf of Erebus and Terror, in mud at 270 fathoms.

Both these species have recently been detected in the Bermuda Islands.

LVII. DESMIDIEÆ, auct. recent.

1. ARTHRODESMUS, Ehrb.

1. ARTHRODESMUS Tania, Ehrb.; Schrift. Berl. Akad. June, 1841. HAB. Falkland Islands; on marine Conferva: Lesson.

LVIII. LICHENES,* L.

1. USNEA, Ach.

1. USNEA melaxantha, Ach.; Lich. Univ. p. 618. Syn. Meth. p. 303. Brown, Plant. Spitz. in Scoresby Voy. vol. i. App. p. 76. Kunth. Synops. vol. i. p. 36. D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 596.

* Since the publication of the "Lichenes Antarctici" in the 'London Journal of Botany,' and of those contained in the first part of this work, I have had the advantage of re-studying all the species with my friend, the Rev. Churchill Babington, whose profound knowledge of the forms of this difficult order, and acquaintance with the most recent writings of European Lichenologists, have been most liberally brought to bear upon this part of the 'Flora Antarctica.'

Bory in Duperrey, Voy. Bot. p. 240. Mont. in Voy. au Pole Sud, Bot. Crypt. p. 201. U. sphacelata, Brown, in Parry, 1st Voy. App. p. 307. Hook. Plant. Arct. in Linn. Soc. Trans. vol. xiv. p. 384. Bot. Miscell. vol. i. p. 15. t. 12. U. fasciata, Torrey, in Silliman Journ. vol. vi. cum ic. Hook. Bot. Miscell. vol. i. p. 14. t. 11. U. aurantiaco-atra, D'Urv. in Mém. Soc. Linn. Paris, vol. iv. p. 596. Cornicularia flavicans, Persoon, in Freye, Voy. Bot. p. 210. Lichen aurantiaco-ater, Jacq. Miscell. (fid. Acharius).

Var. a. Acharii; robusta, thallo scabrido, apotheciorum marginibus nudis .-- U. melaxantha, Ach. I. c.

Var. B. Jacquinii; robusta, thallo lævi.—Lichen aurantiaco-ater, Jacq. l. c.

Var. y. fasciata; robusta, thallo tuberculato, ramulis pluries divisis capillaceo-attenuatis, apotheciis extus tuberculatis.—U. fasciata, Torrey, I. c.

Var. 8. sphacelata; gracilis, thallo lævi pruinoso v. tuberculato fruticuliformi ramosissimo, ramulis capillaribus.—U. sphacelata. Brown, l. c.

HAB. Throughout Tierra del Fuego and the Falkland Islands; on exposed rocks, from the sea to 2,000 feet, most abundant. New Sonth Shetlands; *Webster*, Dr. Eights.

Perhaps the handsomest of all Lichens, whether we regard its colour, stature, or mode of growth, and yet so variable in all these points, that the extremes are scarcely recognizable. In size it varies from a few lines or an inch, with capillary stem and branches, to four inches, with a woody stem a quarter of an inch in diameter. The colour is sometimes wholly (especially in var. δ) black, at others uniformly yellow, but most frequently banded with black, especially towards the apices of the ramuli. Old specimens turn tawny red, as do all when, after being dried, they are soaked in warm water. The apothecia are yellow, grey or deep black. The surface of both the stems and the back of the apothecia is more or less tuberculated or pruinose, smooth or much wrinkled, naked or more or less covered with longer or shorter horizontal terete acuminated ramuli. The apothecia vary exceedingly in breadth, from two or three lines to almost half an inch; their margins are smooth, tuberculated or beset with branching ramuli; they are generally terminal, but the younger appear lateral when subtended by a branch.

It is in the Falkland Islands that this species most abounds, covering the surface of the Quartz rocks with a miniature forest, seeking the most exposed situations, and there attaining its greatest size and beauty. In these Islands, too, all the five varieties I have enumerated may be collected within a few feet of one another, and so associated as to leave little doubt that they are states depending on age, rather than marked races. The var. δ . especially, is certainly only an undeveloped state, which does not bear soredia in the Antarctic regions; or apothecia either in Tasmania or in the Arctic latitudes, where soredia are produced.

The structure of the stem of this, and probably of some other Lichens, presents a marked analogy with that of the Laminarioid Algæ described at p. 460. The central thread is very large, composed of concentric layers of dense, horny, red cellular tissue, gradually passing into a soft white pith, enclosing a cavity. Around this horny thread, whose edges are sharply defined, are arranged concentric layers of a spongy cellular substance, which again are enclosed in a cortical layer, as dense as the central, and to which the yellow and black hues of this Lichen are always confined. Thus, proceeding from the circumference, there is—1st, a horny, coloured, cortical layer, answering to what is called the cortical layer of *Lessonia*, and to which, in that plant also, the coloured chromule is chiefly confined; 2nd, the layers of intermediate lax tissue, successively deposited, though much more obscurely so than in *Lessonia*; 3rd, the central thread of *Usneæ* which is a stout axis, answering to the elliptic core of *Lessonia*, but in this Lichen becoming so lax towards the centre as to enclose a cavity in the older stems.

We have never observed spores in any specimens of this Lichen, from whatever place collected, though we have examined very many apothecia in a live state, as well as after being dried, both young and old, and of all colours, both black, grey, or yellow. Dr. Montagne (Voy. au Pole Sud, l. c.) has been similarly unsuccessful.

521

The range of this Lichen is very wide. It is found in the Arctic regions of both the New and Old World; on the Andes under the Equator, at an elevation of 11,000 feet; on the mountains of Tasmania at 3,000 and 4,000 feet; in Chili, the Falkland Islands, Fuegia, and the New Sonth Shotlands. Now it is worthy of notice, that in none of the Arctic, the equatorial, or south temperate latitudes, does this plant produce apothecia; and that in the Antarctic, where alone apothecia have hitherto been found, these are always barren. Further it is remarkable, that this Lichen grows only where no other Usnea is found in fruit; and is, perhaps, the only species of that genus which universally inhabits rocks; circumstances which, taken together with its increasing in luxuriance with the exposure it is subjected to, suggests the possibility of its being a state of some other species of this highly variable and universally diffused genus, and that, distinct as the Autarctic specimens of U, melaxantha appear, they may owe their characters to the climate, for there is very great difficulty in defining the species so as to exclude states of U. florida. In South Chili, where the U. florida commences (proceeding southwards), we have specimens of U. ceruchia, Mont., which are, perhaps, states of U. melaxantha. Again, in Tasmania I am unable to distinguish some specimens of U. barbata and U. florida (?), which grow on the trunks of dead trees in the higher parts of the island, from the U. melaxantha of the summit of Mount Wellington and other elevated mountains.

2. USNEA Taylori, Hook. fil.; thallo erecto lævi pallide citrino dichotomo, ramis erectis gemmis papulosis sparsis, apotheciis terminalibus majoribus demum planiusculis, disco atro, margine integerrimo, dorso lævigato. Lond. Journ. Bot. vol. iii. p. 657. (TAB. CXCV. Fig. I.)

HAB. Kerguelen's Land; on rocks from 100 to 1,200 feet of elevation; very abundant.

U. melaxanthæ simillima, differt præcipue thallo lævi polito et colore lætiore nitente.

So closely allied to the U. melaxantha, that I advance this species with much hesitation. None of the Kergnelen's Land specimens exactly tally with any of its congeners from other Antarctic localities, but approach them very nearly indeed; so much so, that the present should be perhaps regarded as a permanent variety only. If it he so, it is singular that it is the only one in which asci have hitherto been detected; these are abundant in all the apothecia, and vary much in size, in the form of their contained spores, and in the arrangement or grouping of these, as shown in the accompanying plate.

This is by far the most handsome vegetable production of Kerguelen's Land.

PLATE CXCV. Fig. I.-1, young, and 2, full grown plants of the natural size; 3, vertical section of young and 4, of old apothecium; 5, lamina proligera, with asci and spores :- very highly magnified.

3. USNEA barbata, Ach.; Lich. Univ. p. 624. Fl. Antarct. Pt. 1. p. 194.

Var. c. articulata, Ach.; Syn. Meth. p. 306.

HAB. Hermite Island, Cape Horn, and the Falkland Islands; abundant on the stems of Empetrum and in heathy and rocky places. Var c. Falkland Islands, Mr. Wright. Hermite Island, Cape Horn; top of Mount Foster and Kater's Peak, &c.

Also a native of Lord Auckland's group, Campbell's Island, and Tasmania. In Europe it is found as far north as Lapland.

4. USNEA plicata, Ach.; Lich. Univ. p. 622. Fl. Antarct. Pt. 1. p. 194.

HAB. Falkland Islands, and Hermite Island, Cape Horn; on twigs of bushes, &c. Strait of Magalhaens, Jacquinot.

A more Arctic and Antarctic plant than the following, reaching Cape Horn in 57°S., and the shores of the Polar Sea in Arctic America, or 69° N., beyond which it is succeeded by the U. melaxantha in both extremities of the globe.

5. USNEA florida, Ach.; Lich. Univ. p. 304. Engl. Bot. t. 872.

HAB. Chonos Archipelago, on trees; C. Darwin, Esq.

A plant evidently impatient of perennial cold; inhabiting none of the Antarctic islands south of New Zealand and Tasmania in the Old World, nor of the Chonos Archipelago in the New. In the northern regions, again, it does not pass the wooded regions (63° N.), in America, or reach Lapland in Europe.

2. EVERNIA, Ach.

1. EVERNIA Magellanica, Mont.; in Voy. au Pole Sud, Bot. Crypt. p. 198.

HAB. Strait of Magalhaens; D'Urville and Jacquinot.

A very beautiful and distinct species, which I have seen from no other locality than that indicated by Montagne.

3. RAMALINA, Ach.

1. RAMALINA scopulorum, Ach.; Lich. Univ. p. 604. Engl. Bot. t. 688.

Var. a. thallo $1-1\frac{1}{2}$ pollicari lineari rigido polito utrinque glabro, apotheciis plurimis primum concavis demum convexis marginibus reflexis. R. verrucosa, *nob. in Lond. Journ. Bot.* vol. iii. p. 655.

Var. β . thallo fastigiato sub 2-pollicari cartilagineo pallide stramineo lineari v. lineari-obovato laciniato polito obscure lacunoso hic illic terebrato apicibus sub-acutis, apotheciis nullis.

Var. γ . thallo elongato 2–6-unciali rigide cartilagineo planiusculo v. lacunoso parce pruinoso pertuso laciniis $\frac{1}{2}$ unc. latis lobulis acutis, apotheciis nullis.

Var. δ. thallo flavo dense fastigiato lineari-ligulato 1–5-unciali laciniis flaccidis nunc pertusis pluries divisis acuminatis glabratis punctisve pruinosis sparsis, apotheciis nullis.

Var. ϵ . omnia varietatis δ ., sed rigida, apotheciis apices versus laciniarum confertis corrugatis plerisque monstrosis.

Var. ζ . *tercbrata*; thallo elongato 8 unc. ad pedalem flaccido lacunoso et corrugato pertuso pruinoso v. glabrato, margine integerrimo eroso lobulato v. prolifero, apotheciis nullis:—inter var. γ . and δ . media sed statura proceriore. R. tcrebrata, *nob. in Lond. Journ. Bot.* vol. iii. p. 654.

Var. η . truncata; fastigiata, thallo cuneato in lacinias breves truncatas flabellatim expanso laciniis pertusis pruinosis griseis apicibus involutis :----an forma incompleta?

HAB. Var. α . Dry quartz rocks, Falkland Islands. Var. β . and γ . Dry rocks, chiefly of clay slate, Falkland Islands. Var. δ . Falkland Islands and Cape Horn; on rocks near the sea. Var. ϵ . Dry granite rocks, Cape Horn; Kerguelen's Land, *Anderson*. Var. ζ . Falkland Islands; on moist exposed rocks, near the sea, most abundant. Var. η . with var. ζ . but in more shady places.

On these varieties, or rather forms, we are enabled to offer the following observations :----

The var. a. is exactly the English R. scopulorum, and the only one which produces perfect fruit in the Falkland Islands. Var. β . is found in New Zealand, and in fruit; also abundantly in Brazil. Var. γ . and ζ . are scarcely distinguishable from R. fraxinea of Norfolk: small speciments exactly resemble R. membranacea, Laurer, of New Holland; it is a very common form in the Southern Hemisphere. Var. δ . is very similar to the Uraguay R. prolifera, of Taylor. Var. ϵ . is a small form of a Chilian variety. The var. η . inhabits Chiloe, and approaches near to R. pollinaria. Dissimilar though the states of this plant here brought together are, no one who has collected them together can doubt their all belonging to one species, which, however, seldom fruits.

M. Fries states his conviction of the probability that all the European Ramalinæ are varieties of one species, an opinion in which we certainly concur, and we would further add many of the Exotic (except R. inanis) to it. The above varieties certainly all belong to one species, as abundant in Cape Horn and Fuegia as the ordinary states of R. scopulorum are in Europe, and, however unlike some of these forms are to the English plant of that name, the one called a here, and which is the only one that fruits, is in no way to be distinguished from that plant. Considering how plastic the Lichens are in form and texture, and how amenable to the different climatal conditions, it must be admitted that if the R. scopulorum of England were to inhabit the maritime rocks of the Falklands, its aspect would be changed; the humidity of the atmosphere near the sea of these islands, being much greater than that of similar situations in our own country. Again, the locality inhabited by the var. a., namely, rocks at a considerable elevation and distance from the occan, possesses a climate more assimilated to the British habitats of R. scopulorum than are the moist rocks at a lower level, and hence it is only natural to suppose, that there the Falkland Island form would assume the English. Lastly, the universally acknowledged difficulty of defining the European species. and the singular abundance of forms of the genus exactly similar to these in all parts of the world, between Lat. 60° N. and 57° S., together with the fact that many other Lichens are equally protean and widely distributed, are all arguments in favour of the Antarctic species having a common origin with other forms of the genus inhabiting the Arctic, Temperate, and Tropical regions.

The genus *Ramalina*, in the Arctic zone, attains the parallel of 69°, on the shores of the Polar Sea in North America, and of Lapland in Europe.

4. CETRARIA, Ach.

1. CETRARIA Islandica, Ach.; Lich. Univ. p. 509. Engl. Bot. t. 1330.

HAB. Hermite Island, Cape Horn; amongst moss on the tops of mountains only, 1,500-1,700 feet, barren.

One of the most Arctic of plants, having been collected on Ross Islet, the northernmost known land in Europe (81° N.), and in Melville Island (76°), on the limits of Arctic American vegetation. It inhabits the level of the ocean only within the Arctic circle, or in the extremely cold plains of Central Russia (as Moscow, 55° N.) Dahuria in Asia, 50° N., and in North America (as Labrador, 55° N.); thence, in progressing south, it ascends; attaining the tops of our Scotch Alps, 4,000 feet (56° N.), about 10,000 feet on the Swiss Alps (46° N.), 9,000 feet on the top of the Pyrenees, and 4,000 feet on the mountains of North Carolina and Virginia (in 36° N.). The last locality is the lowest latitude it attains in the Northern Hemisphere; in the Southern it re-appears only on the extreme point of America, and there is confined to the pinnacles of the very highest mountains. There is perhaps no vegetable common to both hemispheres more typical of extreme cold than this Lichen, which is further interesting from being the reputed eure for consumption, and the only plant of that order extensively used in medicine.

2. CETRARIA glauca, Ach.; Lich. Univ. p. 509. Mont. in Voy. au Pole Sud, Bot. Crypt. p. 194.

HAB. Hermite Island, Cape Horn; top of Mount Kater, 1,700 feet; on rocks, sparingly. Straits of Magalhaens, D'Urville and Jacquinot.

This, again, is an instance of the re-appearance of a Northern and Arctic Lichen in the Sonthern Hemisphere only under Antarctic skies. The *C. glauca* finds its principal parallel in Scotland, central and northern Europe, and sub-arctic America, wandering as far south as the Swiss Alps and mountains of the Canary Islands. It is not nearly so Arctic as the *C. Islandica*, not being found in Spitzbergen or Melville Island, or, according to Richardson anywhere to the northward of 54° in Arctic America. Wahlenberg states it to be rare in Lapland. 3. CETRARIA sepincola, Ach.; Lich. Univ. p. 507. Mont. in Voy. au Pole Sud, Bot. Crypt. p. 195. HAB. Strait of Magalhaens, on bark of Berberis ilicifolia; M. Jacquinot.

4. CETRARIA aculeata, Fries, Lich. Europ. p. 35. Mont. l. c. p. 194. Cornicularia aculeata, Ach. Lich. Univ. p. 612.

HAB. Hermite Island, Cape Horn; on the summit of Kater's peak, and other mountains. Falkland Islands, *Gaudichaud*, &c.; but not common. Strait of Magalhaens; *D'Urville and Jacquinot*.

A very Arctic plant, inhabiting Melville Island in Arctic America, and Lapland in Europe; in the latter continent it extends as far south as the Alps and Pyrenees, and to the Canary Islands in the Atlantic Ocean.

5. NEPHROMA, Ach.

1. NEPHROMA polaris, Ach.; Lich. Univ. p. 523. N. arctica, Mont. in Voy. au Pole Sud, Bot Crypt. p. 192.

HAB. Hermite Island, Cape Horn; moist exposed places on the margins of the woods bordering the sea, abundant. Strait of Magalhaens; *MM. Hombron* and *Jacquinot*.

The most magnificent of Lichens, whether we regard the size of the thallus, which often is a foot and upwards across, the general aspect, or the size and beautiful colour of the frond and copious apothecia. Its European range is very northern, being confined to Scandinavia, reaching Lapland but not the islands beyond, nor inhabiting any countries south of the Baltic. In North America it ranges over the wooded regions and barren lands, $54^{\circ}-69^{\circ}$, and to the west of the Rocky Mountains descends to Fort Vancouver, in Lat 46° N.

2. NEPHROMA cellulosa, Ach.; Lich. Univ. p. 523.

HAB. Staten Land; Menzies (in Hook. Herb.)

A very much smaller species than the former, approaching very closely to the *N. resupinata* of Europe. It also inhabits Juan Fernandez and Tasmania.

6. PELTIDEA, Ach.

1. PELTIDEA polydactyla, Ach.; Lich. Univ. p. 517. Mont. in Voy. au Pole Sud, Bot. Crypt. p. 193.

HAB. Strait of Magalhaens; MM. Hombron and Jacquinot.

Very abundant in Tasmania and New Zealand, also found in Lord Auckland's group and Campbell's Island. In Europe, it ranges from Switzerland to Sweden and Norway; in America, from Mexico to Sitka, but does not proceed so far north on the east of the Rocky Mountains; it also inhabits the West Indies, Colombia, the Cape of Good Hope, and other warm climates. Altogether it is a plant which does not shun the cold so markedly as do either of the following species, for it (the var. *sentata*) is also found as far as the northern limits of the forest regions of Arctic America.

2. PELTIDEA canina, Ach.? Lich. Univ. p. 517. Engl. Bot. t. 2299.

HAB. Falkland Islands; on tufts of Bolax glebaria, rarc.

In the Southern Hemisphere the *P. canina* has been found in Juan Fernandez only. Its range in the Northern is very wide, commencing in Mexico it is dispersed as far north as Canada on the east coast, and Sitka on the west. In Europe it is frequent from the Alps to Lapland, but does not inhabit the Arctic Island of Spitzbergen in Europe, or the shores of the Polar Sea and Islands beyond in America, where the *P. apthosa* abounds, a species not hitherto found in the Antarctic regions. The Falkland Island specimens are in a very imperfect state.

3. PELTIDEA venosa, Ach.? Lich. Univ. p. 514. Engl. Bot. t. 887.

HAB. Kerguelen's Land; on tufts of moss on the hills.

The specimens are in a very unsatisfactory state, being stunted and barren. They, however, closely resemble dwarf Scotch and Arctic individuals of the plant in question, differing chiefly in the smoother thallus and occasional buds. The range of the species is not wide: Lapland and Switzerland are its Northern and Southern European limits. In North America it inhabits the United States, Canada, and the Columbia river. Kerguelen's Land is the only recorded habitat in the Southern Hemisphere.

4. PELTIDEA horizontalis, Ach.; Lich. Univ. p. 515. Engl. Bot. t. 883.

HAB. Christmas Harbour, Kerguelen's Land; on wet moss, abundant.

Both in characters and in locality the specimens agree with the European P. horizontalis.

This, again, has not been observed elsewhere in the Southern Hemisphere. In the Northern it is confined to the temperate latitudes, both of Enrope and North America, ranging in the latter from the middle United States to Canada, and in Europe from Lapland to Switzerland and the Pyrenees, where it is the only species of the genus inhabiting the top of the Pic du Midi (9,000 feet).

7. STICTA, Ach.

1. STICTA crocata, Linn.; Engl. Bot. t. 2110. S. citrina, Pers. in Freye. Toy. Bot. p. 201. S. fuliginosa, nobis in Lond. Journ. Bot. vol. iii. p. 646.

Var. B. gilva, Ach. Synops. Lich. p. 232. S. impressa, quoad exempt. Falkland. nobis in Lond. Journ. Bot. l. c. S. Gaudichaudii, Delise, Monogr. Stict. p. 80. t. vii. f. 23. Bory in Duperrey Voy. Bot. p. 236.

HAB. Strait of Magalhaens; D'Urville, Hombron. Hermite Island, Cape Horn, and the Falkland Islands; both varieties on rocks and twigs of shrubs, near the sea.

This species is also found on the west coast of South America, in Tasmania, Swan River, and the Cape of Good Hope. In the Northern Hemisphere it inhabits the United States and West Indian Islands, the Sandwich group, Great Britain and Ircland, attaining its northern limit at Inverary in Scotland (Lat. 56° N.), which singularly coincides with the latitude of the most southern habitat, namely, Cape Horn. The var. *gilva* is certainly only a variety, without the pulverulent lines on the upper surface. The rimæ are sometimes white in this species, when it becomes exceedingly difficult to distinguish it from some of its congeners.

2. STICTA endochrysa, Delise; thallo late expanso cartilagineo glauco intus aurato glabro lobato, lobis rotundatis subintegerrimis marginibus gemmis confertis auratis obsitis super cæsiis flavo-virescentibusve subter rufo-flavis glabratis, cyphellis parvis prominulis citrinis, apotheciis sparsis concavis stipitatis, disco atro-rufo, margine elevato inflexo primum lanuginoso demum glabrato et crenulato. S. endochrysa, Delise, Monogr. p. 43. t. 1. f. 1. S. D'Urvillei, Delise, l. c. p. 170. S. flavicans, nobis in Lond. Journ. Bot. vol. iii. p. 648. S. ochracea, Menzies, MSS. in Hb. Hook. Parmelia pubescens, Pers. in Freye. Voy. Bot. p. 199. (TAB. CXCV. Fig. II.)

HAB. Hermite Island, Cape Horn; from the sea to the tops of mountains, abundant on rocks and trees. Falkland Islands; very abundant. Staten Land; *Menzies*.

Thallus late expansus, pedalis et ultra, glaucescens. Lobi lati, obscure undulati, flavido-marginati v. immarginati; marginibus isidiophoris granulis subfoliaccis dense onustis, rarius denudatis et crenatis; subter lacunosi v. plani, medium versus fusco-flavidi parce tomentosi, ad apices pallidiores et glabrati. Cyphellæ plurimæ, papillæformes, rimæque thalli citrimæ. Apothecia sparsa, 2-4 lin. lata. Variat colore superficiei superioris flavido v. cæsio, nunc glauco-virescente; foliis planis v. sublacunosis; marginibus loborum planis v. elevatis, lævibus v. crispatis, nudis v. granulis corniculatis dense obsitis.

This, and the Nephroma arctica, are the most noble foliaceous Lichens of the Antarctic regions. The present is particularly so, the brightness of its golden thallus, and its great abundance on the sea-ward edges of the Fuegian forest, rendering it a very conspicuous plant. It is also found in Chiloe and Juan Fernandez, and in the Old World it inhabits New Zealand. Though an ally of *S. aurata*, it is perfectly distinct from that plant, especially in the granular and not powdery margins of the thallus, also in size and general features. It is certainly as well marked as any species of Lichen confined to the southern latitudes.

PLATE CXCV. Fig. II.—1, plant in usual state; 2 and 3, portions of thallus; 4, plant as covered with granular tubercles of the natural size; 5, granular surface of thallus; 6, apothecia; 7, under surface of thallus; 8, lamina proligera; 9, spores:—all highly magnified.

3. STICTA orygmæa, Ach.; Fl. Antarct. Pt. 1. p. 197.

HAB. Staten Land, (fid. Acharius). Strait of Magalhaens, D'Urville.

This species was not seen at Cape Horn or the Falkland Islands: but we have specimens from Chiloe and Juan Fernandez. It is singular that a Lichen, which abounds in Lord Auckland's group and Campbell's Island, should not be also found at Cape Horn; and that the *S. endochrysa* of Cape Horn is found in New Zealand, but does not spread so far south as Lord Anckland's group. The much greater cold of Fuegia might be prejudicial to the *S. orygmæa*, but there is no apparent reason for *S. endochrysa* being foreign to Lord Auckland's group. I suspect that the Acharian plant from Staten Land, referred to as *S. orygmæa*, is the *S.endochrysa*, Del.; and that the specimens were received from Menzies.

There are very remarkable differences between the Lichens of those two divisions of Antarctic botany which are here adopted: the most salient features of which consist in the presence or abundance of the following Lichens in only one of the divisions.

LORD AUCKLAND'S GROUP AND CAMPBELL'S ISLAND.

	Usnea melaxantha.
Ramalina inflata.	Ramalina scopulorum.
Stereocaulon ramulosum.	Cetraria Islandica.
Argus.	Nephroma polaris.
Sticta orygmæa.	Stereocaulon corallinum.
Menziesii.	Sticta endochrysa.
— Delisea ?	crocata.
——— faveolata.	Freycinetii.

CAPE HORN.

This remarkable predominance of certain very widely distributed forms in the southern extremity of the Western Hemisphere, and the absence of the same in similar positions in the Eastern, admit of no explanation, beyond what climate will afford.

4. STICTA obvoluta, Ach.; Lich. Univ. p. 452 (vix Delise). S. hirsuta, Mont. in Voy. au Pole Sud, Bot. Crypt. p. 188. t. 15. f. 2.

HAB. Staten Land; Menzies. Strait of Magalhaens; M. Hombron.

Of this we possess an anthentically named specimen, gathered and labelled by Menzies himself: in it the cyphellæ are pale yellow; as Montagne describes those of his *S. hirsuta*. M. Delise does not seem to have understood the species, and describes apparently a very different plant under this name. It is closely allied to the *S. cometia* of Peru, in which the apothecia are fringed with much longer hairs, and also to the *S. Humboldtii*, Hook., another Peruvian plant with fringed apothecia; but both of these have white cyphellæ.

Sticta obvoluta is also a native of South Chili and Juan Fernandez.

5. STICTA Billardieri, Delise, Monogr. Stict. p. 99. t. 8. f. 35. S. Richardi, nobis in Fl. Antarct. Pt. 1. p. 198. (non Mont.)

Var. B. lobis thalli subter pallidioribus. S. divulsa, Tayl. in Lond. Journ. Bot. vol. vi. p. 182.

HAB. Chonos Archipelago; C. Darwin, Esq.

A New Zealand and Tasmanian species, agreeing remarkably well with the figure and description of Delise, except in the under surface being more pale than is described by that author. It differs from *S. faveolata*, Delise, in the much longer and narrower lobes and paler under surface, and in the absence of the granulations on the upper. The apothecia are all marginal when young, deeply concave, the older in these specimens convex, with a very narrow margin: this convexity is not a character to which too much importance should be attached, because the apothecia of many allied species vary extremely on the same specimen, from concave to convex; and these differences do not in such species depend on age.

This is the S. Richardi of the first part of this work, but not of Montagne, the true S. Richardi having much broader lobes to the thallus, more or less rough or even scrobiculate on the upper surface, and almost wholly smooth on the under. It is very nearly allied to the true S. carpoloma, but differs in the white cyphellæ, and like many others of the faveolate group it is possibly only a variety of S. damæcornis.

We have specimens from the Chonos Archipelago, from Chiloe and the Island of Huaffa (on the coast of South Chili), agreeing entirely with one another in the (when dry) pale yellow-brown shining colour of the upper surface. The New Zealand and Lord Auckland's Island examples are rather paler: in all, the under surface of the lobes is naked and tawny yellow towards the apex.

6. STICTA faveolata, Delise; Monogr. Stict. p. 101. t. 8. f. 36. Mont. in Voy. an Pole Sud, Bot. Crypt. p. 186.

HAB. Strait of Magalhaens; Hombron.

We have thought it possible that what we have called *S. Billardieri*, Del., may be the *S. faveolata* of Montagne (in Voy. au Pole Sud); especially as that author makes no mention of the granular lines on the upper surface so expressly alluded to by Delise (under *S. carpoloma*); and also from this *S. faveolata* of Montagne being found by the officers of the French South Polar Expedition, both in the Strait of Magalhaens and Lord Auckland's group. In the absence, however, of authentic specimens, and of any explanatory note by Montagne (whose specimens were very insufficient) on the species he calls *S. faveolata*, we are obliged to abide by the published figures of Delise.

The lacunose fronded *Stictæ* with white soridia, form a most natural group of species, so inextricably confounded by nature, if they be really and truly species at all, that we have failed in discovering limits to the variations of any of them. Even the colour of the cyphellæ is very unsatisfactory, there being, between the snowwhite of the *S. Billardieri* and dirty yellow of some others, all intermediate tints; and in the same species, almost on the same specimen, of Tasmanian examples of *S. faveolata* (?), it is impossible to say whether the cyphellæ are dirty white or pale yellow; whilst other specimens, in all respects similar, have their colour well pronounced.

7. STICTA variabilis, Ach.; Delise, Monogr. Stict. p. 119. t. 11. f. 48. Mont. in Voy. au Pole Sud, Bot. Crypt. p. 185.

HAB. Strait of Magalhaens; Jacquinot.

Possibly these specimens are referable to a state of that highly variable plant which we have referred to the *S. Fregcinetii*, Del.

8. STICTA Thouarsii, Delise, Monogr. Stiet. p. 90. t. 8. f. 29. S. scrobiculata, nobis in Lond. Journ. Bot. vol. iii. p. 646 (non Ach.).

HAB. Hermite Island, Cape Horn, and the Falkland Islands; on rocks, &c. Tristan d'Acunha; Petit Thouars.

The specimens examined, when preparing the short notice of the Antarctic Lichens for the London Journal of Botany, were very imperfect; and their under-surface exhibiting no cyphellæ, we referred them to the *S. scrobiculata*, which they considerably resemble, especially in colour, and in their fetid scent when moistened. Other specimens showed white cyphellæ in abundance, and allied the plant so closely to the European *S. limbata*, that we can detect no marked difference between them, beyond what is afforded by the colour of the powdery granulations on the surface.

Delise's description of S. Thouarsii leaves no doubt in our mind of this being his plant. The apothecia are unknown. Fuegian specimens are of a paler colour than the Falkland Island ones.

9. STICTA Freycinetii, Delise; Monogr. Stict. p. 124. t. 14. f. 51 (non Flor. Antarct. Pt. 1. p. 196). S. fulvo-cinerea, Mont. in Voy. au Pole Sud, Bot. Crypt. p. 184? S. glabra, nobis in Lond. Journ. Bot. vol. iii. p. 647 (in part). Parmelia lactucæfolia, Pers. in Freyc. Voy. Bot. p. 200. (TAB. CXCVI.)

HAB. Hermite Island, Cape Horn; trunks of trees and rocks, from the sea to the mountain tops. Falkland Islands; very abundant on maritime rocks, &c. Strait of Magalhaens, Port Famine; *Capt. King.* Staten Land; *Menzies.*

We have added a figure of this much-disputed species, concerning which we have fallen into an error in the previous part of this work, having regarded it as synonymous with the *S. glabra* of Lord Auckland's group and Tasmania (probably the *S. Delisea* Fée,), and which differs from the *S. Freycinetii* principally in the very shallow, not deeply cupped apothecia.

Fuegia and the Island of Juan Fernandez are the only localities in which we know this species to occur.

PLATE CXCVI. Fig. 1 and 2, portions of thallus of the natural size; 3, apex of ditto, with undeveloped and mature apothecia; 4, ditto with abortive (?) ditto; 5, slice of lamina proligera; 6, ascus; 7, spores :--very highly magnified.

10. STICTA filicina, Ach.; Lich. Univ. p. 145. Platisma Filix, Hoffm. Plant. Lich. t. 55.

HAB. Hermite Island, Cape Horn; on dead wood.

Our specimens, which are small and barren, differ in colour and in the less decidedly marked costæ, from those of New Zealand; the lobes also are occasionally furnished with an isidiophorous border. The thallus is about an inch and a half high, the upper surface of a dirty greenish-brown, the under pale yellow-brown and uniformly covered with a short tomentum, into which the concolorous and rather large cyphellæ are sunk. They may, indeed, belong to a state of *S. obvoluta*, Ach., with the upper surface glabrous; but hardly to any of the other species enumerated here.

S. STEREOCAULON, Ach.

1. STEREOCAULON corallinum, Fries; Lich. Europ. p. 201. Moug. et Nestl. n. 73. S. paschale, nobis in Lond. Journ. Bot. vol. iii. p. 653 (non Ach.).

HAB. Hermite Island, Cape Horn; on rocks near the sea. Kerguelen's Land; on alpine rocks, 600-1200 feet.

We have before pointed out the singular scarcity, in the Southern Hemisphere, of some of those Lichens which are most abundant in all latitudes of the North Temperate and Arctic Zones. Stereocaulon corallinum affords another remarkable instance of this anomalous distribution. Except, perhaps, the Cenomyce rangiferina, it is the very commonest of all Lichens in the subalpine districts of Britain and Central Europe, in the Alpine

regions of Southern Europe ascending to the summit of the Pyrenees, and to the level of perpetual snow on the Alps. Again, in the Arctic zone it is found carpeting the otherwise naked steppes of Asia and the barren lands of America, thence reaching the ultima thule of vegetable life in Melville Island and Ross Islet. To the south of its principal parallel it inhabits the Canary Islands, and a variety is seen on the Andes of Mexico and Colombia. Still further south it is replaced in all longitudes by the following species, being itself unknown in the Southern Hemisphere except at Cape Horn and Kerguelen's Land, where it re-appears in abundance. To reconcile this singular fact with the views of those who suppose it to have migrated into Kerguelen's Land, it is almost necessary to consider the *S. ramulosum*, which inhabits Lord Auckland's group, Campbell's Island, Tasmania, and the northern parts of Fucgia, as a southern variety of *S. corallinum*, which has, in Kerguelen's Land and Cape Horn, reverted to the northern form.

2. STEREOCAULON ramulosum, Ach. Fl. Antarct. Pt. 1. p. 195. t. lxxx. f. 1.

HAB. Strait of Magalhaens, Capt. King. Chonos Archipelago, C. Darwin, Esq.

This widely distributed species replaces in the Southern Hemisphere, to a considerable degree, the *S. paschale* and *corallinum* of the Northern, hut not fully; for it only enters what we have elsewhere defined to be the Antarctic zone of vegetation, not reaching the Falkland Islands, the southern parts of the Fuegian Islands or Kerguelen's Land. In the Old World it first appears in Bourbon, thence ranging from the Philippines, through Java, Australia, the South Sea Islands, Tasmania and New Zealand, to Lord Auckland's group and Campbell's Island, abounding in rocky and damp places, also on the trunks of large trees. In the New World it commences in the West Indian Islands, whence Swartz originally procured it, and runs through every parallel of latitude to the Strait of Magalhaens.

As a species S. ramulosum appears, at first sight, abundantly distinct from S. corallinum, nor does it display a tendency to assume any northern form of the genus in the Strait of Magalhaens. In Tasmania, again, where it ascends the mountains and becomes dwarfish, its lateral ramuli are still slender and fibrous, typical of the species. On the other hand, some of the tropical specimens, especially those from the Equatorial Andes (where both species occur), appear intermediate between S. ramulosum and corallinum; insomuch that it becomes a matter of opinion alone, whether the S. ramulosum should be considered a southern state of S. corallinum, owing its greater development to the more uniform temperature and humidity of the localities it affects in the Southern Hemisphere; or whether these are two species, one originating in the Southern Hemisphere, and one in the Northern, meeting under the Line, and there varying into the similitude of one another.

3. STEREOCAULON alpinum, Fries; Lich. Europ. p. 204.

HAB. Hermite Island, Cape Horn; on the summits of the hills.

A native of all the European Alps, also of the Andes of Peru.

4. STEREOCAULON denudatum, Somm.; Lapp. p.126. Fries, Lich. Europ. p.204. Moug. et Nestl. n. 466.

HAB. Hermite Island, Cape Horn; on rocks near the sea.

Also an inhabitant of the Alps of Europe and the Mexican Andes.

We are indebted to the Rev. Churchill Babington for the identification of the species of this difficult genus.

9. SPHÆROPHORON, Ach.

1. SPHEROPHORON coralloides, Ach.; Lich. Univ. p. 585. Engl. Bot. t. 115. Moug. et Nestl. n. 262.

HAB. Strait of Magalhaens, throughout Fuegia and the Falkland Islands; on the ground and on trunks of trees, most abundant, ascending to the tops of the mountains.

6 н

529

A common Antarctic American plant, rarer in Tasmania, and replaced in Lord Auckland's group by *S. tenerum*. Its range is very wide in both hemispheres, from within the Arctic circle of the New and Old Worlds, attaining Walden Island north of Spitzbergen, within 9° of the North Pole, stretching south, throughout Europe, to the Asturias, Switzerland, and Madeira, and in America to Newfoundland.

2. SPHEROPHORON tenerum, Laurer. Fl. Autarct. Pt. 1. p. 195. Mont. in Voy. au Pole Sud, Bot. Crypt. p. 172. (TAB. CXCVII. Fig. I.)

HAB. Hermite Island, Cape Horn; most abundant on the hills. Chonos Archipelago, C. Darwin, Esq.

In the former part of this work we have pointed out the characters which distinguish this species from the *S. coralloides*. It is much more frequent in Tasmania and New Zealand than in South America, in the latter country having been only found at Cape Horn, Fuegia, Chiloe, and the Chonos Archipelago.

I know of no Lichen which exhibits so well the successive development of "laminæ proligeræ" in the same apothecium. A vertical section of the youngest fruit shows two strata, parallel to, or rather concentric with, one another. Of these, the upper is fully ripe long before the bursting of the apothecium. It consists of immunerable filiform asci, containing from eight to thirty and more sporules. The sporules are vertically arranged and so densely packed that each ascus resembles a moniliform filament: the lower are smaller, the upper gradually larger; none however, attain their full size till after the absorption or disappearance of the walls of the ascus; when they escape as spherical bodies, surrounded by a narrow transparent margin.

The thallus of this genus consists of a firm crustaceous transparent cortex, whose inner edge is sharply defined, enclosing a mass of longitudinally arranged, matted, curved, dry filaments. These filaments are cylindrical, terete, sparingly supplied with very short ramuli, and truncate or obtuse at either extremity: they entirely surround the nucleus of the very immature apothecium.

PLATE CXCVII. Fig. I.—1, fertile, and 2, barren specimens, of the natural size; 3, young, 4, mature, and 5, aged apothecia; 6, 7, and 8, vertical sections of 3, 4, and 5, respectively, showing the formation of successive laminae proligera; 9, asci and spores; 10, young (or possibly abortive) asci; 11, mature ascus; 12, spores; 13, cortical and filamentous substance of thallus; 14, filaments from the latter :—all highly magnified.

3. SPHÆROPHORON compressum, Ach. Fl. Antaret. Pt. 1. p. 196.

HAB. Hermite Island, Cape Horn, and Falkland Islands; on turfy ground, abundant.

These specimens are identical with the English plant so called. It is also an Auckland Island species, and is found in various countries, both within and without the tropics, as far north as the barren lands bordering the Polar Sea in Arctic America. In Europe, Wahlenberg remarks, that it does not occur in any part of Scandinavia. In the Southern Hemisphere it grows on the South American Andes and in Van Diemen's Land.

4. SPHEROPHORON australe, Laurer. Fl. Antaret. Pt. 1. p. 195.

HAB. Strait of Magalhaens; Port Famine; Capt. King.

Manifestly identical with the Tasmanian, New Zealand, and Lord Auckland's group species of this name, but not hitherto found elsewhere in the New World.

5. SPHÆROPHORON fragile, Ach.; Lich. Univ. p. 585. Engl. Bot. t. 2474. Mont. in Voy. au Pole Sud, p. 172.

HAB. Strait of Magalhaens; D'Urville.

A frequent Arctic and North Temperate zone plant, reaching the latitude of Igloolik in the American Polar Sea, and, in Europe, Lapland, Spitzbergen and even Ross Islet, the most northern known land in the world.

10. CENOMYCE, Ach.

1. CENOMYCE pyridata, Ach.; Lich. Univ. p. 534. Engl. Bot. t. 1393. Schar. Lich. Helvet. n. 53-55.

HAB. Strait of Magalhaens; Port Famine, *Capt. King.* Port Gallant; *MM. Hombron* and *Jacquinot*. Abundant throughout the Tropics, but not observed south of the Strait of Magalhaens in extra-tropical South

America. In the Arctic regions it reaches to the very termination of vegetable life at Ross Islet, in 82°N.

2. CENOMYCE gracilis, Ach.; Lich. Univ. p. 550. Engl. Bot. t. 1824.

HAB. Falkland Islands; on the ground, not uncommon.

Possibly a state of *C. sparassa*, there being a decided tendency in the podetia to become squamulose. It is also a native of the extreme north, Spitzbergen, Walden, and Ross Islets.

3. CENOMYCE fimbriata, Ach.; Lich. Univ. p. 535. Engl. Bot. t. 2438.

Var. ustulata; podetiis brevibus lanceolatis fistulosis curvato-decumbentibus basi concoloribus apice nigrescentibus, gemmis pulverulento-granulosis. C. ustulata, nobis in Lond. Journ. Bot. vol. iii. p. 652.

HAB. Falkland Islands; abundant on dry heathy soils : Var. ustulata, on sand-hills, near Uranie Bav.

The apothecia in these specimens copiously fringe the margins of the cups, and becoming coalescent form a broad lobed mass.

4. CENOMYCE verticillata, Ach.; Lich. Univ. p. 555. Dill. Hist. Musc. t. 14. f. 6 G.

HAB. Falkland Islands; in heathy places, abundant.

Our specimens entirely agree with others of British growth and with the figure of Dillenius.

5. CENOMYCE cornuta, Ach.; Lich. Univ. p. 545. Fries, Lich. Europ. p. 225.
Var. γ. ramosa, Delise; Mont. in Foy. an Pole Sud, Bot. Crypt. p. 174.
HAB. Strait of Magalhaeus; on rocks and trunks of trees, M. Jacquinot.

6. CENOMYCE furcata, Ach.; Lich. Univ. p. 560.

Var. squamulosa, Delise; Mont. in Voy. au Pole Sud, Bot. Crypt. p. 175.HAB. Strait of Magalhaens, Port Famine; on dead trunks of trees, M. Jacquinot.

7. CENOMYCE coccifera, auct.; Engl. Bot. t. 2051. Cladonia cornucopioides, Fries, Lich. Europ. p. 236. HAB. Hermite Island, Cape Horn, and the Falkland Islands; abundant on the hills.

8. CENOMYCE deformis, Ach.; Lich. Univ. p. 538. Engl. Bot. t. 1349. Schærer, Lich. Helvet. n. 47-49. HAB. Hermite Island, Cape Horn, and the Falkland Islands; ascending to the tops of the mountains.

9. CENOMYCE rangiferina, Ach.; Lick. Univ. p. 564. Engl. Bot. t.173. Scharer, Lich. Helvet. n.76,77.
Var. alpestris, Eschw.; Dill. Hist. Musc. t. 16. f. 30 A. B. Fries, Lich. Europ. p. 243.
Var. sylvatica, Hoff.; Dill. l. c. f. 29 E. F. Fries, Lich. Europ. p. 243. Scharer, Lich. Helvet. n. 78.
HAB. Throughout South Chili, Fuegia, and the Falkland Islands; most abundant.

Though so widely distributed a Lichen, and, as Fries remarks, "omnium Lichenum copiosissima", this species has its limits within the parallels attained by its congeners and other plants. In the south it is stunted at Cape Horn, and neither inhabits Kerguelen's Land or the South Shetlands; whilst, towards the Northern Pole, it was not detected in Melville Island, though attaining a much higher latitude in Spitzbergen.

10. CENOMYCE uncialis, Ach.; Lich. Univ. p. 559. Engl. Bot. t. 174. Scharer, Lich. Helvet. n. 84.

HAB. Strait of Magalhaens; Port Famine, Capt. King.

Not observed in Hermite Island or the Falklands; nor does it extend in the Arctic regions beyond the continents of Europe and America.

11. CENOMYCE? vermicularis, Ach.; Lich. Univ. p. 566. Engl. Bot. t. 2029.

HAB. Hermite Island, Cape Horn, and the Falkland Islands; local, but very abundant where it occurs.

We cannot but regard this as the abnormal state of some *Cenomyce* (possibly of *alcicornis* or *endiviæfolia*?); though we have never succeeded in identifying the species. It is a highly Arctic and Antarctic plant, in the northern regions advancing to the extreme limits of vegetation, in islands beyond Spitzbergen. It also has been collected on the Andes of Peru and of Colombia.

12. CENOMYCE aggregata, Ach.; Lich. Univ. p. 563. Fl. Antarct. Pt. 1. p. 197. t. lxxx. f. 2.

HAB. South Chili, throughout Fuegia and the Falkland Islands; from the sea to the hill-tops, very abundant.

A very abundant plant in the higher latitudes of the Southern Hemisphere, from Monte Video on the east, Mendoza in Central Chili, Colombia on the Andes, and Juan Fernandez on the west coast of South America to Cape Horn. Its various northern limits in the Old World are the Cape colony in South Africa, Nepaul in Asia, Swan River in Australia, and Norfolk Island in the Pacific. In Tasmania and New Zealand it abounds, reaching 52°S. in Campbell's Island.

13. CENOMYCE bacillaris, Ach.; Synops. p. 266. Cladonia macilenta, Fries, Lich. Europ. p. 241.

HAB. Falkland Islands; dry places on the hills.

Probably a state of *C. coccifera*, and the original *C. coccifera*, *a.*, Linn. It is a native both of the Tropics, and north Temperate zones.

14. CENOMYCE sparassa, Ach.; Synops. p. 273. Engl. Bot. t. 2362. Clad. squamosa and ventricosa Fries, Lich. Europ. p. 231.

HAB. Hermite Island, Cape Horn; on the mountain tops.

The C. ecmocina, var. gracilis, of Lord Auckland's group, should be referred here; its podetia being squamulose, though but slightly so.

11. PARMELIA, Ach.

1. PARMELLA enteromorpha, Ach.; Synops. p. 219. P. physodes, β. vittata, Mont. in Voy. au Pole Sud, Bot. Crypt. p. 182. P. lugubris, Pers. in Freyc. Voy. Bot. p. 196.

Var. β . deusta; parvula, rigida, thallo subcrecto brevi subflabellatim diviso, lobis atris patulis angustis canaliculatis utrinque concoloribus.

HAB. Hermite Island, Cape Horn; from the sea to the mountain tops. Falkland Islands; very abundant. Var. β . barren rocks near the top of Kater's peak.

This species was also found abundantly in Lord Auckland's group and Campbell's Island, though omitted in the first part of the Flora Antarctica. It is further a native of New Zealand and Tasmania; of North-west America, from California to Sitka, and we possess a specimen labelled as from the Mauritius.

Specifically, this differs from P. physodes only in the length of the lobes of the thallus, and these are so variable as to lead to some doubts of the validity of the species. In Tasmanian specimens the lobes are often much dilated and plane, the membranes of which it is composed, and which are normally inflated, being here, not only in contact, but firmly united together; thus effecting a passage between this species and the forms to which P. perlata, &c. belong.

We have authentically named specimens of the North-west American *P. physodes*, β . *vittata*, which is only a narrower state of *P. enteromorpha*. Norwegian specimens also of the latter plant appear to be clearly referable to this.

2. PARMELIA diatrypa, Ach.; Syn. Lich. p. 219. Engl. Bot. t. 1248. Moug. et Nestl. n. 65.

HAB. Hermite Island, Cape Horn; on stems of bushes and on branches of trees, on the mountains. Chonos Archipelago, C. Darwin, Esq.

Probably only a small, or alpine, form of *P. enteromorpha*; it was found in similar situations in Lord Auckland's group, and on the top of Mount Wellington in Tasmania. Besides being a native of Great Britain and alpine situations in northern and midland Europe and of the Sandwich Islands.

3. PARMELIA cincinnata, Ach.; Lich. Univ. p. 495. Syn. Lich. p. 219. (TAB. CXCVII. Fig. II.)

HAB. Staten Land, *Menzies*. Hermite Island, Cape Horn; on rocks and trunks of trees above the limit of the evergreen Beech.

By the apothecia this beautiful species may be distinguished, both from *P. diatrypa* and *P. enteromorpha*, some of our specimens, indeed, are on the same piece of wood with *P. diatrypa*, both retaining their characters. They entirely agree in every other respect with one collected by Menzies, except in being of a pale lemon colour.

PLATE CXCVII. Fig. II.—1, specimen of the natural size; 2, portion of ditto; 3, vertical section of apothecium; 4, slice of lamina proligera; 5, ascus; 6 and 7, spores :--highly magnified.

4. PARMELIA saxatilis, Ach.; Synops. p. 203. Engl. Bot. t. 603. Mougeot ct Nestler, n. 347 and 738.

HAB. Hermite Island, Cape Horn, and the Falkland Islands; abundant on alpine rocks. Cockburn Island, Graham's Land; very scarce.

None of these specimens are in fruit, but they accord perfectly with Scottish and other European examples. The lobes of the thallus vary a good deal in size and colour, according to exposure. What is believed to be this plant was seen at Cockburn Island, on the verge of Antarctic vegetation, but, as the specimens were lost previous to comparison, some doubt may be entertained of the correctness of this habitat. Besides being abundant throughout Europe, advancing as far north in Spitzbergen as vegetation extends, and in Temperate and North America, this species has been found on the Mexican Andes, on the barren grounds bordering the Polar Sea, and also in the Arctic Islands.

5. PARMELIA rubiginosa, Ach.; Lich. Univ. p. 467. Engl. Bot. t. 983.
Var. β. sphinetrina. P. sphinetrina, Mont. in Voy. au Pole Sud, Bot. Crypt. p. 180. t. 45. f. 3.
HAB. Var. β. Hermite Island, Cape Horn; on trunks of trees.

6 I

Chiefly characterized by its much more continuous and leafy thallus. The *P. rubiginosa* is a sub-Arctic species, extending as far north as the region of Willows and Birch in Norway.

6. PARMELIA stellaris, Ach.; Lich. Univ. p. 476. Engl. Bot. t. 1351.

HAB. Cockburn Island, Graham's Land; on rocks.

A specimen apparently of this species was found; but in a very insufficient state for determination.

12. LECANORA, Ach.

§ I. Psoroma, Fries.

1. LECANORA microphylla, Ach.; Lich. Univ. p. 420. Engl. Bot. t. 1247. Scharer, Lich. Helvet. n. 160.

HAB. Staten Land; on dead wood, A. Menzies, Esq.

Possibly the *L. triptophylla*, Fries, but the specimens are not very satisfactory; they agree tolerably with the plate and specimens quoted. *C.Babington*.

2. LECANORA paleacea; (Parmelia), Frics, Lich. Europ. p. 97. (TAB. CXCVII. Fig. III.)

HAB. Falkland Islands; on the ground and on Tussock mounds, rare.

 Λ very rare and curious species, hitherto known only as a native of Denmark. The paleaceous apothecia resemble a *Peziza*. We have seen no authentic specimens, and add a figure of the Falkland Island plant.

PLATE CXCVII. Fig. III.—1, plant of the natural size; 2 and 3, young and mature apothecia; 4, vertical section of portion of apothecium; 5, asci; 6, spores :--highly magnified.

3. LECANORA muscorum, Ach.; Syn. Mcth. Lich. p. 193. Lich. carnosus, Engl. Bot. t. 1684. HAB. Falkland Islands; on the ground and on decaying roots of Ferns.

4. LECANORA Hypnorum, Ach.; Syn. Meth. Lich. p. 193. Engl. Bot. p. 740. Fl. Antarct. Pt. 1. p. 199. HAB. Hermite Island, Cape Horn; on mossy trunks of trees. Falkland Islands; on the ground, &c.

§ II. Placodium, Fries.

5. LECANORA chrysoleuca, Ach.; Lich. Univ. p. 411.

Var. β . Dultoni; thallo centro affixo, lobis radiantibus cuneatis, gemmis marginalibus granuliferis. Lecanora Daltoniana, nobis in Lond. Journ. Bot. vol. iii. p. 641. (TAB. CXCVIII. Fig. I.)

Var. y. lignicola; thallo adnato, lobis cortice appressis.

HAB. Var. β . Cockburn Island, Graham's Land. Var. γ . Hermite Island, Cape Horn; on trunks of trees.

A very rare inhabitant of the Southern Hemisphere, and there confined to the Antarctic regions. The two varieties are certainly not distinct from the European *L. chrysoleuca*, which inhabits mountainous regions from Norway to the Alps and Pyrenees.

PLATE CXCVIII. Fig. I.—1, mature, and 2, immature specimens of var. β . of the natural size; 3, upper, and 4, lower view of thallus; 5, central portion of ditto, with young apothecia; 6, mature apothecium; 7. vertical section of ditto; 8 and 9, portions of ditto showing the lamina proligera; 10, asci; 11, spores :—all magnified.
6. LECANORA *Babingtoni*, Hook. fil. et Tayl.; thallo crustaceo adnato orbiculari subsquamuloso areolato areolis radiantibus albido-glaucescente demum virescente, squamulis diffractis ambitu sub-continuis crenulatis, apotheciis adnatis margine thallode tenuissimo evanescente, disco atro primitus tumido margine subelevato demum planiore immarginato. Lecidea atro-alba, *nobis in Lond. Journ. Bot.* vol. iii. p. 636 (quoad exempl. Ins. Cockburn). (TAB. CXCVIII. Fig. II.)

HAB. Cockburn Island, Graham's Land; on volcanic rocks.

Thallus inconspicuus $\frac{1}{2}$ -1 unc. diametro, margine definito crenulato, totus in areolas minimas (non nisi ope lentis conspicuas) diffractus ; areolis angulatis, albidis, saxæ adnatis, centralibus fertilibus, reliquis radiantibus, extimis subfoliaceis lobatis crenulatisve. Apothecia punctiformia, interiora majora subconfluentia.

Although the specimens of this plant brought from Cockburn Island are very perfect and well developed, they belong to so difficult a group of Lichens as to have baffled the Rev. Mess. Babington, Berkeley, and ourselves, in our attempts to reduce it to any known species. Though closely resembling a *Lecidea* in habit, and, indeed, in characters too, it is certainly not of that genus, for though, as Mr. Berkeley remarks, the apothecia of *Lecid. rivulosa* and *confluens* are sometimes obscurely margined (as in this species), yet, Mr. Babington observes, that the thallus here is radiating, which is not the case with the areolate *Lecidea*, nor has it the carbonaceous margin to the apothecium and substratum of that genus.

Of the tribe in which it should be placed (as a *Lecanora*) there is some doubt: Mr. Berkeley regards it as belonging to the section "glaucescentes" of *Placodium*, Fries, and allied to *L. coarctata*, in which the thallodal border of the apothecium is evanescent. Mr. Babington, on the other hand, remarks, that the thallodal border and that of the disc itself place it in *Psora*, Fries, and that it will rank amongst the section "glaucescentes" near *L. melanaspis*, of which it may possibly be a variety, or a depauperated and crustaceous form. The figure represents the plant as freshly gathered, before drying; it has since assumed a more obscure, somewhat leaden colour, and the oldest portions of the thallus break up into a greenish mass, which is not given in our plate.

PLATE CXCVIII. Fig. II.—1, plant of the natural size; 2, portion of ditto; 3, central part of thallns and apothecia; 4 and 5, lateral views of areola and apothecia; 6, vertical slice of two apothecia; 7, portion of lamina proligera:—highly magnified.

7. LECANORA gelida, Ach.; Lich. Univ. p. 428. Engl. Bot. t. 699. Urccolaria macropthalma, nobis in Lond. Journ. Bot. vol. iii. p. 640.

Var. β . vitellina, thallo vitellino.

HAB. Kerguelen's Land; both varieties very common.

The variety β is probably dependent on the thallus having changed colour. What was described as Urceolaria macrophalma is a state noticed by Fries (Lich. Europ. p. 104). C.Babington.

8. LECANORA murorum, Ach.; Lich. Univ. p. 433. Engl. Bot. t. 2157.

Var. farcta; apotheciis substantia granulata fere clausis.

HAB. Hermite Island, Cape Horn, the Falkland Islands, and Cockburn Island, Graham's Land; on maritime rocks. Var. β . Kerguelen's Land; on rocks near the sea.

9. LECANORA miniata, Ach.; Lich. Univ. p. 434. Hoffm. Plant. Lich. t. 60. f. 1.

HAB. Falkland Islands, and Cockburn Island, Graham's Land; very abundantly on rocks near the sea.

This plant forms the most curious feature in the botany of Cockburn Island, a desolate spot of land on the extreme limit of southern vegetation; for there it abounds so as to stain the rocks, and render the colour thus

[Fuegia, the

produced visible for many miles. It is partial to the efflurium from decaying animal matter, as is the case with other *Parmelia* belonging to the citrinous series.

10. LECANORA citrina, Ach.; Lich. Univ. p. 402. Engl. Bot. t. 857. Moug. et Nestl. n. 742.

HAB. Kerguelen's Land; on rocks near the sea.

Specimens very imperfect, but, we think, referable to this species.

11. LECANORA erythrocarpia, Fries, Lich. Europ. p. 119. L. theioclyta, Ach. Lich. Univ. p. 425. HAB. Kerguelen's Land; in a cave near the sea. Rather a doubtful determination. C.Babington.

§ III. Psora, Fries.

12. LECANORA melanaspis, Ach.; Lich. Univ. p. 427. Fries, Lich. Europ. p. 122. L. dichroa, nobis in Lond. Journ. Bot. vol. iii. p. 643.

HAB. Kerguelen's Land; on hard earth and stones, rare.

13. LECANORA molybdina, Ach.; Lich. Univ. p. 430. Fries, Lich. Europ. p. 126.

HAB. Kerguelen's Land; on bare and hardened earth.

§ IV. Patellaria, Fries.

14. LECANORA tartarea, Ach.; Lich. Univ. p. 172. Engl. Bot. t. 156.

HAB. Hermite Island, Cape Horn, and the Falkland Islands; abundant. Kerguelen's Land; rare.

15. LECANORA parella, Ach.; Lich. Univ. p. 370. Engl. Bot. t. 727.

Var. y. Upsaliensis, Ach.; Lich. Univ. p. 371. Engl. Bot. t. 1634.

HAB. Falkland Islands; on quartz rocks. Var. γ . Hermite Island, Cape Horn, and the Falkland Islands; on the ground.

16. LECANORA subfusca, Ach.; Lich. Univ. p. 393. Engl. Bot. t. 2109.

Var. epibryon; Lecanora epibryon, Ach. l. c. Moug. et Nestl. n. 120.

Var. albella, Fries; Lecanora albella, Ach. l. c. Engl. Bot. t. 2157.

HAB. Strait of Magalhaens; on wood, *Capt. King.* Falkland Islands; on rocks, and on dead twigs of *Acana.* Var. *epibryon*, Kerguelen's Land; on decayed *Azorella.* Var. *albella*, Hermite Island, Cape Horn; on Winter's bark.

17. LECANORA atra, Ach.; Lich. Univ. p. 344. Engl. Bot. t. 949. Moug. et Nestl. n. 458.

Var. B. confragosa, Ach.; l. c. p. 345.

HAB. Hermite Island, Cape Horn, and the Falkland Islands; on rocks. Var. β . Hermite Island, Cape Horn; on rocks.

18. LECANORA sophodes, var. c. exigua, Fries, Lich. Europ. p. 149. Engl. Bot. t. 1849. HAB. Falkland Islands; on rocks.

19. LECANORA ventosa, Ach.; Lich. Univ. p. 399. Engl. Bot. t. 906. Moug. et Nestl. n. 256. HAB. Falkland Islands; on alpine quartz rocks.

20. LECANORA hæmatomma, Ach.; Lich. Univ. p. 388. Engl. Bot. t. 486.

HAB. Falkland Islands; on clay-slate near the sea and on quartz rocks on the hills.

21. LECANORA candelaria, Ach.; Lich. Univ. p. 416. Engl. Bot. t. 1794.

HAB. Falkland Islands; on twigs of Acana. Kerguelen's Land; on maritime rocks.

22. LECANORA erythrella, Ach.; Lich. Univ. p. 401. Engl. Bot. t. 1993.

HAB. Falkland Islands; on very arid quartz rocks at Port William.

13. URCEOLARIA, Ach.

1. URCEOLARIA sordida, Fries; Lich. Europ. p. 178. Lecanora glaucoma, Engl. Bot. t. 2156. HAB. Falkland Islands; on rocks.

2. URCEOLARIA endochlora, Hook. fil. et Tayl.; crusta cinereo-albida tenui-rimosa inæquabili nigrolimitata et punctata, apotheciis immersis planis atris retundato-difformibus, margine thallode tenui madore obsoleto, lamina tenuissima virescenti. Nobis in Lond. Journ. Bot. vol. iii. p. 640.

HAB. Kerguelen's Land; on rocks.

Crusta 2-unc. lata, eburnea, demum virescens. Apothecia obconica, disco atro-pruinoso.

The lamina proligera rests upon a very thin layer of green matter, which, again, is placed on a black hypothallus. The plant approaches the *Bæomyces anomalus*, Tayl. (in Flor. Hib.)

3. URCEOLARIA scruposa, var. β. bryophila, Fries; Lich. Europ. p. 101.

HAB. Hermite Island, Cape Horn; rocks on the top of Kater's peak.

4. URCEOLARIA erubescens, Hook. fil. et Tayl.; thallo crustaceo rimoso arcolato inæquabili ruguloso albido plerumque rufescente nigro-limitato, apotheciis confertis immersis concaviusculis difformibus subpellucidis olivaceis, disco scabrido nigro-punctato, margine thallode lacero-crenulato. *Nobis in Lond. Journ. Bot.* vol. iii. p. 640.

HAB. Falkland Islands; on rocks.

Thallus albidus, superficie plerumque colorato. Lamina proligera pellucida, strato albido insidens.

5. URCEOLARIA calcarea, Ach.; Lich. Univ. p. 340. Lichen cinereus, Engl. Bot. t. 820. Porina fallax (in part), nobis in Lond. Journ. Bot. vol. iii. p. 639.

HAB. Hermite Island, Cape Horn; on rocks and stones.

14. BIATORA, Fries.

1. BIATORA corallina. Lecidea corallina, Eschweiler in Mart. Flor. Bras. p. 256. L. mamillata, nobis, et L. geomæa, Tayl.; nobis in Lond. Journ. Bot. vol. iii. p. 636 and 637.

HAB. Hermite Island, Cape Horn; on the ground. Falkland Islands; on tufts of Bolax.

6 к

A true *Biatora*, having the margins of the young apothecia both coloured and soft. It is not a European species, though belonging to Fries' section "fuscescentes" and allied to *B. uliginosa*. I have little doubt of its being the plant of Eschweiler. *C. Babington*.

2. BIATORA pulverca, (Lecidea) Borr.; in Engl. Bot. Suppl. t. 2726.

HAB. Falkland Islands; on rocks.

15. LECIDEA, Ach.

1. LECIDEA mamillaris, Fries? Lich. Europ. p. 285. Lichen tumidulus, Smith in Linn. Soc. Trans. vol. i. p. 82. t. 4. f. 3.

HAB. Hermite Island, Cape Horn; on the top of Kater's peak.

It is possible that these imperfect specimens may be referable to some paradoxical form of L. vesicularis. C.B.

2. LECIDEA candida, Ach.? Lich. Univ. p. 212. Engl. Bot. t. 1138. Schærer, Lich. Helvet. n. 167. HAB. Hermite Island, Cape Horn; on hard soil.

I am not satisfied with the identification of this with the British L. candida: it may be merely a form of L. vesicularis. The hypothallus is black. C. Babington.

3. LECIDEA vesicularis, Ach.; Lich. Univ. p. 212. Engl. Bot. t. 1139. Scharer, Spicel. p. 120. Lich. Helvet. n. 168.

HAB. Hermite Island, Cape Horn, and the Falkland Islands; on the ground, abundant.

These specimens resemble *Isidium oculatum* when in a young state. I am in doubt whether some may not be referable to *L. epigæa*, which, as well as *L. vesicularis*, is a very polymorphous plant. *C. Babingtoa*.

4. LECIDEA aromatica, Ach.; Lich. Univ. p. 168. Engl. Bot. t. 1777.

HAB. Kerguelen's Land; on moist rocks.

Specimens greener and darker than in Schærer's, but still I think referable to that species. C.B.

5. LECIDEA albo-cærulescens, Ach.? Lich. Univ. p. 188. Lichen pruinosus, Engl. Bot. t. 2244. L. confluens, nob. (in part), Lond. Journ. Bot. vol. iii. p. 636.

HAB. Kerguelen's Land; on rocks near the sea and on the hills.

Specimens very fine, altogether resembling what I have gathered on the Austrian Alps. C. Babington.

6. LECIDEA spilota, Fries; Lich. Europ. p. 297. L. rivulosa, nobis, in Lond. Journ. Bot. vol. iii. p. 636.

HAB. Kerguelen's Land; on moist rocks.

The specimens, which are rather young, certainly do not belong to *L. rivulosa*, and are, I think, referable to *L. spilota*, with which species, however, I am not sufficiently acquainted. *C. Babington*.

7. LECIDEA contigua, Fries; Lich. Europ. p. 298. L. speirca, var. hydrophila, Fries, l. c. Lec. confluens, (in part), Engl. Bot. t. 1864.

HAB. Hermite Island, Cape Horn; on rocks. Var. hydrophila, Kerguelen's Land; also on rocks.

The disc does not appear to be pruinose when young, in which respect only this differs from Fries' plant. It is allied to L. *nitidula*, differing from it only in the apothecia springing from the crust; which is the case also in Schwere's specimens of that plant. *C. Babington*.

8. LECIDEA atro-alba, Ach.; Lich. Univ. p. 162. Lichen Ederi, Engl. Bot. t. 1117. Schærer, Lich. Helvet, n. 178.

HAB. Hermite Island, Cape Horn; on rocks.

The Cockburn Island plant, referred (in Lond. Journ. Bot.) to this, we have elsewhere described as *Lecanora* Babingtoni.

9. LECIDEA lugubris, Sommf.; Lapp. p. 143. Fries, Lich. Europ. p. 314.

HAB. Hermite Island, Cape Horn; on rocks.

I have little hesitation in considering this plant to be that described by Fries; though I know the latter from description alone. It approaches Schærer's L. atro-alba (n. 178); but the apothecia do not appear to rise from the thallus. C. Babington.

10. LECIDEA fusco-atra, Ach.; Lich. Europ. p. 359. Lichen athrocarpus, Engl. Bot. t. 1929. HAB. Kerguelen's Land; on rocks.

11. LECIDEA stellulata, Tayl. in Flora Hib. p. 118.

HAB. Kerguelen's Land; on rocks.

12. LECIDEA glacialis, Scherer; Spieil. p. 147. Fries, Lich. Europ. p. 323. L. sulphurea, Ach., nobis in Lond. Journ. Bot. vol. iv. p. 636.

HAB. Falkland Islands, on alpine quartz rocks.

At first sight this resembles *Lecanora subfusca*, (or rather the var. *orosthea*,) but the hypothallus is quite black. I feel no doubt of its being the *L. gtacialis*, which is compared with the above-mentioned species, both by Fries and Schærer. The only difference I can detect between the European and Antarctic specimens, lies in the apothecia of the former being dusky within, and not white.

13. LECIDEA geographica, Schærer; Spicil. p. 124. Engl. Bot. t. 245. Var. urceolata, Schærer, l. c.
HAB. Hermite Island, Cape Horn, the Falkland Islands, and Kerguelen's Land; very abundant on

14. LECIDEA parasema, Ach.; Lich. Univ. p. 175. Schærer, Lich. Helvet. n. 197-199.

rocks from the sea to the mountain tops. Var. urceolata, Kerguelen's Land; on maritime rocks.

HAB. Hermite Island, Cape Horn; on bark of trees.

15. LECIDEA sabuletorum, Ach.; Synops. Lich. p. 20. L. quadricolor, Borr., nobis, in Lond. Journ. Bot. vol. iii. p. 637.

Var. y. coniops, Fries, Lich. Europ. p. 340. L. scabra, Tayl. in Herb. Hib. p. 121.

HAB. Falkland Islands; on the ground. Var. coniops; Hermite Island, Cape Horn; on greenstone. Falkland Islands; on clay-slate rocks.

I am hardly satisfied with the determination of the Hermite Island specimens: they are certainly allied to the *L. sabuletorum* and also to *L. arctica*. They further resemble *Biatora vernalis*, var. sanguineo-atra, Fries; but this *L. sabuletorum* is scarcely a *Biatora*, and may be an undescribed species of *Lecidea*. *C. Babington*.

16. LECIDEA arctica, Sommerf., Lapp. p. 156. Fries, Lich. Europ. p. 342. HAB. Hermite Island, Cape Horn; on hard gravelly soil.

17. LECIDEA milliaria, var. c. ligniaria, Fries, Lich. Europ. p. 343. Lichen dubius, Engl. Bot. t. 2347. Lecidea elæochroma, Ach.; nobis, in Lond. Journ. Bot. vol. iii. p. 636.

HAB. Falkland Islands; on dead twigs of Acana.

A very puzzling species, differing from *L.elæochroma* in the pale hypothallus. I am doubtful if the English Botany *L. dubius* be the same plant, or *L. milliaria*, Fries. The only others to which the Falkland Island one ean be referred, are *L. dolosa*, Fries, and *L. sabuletorum*, Fr.: but after a careful examination of Schærer's specimen of the former, and Reichenbach's of the latter, I have concluded that this belongs to neither of them. *C.B.*

18. LECIDEA abietina, var. rubens, Eschweiler, in Mart. Fl. Bras. p. 251.

HAB. Hermite Island, Cape Horn; on bark.

I doubt not this being Eschweiler's plant, but am not convinced of that being the same with the European L. abietina. The hypophloidal erust appears to indicate its not being a true *Lecidea*, whence it may prove to be a Lecideal form of some *Pyrenotheca*; to which genus the *L. abietina* properly belongs.

16. GYROPHORA, Ach.

1. GYROPHORA ænea, var. a. Schærer, Lich. Helvet. n. 149.

HAB. Falkland Islands; on quartz rocks; very rare and barren.

It is remarkable that the Antarctic regions should present us with but a solitary species of this curious genus, which abounds so strikingly in the Arctic. In one respect they are replaced by Sticta, which are almost equally rare in the high northern latitudes. These latter affect an equable, as decidedly as the *Gyrophora* do an extreme elimate; and it is in the Falkland Islands, of all the Antarctic localities, that the Liehens are exposed to the greatest and most sudden viewsitudes. The *G. enea*, considered by Fries as a variety of *G. hyperborca*, is a Scotch and American plant.

17. OPEGRAPHA, Ach.

1. OPEGRAPHA atra, Pers.; Schærer, Lich. Helvet. n. 93.

HAB. Hermite Island, Cape Horn; on the bark of trees. Falkland Islands; on stems of Acana.

18. ARTHONIA, Ach.

1. ARTHONIA polymorpha, Ach.; Syn. Lich. p. 7. Eschweiler, in Mart. Fl. Bras. Crypt. p. 14. t. 9. f. 3. (tabula sub. nom. A. tremellosa.) Lecanora microphalma, nobis in Lond. Journ. Bot. vol. iii. p. 636.

HAB. Hermite Island, Cape Horn; on Winter's bark.

19. PERTUSARIA, DC.

1. PERTUSARIA communis, DC.; Engl. Bot. t. 677. Schærer, Lich. Helvet. n. 118.

HAB. Hermite Island, Cape Horn; encrusting the bark of trees, abundant. Kerguelen's Land; on rocks near the sea. Cockburn Island, Graham's Land; on rocks.

The Cockburn Island specimens are very imperfect, and may possibly helong to Umbilicaria sordida.

2. PERTUSARIA Wulfenii, DC.; Fries, Lich. Europ. p. 424. Porina fallax, Pers.; Ach. Synops. Lich. p. 110. Lichen hymenius, Engl. Bot. t. 1731.

HAB. Falkland Islands; on rocks.

20. THELOTREMA, Ach.

1. THELOTREMA lepadinum, Ach.; Lich. Univ. p. 312. t. 6. f. 1. Schærer, Lich. Helvet. u. 121. Fl. Antaret. Pt. 1. p. 200.

HAB. Hermite Island, Cape Horn; on Winter's-bark.

21. VERRUCARIA, Pers.

1. VERRUCARIA umbrina, Ach.; Lich. Univ. p. 291. Engl. Bot. t. 1499. V. gelida, nobis in Lond. Journ. Bot. vol. iii. p. 639. (TAB. CXCVIII. Fig. IV.)

HAB. Cockburn Island, Graham's Land; on rocks.

The difference between the apothecia of the Antaretic and European specimens is the same as exists between *V. maura* and *V. umbrina*, plants which I consider as specifically the same. *C. Babington*.

PLATE CXCVIII. Fig. IV.—1, plant of the natural size; 2, portion of crusti; 3, do with apothecia; 4 and 5, apothecia; 6 and 7, vertical slices of do; 8, portion of lamina proligera; 9 and 10, sporules:—all very highly magnified.

22. COLLEMA, Ach.

1. Collema erispum, Ach.; Synops. Lich. p. 311. Engl. Bot. t. 834. Parmelia pulposa, Schærer. HAB. Cockburn Island, Graham's Land; on wet earth.

Miserably depanperated specimens, referred both by the Rev. Mr. Berkeley and Babington to this plant.

2. COLLEMA tremelloides, Ach.; Lich. Univ. p. 455. Engl. Bot. t. 1981.

HAB. Hermite Island, Cape Horn; on wet banks.

Possibly the C. palmatum, Sm.; my only specimen of which plant (received from Mr. Borrer), may be a dwarf and brown one of C. tremelloides. C. Babington.

3. COLLEMA saturninum, Ach; Lich. Univ. p. 644. Engl. Bot. t. 1980. C. myochroma, Scheerer, Lich. Helvet.

Var. australe, thallo subferrugineo. Collema australis, nobis in Lond. Journ. Bot. vol. iii. p. 656.

ILAB. Hermite Island, Cape Horn; on wet banks in dense woods; abundant.

There is a redder hue about these specimens than I am accustomed to see in British ones of *C. saturninum*, but according to Schærer's description, this is evidently a very variable plant. The characters drawn from the powdery buds, are not available. *C. Babington*.

6 L

(Lichenes imperfecti.)

23. ISIDIUM, Ach.

1. ISIDIUM oculatum, Ach.; Lich. Univ. p. 570. Engl. Bot. t. 1833.

HAB. Hermite Island, Cape Horn, the Falkland Islands and Kerguelen's Land; on the earth.

Various *Parmeliæ* and *Lecanoræ* in a young state, are scarcely distinguishable from one another, and have been referred to *Isidium oculatum*.

2. ISIDIUM lutescens, Turn. and Borr. Lepraria lutescens, Engl. Bot. t. 1529.

HAB. Kerguelen's Land; on rocks near the sea.

Very probably a state of Lecanora murorum.

24. LEPRARIA, Ach.

1. LEPRARIA flava, Ach.; Lich. Univ. p. 663. Engl. Bot. t. 1350.

HAB. Hermite Island, Cape Horn, the Falkland Islands and Kerguelen's Land; abundant near the sea. Evidently the powdery state of some *Parmelia*, belonging to the citrinous series.

.

ADDENDA ET CORRIGENDA.

PART I.

- p. 8. DROSERA sp.—I have examined a specimen of this plant, collected by one of the officers of Admiral D'Urville's expedition, and by him given to my friend Mr. Gunn of Tasmania. It is certainly very nearly allied to the *D. uniflora* of Cape Horn, but differs from that and from all its congeners in the perigynous insertion of the stamina.
- p. 10. EPILOBIUM confertifolium.—Mr. Watson has given me cultivated specimens of *E. alpinum*, entirely according in habit and foliage with this plant.
- p. 10. ACENA adscendens.-The Kerguelen's Land species differs from this, see Pt. 2. p. 268. t. 96 B.
- p. 14. COLOBANTHUS subulatus.—For an explanation of the monstrous appearance of the flowers alluded to, see Pt. 2. p. 248.
- p. 19. ARALIA polaris .- For analysis of the flowers, &c., see Hook. Ic. Plant. t. 747.
- p. 22. COPROSMA repens.—I have, since the publication of this species, received from Mr. Gunn fruiting specimens of the Tasmanian plant included under this, and figured at Pt. I. t. 16 B. In them the berry has but two nucules; and I am obliged to regard it as a distinct species, to which the name of C. pumila has been given.
- p. 30. HELICHRYSUM prostratum :----This is the true *H. bellidioides* of Forster, though not of Banks and Solander (Hb. Banks): in its prostrate straggling habit it resembles a Cape species.
- p. 32. PLEUROPHYLLUM criniferum.—I have observed the same rigid setæ amongst the tomentum of the foliage in this plant as exist in the *P. speciosum*.
- p. 37. To notes upon CELMISIA vernicosa, add :--- The generic distinction between this genus and Chiliotrichum rests on the presence of scales on the receptacle of the latter.
- p. 37. Of the two Compositæ alluded to as "dubii generis" I have had the opportunity of examining specimens, they are

1. GNAPHALIUM luteo-album, Linn.

This abounds throughout New Zcaland, from the Bay of Islands to Stewart's Island in the extreme south: I have seen Auckland Island specimens collected by the French Antarctic Expedition, with which Dr. Lyall's barren ones entircly accord.

2. EURYBIA (Brachyglossa) Lyallii, Hook. fil.; foliis amplis alternis breviter petiolatis ellipticoovatis obovatisve acutis plerumque argute subduplicato-dentatis coriaceis super glaberrimis venosis subter tomento dense appresso lanatis, paniculis terminalibus, pedunculis validis lanatis, capitulis majnsculis, fl. radii paucis inconspicuis, corollæ tubo achæniisque villosis. (*Charact. ex exempl. Nov. Zealand.*)

Perhaps most nearly allied to the *Eurybia erubescens* of Tasmania in the foliage, though a very different plant. The pappus is in a measure double, for I observe small setæ mixed with or external to the longer : this is the case with various Tasmanian species, and much diminishes the value of the characters distinguishing *Olearia* from *Eurybia*.

The Auckland Island specimens have large and very obtusely dentate leaves, but are in other respects so similar to those from the mountains of the north Island of New Zealand (collected by Mr. Colenso), that I have little hesitation in considering them specifically the same.

p. 29. Add

1. ERECHTITES prenanthoides, DC.; Prodr. vol. vi. p. 296 (in Hb. Gunn).

HAB. Auckland Island, M. le Guillou.

I have examined a specimen of this plant in Mr. Gunn's herbarium (collected by M. Le Guillou, an officer of Admiral D'Urville's expedition): it is small but similar; and much larger ones from various parts of New Zealand appear to be specifically the same with others from Tasmania.

p. 63. Add

4. VERONICA salicifolia? Forst.; Prodr. p. 3. Benth. in DC. Prodr. vol. x. p. 459.

HAB. Auckland Island : M. le Guillou (in IIb. Gunn).

Specimen in fruit only, but I think referable to this species: it is not an uncommon plant in various parts both of the Northern and Southern Islands of New Zealand.

p. 68. URTICA australis, Hook. fil. Add :- foliis nunc omnibus oppositis, pedunculis floriferis petiolo æquilongis simplicibus ramosisve, floribus laxe densiusve aggregatis, masculis subterminalibus pilosiusculis, fœmineis glabratis.

The above additions to the published characters are supplied from a specimen gathered in Lord Auckland Island by M. le Guillou, and given by him to Mr. Gunn of Tasmania.

p. 69. THELIMYTRA? uniflora.-After the description add :-folio solitario tereti canaliculato.

A specimeu of this plant, collected by M. le Guillou in Lord Auckland Island, is in no better state of flower than those I gathered, the leaf is, however, in good preservation and similar to that of some other *Thelimytræ*.

p. 71. Orchid. "dubii generis" n. 8.

1. LYPERANTHUS Antarcticus, Hook. fil.; folio lauceolato acuminato, perianthii foliolo dorsali galeato acuto lateralibus interioribusque parvis anguste linearibus, labello recurvo marginibus erectis disco plicato sub 6-glanduloso.

Though somewhat different in habit from the New Holland species, I do not think that this can be generically separated from them. The flowers are shorter and less expanded, and the upper sepal larger and more galeate, the remainder smaller than in its congeners. The description is completed from M. le Guillou's specimens, in which the foliage is imperfect.

p. 80. 3rd line from bottom :- for "Juneus exiguus" read "Juneus inconspicuus."

p. 80. To Juncus, add

3. JUNCUS planifolius, Br.; Prodr. p. 259. Fl. Antarct. Pt. 2. p. 358. HAB. Auckland Island: M. le Guillou (in Hb. Gunn).

Specimens very diminutive, but, I think, clearly referable to this species, which is not uncommon throughout the Islands of New Zealand.

p. 84. LUZULA crinita.-Add to Habitats :--M'e Quarrie Island. (Hb. Hook.)

p. 119. 5th line from bottom, for "Bruch and Schimper" read "Nees and Hornschuch."

p. 122. In remarks on Leptostomum gracile ;---the L. Bridgesii, Wils. MS., is L. splachnoides, Hook. and Arn.

p. 123. SPLACHNUM octoblepharum, add synon .: -- S. plagiopus, Mont. in Voy. au Pole Sud, Bot. Crypt. p. 285.

p. 124. To RACOMITRIUM, add

2. RACOMITRIUM microcarpum, Brid.; Mont. l. c. p. 284.

HAB. Auckland Island; barren: M. Hombron.

p. 128. Genus 11. SPRUCEA, for "Brid." read Hook. fil. et Wils.

, p. 130. After DICRANUM add

5. DICRANUM dichotomum, Brid.; Mont. l. c. p. 298.

HAB. Auckland Island; barren: M. Hombron.

p. 130. To CAMPYLOPUS, add

3. CAMPYLOPUS atro-virens, De Notaris; Mont. l. c. p. 300.

HAB. Auckland Island; barren : M. Hombron.

p. 132. To POLYTRICHUM, add

2. POLYTRICHUM juniperinum, Willd.; Mont. l. c. p. 313.

HAB. Auckland Island: M. Hombron.

p. 142. For HYPNUM Terræ-Novæ, Brid., var. B., substitute

20. HYPNUM *limatum*, Hook. fil. et Wils.; caule humili prostrato vage ramoso, ramis subfastigiatis, foliis subsecundis lanceolato-acuminatis integerrimis enerviis, capsula subcrecta, operculo conico.

Dioienm. Rami breves, erecti. Folia conferta, suberceta, membranacea, e basi lata gradatim angustata, longe acuminata, subpilifera, siecitate mitida, luteola; perichætialia erecta, acuminata, pilifera. Seta 3-4 liu. longa, rubra. Capsula suberceta, curvula. Operculum majusculum, conicum, acutinsculum, rubellum.

Allied to *H. acutifolium*, nob.; but the leaves are narrower, more membranaceous, tapering gradually upwards from a broad base; and the areolæ are larger.

p. 143. To HOOKERIA, add

5. HOOKERIA crispula, Hook. fil. et Wils.; Lond. Journ. Bot. vol. fii. p. 550. Mont. l.e. p. 320. HAB. Auckland Island; barren: M. Hombron.

p. 153. After JUNGERMANNIA vertebralis, add

27 bis. JUNGERMANNIA punicea, Nees; Mont. l. c. p. 261.

HAB. Auckland Island: M. Hombron.

Fuegia, the

- p. 153. To JUNGERMANNIA Urvilleana, add syn :- J. abbreviata, Hook.fil. et Tayl. in Lond. Journ. Bot. vol. iii. p. 374.
- p. 156. After JUNGERMANNIA planiuscula, add 37 bis. JUNGERMANNIA connata, Sw.; Mont. l. c. p. 256. HAB. Auckland Island : M. Hombron.
- p. 157. After JUNGERMANNIA fissistipa, add 43 bis. JUNGERMANNIA amphibolius, Nees; Mont. l. c. p. 352. HAB. Auckland Island : M. Hombron.
- p. 159. JUNGERMANNIA hippuroides is J. capillaris, Sw., B. minor, Lchm. Lind. et Gottsche, Syn. Hep. p. 213.
- p. 159. After JUNGERMANNIA albula, add

50 bis. JUNGERMANNIA filamentosa, Lehm. et Lind.; Mont. l. c. p. 246.

HAB. Auckland Island : M. Hombron.

p. 160. After JUNGERMANNIA nutans, add

54 bis. JUNGERMANNIA adnexa, Lehm. ct Lind.; Mont. l. c. p. 243.

54 ter. JUNGERMANNIA decrescens, Lehm. et Lind.; Mont. l. c. p. 243. t. 19. f. 4.

HAB. Auckland Island: M. Hombron.

- p. 160. JUNGERMANNIA hirsuta is J. ochroleucu, Spr.; Gottsche, Nees et Lind. Syn. Hep. p. 240.
- p. 160. JUNGERMANNIA mollissima, is J. tomentella, y. Gottsche, Necs and Lind. Syn. Hep. p. 237.
- p. 162. JUNGERMANNIA elegantula is MADOTHECA Stangeri, Gottsche, Nees, and Lind. Syn. Hep. p. 280.
- p. 165. After JUNGERMANNIA scandens, add

71 bis. JUNGERMANNIA gracilis, Nees; Mont. l. c. p. 223.

HAB. Auckland Island; D'Urville.

p. 167. After JUNGERMANNIA plicatitoba, add

77 bis. JUNGERMANNIA cucullata, Nees; Mont. l. c. p. 218.

HAB. Auckland Island : M. Hombron.

- p. 177. Amongst synonyms to XIPHOPHORA Billardieri, dele "CTENODUS, Kütz."
- p. 180. RHODOMELA glomerulata, Mont., is POLYSIPHONIA botryocarpa, nobis.
- p 184. After JANIA insert

1. MELOBESIA verrucata, var. Antarctica, vide Part II. p.482.

p. 191. After CALLITHAMNION gracile, add

PLATE LXXXVIII. Fig. 1.—1, plant of the natural size; 2, ramulus; 3, ditto with spharospores: 4, articuli of ditto :—very highly magnified.

p. 193. After ULVA latissima add

1. ZIGNOA clathrata, Trevis.; Mont. l. c. p. 30. Enteromorpha, auct.

HAB. Auckland Island; D'Urville.

p. 196. STEREOCAULON Argus.

I have examined specimens of *S. ramulosum*, approaching this so very closely, that Mr. Churchill Babington inclines to consider the plants as varieties of one species.

546

p. 197. CENOMYCE ecmocyna, var. gracilis, is rather the C. sparassa; but the specimens are not very satisfactory.

p. 198. STICTA Freycinetii; these specimens probably belong to the S. Delisea, Fée., and differ from the true S. Freycinetii in the flatter (not concave) sessile apothecia.

p. 198. After STICTA Menziesii, add

5 bis. STICTA Richardi, Mont.; in Voy. au Pole Sud, p. 187.

HAB. Lord Auckland's group : M. Hombron.

p. 198. For 6. STICTA Richardi, substitute

6. STICTA Billardieri, see p. 527.

I very much doubt the *S. Richardi*, Mont., being anything more than a larger state of this plant. It is a very frequent and most variable inhabitant of New Zealand.

p. 199. After PARMELIA sphinctrina, add

3. PARMELIA enteromorpha, Ach.; P. physodes, var. vittata, Mont. in Voy. au Polc Sud, Bot. Crypt. p. 183.

HAB. Lord Auckland's group and Campbell's Island; not uncommon.

4. PARMELIA diatrypa, Ach.; P. physodes, var. Mont. l. c.

HAB. Lord Auckland's group; on trunks of trees.

p. 199. For PARMELIA rubiginosa, Ach., read

PARMELIA Mariana, Fries? Syst. Orb. Veg. pp. 245 and 284 (fid. Bab.).

Of the present plant the Rev. C. Babington remarks, "This seems to differ from *P. rubiginosa*, not only in general habit, but most especially in the apothecia being black : the scales, too, are singularly appressed; and the hypothallus is more carbonaceous. If a described plant, it is either *P. pellita*, Ach., or *P. Mariana*, Fries. The Acharian plant is barren, whence the thallus of the fertile might differ from this. Fries' plant exactly agrees in the apothecia and hypothallus; whilst the variation of the thallus to me seems caused by his specimens being more perfect."—*C. Babington*.

p. 199. After LECANORA Parella, add

5. LECANORA varia, Ach.; Lich. Univ. p. 377. Engl. Bot. t. 1666.

HAB. Lord Auckland's group; on bark of trees.

p. 200. For LECIDEA geomæa, substitute

1. LECIDEA papillata, Fries; Lich. Europ. p. 336.

"I have little doubt of this being the plant of Fries, judging from the description."-C.Babington.

p. 200. Add

3. LECIDEA parasema, Ach.; Syn. Lich. p. 17. Schærer, Lich. Helvet. n. 197-199. L. Lightfootii, Engl. Bot. t. 1457.

HAB. Lord Anekland's group; on trunks of Dracophyllum.

p. 200. Add

1. VERRUCARIA *punctiformis*, Ach.; *Syn. Lick.* p. 87. V. stigmatella, *Engl. Bot.* t. 1891. HAB. Lord Auckland's group; on trunks of trees.

p. 200. PORINA granulata, Hook. fil. and Tayl., is probably a state of Lecanora tartarea.

PART II.

- p. 212. 23rd line, for "granitic" read tertiary.
- p. 228. To HAMADRYAS, add

4. HAMADRYAS *paniculata*, Hook. fil.; foliis longe petiolatis rotundatis sub-5-lobis, lobis obtusis crenatis supra glabratis reticulatis subter parce sericeo-tomentosis, scapo gracili masculo laxe paniculatim ramoso multifloro superne tomentoso.

HAB. Staten Land; Webster.

Petioli (exemplare unico) 4 une. longi, glabri, validi. Folia concava, 2 une. lata, coriacea, inæqualiter lobata, vix ad medium fissa, lobis grosse et obtuse erenatis. Scapi folio longiores; masculi parce sericei, ramis paucis inæquilongis multifloris; fœminei (manci) pauciflori. Flores ut in II. argentea.

A distinct looking plant, both in the foliage and compound panicle, from any of its congeners. The specimens are in a very poor state.

- p. 241. 4th line from bottom, for "S. australis" read S. pinnatifida.
- p. 253. OXALIS enneaphylla, Cav.; add to the Habitat :- Strait of Magalhacns; MM. Hombron and Jacquinot.
- p. 274. GUNNERA Magellanica:—the Colombian plant alluded to as probably identical with this, is the G. pilosa, H.B.K.
- p. 277. 13th line from bottom, for "not one" read but one.
- p. 278. After MONTIA, add

2. LYALLIA, Hook. fil.

Calyx persistens, 4-partitus, lobis subæqualibus obtusis. Petala, stamina, ovariumque non suppetebant. Fructus, utriculus globosus, carnosus, apiculatus, venosus (stomatibus instructus), 1-locularis. Semina 3, quorum 2 abortiva, e fundo utriculi orta, funiculis validis ascendentibus affixa: semen maturum orhicularireniforme, compressum; testa crustaeca, subtiliter tuberculata; albumine carnoso v. subfarinaeco; embryone peripherico.—Herba sufficutescens, Insulæ Kerguelen, dense cæspitosa, glaberrima, ramosissima; ramis teretibus, foliis densissime imbricatis tectis. Pedunculi breviusculi, solitarii, terminales, basi bracteis connatis lanceolatosubulatis aucti. Flores verosimiliter valde inconspicui. Fructus parvus inter folia fere occlusus.—Genus dicatum Doetori D. Lyall, amico meo periplique Antaretici participi, assiduo solertique plantarum indagatori.

1. LYALLIA Kerguelensis, Hook. fil. (TAB. CXXII.)

HAB. Kerguelen's Land; forming large tufts in barren places, but very local.

Radices lignosi, descendentes. Rami perphurimi, densissime fastigiati, in cæspites globosas dispositi, 3-5 une. longi, strieti v. eurvati, $\frac{1}{4}$ une. diametro. Folia seeus ramos densissime imbrieata, pluriseriata, alterna, ramo multoties angustiora, ereeta, incurva, linearia, sessilia, subconcava, apice rotundata, obscure irregulariter sinuato-dentata, 3-5-costata (costa in nervos validos parallelos infra apicem evanidos divisa), coriacea v. subchartacea, persistentia, sieca pallida, terminalia pallide virescentia, 2-3 lin. longa, ter quaterve longiora quam lata. Pedunculi terminales, solitarii, folio breviores, erecti, infra florem dilatati, basi bibraeteati. Bracteæ pedunculis æquilongæ, basi connatæ, acuminatæ. Calyæ fructiferus utriculo brevior, e foliolis 4 (mnc 5?) ovatis obtusis 5-nerviis utriculo appressis constans. Petala et stamina ignota, sed (ab indole calyeis discique) verosimiliter perigyna. Utriculus $\frac{1}{2}$ -1 lin. longus. Semen utriculum fere implens.

This is certainly, in its present state, the most obscure and, except the *Pringlea*, the most remarkable plant of Kergnelen's Land. I have placed it provisionally amongst *Portulaceæ*, knowing no other order with which it has any equally direct affinity. There is one plant to which it bears, at any rate, a very close resemblance if nothing more, the *Pycnophyllum molle*, Remy (Ann. Sc. Nat., 3rd Ser. vol. v. p. 355. t. 20. f. 2–8), of the Bolivian Andes; for a fragment of the original specimen of which (preserved in Herb. Mus. Paris) I am indebted to the liberality of M. Decaisne. In *Pycnophyllum*, however, the leaves are truly opposite and connate, and the capsule three-lobed, if not three-valved.

This highly curious genus, coming from the most interesting island visited by the Antarctic Expedition, will serve to commemorate in some slight degree the important services rendered to Botany by my zealous friend and co-operator, Dr. Lyall, R.N.

PLATE CXXII.—*Fig.* 1, a leaf; *fig.* 2, bracteæ, peduncle, and fruit; *fig.* 3, utriculus removed from the calyx; *fig.* 4, vertical section of the same, showing the ripened and abortive seed; *fig.* 5, seed and funiculus; *fig.* 6, seed, with the testa removed; *fig.* 7, embryo:—all *magnified*.

- p. 292. Of the Valdivian specimens alluded to as belonging apparently to this species, I have recently examined complete individuals, which prove them to be *M. imbricatum*, Poepp. The female flowers do not probably differ materially from those of *M. punctulatum*.
- p. 296. In description of Plate CIV. the figures of 9, 10, 11, and 12, referred to as taken from Bridges' Valdivian specimens, belong to *M. imbricatum*, Poepp.
- p. 304. CHILIOTRICHUM humile :---M. Planchon considers this species, together with the C. Kingii and C. Darwinii, as referable to the genus Nardophyllum, DC.
- p. 327. LEBETANTHUS Americanus:—this appears to be a true Prionotes, the placentation being the same as in the original *P. cerinthoides* of Tasmania, and the position of the braeteæ on the pedicel not affording a generic character. In two undoubted congeners from Tasmania, the ovules are attached to erect basal columns.
- p. 341. After CHENOPODIUM, add

2. BLITUM, L.

1. BLITUM (Orthosporum) *Antarcticum*, Hook. fil.; caule prostrato ascendente parce papilloso, foliis petiolatis deltoideo-ovatis obtusiusculis profunde irregulariter sinuato-dentatis lobis lobulatis utrinque petioloque papillosis, glomerulis sessilibus compositis axillaribus et in spicam terminalem foliosam dispositis, perigonii 3-phylli foliolis herbaceis post anthesin immutatis utriculum superantibus lineari-spathulatis dorso grosse papulatis, semine verticali orbiculari punctulato margine obtuso.

HAB. Staten Land; Webster (in Mus. Hort. Soc. Lond.).

Ramus 10-uncialis simplicissimus solum mihi notus. Pelioli unciales, foliis æquilongi. Flores minimi, dense aggregati.

Described from a solitary specimen in the Herbarium of the Horticultural Society of London, in the absence of any means of comparing it with its congeners in the Herb. Hook., of which the *Chenopodiaceæ* are now in the hands of M. Moquin Tandon for examination.

- p. 343. NANODEA muscosa :--- I am inclined to suspect, from certain circumstances connected with the locality of this plant, that it may be parasitical, like the *Thesium linophyllum*.
- p. 359. LUZULA, sp.—Of this species I have recently found more complete (hitherto mislaid) specimens in my collection, they may be thus described :—

2. LuzuLA Antarctica, Hook. fil.; pusilla, cæspitosa, foliis late lineari-subulatis concavis basin versus ciliatis, culmo gracili filiformi arcuato v. erecto, panieula ovata densissime lanata, bracteolis foliolisque perianthii subæqualibus superne searioso-membranaceis inferne medioque coloratis marginibus in lacinias piliformes fimbriato-laceras apicibus hyalinis, capsula elliptico-subrotundata perianthio dimidio breviore, stigmatibus 3 sessilibus filiformibus.

HAB. Hermite Island, Cape Horn; alpine rocks.

llabitus *L. spicatæ*, statura *L. arcuatæ* humilior. *Folia* uncialia, basi fere $\frac{1}{4}$ unc. lata, pleraque exemplaribus meis mancis superne glabrata v. glaberrima, basin versus ciliata. *Culmus* filiformis, 2-uncialis. *Panicula* $\frac{1}{3}$ unc. longa, late ovata. *Perianthii foliola* per totam longitudinem in lacinias foliolum longe superantes fissa; parte inferiore mediaque brunnea, coriacca, superiore hyalina.

Allied to *L. Peruviana* (of the Andes) in habit; but the leaves (in my specimens) are not ciliated and the perianth is of a different form, its leaflets being shorter, broader, not coriaceous, and subulate at the apices, and with very much more copious and longer ciliæ.

- p. 371. MUHLENBERGIA rariflora, Hook. fil.:—Nees (in Herb. Arnott) regards this as a species of Streptachne, H.B.K.
- p. 375. ARUNDO pilosa ; add to Habitats :- Fuegia, Good Success Bay ; Banks and Solander.
- p. 381. FESTUCA Fuegiana, a., is considered by Nees (fid. Herb. Arnott) a variety of Poa lanigera, Necs, in Martius Fl. Bras. p. 490.
- p. 392. LOMARIA alpina, Br.; add to synonyms:-L. pumila, Raoul, Choix de Plantes de la Nouvelle Zelande, t. 10. t. 2. f. A.
- p. 393. GLEICHENIA aculifolia; add to the Habitats :---Staten Land; Mr. Webster.
- p. 394. Lycorodium clavatum, var. Magellanicum; add synonym :- L. confertum, Willd. Sp. Pl. vol. v. p. 27. Hook, et Grev. in Bot. Misc. vol. ii. p. 372.
- p. 394. After Schlzæa, add

10. BOTRYCHIUM, Swartz.

1. BOTRYCHIUM Lunaria, Sw.; Syn. Fil. p. 171. Engl. Bot. t. 318.

HAB. Fuegia, Good Success Bay; in sandy places : Banks and Solander (in Herb. Mus. Brit.).

Identical with the European plant, which ranges in Europe from Iceland and Lapland to the Asturias. In North America it is only found in Hudson's Bay, Newfoundland, Canada, and the Rocky Mountains. I know of no habitat except this of Fuegia and Tasmania anywhere south of the north of Spain. It is apparently a very rare Fuegian plant.

- p. 403. ORTHOTRICHUM luteolum, Hook. fil. et Wils.—This approaches very closely the description of O. germanum, Mont. (in Ann. Sc. Nat., 3rd Ser. vol. iv. p. 121), a Chilian plant, but the leaves of which are said to be rather obtuse and reflexed at their margins.
- p. 408. After CAMPYLOPUS flexuosus, add

3. CAMPYLOPUS *rigidus*, Hook. fil. et Wils.; caule erecto subramoso rigidiusculo, foliis ovatolanceolatis acuminatis integerrimis, nervo latissimo.

HAB. Hermite Island, Cape Horn; on the summits of the hills.

ø

p. 409. TORTULA densifolia, Hook. fil. ct Wils., is evidently closely allied to Barbula mnoides, Schwaeg. Suppl. 1.310.

p. 410. POLYTRICHUM compressum, Hook. fil. et Wils.

Var. β ., foliis apices versus obscure serratis lamellatis, capsula longiore.

HAB. Hermite Island, Cape Horn; with var. a.

p. 418. HYPNUM subpilosum, Hook. fil. et Wils.; (character reformata) caule arcuato parce ramoso, ramis recurvis attenuatis, foliis cordato-ovatis acuminatis subpiliferis concavis striatis serrulatis ruptinerviis, capsula subrotunda cernua, operculo conico, seta scabra.

p. 449. After EXIDIA Auricula-Judæ, add

2. EXIDIA *flammea*, Berk.; aurantiaca, hemispherica, depressa, substipitata, margine crenulata subtus rugulosa minutissime verrucoso-spiculata, sporis oblongis basi curvatis. (TAB. CLXIV. Fig. III. left hand specimen.)

HAB. Hermite Island, Cape Horn; on dead wood amongst the snow.

Hemisphærica, leviter depressa, vel humore saturata planiuscula, brevissime stipitata; margine læviter erenata; subtus rugulosa, sub lente maxime augente subtiliter verrucosa, hic illic spiculata. *Sporæ* oblougæ, basi curvatæ.

Allied to *Exidia truncata*, but differing remarkably in its bright colour. When first taken out of spirit the hymenium is quite plane, but becomes depressed afterwards. It is doubtful whether the margin be crenate in the living plant, for it is not represented in the drawing made from the fresh specimen.

PLATE CLXIV. Fig. III .--- 2, (left hand figure) E. flammea, of the natural size.

p. 451. PEZIZA Kerguelensis.—The Hermite Island plant is *Exidia flammea*, Berk.; to which also the left hand figure of Plate CLX1V. Fig. III. 2, is referable. The right hand figure (1), which, however, is not represented sufficiently adnate, and the dissections, belong to *P. Kerguelensis*.

PLATES.

PLATE XVI.—The Tasmaniau flowering plant, figured at B, is another species, C. pumila, mihi (see Supplement).

PLATE XXI.-HELICHRYSUM prostratum, is II. bellidioides, Forst. (see Suppl.)

PLATE LXI. Fig. IV .- HYPNUM Terræ-Novæ is II. timatum, Hook. fil. et Wils. (see Suppl.)

PLATE LXXXII. B .- RANUNCULUS hydrophyllus, should be R. hydrophilus.

PLATE LXXXV.-HAMADRYA's tomentosa is H. argentea, Ilook. fil. (see Suppl.)

PLATE LXXXVII.—BERBER1S microphylla is a synonym of B buxifolia, Lam. (see Suppl.)

PLATE XCIII.—SAGINA subulata, D'Urv., is a synonym of Colobanthus subulatus (see Suppl.)

PLATE CI.-CALDASIA daucoides, Hook. fil., is a synonym of Oreomyrrhis andicola, Endl. (see Suppl.)

PLATE CIV.-Figs. 9, 10, 11, and 12, Myzodendron imbricatum, Poepp. (see Suppl.)

PLATE CXII.-MACRORHYNCHUS coronopifolius should be M. pumilus, DC. (see p. 324.)

PLATE CXV.—GENTIANA Magellanica should be G. Patagonica (see p. 328).

PLATE CXVI.-GAULTHERIA Antarctica, Hook. fil., is synonymous with G. microphylla, Hook. fil. (see p. 327.)

PLATE CXVIII.-OURISIA Antarctica, Hook. fil., is synonymous with O. breviflora, Benth. (see p. 335.)

PLATE CXX.-PRIMULA Magellanica, Lam., is a variety of P. farinosa (see p. 337).

PLATE CXXIX .- SISYRINCHIUM pumilum, Hook. fil., is a synonym of Tapeinia Magellanica, Juss. (see p. 353.)

PLATE CXXX.-ALOPECURUS Antarcticus, Vahl, is a variety of A. alpinus (see p. 370).

PLATE CXXXVIII.-POA Kerguelensis, Hook. fil., is TRIODIA Kerguelensis, Hook. fil. (see p. 379).

PLATE CLVII. Fig. VII.—JUNGERMANNIA cavispina, Hook. fil. et Tayl., is a variety of J. austrigena, Hook. fil. et Tayl. (see p. 431.)

PLATE CLXI. Fig. III.—For "JUNGERMANNIA reclinata," read "J. retusata (see p. 441).

PLATE CLXIV. Fig. III.-1, PEZIZA Kerguelensis, Berk., to which also the dissections, 3, 4, and 5, belong; 2, is EXIDIA flammea, Berk. (see Suppl.)

PLATE CLXIX. and CLXX.-MACROCYSTIS luxurians is a variety of M. pyrifera, Ag. (see p. 461.)