



JOHN EDWARD TAYLOR, PRINTER,  
LITTLE QUEEN STREET, LINCOLN'S INN FIELDS.

TO

SIR WILLIAM JACKSON HOOKER, K.H.,

D.C.L. OXON., F.R.S., F.A.S., F.L.S., ETC.,

This Little Work is Dedicated,

IN TESTIMONY OF THE MOST SINCERE RESPECT

AND AFFECTION,

BY HIS OBLIGED FRIEND,

THE AUTHOR.

## PREFACE.

---

SINCE the publication of Linnæus's 'Flora Zeylanica' and Burmann's 'Thesaurus Zeylanicus,' no work has appeared, with the exception of Moon's 'Catalogue,' professing to give a list of the plants indigenous to Ceylon.

Several species of Ceylon plants have been figured and described by Sir William Hooker in his 'Icones Plantarum' and other valuable publications, from specimens collected by the late Colonel and Mrs. Walker; and a good many Ceylon plants have been figured by Dr. Wight in his excellent 'Icones Plantarum Indiæ Orientalis.' Dr. Walker-Arnott has published descriptions of several kinds in his 'Pugillus' and elsewhere.

Most of the *Orchidaceæ* of Ceylon have been described by Dr. Lindley from specimens and drawings sent to him by Mr. Macrae.

My predecessor, Dr. Gardner, contributed to the 'Calcutta Journal of Natural History' descriptions of several interesting Ceylon plants.

References throughout the present work indicate where and by whom other species have been described.

Owing to the physical conformation of the island, which geologically belongs to the Metamorphic system, there is great diversity of climate.

The hills of the Central Province rise to an elevation of 8300 feet, with an annual rainfall of about 100 inches.

The low country of the south of the island possesses a humid climate, with a very equable temperature, and the rainfall is considerable, but not equal to that on the hills.

The low northern half of the island, shut off by the hills from the rains of the south-west monsoon, has a rainy season of very short duration, and is usually extremely dry during the remainder of the year.

The character of the island vegetation varies, as might be expected, in accordance with this difference of climatal conditions; for, whilst on the hills the flora resembles very much that of the Neilgherrries, in the humid south it is more akin to that of Sumatra and the Malay Archipelago; and in the dry north of the island it would seem to be nearly identical with that of the Coromandel coast.

The indigenous species enumerated in the present work amount to 2832, viz. :—

Dicotyledones . . . . .	1959
Monocotyledones . . . . .	648
Filices, Lycopodiaceæ, and Marsileaceæ . . . . .	225

Care has been taken not to multiply species unnecessarily, for observation has shown that the amount of variation is often considerable in plants affecting a large range of variation, and consequently of climate. Instances occur, as in the genera *Hortonia*, *Mappia*, *Turpinia*, *Evonymus*, *Elæodendron*, in which a more elevated locality produces a form or variety possessing a stouter habit and larger flowers than are observed in the same species when growing only a little above the level of the sea. These forms or varieties would probably be viewed by some botanists in the light of distinct though closely allied species, and they occupy, in fact, that debatable ground the difficulties and perplexities of which the practical naturalist alone knows, and which, in the opinion of many (and I may include myself among the number), are only to be got rid of by the adoption of the views enunciated by Mr. Darwin as regards the relationship of allied forms or species by descent from a common ancestor.

From the large extent of forest-land which has been and is now being appropriated to coffee cultivation, there is little doubt that some of the indigenous plants will in time become exceedingly rare, if not altogether extirpated, or exist only in the botanical garden, into which as many kinds as possible are being introduced. The obtrusive character, too, of a plant brought to the island,

about forty years ago, is also helping to alter the character of the vegetation up to an elevation of 3000 feet. The plant alluded to is the *Lantana mixta*, Linn., which appears to have found in Ceylon a soil and climate exactly suited to its growth, for it now covers thousands of acres with its dense masses of foliage, taking complete possession of land where cultivation has been neglected or abandoned, preventing the growth of any other plants, and even destroying small trees, the tops of which its subscandent stems are able to reach. The fruit of this plant is so acceptable to frugivorous birds of all kinds, that through their instrumentality it is spreading rapidly, to the complete exclusion, in spots where it becomes established, of the indigenous vegetation.

*Brucea Sumatrana*, *Crotalaria incana*, *Lagascea mollis*, *Cosmos caudatus* and *Asclepias Curassavica*, have become perfectly naturalized, and are common weeds in the warmer parts of the island.

In Newera Ellia, a few exotic plants, which were introduced with grass seed, have become naturalized; those I have observed are *Cerastium vulgatum*, *Hypericum humifusum*, *Trifolium repens*, *Verbascum Thapsus*, *Stachys arvensis*, *Plantago lanceolata*, and *Dactylis glomerata*.

It now remains to express my grateful acknowledgment for assistance rendered to me during the progress of this work.

Dr. Wight, immediately on my arrival in the island, gave me his most valuable help in naming the numerous specimens I found in the herbarium, which had been collected by my predecessors, and principally by Dr. Gardner.

Sir William Hooker has most kindly assisted me with his profound knowledge of the subject in determining the names of the Ferns.

To Dr. Joseph D. Hooker I am most deeply indebted for the invaluable aid he has been uninterruptingly giving me during the whole progress of this work, and without this aid I could not with any satisfaction have undertaken it.

My cordial thanks are due to Dr. Thomas Anderson for having undertaken the identification and description of the *Acanthaceæ*, an Order of plants to which he was at the time devoting particular attention, and was studying in the herbarium at Kew.

To the Rev. S. O. Glenie, F.L.S., Episcopal Colonial Chaplain at Trincomalie, and to Mr. W. Ferguson, F.L.S., Civil Engineer, I am

under great obligations for frequent valuable communications, and for specimens of plants from their respective places of residence. It gives me great pleasure, too, to record the uniform courtesy and kindness with which the gentlemen, holding the important and influential positions of Government Agents, have assisted me in carrying out the objects I had in view, when making botanical tours through their respective provinces or districts.

I would wish to associate with this little book the names of Harmanis de Alwis Modliar, and of Don Seman De Silva Aratchy, native officers, employed during many years in this establishment, the former as Draftsman, and the latter as Principal Plant-collector, to whose intelligent and hearty co-operation is due much of my success in procuring new and rare species of plants from the jungles.

Some apology may be considered necessary for the length of time that has elapsed since the present work was commenced, but a large portion of that time has been occupied in researches connected with it, and these researches have, it is hoped, tended to make it more useful and more worthy of being laid before the scientific world.

Part 1, containing 80 pages, was published in 1858.

Part 2, from p. 81 to p. 160 inclusive, was published in 1859.

Part 3,     "   161     "   240             "             "             1860.

Part 4,     "   241     "   320             "             "             1861.

Part 5,     "   321     "   483             "             "             1864.

*Royal Botanic Garden, Peradenia, Ceylon,  
August, 1864.*

# ENUMERATION OF CEYLON PLANTS.

## I. RANUNCULACEÆ.

### 1. NARAVELIA, DC.

1. **N. Zeylanica**, DC. (H.f. et T. Fl. Ind. p. 3, cum syn.)—c.p. 1009.  
HAB. Not very uncommon in the warmer parts of the island. Nom. vulg. "Wayoopaloo" or "Narawella."

### 2. CLEMATIS, Lin.

1. **C. smilacifolia**, Wall. (H.f. et T., l. c. p. 6, cum syn.)—c.p. 2690.  
HAB. At an elevation of about 3000 feet, at Allagalla and Ambagamowa, in the Central Province.
2. **C. Gouriana**, Roxb. (H.f. et T., l. c. p. 8, cum syn.)—c.p. 680.  
HAB. Newera Ellia, and other elevated parts of the Central Province.

### 3. THALICTRUM, Lin.

1. **T. glyphocarpum**, W. et A. (H.f. et T., l. c. p. 16, cum syn.)—c.p. 2574.  
HAB. On rocks, Newera Ellia, Horton Plains, etc.

### 4. ANEMONE, Lin.

1. **A. rivularis**, Ham. (H.f. et T., l. c. p. 23, cum syn.)—c.p. 1011.  
HAB. Newera Ellia.

### 5. RANUNCULUS, Lin.

1. **R. sagittifolius**, Hook. (H.f. et T., l. c. p. 30, cum syn.)—c.p. 520.  
HAB. Swampy spots at Newera Ellia and Horton Plains.
2. **R. Wallichianus**, W. et A. (H.f. et T., l. c. p. 38, cum syn.)—c.p. 1012.  
HAB. Very common on the more elevated parts of the Central Province, from 5000 to 8000 feet.



## II. DILLENiaceæ.

## 1. DELIMA, Lin.

1. **D. sarmentosa**, Lin. (H.f. et T., l. c. p. 61, cum syn.)—c.p. 1015.

HAB. Southern parts of the island, up to an elevation of 1000 feet.  
Nom. vulg. "Korasawel."

## 2. TETRACERA, Lin.

1. **T. lævis**, Vahl, (H.f. et T., l. c. p. 62, cum syn.)—c.p. 1016.

HAB. Between Colombo and Galle; not very common.

## 3. ACROTREMA, Jack.

CHAR. EMEND.—*Sepala et petala* 5. *Stamina* indefinita, 15 vel plura; *filamentis* liberis, erectis, pluriserialibus, in 3 fasciculis cum ovariis alternantibus plus minus aggregatis; *antheris* sublinearibus, ab apice ad medium vel ultra lateraliter dehiscentibus. *Ovaria* 3, 2-vel multi-ovulata, sutura ventrali plus minus cohærentia; *stylis* subulatis, recurvatis; *stigmatibus* minuto. *Carpella* irregulariter dehiscentia. *Semina* membranaceo-arillata; *testa* crustacea, cancellata.—*Herbæ* perennantes, subacaules; rhizomate sublignoso; *foliis* magnis, parallelis-penninerviis, parce glanduloso-punctatis, denticulatis; *petiolis* marginibus magnis, vaginantibus, stipularibus, denique deciduis munitis; *floribus* racemosis, flavis, expansis, 4-8 lin. latis; *pedicellis* plus minus elongatis; *bracteis* magnis, sæpissime bidentatis.

*Folliculis* polyspermis.

† *Racemis* brevi-pedunculatis.

‡ *Foliis* indivisis, obovato-oblongis.

1. **A. uniflorum**, Hook. (Ic. Pl. t. 157; H.f. et T., Fl. Ind. p. 64); *foliis* superne glabris vel inter nervos pilis paucis rigidis munitis, subtus pallidioribus, glabris, costa nervisque dense strigosis; *pedicellis* (siccis) adpresse pilosis.

Var. *a. petiolaris* (c.p. 265); *foliis* angustioribus, crenatis vel crenatodentatis, apice acuto rotundato vel retuso; *staminibus* 30-50.

Var. *β. rotundata* (c.p. 1014 et 3484); *foliis* rotundatis brevi-petiolatis, margine subintegro; *staminibus* circiter 30.

HAB. Not uncommon in the Central and Southern Provinces, at an elevation of from 2000 to 4000 feet, occurring, like its congeners, in damp shady places, generally on the banks of streams.

*Folia* 4-9 poll. longa, 1-3 poll. lata, *petiolis* (in var. *a*) 1-1 $\frac{3}{4}$  poll. longis.

A very variable species, both as to the size as well as the form of the leaves, and the number of their primary veins. I should not be surprised if a more critical examination of fresh specimens proved vars. *a* and *β* to be distinct species.

2. **A. bullatum**, Thw. (c.p. 239); *foliis* bullatis, utrinque (præcipue superne inter nervos et ad marginem crenato-serratum) mollissime sericeo-pubescentibus; *pedicellis* (siccis) adpresse sericeo-pilosis.

HAB. In similar places with the foregoing, and often mixed with it.

*Folia* 2-6 poll. longa,  $\frac{3}{4}$ -2 poll. lata, *petiolis*  $\frac{1}{2}$ - $\frac{3}{4}$  poll. longis.

This species is well distinguished by its bullate, softly pubescent leaves, characters better observed in fresh than in dried specimens.

3. **A. sylvaticum**, Thw. (c.p. 2659); *foliis* utrinque minute pilosis,

subtus costa nervisque pilis rigidis patentibus hirsutis, apice rotundatis, basi profunde cordatis, margine subintegro; *petiolo* brevi; *pedicellis* brevibus, patentim pilosis; *staminibus* circiter 50.

HAB. Damp forest at Eknalagodde, near Ratnapoora, March, 1853.

*Rhizoma* breve; *folia* membranacea, 5-8 poll. longa, 3-4 poll. lata.

4. **A. Walkeri**, Wight, MSS. (c.p. 345, 446, 694); *foliis* crenato-dentatis, subauriculatis, utrinque præcipue superne versus marginem et subtus ad costam nervosque pilosis; *pedicellis* patentim pilosis; *staminibus* circiter 15.

HAB. Not uncommon in the Central Province, at an elevation of from 2000 to 4000 feet.

*Folia* 2-5 poll. longa,  $\frac{3}{4}$ -2 poll. lata, *petiolo*  $\frac{1}{2}$  poll. longo; *pedicelli* elongati.

The leaves of this pretty species are frequently marked on the upper surface with a paler area occupying a more or less wide space on either side of the costa and primary veins; the same peculiarity is sometimes observed in other species.

‡‡ *Foliis indivisis, lanceolatis, subauriculatis.*

5. **A. intermedium**, Thw. (c.p. 3114); *foliis* acutis, crenato-dentatis, supra medium versus basin graduatim valde angustatis, superne ad costam marginemque et parce inter nervos hirsutis, subtus costa nervisque adpresse pilosis; *pedicellis calycibusque* patentim rigido-hirsutis; *staminibus* circiter 40.

HAB. Banks of streams in the Ambagamowa District, at no great elevation, February, 1854.

*Rhizoma* elongatum; *folia* 5-12 poll. longa,  $1\frac{1}{2}$ - $2\frac{1}{4}$  poll. lata, *petiolo*  $1-1\frac{1}{2}$  poll. longo.

6. **A. lanceolatum**, Hook. (H.f. et T. Fl. Ind. p. 65.)—c.p. 2660; *foliis* angustissime lanceolatis, acutis, sinuato-dentatis, subglabris, subtus costa nervisque adpresse brevi-hirsutis; *pedicellis* patentim parce-hirsutis; *staminibus* circiter 40.

HAB. Kittool Gulle, March, 1853. I do not know where Dr. Wight found his specimens.

*Rhizoma* breve; *folia* 4-12 poll. longa,  $\frac{1}{2}$ - $1\frac{1}{4}$  poll. lata, *petiolo*  $\frac{1}{2}$ -1 poll. longo.

7. **A. Gardneri**, Thw. (c.p. 253); *foliis* anguste-lanceolatis vel subspathulatis, glabris, margine subintegro, subtus costa nervisque adpresse pilosis; *pedicellis* elongatis, adpresse-pilosis; *staminibus* circiter 15.

HAB. Common in the district between Galle and Ratnapoora, on the banks of streams.

*Folia*  $1\frac{1}{2}$ -3 poll. longa, 3-7 lin. lata, *petiolo* 3-6 lin. longo.

‡‡‡ *Foliis decompositis.*

8. **A. Thwaitesii**, H.f. et T., Kew Journ. Bot. vol. 8, p. 241, t. iv. A.—c.p. 3364.—Vars. *a* and *β*.

HAB. Dolosbage District, at no great elevation, February, 1855.

9. **A. dissectum**, H.f. et T., l. c. p. 242, t. iv. B.—c.p. 3393; et var. *pinnatifida*.

HAB. Near Hellessee, in the Pas-dun Corle, at no great elevation, May, 1855.

‡‡ *Racemis longe-pedunculatis.*

10. **A. lyratum**, H.f. l. c. p. 242.—c.p. 3392.

HAB. Near Nellowe, in the Hinidoon Corle, at an elevation of about 1000 feet, May, 1855.

4. **SCHUMACHERIA**, Vahl.

*Sepala* 5 (2 exteriora minora). *Petala* 5. *Stamina* indefinita, pluriserialia, monadelphica, unilaterali-posteriora; *filamentis* in columnam brevem obliquam semicylindricam coalitis, apicibus supremis liberis; *antheris* lineari-oblongis, obtusis, apiculatis, bilocularibus, lateraliter dehiscentibus. *Ovaria* 3 (2 anteriora), discreta, dense pilosa, uniovulata; *stylis* filiformi-subulatis; *stigmatibus* minuto. *Carpella* indehiscentia. *Semen* erectum, subglobosum, basi arillo parvo membranaceo munitum; *testa* crustacea.—*Frutices erecti*; ramis *rigidis, flexuosis*; foliis *coriaceis, conspicue penniveniis*; petiolo *basi latiuscula, vaginante*. *Spicis axillaribus vel terminalibus*; floribus *sessilibus, secundis, bibracteolatis, flavis, expans. 6–9 lin. latis*.

*Note*.—The generic characters of this and the foregoing genus have been principally copied from the 'Flora Indica' of Drs. Hooker and Thomson, such additions or slight alterations only having been made as the examination of fresh and more abundant specimens of the several species afforded the favourable opportunity of doing.

1. **S. castaneæfolia**, Vahl, (H.f. et T., l. c. p. 66, cum syn.)—c.P. 232, 248.

*HAB.* Common in the Saffragam and Galle Districts, at no great elevation. *Nom. vulg.* "Kakareewara."

2. **S. angustifolia**, H.f. et T., l. c. p. 66.—c.P. 2992.

*HAB.* Abundant in the Hinidoon Corle.

Not so straggling in its growth as the preceding species, and with a good deal smaller flowers.

3. **S. alnifolia**, H.f. et T., l. c. p. 66.

*Var. a. scabra*; foliis hirsutis rigida pauca scabris.—c.P. 9.

*Var. β. dentata* (Wight, MSS.); foliis grosse et acute dentatis, subtus petiolisque hirsutis.—c.P. 339.

*Var. γ. subglabra*; foliis subtus subglabris.—c.P. 397, 2416.

*HAB.* Vars. *a* and *β*, in forests in the Ambagamowa District. *Var. β*, which has sharply-dentate leaves, and these more hirsute beneath and with coarser reticulation, owes its peculiarity, I believe, to growing in a more damp and shady locality. *Var. γ* occurs in forests in the Hunasgiri District.

5. **WORMIA**, Rottb.

1. **W. triquetra**, Rottb. (H.f. et T., l. c. p. 67, cum syn.)—c.P. 1013.

*HAB.* Common in the moist, warmer parts of the island, up to an elevation of 2000 feet. *Nom. vulg.* "Deeyapara."

*Arbor* mediocris; foliorum *petiolis* canaliculatis, stipulari-marginatis (marginibus vaginantibus, cito deciduis), basi externe appendicula carnosa prominente hippocrepiformi semicinctis; *floribus* secundis; *sepalis* post anthesin parce accrescentibus, imprimis arcte imbricato-convergentibus, denique fructu maturo apertis vel reflexis; *petalis* plicato-rugosis, albis, basi pallide viridescens, caducis; *staminibus* erectis, luteis, circiter 3-serialibus, subæquilongis, persistentibus; *ovariis* 5, circiter 8-ovulatis; *ovulis* biserialibus; *carpellis* indehiscentibus, 1-2-spermis; *seminibus* arillatis, in pulpa pellucida gelatinosa nidulantibus.

The carpels of this, the original species of the genus, being indehiscent, it would seem doubtful whether some of the other species now included in the genus should be retained in it, or Blume's genus *Capellia* be restored for their reception.

## 6. DILLENIA, Lin.

1. **D. speciosa**, Thunb. (H.f. et T., l. c. p. 69, cum syn.)—c.p. 2961.

HAB. Common in the warmer parts of the island, up to an elevation of 2000 feet; most frequent on the banks of streams. Nom. vulg. "Hondapara."

## Sect. COLBERTIA, Salisb.

2. **D. retusa**, Thunb. (*Wormia retusa*, H.f. et T., l. c. p. 68, cum syn.)—c.p. 2960.

HAB. In similar situations with the preceding species, but by no means so abundant. Nom. vulg. "Godapara."

*Arbor mediocris; petalis albis; staminibus numerosis, 5-6-serialibus (externis graduatim brevioribus), pallide sanguineis; antherarum loculis lateralibus, albis; ovariis 5, 30-35-ovulatis; stylis filiformi-subulatis, recurvatis; ovulis 4-serialibus; fructu maturo pendulo, croceo, 1¼-1½ poll. diam.; carpellis indehiscentibus, calyce valde aucto carnosio arcte inclusis; seminibus exarillatis, in pulpa pellucida gelatinosa nidulantibus.*

There can be no doubt that this handsome species belongs to the *Colbertia* section of *Dillenia*, though it seems to differ from the others in having white instead of yellow petals. There is an indication of the stamens being collected into five groups alternating with the carpels.

† 3. **D. integra**, Thunb. (*Wormia integra*, H.f. et T., l. c. p. 68, cum syn.)

I know nothing of this plant, and suspect there must be some error in a fourth species of these large Dilleniaceous trees being attributed to Ceylon.

## III. MAGNOLIACEÆ.

## 1. MICHELIA, Lin.

1. **M. Nilagirica**, Zenk. (H.f. et T., l. c. p. 82, cum syn.)

Var. *a. ovalifolia*, Wight; petalis cum sep. sæpissime 9.—c.p. 141, 424.

Var. *β. Walkeri*, H.f. et T. (*M. Walkeri* and *M. glauca*, Wight); petalis cum sepalis sæpissime 12.—c.p. 2636 and 576, 299, 338, 423, 617.

HAB. Central Province, at an elevation of 3000 to 8000 feet. Nom. vulg. "Walsappoo."

In Ceylon the most typical form of this plant is that called *M. Walkeri* by Dr. Wight, having lanceolate leaves more or less glaucous on the under surface, and most commonly twelve petals. Between this and the oval-leaved form a gradual transition may apparently be observed, and, although the latter has most commonly nine petals, and these wider, yet, like the authors of the 'Flora Indica,' I have been quite unable to discover any permanent characters on which two separate, clearly-defined species can be founded. The form occurring on the Neilgherries, as shown in a specimen sent to me by Dr. Hooker, differs from any I have seen growing in Ceylon in its larger leaves and much larger petals.

*M. Champaca*, L., is very common in cultivated ground, but I have never seen it undoubtedly wild in the forests.

## IV. SCHIZANDRACEÆ.

## 1. KADSURA, Juss.

1. **K. Wightiana**, Arn. (H.f. et T., l. c. p. 84, cum syn.)—c.p. 1028.

HAB. Central Province, at an elevation of 1500 to 4000 feet.

*Ramulis foliisque junioribus rufescentibus; pedunculis sæpe petiolos multo superantibus; floribus pallide viridi-ochraceis; ovariis 3-5, sæpissime 4-ovulatis; toro conico.*

## V. ANONACEÆ.

### 1. SAGERÆA, Dalzell.

1. **Thwaitesii**, H.f. et T., l. c. p. 93.—C.P. 2702.

HAB. Ambagamowa and Saffragam Districts, at an elevation of about 1000 feet, February, 1853.

*Flores pallide crocei. Antheræ rufescentes. Carpella subsessilia, nucis Avellanæ magnitudine vel minora, pluri- vel pauci-sperma. Semina compressa, sulco circumcincta.*

### 2. UVARIA, Lin.

1. **U. semecarpifolia**, H.f. et T., l. c. p. 97.—C.P. 244.

HAB. Central and south-eastern parts of the island, at an elevation of 1000-3000 feet.

*Flores fulvo-viridescentes. Sepala interiora convergentia.*

2. **U. macrophylla**, Roxb. (H.f. et T., Fl. Ind. l. c. 97, cum syn.)—C.P. 3527.

HAB. South-east of the island, not uncommon.

The fruit is eaten by the Cinghalese, and was described to me as nearly black when ripe, with the interior red. The unripe fruit in the Ceylon specimen is sparingly tomentose.

3. **U. sphenocarpa**, H.f. et T., l. c. p. 99.—C.P. 1045.

HAB. Central and eastern parts of the island, up to an elevation of 1500 feet.

*Flores fulvo-viridescentes. Ovaria elongata, prismatica, dimidio inferiore circiter 10 ovula biserialia includentia. Stigmata subplana, lateribus involutis.*

4. **U. macropoda**, H.f. et T., l. c. p. 101.—C.P. 2544.

HAB. Similar localities with the foregoing.

*Flores rubiginosi. Carpella matura sanguineo-coccinea. Semina compressa, lævia, albida, oblonga, 5 lin. longa.*

5. **U. Narum**, Wall. (H.f. et T., l. c. p. 102, cum syn.)—C.P. 1032.

HAB. Common in the Central Province, up to an elevation of 4000 feet.

*Flores rubiginosi, plus minus viridescentes, vel sæpe virides. Carpella ovalia, lævia, nitida, coccinea.*

6. **U. Zeylanica**, Lin. (H.f. et T., l. c. p. 102, cum syn.)—C.P. 1030, 1031.

HAB. Southern parts of the island, at no great elevation.

### 3. GONIOTHALAMUS, Blume.

Sect. 1.—*Sepalis persistentibus; stylis subulatis, indivisis.*

1. **G. Hookeri**, Thw. (C.P. 3399); *foliis magnis, ovato-oblongis, breviter et obtuse acuminatis, basi parce angustatis, lævibus; pedunculis petiolos æquantibus, solitariis axillaribus et fasciculatis, caulinis.*

HAB. Hinidoon and Reigam Corles, at an elevation of about 1000 feet, April, 1855.

*Arbor* mediocris, circiter 30-pedalis. *Folia* 6–13 poll. longa,  $2\frac{1}{2}$ – $4\frac{1}{4}$  poll. lata; *petiolo*  $\frac{1}{2}$ – $\frac{3}{4}$  poll. longo. *Flores* pallide virides, parcissime rufo-hirsuti. *Sepala* rotundato-acuta,  $\frac{3}{4}$  poll. longa. *Petala exteriora* circiter 2 poll. longa,  $\frac{3}{4}$  poll. lata, basi rotundata, supra medium parce contracta; *interiora* in mitram pyramidalem, intus tomentosam, circiter  $\frac{3}{4}$  poll. altam coalita. *Carpella* numerosa, fabæ minoris magnitudine, brevissime pedicellata. *Semina* nitida; *testa* brunnea.

2. **G. Thwaitesii**, H.f. et T., l. c. p. 106.—C.P. 1040.

HAB. Not uncommon in the Central Province, at an elevation of 2000 to 4000 feet. Nom. vulg. "Caloocaara."

*Arbor* mediocris. *Flores* virides.

3. **G. Gardneri**, H.f. et T., l. c. p. 107.—C.P. 1029.

HAB. In similar localities with the preceding, not uncommon.

*Frutex* erectus, 8–10-pedalis. *Flores* virides, flavescentes. *Carpella* brevi-pedicellata, sæpissime 2-sperma, subpollicaria, oblonga.

Sect. 2.—*Sepalis* persistentibus; *stylo* in segmentis 2 longis filiformibus diviso.

4. **G. Thomsoni**, Thw. (C.P. 3398); *foliis* obovatis, longe et abrupte acuminatis, supra medium ad *petiolum* brevem graduatim angustatis; *pedunculis* caulinis, solitariis vel binis.

HAB. Forests in the district between Galle and Ratnapoora, April, 1855.

*Frutex* erectus,  $1\frac{1}{2}$ –3-pedalis. *Folia* flavescentia, 5–12 poll. longa,  $1\frac{1}{2}$ –3 poll. lata; *acumine*  $\frac{3}{4}$ – $1\frac{1}{2}$  poll.; *petiolo* circiter  $\frac{1}{2}$  poll. longo. *Pedunculi*  $\frac{3}{4}$  poll. longi, adscendentes, prope apicem decurvati. *Flores* ochracei. *Sepala* lanceolata,  $\frac{1}{4}$  poll. longa. *Petala exteriora* lanceolata, apicem versus valde angustata, basi rotundata,  $1\frac{1}{2}$  poll. longa; *interiora* in mitram ovalem acutam 6 lin. altam coalita. *Ovaria* elongata, *stylis* æquilonga. *Carpella* monosperma, ovalia, apiculata,  $\frac{1}{2}$  poll. longa; *pedicello* 2 lin. longo.

A very distinct species, remarkable for the small size of the plant, its cauline flowers often close to the ground, and its deeply bisected styles.

Sect. 3.—*Sepalis* deciduis. *Stylo* cylindrico vel clavato; *stigmatibus* dilatato, bilobo.

5. **G. salicinus**, H.f. et T., l. c. p. 106.—C.P. 373.

HAB. Forests in the neighbourhood of "Adam's Peak," at an elevation of 2000 to 3000 feet.

*Frutex* erectus, 10–12-pedalis. *Flores* ochraceo-rufescentes vel fusciscentes.

6. **G. reticulatus**, Thw. (C.P. 3397); *foliis* lanceolatis vel lineari-lanceolatis, plus minus abrupte acuminatis, basi subrotundatis, subtus conspicue reticulatis.

HAB. Carawitta Kanda and other forests between Galle and Ratnapoora.

*Frutex* erectus, 6–8-pedalis. *Ramuli* nigricantes, juniores strigosi. *Folia* membranacea, densissime glanduloso-punctata, subtus sparse rufro-nigro-pilosa, 6–11 poll. longa,  $1$ – $2\frac{1}{4}$  poll. lata. *Petioli* nigricantes, subtumidi, 2–3 lin. longi, superne sulcati, alati (lamina decurrente), subtus cum costa minute tuberculati. *Pedunculi* uniflori, paullo supra-axillares, decurvati,  $\frac{1}{4}$  poll. longi, basi bracteati. *Styli* clavati, *ovariis* oblongis dense rufo-strigosis dimidio longiores. *Carpella* globose 1-sperma, oblonge 2-sperma, rubiginosa; *pedicello*  $\frac{1}{2}$  lin. longo.

Closely allied to *G. salicinus*, but differing in its very much larger and differently-shaped leaves. I have not seen the sepals and petals of *G. reticulatus*, as they had fallen when the specimens were gathered.

7. *G. Walkeri*, H.f. et T., l. c. p. 109.

I have never been so fortunate as to meet with this species, so can add nothing to the description given of it by the authors of the 'Flora Indica.' Nor can I tell, without an examination of specimens, to which section of this genus it is to be referred.\*

4. *OROPHEA*, Blume.1. *O. Zeylanica*, H.f. et T., l. c. p. 111.—c.p. 2406.

HAB. Central Province, at an elevation of 2000 to 3000 feet.

*Flores virides, fuscescentes.*

2. *O. coriacea*, Thw. (c.p. 2612); *foliis* valde coriaceis, ovato-lanceolatis, subacutis, basi rotundatis, brevi-petiolatis; *pedunculis* unifloris; *staminibus* 9 (6 exterioribus 3 interioribus), *ovarium* solitarium oblongum 2-ovulatum cingentibus.

HAB. Dimboola and Raxawa, in the Central Province, at an elevation of about 3000 feet, April, 1852.

*Arbor* mediocris. *Ramuli* rigidi, nigricantes. *Folia* intense viridia, 2-5 poll. longa, 1-2 $\frac{1}{4}$  poll. lata, costa prominenti, nervis inconspicuis. *Petioles* vix 2 lin. longi. *Pedunculi* circiter  $\frac{1}{4}$  poll. longi, basin versus 3-4 bracteis parvis strigosis muniti. *Flores* parvi, 4 lin. in exp. *Sepala* minuta, deltoidea, ciliata. *Petala exteriora* rotundata, obscure acuminata, externe purpurea, interne ochraceo-rufescentia; *interiora* purpurea, in mitram obtusam 1 $\frac{1}{2}$  lin. altam coalita. *Ovula* superposita. *Carpella* subsessilia, subglobosa, obliqua, 1-2-sperma.

3. *O. ? obliqua*, H.f. et T., l. c. p. 112.—c.p. 2703; *floribus* terminalibus caulisque fasciculatis; *staminibus* 6, seriebus duabus, *ovaria* 3, 2-ovulata, *stigmata* depresso coronata, cingentibus; *toro* plano; *carpellis* sessilibus, globosis, 1-2-spermis.

HAB. In the Galle and Ratnapoora Districts, at no great elevation.

*Arbor* mediocris. *Flores* purpurei, parvi. *Sepala* minuta, ciliata. *Petala* 6, subæqualia, carnosia, *exteriora* aperta, *interiora* convergentia, imbricata v. subvalvata. *Carpella* semipollicaria, rufescentia.

This somewhat departs from the character of *Orophea* in the sepals and petals being frequently (not always) imbricated, as in *Sageræa*, but it is too closely allied to the previous species to be separated generically from it. It would seem to point out, however, that *Sageræa* and *Orophea* should be arranged in juxtaposition.

5. *MITREPHORA*, Blume.

1. *M. Heyneana*, (*Orophea Heyneana*, H.f. et T., l. c. p. 110, cum syn.)—c.p. 1033.

HAB. Haragam and other places on the Lower Badulla Road from Kandy, up to an elevation of 1500 feet.

*Arbor* mediocris. *Pedunculi* 1-3-flori. *Petala* lutea, *interiora* rubro-maculata.

The authors of the 'Flora Indica' must have had imperfect specimens of the flower of this plant for examination, for in all that I have dissected I have found the stamens far more numerous than they have described. Is the Courtalam plant distinct from the Ceylon one? †

\* The sepals appear to be persistent, but the specimens being in fruit only, I cannot determine whether it should be referred to section 1 or 2.—*J. D. H.*

† We have no flowering specimens of the Ceylon plant.—*J. D. H.*

## 6. XYLOPIA, Lin.

1. **X. parvifolia**, H.f. et T., l. c. p. 125.—c.P. 3335.

HAB. Southern parts of the island, at no great elevation. Nom. vulg. "Nattou."

*Arbor*. Flores ochracei. *Carpella* columbæ ovi magnitudine vel minora, oblonga. *Semina* biserialia, oblonga, in pulpa carnosâ rufa nidulantia.

The flowers and bark of this tree are used by the Cinghalese for chewing with their betel.

2. **X. nigricans**, H.f. et T., l. c. p. 125.—c.P. 615.

HAB. This grows in the Botanic Garden, and I have not met with it elsewhere, though it probably occurs in the southern parts of the island where the other two species are found.

*Arbor* mediocris. Flores ochracei. *Carpella* ut in præcedente. *Semina* fusconigra, lævia, hilum versus attenuata, 4 lin. longa, biserialia.

3. **X. Championii**, H.f. et T., l. c. p. 126.—c.P. 1038.

HAB. Ambagamowa and Ratnapoora Districts, up to an elevation of 2000 feet. Nom. vulg. "Datkateya."

Flores ochracei, plus minus rubro tincti. *Carpella* pedicellata, longitudinaliter 5-costata. *Semina* uniserialia, in pulpa nidulantia.

## 7. CYATHOCALYX, Champion.

1. **C. Zeylanicus**, Champ. (H.f. et T., l. c. p. 127.)—c.P. 1044.

HAB. Central and Southern Provinces, at an elevation of 1500 to 3000 feet. Nom. vulg. "Kakala."

*Ovarium* solitarium, uni calycis dentium oppositum. *Carpella* subglobosa, plurisperma, 2½ poll. longa, drupacea. *Semina* compressa, 1 poll. longa, lateribus transverse rugosa; *testa* ossea, rufo-fusca.

## 8. ARTABOTRYS, R. Br.

1. **A. odoratissimus**, R. Br. (H.f. et T., l. c. p. 128, cum syn.)—c.P. 1042.

HAB. Northern parts of the island, *Gardner*.

2. **A. Zeylanicus**, H.f. et T., l. c. p. 128.—c.P. 1043.

HAB. Central and southern parts of the island, at an elevation of 3000 to 4000 feet.

## 9. UNONA, Lin.

1. **U. Zeylanica**, H.f. et T., p. 132.—c.P. 1037.

HAB. Central Province, at an elevation of 1500 to 3000 feet.

*Frutex* erectus, 3-6-pedalis. *Pedunculi* sæpe caulini. Flores viridi-ochracei, intus basi rufescentes. *Carpella* rufescentia. *Semina* ovalia, lævia, 4 lin. longa.

2. **U. discolor**, Vahl, (H.f. et T., l. c. p. 132, cum syn.)

I have never met with this species wild in the island, nor are there any specimens of it in this Herbarium but from cultivated plants.

## 10. POLYALTHIA, Blume.

1. **P. Moonii**, Thw. (c.P. 437); *foliis* lanceolatis, basin versus rotun-



datam obliquam angustatis, brevi-petiolatis; *pedunculis* longis, unifloris, caulinis et oppositifoliis; *petalis* carnosis, conchoideis; *carpellis* pedicellatis, 1-2-spermis; *seminibus* oblongis, lævibus, sulco circumcinctis.

HAB. Caltura, *Moon*. Reigam Corle, September, 1856.

*Frutex* 8-10-pedalis, *partibus novellis* fulvo-strigosis. *Ramuli* rigidi, cortice fusco, rugoso, albo-maculato. *Folia* 3-6 poll. longa,  $\frac{3}{4}$ -2 poll. lata. *Petioli* circiter 1 lin. longi, transverse rugulosi. *Pedunculi* 1-1 $\frac{1}{2}$  poll. longi, basi bracteis 3 parvis lanceolatis muniti, fulvo-strigosi. *Flores* rubri, 4 lin. lati, externe sparse pilosi. *Sepala* lata, acuminata, 1 lin. longa. *Petala exteriora* rotunda, acuminata, *interioribus* obtusis minora. *Stamina* numerosa, cuneata, arcte compacta. *Ovaria* numerosa, oblonga, pilosa, *stylo* hirsuto clavato dimidio longiora. *Carpella* rufescentia, globose 1-sperma vel oblonge 2-sperma, 2-4 lin. longa, *pedicello* circiter 2 lin. longo suffulta.

This is probably the species referred to at the bottom of page 137 of the 'Flora Indica.'

### 11. GUATTERIA, Ruiz et Pavon.

1. **G. longifolia**, Wall. (H.f. et T., l. c. p. 139, cum syn.)—c.P. 1041.

HAB. Northern parts of the island, at no great elevation.

2. **G. suberosa**, Dun. (H.f. et T., l. c. p. 140, cum syn.)—c.P. 1035.

HAB. Near Caltura and Colombo.

3. **G. Corinti**, Dun. (H.f. et T., l. c. p. 141, cum syn.)—c.P. 1034.

HAB. Very common in the warmer parts of the island, ascending to an elevation of 1500 feet.

4. **G. coffeoides**, H.f. et T., l. c. p. 141.—c.P. 2503.

HAB. Haragam and other places on the Lower Badulla Road from Kandy, up to an elevation of 1500 feet.

*Flores* flavo-virides. *Carpella* nigro-purpurea. *Semina* oblonga, 7 lin. longa, 5 lin. lata, lævia, fusca, sulco undulato circumcincta.

5. **G. persicæfolia**, H.f. et T., l. c. p. 140.

This species, found by Captain Champion near Galle, I have never been so fortunate as to meet with. It would appear, from the description in 'Flora Indica,' to bear considerable resemblance to *Polyalthia Moonii*.

### 12. ANAXAGOREA, St. Hilaire.

1. **A. Zeylanica**, H.f. et T., l. c. p. 144.

I have not yet met with this species, nor are there any specimens of it in this Herbarium; so I can add nothing to the description given of it in 'Flora Indica.'

### 13. MILIUSA, Lesch.

1. **M. montana**, Gardn. MSS. (H.f. et T., l. c. p. 148.)

Vars.  $\alpha$  and  $\beta$  (c.P. 194); *ramulis foliisque junioribus* parce strigosis.

Var.  $\gamma$ . *tomentosa* (c.P. 1036); *ramulis foliisque subtus* plus minus tomentosus.

HAB. Central Province, at an elevation of 1500 to 3000 feet.

The extreme forms of the above varieties look very distinct, but they pass into each other by such insensible gradations, that it is quite impossible to separate them by well-defined characters. The small *exterior petals* vary from being subequal to nearly three times the length of the sepals; the carpels, too, are either sessile or on pedicels nearly two lines in length. The leaves are occasionally subrotund and cordate at the base.

2. **M. Zeylanica**, Gardner, MSS. (H.f. et T., l. c. p. 149.)—c.p. 2678.  
HAB. South of the island, at no great elevation.

*Carpella* subsessilia; glabra, minute granulata.

Closely allied to the preceding, but distinguished by its larger and more acute leaves, shorter peduncles, and narrower petals.

#### 14. ALPHONSEA, H.f. et T.

1. **A. Zeylanica**, H.f. et T., l. c. p. 153.—c.p. 1039.

HAB. Central Province, at an elevation of 3000 to 4000 feet.

*Flores* viridi-ochracei. *Carpella* breviter fulvo-tomentosa, dura, lignosa.

2. **A. sclerocarpa**, Thw. (c.p. 2727); *foliis* lanceolatis, obtusis, basi angustatis, coriaceis, reticulatis, glaberrimis, superne nitidis, subtus opacis, pallidioribus; *fasciculis* multifloris; *carpellis* subosseis, grosse verrucosis, brevi-pedicellatis.

HAB. Haragam, etc., on the Lower Badulla Road from Kandy, up to an elevation of 1500 feet.

*Arbor*. *Ramuli* cinerei, rugosi; *gemmae* fulvo-pilosæ. *Folia* 2-3½ poll. longa, ¾-1 poll. lata, *petiolo* 2-lin. longo. *Pedunculi* vix ½ lin. longi, extra-alares. *Pedicelli* ¼ poll. longi, medio 1-bracteati, fusco-pilosi. *Sepala* minuta. *Petala* ¼ poll. longa, pallide ochracea, externe pilosa. *Stamina* 15, 2-3-serialia. *Ovaria* 3-5, *stylo* depresso. *Ovula* circiter 8. *Carpella* 3-5, stellatim disposita, 1-2 poll. longa, subglobosa, breviter fulvo-tomentosa, *pedicello* robusto, ¼ poll. longo. *Semina* biserialia.

### VI. MYRISTICACEÆ.

#### 1. MYRISTICA, Lin.

1. **M. Irya**, Gært. (H.f. et T., Fl. Ind. p. 159, cum syn.; Alph. DC. Prod. xiv., p. 202.)—c.p. 2620.

HAB. Not uncommon in damp places, up to an elevation of 2000 feet. Nom. vulg. "Eereya-gass."

2. **M. Zeylanica**, Alph. DC. Ann. Sc. Nat. ser. 4, vol. iv., p. 29; Prod. xiv., p. 190.—c.p. 2923.

HAB. Banks of rivers, at no great elevation. Nom. vulg. "Malabodde."

3. **M. laurifolia**, H.f. et T., l. c. p. 163; Alph. DC. Prod. xiv., p. 191.—*M. diospyrifolia*, Alph. DC., l. c.—c.p. 416.

HAB. Central Province, in the forests, at an elevation of 1500 to 4000 feet. Nom. vulg. "Malabodde."

4. **M. Horsfieldii**, Bl. (H.f. et T., l. c. p. 163, cum syn.; Alph. DC. Prod. xiv., p. 200.)—c.p. 221.

HAB. Abundant in the Ambagamowa and Ratnapoora districts, up to an elevation of 1000 feet. Nom. vulg. "Rook-gass."

The very fragrant male flowers are used by the natives as a perfume for scenting their clothes.

### VII. MONIMIACEÆ.

#### 1. HORTONIA, Wight.

1. **H. floribunda**, Wight, (H.f. et T., l. c. p. 166.)

Var. *a. acuminata*, H.f. et T., l. c. (*Hort. floribunda* and *H. acuminata*, Wight, Ic.)—c.p. 1027.

Var. *β. ovalifolia* (*Hort. ovalifolia*, Wight, Ic.)—c.p. 159, 161, 175.

Var. *γ. angustifolia*; foliis lineari-lanceolatis; floribus multo minoribus quam in varr. *a* et *β*.—c.p. 1026.

HAB. Vars. *a* and *β*. Central Province, at an elevation of 4000–7000 feet. Var. *γ*. In the neighbourhood of Galle, at no great elevation. Nom. vulg. "Waaweya."

*Flores* pallide viridi-flavi. *Drupæ* carnosæ, pulposæ, rubro-nigricantes.

Between varieties *a* and *β* of this species there are intermediate forms, by which they pass insensibly into one another, and they both occur at considerable elevations on the hills of the Central Province. Variety *γ*, however, may be considered a permanent or sub-permanent one, as it is a much more delicate plant, with far smaller flowers, and occurs at no great elevation above the level of the sea. I shall have occasion to notice, in this "Enumeration," similar varieties of several other species.

## VIII. MENISPERMACEÆ.

### 1. COSCINIUM, Colebr.

1. **C. fenestratum**, Colebr. (H.f. et T., l. c. p. 178, cum syn.)—c.p. 1055.

HAB. Central and Southern Provinces, up to an elevation of 1500 feet. Nom. vulg. "Weniwel."

An infusion of the wood of this plant is employed by the Cinghalese as a bitter tonic.

### 2. TINOSPORA, Miers.

1. **T. tomentosa**, Miers, (H.f. et T., l. c. p. 183, cum syn.)—c.p. 2804.

HAB. On the Lower Badulla Road from Kandy, at no great elevation. Nom. vulg. "Bookinda."

2. **T. cordifolia**, Miers, (H.f. et T., l. c. p. 184, cum syn.)—c.p. 1053.

HAB. Warm, drier parts of the island, at no great elevation. Nom. vulg. "Rassakinda."

An infusion of the young stems of this plant is used as a remedy in fevers.

### 3. ANAMIRTA, Colebr.

1. **A. Cocculus**, W. et A. (H.f. et T., l. c. p. 185, cum syn.)—c.p. 2722.

HAB. Warmer parts of the island, up to an elevation of 1500 feet. Nom. vulg. "Tetti-wel."

### 4. TILIACORA, Colebr.

1. **T. acuminata**, Miers, (H.f. et T., l. c. p. 187, cum syn.)—c.p. 1056.

HAB. North-eastern parts of the island, at no great elevation. Gardner.

### 5. LIMACIA, Lour.

1. **L. cuspidata**, H.f. et T., l. c. p. 189.—c.p. 1051.

HAB. Common up to an elevation of 4000 feet. Nom. vulg. "Nirriwel."

### 6. COCCULUS, DC.

1. *C. macrocarpus*, W. et A. (H.f. et T., l. c. p. 191, cum syn.)—c.p. 1052.

HAB. Central Province, not uncommon.

### 7. STEPHANIA, Lour.

1. *S. hernandifolia*, Walpers, (H.f. et T., l. c. p. 196, cum syn.)—c.p. 2757.

HAB. Common in the Central Province.

### 8. CISSAMPELOS, Lin.

1. *C. Pareira*, Lin. (H.f. et T., l. c. p. 198, cum syn.)—c.p. 169.

HAB. Very common throughout the island. Nom. vulg. "Deyamitta."

2. *C. subpeltata*, Thw. (c.p. 168); *foliis* cordatis vel rotundato-acutis obtusatisve, subpeltatis, apiculatis, superne parce pilosis, subtus cinereis, subtomentosis; *FÆM. bracteis* parvis, oblongis oblatisve, utrinque hirsutissimis, apiculo subæquilongo; *drupis* oblongis.

HAB. Ambagamowa and Ratnapoora Districts, at no great elevation.

*Ramuli* striati, parce pubescentes. *Folia* 2–3 poll. longa et æquilata. *Petioli* 1½–2½ poll. longi, striati, pubescentes, prope marginem folii affixi. *Paniculæ* plus minus elongatæ, subtomentosæ. *Drupæ* 2 lin. longæ, 1½ lin. latæ.

This seems to be a very distinct species, the female inflorescence wanting the large foliaceous bracts so characteristic of *C. Pareira*. The drupes of *C. subpeltata* are also much larger than those of its congener.

### 9. CYCLEA, Arn.

1. *C. Burmanni*, Miers, (H.f. et T., l. c. p. 201, cum syn.)—c.p. 1049.

HAB. Very abundant throughout the island. Nom. vulg. "Caahe-pittang."

2. *C. peltata*, H.f. et T., p. 201, cum syn.

I know nothing of this plant, never having collected any specimens of the genus but what seemed to be all referable to one species.

### 10. PACHYGONE, Miers.

1. *P. ovata*, Miers, (H.f. et T., l. c. p. 203, cum syn.)—c.p. 1050, 1054.

HAB. Dry, hotter parts of the island, not uncommon.

## IX. BERBERIDEÆ.

### 1. BERBERIS, Lin.

1. *B. aristata*, DC. (H.f. et T., l. c. p. 222, cum syn.)—c.p. 2405.

HAB. Newera Ellia, and other elevated parts of the island, abundant.

## X. NYMPHÆACEÆ.

## 1. NYMPHÆA, Lin.

1. **N. Lotus**, Lin. (H.f. et T., l. c. p. 241, cum syn.)

Var. *a. Lotus* (c.p. 1020).

Var. *β. pubescens* (c.p. 1019).

HAB. Not uncommon in the lower parts of the island.

2. **N. stellata**, Willd. (H.f. et T., l. c. p. 243, cum syn.)—c.p. 1021.

HAB. Common in the warmer parts of the island.

I have never seen any but the pale, whitish variety in Ceylon.

## XI. NELUMBIACEÆ.

## 1. NELUMBIUM, Juss.

1. **N. speciosum**, Willd. (H.f. et T., l. c. p. 248, cum syn.)—c.p. 1022.

HAB. Colombo, and elsewhere in the island.

## XII. CRUCIFERÆ.

## 1. CARDAMINE, Lin.

1. **C. hirsuta**, Lin.—c.p. 2962.

Var. *major*; sæpissime trifoliolata.—c.p. 1017.

HAB. Not uncommon in the more elevated parts of the island, above 4000 feet. The trifoliolate variety occurs in shady woods.

## XIII. CAPPARIDACEÆ.

## 1. GYNANDROPSIS, DC.

1. **G. pentaphylla**, DC. Prod. i. 238, cum syn.—c.p. 2640.

HAB. Common in cultivated ground. Nom. vulg. "Waila."

## 2. CLEOME, DC.

1. **C. monophylla**, Lin. (DC. Prod. i. 239.)—c.p. 2792.

HAB. Cultivated ground, common. The flowers are of a dull-purple colour.

2. **C. aspera**, Kœnig, (DC. Prod. i. 241.)—c.p. 1068.

HAB. Dry sandy ground, common.

## 3. POLANISIA, Rafin.

1. **P. viscosa**, *β. icosandra*, DC. Prod. i. 242.—c.p. 1073.

HAB. Very common in cultivated ground.

## 4. CRATÆVA, Lin.

1. **C. Roxburghii**, R. Br. (W. et A., Prod. i. 23.)—c.p. 1067.

HAB. North-eastern parts of the island, at no great elevation. Nom. vulg. "Loonoowarana."

### 5. CADABA, Försk.

1. **C. trifoliata**, W. et A. Prod. i. 24.—C.P. 1066.

HAB. Northern part of the island, *Gardner*.

2. **C. Indica**, Lam. (W. et A. Prod. i. 24.)—C.P. 1070.

HAB. Northern part of the island, *Gardner*.

### 6. MÆRUA, Försk.

1. **M. oblongifolia**, A. Rich. Tent. Fl. Abyss. i. 32, t. 5.—*Niebuhrria oblongifolia*, DC. Prod. i. 244; Hook. Ic. ii. t. 128.—C.P. 1064.

HAB. Trincomalee, *Gardner*.

### 7. CAPPARIS, Lin.

#### Sect. 1.

1. **C. brevispina**, DC. Prod. i. 246, (W. et A. Prod. i. 24, cum syn.)

Var. *a. brevispina* (*C. brevispina*, DC.)—C.P. 1062 and 2509.

Var. *β. Rheedii* (*C. Rheedii*, DC.)—C.P. 1060.

Var. *γ. rotundifolia* (*C. rotundifolia*, Rottb.; Willd.)—C.P. 1059.

HAB. Very abundant in the hot, dry northern parts of the island.

May not one of the forms of this species be the *C. Zeylanica* of Linnæus?

2. **C. stylosa**, DC. Prod. i. 246, (W. et A. Prod. i. 25, cum syn.)—C.P. 1072.

HAB. North of the island, *Gardner*.

#### Sect. 2.

3. **C. horrida**, Lin. (W. et A. Prod. i. 26, cum syn.)—C.P. 1058.

HAB. Hot, drier parts of the island, up to an elevation of 1500 feet.

4. **C. tetrasperma**, Thw. (C.P. 614); *stipulis* spinosis, brevibus, uncinatis; *foliis* ovatis, acuminatis, glabris; *junioribus* parce pubescentibus; *sepalis* glabris, margine lanatis; *petalis* interne glabris, externe lanuginosis.

HAB. Lower Badulla Road from Kandy, up to an elevation of 1500 feet.

*Frutex* scandens. *Ramuli* teretes, minute striati, glabri. *Folia*  $1\frac{3}{4}$ – $2\frac{1}{2}$  poll. longa,  $\frac{3}{4}$ – $1\frac{1}{4}$  poll. lata, *petiolo*  $\frac{1}{4}$  poll. longo. *Pedicelli* supra-axillares, sæpissime 3, 1-seriales, 6 lin. longi. *Petala* oblonga, 2 lin. longa, *sepala* ovalia acuta parum excedentia. *Stamina* circiter 20. *Ovarium* oblongum, attenuatum; *gynophoro* pedicellum subæquante; *placentis* 2, 2–3-ovulatis. *Ovula* brevi-stipitata. *Nuculania* globosa, 4 lin. in diam., sæpissime 4-sperma.

This is probably the *C. Zeylanica* of De Candolle, but can scarcely be the plant so called by Linnæus, the leaves of which are described as "subpetiolata;" nor can it well be the *C. acuminata*, Willd. (*C. Zeylanica*, W. et A.), which is stated to have the leaves tomentose beneath, and the spiny stipules straight.

#### Sect. 3.

5. **C. Roxburghii**, DC. Prod. i. 247; W. et A. Prod. i. 26; R. W. Ic. t. 1048.—C.P. 1065, 2480.

HAB. Hot, drier parts of the island, at no great elevation.

*Petala* albida. *Ovarium* oblongum, attenuatum; *placentis* 5, pluriovulatis. *Ovula* subsessilia. *Nuculania* globosa. *Semina* numerosa.

6. **C. Moonii**, R. W. Illust. i. p. 35.—c.p. 2415.

HAB. Central Province, up to an elevation of 3000 feet.

*Frutex* scandens, maximus, glaberrimus; *stipulis* spinosis, uncinatis. *Folia* 4–6 poll. longa,  $1\frac{3}{4}$ – $2\frac{1}{4}$  lata, oblonga, acuminata, acumine reflexo, basi rotundata, *petiolo* circiter  $\frac{1}{2}$  poll. longo. *Petala* alba, parce pilosa, *sepalis* rotundatis circiter 1 poll. longis bis longiora. *Stamina* numerosa, filamentis albis, 3–4 poll. longis. *Ovarium* ampullæforme, *gynophoro* 4 poll. longo, *placentis* 4, pluriovulatis. *Ovula* longe-stipitata. *Nuculania* globosa, circiter 4 poll. in diam. *Semina* numerosa, fabæ magnæ magnitudine.

7. **C. retusella**, Thw. (c.p. 2550); *ramis ramulisque* ferrugineo-tomentosis, *stipulis* spinosis uncinatis armatis; *foliis* parvis, ovatis, versus apicem retusum angustatis; *racemis* terminalibus, 5–6-floris; *pedicellis* horizontalibus; *floribus* parvis.

HAB. Lower Badulla Road from Kandy, up to an elevation of 1500 feet.

*Folia* 1–2 poll. longa, 3–6 lin. lata, glabra, *juniora* ferrugineo-pilosa. *Petioli*  $1\frac{1}{2}$  lin. longi, tomentosi. *Pedicelli* glabri, 6 lin. longi, basi *bractea* minuta lanceolata tomentosa suffulti. *Sepala* rotundata, albo-marginata, glabra, parce ciliata, 2 lin. longa. *Petala* alba, glabra, ciliata, oblonga, 3 lin. longa. *Stamina* circiter 30, *gynophorum* 4 lin. longum æquantia. *Ovarium* oblongum, apice attenuatum; *placentis* 2, pluriovulatis. *Ovula* brevi-stipitata. *Nuculania* globosa, 3 lin. in diam.

8. **C. pedunculosa**, Wall. No. 6999, (W. et A. Prod. i. 27.)—c.p. 1069.

HAB. Lower Badulla Road from Kandy, up to an elevation of 1500 feet.

9. **C. grandis**, Hb. Heyne, (W. et A. Prod. i. 27.)—c.p. 1071.

HAB. North-east of the island, *Gardner*.

10. **C. sepiaria**, Lin. (W. et A. Prod. i. 26, cum syn.)—c.p. 1063.

HAB. Jaffna, *Gardner*.

#### XIV. FLACOURTIACEÆ.

##### BIXEÆ.

##### 1. BIXA, Lin.

1. **B. Orellana**, Lin. (W. et A. Prod. i. 31, cum syn.)—c.p. 2821.

HAB. Kornegalle, Four Korles, etc., but scarcely truly wild.

##### PROCKEÆ.

##### 2. PHOBEROS, Lour.

Sect. 1.—*Foliis ad petioli apicem biglandulosis*.

1. **P. Arnottianus**, Thw. (c.p. 3526); *foliis* ovalibus, apicem versus obscure serratis; *sepalis* 8–10, oblongis.

HAB. I know of this species only from two not very old trees growing in the Botanic Garden, where there is no record of whence they were procured. Nom. vulg. “Katookooroondoo.”

*Arbor* mediocris. *Truncus* spinis magnis compositis armatus. *Ramuli* plerumque

inermes. *Folia*  $1\frac{1}{2}$ – $2\frac{1}{4}$  poll. longa,  $1$ – $1\frac{1}{2}$  poll. lata, apice obscure et brevissime acuminato, rotundato vel retuso. *Petioli*  $\frac{1}{4}$  poll. longi. *Nuculania* pisi magnitudine, subglobosi, pulposi, rubro-nigrescentes.

There can be little doubt, I think, of this being Gærtner's plant, from the number of sepals being frequently 8, as he has figured them. It can scarcely be the *P. macrophyllus*, W. et A. Prod. i. 30, as the leaves are scarcely acuminated, and the fruit is pulpy, not dry.

Sect. 2.—*Foliis basi eglandulosis.*

2. **P. Gærtneri**, Thw.; *trunco* armato; *foliis* utrinque venis prominentibus reticulatis; *sepalis* 10, rotundatis, obtusis.—*Limonia pusilla*, Gærtner. i. 279. t. 58, 4.

Var. *a. oblongifolius* (C.P. 2497); *foliis* oblongis, basi apiceque plerumque rotundatis.

Var. *β. cordifolius* (*Scolopia pusilla*, Moon's Cat.)—C.P. 1076; *foliis* ovato-vel rotundato-cordatis, obtusis vel subacuminatis.

Var. *γ. lanceolatus* (C.P. 64, 181, 211); *foliis* lanceolatis, basi apiceque plus minus angustatis.

HAB. Warmer parts of the island, up to an elevation of 1500 feet.

Var. *β.* Hot parts of the island, and especially near the seacoast.

Var. *γ.* Hot, drier parts of the island. Nom. vulg. "Kattoo-koo-roondoo."

*Arbor* mediocris. *Folia* (varr. *a.* et *γ.*)  $1$ – $2$  poll. longa,  $\frac{1}{2}$ – $\frac{3}{4}$  poll. lata; (var. *β.*) valde variantia. *Nuculania* subglobosa, pisi magnitudine, pulposa, rubro-nigrescentia.

It is impossible to retain Gærtner's specific name of *pusillus* for this, which forms a large tree. The leaves of var. *γ* are less coriaceous and more minutely reticulated than those of varr. *a* and *β*, and it may possibly be a distinct species.

3. **P. acuminatus**, Gardner, MSS. (C.P. 1077); *foliis* glabris, ovato-lanceolatis, longe acuminatis, apicem versus obtusum serratis, basi angustatis; *sepalis* 10–12, late lanceolatis.

HAB. Central Province, at an elevation of 1000 to 3000 feet. Nom. vulg. "Katookenda."

*Arbor* mediocris, junior spinosa, denique inermis. *Folia*  $2$ – $3\frac{1}{2}$  poll. longa,  $1$ – $1\frac{3}{4}$  poll. lata, subtus pallidiora, obscure et delicatule venosa. *Petioli*  $2$ – $3$  lin. longi. *Nuculania* subglobosa, cerasi magnitudine, carnosae, viridia, firma nec pulposa.

4. **P. Hookerianus**, Wight, MSS. (C.P. 629); *foliis* ovatis, subacuminatis, obtusis, basi angustatis, margine parce undulatis; *sepalis* 12–14, oblongis.

HAB. Central Province, at an elevation of 4000 to 7000 feet.

*Arbor* ingens, junior spinosa, denique inermis. *Folia*  $1\frac{1}{4}$ – $2$  poll. longa,  $\frac{3}{4}$ – $1$  poll. lata, valde coriacea, subtus venis prominentibus reticulata. *Petioli* vix  $2$  lin. longi. *Nuculania* cerasi magnitudine vel majora, carnosae, viridia, firma nec pulposa.

FLACOURTIEÆ.

3. FLACOURTIA, Commers.

1. **F. sepiaria**, Roxb. (W. et A. Prod. i. 29, cum syn.)—C.P. 1650.

HAB. Hot, drier parts of the island, up to an elevation of 1500 feet.

2. **F. sapida**, Roxb. (W. et A., l. c.)—C.P. 2583.

HAB. Central Province, at an elevation of 1500 to 3000 feet. Nom. vulg. "Oogoorassa."



*Arbor* mediocris. *Truncus* spinis magnis compositis armatus. *Ramuli* cortice cinerascenti; *spinis* paucis, simplicibus, axillaribus. *Folia* 2-3 poll. longa, 1½-2 poll. lata, superne nitida, reticulatim venosa, subtus pallidiora. *Nuculania* cerasi parvi magnitudine, rubro-nigrescentia.

The fruit of this tree is eaten, when quite ripe, by the natives; but when unripe it is extremely astringent.

#### 4. ROUMEA, Poit.

1. **R. hebecarpa**, Gardner, in *Calcutta Journ. of Nat. Hist.* vol. vii. p. 9 (c.p. 1075); arborea, inermis, *foliis* petiolatis, ovato-oblongis vel oblongo-lanceolatis, acuminatis, basi subobtusis, integris vel dentato-serratis, supra nitidis pubescentibus, subtus pallidis, tomentosus; *nuculanis* globosis, velutinis.—*Gardner*.

HAB. Condasalle, Maturatte, etc. Nom. vulg. "Katambilla."

*Arbor* mediocris, 16-20-pedalis. *Ramuli* teretes, cinerei, tuberculati, *juniores* rufescentes, tomentosi. *Folia* circiter 4 poll. longa, 2 lata, *petiolo* circiter 4 lin. longo, *costa* venis primariisque rufescentibus. MAS. *Pedunculi* brevissimi, sæpe geminati. *Pedicelli* 10-20, umbellati, circiter 4 lin. longi, pubescentes. *Calyx* profunde 5-7-partitus, *lobis* lanceolatis, acutis, pubescentibus. *Stamina* numerosa; *filamentis* filiformibus, *calycis* segmenta æquantibus; *antheris* introrsis, obtusis. *Ovarii* rudimentum nullum vel parvum. *Discus hypogynus* nullus. FÆM. *Pedicelli* 1-3. *Calycis* lobi ovato-lanceolati, acuti, persistentes, circiter 3½ lin. longi. *Ovarium* dense pubescens, subglobosum, *disco* annulari crenulato cinctum, *placentis* 5-7, 2-ovulatis. *Styli* divergentes, 1½ lin. longi, pilosi. *Stigmata* radiatim fimbriata. *Nuculania* circiter 1 poll. in diam.—*A* *descript. cl. Gardner præcipue excerpt.*

The acid fruit of this species is eaten by the Cinghalese.

#### ERYTHROSPERMÆ.

#### 5. ERYTHROSPERMUM, Lam.

1. **E. phytolaccoides**, Gardner, l. c. p. 11 (c.p. 468); *foliis* sparsis, petiolatis, oblongo-lanceolatis, acuminatis, basi acutis, utrinque glaberrimis, integris vel obscure serratis; *racemis* paniculatis, axillaribus terminalibusque, *folia* subæquantibus; *sepalis* 5, ovatis, obtusis, concavis; *petalis* ovato-oblongis, ciliatis; *antheris* sagittatis.—*Gardner*.

HAB. Ambagamowa and Ratnapoora Districts, up to an elevation of 1500 feet.

*Arbor* mediocris, 20-30-pedalis. *Ramuli* glabri, teretes. *Folia* alterna, superne nitida, subtus pallidiora, venis prominentibus reticulata, 3½-8 poll. longa, 1½-3 poll. lata. *Petioli* 3-8 lin. longi, superne sulcati, basi apiceque tumiduli. *Stipulæ* minutæ, lanceolatæ, rufescentes. *Racemi* multiflori. *Pedunculi* glabri, obtusanguli, albidi. *Pedicelli* 2-3 lin. longi, basi 3-bracteolati. *Sepala* albida, rufo-tincta, circiter 2 lin. longa. *Petala* alba, ovato-oblonga, obtusa, ciliata, *sepalis* æquilonga. *Ovarium* oblongum, glabrum. *Capsula* globosa, 3-5 lin. in diam., fere ad basin 3-4-valva, oligosperma.—*A* *descript. cl. Gardner præcipue excerpt.*

### XV. PANGIACEÆ.

#### 1. HYDNOCARPUS, Gærtner.

1. **H. inebrians**, Vahl, (non W. et A. *Prod.* i. 30, et R. W. *Ill.* i. 38. t. 16.)—*H. venenata*, Gærtner, *Fruct.* i. 288. t. 60, 3.—c.p. 1630.

HAB. Banks of rivers, up to an elevation of 2000 feet. Nom. vulg. "Makooloo."

*Arbor magna. Folia lanceolata, acuminata, serrata vel subintegra, basi plus minus angustata, subobliqua. Sepala rotundata, petalis rotundatis, ciliatis, 1½ lin. longis parce longiora. Squamulæ rotundatæ, acuminatæ, ferrugineo-strigosæ. Nuculania circiter 1 poll. in diam., ferrugineo-tomentosa, demum nigrescentia.*

The fruit of this plant is used by the Cinghalese for intoxicating fish. The oil extracted from the seeds is employed as a remedy in cutaneous complaints.

2. **H. octandrus**, Thw. in Hooker's Journ. of Bot. vol. vii. (1855) p. 197.—C.P. 2640.

HAB. Ambagamowa District, at an elevation of about 2500 feet.

3. **H. alpinus**, Wight, Ic. t. 942.—C.P. 2918.

HAB. Saffragam and Bintenne Districts, at no great elevation.

## 2. TRICHADENIA, Thw. l. c.

1. **T. Zeylanica**, Thw. l. c. 196. t. 5.—C.P. 2505.

HAB. Central Province, at an elevation of about 3000 feet, not uncommon. Nom. vulg. "Tettigass."

The oil extracted from the seeds is used for diseases of the skin of children.

## XVI. SAMYDACEÆ.

### 1. CASEARIA, Jacq.

1. **C. tomentosa**, Roxb. (DC. Prod. ii. 51.)—C.P. 1248.

HAB. Jaffna and Sigre, Gardner.

*Ramuli pubescentes. Folia oblonga, utrinque angustata, obtusa, crenata, 3-4 poll. longa, 1-1½ lata, superne subglabra, subtus pubescentia. Petioli pubescentes, ½ poll. longi. Ovarium 3-placentatum.*

2. **C. Zeylanica**, (*C. ovata*, Willd.? *C. esculenta*, Roxb.? *Vareca Zeylanica*, Gærtn. Fruct. i. 290. t. 60, 6.)—C.P. 415, 2603; *foliis lanceolatis, acuminatis, basi acutis, integris vel obscure dentatis, nitidis, reticulatis, junioribus parce hirsutis; staminibus 8; ovario hirsuto, 3-placentato; stigmatate in medio depresso.*

HAB. Hotter parts of the island, very common, up to an elevation of 1500 feet. Nom. vulg. "Wal-wareka."

*Arbor mediocris. Folia 3-6 poll. longa, 1-2 poll. lata. Petioli ¼-½ poll. longi.*

3. **C. Championi**, Thw. (C.P. 2608); *foliis ovatis vel ovato-lanceolatis, acuminatis, glabris, nitidis, reticulatis; staminibus 8; ovario glabro, 3-placentato; stigmatate plano, obscure 3-lobato.*

HAB. Central Province, at an elevation of 3000 to 4000 feet.

*Arbor mediocris. Ramuli, cum foliis junioribus, rufescentes, glabri. Folia 3-6 poll. longa, 1¼-2¾ lata. Petioli ¼-½ poll. longi.*

4. **C. varians**, Thw.; *foliis glabris, subopacis, integris vel rarius parce crenatis, obscure reticulatis, basi acutis; staminibus sæpissime 7; ovario glabro, 2-placentato; stigmatate obscure 2-lobo.*

Var. *a. ovata* (C.P. 2604 and 2657); *foliis ovatis vel ovato-lanceolatis, acuminatis.*

Var. *β. minor* (C.P. 3365); *foliis ut in var. a, sed multo minoribus.*

Var.  $\gamma$ . *obovata* (c.p. 1217, 1247); *foliis* obovatis, apice obtuse sub-acuminato, rotundato vel retuso.

HAB. Vars.  $\alpha$  and  $\beta$  in the Central Province, at an elevation of 3000 to 5000 feet. Var.  $\gamma$  at Newera Ellia, at an elevation of 7000 to 8000 feet.

Variety  $\alpha$  of this species bears considerable resemblance to *C. Championi*, as regards the shape and size of the leaves, but in other respects is sufficiently distinct.

5. **C. coriacea**, Thw. (c.p. 465); *foliis* obovatis, brevissime acuminatis, obtusis, basi angustatis, coriaceis, creberrime reticulatis, glabris, superne (siccis) nigricantibus, subtus cinereis; *staminibus* 6-8, *squamis* truncatis pilosis multo longioribus; *ovario* glabro, 2-placentato.

HAB. Adam's Peak, at an elevation of 6000 to 7000 feet.

*Arbor* mediocris. *Folia* 2-3 poll. longa,  $1\frac{1}{4}$ - $1\frac{3}{4}$  lata. *Petioli* 2-3 lin. longi.

This species a good deal resembles the variety  $\gamma$  of *C. varians*, but the leaves are far more coriaceous, and more densely reticulated.

## 2. OSMELIA, Thw.

CHAR. GEN.—*Calyx* 5-partitus. *Stamina* 10 (5 cum totidem *squamulis* profunde bilobis, dense pilosis alternantia, 5 in earundem *squamularum* sinibus singulatim inserta), omnia fertilia, subæqualia. *Styli* 3, incurvati. *Capsula* subcoriacea, 3-valvis. *Semina* pauca.

1. **O. Gardneri**, Thw. (*Casearia paniculata*, Gardner, MSS.)—c.p. 1246.

HAB. Central Province, at an elevation of about 3000 feet. Not common.

*Arbor* mediocris, *cortice* lævi. *Ramuli* teretes, juniores pilosi. *Folia* integra, ovato-lanceolata, acuminata, glabra, pennivenia,  $2\frac{1}{2}$ -4 poll. longa,  $1-1\frac{1}{2}$  poll. lata. *Petioli* superne sulcati, 2-3 lin. longi. *Stipulæ* minutæ, deciduæ. *Racemi* terminales, elongati, interrupti, multiflori, pubescentes, pauculati. *Flores* brevissime pedicellati, albidii, plus minus rufo-tincti, singuli involucello minuto cupuliformi basi unibracteato suffulti. *Calycis* lobi rotundati, externe parce pilosi. *Stamina* exserta. *Ovarium* oblongum, dense pilosum, 3-placentatum; *placentis* 2-3-ovulatis. *Capsula* subglobosa, 1-4-sperma. *Semina* subglobosa, arillo carnosio, rubro; *testa* membranacea.

## XVII. VIOLACEÆ.

### 1. VIOLA, Lin.

1. **V. Patrinii**, DC. (W. et A. Prod. i. 32, cum syn.)—c.p. 1087.

HAB. Central Province, at an elevation of 5000 to 8000 feet. Not uncommon.

2. **V. Wightiana**, Wall. (W. et A. Prod. l. c.; R. W. Illust. 943.)—c.p. 153.

Var.  $\alpha$ . *pubescens*; *planta tota* plus minus pubescens, *foliis* crenato-serratis.

Var.  $\beta$ . *glabra*; *planta tota* glabra, *foliis* obscure crenato-serratis.

HAB. Var.  $\alpha$ , Central Province, very abundant, at an elevation of 4000 to 8000 feet. Var.  $\beta$  in wet, spongy places, at Newera Ellia.

### 2. IONIDIUM, Ventn.

1. **I. suffruticosum**, Ging. in DC. (W. et A. Prod. l. c.; R. W. Ill. 308.)—c.p. 76.

HAB. Hot, dry, exposed places, up to an elevation of 3000 feet. Common.

2. **I. ramosissimum**, Thw. (c.p. 1084); fruticosum, suberectum, foliis linearibus, apiculatis, lævibus, integerrimis, glaucis, marginibus revolutis.

HAB. Banks of the Mahawelliganga at Haragam.

Closely allied to the preceding species, but besides the characters mentioned above, the stems are much more woody, and the internodes shorter, than in *I. suffruticosum*. In the inflorescence of the two species I can discover no difference; and the glandular-tipped stipules are alike in both.

### 3. ALSODEIA, Thouars.

1. **A. Zeylanica**, Thw. (*Pentaloba Zeylanica*, Arn. Mag. of Zool. and Bot. ii. 543.)—c.p. 2669.

HAB. Ratnapoora District. Not uncommon.

*Frutex* erectus, 10-12-pedalis. *Ramuli* teretes, cinerei, opaci, juniores pilosi. *Folia* lanceolata vel ovato-lanceolata, acuminata, 2½-6 poll. longa, 1-2½ poll. lata, nitida, reticulata, dentata. *Petioli* 2½-5 lin. longi, superne sulcati. *Stipulae* lanceolatae, acutae, appressae, 3-4 lin. longae, rufo-fuscae, subpersistentes. *Fasciculi* pluribracteolati, bracteolis acutis, rufo-fuscis. *Pedicelli* circiter 1 lin. longi. *Flores* albidæ. *Calycis lobi* lanceolati, acuti, petalis apice recurvatis, plus quam dimidio breviores, rufo-tincti. *Urceolus* rufescens, 5-crenatus; crenaturis filamentis supra basin insertis oppositis. *Staminum filamenta* cum ligulis alba, *antherarum loculi* introrsi, flavi. *Stigma* trilobum.

### 4. Gen. Nov. SCYPHELLANDRA, Thw.

*Sepala* 5. *Petala* 5. *Glandulae* 5, minutæ, carnosæ, petalis alternantes, *staminum* 5 dorso appressæ. *Antheræ* sessiles, marginibus coherentes, singulæ connectivo lato in ligulam apicalem, liberam, membranaceam producto. *Ovarium* sessile, 1-loculare; *stylo* lineari; *stigmatate* 3-lobato. *Capsula* 3-valvis, 3-4-sperma.

1. **S. virgata**, Thw.; foliis alternis, ovatis vel oblongis, denticulatis, glabris, basi apiceque obtusis, petiolatis.—c.p. 1085.

HAB. Not uncommon in the hot, drier parts of the island.

*Frutex* parvus. *Ramuli* virgati, cinerei, striati; juniores pilis curvatis pubescentes. *Folia* magnitudine etiam in ramo uno valde varia, 1½-14 lin. longa, 1-6 lin. lata. *Petioli* ½-1 lin. longi, pubescentes. *Stipulae* minutæ, lanceolatae. *Peduncululi* axillares, solitarii, 1-flori, 1-2 lin. longi, basi pluri-bracteolati. *Sepala* ovata, acuminata, *petalis* lineari-oblongis, 5-nervis, circiter 1 lin. longis plus quam dimidio breviora. *Capsula* 2½ lin. longa.

This interesting little plant has the authers of *Hymenantha* and the carpels and numerous bracteolæ of *Alsodeia*, thus forming a connecting link between the two genera.

## XVIII. DROSERACEÆ.

### 1. DROSERA, Lin.

1. **D. Burmanni**, Vahl, (W. et A. Prod. i. 34, cum syn; R. W. Ic. 944.)—c.p. 1089.

HAB. Not uncommon throughout the island.

2. **D. Indica**, Linn. (W. et A. Prod. l. c. cum. syn.)—c.p. 1088.

HAB. South of the island. Not common.

3. **D. lunata**, Ham. (DC. Prod. i. 319.)—c.p. 2553.  
 HAB. Common in the most elevated districts of the island.

## XIX. POLYGALACEÆ.

### 1. POLYGALA, Lin.

1. **P. telephioides**, Willd. (W. et A. Prod. i. 36, cum syn.)—c.p. 1080.  
 HAB. Colombo, Haragam, etc.
2. **P. rosmarinifolia**, W. et A. Prod. i. 37 (non Eck. et Zeyh.).  
 Var. *a. mucronata* (c.p. 715 et 1083); *foliis* glabris vel parce pubescentibus, *bracteolis* longe mucronatis.  
 Var. *β. puberula* (c.p. 1082); *foliis* pubescentibus, *bracteolis* raro mucronulatis.  
 HAB. Not very common, but generally distributed, occurring from the level of the sea up to an elevation of 4000 feet.  
 I have hitherto been disposed to consider vars. *a* and *β* distinct species, but they resemble each other so closely in most respects that it is perhaps hardly safe to separate them on account of such trivial differences as the greater or less pubescence of the foliage, or the bracteoles being mucronate or otherwise.
3. **P. elongata**, Klein, in Willd. Sp. iii. 879; DC. Prod. i. 332.—*P. densiflora*, Blume, Bijd. 59.—(c.p. 7, 260, et 592.)  
 Var. *β. hirsutula*.—*P. hirsutula*, Arn. Pug. 4 (322.)—c.p. 1079, 1081.  
 HAB. Central Province, at an elevation of 4000 to 7000 feet.  
 This is a very variable species, especially in the form and hairiness of the leaves, and length of the racemes. The flowers, which are of a bluish or a purple colour, are very uniform in size and structure throughout the several varieties. The narrow border of the capsule is pubescent with curved hairs, and with these are frequently intermixed straight, longer hairs. Var. *β* occurs on banks at Newera Ellia, and is only a stunted form of the plant.
4. **P. Javana**, DC. Prod. i. 327 (W. et A. Prod. i. 38.)—c.p. 184.  
 Var. *β. angustifolia*: *foliis* lineari-oblongis.  
 HAB. Var. *a*, north of the island, *Gardner*. Var. *β*, Balangodde, *Gardner*, and Maturatte.
5. **P. leptalea**, DC. Prod. i. 325.—*P. oligophylla*, DC., l. c.—c.p. 3550.  
 HAB. Nilgalla, in the Badalla District; amongst high grass, rare.
6. **P. arillata**, Ham. (W. et A. Prod. i. 39; R. W. Icon. 946.)—c.p. 434 et 517.  
 HAB. Central Province, at an elevation of 4000 to 7000 feet.

### 2. SALOMONIA, Lour.

1. **S. oblongifolia**, DC. Prod. i. 334 (*S. leptostachya*, Wall. Cat.)—c.p. 1086.  
 HAB. Swampy ground in the south of the island.  
 Mr. W. Ferguson, who has been examining Hermann's Herbarium in the British Museum, informs me that this plant is there called *Polygala ciliata*.
2. **S. cordata**, Arn. Pugillus, 4 (322.)—*S. cordata* et *S. obovata*, Wight, Illust. 49 et 50. t. 22 B, etc.—c.p. 2906.  
 HAB. Similar places with the preceding.  
 The spikes of flowers are more compact in this species, and the cilia upon the capsules are much longer.

## 3. XANTHOPHYLLUM, Roxb.

1. **X. flavescens**, Roxb. (W. et A. Prod. i. 39, cum syn.)—c.p. 1244.  
HAB. South of the island, at no great elevation.

## XX. TAMARISCINEÆ.

## 1. TRICHAURUS (Arn. MSS.), W. et A.

1. **T. ericoides**, Arn. MSS. (W. et A. Prod. i. 40, cum syn; R. W. Illust.  $\frac{24}{141}$  B.)—c.p. 1254.  
HAB. Negombo, Puttam, and Moolitivoë, *Gardner*.

## XXI. ELATINEÆ.

## 1. BERGIA, Linn.

1. **B. verticillata**, W. et A. Prod. i. 41, cum syn.—c.p. 3551.  
HAB. Swampy places in the Batticaloa District.  
2. **B. ammannoides**, Roxb.—*Elatine* (*Bergia*) *ammannoides*, W. et A. Prod. i. 41, cum syn.; Wight, Illust. p. 54. t. 25 A.—c.p. 1543.  
HAB. North of the island, *Gardner*.

## XXII. SESUVIACEÆ.

## 1. TRIANTHEMA, Sauv.

1. **T. crystallina**, Vahl, (W. et Prod. i. 355, cum syn.)—c.p. 3552.  
HAB. Batticaloa District.  
2. **T. obcordata**, Roxb. (W. et A. Prod. i. 355, cum syn.; R. W. Icon. t. 288.)—c.p. 1102.  
HAB. Very common on the seacoast of the island.  
3. **T. decandra**, Lin. (W. et A., l. c. cum syn.; Wight, Icon. t. 296.)—c.p. 1104.  
HAB. East of the island, *Gardner*; Colombo, etc. Nom. vulg. "Saarana." Used as a medicine.

## 2. SESUVIUM, Linn.

1. **S. repens**, Rottb. (W. et A. Prod. i. 361, cum syn.)—c.p. 1098.  
HAB. Trincomalee, *Gardner*.

## XXIII. PORTULACEÆ.

## 1. PORTULACA, Tourn.

1. **P. oleracea**, Lin. (W. et A. Prod. i. 356, cum syn.)—c.p. 1103.  
HAB. Abundant throughout the island. Nom. vulg. "Genda-kola."  
2. **P. quadrifida**, Lin. (W. et A., l. c. cum syn.)—c.p. 1100.  
HAB. Very abundant. Nom. vulg. "Heen-genda-kola."

3. **P. suffruticosa**, R. W. (W. et A., l. c.)—C.P. 1101.  
 HAB. Damboul, *Gardner*.

## XXIV. CARYOPHYLLACEÆ.

### 1. STELLARIA, Linn.

1. **S. drymarioides**, Thw.; *foliis* rotundatis, apiculatis, basi in petio-  
 lum angustatis, nervo intramarginali conspicuo munitis; *cymis* termi-  
 nalibus, subdichotomis; *floribus* post anthesin cernuis.—C.P. 400.

HAB. Hapootelle Pass, at an elevation of about 4000 feet. April,  
 1856.

*Caulis* tetragonus, procumbens, elongatus, radicans, *ramulis* erectis. *Folia*  $\frac{1}{2}$ – $1\frac{1}{4}$   
 poll. longa et æquilata, sparse pilosa, *petiolo* 1–4 lin. longo. *Cymæ* viscoso-tomen-  
 tosæ, ramis sæpe elongatis. *Pedicelli* flores subæquantes. *Sepala* 5, ovata, sub-  
 acuta, *petalis* albis divaricato-bifidis duplo longiora. *Discus annularis* parvus, 5-  
 lobus; *lobis* rotundatis, cum *petalis* alternantibus.

2. **S. media**, Sm.—C.P. 3090.

HAB. Newera Ellia, probably introduced.

### 2. CERASTIUM, Linn.

1. **C. Indicum**, W. et A. Prod. i. 43.—C.P. 2957.

HAB. Newera Ellia, Horton Plains, and other very elevated parts of  
 the island.

In Ceylon specimens the leaves are lanceolate, and the flowers are of about equal  
 length with their pedicels.

### 3. MOLLUGO, Linn.

1. **M. nudicaulis**, Linn. (W. et A. Prod. i. 43, cum syn.)—C.P.  
 2809.

HAB. Colombo, etc.

2. **M. pentaphylla**, Linn., including *M. triphylla*, Linn.; W. et A.,  
 l. c.—C.P. 1096.

HAB. Common in the warmer parts of the island.

3. **M. stricta**, Lin. (W. et A. Prod. i. 44, cum syn.)—C.P. 1097.

HAB. Not uncommon with the preceding.

4. **M. cerviana**, Ser. (W. et A., l. c. cum syn.)—C.P. 1092.

HAB. Common in hot, sandy places. Nom. vulg. "Pat-paadagan."  
 Used as a medicine in fevers.

5. **M. disticha**, Ser. (W. et A., l. c. cum syn.)—C.P. 1094.

HAB. Near the seacoast; not uncommon.

6. **M. Spergula**, Linn. (W. et A., l. c. cum syn.)—C.P. 1095.

HAB. Common in the warmer parts of the island.

### 4. GLINUS, Linn.

1. **G. dictamnoides**, Linn. (W. et A. Prod. i. 362.)—C.P. 2937.

HAB. Not uncommon in the hot, drier parts of the island.

I have found ten stamens in all the Ceylon specimens I have examined.

## XXV. PARONYCHIACEÆ.

## 1. POLYCARPÆA, Lam.

1. **P. corymbosa**, Lam. (W. et A. Prod. i. 358, cum syn.; R. W. Icon. t. 712.)—C.P. 2383.

HAB. Not uncommon in the drier parts of the island, up to an elevation of 4000 feet.

## 2. HAPALOSIA, Wall.

1. **H. Lœfflingiæ**, Wall. (W. et A. Prod. i. 358, cum syn.)—C.P. 1090.

HAB. Warm, drier parts of the island.

## 3. DRYMARIA, Willd.

1. **D. cordata**, Willd. (W. et A. Prod. i. 359, cum syn.)—C.P. 1091.

HAB. Central Province, up to an elevation of 4000 feet.

## XXVI. LINEÆ.

## 1. LINUM, Linn.

1. **L. Mysorense**, Heyne, (W. et A. Prod. i. 134, cum syn.)—C.P. 3127.

HAB. Abundant between Newera Ellia and Badulla, at an elevation of from 4000 to 6000 feet.

*Folia* trinervia, acuta. *Petala* pallide flavâ, rotundata, obtusa. *Filamenta* a supra-medio dilatata, basi coalita; dentibus intermediis nullis.

## XXVII. HUGONIACEÆ.

## 1. HUGONIA, Linn.

1. **H. Mystax**, Linn. (W. et A. Prod. i. 72, cum syn.; Wight, Illust. 79. t. 32.)—C.P. 1185.

HAB. Not uncommon in the warmer parts of the island. Nom. vulg. "Boogatteya."

2. **H. ferruginea**, W. et A., l. c.—C.P. 1187.

HAB. Hot, drier parts of the island.

## XXVIII. MALVACEÆ.

## 1. URENA, Linn.

1. **U. lobata**, Linn. (W. et A. Prod. i. 46, cum syn.)—C.P. 3328.

HAB. Very common throughout the island. Nom. vulg. "Patta-appelle." Liber used for cordage.

2. **U. sinuata**, Linn. (W. et A., l. c. cum syn.)—C.P. 1129.

HAB. Equally common with the preceding. Nom. vulg. "Heen-appelle."



## 2. PAVONIA, Cav.

1. *P. odorata*, Willd. (W. et A. Prod. i. 47, cum syn.)—c.p. 1125.  
 HAB. Hot, drier parts of the island, up to an elevation of 1500 feet.
2. *P. Zeylanica*, Cav. (W. et A., l. c. cum syn.)

I have not met with a plant corresponding quite with the description given of this, and suspect it is but a form of the preceding, which I find with the leaves of the involucre varying in number from eleven to fifteen. Its flowers, when fresh, are either white or very pale red, becoming of a dull reddish colour when dry.

## 3. HIBISCUS, Linn.

1. *H. Surattensis*, Linn. (W. et A. Prod. i. 48, cum syn.)—c.p. 1118.  
 HAB. Very common in the hotter parts of the island. Nom. vulg. "Naapiritta."
2. *H. furcatus*, Roxb. (W. et A., l. c. cum syn.)—c.p. 727.  
 HAB. Very common up to an elevation of 2000 feet. Nom. vulg. "Heen-naapiritta."
3. *H. Lampas*, Cav. (W. et A. Prod. i. 48, cum syn.)—c.p. 3555.  
 HAB. Near Nilgalla, in the Badalla District.
4. *H. panduriformis*, Burm. (W. et A. Prod. i. 50, cum syn.)—c.p. 3528.  
 HAB. Borders of a tank, at Toonkimy, in the Atakalan Corle. September, 1857.
5. *H. vitifolius*, Linn. (W. et A., l. c. cum syn.)—c.p. 1122.  
 HAB. Not uncommon, up to an elevation of 3000 feet.
6. *H. eriocarpus*, DC. (W. et A., l. c. cum syn.)—c.p. 1119.  
 HAB. In the hot, drier parts of the island, up to an elevation of 1500 feet.
7. *H. micranthus*, Linn. (W. et A. Prod. i. 51, cum syn.)—c.p. 1126.  
 HAB. Hot, drier parts of the island, very abundant.

## 4. PARITIUM, Ad. Juss.

1. *P. tiliaceum*, Ad. Juss. (W. et A. Prod. i. 52, cum syn.)—c.p. 1120.  
 HAB. Not uncommon up to an elevation of 2000 feet. Nom. vulg. "Bellipatta." The liber is used by the Cinghalese for rough ropes.

## 5. ABELMOSCHUS, Med.

Nom. vulg. "Kapookanissa."

1. *A. angulosus*, Wall. (W. et A. Prod. i. 53, cum syn.); *involucello* (calyce, W. et A.) *subpersistente*; *calyce* membranaceo, tenero, *spathæformi*, cum *corolla deciduo*.

Var. *a. grandiflorus*; *petiolis* scabris, nec hispidis; *corolla* majore, flava; *seminibus* subpyriformibus.—c.p. 2567.

Var. *β. purpureus*; *petiolis* pilis rigidis hispidis; *corolla* purpurea, *venis* pallidioribus; *seminibus* globosis.—c.p. 1117.

HAB. Var. *a.* Not uncommon in the Central Province, up to an elevation of 4000 feet. Var. *β.* Newera Ellia.

There appears to have been some misconception with regard to the involucl and calyx of this species. The true calyx is of a very tender membranaceous structure, consisting of five sepals united at their edges into the form of a spathe, which falls at the same time with the corolla.

Vars.  $\alpha$  and  $\beta$  may possibly prove to be specifically distinct.

2. **A. moschatus**, Mœnch. (W. et A. Prod. i. 53, cum syn.)—C.P. 726.

HAB. South of the island; not uncommon.

3. **A. ficulneus**, W. et A., l. c.—C.P. 1157.

HAB. Colombo, Moon. Borders of the Minery Tank.

## 6. THESPESIA, Corr.

1. **T. populnea**, Corr. (W. et A. Prod. i. 54, cum syn.)—C.P. 1121.

HAB. Colombo, Moon. Calpentyne, Gardner; near Vavenery, in the Batticaloa District, apparently wild. Much used for planting as an ornamental and useful tree. Nom. vulg. "Sooreya-gass."

## 7. LAGUNEA, Cav.

1. **L. lobata**, Willd. (W. et A. Prod. i. 55, cum syn.)—C.P. 67.

HAB. Very abundant in the hot, drier parts of the island. Galagama, Gardner.

## 8. ABUTILON, Dill.

Nom. vulg. "Anoda."

1. **A. polyandrum**, W. et A., l. c. cum syn.—C.P. 1186.

HAB. I have not met with this in a wild state, and there is no record of where the herbarium specimen was found.

2. **A. Indicum**, G. Don, (W. et A., Prod. i. 56, cum syn.)—C.P. 1133.

HAB. Not uncommon.

3. **A. Asiaticum**, G. Don, (W. et A., l. c. cum syn.)—C.P. 1135.

HAB. Not uncommon.

4. **A. graveolens**, W. et A., l. c. cum syn.—C.P. 1134.

HAB. Common.

5. **A. tomentosum**, W. et A., l. c. cum syn.—C.P. 3529.

HAB. Near Taugalle, Sept. 1857.

## 9. WISSADULA, Medilo.

1. **W. periplocifolia**, Presl.—*Abutilon periplocifolium*, G. Don; W. et A. Prod. i. 55, cum syn.—C.P. 1132.

HAB. Very common in the south of the island.

## 10. SIDA, Linn.

Nom. vulg. "Babilla."

1. **S. acuta**, Burm. (W. et A. Prod. i. 57, cum syn.; Wight, Icon. t. 95.)—C.P. 2760.

HAB. Hot, drier parts of the island.

2. **S. rhombifolia**, Linn. (DC. Prod. i. 462.)—c.p. 1131 (1130).

HAB. Very abundant.

3. **S. retusa**, Linn. (W. et A. Prod. i. 58, cum syn.)—c.p. 3117.

HAB. Very common. Nom. vulg. "Kotekau-babilla." This is used as a medicine by the Cinghalese.

4. **S. alba**, Linn. (W. et A., l. c. cum syn.)—c.p. 3553.

HAB. Vavenery, in the Batticaloa District.

5. **S. cordifolia**, Linn. (W. et A., l. c. cum syn.)—c.p. 259.

HAB. Hot, drier parts of the island.

6. **S. Mysorensis**, Herb. Madr. (W. et A. Prod. i. 59, cum syn.)—c.p. 1123.

HAB. Not uncommon in the warmer parts of the island.

7. **S. humilis**, Willd. (W. et A., l. c. cum syn.)—c.p. 1124.

HAB. Very abundant.

## XXIX. STERCULIACEÆ.

### 1. ADANSONIA, Linn.

1. **A. digitata**, Linn. (W. et A. Prod. i. 61, cum syn.)—c.p. 1141.

HAB. Naturalized in the north of the island.

### 2. SALMALIA, Sch. et End.

1. **S. Malabarica**, Sch. et End. (Walp. Rep. i. 330, cum syn.; Wight, Illust. p. 72, t. 29.)—c.p. 545.

HAB. Common in the warmer parts of the island. Nom. vulg. "Katoo-imbool-gass."

### 3. ERIODENDRON, DC.

1. **E. orientale**, Steud. Nom. 587.—*E. anfractuosum*, DC.; W. et A., l. c. cum syn.; Wight, Icon. t. 400.—c.p. 1138.

HAB. Very common, up to an elevation of 2000 feet. Nom. vulg. "Imbool-gass."

### 4. CULLENIA, Wight.

1. **C. excelsa**, Wight, Icon. Pl. Ind. Orient. t. 1761-2.—*Durio Zeylanicus*, Gardner, in Calc. Journ. of Nat. Hist. vol. viii.—c.p. 734.

HAB. Common in the Central Province, at an elevation of from 2000 to 5000 feet. Nom. vulg. "Katoo-bodde, Kattooherileya."

### 5. HELICTERES, Linn.

1. **H. Isora**, Linn. (W. et A. Prod. i. 60, cum syn.; Wight, Icon. t. 180.)—c.p. 1137.

HAB. Common in the hotter parts of the island. Nom. vulg. "Leeneya-gass."

### 6. HERITIERA, Ait.

1. **H. littoralis**, Ait. (W. et A. Prod. i. 63, cum syn.)—c.p. 1143.

HAB. South of the island, not uncommon. Nom. vulg. "Attoona."

### 7. STERCULIA, Lin.

1. **S. Balanghas**, Linn. (W. et A. Prod. i. 62, cum syn.; Wight, Illust. p. 78. t. 30.)—C.P. 1142.

HAB. Hotter parts of the island, common.

2. **S. guttata**, Roxb. (W. et A. Prod. i. 62, cum syn.; Wight's Icon. t. 487.)—C.P. 2723.

HAB. Ambagamowa District, at no great elevation.

3. **S. urens**, Roxb. (W. et A. Prod. i. 63, cum syn.)—C.P. 2858.

HAB. Hot, drier parts of the island, not uncommon.

4. **S. foetida**, Linn. (W. et A., l. c. cum syn.; Wight, Icon. t. 181 and 364.)—C.P. 1140.

HAB. Very common in the warmer parts of the island. Nom. vulg. "Telemboo." The seeds are eaten by the Cinghalese.

### 8. PTERYGOTA, Endl.

1. **P. alata**, Br. in Brown et Bennett, Pl. Jav. Rar. 234.—*P. Roxburghii*, Sch. et Endl., Walp. Rep. v. 103. *Sterculia alata*, Roxb. Cor. iii. p. 84. t. 287.—C.P. 2963.

HAB. Banks of the Mahawelliganga, at Haragam.

### 9. FIRMIANA, Marsili.

1. **F. colorata**, Br. l. c. 235, cum syn.—C.P. 2566.

HAB. Hot, drier parts of the island.

## XXX. BYTTNERIACEÆ.

### 1. GUAZUMA, Plum.

1. **G. tomentosa**, H.B.K. (W. et A. Prod. i. 64; Wight, Illust. 78. t. 31.)—C.P. 1114.

HAB. Jaffna, *Gardner*.

### 2. Gen. Nov. PITYRANTHE, Thw.

*Calyx* urceolatus, 3-4-lobatus. *Petala* 5, æqualia, subcuneiformia. *Stamina fertilia* 15, ternatim cum alteris 5 *sterilibus* acutis spathulatis æquilongis alternantia; *filamentis* liberis vel imo basi cohærentibus; *antheris* adnatis, loculis divaricatis. *Ovarium* sessile, 5-loculare, loculis 2-ovulatis; *ovulis* superpositis; *stylis* 5, arcte coalitis; *stigmatibus* minutis, simplicibus. *Capsula* inflata, turbinata, sæpissime 1-sperma, loculicide dehiscens.

Genus *Kleinhovia* affine, sed calyce urceolato, petalis æqualibus, filamentis subliberis et ovario sessili differt.

1. **P. verrucosa**, Thw.—*Kleinhovia verrucosa*, Gardn. MS.—C.P. 1144.

HAB. Abundant in the Batticaloa District; Jaffna and Trincomalee, *Gardner*; Putlam, *A. O. Brodie, Esq.*

*Arbor*. *Folia* obovata, crenata, basi rotundata, 3-nervia, 2-6 poll. longa, 1-3

poll. lata, superne nitida, subtus pallida, pubescentia stellari tomentosa, *petiolo*  $\frac{1}{2}$ – $\frac{3}{4}$  poll. longo. *Paniculæ* terminales, lepidotæ. *Flores* circiter  $1\frac{1}{2}$  lin. longi. *Capsulæ* furfuraceæ, verrucosæ. *Semina* nigra, oblonga, 2 lin. longa, minutissime reticulata, pubescentia stellari pallida decidua vestita.

### 3. WALTHERIA, Linn.

1. **W. Indica**, Linn. (W. et A. Prod. i. 67, cum syn.)—c.p. 1128.  
HAB. Batticaloa, *Gardner*.

### 4. MELOCHIA, Linn.

1. **M. corchorifolia**, Linn.—*Riedleia corchorifolia*, DC. (W. et A. Prod. i. 66, cum syn.)—c.p. 1127.  
HAB. Very abundant.

### 5. PTEROSPERMUM, Schreb.

1. **P. suberifolium**, Lam. (W. et A. Prod. i. 68, cum syn.)—c.p. 1136.  
HAB. Common up to an elevation of 2000 feet, especially in the drier parts of the island. Nom. vulg. "Velenge." The wood is useful for many purposes where toughness is required.

### 6. KYDIA, Roxb.

1. **K. axillaris**, Thw.; *foliis* integris, ovato-lanceolatis, lævibus; *floribus* solitariis, axillaribus.—c.p. 3126.

HAB. Near Badulla, at an elevation of about 2000 feet.

*Arbor* mediocris. *Folia*  $2\frac{1}{2}$ – $4\frac{1}{2}$  poll. longa,  $1\frac{1}{4}$ –2 poll. lata, subtus pallidiora, basi rotundata, *petiolo* 3–4 lin. longo. *Involucelli* fructiferi segmenta lineari-oblonga, 1 poll. longa, 3– $3\frac{1}{2}$  lin. lata. *Calyx* cupuliformis, 5-dentatus,  $\frac{1}{2}$  poll. longus. *Ovarium* biloculare. *Fructus* hispidus.

This species I have only seen in fruit, and am therefore not quite certain that it is a true *Kydia*. The axillary inflorescence and the character of the foliage give it a very different aspect from that of *K. calycina*. The latter species has not, so far as I am aware, been detected wild in Ceylon, though it most probably occurs so in the north of the island.

### 7. Gen. Nov. JULOSTYLIS, Thw.

*Involucellum* 4-phyllum, persistens. *Calyx* 5-partitus, persistens. *Corollæ* petala 5, basi coalita, subtus ad sinus plicato-appendiculata. *Tubus stamineus* cylindricus, apice dentatus, anantherus, medium versus *filamentis* 10, biserialibus, æquilongis, *monantheriferis* cinctus, basi corollæ adnatus. *Antheræ* reniformes. *Ovarium* biloculare; *loculis* 2-ovulatis; *ovulis* collateralibus, adscendentibus. *Stylus* supra medium bifidus. *Stigmata* disciformia.

1. **J. angustifolia**, Thw.—*Kydia angustifolia*, Arn. Pug. Pl. Ind. Orient. 4 (322).—c.p. 1146.

HAB. South of the island, not uncommon.

*Arbor* mediocris, parce ramosa. *Folia* 5–8 poll. longa,  $1\frac{1}{2}$ –3 poll. lata, pubescentia stellari-hirsuta, subtus pallidiora, *petiolo* 3–5 lin. longo. *Paniculæ* terminales, pendulæ. *Flores*  $\frac{1}{2}$  poll. lati. *Corolla* cupuliformis, pallida, interne dimidio inferiore rufa. *Stylus* versus apicem dense lanuginosus.

The anthers of this species are quite those of a Malvaceous plant, but from its near affinity with *Kydia* it must remain in the same Natural Order in which that is located.

## XXXI. TILIACEÆ.

## SECT. TILIEÆ.

## 1. CORCHORUS, Linn.

1. *C. urticæfolius*, W. et A. Prod. i. 73, cum syn.—c.p. 2758.  
HAB. Trincomalee, *Moon*.
2. *C. olitorius*, Linn. (W. et A., l. c. cum syn.)—c.p. 1107.  
HAB. Hotter parts of the island, common.
3. *C. acutangulus*, Lam. (W. et A., l. c. cum syn.; Wight, Icon. t. 739.)—c.p. 1106.  
HAB. Same situations as the last, common.
4. *C. capsularis*, Linn. (W. et A., l. c. cum syn.; Wight, Icon. t. 311.)—c.p. 1105.  
HAB. Very common.

## 2. TRIUMFETTA, Plum.

Nom. vulg. "Appela."

1. *T. pilosa*, Roth. (W. et A. Prod. i. 74, cum syn.)  
Var. *a*; capsularum echinis firmis, 2 lin. longis, prope basin parce pilosis, apiculo vix curvato.—c.p. 2902.  
Var. *β*; capsularum echinis debilibus, 4½ lin. longis, dense longe pilosis, apiculo hamato.—c.p. 730.  
HAB. Var. *a*. Common in cultivated ground. Var. *β*. Maturatte.
2. *T. angulata*, Lam. (W. et A., l. c. cum syn.; Wight, Icon. t. 320.)—c.p. 2759.  
HAB. Common everywhere.

## 3. GREWIA, Juss.

Nom. vulg. "Caaleya."

1. *G. orientalis*, Linn. (W. et A. Prod. i. 76, cum syn.)—c.p. 1110, 1112, 2419, and 2997.  
HAB. Warmer parts of the island, up to an elevation of 1500 feet.
2. *G. columnaris*, Sm. (W. et A., l. c. cum syn.)—c.p. 1111.  
HAB. Hot, drier parts of the island, not uncommon.
3. *G. hirsuta*, Vahl, (W. et A., l. c. i. 78, cum syn.)—c.p. 3554.  
HAB. Open, grassy places in the east of the island.
4. *G. helicterifolia*, Wall. MSS. (Walp. Rep. ii. 799.)—c.p. 1113.  
HAB. In similar situations with the last, and equally common.  
Dr. Wight named this "*G. rotundifolia*, MSS., var. fol. obliquis acuminatis." It differs from the description of *G. helicterifolia* in the peduncles being commonly much longer than the petioles.
5. *G. diplocarpa*, Thw.; foliis lanceolatis, acuminatis, crenato-serratis, 3-nerviis, sublaevibus; fructibus parce strigosis, bipartitis, segmentis bilobis.—c.p. 2568.

HAB. Maturatte and Oova, at an elevation of about 3000 feet.

*Folia* 3-4 poll. longa,  $1\frac{1}{4}$ - $1\frac{3}{4}$  lata, juniora parce hirsuta, *petiolo* 4-5 lin. longo, hirsuto. *Stipulæ* parvæ, lanceolatae, hirsutæ. *Pedunculi* solitarii, axillares, petiolis æquilongi, 3-flori. *Bracteoli* parvi. *Sepala* linearia, extus cinereo-tomentosa, petalis lanceolatis acutis plus quam duplo longiora. *Stigma* dilatatum, 4-lobatum.

6. **G. tiliæfolia**, Vahl, (W. et A. Prod. i. 80, cum syn.)—c.p. 621.

HAB. Warmer parts of the island, common.

7. **G. Microcos**, Linn. (W. et A. Prod. i. 81, cum syn.; Wight, Illust. 81. t. 33.)—c.p. 1109.

HAB. Very abundant, up to an elevation of 3000 feet.

#### 4. BERRYA, Roxb.

1. **B. Ammonilla**, Roxb. (W. et A., l. c. cum syn.; Wight, Illust. 82. t. 34.)—c.p. 1108.

HAB. Hot, drier parts of the island. Nom. vulg. "Hal-mililla." The wood of this fine tree is very valuable for building and other purposes.

#### SECT. ELEOCARPEÆ.

#### 5. ELÆOCARPUS, Linn.

Nom. vulg. "Weraloo, Gal-weraloo."

1. **E. serratus**, Linn. (Fl. Zeyl. 206; Burm. Th. Zeyl. t. 40.)—c.p. 539 (554, 1115).

HAB. Warmer parts of the island, up to an elevation of 2000 feet.

*Stamina* 30-35, omnia ad valvæ exterioris apicem setis 3-4 longis rigidis patentibus instructa. *Drupa* oblonga.

The fleshy exterior of the drupes of this and the following species are eaten by the Cinghalese.

2. **E. obovatus**, Arn. Pug. Pl. Ind. Or. 5 (323).—*E. coriaceus*, Hook. Ic. Plant. v. 2. t. 154.—c.p. 316.

HAB. Newera Ellia and other elevated parts of the island, at an elevation of from 6000 to 8000 feet.

*Drupa* sphaerica, putamine ruguloso-tuberculato.

3. **E. montanus**, Thw.; *foliis* ovatis vel rotundatis, obtusis, denticulatis, parce hirsutis (junioribus subtus tomentosis), *venis* primariis ad axillas ramificationes divaricatasque conspicue glanduliferis; *antheris* nudis; *drupis* sphaericis.—c.p. 624 (630, 642).

HAB. In similar localities with the last.

*Arbor* mediocris. *Folia* 2-3½ poll. longa, *petiolo* ½-1 poll. longo. *Racemi* axillares, foliis æquilongi, tomentosi.

4. **E. amœnus**, Thw.; *foliis* lævibus, lanceolatis vel ovato-lanceolatis, brevi-acuminatis, obtusis, crenato-serratis, denticulatis, subtus ad nervarum primariarum axillas glanduliferis; *antheris* nudis; *drupis* sphaericis.—c.p. 2587 and 2651 (? c.p. 2524).

HAB. Central Province, up to an elevation of 4000 feet.

*Arbor* mediocris. *Gemmæ* pilosæ. *Folia* 2-4 poll. longa,  $\frac{3}{4}$ -2 poll. lata, *petiolo* 4-8 lin. longo. *Racemi* numerosi, parce pilosi, e foliorum delapsorum axillis, foliis breviores.

Some specimens (c.p. 2524) in the herbarium are somewhat different in appear-

ance, but they are not sufficiently perfect to enable me to determine whether they are really distinct from or are a mere state of the present species.

5. **E. subvillosus**, Arn. Pug. 5 (323).—*E. pubescens*, Hooker, Ic. Pl. ii. t. 155. *E. integrifolius*, Moon's Cat.—c.p. 704.

HAB. Saffragam District, up to an elevation of 2000 feet.

*Drupa* oblonga, putamine tuberculato, apice acuto.

## 6. MONOCERA, Jack.

1. **M. glandulifera**, Hook. (Wight, Illust. p. 84.)—c.p. 675 (2476).

HAB. Central Province; not uncommon, at an elevation of from 4000 to 6000 feet.

2. **M. Walkeri**, Wight, l. c.—? *M. Ceylanica*, Arn. Pug. p. 5 (323).—c.p. 8 and 166 (172, 421).

HAB. Saffragam District, not very common, at an elevation of about 5000 feet.

I suspect this must be Dr. Arnott's *M. Ceylanica*, which, having been described, as he states, from a single specimen, may not indicate the typical character of the species.

## XXXII. DIPTEROCARPEÆ.

### 1. DIPTEROCARPUS, Gærtner.

1. **D. Zeylanicus**, Thw.; *foliis* magnis, ovatis vel oblongis, brevi-acuminatis, glabris, petiolo longo costaque leprosis; *gemma* villosis; *calycis fructiferi* tubo urceolato, 5-costato.—c.p. 1921.

HAB. Abundant, up to an elevation of 3000 feet. Nom. vulg. "Hora-gass."

*Arbor* ingens. *Racemi* axillares, solitarii, prope basin bifidi, 6-12-flori, ad ramulorum apices aggregati, leprosi. *Pedicelli* tubo calycis campanulato paulo breviores.

The fruit of this species closely resembles Gærtner's *D. costatus* (De Fruct. tab. 187) as regards the size and form of the segments of the enlarged calyx, but differs from it essentially in the constriction beneath these. From *D. laevis*, Hamilton, the only other species with which it can be confounded, its costate or angled fruit would seem sufficiently to distinguish it.

A gum-resin and wood-oil are obtained from this and the species following.

2. **D. hispidus**, Thw.; *foliis* maximis, oblongis, brevi-acuminatis, subcordatis, superne ad costam et subtus *petioloque* pilis rigidis fasciculatis patentibus hirsutis; *calycis fructiferi* tubo spherico vel subpyriformi, pedicellato, leproso, apice 5-angulo, contracto.—c.p. 2903. (c.p. 3405 *in parte*.)

HAB. Saffragam District, at no great elevation. Nom. vulg. "Boo-hora-gass."

This species is well distinguished by its large hispid leaves, one of which in my collection, from a young tree, is 19 inches long and 9 inches wide, with a petiole  $2\frac{1}{2}$  inches long. The enlarged calyx-lobes are 6-7 inches long.

3. **D. oblongifolius**, Thw.; *foliis* oblongis, brevi-acuminatis, basi rotundatis, superne ad costam et subtus ad costam venas primariasque hirsutis; *ramis junioribus*, *gemma* *petiolisque* pilis rigidis nitentibus subpatentibus hispidis; *racemis* supra-axillaribus, solitariis, simplicibus,



2-4-floris, hispido-leprosis; *calyce* incano-leproso, 5-angulato, in *pedicello* brevi desinente.—C.P. 3405.

HAB. Near Ratnapoora, April, 1855.

*Arbor* ingens. *Folia* 5-7 poll. longa,  $2\frac{1}{2}$ -4 poll. lata, *petiolo* 1-2 poll. longo.

4. **D. scabridus**, Thw.; *foliis* lineari-oblongis, brevi-acuminatis, basi parce angustatis, superne glabris, subtus pubescentia stellari sparsa scabridis, *costa* subtus *petioloque* pilis rigidis adpressis hirsutis; *calycis fructiferi* tubo parce leproso, subsphærico, angulis 5 singulis in membranam undulatam *pedicello* brevi attingentem productis.—C.P. 2692. (C.P. 3406, *in parte*.)

HAB. Near Ratnapoora, September, 1855.

*Arbor* ingens. *Folia* circiter 7 poll. longa,  $2\frac{1}{2}$  lata, *petiolo*  $\frac{3}{4}$  poll. longo. *Calycis fructiferi* laciniæ majores  $3\frac{1}{2}$ - $4\frac{1}{2}$  poll. longæ,  $\frac{3}{4}$ -1 poll. latæ.

The adpressed pubescence and the shape of the leaves distinguish this species.

5. **D. glandulosus**, Thw.; *foliis* ovatis, acuminatis, basi rotundatis, superne lævibus, *costa* venisque primariis hirsutis, subtus pilis minutis coloratis conspersis; *ramis junioribus*, *gemmis petiolisque* piloso-leprosis; *racemis* axillaribus, solitariis, 2-3-floris, leprosis; *calyce* 5-angulato, in *pedicello* brevi desinente, leproso; *calycis fructiferi* tubo subsphærico, angulis in membranam medio latissimam *pedicello* tenui sæpissime non attingentem productis.—C.P. 2590.

HAB. Saffragam and Ambagamowa Districts, at no great elevation. Nom. vulg. "Dorana-gass."

*Arbor* ingens. *Folia* 4-6 poll. longa,  $2-2\frac{1}{2}$  poll. lata, *petiolo* 6-8 lin. longo. *Calycis fructiferi* laciniæ majores 3-4 poll. longæ, 7-8 lin. latæ.

This species is easily recognized by the glandular pubescence on the under side of its leaves, which is at first pale-yellow, then red, and in the old leaves nearly black.

6. **D. insignis**, Thw.; *foliis* ovatis, acuminatis, lævibus, basi rotundatis, subtus *costa* venisque primariis hirsutis; *ramis junioribus*, *gemmis petiolisque* pubescentibus; *racemis* supra-axillaribus, solitariis, 3-5-floris, basi pubescentibus, apicem versus lævibus; *pedicellis* subnullis; *calyce* 5-angulato, angulis per totam longitudinem late membranaceo-alatis.—C.P. 3406.

HAB. Saffragam District, April, 1855.

*Arbor* ingens. *Folia*  $4\frac{1}{2}$ - $5\frac{1}{2}$  poll. longa,  $2-2\frac{1}{2}$  poll. lata, *petiolo* 7-9 lin. longo.

This species is easily distinguishable from the others by its smooth, longitudinally 5-bordered calyx, sessile upon the rachis of the raceme.

~~*D. turbinatus*, Roxb., cultivated in Ceylon, is C.P. 2398.~~

## 2. DOONA, Thw.

Hook. Journ. of Bot. vol. iv. (1852) p. 7.

*Stamina* 15, biserialia. *Embryo* cotyledonibus foliaceis valde inæqualibus; harum maxima inter stratum oleoso-albuminosum contorto-convoluta, in germinatione inclusa; altera brevissima, in germinatione sæpe emergente.

1. **D. Zeylanica**, Thw. l. c. vol. iii. t. 12; *foliis* lanceolatis vel ovato-lanceolatis, acuminatis, basi angustatis, venis inconspicuis; *pedicellis* calyce longioribus.—C.P. 2423.

HAB. Central Province, up to an elevation of 4000 feet. Nom. vulg. "Doon-gass."

A large quantity of colourless gum-resin exudes from the trunk and branches of this fine tree, which, when dissolved in spirits of wine or turpentine, makes an excellent varnish.

2. **D. affinis**, Thw.; *foliis lanceolatis vel lineari-lanceolatis, longe acuminatis, basi rotundatis, venis subinconspicuis; pedicellis calyce longioribus.*—C.P. 3409.

HAB. Between Ratnapoora and Galle, at no great elevation.

*Arbor ingens. Folia 2-3 poll. longa (acumine 4 lin. longo), 9-11 lin. lata, petiolo circiter 4 lin. longo.*

Closely allied to the preceding, but the leaves are larger and differently shaped, and the petioles much stouter.

3. **D. nervosa**, Thw.; *foliis ovatis, ovato-lanceolatis vel lanceolatis, longe acuminatis, opacis, venis venulisque prominulis, basi rotundatis; pedicellis calyce subæquilongis; calycis fructiferi laciniis majoribus spathulato-dilatatis.*—C.P. 3410.

HAB. Ekualagodde, near Ratnapoora, April, 1855.

*Arbor ingens. Folia 3-4 poll. longa (acumine  $\frac{1}{2}$  poll. longo),  $1\frac{1}{4}$ - $1\frac{3}{4}$  lata.*

4. **D. trapezifolia**, Thw.; *foliis trapezoideo-lanceolatis, acuminatis, basi rotundatis, superne minute reticulatis, subtus venis inconspicuis; pedicellis calyce æquilongis; calycis fructiferi laciniis majoribus spathulato-linearibus.*—C.P. 3341.

HAB. Central and southern parts of the island, up to an elevation of 1500 feet; a common forest tree. Nom. vulg. "Tuccahaaloo-gass."

*Arbor ingens. Folia  $2\frac{1}{2}$ - $3\frac{1}{2}$  poll. longa (acumine circiter 3 lin. longo), 9-14 lin. lata, petiolo 4 lin. longo.*

5. **D. congestiflora**, Thw.; *foliis ovato-lanceolatis, acuminatis, basi rotundatis vel subcordatis; floribus ad apices panicularum congestis; pedicellis brevissimis.*—C.P. 3411.

HAB. Hinidoon and Pasdoon Corles, April, 1855. Nom. vulg. "Tinneya-gass."

*Arbor ingens. Folia  $3\frac{1}{2}$ -5 poll. longa,  $1\frac{1}{4}$ -2 poll. lata, petiolo 4-5 lin. longo.*

This fine species is characterized by its numerous shortly-stalked flowers collected towards the ends of the panicles. The fruit, with its enlarged calyx, closely resembles that of the preceding and following species.

6. **D. Gardneri**, Thw.; *foliis ovatis vel ovato-lanceolatis, acuminatis, basi rotundatis, longe petiolatis; floribus cernuis, pedicellis calyce brevioribus.*—C.P. 1919.

HAB. Central Province, at an elevation of from 3000 to 5000 feet.

*Arbor ingens. Folia 2-3 poll. longa,  $1\frac{1}{4}$ - $1\frac{3}{4}$  lata, petiolo 6-7 lin. longo.*

A very distinct and beautiful species.

7. **D. cordifolia**, Thw.; *foliis oblongis, abrupte acuminatis, cordatis, subtus nitidis, venis venulisque prominentibus.*—C.P. 3340.

HAB. Pasdoon Corle and Ambagamowa, at no great elevation. Nom. vulg. "Bairaleya."

*Arbor ingens. Folia 3-4 poll. longa (acumine circiter  $\frac{1}{2}$  poll. longo),  $1\frac{1}{2}$ - $2\frac{1}{2}$  poll. lata, petiolo 3-5 lin. longo.*

I have not yet been able to obtain the flowers of this species, which is said to produce them only once in three years. Some seeds which were given to me

were unfortunately all sown, and none reserved for the herbarium. The Cinghalese eat the ripe seeds after toasting or frying them.

### 3. SHOREA, Roxb.

1. **S. oblongifolia**, Thw.; *foliis* lineari-oblongis, abrupte acuminatis, basi truncato-rotundatis vel cordatis; ramis *paniculæ* secundifloris.—C.P. 3005.

HAB. Saffragam and other districts in the south of the island, at no great elevation.

*Arbor* ingens. *Folia* 4-6 poll. longa,  $1\frac{3}{4}$ - $2\frac{1}{2}$  poll. lata, *petiolo* 6-8 lin. longo. *Ramuli* juniores *petiolique*, cum *paniculis* supra-axillaribus, minute stellato-incani. *Pedicelli* brevissimi. *Flores* flavi, odorati, dorso strigosi. *Calyx fructiferus* minute tomentosus, laciniis sursum dilatatis. *Fructus* tomentoso-incanus.

This is evidently closely allied to, if it be not a variety of, *Vatica Tumbugaia*, W. and A., Wight, Icon. tab. 27, with which it would seem to correspond very nearly in the inflorescence and in the structure of the fruit, but the leaves are much longer compared with their width.

2. **S.? stipularis**, Thw.; *foliis* oblongis, brevi-acuminatis, basi rotundatis vel subcordatis, longe petiolatis, lævibus; *stipulis* magnis, ovatis, subobliquis, striatis, diu persistentibus.—C.P. 3408.

HAB. Between Ratnapoora and Galle, at no great elevation.

*Arbor* ingens, *ramulis* teretibus. *Folia* 4-6 poll. longa, 2-4 poll. lata, *petiolo* 12-14 lin. longo. *Stipulæ* 6-11 lin. longæ, 4-7 lin. latæ. *Calycis fructiferi* laciniæ sursum dilatatæ; tres majores 5-5½ poll. longæ,  $1\frac{1}{4}$  poll. latæ, 11-14-costatæ; duæ minores  $3\frac{1}{2}$ -4 poll. longæ, 5-6 lin. latæ, 6-8-costatæ. *Fructus* subsphæricus, apice conico acuto.

I have not seen the flowers, nor have I been able to ascertain the character of the inflorescence of this fine species, but, judging from the structure of the fruit, it must be very nearly allied to this genus. The large size and persistence of the stipules form a very striking character.

In the herbarium are specimens of the foliage of what appears to be another species of *Shorea* (C.P. 3407), but the description must be deferred until better materials can be collected.

### 4. HOPEA, Roxb.

1. **H. discolor**, Thw.; *foliis* lanceolatis, longe acuminatis, basi rotundatis, superne lævibus, ad costam venasque primarias canaliculatis, subtus brunneo-leprosis; *paniculis* axillaribus, ramis secundifloris; *pedicellis* brevissimis.—C.P. 3125.

HAB. Saffragam and Ambagamowa Districts, at no great elevation.

*Arbor* ingens, *ramulis* teretibus, nigro-rufis, leprosis. *Folia*  $2\frac{1}{2}$ - $3\frac{1}{2}$  poll. longa (acumine circiter  $\frac{1}{2}$  poll. longo), 8-13 lin. lata, *petiolo* 4-5 lin. longo. *Paniculæ* solitariae vel 2-3-fasciculatæ, foliis multo breviores, rufescentes, parce leprosæ. *Flores* flavo-rufescentes, odoratissimi. *Calycis fructiferi* laciniæ majores spathulatæ,  $2\frac{1}{4}$  poll. longæ. *Fructus* subconicus, lævis.

The rich brown colour of the under side of the leaves well distinguishes this beautiful species.

### 5. VATERIA, Linn.

*Stamina* numerosa, indefinita, multiserialia, apice bifida.

1. **V. Indica**, Linn. (W. et A. Prod. i. 83, cum syn.; Wight, Ill. t. 36; Gærtner, De Fruct. iii. 53. t. 189.)—c.p. 1918.

HAB. Common in the hotter parts of the island, up to an elevation of 2000 feet. Nom. vulg. "Hal-gass." A gum-resin exudes in considerable quantity from this handsome tree, and is much used by the Cinghalese for burning during some of their superstitious ceremonies.

*Arbor* ingens, trunco lævi. *Ramuli* teretes, *juniores* stellato-furfuracci. *Folia* oblonga vel lineari-oblonga, apiculata, basi cordata, 6-18 poll. longa, 3-6 poll. lata; *petiolo* tereti,  $1\frac{1}{2}$ - $2\frac{1}{2}$  poll. longo, apice incrassato. *Stipulæ* lineari-lanceolatae, dorso furfuraceæ, deciduæ, circiter 1 poll. longæ. *Paniculæ* ad apices ramorum, foliis subæquilongæ, flocculoso-furfuraceæ, pallide-ochraceæ, multifloræ. *Flores* brevipedicellati, bibracteolati. *Bracteolæ* ovatae, acutæ, 4 lin. longæ. *Stamina* numerosa, *filamentis* brevissimis, basi coalitis; *antheris* introrsis, longitudinaliter dehiscentibus, basi bi-auriculatis, loculis ad apicem divergentibus, acutis. *Fructus* 4 poll. longi,  $2\frac{1}{2}$ -3 poll. lati.

Blume, in his Mus. Bot. Lugd. Bat. ii. 29, makes two species of the Linnæan *Vateria Indica*, reserving this name for the Ceylon plant, and calling the Peninsular one *V. Malabarica*. I have no means of determining whether this view is correct, but in order to assist in clearing the matter of doubt, I have given above a pretty full description of the Ceylon plant.

The figure of this species in Dr. Wight's Illust. tab. 36 is not satisfactory. Gærtner's figure of the fruit is very characteristic, though it is rather smaller than ordinary specimens.

## 6. ISAUXIS, Arn.

*Stamina* 15, biserialia. *Calyx fructifer* multum auctus.

1. **I. Roxburghiana**, Wight, Illust. p. 88; Icon. Pl. Ind. Or. t. 26.—c.p. 3416.

HAB. South of the island, at no great elevation. Nom. vulg. "Mandora-gass." This tree yields a considerable quantity of gum-resin.

*Antheræ* planæ, obtusæ, connectivo in apiculum brevem producto. *Calyx fructifer* basi depressus, laciniis 5-nerviis. *Fructus* 3-sulcatus, apice depresso.

I have little doubt of this being Dr. Wight's plant, though there is a slight discrepancy between my specimens and the figure in the 'Icones;' but this is most likely due to a little want of accuracy on the part of the draftsman.

## 7. STEMONOPORUS, Thw.

Hook. Journ. of Bot. vol. vi. (1854) p. 67.

*Stamina* 15, biserialia, *antheris* apice per laterum involuptionem tubulatis. *Calyx fructifer* non auctus.

Sect. 1.—*Inflorescentia paniculata*.

1. **S. Wightii**, Thw.; *foliis* oblongis vel lineari-oblongis, apiculatis, basi rotundatis vel subcuneatis, subtus scabridis, longe petiolatis; *antheris* brevi-apiculatis.—*S. macrophyllus*, Thw. MSS. in Herb. Hook. ? *Vateria Ceylanica*, Wight, Illust. p. 88.—c.p. 3415.

HAB. In forests between Galle and Ratnapoora, and at Palmadoolla, near the latter place.

*Arbor* ingens. *Folia* 5-12 poll. longa, 3-6 poll. lata, *venis primariis* circiter 20, *petiolo*  $1\frac{1}{2}$ -3 poll. longo. *Paniculæ* foliis multo breviores.

2. **S. Gardneri**, Thw. l. c. tab. 2 A. ; *foliis* ovato-acuminatis, basi rotundatis vel subcordatis.—C.P. 1920.

HAB. Rambodde, *Gardner*; near Adam's Peak, at an elevation of about 5000 feet.

*Arbor* magna. *Folia* 3-5 poll. longa,  $1\frac{1}{4}$ - $2\frac{1}{2}$  poll. lata, *venis* *primariis* circiter 10, *petiolo*  $\frac{1}{2}$ -1 poll. longo. *Paniculæ* numerosæ ad apices ramulorum.

Sect. 2.—*Floribus* paucis, racemosis vel solitariis; *pedunculis* *pedicellis*que brevibus.

3. **S. affinis**, Thw. l. c. ; *foliis* ovato-oblongis, acuminatis, basi rotundatis; *pedunculis* sæpissime 1-floris.—C.P. 2430.

HAB. Hunasgiria District, at an elevation of about 4000 feet.

*Arbor* magna. *Folia* 3- $4\frac{1}{2}$  poll. longa (acumine circiter  $\frac{1}{2}$  poll. longo),  $1\frac{1}{2}$ - $2\frac{1}{2}$  poll. lata, *venis* *primariis* circiter 8, *petiolo* 6-8 lin. longo.

4. **S. lanceolatus**, Thw. l. c. ; *foliis* lanceolatis vel oblongo-lanceolatis, acuminatis, basi cuneatis, superne glabris, *venis* prominentibus reticulatis, subtus scabridis; *pedunculis* sæpissime 1-floris.—C.P. 2658.

HAB. Near Ratnapoora, at no great elevation.

*Arbor* parva. *Folia*  $4\frac{1}{2}$ - $8\frac{1}{2}$  poll. longa (acumine 4-8 lin. longo),  $1\frac{1}{4}$ -3 poll. lata, *venis* *primariis* circiter 8, *petiolo* 5-7 lin. longo.

5. **S. canaliculatus**, Thw. ; *foliis* oblongis, lanceolatis vel lineari-lanceolatis, plus minus abrupte acuminatis, basi rotundatis vel subcuneatis, superne ad costam canaliculatis, subtus scabridis lævibusve; *pedunculis* sæpissime 1-floris.—C.P. 3413.

HAB. Hinidoon and Reigam Corles, at no great elevation.

*Arbor* mediocris. *Folia* 4-6 poll. longa (acumine circiter  $\frac{1}{2}$  poll. longo),  $1\frac{1}{4}$ - $2\frac{1}{2}$  lata, *venis* *primariis* circiter 12, *petiolo*  $\frac{1}{2}$ - $1\frac{1}{2}$  poll. longo, apicem versus incrassato. *Fructus*  $1\frac{1}{4}$  poll. in diam.

This bears considerable resemblance to the species preceding and following it, but its canaliculated leaves distinguish it from both.

6. **S. petiolaris**, Thw. ; *foliis* lineari-oblongis, abrupte acuminatis, basi rotundatis vel subcuneatis, superne lævibus, subtus scabridis lævibusve, longe petiolatis; *pedunculis* 1-3-floris.—C.P. 3151.

HAB. Kittool Galle, at no great elevation.

*Arbor* mediocris. *Folia* 6-10 poll. longa (acumine  $\frac{1}{2}$ - $\frac{3}{4}$  poll. longo), 2- $2\frac{3}{4}$  poll. lata, *venis* *primariis* circiter 12, *petiolo* 1-3 poll. longo, apicem versus incrassato.

7. **S. rigidus**, Thw. l. c. ; *foliis* rigidissimis, scabridis, oblongis, apice rotundatis vel subapiculatis, basi cuneatis vel cuneato-rotundatis; *pedunculis* sæpissime 1-floris.—C.P. 2645.

HAB. Ambagamowa District, at an elevation of about 3000 feet.

*Arbor* magna. *Folia*  $3\frac{1}{2}$ - $5\frac{1}{2}$  poll. longa,  $1\frac{1}{2}$ - $2\frac{1}{4}$  lata, *venis* *primariis* circiter 9, *petiolo* 6-8 lin. longo.

8. **S. oblongifolius**, Thw. l. c. ; *foliis* lævibus, oblongis, subacuminatis, obtusis, basi rotundatis; *pedunculis* 1-4-floris.—C.P. 2646.

HAB. Ambagamowa District, at an elevation of about 4000 feet.

*Arbor* magna. *Folia* 4- $6\frac{1}{2}$  poll. longa,  $1\frac{3}{4}$ - $2\frac{1}{2}$  lata, *venis* *primariis* circiter 8; *petiolo* 4-6 lin. longo.

9. **S. reticulatus**, Thw. ; *foliis* oblongis vel oblongo-lanceolatis, longe acuminatis, basi rotundatis, superne nitidis, subtus opacis inter *venas* *primarias* reticulato- nec parallelo-venosis, longe petiolatis; *pedunculis* sæpissime 1-floris.—C.P. 3414.

HAB. Hinidoon Corle, at an elevation of about 1000 feet.

*Arbor mediocris. Folia*  $2\frac{1}{2}$ –5 poll. longa (acumine  $\frac{1}{2}$ –1 poll. longo),  $1\frac{1}{4}$ – $2\frac{1}{4}$  lata, *venis primariis* circiter 8, *petiolo* 7–15 lin. longo. *Fructus* rotundatus, apice parce acuminatus,  $1\frac{1}{2}$  poll. in diam.

10. **S. nitidus**, Thw.; *foliis* ovatis vel ovato-lanceolatis, acuminatis, basi rotundatis vel subcuneatis, utrinque nitidis, subtus reticulatis; *pedunculis* 1-floris.—*S. lucidus*, Thw. MSS. in Herb. Hook.—c.p. 3483.

HAB. Pasdoon Corle, at no great elevation.

*Arbor mediocris. Folia*  $2\frac{1}{2}$ – $3\frac{1}{2}$  poll. longa (acumine 3–6 lin. longo), 8–13 lin. lata, *venis primariis* circiter 6, *petiolo* 3–4 lin. longo.

Closely allied to but perfectly distinct from the preceding.

11. **S. ? Moonii**, Thw.; *foliis* anguste lanceolatis, basi cuneatis, superne lævibus, subtus ad costam venas primariasque, cum *petiolo* tereti, pilosis; *stipulis* longis, lineari-lanceolatis, rigidis, persistentibus; *pedunculis* 1-floris, in foliorum delapsorum axillis inter stipulas persistentes.—c.p. 1792.

HAB. Maturatte?, Moon.

*Folia* 6–10 poll. longa, 1–2 poll. lata, *petiolo* circiter 1 poll. longo. *Stipulae* 8–10 lin. longæ.

This interesting species is so unlike in general appearance those that precede it in this genus, that it was with some surprise I found, upon an analysis of an unopened flower, that it corresponded with them in the number, structure, and arrangement of the stamens, and in the external characters of the ovary and style. There are only two indifferent specimens in the herbarium, and one of these without flowers, collected by Mr. Moon many years ago, so that I have not materials for such a critical examination of the plant as would be desirable.

## 8. MONOPORANDRA, Thw.

Hook. Journ. of Bot. vol. vi. (1854) p. 69.

*Stamina* 5, *antheris* apice per laterum involuptionem tubulatis. *Ovarium* 2-loculare.

1. **M. cordifolia**, Thw. l.c.; *foliis* ovatis vel oblongis, longe acuminatis, basi cordatis vel rotundatis, superne costa et sæpissime *venis primariis* depressis, subtus *venis secundariis* parallelis; *paniculis* multifloris.—c.p. 2647.

HAB. Ambagamowa and Saffragam Districts, at an elevation of about 3000 feet:

*Arbor mediocris, ramosissima. Folia* 2– $4\frac{1}{2}$  poll. longa (acumine  $\frac{1}{2}$ –1 poll. longo), 1– $2\frac{1}{4}$  poll. lata, *petiolo* 4–11 lin. longo.

2. **M. elegans**, Thw. l.c.; *foliis* ovatis vel oblongis, longe acuminatis, basi rotundatis, superne ad costam depressis, *venis* prominentibus, subtus inter *venas primarias* reticulato-venosis; *pedunculis* paucifloris.—c.p. 371.

HAB. Saffragam District, at an elevation of about 2000 feet.

*Arbor mediocris, ramosissima. Folia*  $1\frac{1}{2}$ – $2\frac{1}{2}$  poll. longa (acumine 4–6 lin. longo), 6–10 lin. lata, *petiolo* 2–5 lin. longo.

3. **M. lancifolia**, Thw.; *foliis* lanceolatis, valde acuminatis, basi rotundatis, superne costa *venisque* prominentibus; *pedunculis* paucifloris.—c.p. 3412.

HAB. Hellessee, in the Pasdoon Corle, at no great elevation. April, 1855.

*Arbor parva, ramosissima. Folia*  $2\frac{1}{4}$ – $3\frac{1}{2}$  poll. longa (acumine 4–8 lin. longo), 7–11 lin. lata, *petiolo* 3–4 lin. longo.

### XXXIII. TERNSTROEMIACEÆ.

#### 1. CLEYERA, Thunb.

1. *C. gymnantha*, W. et A. Prod. i. p. 87.—C.P. 778 (779).

HAB. Common in forests of the Central Province, at an elevation of 4000 feet and upwards.

2. *C. emarginata*, Gard. in Calc. Journ. of Nat. Hist. vol. vii. p. 447.—C.P. 782.

HAB. Not uncommon near Adam's Peak and other more elevated parts of the island.

*Frutex glaber, 10–16-pedalis. Ramuli teretes. Folia coriacea, spathulata, apicem versus crenato-serrata, emarginata, subtus pallidiora, 1½–2½ poll. longa, 9–15 lin. lata. Pedunculi axillares, solitarii, 1-flori, circiter 8 lin. longi. Flores flavido-purpurascens, cernui, bracteolis 2 minutis suffulti. Petala sepalis duplo longiora, circiter 4 lin. longa. Stamina numerosa, glabra, filamentis brevibus, planis; antherarum introrsarum loculis linearibus, connectivo lato, filamento continuo, apice producto separatis. Ovarium glabrum, biloculare; loculis 1-ovulatis; stylo brevi, bifido; stigmatibus 2, dilatatis, lobulatis. Fructus conicus, 2-spermus.*

I have examined a great many specimens of this plant, and have invariably found two minute bracteolæ below each flower, and the loculi of the ovary one-ovuled. The leaves must be considered sessile, since the lamina is continued to their very base, though it forms but a narrow border to each side of the costa in the lower part.

#### 2. GORDONIA, Ellis.

1. *G. Zeylanica*, Wight, Illust. i. p. 99.

Var. *a. lanceolata*; ramulis pilosis, foliis lanceolatis vel oblongo-lanceolatis.—C.P. 788.

Var. *β. elliptica*; ramulis lævibus vel sublævibus, foliis latioribus.—*G. elliptica*, Gard. l. c. vol. vii. p. 448.—C.P. 2417.

HAB. Forests of the Central Province, at an elevation of 4000 to 7000 feet.

This species, according to the situation in which it occurs, varies very much in the size of the flowers and size and shape of the leaves. In exposed places where the soil is poor, the var. *α* is found, whilst in more sheltered localities and richer soil the plant is much more luxuriant, and puts on the form of var. *β*. In a series of specimens the two forms may be seen to pass imperceptibly into each other.

2. *G. speciosa*, Thw.—*Carria speciosa*, Gardn. l. c. vol. vii. p. 7.—C.P. 2403.

HAB. Rather uncommon; in damp forests of the Central Province, at an elevation of 5000 feet and upwards.

*Arbor magna, 40–50-pedalis. Ramuli teretes, glabri. Folia integra, sessilia, coriacea, glabra, elliptica vel ovata, retusa, 3–4½ poll. longa, 1½–2¾ poll. lata. Flores magni, axillares, sessiles vel brevi-pedunculati; petalis cordatis, sanguineo-purpureis, 1½–2 poll. longis.*

I regret being obliged to include in the genus *Gordonia* this fine species, which Dr. Gardner dedicated, under the generic name of *Carria*, to the late excellent Sir Wm. Carr; but I find that the characters upon which he based his generic distinctive diagnosis are common to the other Ceylon species of *Gordonia*. The structure of the anthers is the same in both, and the analysis of the seed of *Gordonia Zeylanica*

corresponds with that which he gives of *Carria speciosa*, the cotyledons being perfectly flat, and not plaited.

### 3. SAROSANTHERA, Korthals.

1. **S. lasiopetala**, Thw.—*Cleyera lasiopetala*, Wight, Ill. i. p. 99. *Eurya lasiopetala*, Gardner, l. c. vol. vii. p. 446.—c.p. 775 (776, 783).

HAB. Common in the forests of the Central Province, at an elevation of 6000 feet and upwards.

*Arbor* mediocris. *Ramuli* teretes, minute strigosi. *Folia* oblongo-lanceolata, obtusa vel subacuminata, retusa, margine revoluta, minute denticulata, superne glabra, subtus minute et parce strigosa,  $2\frac{1}{2}$ –4 poll. longa,  $\frac{3}{4}$ –1 poll. lata. *Pedunculi* axillares, solitarii, 1-flori, cernui, circiter 8 lin. longi, apice incrassati. *Flores* hermaphroditi, 6–8 lin. in diam., externe sericeo-strigosi; *petalis* albidis. *Fructus* globosus, carnosus nec pulposus, circiter 5 lin. in diam.

This description is taken principally from the one published by Dr. Gardner.

### 4. EURYA, Thunb.

1. **E. Japonica**, Thunb. Fl. Jap. p. 191, t. 25.

Var. *a*. *Thunbergii*; ramulis teretibus, glabris.—*E. Japonica*, DC. Prod. i. 525. *E. elliptica*, Gardner, l. c. vol. vii. p. 443.—c.p. 777 (787).

Var. *β*. *acuminata*; ramulis teretibus, pilosis.—*E. acuminata*, DC. Prod. i. 525. *E. membranacea*, Gard. l. c. vol. vii. p. 444.—c.p. 1078 (785, 786).

Var. *γ*. *Chinensis*; ramulis angulosis, pilosis.—*E. Chinensis*, R. Br. Abel's Voy. App. p. 379; DC. l. c.; Bentham in Hook. Kew Journ. of Bot. vol. iii. (1851) p. 307. *E. Ceylanica*, Wight, Ill. i. 98; Gardner, l. c. vol. vii. p. 444.—c.p. 2600.

Var. *δ*. *parvifolia*; ramulis angulosis, pilosis; *foliis* minoribus, pallidis.—*E. parvifolia*, Gardner, l. c. vol. vii. p. 445.—c.p. 784.

HAB. Vars. *a*, *β*, and *δ* in the more elevated parts of the island, up to 8000 feet; *δ* in exposed situations; var. *β* from a little above the sea-level, up to an elevation of 5000 feet, very abundant. Nom. vulg. "Neyadasse-gass."

It is only after a very careful and long-continued study of these plants that I have ventured to unite them as forms of one variable species. I have been quite unable to discover, in those parts where other botanists have thought they existed, any definite difference of structure sufficient to constitute a distinction of species. The number of the stamens and apparent length of the style vary in the same variety just as much as between the so-called species, and it seems to me quite impossible to draw a clear line of distinction between the several varieties, since specimens are to be met with combining them all by intermediate links.

*Cochlospermum Gossypium*, DC., is not found in a wild state in the island, but only, as Dr. Gardner has remarked, in the vicinity of temples.

## XXXIV. OLACACEÆ.

### TRIBE OPILEÆ.

#### 1. OPILIA, Roxb.

1. **O. amentacea**, Roxb. Corom. Pl. t. 158; Wight, Illust. p. 103, t. 40.—c.p. 2916 (1147).

HAB. Hot, drier parts of the island, at no great elevation.



TRIBE *OLACEÆ*.2. *OLAX*, Lin.

1. *O. Zeylanica*, Lin. (W. et A. Prod. i. 88, cum syn.)—c.p. 1214.  
 HAB. Southern part of the island; abundant. Nom. vulg. "Malla."  
 The Cinghalese eat the leaves of this plant in their curries.
2. *O. scandens*, Roxb. (W. et A., l. c. p. 89, cum syn.)—c.p. 1216.  
 HAB. Hot, drier parts of the island.
3. *O. Wightiana*, Wall. (W. et A., l. c. cum syn.)—c.p. 1215.  
 HAB. Warmer parts of the island; not uncommon.

3. *STROMBOSIA*, Blume.

1. *S. Javanica*, Bl. Bijdr. p. 1154; Mus. Bot. Lugd. Bat. p. 251, t. 47.—*S. Ceylanica*, Gardner, in Calc. Journ. of Nat. Hist. vol. vi. p. 350.—c.p. 1237.  
 HAB. In forests of the Central Province, up to an elevation of 3000 feet.

Blume's figure and description leave no room for doubt that this is his plant.

TRIBE *ICACINEÆ*.4. *BURSINOPETALUM*, Wight, Icones, 956.

1. *B. arboreum*, Wight, l. c. Var. *macrophyllum*.—c.p. 637, 2440.  
 HAB. Forests of the Central Province, at an elevation of 4000 to 7000 feet.

*Arbor ingens. Folia* ovato-oblonga, acuminata, 4-9 poll. longa,  $1\frac{3}{4}$ -4 poll. lata, *petiolo* 1-2 poll. longo.

I can discover no character to distinguish this from Dr. Wight's plant but the larger size of the leaves, and do not venture to separate it on that account only. A comparison with authentic specimens of Dr. Wight's *B. arboreum* may possibly prove this to be distinct from it, and in that case it might be called *B. macrophyllum*.

2. *B. tetrandrum*, Wight, MSS.; *foliis* oblongis vel ovato-oblongis, *floribus* tetrameris.

Var. *a*; *foliis* longe acuminatis.—c.p. 2441.

Var. *β*; *foliis* minoribus, apice rotundatis vel abrupte et brev-acuminatis.—c.p. 2542.

*Arbor ingens. Folia* 2-4 poll. longa,  $\frac{3}{4}$ -2 poll. lata, *petiolo* 4-8 lin. longo.

Distinguished most readily from the preceding by its smaller, tetramerous flowers; the leaves, also, are not nearly so large, and the ripe fruit is more cylindrical in form, and smaller.

5. *APODYTES*, E. Mey.

1. *A. Gardneriana*, Miers, Ann. and Mag. of Nat. Hist. ser. 2. vol. ix. p. 389.—c.p. 204 (495, 1218, 1219).

HAB. Forests of the Central Province, at an elevation of from 5000 to 7000 feet; not uncommon.

*Arbor mediocris. Folia* 2-4 poll. longa,  $\frac{3}{4}$ -2 poll. lata, *petiolo* 3-5 lin. longo.

I have not had an opportunity of comparing this with *A. Benthamiana*, Wight, Icon. t. 1153, which is described as having larger flowers than those of this species. The Ceylon plant invariably turns blackish in drying.

## 6. MAPPIA, Jacq.

1. *M. foetida*, Miers, l. c. p. 395.—*M. ovata*, Miers, l. c. p. 396. *Stemonurus? foetidus*, Wight, Icon. t. 955; Spicil. Neilgh. t. 23.—C.P. 492.

Var. *a. Gardneriana*; *alabastris* rotundatis.—*M. Gardneriana*, Miers, l. c. p. 396.

Var. *β. Championiana*; *foliis* membranaceis, *alabastris* oblongis.—*M. Championiana*, Miers, l. c. 397.

HAB. One of the commonest plants in the island, and to be met with from the level of the sea, up to an elevation of 7000 feet. Nom. vulg. "Gandapaana-gass."

Owing to its great range in elevation, this plant varies a great deal in appearance and in the texture of its growing parts, these being of a much firmer consistence on the hills than in the neighbourhood of the sea, and more palpably so in dried than in fresh specimens.

## 7. STEMONURUS, Blume.

*Stemonurus* (in parte), Miers. *Urandra*, Thw.

1. *S. apicalis*, Thw.—*Urandra apicalis*, Thw. in Hook. Kew Journ. of Bot. vol. vii. (1855) p. 211.—C.P. 2569.

HAB. Damp forests, at an elevation of from 1000 to 2000 feet. Nom. vulg. "Oorookannoo-gass."

*Arbor* ingens. *Ramuli* teretes, glabri. *Folia* integra, coriacea, oblonga vel obovato-oblonga, abrupte brevi-acuminata, basi angustata, subtus pallidiora, 4–6 poll. longa, 2–3 poll. lata, *petiolo* 5–8 lin. longo. *Pedunculi* axillares, solitarii vel bini, *petiolo* breviores vel subæquilongi. *Capitula* 7–12-flora. *Flores* hermaphroditæ, *petalis* viridibus, basin versus purpurascens, circiter 2 lin. longis. *Drupa* oblonga, subacuta, 15 lin. longa, 8 lin. lata, viridi-purpurascens, dimidio apicali albido; *putamine* ligneo, cancellato, externe fasciculato-fibroso. *Embryo* in axi albuminis copiosi, carnosi; *radicula* supera, cylindrica, *cotyledonibus* parvis cordatis æquilonga; *testa* pericarpio consolidata.

When describing this plant, two years ago, under the name of *Urandra apicalis*, I had not the opportunity of referring to Blume's Museum Bot. Ludg. Batav., and took it for granted that Mr. Miers was correct in determining Dr. Wallich's genus *Gomphandra* to be identical with *Stemonurus* of Blume. Now, however, I am satisfied that that learned botanist was in error, and that to *Platea* of Blume is to be referred the genus *Gomphandra*. Blume's definition of *Stemonurus* as "drupa baccata, nucleo fibroso," does not apply to *Gomphandra*, though it does to the species figured in the Mus. Bot. Lugd. Bat., and to the plant now under description. By a strange oversight, Mr. Miers has read my account of the structure of the seed of *Gomphandra polymorpha*, Wight (Hooker's Journal of Bot. vol. vii. p. 211), as if it were that of my *Urandra apicalis*, and has thus been under the erroneous impression (Linnaean Trans. vol. xxii. p. 97 *et seq.*) that the structure is the same in the two plants. It is very different, for in *Stemonurus apicalis* there is no appearance of the longitudinal division of the albumen into two equal parts; the testa is firmly adherent to the pericarp, and the putamen is very peculiar in character, being ligneous, with several longitudinal cavities filled with soft cellular matter, and externally, when the flesh of the

drupe has decomposed or been removed, exhibiting a fasciculate-fibrous structure, which in appearance is not very unlike an arillus, but of course essentially very different from one. Mr. Miers is right in saying that *Urandra apicalis* is a *Stemonurus*, and I have now called it so; but it is not so for the reasons he gives, as has been shown above.

## 8. PLATEA, Blume.

*Gomphandra*, Wallich, Wight. *Stemonurus* (in parte), Miers.

1. **P. axillaris**, Thw.—*Gomphandra axillaris*, Wall. Cat. n. 3718. *Lasianthera? tetrandra*, Wall. in Roxb. Fl. Ind. ii. p. 328. *Gomphandra polymorpha*, Wight, Illust. i. p. 103; Icones, t. 954. *Stemonurus axillaris*, Miers, in Ann. of Nat. Hist. ser. 2. vol. x. p. 41. *St. polymorphus*, Miers, l. c. p. 37. *St. Ceylanicus*, Miers, l. c. p. 42. *St. Heyneanus*, Miers, l. c. p. 40.—c.p. 251 (762).

HAB. From the level of the sea, up to an elevation of 4000 feet; very abundant.

After reading the description of Dr. Wallich's plant in 'Flora Indica,' cited above, it would seem almost impossible to doubt that this is the species intended by him. It is a very protean one as regards the shape and size of the leaves, and has consequently been overwhelmed with synonyms. I have not the least doubt that those quoted above, if not more, are referable to it in one or other of its forms.

2. **P. coriacea**, Thw.—*Gomphandra coriacea*, Wight, Ill. i. p. 103. *G. polymorpha*, ♂, Wight, Icon. t. 953. *Stemonurus Gardneri*, Miers, l. c. p. 38. *St. Walkeri*, Miers, l. c. p. 43. *Platea Wightiana*, Miers, l. c. p. 111.—c.p. 375.

HAB. Damp forests of the Central Province, at an elevation of 3000 to 6000 feet; not uncommon.

Equally protean, or even more so than the last species, in the size and shape of its leaves, which vary from narrow-lanceolate to orbicular, with a long, narrow acumination. The hairs upon the filaments are very short, and may easily have been overlooked by Dr. Wight's draughtsman; and the petals and stamens of the fertile plant are so very caducous, that it is not surprising to find Blume describing the flowers as having none. The peduncle is at first axillary, but a leaf being most commonly produced upon it at about its middle, its upper part of course appears opposite to this new leaf, and more strikingly so when a leaf-bud has become developed in the axil of the latter.

This species differs most obviously from the last in its coriaceous, opaque leaves, and longer, dichotomous peduncles, and in these latter appearing most commonly opposite to a leaf. *Gomphandra coriacea* of Dr. Wight's 'Illustrations' is, no doubt, the tetramerous form of this.

## XXXV. AURANTIACEÆ.

### 1. ATALANTIA, Corr.

1. **A. monophylla**, DC. (W. et A. Prod. i. 91; Wight, Illust. i. 108.) —c.p. 497 (1198).

HAB. Kornegalle and towards the north of the island, at no great elevation; not uncommon.

## 2. LIMONIA, Lin.

1. *L. alata*, Herb. Madr. (W. et A., l. c. p. 92.)—c.p. 1191.

HAB. Hot, drier parts of the island; common. Nom. vulg. "Toompat-koo-roondoo-gass."

2. *L. Missionis*, Wall. (W. et A., l. c.)—c.p. 1194.

HAB. Hotter parts of the island. Nom. vulg. "Pambooroo-gass."

## 3. GLYCOSMIS, Corr.

1. *G. arborea*, DC. (W. et A., l. c. cum syn.)—c.p. 180.

HAB. Abundant in the hot, drier parts of the island. Nom. vulg. "Dodang-paana-gass."

*Filamenta* dilatata, supra medium latissima, ad apicem angustata. *Ovarium* glandulosum, breve, gynophoro sphaeroidali angustius. *Stylus* depressus, brevis, stigmatibus discoideo distincto æquilatus.

This must certainly be the *Limonia arborea* of Roxburgh (Cor. Plant. t. 85), though the analysis of the flower accompanying his figure expresses far from correctly the character of the species.

2. *G. pentaphylla*, DC. (W. et A., l. c. 93, cum syn.)—c.p. 1192.

HAB. Very common; up to an elevation of 3000 feet.

*Filamenta* ut in specie præcedenti. *Pistillum*, una cum gynophoro mediocri, subcylindricum, truncatum. *Fructus* pulposus.

This is a very variable species as regards the shape and size of its leaves, but may always be distinguished from the preceding by the much smaller panicles and different shape of the pistil and gynophore.

3. *G. bilocularis*, Thw.; *foliis* pinnatis; *foliolis* alternis, brevipetiolulatis, trapezoideo-lanceolatis, apice obtusis; *paniculis* axillaribus, brevissimis, paucifloris; *ovario* biloculari.—c.p. 3362 (olim cum c.p. 1192 mixta).

HAB. Maturatte District, at no great elevation.

*Frutex*. *Foliola*  $1\frac{3}{4}$ – $2\frac{1}{2}$  poll. longa, 5–8 lin. lata, subtus pallidiora. *Flores* subsessiles. *Calyx* 5-fidus, segmentis triangularibus. *Petala* 5, albida, minute rufopunctata, oblonga. *Filamenta* alterna breviora, multum et irregulariter dilatata, interne concava. *Pistillum*, una cum gynophoro, conicum, truncatum.

This bears a considerable resemblance to some forms of the preceding species, but its two-celled ovary, besides other differences, proves it to be quite distinct. Its technical characters would perhaps bring it within the genus *Sclerostylis*, but its nearest affinity is certainly with the species of *Glycosmis*.

## 4. MURRAYA, Kœnig.

1. *M. exotica*, Lin.

Var. *a. buxifolia*; *ramulis* confertis, glabris vel tomentosis; *foliis* subcoriaceis; *calycis* segmentis triangularibus.—*M. exotica*, W. et A., l. c. 94, cum syn.—c.p. 1200.

Var. *β. paniculata*; *foliis* majoribus, membranaceis, venis subtus prominentibus; *calycis* segmentis oblongis.—*M. paniculata*, W. et A., l. c. cum syn.—c.p. 1199.

HAB. Var. *a* in gardens only. Var. *β* common in the warmer parts of the island, and up to an elevation of 3000 feet. Nom. vulg. "Attairoya-gass."

With fresh specimens of these two forms before me, I have been unable to discover anything like a good specific difference between them.

### 5. SCLEROSTYLIS, Blume.

1. **S. Ceylanica**, Wight, Ill. i. p. 109.—*S. Arnottiana*, Wight, l. c. *Rissoa Ceylanica*, Arn. Pug. p. 6 (324).—C.P. 1196.

HAB. Warmer parts of the island, not uncommon. Nom. vulg. "Yucca-naara-gass."

A very variable species in the size and shape of its leaves, and the number of flowers in the axils. I have always found the flowers tetramerous, and two ovules in each loculus of the ovary.

2. **S. rotundifolia**, Thw.; *inermis?*, *foliis* rotundis vel ovato-rotundis, retusis, coriaceis; *racemis* ad apices ramulorum confertis.—C.P. 3295.

HAB. Maturatte, at an elevation of 4000 feet and upwards; not common.

*Arbor* parva, ramosissima, glabra. *Folia*  $\frac{3}{4}$ – $1\frac{3}{4}$  poll. longa,  $\frac{1}{2}$ – $1\frac{1}{2}$  poll. lata, *petiolo* 1–2 lin. longo. *Racemi* axillares et terminales, solitarii vel 2–3-fasciculati, 5–10-flori, foliis æquilongi. *Flores* tetrameri. *Calyx* profunde 4-fidus, lobis rotundatis. *Ovarium* oblongum, brevi-stipitatum, in *stylum* cylindricum apice compresso-dilatatum margine stigmatosum æquilongum attenuatum; *loculis* 2, biovulatis.

Closely allied to the preceding species, but distinguished at once by the absence (apparently) of spines, and the different shape of the leaves. The calyx is deeply four-lobed, whereas in *S. Ceylanica* it is scarcely more than undulated; the style is also more slender in the present species, and the ovary more decidedly stalked. In both species the latter organ is surrounded at its base by a free, glandular annulus.

### 6. BERGERA, Kœnig.

1. **B. Kœnigii**, Lin. (W. et A. Prod. i. 94, cum syn.; Wight, Icones, t. 13.)—C.P. 1203, 2547.

HAB. Hot, drier parts of the island; not uncommon. Nom. vulg. "Karapinchee-gass."

The Cinghalese use the leaves of this plant as a seasoning for their curries.

2. **B. nitida**, Thw.; *foliolis* 5–14, oblongis, obtuse acuminatis vel retusis, valde obliquis, obscure crenatis, glabris, nitidis; *ovarii* loculis 2, biovulatis.—C.P. 2421.

HAB. Warmer parts of the island; not very uncommon. Nom. vulg. "Meegong-karapinchee-gass."

*Arbor* mediocris. *Ramuli* minute pilosi. *Foliola*  $1\frac{1}{4}$ –3 poll. longa,  $\frac{3}{4}$ – $1\frac{1}{4}$  poll. lata, *petiolulo* puberulo, 1–2 lin. longo. *Paniculi* terminales, puberuli. *Flores* parvi, albidi, pentameri. *Staminum filamenta* medio dilatata; *antheræ* cordato-rotundatæ. *Ovarium* oblongum; *stylo* brevi, cylindrico, deciduo; *stigmatæ* capitato. *Fructus* sphaericus, pulposus, flavus, 1-spermus, 4–5 lin. in diam.

This species has evidently a near affinity with *Piptostylis Indica*, Dalzell (Hook. Journ. of Bot. vol. iii. p. 33. t. 2), the style being wholly deciduous as in that species, and the loculi of the ovary always two-ovuled; but it differs from it in its very oblique leaves, smaller flowers, capitate stigma, and two-celled ovary. It may be considered doubtful, perhaps, whether it is desirable to retain the genus *Piptostylis* as distinct from *Bergera*. I am able to confirm Mr. Dalzell's observation that the loculi of the ovarium of *B. Kœnigii* are frequently two-ovuled.

### 7. MICROMELUM, Blume.

1. **M. pubescens**, Blume, Bijl. p. 138?—C.P. 188.

HAB. Very common in the hotter parts of the island, up to an elevation of 2000 feet. Nom. vulg. "Wal-karapinchee-gass."

This is a very variable plant in the size and shape of its leaves, which are always very oblique at the base. The small-leaved state bears very considerable resemblance to *Bergera Koenigii*. There seems to me to be scarcely sufficient grounds for excluding the present species from the genus *Cookia*.

### 8. CLAUSENA, Burm.

1. **C. Willdenovii**, W. et A. Prod. i. 96, cum syn.; Wight, Icones, t. 339.—C.P. 1204.

HAB. Hot, drier parts of the island.

*Foliorum juniorum* rachis, cum foliolorum petiolulis, costulis marginibusque superne puberula.

2. **C. pubescens**, W. et A., l. c.—C.P. 2546.

HAB. Near Hanguranketty, at an elevation of about 3000 feet; not common.

I suspect that this and the preceding will prove to be but varieties of one species, for I can discover no character of any importance to distinguish them except the dense pubescence upon the ramuli and under surface of the leaves of the present one.

### 9. PARAMIGNYA, Wight.

1. **P. monophylla**, Wight, Illust. p. 108. t. 42.—C.P. 1201, 1202.

HAB. Hot, drier parts of the island, up to an elevation of 2000 feet. Nom. vulg. "Wallangereya."

### 10. Gen. Nov. ARTHROMISCHUS, Thw.

*Calyx* brevis, 4-dentatus. *Corollæ* petala 4, oblonga. *Stamina* 8, æqualia; *filamentis* medio dilatatis; *antheris* oblongis. *Ovarium* gynophoro conspicuo suffultum, 4-loculare; *loculis* 1-ovulatis. *Ovula* pendula. *Stylus* longus, cylindricus. *Stigma* dilatatum. *Fructus* 1-4-spermus.

1. **A. armatus**, Thw.—C.P. 1197, 3115.

HAB. Hotter parts of the island, at no great elevation; not common.

*Frutex* scandens, spinis rectis vel apice curvatis armatus. *Ramuli* glabri vel minute puberuli. *Folia* 1-foliolata, *petiolo* 2-4 lin. longo; *foliolo* ovato-vel oblongo-acuminato, basi rotundato,  $1\frac{1}{2}$ - $3\frac{1}{2}$  poll. longo, 7-15 lin. lato; *petiolulo*  $1\frac{1}{2}$  lin. longo. *Fasciculi* axillares, 1-8-flori, *pedicellis* 3-4 lin. longis. *Petala* 2 lin. longa. *Ovarium* subsphæricum, gynophoro rotundato æquilongo subæquilato suffultum. *Fructus* pulposus, rotundatus, flavescens, 4-8 lin. in diam.

This plant differs from *Luvunga* in the greater number of cells to the ovary, and these being one-ovuled; from *Paramignya* its jointed leaf-stalk, small calyx, and one-ovuled loculi distinguish it, whilst from *Atalantia* it is separated by its scandent habit, free filaments, and stalked ovary.

### 11. LUVUNGA, Ham.

1. **L. scandens**, Ham. (Wight, Illust. i. 108.)—*Limonia scandens*, Roxb. Fl. Ind. ii. 380.—C.P. 1195.

HAB. Central Province, at an elevation of 3000 to 5000 feet.

In all the specimens of this plant I have examined, the filaments were free to their very base, even in the unexpanded flowers, and the species appears to be somewhat dioecious, as in some plants the ovaries of all the flowers are entirely without ovules.

If Roxburgh did not make a mistake in his description of this plant, it will be necessary to call the Ceylon *Luvunga* in this 'Enumeration,' *L. eleutherandra*, Dalzell, or what is the *L. Tavoyana*, Wall., referred to by Wight in the *Illust.* i. p. 108?

~~///~~ **L. eleutherandra**, Dalz. Hook. Journ. of Bot. ii. (1850) p. 258; Walp. Ann. ii. p. 184.—c.p. 1195.

HAB. Central Province, at an elevation of 3000 to 5000 feet.

This species would appear to be dioecious, as in some plants the ovaries of all the flowers are entirely without ovules. It varies much in the shape and size of the leaves.

## 12. FERONIA, Corr.

1. **F. Elephantum**, Corr. (W. et A. Prod. i. 96, cum syn.; Wight, *Icones*, t. 15.)—c.p. 1205.

HAB. Hot, drier parts of the island, very abundant. Nom. vulg. "Diwool-gass."

## XXXVI. HYPERICACEÆ.

### 1. HYPERICUM, Lin.

1. **H. Mysorense**, Herb. Mad. (W. et A. Prod. i. 99.)—*Norysca Mysorensis*, Wight, *Icones*, t. 56.—c.p. 48.

HAB. Central Province, at an elevation of 6000 to 8000 feet; not uncommon.

2. **H. Japonicum**, Thunb. (W. et A., l. c. cum syn.)—c.p. 2420.

HAB. Abundant in paddy-fields and other wet places throughout the island.

## XXXVII. GUTTIFERÆ.

### 1. GARCINIA, Lin.

1. **G. Cambogia**, Desrous. (DC. Prod. i. 561, cum syn.; Wight, *Illust.* i. 125, cum syn.)—*Cambogia Gutta*, Lin. (in parte), et forsan *Garc. Roxburghii*, Wight, l. c. cum syn.—c.p. 1172.

Var.  $\alpha$ ; fructu maturo flavo.

Var.  $\beta$ ; fructu maturo rubro.

HAB. Not uncommon up to an elevation of 1500 feet. Nom. vulg. "Goraka-gass."

*Florum fasciculi*:—MASC. 3-8, pedicellati, *pedicellis* 7-10 lin. longis, apice parce incrassatis. FÆM. 1-2, pedicellati, *pedicellis* 2-5 lin. longis. *Fructus* 2½-3 poll. in diam.

There appears to be in Ceylon but one species of *Garcinia* with sulcated fruit, but in all the figures and descriptions of it I have seen, no notice is taken of a very marked character it has, namely, that the sulci or furrows are not continued to the apex, but terminate at some distance from it, the apex itself being smooth and depressed, and somewhat mamillæform. The gum which exudes from the tree

is semitransparent, very adhesive, and is insoluble in water, and therefore quite unsuitable for the purposes of the draughtsman. The Cinghalese dry the acid rinds of the ripe fruit, both the yellow and red varieties, and use them as a condiment for eating with their curries.

2. **G. echinocarpa**, Thw. (Hook. Journ. of Bot. vol. vi. (1854) p. 71.)

Var. *a*; *foliis* oblongis vel lanceolatis, acuminatis.—c.p. 2445.

Var. *β*; *foliis* valde coriaceis, obovatis vel oblongis, retusis vel acumine brevi obtuso; *floribus* majoribus.—c.p. 335.

HAB. Var. *a*, in the forests between Ratnapoora and Galle, at no great elevation. Var. *β*, in the Central Province, at an elevation of 4000 to 6000 feet; common. Nom. vulg. "Madol-gass."

*Flores* terminales, sessiles: MASC. numerosi, aggregati; FÆM. solitarii. *Ovarium* squamis carnosis numerosis imbricatis tectum. *Fructus* subsphæricus, echinulato-muricatus.

A thick oil is extracted from the seeds, which is used by the Cinghalese for burning in their lamps, but it gives a very indifferent light.

3. **G. Morella**, Desrous.—*Hebradendron cambogioides*, Graham, in Hook. Comp. to Bot. Mag. vol. ii. p. 199. t. 27, cum syn. *Garcinia Gutta*, Wight, Illust. i. 126. *Cambogia Gutta*, Linn. Fl. Zeyl. p. 87, in parte.—c.p. 322.

HAB. Warmer parts of the island, up to an elevation of 2000 feet. Nom. vulg. "Gokatoo-gass," "Kana-goraka-gass."

FÆM. *Ovarium* glabrum, apice stigmatè sessili imbricatim lobato tectum; basi *staminibus sterilibus* circiter 12, 1-serialibus, basi cohærentibus cinctum; *antheris* effatis, subquadratis, apice emarginatis.

This is the only species growing in Ceylon from which gamboge is obtainable, and as the tree is not uncommon, the pigment might be collected in considerable quantities.

## 2. TERPNOPHYLLUM, Thw.

Hook. Journ. of Bot. vol. vi. (1854) p. 70.

? *Discostigma*, Hassk. (Endl. Gen. Plant. Suppl. iii. p. 95.)

*Flores* dioici. *Calyx* 4-phyllus, deciduus. MASC. *Stamina* plurima in discum mamillatum 2- vel 4-lobatum subtus petalis adhærentem coalita; mamillis singulis, *antheras* 2-4, loculis horizontalibus subimmersis gerentibus; *ovarii rudimento* minuto.

FÆM. *Ovarium* biloculare, *staminibus sterilibus* paucis squamæformibus cinctum.

1. **T. Zeylanicum**, Thw. l. c. t. 2.

Var. *a*. *lanceolatum*; *foliis* lanceolatis; *disco stamineo* 2-lobato, *lobis* bifidis, medio sulcatis.—c.p. 2695.

Var. *β*. *acuminatum*; *foliis* lanceolatis, obtuse rostrato-acuminatis; *disco stamineo* 4-lobato.—c.p. 3400.

HAB. Var. *a*, Central Province, at an elevation of 2000 to 4000 feet. Var. *β*, in forests between Ratnapoora and Galle, at no great elevation.

The *Garcinia Merguensis*, Wight, Icones, t. 116, will probably prove to be a species of this genus.

## 3. XANTHOCHYMUS, Roxb.

1. **X. ovalifolius**, Roxb. Fl. Ind. ii. p. 632; W. et A. Prod. i. 102 (excl. syn. a Burm. Fl. Ind. et Fl. Zeyl.).—? *X. spicatus*, W. et A., l. c.



Var.  $\alpha$ ; *foliis* ovatis, apice rotundatis vel retusis.—C.P. 1173.

Var.  $\beta$ ; *foliis* lanceolatis vel ovato-lanceolatis, plus minus acuminatis.—C.P. 731 (743, 755).

HAB. Var.  $\alpha$ , hot, drier parts of the island. Var.  $\beta$ , Central Province, up to an elevation of 3000 feet; abundant. Nom. vulg. "Ella-gokatoo-gass."

A most variable species, both as to the shape of the leaves and length of the peduncles and pedicels. Although the extreme forms of vars.  $\alpha$  and  $\beta$  look very dissimilar, I am unable to find any constant character by which to separate them. A tenacious gummy matter exudes from the trunk, but it is of no value.

#### 4. MESUA, Linn.

1. **M. ferrea**, Linn. Sp. 734; DC. Prod. i. 562; Wight, Ill. i. 127; Icones, t. 118.—*M. Nagaha*, Gard. in Calc. Journ. of Nat. Hist. vol. viii. p. 4. *Arbor Naghas*, Burm. Zeyl. p. 25. *Mesua foliis lanceolatis, etc.*, Fl. Zeyl. n. 203. *Nagassarium*, Rumph. Amb. vii. 3. t. 2.—C.P. 602.

Var.  $\alpha$ ; *foliis* lanceolatis, acuminatis.

Var.  $\beta$ ; *foliis* lineari-lanceolatis.

HAB. Vars.  $\alpha$  and  $\beta$ , common about Galagama, especially the latter. Var.  $\alpha$  occurs very commonly in the neighbourhood of temples as one of the favourite trees planted for ornament. Nom. vulg. "Nā-gass."

Dr. Gardner has given, in the work quoted above, a very full description of the ordinary form of this species. The natives express an oil from the seeds for burning. The wood is very hard and durable.

2. **M. speciosa**, Chois. in DC. Prodr. i. 562; Rheede, Hort. Malab. iii. 63. t. 53. ? Wight, Icones, t. 961.—C.P. 2675.

Var.  $\beta$ ; *foliis* multo minoribus.—C.P. 603.

HAB. Banks of streams, in the Ratnapoora District. Nom. vulg. "Deya-nā-gass."

The figure in Hort. Mal., quoted above, and from which M. Choisy appears to have derived his character of the species, is a very fair representation of the usual form of this plant, which differs from the preceding in its much larger leaves, most commonly rounded at the base, its sessile or subsessile flowers, and larger fruit.

3. **M. Coromandelina**, Wight, Icones, t. 117; Illust. i. 127.—C.P. 3404.

HAB. Forests between Galle and Ratnapoora, at no great elevation.

Dr. Wight's figure in the 'Icones' well represents this plant, which is distinguished from *M. ferrea* by its more suddenly acuminated leaves, their base more rounded, and in these being without the white coating on the under side, so conspicuous in the other two species, and in its much smaller flowers on longer pedicels.

#### 5. KAYEA, Wall.

*Capsula* rotundata, coriacea, irregulariter fissa, 1-4-sperma, *calyce* multum aucto coriaceo inclusa. *Semina cotyledonibus* distinctis, *car-nosis*; *testa* membranacea, colorata.

1. **K. stylosa**, Thw; *foliis* ovato-lanceolatis, acuminatis.—C.P. 2708.

HAB. South of the island, at no great elevation. Nom. vulg. "Soovanda-gass."

*Arbor magna. Ramuli teretes, glabri, rufescentes. Folia 2-3 poll. longa, 7-10 lin. lata, subtus minutissime reticulata, petiolo 2-3 lin. longo. Stylus staminibus numerosis multum longior.*

This differs from *Kayea floribunda* of Wallich's Pl. Asiat. Rarior. vol. iii. p. 5. t. 210, in its much smaller and differently-shaped leaves, and long, protruded style. The flowers are very fragrant. The timber is useful for building purposes.

## 6. CALOPHYLLUM, Linn.

Nom. vulg. "Keena-gass," "Walloo-keena-gass."

### Sect. 1. *Sepala 4. Petala 8.*

1. **C. Walkerii**, Wight, Illust. i. 128. t. 45.—C.P. 1170.

HAB. Newera Ellia, Adam's Peak, and other of the most elevated parts of the island.

An oil is extracted from the seeds of this and the other species of the genus, which is used for burning.

### Sect. 2. *Sepala 4. Petala 4.*

† *Racemi folia æquantés vel longiores.*

2. **C. decipiens**, Wight, Illust. i. 128; Icones, t. 106.—C.P. 3403.

Var. *a*; *foliis* obovato-oblongis, basi rotundatis.

Var. *β*; *foliis* cordato-orbiculatis.

HAB. Var. *a*, Ambagamowa District. Var. *β*, Hinidoon Corle, at an elevation of 1000 to 2000 feet.

The leaves of this plant vary much in shape, and the foliage often strongly resembles that of the preceding. In very young plants (C.P. 3401) of what I believe to be this species, the leaves are lanceolate or cordate-lanceolate.

3. **C. trapezifolium**, Thw.; *foliis* coriaceis, parvis, ovato-trapezoides, obtuse et breve acuminatis vel retusis; *racemis* folio multum longioribus; *floribus* majusculis.—C.P. 2446.

HAB. Hunasgiri District, in the Central Province, at an elevation of 4000 to 5000 feet.

*Arbor magna. Ramuli glabri. Folia 1½-2 poll. longa, 7-11 lin. lata, petiolo 3-4 lin. longo. Fructus subsphæricus, 8 lin. longus, 7 lin. latus.*

4. **C. cuneifolium**, Thw.; *foliis* parvis, cuneato-spathulatis, obtuse brevi-acuminatis vel retusis, coriaceis.—C.P. 2917.

HAB. Madamahanevewa, at an elevation of 3000 to 4000 feet.

*Arbor magna. Ramuli glabri. Folia 1¼-1¾ poll. longa, 5-9 lin. lata, petiolo 3-4 lin. longo. Flores mediocres. Fructus sphæricus, 8 lin. in diam.*

†† *Racemi foliis breviores.*

5. **C. Inophyllum**, Linn. (W. et A. Prod. i. 103; Wight, Illust. i. 128; Icones, t. 77.)—C.P. 2764.

HAB. Hotter parts of the island, up to an elevation of 2000 feet. Nom. vulg. "Domba-gass."

6. **C. tomentosum**, Wight, Illust. i. 128; Icones, t. 110.—C.P. 1171.

HAB. Central Province, at an elevation of 3000 to 5000 feet; common.

The timber of this abundant tree is valued for building purposes, and the seeds are collected in considerable quantities for the sake of the oil they contain, which is called "Keena-tel."

7. **C. bracteatum**, Thw.; *foliis* lanceolatis, acuminatis, a medio

versus basin subrotundatam sensim angustatis; *racemis* conspicue bracteatis, tomentosus.—C.P. 2674.

HAB. Saffragam District, by the sides of streams, at no great elevation.

*Arbor* ingens. *Ramuli* quadranguli, penduli, tomento ferrugineo caduco vestiti. *Folia*  $3\frac{1}{2}$ –8 poll. longa,  $1-2\frac{1}{4}$  lata; *petiolo* 2–4 lin. longo; *juniora* tomento fugaci pilosa. *Racemi* foliis multo breviores, 4–8-flori. *Pedicelli* basi 1-bracteati, *bracteis* oblongis, tomentosus, 2–3 lin. longis. *Flores* 4 lin. in expans. *Fructus* immaturus oblongus.

The bracteated racemes, much smaller flowers, and oblong fruit, abundantly distinguish this from the preceding species.

### Sect. 3. *Sepala* 4. *Petala* 0.

*Apoterium*, Blume, Bijdr. p. 218.

8. **C. Moonii**, Wight, Illust. i. 129; Icones, t. 111.—C.P. 3402.

HAB. Forests in the district between Galle and Ratnapoora; not uncommon. Nom. vulg. “Domba-keena-gass.”

*Arbor* magna. *Ramuli* juniores tomento ferrugineo fugaci vestiti. *Fructus* sphaericus.

9. **C. Burmanni**, Wight, Illust. i. 129; Icones, t. 107, 108; Burm. Fl. Zeyl. t. 60.—C.P. 242, 2447.

HAB. Hotter parts of the island, at no great elevation.

*Fructus* subsphaericus, rufo-brunneus,  $\frac{1}{2}$  poll. in diam.

## XX·XVIII. HIPPOCRATEACEÆ.

### 1. HIPPOCRATEA, Linn.

1. **H. Indica**, Willd. (W. et A. Prod. i. 104, cum syn.)—C.P. 1164.

HAB. Hot, drier parts of the island; not uncommon.

2. **H. obtusifolia**, Roxb. (W. et A., l. c. cum syn.; Wight, Icon. t. 963.)—C.P. 3158.

HAB. South of the island. Nom. vulg. “Deyakirindi-wel.”

### 2. KOKOONA, Thw.

Hook. Journ. of Bot. vol. v. (1853) p. 379.

*Stamina* 5, *petalis* alterna; *filamentis* subulatis, *disco* magno annulari expanso singulatim insertis. *Ovarium* semi-immersum, 3-loculare; *loculis* 4–12-ovulatis. *Ovula* erecta. *Capsula* triangularis, loculicide dehiscens, *valvis* 3, lignosis. *Semina* imbricata, longe alata.

1. **K. Zeylanica**, Thw. l. c. t. 6.—C.P. 2584.

HAB. Not uncommon on the banks of streams in the Saffragam and Ambagamowa Districts, at an elevation of 2000 to 4000 feet. Nom. vulg. “Kokoon-gass.”

The inner yellow bark of this tree is employed by the natives medicinally as a sternutatory, and an oil is expressed from the seeds, which is used for burning in lamps.

## 3. SALACIA, Linn.

1. **S. oblonga**, Wall. (W. et A. Prod. i. 106, cum syn.; Wight, Icones, t. 97; Illust. t. 47 B.)—c.p. 1061, 3148.

HAB. Common in the hot, drier parts of the island.

2. **S. reticulata**, Wight, Illust. i. 134.—c.p. 658.

HAB. Central Province, at an elevation of 1000 to 3000 feet. Nom. vulg. "Himbootoo-wel."

The natives eat the pulp of the ripe, rather large, spherical fruit.

3. **S. prinoides**, DC. (W. et A. Prod. i. 105; Wight, Icones, t. 321.)—c.p. 1165.

HAB. Central Province, up to an elevation of 2000 feet. Nom. vulg. "Heen-himbootoo-wel."

4. **S. diandra**, Thw.; *foliis* integris, ovatis, obtuse acuminatis, basi angustatis, brevi-petiolatis, subtus nitidis, delicatule reticulatis; *floribus* tetrameris, diandris.—c.p. 2720.

HAB. Central Province and Saffragam District, up to an elevation of 2000 feet; not common.

*Frutex* scandens, siccitate nigrescens. *Folia* 2–4½ poll. longa, 1–1¾ lata; *petiolo* 1–2 lin. longo. *Flores* axillares, solitarii; *pedicello* petiolo æquilongo. *Petala* rotundata, unguiculata, in anthesi horizontalia, lateribus reflexis; *calycis segmentis* triangularibus multo longioribus. *Staminum filamenta* brevia; *antheræ* extrorsæ, loculis subsphæricis, horizontaliter dehiscentibus. *Ovarium* biloculare, loculis bi-ovulatis.

## XXXIX. MALPIGHIACEÆ.

## 1. HIPTAGE, Gært. n.

1. **H. Madablota**, Gært. n. (W. et A. Prod. i. 107, cum syn.; Wight, Illust. i. 139. t. 50.)—c.p. 659.

HAB. Common up to an elevation of 3000 feet.

2. **H. parvifolia**, W. et A., l. c.—c.p. 1166, 1167.

HAB. Hot, drier parts of the island, *Gardner*.

*Platynema laurifolium*, W. et A., described as from Ceylon, is probably cultivated there only. Specimens in Hook. Herb. are from Singapore, but it may not be wild there.

## XL. ERYTHROXYLEÆ.

## 1. SETHIA, Kunth.

1. **S. Indica**, DC. (W. et A. Prod. i. 106, cum syn.; Wight, Illust. i. 136. t. 48.)—c.p. 1168.

HAB. Hot, drier parts of the island.

Mr. Ondaatze informs me that an empyreumatic oil or wood-tar, used for preserving the wood employed in the construction of native boats, is obtained from the wood of this tree.

2. **S. lanceolata**, Wight, *Illust.* i. 136; *foliis lanceolatis vel lineari-lanceolatis, obtusis*.—C.P. 493.

HAB. On the banks of streams, Galagama, at an elevation of 2000 to 3000 feet.

*Frutex* erectus, ramosus. *Folia* 2–3½ poll. longa, ½–1 poll. lata; *petiolo* 1½–2½ lin. longo.

This is, as truly observed by Dr. Wight, very closely allied to the next species; but, in addition to the difference in their shape, the under side of the leaves of the present plant is less shining, and the large areolæ formed by the inosculation of the primary veins approach nearer to the margin than they do in *S. obtusifolia*. If these are, as Dr. Wight suggests, to be considered only varieties of one species, *S. acuminata* would also have to be joined with them, and perhaps this would best express their real character, though I can hardly venture to unite them without further investigation.

3. **S. obtusifolia**; *foliis ovato-lanceolatis, ellipticis vel obovatis, apice obtuse brevi-acuminatis vel rotundatis, basi plus minus angustatis vel rotundatis*.

a. *stylosa*; *staminibus alternis longioribus, stylo brevioribus*.—*S. lanceolata*, var.  $\beta$  *obtusifolia*, Wight, *Illust.* i. 136.—C.P. 2613.

b. *staminea*; *staminibus æqualibus, stylo longioribus*.—*S. erythroxyloides*, Wight, l. c.—C.P. 25.

HAB. Central Province, at an elevation of 2000 to 4000 feet.

*Folia* 1½–3½ poll. longa, ¾–1¾ lata; *petiolo* 2–3½ lin. longo.

Until I found the same peculiarity existing in the next species, I considered, with Dr. Wight, that the different states indicated above under the letters *a* and *b* were distinct species: I am now disposed to look upon them rather as the foreshadowing of a dioicous character, though fruit is produced apparently in equal abundance upon the plants of either state.

4. **S. acuminata**, Arn. *Pug.* p. 6 (324); Wight, *Illust.* i. 135.—*Erythroxylon lucidum*, Moon's *Cat.*—C.P. 222.

Var.  $\beta$ ; *foliis minoribus, longius acuminatis; petiolo breviori*.—C.P. 3488.

a. *stylosa*; *staminibus alternis longioribus, stylo brevioribus*.

b. *staminea*; *staminibus æqualibus, stylo longioribus*.

HAB. Ambagamowa and Saffragam Districts, at an elevation of 1000 to 2000 feet. Nom. vulg. "Batta-kerilla-gass."

The smaller and much acuminate leaves, and the more acute segments of the calyx, seem alone to distinguish this from the preceding species.

The Cinghalese attach much value to this plant as an anthelmintic for children, giving the juice expressed from the fresh leaves.

## XLI. SAPINDACEÆ.

### 1. CARDIOSPERMUM, Linn.

1. **C. Halicacabum**, Linn. (*W. et A. Prod.* i. 109, cum syn.)—C.P. 2958.

HAB. Very abundant in the warmer parts of the island. Nom. vulg. "Painaira-wel."

It is probable that *C. canescens*, Wall., occurs also in the island, but I have never yet met with it.

## 2. SCHMIDELIA, Lin.

1. **S. Cobbe**, DC. Prod. i. 610, cum syn.; Wight, Icon. t. 964, 2; W. et A. Prod. i. 110, cum syn.—c.p. 1156.

HAB. Very abundant up to an elevation of 3000 feet. Nom. vulg. "Cobbai-gass."

2. **S. villosa**, Wight, Icon. t. 401.—? *S. Rheedii*, Wight, Ic. t. 964.—c.p. 3549.

HAB. Central Province, at an elevation of 3000 to 5000 feet.

I suspect that this will prove to be but a variety of the preceding species, from which it seems scarcely to differ except in being much more hairy.

3. **S. allophylla**, DC. Prod. i. 611, cum syn.; *foliis* lanceolatis vel ovato-lanceolatis, acuminatis, glabris, integris vel versus apicem grosse dentatis, basi angustatis.

Var. *a*; *racemis* brevibus, hispidis; *bracteolis* confertis.—c.p. 2679. *S. allophylla*, Arn. Pug. 8 (326).

Var. *β*; *racemis* elongatis, minute pilosis, florum fasciculis sparsis; *bracteolis* inconspicuis.—c.p. 562.

HAB. Var. *a*, Ambagamowa and Hinidoon Districts, up to an elevation of 3000 feet. Var. *β*, Central Province, at an elevation of 2000 to 5000 feet.

*Arbor* parva. *Folia* 4–10 poll. longa,  $1\frac{1}{2}$ – $3\frac{1}{2}$  poll. lata; *petiolo*  $\frac{1}{2}$ –1 poll. longo. *Flores* parvi; *petalis* longe unguiculatis; *squamulis* villosis, lamina æquilongis. *Baccæ* rotundatæ, rubræ.

4. **S. hispida**, Thw.; *ramulis* hispidis; *foliis* lanceolatis, acuminatis, basi rotundatis vel parce angustatis, glabris, margine costa venis primariisque utrinque cum *petiolo* longo hispidis; *racemis* brevissimis; *floribus* confertis.—c.p. 3294.

HAB. Ambagamowa District, at an elevation of 1000 to 2000 feet.

*Arbor* parva. *Folia* 5–14 poll. longa, 2–5 poll. lata; *petiolo*  $\frac{3}{4}$ –2 poll. longo. *Petala* unguiculata; *squamulis* longe villosis, lamina brevioribus.

5. **S. acuminata**, Thw.; *foliis* ovatis, obtuse rostrato-acuminatis, glabris; *racemis* elongatulis, subglabris, sparsifloris.—c.p. 3001.

HAB. Galagama, on the banks of streams, at an elevation of 2000 to 3000 feet.

*Arbor* parva. *Ramuli* teretes, glabri. *Folia*  $2\frac{1}{2}$ – $4\frac{1}{2}$  poll. longa, 1–2 poll. lata; *petiolo* 4–7 lin. longo. *Petalorum squamulæ* densissime villosæ, lamina æquilongæ.

## 3. SAPINDUS, Linn.

1. **S. laurifolius**, Vahl, (Wight et Arn. Prod. i. 111, cum syn.)—c.p. 2545.

HAB. Southern and central parts of the island, up to an elevation of 3000 feet.

The specimens in this Herbarium have the leaves tomentose underneath, and the petals, which are hairy over their entire inner surface, have two hairy glandular appendages, as in the following species.

2. **S. emarginatus**, Vahl, (W. et A., l. c. cum syn.; Wight, Illust. i. 141. t. 51.)—c.p. 1151.

HAB. Hot, drier parts of the island. Nom. vulg. "Penella-gass."

My specimens have the under side of the leaves quite glabrous, so that it would

appear that this and the foregoing species vary in having their leaves either glabrous or hairy beneath.

3. **S. unijugus**, Thw.; *foliis* bifoliolatis, glabris; *petiolo* tereti, basi tumido; *foliolis* lanceolatis, reticulatis, obtuse acuminatis, brevi-petiolumatis; *racemis* simplicibus vel ramosis, subterminalibus, oppositifoliis, minute pilosis.—C.P. 2465.

HAB. Hewahette District, at an elevation of 3000 to 4000 feet.

*Arbor* magna. *Folia* siccitate nigrescentia; *petiolo* 7-12 lin. longo; *foliolis* 3-5 poll. longis,  $1\frac{1}{4}$ -2 poll. latis; *petiolulo* tumido 2-3 lin. longo. *Calycis* segmenta acuta. *Stamina* 8. *Ovarium* hirsutum, 3-loculare; *stigmatibus* sessilibus, 3-radiatis. *Bacca* 1-3-sperma,  $\frac{3}{4}$ - $1\frac{1}{2}$  poll. in diam., subrotundata.

The specimens I have obtained of this plant do not enable me to give so full a description of the flower and fruit as I could wish. In appearance it a good deal resembles *Nephelium bifoliolatum* of this work, but the structure of the fruit is very different.

#### 4. HEMIGYROSA, Blume.

Rumphia, vol. iii. p. 165. Walpers, Ann. vol. ii. p. 212.

1. **H. trichocarpa**, Thw.; *foliis* abrupte pinnatis, subglabris; *foliolis* suboppositis, 2-4-jugis, lanceolatis, acuminatis, integris; *racemis* tomentosis, plus minus ramosis; *bracteolis* subulatis, curvatis, *pedicellis* longioribus.

Var.  $\beta$ ; *foliis* minoribus, angustioribus.—C.P. 607.

HAB. Var.  $\alpha$ , Central Province, up to an elevation of 3000 feet. Var.  $\beta$ , hot, drier parts of the island.

*Arbor* mediocris. *Petala* 4 (uno deficiente). *Ovarium* hirsutum, trigonum; *stylo* brevi. *Fructus* hirsutus, indehiscens, trigonus, trilobularis; *pericarpio* sublignoso, interne piloso. *Semina* oblonga; *testa* membranacea; *cotyledonibus* carnis, æqualibus, incumbentibus; *radicula* parva, ab hilo distante.

2. **H. canescens**, Thw.—*Cupania canescens*, W. et A. Prod. i. 113. *Molinæa canescens*, Roxb. Cor. i. 43. t. 60; *foliis* ovato-lanceolatis, glabris, subtus fusciscentibus; *racemis* plus minus ramosis, minute tomentosis; *bracteolis* minutis, triangularibus, *pedicellis* brevioribus.—C.P. 3508.

HAB. Lower Badulla Road from Kandy, at no great elevation.

This plant is so closely allied to the preceding that I suspect there must be some mistake in its having been described with dehiscent fruit and arillate seeds. The apparent arillus in Roxburgh's figure is, I believe, the hairy coating of the inside of the pericarp. The fruit in my specimens is very immature. I find the fifth (small) petal sometimes present, as figured by Roxburgh.

#### 5. HARPULLIA, Roxb.

*Otonychium*, Blume. *Streptostigma*, Thw.

1. **H. imbricata**.—*Otonychium imbricatum*, Bl. Rumphia, iii. 180; Walpers, Ann. ii. 213. *Streptostigma viridiflorum*, Thw. in Hook. Journ. of Bot. vol. vi. (1854) p. 298. t. 9 A. ? *Harpullia cupanioides*, Roxb. Fl. Ind. ii. p. 442.—C.P. 605, 736.

HAB. Central Province; not uncommon, up to an elevation of 3000 feet.

This must be closely allied to, if not identical with, Roxburgh's *Harpullia cupanioides*, but, if that eminent botanist's description be correct, the Ceylon plant

differs from his in having most frequently two instead of one seed in each locus of the seed-vessel; the seeds of a different colour and shape, and with an aril very minute, instead of reaching nearly to their apex: the position of the radicle would also appear to be different. The peculiar twisted character of the stigma, so evident in the Ceylon plant, would scarcely have been left unnoticed by Roxburgh and Wallich had it been present in the species described by them, which on the other hand is said to have a bifid style.

## 6. NEPHELIUM, Linn.

† *Petala appendiculata.*

1. **N. erectum**, Thw.; *foliis* majusculis, abrupte pinnatis; *foliolis* 3-4-jugis, suboppositis, lanceolatis, acuminatis, integris, glabris, reticulatis; *paniculis* terminalibus, multifloris, foliis multo brevioribus; *fructibus* 1-3- raro 4-lobis, minute tomentosis, lobis oblongis.—C.P. 1150 (in parte).

HAB. Central Province, up to an elevation of 3000 feet.

*Frutex* erectus, non vel parce ramosus, 6-10-pedalis. *Foliorum rachis* striata, 10-14 poll. longa; *foliolis* 6-10 poll. longis, 2-3½ latis, brevi-petiolaris. *Flores* 5-meri. *Calycis segmenta* rotundata. *Petala* late unguiculata, medio squamata; *lamina* rotundata, reflexa, lævi, margine involuta; *squama* erecta, bifida, villosa. *Discus* annuliformis, lævis. *Stamina* 8. *Stylus* brevissimus. *Stigma* minutum, 3-lobum. *Fructus* carnosus, luteus; *lobis* circiter 1 poll. longis. *Semina* ovoidea, arillo carnoso hyalino integro inclusa; *testa* membranacea, rufa; *cotyledonibus* crassis, incumbentibus.

2. **N. simplicifolium**, Thw.; *foliis* integris, cuneato-lanceolatis, acuminatis, lævibus, membranaceis, reticulatis; *paniculis* brevibus, subterminalibus, minutissime tomentosis.—C.P. 443.

HAB. South of the island, at no great elevation; rare.

*Frutex* erectus, 6-10-pedalis, parce ramosus. *Folia* 8-12 poll. longa, 2½-4 poll. lata; *petiolo* ½-2 poll. longo. *Flores* 5-meri. *Calycis segmenta* rotundata. *Petala* exunguiculata, rotundata, ciliata, basi *squamula* parva villosa munita. *Discus* regularis, lævis. *Stamina* 8, *ovariorum rudimentum* 3-lobatum cingentia.

Until the fruit of this species is known, there must remain some little doubt as to whether it is a true *Nephelium*. I have as yet seen only very young male flowers of it.

3. **N. bifoliolatum**, Thw.; *foliolis* unijugis, integris, lanceolatis, obtusiusculis, basi angustatis, glabris; *paniculis* terminalibus, foliis æquilongis vel longioribus.—C.P. 1721.

HAB. Lower Badulla Road from Kandy, at no great elevation. (Flor. Aprili.)

*Arbor* mediocris, 20-30-pedalis, ramosissima. *Foliola* superne nitida, reticulata, subtus opaca, fuscescentia, 3-5 poll. longa, ¾-1½ poll. lata, *petiolulo* 2-3 lin. longo, *petiolum* ½-1 poll. longum terminalata. *Paniculæ* pilosulæ. *Flores* 5-meri. *Calycis segmenta* rotundata, glabra, ciliata. *Petala* parce longiora, oblonga, glabra, ciliata, abrupte brevi-unguiculata, basi *squamula* parva rotundata longe ciliata munita. *Discus* carnosus, crenulatus. *Stamina* sæpissime 7; *masculinis filamentis* linearibus, *petala* æquantibus, basi pilosis; *fœm.* brevibus, *ovarium* glabrum bilobum cingentibus. *Stylus* linearis. *Stigma* minutum, obscure bilobum.

4. **N. eximium**.—*Eccremanthus eximius*, Thw. in Hook. Journ. of Bot. vii. p. 272. t. 9; *foliis* magnis, abrupte pinnatis, subsessilibus; *foliolis* 5-13-jugis, oblongo-lanceolatis, serratis, subsessilibus, subtus hirsutulis, parte inferiore minore, obliquo, curvato, sæpe deciduo; *pani-*



*culis* tomentosis, ramis elongatis, pendulis, multifloris, cylindricis; *floribus* minutis.—C.P. 1153.

HAB. Central Province, at an elevation of 1000 to 2000 feet. (Flor. Maio; fruct. Julio.)

*Arbor* magna, 40–60-pedalis. *Ramuli* juniores rufo-hirsuti. *Flores* 5-meri. *Calycis segmenta* brevia, subacuta. *Petala* minuta, transverse oblonga, exunguiculata, *squama* adnata subæquilonga margine dense villosa munita. *Discus* carnosus, rufescens. *Stamina* 5, *petalis* multo longiora. *Ovarium* bilobum, hirsutulium; *stylo* lineari; *stigmatibus* minuto, bilobo. *Fructus* carnosus, subglaber, sæpissime 1-lobus. *Semen* oblongum, *arillo* hyalino fere tectum, hilo magno obliquo; *testa* rufa.

†† *Petala exappendiculata.*

5. **N. Longanum**, Camb. (W. et A. Prod. i. 113, cum syn.)—*N. Mora*, Gardner, MSS. *Dimocarpus pupilla*, Moon's Cat. p. 31.—C.P. 1152.

HAB. Abundant up to an elevation of 2000 feet. Nom. vulg. "Mora-gass."

*Ramuli*, cum partibus aliis junioribus, tomento stellari rufescente vestiti. *Fructus* immaturus tuberculatus, stellari-tomentosus, demum glaber, lævis; *arillo* plus minus copioso.

The Cinghalese eat the fleshy aril of the fruit of this tree, and the wood is useful for building purposes.

6. **N. Gardneri**, Thw.; *foliis* abrupte pinnatis, subglabris; *foliolis* 2–3-jugis, lanceolatis, grosse et obtuse dentatis, brevi-petiolulatis, subtus glaucis.—C.P. 1154.

HAB. Putlam, Gardner.

Præcedenti valde similis affinisque, sed foliis dentatis, sepalis petalisque acutioribus, et pubescentia vix stellari satis distincta videtur.

††† *Petala nulla.*

7. **N. fuscatum**, Thw.; *foliis* abrupte pinnatis, glabris; *foliolis* 2-jugis, lanceolatis, obtusis vel retusis, basi cuneatis, reticulatis, brevi-petiolulatis; *racemis* terminalibus, minute fusco-pilosis, ramosis; *ramis* densifloris, spicæformibus.—C.P. 2577.

HAB. Putlam, Gardner.

*Foliola* 3–4 poll. longa,  $\frac{3}{4}$ – $1\frac{1}{4}$  poll. lata. *Flores* minuti. *Calycis segmenta* acuta. *Discus* complanatus, crenatus. *Stamina* 8–10, glabra.

I have not seen the fertile flowers or fruit of this plant.

7. **SCHLEICHERA**, Willd.

1. **S. trijuga**, Willd. (W. et A. Prod. i. 114, cum syn.)—C.P. 1155.

HAB. Common in the warmer parts of the island, up to an elevation of 2000 feet. Nom. vulg. "Cōng-gass."

The Cinghalese express an oil from the seeds of this tree, which they use for burning, and a considerable quantity of lac is often produced upon the young branches.

8. **PTERIDOPHYLLUM**, Thw.

Hook. Journ. of Bot. vol. vi. (1854) p. 65.

*Flores* polygamo-monoici. *Stamina* 5, *disco* intus complanato crenato inserta. *Ovarium* rotundatum, biloculare. *Ovula* in loculis solitaria,

pendula, anatropa. *Stylus* linearis. *Stigma* minutum, bilobum. *Fructus* drupaceus, sæpissime 1-spermus; *putamine* membranaceo. *Semen* exarillatum; *testa* membranacea; *cotyledonibus* foliaceis, plicatis; *radicula* dorsali, hilo fere attingente.

1. **P. decipiens**, Thw. l. c. t. 1 *B.*—*Rhus decipiens*, W. et A. Prod. i. 172; Wight, Illust. i. 186. t. 75.—C.P. 536.

HAB. Central Province, up to an elevation of 3000 feet. Nom. vulg. "Pehimbia-gass." (Flor. Jan.; fruct. Mar.)

The wood of this very ornamental tree is useful for building purposes.

### 9. DODONÆA, Linn.

1. **D. Burmanniana**, DC. (W. et A. Prod. i. 114, cum syn.; Wight, Illust. i. 142. t. 52.)—C.P. 1158.

HAB. Central Province, up to an elevation of 7000 feet. Nom. vulg. "Attawarella-gass."

### 10. MELIOSMA, Blume.

*Millingtonia*, Roxb., W. et A.

1. **M. pungens**, Walp. Rep. i. 423.—*Millingtonia pungens*, Wall.; W. et A. Prod. i. 115, cum syn.; Wight, Icones, t. 964/3.—C.P. 300 (580).

HAB. Central Province, at an elevation of 5000 to 7000 feet.

*Folia* subintegra vel distanter serrulata, superne, costa venis primariisque exceptis, glabra (juniora sparse pilosa), subtus plus minus hirsuta.

2. **M. simplicifolia**, Walp. l. c.—*Millingtonia simplicifolia*, Roxb.; W. et A., l. c. cum syn.—C.P. 703.

HAB. Abundant up to an elevation of 3000 feet. Nom. vulg. "Albadda-gass," "Rawaniddala-gass."

3. **M. Arnottiana**, Walp. l. c.—*Millingtonia Arnottiana*, Wight, Illust. i. 144. t. 53.—C.P. 293.

HAB. Central Province, at an elevation of 4000 to 7000 feet.

## XLII. MELIACEÆ.

### 1. MUNRONIA, Wight.

1. **M. pumila**, Wight, Icones, t. 91.—C.P. 177.

HAB. Not uncommon in the warmer parts of the island. Nom. vulg. "Bin-kohōmba-gass."

This is much valued by the Cinghalese as a medicinal plant.

### 2. MELIA, Linn.

1. **M. composita**, Willd. (W. et A. Prod. i. 117, cum syn.)—C.P. 699.

HAB. South of the island. Nom. vulg. "Loonoomidella-gass."

### 3. AZADIRACHTA, Adr. Juss.

1. **A. Indica**, Adr. Juss. (W. et A., l. c. 118, cum syn.; Wight, Icon. 396.)—C.P. 1159.

HAB. Hot, drier parts of the island, at no great elevation. Nom. vulg. "Kohōmba-gass."

The juice of the leaves of this plant is employed as an anthelmintic for cattle.

#### 4. MALLEA, Adr. Juss.

1. *M. Rothii*, Adr. Juss. (W. et A., l. c. cum syn.)—C.P. 496.

HAB. Very abundant, up to an elevation of 2000 feet.

#### 5. MILNEA, Roxb.

1. *M. Roxburghiana*, W. et A. Prod. i. 119; Wight, Icon. t. 166.—C.P. 1148.

HAB. Common in the Central Province, up to an elevation of 3000 feet.

This and the next species are polygamo-dioicous, the barren flowers being smaller and in larger panicles than the fertile ones.

2. *M. apiocarpa*, Thw.; *foliis* imparipinnatis, *foliolis* 5-9, ovatis, obtuse acuminatis, basi cuneatis; *junioribus* subtus spadiceo-furfuraceis; *floribus* pedicellis brevioribus; *fructibus* pyriformibus.—C.P. 405, 2448.

HAB. Central Province, at an elevation of 3000 to 6000 feet.

*Arbor* parva. *Ramuli* juniores spadiceo-furfuracei. *Petala* externe furfuracea. *Cotyledones* singulæ in duas partes plus minus inæquales fere ad basin divisæ.

Closely allied to the preceding, but differing in its larger and more furfuraceous leaves and flowers. In the present species the flowers are shorter than their pedicels, whereas in *M. Roxburghiana* they are longer.

#### 6. AMOORA, Roxb.

1. *A. Rohituka*, W. et A. Prod. i. 119—C.P. 558, 2479.

HAB. Central Province, up to an elevation of 3000 feet. Nom. vulg. "Hingool-gass."

*Fructus* subglobosus, rufescens, 1-1½ poll. in diam., basi parum attenuatus. *Semina* oblonga, brunnea, arillo coccineo vestita.

#### 7. DYSOXYLON, Blume.

1. *D. macrocarpum*, Blume, Bijdr. p. 175.—*Guarea binectarifera*, Roxb. Cat. ? *Amoora ficiformis*, Wight, Illust. i. 147.—C.P. 2637 (1150 in parte).

Var.  $\beta$ ; *foliis* minoribus, angustioribus.

HAB. Central Province, up to an elevation of 3000 feet. Var.  $\beta$ . Batticaloa, Gardner.

*Arbor* ingens. *Foliola* 5-8 poll. longa, 1½-3 poll. lata, *petiolulo* 4 lin. longo. *Paniculæ* supra-axillares, 3-5 poll. longæ, minutissime incano-tomentosæ, sparsifloræ; *pedicellis* articulatis. *Flores* pallide virides, octandri, circiter 3 lin. longi. *Calyx* cupuliformis, 4-emarginulatus. *Petala* reflexa, acuta, externe minute tomentosa. *Tubus stamineus* 4-gonus, dentibus 8, obtusis, bifidis. *Tubus interior ovario* rotundato, piloso æquilongus, denticulatus. *Fructus* rubescens, rotundatus, 4-ocularis, 4-spermus, circiter 2½ poll. in diam., brevi-pedunculatus.

I suspect I have not seen the fertile flowers of this plant, for, judging from the

fruit being so shortly peduncled, it would seem probable that they are on much shorter panicles or racemes.

2. **D. Championii**, H.f. et T. MSS.; *foliis* glabris, abrupte pinnatis; *foliolis* 2-3-jugis, oblongo-spathulatis, subcoriaceis, aveniis, apice rotundatis.—C.P. 1193, 3596.

HAB. Central Province, up to an elevation of 4000 feet.

*Arbor* magna. *Foliola* 2-4 poll. longa,  $\frac{3}{4}$ -1 $\frac{1}{2}$  poll. lata. *Paniculae* supra-axillares, 2-3 poll. longæ, minutissime pilosulæ. *Flores* octandri, flavescentes, circiter 3 lin. longi. *Calyx* brevis, 4-lobus. *Petala* glabra, acuta. *Tubus stamineus* dentibus 8, obtusis, bifidis. *Tubus interior* nullus. *Fructus* pyriformis, pollicaris, *loculis* 1-2-spermis.

This differs from the other species of *Dysoxylon* in having no inner tube surrounding the ovary.

## 8. WALSURA, Roxb.

1. **W. piscidia**, Roxb. (W. et A. Prod. i. 120, cum syn.; Wight, Illust. i. 148. t. 55.)—C.P. 1162.

HAB. Common up to an elevation of 3000 feet. Nom. vulg. "Kirrekōng-gass."

2. **W. Gardneri**, Thw.; *foliis* unifoliolatis, glabris; *foliolo* integro, ovato, obtuse-acuminato; *petiolulo* brevi, cum *petiolo* multo longiore articulado.—C.P. 1163.

HAB. Central Province, at an elevation of 2000 to 4000 feet.

*Arbor* parva, 10-12-pedalis. *Foliola* 3-6 poll. longa, 1 $\frac{1}{2}$ -2 $\frac{1}{2}$  lata, *petiolulo* 1-3 lin. longo. *Fructus* oblongus, parum apiculatus, 6-8 lin. longus.

The flowers of this are very similar to those of the preceding species, but the fruit is larger and more oblong. The axis of the panicle is most frequently produced into a leaf-branch, in which case the inflorescence appears scattered along the stem.

## 9. XYLOCARPUS, König.

1. **X. Granatum**, Kön. (W. et A. Prod. i. 121, cum syn.)—C.P. 2650.

HAB. South of the island; not common.

## XLIII. CEDRELACEÆ.

### 1. CHICKRASSIA, Adr. Juss.

1. **C. tabularis**, Adr. Juss. (W. et A. Prod. i. 123; Wight, Illust. i. 149. t. 56.)

Var.  $\beta$ ; *foliis* subtus tomentosus.—C.P. 2454.

HAB. Warmer parts of the island. Nom. vulg. "Hoolanghik-gass."

The wood of this tree is useful for building purposes.

### 2. CHLOROXYLON, DC.

1. **C. Swietenia**, DC. (W. et A. Prod. i. 123, cum syn.; Wight, Illust. i. 149. t. 56 *bis*.)—C.P. 1160.

HAB. Hot, drier parts of the island. Nom. vulg. "Booroota-gass."

This tree yields the satin-wood of commerce.

## XLIV. AMPELIDEÆ.

## 1. CISSUS, Linn.

1. **C. edulis**, Dalz. in Hook. Lond. Journ. of Bot. ix. p. 248.—*C. quadrangularis*, Roxb. (non Linn.). *Vitis quadrangularis*, Wall.; W. et A. Prod. i. 125; Wight, Icones, t. 51.—c.p. 1174.

HAB. Common in the hot, drier parts of the island.

The Cinghalese use the succulent stems of the plant as a medicine.

2. **C. glyptocarpa**, Thw.; *ramulis* tenuibus, 4-alatis; *foliis* truncato-cordatis, acuminatis, serratis, glabris, subtus basi, costa nervisque adpresse pilosis; *stipulis* cymbiformibus; *cymis* petiolo longioribus, medium versus 3-5-radiatis, glabris.—c.p. 1183.

HAB. Hot, drier parts of the island.

*Folia* 2-3 poll. longa, 1-1 $\frac{3}{4}$  lata; *petiolo* alato,  $\frac{3}{4}$ -1 $\frac{1}{4}$  poll. longo. *Flores* parvi, rufescentes. *Petala* acuta, reflexa. *Stylus* cylindricus. *Stigma* minutum, bilobum. *Nuculania* oblonga, 3 $\frac{1}{2}$  lin. longa, 1-sperma, nigro-purpurea. *Semen* pyriforme, dorso 6-seriatim excavato-punctatum, antice 2-sulcatum.

3. **C. repens**, Lam. (W. et A. Prod. i. 125, cum syn.)—c.p. 1180.

HAB. Warmer parts of the island; not uncommon.

4. **C. acuminata**, Thw.; *ramulis* teretibus; *foliis* membranaceis, truncato-cordatis, rostrato-acuminatis, denticulatis, glabris, superne costa venis primariisque appresse pilosis, longe petiolatis; *cymis* spadicis tomentosus, petiolo multo brevioribus.—c.p. 3451.

HAB. Ratnapoora District.

*Folia* 5-7 poll. longa, 2 $\frac{1}{2}$ -4 poll. lata; *petiolo* 2 $\frac{1}{2}$ -3 $\frac{1}{2}$  poll. longo. *Stipulae* reniformes, medio adnatæ. *Petala* externe pilosa.

Closely allied to the preceding species, but differing from it in the shape and texture of the leaves, and in these being usually five-nerved at the base, whereas in *C. repens* they are most commonly three-nerved; in the latter, too, the lamina is produced below the point of attachment of the petiole, giving the leaf a subpeltate character, but it is not so in the present species.

5. **C. glauca**, Roxb. Fl. Ind. i. 425.—*Vitis glauca*, W. et A. Prod. i. 126.—c.p. 2939.

HAB. Abundant, up to an elevation of 2000 feet.

6. **C. lonchiphylla**, Thw.; glabra, *foliis* oblongo-lanceolatis, acuminatis, remote denticulatis, basi truncato-rotundatis, 3-nerviis, parum subpeltatis; *cymis* brevi petiolo longioribus.—c.p. 1177.

HAB. Ambagamowa District, at no great elevation; rare. Muruta, Gardner.

*Folia* 4-7 poll. longa, 1 $\frac{1}{2}$ -2 $\frac{1}{2}$  poll. lata, *petiolo* 3-6 lin. longo. *Stipulae* cordatæ, medio adnatæ. *Flores* parvi. *Nuculania* rotundata, depressa.

7. **C. adnata**, Roxb.—*Vitis adnata*, Wall.; W. et A. Prod. i. 126, cum syn.; Wight, Icones, t. 144.—c.p. 3450.

HAB. Warmer parts of the island; not uncommon.

8. **C. angulata**, Lam.—*Vitis Linnæi*, Wall.; W. et A. Prod. i. 126, cum syn.—c.p. 1181.

HAB. Hot, drier parts of the island; common.

9. **C. trilobata**, Lam.—*Vitis Rheedii*, W. et A. Prod. i. 127, cum syn.—c.p. 1182.

HAB. Warmer parts of the island; common.

10. **C. setosa**, Roxb.—*Vitis setosa*, Wall.; W. et A. Prod. i. 127, cum syn.; Wight, Icones, t. 170.—c.p. 1179.

HAB. North of the island, Gardner.

11. **C. carnososa**, Lam.—*Vitis carnososa*, W. et A. Prod. i. 127, cum syn.—c.p. 2938.

HAB. Warmer parts of the island; common.

12. **C. muricata**.—*Vitis muricata*, Wall.; W. et A. Prod. i. 128, cum syn.; Wight, Icones, t. 740. ? *C. papillosa*, Bl. Bijdr. p. 183.

Var.  $\beta$ . *minor*.—c.p. 1178.

HAB. Central Province, up to an elevation of 3000 feet; not uncommon.

*Folia* simplicia vel 2-3-foliolata; *foliolis* 2-3½ poll. longis. *Nuculania* albida, 3-5 lin. in diam., 1-4-sperma.

I can find no character but its smaller size to distinguish this from the plant figured by Dr. Wight.

13. **C. Gardneri**, Thw.; glabra, *foliis* 3-foliolatis; *foliolis* ovato-lanceolatis, repando-dentatis, brevi-petiolatis, basi cuneatis; *lateralibus* minoribus obliquis; *stipulis* cordato-oblongis, adnatis; *cymis* petiolo multo longioribus.—c.p. 1176.

HAB. Matele and Hunasgiriya Districts; not common.

*Ramuli* teretes, rufescentes. *Folia* petiolo circiter 1 poll. longo; *foliolo* intermedio 2-3 poll. longo, 1-1½ poll. lato. *Cymæ* paucifloræ, rufescentes. *Nuculania* pisi magnitudine, nigro-purpurea.

14. **C. lanceolaria**, Roxb.—*Vitis lanceolaria*, Wall.; W. et A., l. c. cum syn.; Wight, Icones, tt. 28 et 177.—c.p. 3597.

HAB. Near Nalande.

It seems probable that *C. muricata* is only a form of this species, as suggested by Wight and Arnott.

15. **C. pedata**, Lam.—*Vitis pedata*, Wall.; W. et A. Prod. i. 128, cum syn.—c.p. 1175, 2725.

HAB. Very abundant. Nom. vulg. "Madeya-wel."

Varies very much in the amount of pubescence, sometimes being nearly glabrous.

16. **C. reticulata**, Thw.; glabra, *foliis* 3-foliolatis; *foliolis* subæqualibus, ovatis, cuspidatis, obscure denticulatis, subtus prominenti-reticulatis, basi angustatis; *lateralibus* obliquis; *cymis* petiolo longioribus.—c.p. 2857.

HAB. Ambagamowa District; rare.

*Folia* petiolo 1½-2½ poll. longo; *foliolis* 4-5 poll. longis, 2-2½ poll. latis; *petiolulis* *lateralibus* 3-6 lin. longis; *intermedio* pollicari. *Nuculania* ovoidea, pallide rosea, cerasi magnitudine. *Semina* oblonga, 4 lin. longa.

Closely allied to some forms of the preceding, but its differently-shaped and reticulated leaves at once distinguish it. The fruit is larger and somewhat attenuated at the apex, and the seeds, though with similar markings, are half as long again as those of *C. pedata*. The ternate character of the leaflets will probably be found not to be constant.

## 2. VITIS, Linn.

1. **V. Indica**, Linn. (W. et A. Prod. i. 131, cum syn.)—c.p. 1184.

HAB. Abundant, up to an elevation of 3000 feet.

3. **LEEAE**, Linn.

1. **L. Staphylea**, Roxb. (W. et A. Prod. i. 132, cum syn.; Wight, Illust. i. 153. t. 58; Wight, Icon. t. 78.)—c.p. 732.

HAB. Very common, up to an elevation of 3000 feet. Nom. vulg. "Booroola-gass."

## XLV. GERANIACEÆ.

1. **GERANIUM**, Linn.

1. **G. Nepalense**, Sweet, Ger. t. 12; DC. Prod. i. 643.—*G. affine*, W. et A. Prod. i. 133.—c.p. 2788.

HAB. Between Newera Ellia and Maturatte, at an elevation of 7000 feet; not common.

## XLVI. OXALIDEÆ.

1. **OXALIS**, Linn.

1. **O. corniculata**, Linn. (W. et A. Prod. i. 142, cum syn.)—*O. repens*, Thunb.; DC. Prod. i. 693.—c.p. 1189.

Var.  $\beta$ . *minor*.—c.p. 1188.

HAB. Central Province, abundant up to an elevation of 7000 feet.

2. **BIOPHYTUM**, DC.

1. **B. sensitivum**, DC. Prod. i. 690, cum syn.—c.p. 1190.

HAB. Very abundant in the warmer parts of the island.

2. **B. nudum**, Arn. Pug. viii. 326; Wight, Illust. i. 161.—c.p. 464 (6 in parte).

HAB. Central Province, at an elevation of 4000 to 6000 feet.

The leaflets of this species are less falcate and of a firmer texture than those of the preceding, of which it is perhaps only a variety, and which it resembles in its subglabrous styles, shorter than those of the two following species.

3. **B. proliferum**, Arn. Pug. ix. 327; Wight, Illust. i. 162.—c.p. 2418.

Var.  $\beta$ ; *foliis* majoribus; *foliolis* plus numerosis.—*B. intermedium*, Wight, l. c.—c.p. 2419 (6 in parte).

HAB. Central Province, at an elevation of 4000 to 7000 feet.

Besides the difference of habit, the longer pilose styles distinguish this from the two preceding species.

4. **B. nervifolium**, Thw.; *caule* simplici; *foliolis* circiter 12-jugis, obovato-oblongis, obtuse apiculatis; *venis* primariis paucis divaricatis, subtus prominentibus; *pedicellis* floribus brevioribus; *stylis* longis, pilosis, apice bifidis.—c.p. 2787.

HAB. Putlam, A. O. Brodie, Esq.

In habit and general appearance this a good deal resembles *B. sensitivum*, but the different shape and venation of the folioles, and the longer, hairy styles, seem satisfactorily to distinguish it.

## XLVII. BALSAMINEÆ.

1. **IMPATIENS**, Linn.

(Nom. vulg. "Koodaloo-mal.")

Sect. 1. *Foliis oppositis.*

1. **I. rosmarinifolia**, Retz, (Arn. in Comp. to Bot. Mag. i. 325, cum syn.; Walp. Rep. i. 475.)—c.p. 31.

HAB. Very abundant, especially amongst high grass, from the level of the sea to an elevation of 7000 feet.

Sect. 2. *Foliis alternis. Pedicellis axillaribus, solitariis vel pluribus.*

2. **I. Balsamina**, Linn. (W. et A. Prod. i. 135, cum syn.)—c.p. 1213.

HAB. Central Province, at an elevation of 3000 to 6000 feet.

3. **I. cuspidata**, W. et A. Comp. to Bot. Mag. i. 321; Walp. Rep. p. 468; Wight, Icones, t. 741.—c.p. 156.

Var.  $\beta$ . *foliis longioribus*.—*I. bipartita*, Arn. l. c. i. 322; Walp. Rep. p. 468.—c.p. 503.

HAB. Var. *a*. Above Galagama, Gardner. Var.  $\beta$ . Central Province, at an elevation of 4000 to 7000 feet.

I can find nothing to distinguish var.  $\beta$  from the other form of the plant but its longer, narrower leaves, and these with rather more numerous and less distinct primary veins. The flowers are of a dull reddish-purple in both: the relative size of the anterior lobe of the petals varies a good deal in different specimens.

4. **I. Henslowiana**, Arn. l. c.; Walp. l. c. p. 469.—c.p. 49.

Var.  $\beta$ . *foliis tomentosus*.—*I. albida*, Wight, Icones, t. 743.—c.p. 547.

HAB. Central Province, at an elevation of 4000 to 6000 feet; on rocks.

The smoother variety of this species agrees very completely with Dr. Arnott's description, except that the posterior sepal is not three-lobed, but two-lobed, with a wide sinus. The flowers are white, or slightly tinged with red.

5. **I. flaccida**, Arn. l. c. p. 322; Walp. l. c. p. 468.—*I. latifolia*, Moon's Cat. p. 18.—c.p. 2791.

HAB. Abundant, up to an elevation of 3000 feet.

Herbacea, decumbens. Flores rubro-purpurei, pallidi vel rarius candidi, 1-1½ poll. in exp. Sepalo posteriore obcordato, carinato; petalorum lobis subæqualibus, rotundatis, emarginatis. Semina juniora pilis patentibus deciduis vestita.

I do not find the spur in this species thicker towards the middle, and the appearance was probably caused in Dr. Arnott's specimen by unequal pressure in drying. This must be closely allied to, if not a variety of, *I. latifolia*, Linn., judging from the figure in Rheede, Hort. Mal. ix. 91. t. 48; and the last paragraph in the description of t. 52, page 101, in the same volume, would seem to imply that the former has smooth capsules, though it is referred by Messrs. Wright and Arnott to a species with them tomentose.

6. **I. leptopoda**, Arn. l. c. p. 321.—*I. brevicornu*, Arn. l. c. *I. gibbosa*, Arn. l. c.; Walp. l. c. pp. 468, 469. ? *I. Leschenaultii*, Wight, Icon. t. 970 bis.—c.p. 140 (455).

HAB. Central Province, at an elevation of 5000 to 8000 feet.

Erecta, 1-3-pedalis. Petala rosea, divergentia, lobis subæquilongis, posteriore parum latiore, emarginato. Capsulæ glabræ.

A very common, and also a very variable species, and the flowers are frequently quite ecalcarate. The scattered, brittle, yellow pubescence sometimes upon the



plant, would seem to be of an abnormal character, and perhaps the absence or reduced size of the spur arises from disease or injury. *I. Leschenaultii* of Dr. Wight's 'Icones,' closely resembles some states of the present; but in that the petals appear to be approximated, whilst in this they are divergent.

7. ***I. truncata***, Thw.; *foliis* ovato-lanceolatis, acuminatis, denticulatis, sparse utrinque pilosis, basi angustata petioloque parce glanduloso-setigeris; *pedicellis* 1-3-nis, tenuibus, una cum alabastris pilosis, foliis brevioribus; *sepalis lateralibus* minutis, lanceolatis, *posteriore* multo majore, rotundato, cuspidato, *anteriore* hoc æquilongo, cuspidato, concavo, *calcare* recto, brevi, cylindrico, truncato; *petalis* approximatis, inæqualiter bilobis, lobo anteriore multo majore, subsemicirculari; *capsulis* junioribus pilosis.—C.P. 2426.

Var.  $\beta$ . *subglabra*; planta fere tota glabra.

HAB. Central Province, at an elevation of 4000 to 6000 feet.

Herbacea, 1-2-pedalis, basi decumbens, radicans. *Ramuli* rufescentes. *Folia* 1-2 poll. longa,  $\frac{1}{2}$ -1 $\frac{1}{2}$  poll. lata, subtus pallidiora, *petiolo*  $\frac{1}{2}$ -1 $\frac{1}{2}$  poll. longo. *Flores* pallide rosei, externe rufescentes, *petalorum lobis* basi rufo-tinctis.

Apparently very closely allied to *I. inconspicua*, Benth., Wight, Icon. t. 970, but is distinguished by its alternate leaves, spurred flowers, and much smaller, lateral sepals.

8. ***I. macrophylla***, Gardner, MSS.; Hook. Bot. Mag. 4662.—C.P. 1212.

HAB. Central Province, at an elevation of 4000 to 6000 feet.

*Sepala lateralibus* anguste oblonga, *posteriore* 3-4-plo breviora. *Petala* lutea, subbiloba, apice exsecta, inclusa.

Nearly allied to, and in its foliage closely resembling the next species, but distinguished by its more numerous flowers, shorter and less saccate lower sepal, and included petals.

9. ***I. glandulifera***, Arn. in Hook. Comp. Bot. Mag. ii. 322.—*I. cornigera*, Hook. Bot. Mag. t. 4623, non Arn.—C.P. 2427.

HAB. Central Province, at an elevation of 4000 to 6000 feet.

*Caulis* erectus, 4-6-pedalis. *Folia* majuscula, denticulata, longe petiolata. *Pedicelli* fasciculati, petiolis glanduloso-setigeris breviores. *Sepala lateralibus* minuta, viridia; *posterius* cucullatum, viride, margine rufescente; *anterius* luteo-rufescens, profunde saccatum, in calcar breve incurvum apice incrassatum angustatum. *Petala* rosea, lobis subæqualibus, rotundatis; *posteriore* interne appendiculato; *anteriore* porrecto, exserto, apice emarginato.

10. ***I. repens***, Moon's Cat.; Wight, Illust. i. 160. t. 61; Walp. l. c. p. 469.—C.P. 2790.

HAB. Southern and central parts of the island, at no great elevation.

Sect. 3.—*Foliis alternis; pedunculis flores plures gerentibus.*

11. ***I. Walkeri***, Hook. (Arn. l. c. p. 324. t. 18; Walp. l. c. p. 471.)—C.P. 1211.

HAB. Central Province, at an elevation of 4000 to 6000 feet.

12. ***I. Hookeriana***, Arn. l. c. p. 324; Walp. l. c. p. 471.—? *I. grandis*, Heyne; Roxb. Fl. Ind. (ed. Wallich) ii. 464.—C.P. 2425.

HAB. Central Province, at an elevation of 3000 to 5000 feet.

I strongly suspect that this beautiful species is identical with *I. grandis*, Heyne. It varies a good deal in the shape of the lateral sepals, and in the size and colour of the flowers, which latter are either quite white or more or less blotched with pale or deep blood-colour.

13. **I. elongata**, Arn. l. c. p. 324; Walp. l. c. p. 473.—c.p. 436 (453).

HAB. Near Adam's Peak, at an elevation of 3000 to 4000 feet.

*Flores* rubri. *Petalorum lobus posterior* parvus, sepalis lateralibus æquilongus; *anterior* multo major, oblique rotundatus, obtusus.

14. **I. subcordata**, Arn. l. c. p. 323; Walp. l. c. p. 471.—c.p. 1230 and 2789.

HAB. Central Province, at an elevation of 3000 to 6000 feet.

Herbacea, 1-2-pedalis, basi decumbens, radicans. *Folia*  $1\frac{1}{2}$ - $2\frac{3}{4}$  poll. longa,  $\frac{3}{4}$ - $1\frac{1}{2}$  poll. lata, *petiolo*  $\frac{3}{4}$ -2 poll. longo. *Flores* albidii, sepalis lateralibus posterioreque rubescentibus.

15. **I. cornigera**, Arn. l. c. p. 323; Walp. l. c. p. 473.—c.p. 332.

HAB. Warmer parts of the island, at no great elevation, in damp, shady woods.

Glaberrima. *Caulis* simplex,  $\frac{1}{2}$ -2-pedalis. *Folia* pallide vel nigro-viridia, sæpe ad costam albido-areolata, subtus pallida vel rufo-plumbea, *petiolo* mediocri. *Flores* citrini, plus minus viridi vel rubro tincti. *Sepala lateralibus* parva, ovata, subcordata, acuminata; *posterior* rotundatum, subcucullatum, apiculatum; *anterior* in calcar longum subcylindricum apice attenuato sursum arcte uncinato productum. *Petalorum lobus posterior* minutus; *anterior* multo major, subquadratus, pedicello æquilongo suffultus.

16. **I. Arnottii**, Thw.; *foliis* ovato-lanceolatis, acuminatis, crenatis, denticulatis, superne sparsissime pilosis, subtus glabris, basi angustatis, *petiolo* eglanduloso; *sepalis lateralibus* parvis, cordatis, valde obliquis, obtuse apiculatis, *anterior* in calcar gracile incurvum sensim attenuatum flore duplo longius productum; *petalorum lobo posteriore* parvo, *anterior* multo majore, subsemicirculari, pedicellato, porrecto.—c.p. 725.

HAB. Central Province, at an elevation of 1000 to 3000 feet; not common.

Præcedente valde affinis similisque, sed sepalorum lateralium, calcaris petalorumque forma satis differt.—*Flores* rubro-purpurei, medio luteo-tincti. *Capsula* 1-3-spermæ.

17. **I. linearis**, Arn. l. c. p. 323; Walp. l. c. p. 473.—c.p. 372.

HAB. Adam's Peak, at an elevation of 4000 to 6000 feet.

18. **I. appendiculata**, Arn. l. c. p. 323; Walp. l. c. p. 473.

Var. *a.*—? *I. candida*, Lindl. Bot. Reg. new series, vol. xiii.; Misc. 85. n. 204; Walp. l. c. p. 475.—c.p. 1207.

Var. *β.*—c.p. 1208.

Var. *γ*; *foliis* majoribus, pilosioribus.—c.p. 1209.

HAB. Southern and central parts of the island, at an elevation of 2000 to 5000 feet.

Species valde variabilis. *Folia* ad caulis sæpissime simplicis apicem aggregata, vel sparsa,  $\frac{1}{2}$ -6 poll. longa, subglabra vel pilosa. *Flores* albidii vel pallide rosei, *sepalo anteriore* transverse rubro-striato. *Petalorum lobus posterior* parvus, lanceolatus, longe caudatus; *anterior* major, semicordatus, acuminatus.

19. **I. leucantha**, Thw.; herbacea, pusilla, simplex; *foliis* ad caulis apicem approximatis, lanceolatis, acuminatis, denticulatis, parce pilosis, versus basin petioloque subalato glanduloso-setigeris; *sepalo anteriore* in calcar æquilongum rectum conicum apice (siccum) subabrupte attenuatum productum; *petalorum lobo posteriore* parvo, lanceolato, *anterior* multo majore, inæqualiter 3-partito, apicibus rotundatis.—c.p. 3548.

HAB. Hinidoon and Kittool Galle.

A præcedente, cui valde similis affinisque, calcaris petalorumque forma præcipue differt. *Flores albi, sepalo anteriore rufo-guttato.*

20. **I. janthina**, Thw.; herbacea, pusilla, simplex, glaberrima; *foliis* 1-4, sæpissime caulem terminantibus, membranaceis, ovatis vel rotundatis, acuminatis, denticulatis, *petiolo* eglanduloso; *sepalis lateralibus* minoribus, oblongis, subacutis, *posteriore* rotundato, cucullato, *anteriore* infundibuliformi, in calcar æquilongum sensim attenuatum incurvatum producto; *petalorum lobis* subæqualibus, *posteriore* acuto, *anteriore* subrotundato, obtuso.—c.p. 3003.

HAB. Hinidoon and Pasdoon Corles, at no great elevation; rare.

*Folia* 1-4 poll. longa,  $\frac{3}{4}$ -3 poll. lata, *petiolo*  $\frac{1}{2}$ -2 poll. longo. *Pedicelli* pollicares, ad *pedunculi* elongati apicem in racemum brevem aggregati; *bracteis* ovalibus, persistentibus. *Flores* violacei, 6-8 lin. in exp. *Capsulæ* polyspermæ.

21. **I. acaulis**, Arn. l. c. p. 325.—? *I. rivalis*, Wight, Icon. t. 751.—c.p. 490 (2595).

HAB. Central Province, at an elevation of 3000 to 5000 feet, on wet rocks; abundant.

*Folia* rotundata, ovata vel lanceolata, *petiolo* rufescente æquilonga vel multo breviora, parce pilosa vel subglabra. *Scapi* rufescentes. *Flores* rubri, 1-1 $\frac{1}{4}$  poll. in exp., *calcare* æquilongo vel parum longiore. *Petalorum lobi* æquilongi; *anterior* latior, oblique truncatus vel emarginatus.

Dr. Wight's *I. rivalis*, figured with the anterior lobes of the petals bipartite, is probably only a form of the present species. *I. scapiflora*, Heyne, is described by Roxburgh as having a spur several inches in length, and therefore must certainly be different both from this and from the *I. scapiflora* of Wight's 'Icones,' t. 967.

## 2. HYDROCERA, Blume.

1. **H. triflora**, W. et A. Prod. i. 140, cum syn.—c.p. 1210.

HAB. Warmer parts of the island, in watery places; very abundant.

## XLVIII. PITTOSPOREÆ.

### 1. PITTOSPORUM, Sol.

1. **P. Ceylanicum**, Wight, Illustr. i. 173.—? *P. floribundum*, W. et A. Prod. i. 154. ? *Celastrus verticillatus*, Roxb. Fl. Ind. i. 391. ? *Senacia Napaulensis*, DC. Prod. i. 347.—c.p. 476.

HAB. Oova District, at an elevation of 3000 to 5000 feet. Nom. vulg. "Katteya-gass."

*Arbor* mediocris, 15-20-pedalis. *Folia* oblongo-lanceolata, cuneata, apice subacuta vel rotundata, glabra, 3-4 poll. longa,  $\frac{3}{4}$ -1 poll. lata, *petiolo* circiter  $\frac{1}{2}$  poll. longo. *Flores* albidæ. *Ovarium* pilosum. *Stylus* cylindricus, glaber. *Capsulæ* uniloculares, viridi-fuscescentes.

2. **P. tetraspermum**, W. et A. Prod. i. 154; Wight, Icon. t. 971.—c.p. 572 (685).

HAB. Central Province, at an elevation of 5000 to 7000 feet, on the banks of streams.

## XLIX. ZYGOPHYLLÆ.

### 1. TRIBULUS, Tourn.

1. **T. terrestris**, Linn. (Walp. Rep. ii. 822, cum syn.)—*T. lanuginosus*,

Linn.; DC. Prod. i. 704, cum syn.; Wight, Icon. t. 98; Illust. p. 164.—C.P. 1252.

HAB. North of the island, *Gardner*.

## L. XANTHOXYLACEÆ.

### 1. XANTHOXYLON, Linn.

1. **X. Rhetsa**, DC. (W. et A. Prod. i. 148, cum syn.)—C.P. 3490.

HAB. Central Province, *W. H. Wright, Esq.*; near Colombo, *W. Ferguson, Esq.* Nom. vulg. "Kattoo-keena-gass."

This tree so precisely corresponds with Roxburgh's description, that there can be no doubt of its being the *Fagara Rhetsa* of that author.

2. **X. triphyllum**, Juss. (Wight, Illust. i. 169; Icon. t. 204.)—*X. Zeylanicum*, DC. Prod. i. 728, cum syn. *Evodia triphylla*, DC. l. c. p. 724, cum syn.—C.P. 538, 2501.

Var.  $\beta$ ; *foliis* digitatis, 5-foliolatis.

HAB. Very abundant, up to an elevation of 5000 feet. Var.  $\beta$ . Near Ratnapoora. Nom. vulg. "Loonoo-ankenda-gass."

3. **X. tetraspermum**, W. et A. Prod. i. 148.—C.P. 386.

HAB. Central Province, at an elevation of 3000 to 5000 feet.

*Frutex* scandens, aculeatus. *Folia* impari-pinnata, rachi aculeata. *Foliola* 2-3-juga, cum impari, subsessilia, costulis sæpissime aculeatis. *Flores* parvi, abortu dioici, tetrameri, albidi, roseo-tincti. MASC. *Petala* oblonga, acuta, calycis segmentis multo longiora. *Stamina* in lobis disci ovarium abortivum eingentis inserta. *Ovarii* abort. lorum apices recurvato-uncinati.

### 2. TODDALIA, Juss.

1. **T. aculeata**, Pers. (W. et A. Prod. i. 149, cum syn.)—*T. floribunda*, Wall. Plant. As. Rar. ii. 17. t. 232; Wight, Illust. i. 169. t. 66.—C.P. 1250, 2413 (1251).

HAB. Very abundant, up to an elevation of 4000 feet. Nom. vulg. "Koodoomirris-wel."

A most variable plant, as regards the shape and size of the leaflets, the extent of aculeation, and the amount of the red pubescence with which the younger parts are clothed. The large-based aculei are common to the stems of all the varieties, and therefore furnish no character for their separation.

### 3. CYMINOSMA, Gærtn.

1. **C. pedunculata**, DC. (W. et A. Prod. i. 147, cum syn.; Wight, Illust. i. 165. t. 65.)—C.P. 1249.

HAB. Common, up to an elevation of 4000 feet. Nom. vulg. "Ankenda-gass."

*Nuculania* rotundata, 5-7 lin. in diam., albida.

### 4. AILANTHUS, Desf.

1. **A. Malabaricus**, DC. Prod. ii. p. 89, cum syn.; Wight, Icon. t. 1604.—C.P. 3456.

HAB. Near Colombo, *Mr. W. Ferguson*; district of the Seven Corles. Nom. vulg. "Koombaloo-gass," "Wal-biling-gass."

*Brucea Sumatrana*, Roxb., is become a troublesome weed, but it does not appear to be indigenous.

## LI. SIMARUBEÆ.

## 1. SAMADERA, Gærtn.

1. **S. Indica**, Gærtn. (W. et A. Prod. i. 151, cum syn.)—C.P. 2766.

HAB. South of the island; not uncommon. Nom. vulg. "Samadara-gass."

The root of this plant is used as a medicine by the Cinghalese, and so also is the fruit.

## LII. OCHNACEÆ.

## 1. OCHNA, Linn.

(Nom. vulg. "Mal-kaara-gass.")

1. **O. squarrosa**, Linn. (Wight, Illust. i. 172. t. 69; Roxb. Cor. i. t. 89.)—C.P. 1223.

HAB. Batticaloa and Jaffna, *Gardner*.

2. **O. cordata**, Thw.; *foliis* ellipticis vel obovatis, obtusis, dentatis, basi cordatis; *racemis* ramulos foliosos terminantibus, brevibus, 3-6-floris; *sepalis* 3-5; *petalis* carpellisque circiter 7.—C.P. 1222.

HAB. Jaffna, *Gardner*.

*Folia* 1-2 poll. longa,  $\frac{1}{2}$ - $1\frac{1}{4}$  lata, *petiolo* circiter 1 lin. longo. *Pedicelli* pollicares, prope basin articulati. *Petala* rotundata, longe unguiculata,  $\frac{3}{4}$  poll. longa. *Antheræ* filamentis plus quam duplo longiores. *Stigma* capitatum.

But for the cordate base of the leaves I should have referred this, without much doubt, to the *O. obtusata*, DC., but that author would scarcely have passed over, unnoticed, so striking a characteristic of the present species. I have not sufficient materials to give so detailed a description as would be desirable.

3. **O. Moonii**, Thw.—*O. squarrosa*, Moon's Cat.; *foliis* lanceolatis, utrinque angustatis, denticulatis, *venis* *venulisque* prominentibus; *racemis* 6-12-floris, ramulos inferiores breves denudatos terminantibus; *pedicellis* floribus subduplo longioribus, ad partem inferiorem tertiam vel quartam articulatis.—C.P. 1224.

Var.  $\beta$ ; *foliis* minoribus, utrinque subobtusis.—C.P. 2554.

HAB. Warmer parts of the island, up to an elevation of 2000 feet.

Var.  $\beta$ . Jaffna, *Gardner*.

*Folia* 3-7 poll. longa,  $1\frac{1}{4}$ -2 poll. lata, *petiolo* circiter 1 lin. longo. *Flores*  $4\frac{1}{2}$ -5 lin. longi. *Petala* 5, oblonga, *sepalis* æquilonga. *Antheræ* filamentis subduplo longiores. *Carpella* 5-7.

4. **O. rufescens**, Thw.; *foliis* lanceolatis, acuminatis, argute dentatis, nitidis, conspicue et minutissime venulosis, basi angustatis; *racemis* 6-12-floris, ramulos inferiores breves denudatos terminantibus; *pedicellis* floribus subtriplo longioribus, prope basin articulatis.—C.P. 3455.

HAB. Hinidoon Corle, at an elevation of about 1000 feet.

*Folia* 4-6 poll. longa, circiter  $1\frac{1}{2}$  poll. lata, *petiolo* 2 lin. longo, juniora rufescentia. *Flores* 3- $3\frac{1}{2}$  lin. longi. *Carpella* 7-8.

Closely allied to the preceding, but distinguished by its much smaller flowers, and the relatively much shorter inferior articulation of the pedicel. In many respects it would seem to resemble the *O. nitida*, Thunb., also said to occur in Ceylon, but the character of the inflorescence is different.

5. **O. Wightiana**, Wall. (W. et A. Prod. i. 152: ? Wight, Icon.

t. 223.)—*O. Heyneana*, W. et A., l. c. *O. Walkerii*, Planchon, Hook. Lond. Journ. v. 653; Walp. Ann. i. 180.—c.p. 1221.

HAB. Hot, drier parts of the island, at no great elevation.

*Folia* 1-2 poll. longa,  $\frac{1}{2}$ - $\frac{3}{4}$  lata, *petiolo* 1 lin. longo. *Pedicelli* 4-7 lin. longi, prope medium articulati. *Flores*  $3\frac{1}{2}$ -4 lin. longi.

## 2. GOMPHIA, Schreb.

1. *G. angustifolia*, Vahl, (W. et A. Prod. i. 153, cum syn.; et, fere sine dubio, *Walkera serrata*, Willd. DC. Prod. i. 737, cum syn.)—c.p. 2412.

HAB. Common, up to an elevation of 3000 feet. Nom. vulg. "Bo-kaara-gass."

The wood of this tree, which grows to the height of thirty feet, is useful for building purposes.

## LIII. STAPHYLEACEÆ.

### 1. TURPINIA, Ventn.

1. *T. Nepalensis*, Wall. (W. et A. Prod. i. 156; Wight, Icon. t. 972.)—c.p. 218 (563, 569).

Var.  $\beta$ . *montana*; *floribus* majoribus, paucioribus.—? *T. pomifera*, DC. Prod. ii. p. 3, cum syn.

HAB. Var. *a*. Common in the warm, moister parts of the island. Var.  $\beta$  occurs abundantly in more elevated situations, up to 6000 feet. Nom. vulg. "Kankombala-gass," "Attakirilla-gass."

*Stipulae interpetiolares* triangulares, cito deciduæ. *Nuculania* firma, subglobosa, viridia, nigro-purpurascens, deinde cinerascens,  $\frac{1}{2}$ -1 poll. in diam.

The extreme forms of this plant might easily be taken for two distinct species; but this is another instance of an elevated humid locality producing a form with fewer and larger flowers than are exhibited in the same species when growing nearer the level of the sea.

## LIV. CELASTRACEÆ.

### Sect. 1. *Ovarium liberum vel subliberum.*

#### 1. PLEUROSTYLIA, W. et A.

1. *P. Wightii*, W. et A. Prod. i. 157, cum syn.; Wight, Icones, t. 155.—c.p. 329.

HAB. Hot, drier parts of the island, up to no great elevation.

#### 2. MICROTROPIS, Wall.

*Semina* exarillata; *testa* colorata.

1. *M. Wallichiana*, Wight, MSS.; *foliis* obovato-vel oblongo-lanceolatis, acuminatis, petiolatis; *floribus* axillaribus ramulinisque, subsessilibus, confertis.—c.p. 43 (202, 374, 402).

HAB. Forests of the Central Province and Saffragam, at an elevation of 2000 to 5000 feet; abundant.

*Arbor mediocris. Folia glabra, 2½-6 poll. longa, 1-3 poll. lata, petiolo 2-4 lin. longo.*

This would appear to differ from *M. garciniaefolia*, Wall. (Wight, Icon. t. 761), principally in its subsessile, more numerous flowers, and it will probably prove to be merely a variety of that species.

2. **M. ramiflora**, Wight, Icones, t. 977.—C.P. 148, 1228.

Var.  $\beta$ ; *foliis minoribus, oblongo-lanceolatis, basi angustatis.*

HAB. Abundant at Newera Ellia and other of the more elevated parts of the Central Province. Var.  $\beta$  occurs in rather dry situations.

*Folia 1-3 poll. longa, ½-2 poll. lata, petiolo 1-2 lin. longo. Pedunculi ½-2 lin. longi.*

This, like the preceding, is a very variable species, the size, form, and texture of the leaves being modified according to the kind of situation in which it occurs. Var.  $\beta$  bears a considerable general resemblance to *M. microcarpa*, Wight, Icones, t. 975, but I have never seen the Ceylon plant with the peduncles and pedicels so elongated as represented in that figure.

### 3. KURRIMIA, Wall., Arnott.

*Bhesa*, Ham., Arn., Endlich. *Rhesa*, Walpers.

*Ovarium liberum, 2-loculare; loculis 2-ovulatis. Ovula basi affixa, erecta, collateralia. Capsula 1-locularis, bivalvis, 1-2-sperma. Semina erecta, arillo carnosio.*

1. **K. Ceylanica**, Arn. Pug. x. 328.—C.P. 1225.

Var.  $\beta$ . *montana; paniculis minus elongatis; floribus majoribus.*—C.P. 409.

HAB. Var.  $\alpha$ . In warm, moister parts of the island. Var.  $\beta$ . Central Province, up to an elevation of 5000 feet. Nom. vulg. "Palang," "Hoorakandoo," and "Alareya-gass."

*Arbor magna. Ramuli teretes, glabri; juniorum apices acuti, stipulis magnis circumvolutis cito deciduis tecti. Folia glabra, 3-6 poll. longa, ½-3 poll. lata, petiolo ½-1½ poll. longo. Flores parvi, pallide virides. Capsulae oblongae, carnosocoriaceae, ¾ poll. longae, rufescentes. Semina oblonga, fere ad apicem arillo carnosio albido inclusa; testa membranacea, brunnea.*

### 4. CELASTRUS, Linn.

1. **C. paniculata**, Willd. (W. et A. Prod. i. 158, cum syn.; Wight, Illust. i. 172. t. 72; Icones, t. 158.)—C.P. 1232.

HAB. Common up to an elevation of 2000 feet.

Sect. 2.—*Ovarium disco immersum et cum eo consolidatum.*

### 5. CATHA, Försk.

1. **C. emarginata**, G. Don.—*Celastrus emarginatus*, Willd.; W. et A. Prod. i. p. 160, cum syn.—C.P. 1235.

HAB. Hot, drier parts of the island. Nom. vulg. "Deya-eringoo."

2. **C. fruticosa**, Thw.; *foliis parvis, oblongo-spathulatis vel subtrapezoides, crenato-serratis, glabris, petiolatis, subtus prominenti-reticulatis; corymbis axillaribus, multifloris, folio brevioribus.*—C.P. 3386.

HAB. Banks of the Mahawelliganga, near Nawalapittya, in the Central Province.

*Frutex* inermis, 8-10-pedalis. *Ramuli* angulosi. *Folia*  $\frac{3}{4}$ -1 $\frac{1}{2}$  poll. longa, 4-7 lin. lata, *petiolo* circiter 1 lin. longo. *Flores* parvi, 1 lin. in exp. *Petala* 5, albida. *Stylus* brevis. *Stigmata* 3, brevia, recurvata. *Capsulæ* 3-loculares, pisi magnitudine, rubiginosæ, transverse striatæ. *Semina* coccinea, nitida; *arillo* parvo, albido.

## 6. EVONYMUS, Linn.

1. **E. revolutus**, Wight, *Illust.* i. 178; Walp. *Rep.* i. 531.—C.P. 579.

HAB. Newera and other very elevated parts of the island.

*Arbor* mediocris. *Folia* opposita vel terna, sæpe rotundata. *Flores* virides, purpurascens.

2. **E. dichotomus**, Heyne, (*W. et A. Prod.* i. 160, cum syn.)—C.P. 171.

HAB. Galagama, on the banks of streams.

3. **E. Walkerii**, Wight, l. c.; Walpers, l. c.—*E. Zeylanicus*, Moon, *Cat.* p. 17.—C.P. 478.

HAB. Very abundant in the southern and central parts of the island, up to an elevation of 3000 feet.

Closely allied to the preceding, from which it differs principally in its shorter petioles and flower-stalks.

## 7. GLYPTOPETALUM, Thw.

Hook. *Journ. of Bot.* vol. viii. (1856) p. 267. t. 7 B.

*Flores* tetrameri. *Petala* superne bifoveolata. *Antherarum loculi* ad apicem connectivi maximi innati siti, basi divergentes. *Ovarium* pyramidato-tetragonum, 4-loculare. *Ovula* in loculis singula, ab apice pendula. *Capsulæ* rotundatæ, 1-4-spermæ. *Semina* arillata.

1. **G. Zeylanicum**, Thw. l. c.—C.P. 589.

HAB. Central Province, up to an elevation of 3000 feet.

*Folia* opposita, lanceolata, serrata, glabra, 4 $\frac{1}{2}$ -6 poll. longa, 1 $\frac{1}{2}$ -2 $\frac{1}{2}$  poll. lata; *petiolo*  $\frac{1}{4}$  poll. longo. *Cymarum* supra-axillarium *pedunculi* 1-1 $\frac{1}{2}$  poll. longi; *pedicelli* divaricati,  $\frac{1}{2}$  poll. longi. *Flores* albido-virescentes, 4 lin. lati. *Semina*  $\frac{1}{2}$  poll. longa, ovoidea; *arillo* rubro ultra medium attingente; *testa* flavo-rufescente.

## 8. ELÆODENDRON, Jacq.

1. **E. glaucum**, Pers. (*Roxb. Fl. Ind.* (Wallich) ii. 432, cum syn.)—C.P. 1227.

Var.  $\beta$ . *montanum*; *floribus nuculanisque* majoribus.—C.P. 2520.

HAB. Var.  $\alpha$ . Hot, drier parts of the island. Var.  $\beta$ . Central Province, at an elevation of 3000 to 4500 feet.

I can find nothing to distinguish the two varieties of this species except the larger flowers and fruit of var.  $\beta$ , and the petioles of this form are usually somewhat shorter than in var.  $\alpha$ .

## LV. RHAMNACEÆ.

### 1. ZIZYPHUS, Tourn.

1. **Z. rugosa**, Lam. (*W. et A. Prod.* i. 162, cum syn.; Wight, *Icon.* t. 339.)—C.P. 2719.



HAB. Very abundant, up to an elevation of 2000 feet. Nom. vulg. "Maha-erramina."

2. *Z. xylopyra*, Willd. (W. et A., l. c. cum syn.)—c.p. 1243.

HAB. Hot, drier parts of the island.

3. *Z. Jujuba*, Lam. (W. et A., l. c. ; Wight, Icon. t. 99.)—c.p. 1242.

HAB. Anooradhapoorra, *Gardner*.

4. *Z. Cœnopia*, Mill. (W. et A., l. c. p. 163.)—c.p. 1240.

HAB. Common, up to an elevation of 3000 feet. Nom. vulg. "Erramina-wel."

5. *Z. lucida*, Moon, Cat. p. 17; *foliis* ovato-lanceolatis, crenato-serrulatis, obtuse acuminatis, minute apiculatis, basi obliquis, 3-5-nerviis, transverse parallelo-venulosis, glabris, petiolo nervisque pubescentibus; *cymis* folio multo brevioribus, tomentosis; *stylis* 3-4, ad medium coalitis.—? *Z. horrida*, Roth, nov. sp. p. 159.—c.p. 1241.

HAB. Matele, *Moon*; Kornegalle District, abundant.

*Fruticosa*. *Ramuli* teretes, nigrescentes, juniores tomentosi; *aculeis* solitariis vel binis. *Folia* 2-3 poll. longa,  $\frac{3}{4}$ -1 $\frac{1}{4}$  poll. lata, *petiolo* 3 lin. longo. *Ovarium* 3-4-loculare. *Nuculaniam* globosa, circiter 8 lin. in diam., carnosam, viridi-purpurascentiam, 2-4-sperma; *putamine* osseo, ruguloso.

It is difficult to surmise what can be the *Rhamnus Napeca*, Linn. Fl. Zeyl. 87, the description of which agrees tolerably well with the characters of the present plant, except as regards the number of styles, which seems decisive against their being identical.

## 2. BERCHEMIA, Neck.

1. *B. parviflora*, Wall. (W. et A. Prod. i. 163, cum syn.; Wight, Icones (19), t. 513.)—c.p. 2477.

HAB. Maturate District, in the Central Province, up to an elevation of 3000 feet; not common.

In my specimens the leaves are oblong-lanceolate, and not cordate at the base as in Dr. Wight's figure, but in other respects the characters correspond.

## 3. VENTILAGO, Gærtn.

1. *V. Maderaspatana*, Gærtn. (W. et A. Prod. i. 164, cum syn.; Wight, Icones, t. 163.)—c.p. 1236.

HAB. Common in the hot, drier parts of the island, up to an elevation of 1500 feet.

## 4. RHAMNUS, Linn.

1. *R. Wightii*, W. et A. Prod. i. 164, cum syn.; Wight, Icones, t. 159.—c.p. 2525.

HAB. Newera Ellia and other of the most elevated parts of the Central Province.

2. *R. Arnottianus*, Gardn. MSS.; *foliis* glabris, ovato-lanceolatis, crenatis, glandulose serrulatis, obtuse acuminatis, superne depresso-reticulatis; *pedunculis* unifloris, axillaribus, ternis, puberulis, *petiolo* superne puberulo longioribus; *petalis* nullis.—c.p. 201.

HAB. Most elevated parts of the Central Province.

*Arbor* parva. *Ramuli* juniores puberuli. *Folia* 1 $\frac{1}{2}$ -2 $\frac{1}{2}$  poll. longa,  $\frac{3}{4}$ -1 $\frac{1}{4}$  poll.

*lata*, *petiolo* 2-4 lin. longo. *Stipulæ* lineari-lanceolatæ, 2½ lin. longæ, basi interne glandulosæ, cito deciduæ; *glandulis* minutis, erectis, clavatis, rubris, persistentibus. *Calycis laciniæ* 4, acutæ, externe pilosæ. *Stamina* 4, filamentis brevibus. *Styli* 2-3, basi coaliti. *Nuculania* pisi parvi magnitudine, nigro-purpurea.

### 5. SCUTIA, Comm.

1. **S. Indica**, Brongn. (W. et A. Prod. i. 165, cum syn.; Wight, *Illustr.* 180. t. 73.)—C.P. 1233.

HAB. Abundant in the hot, drier parts of the island.

### 6. COLUBRINA, Rich.

1. **C. Asiatica**, Brongn. (W. et A. Prod. i. 166, cum syn.; Wight, *Illustr.* 180. t. 74.)—C.P. 1239.

HAB. North of the island, *Gardner*.

### 7. GOUANIA, Jacq.

1. **G. microcarpa**, DC. (W. et A. Prod. i. 167, cum syn.)—C.P. 1238.

HAB. Common, up to an elevation of 2000 feet.

## LVI. TEREBINTHACEÆ.

### 1. MANGIFERA, Linn.

1. **M. Indica**, Linn. (W. et A. Prod. i. 170, cum syn.)—C.P. 1256, 2614.

HAB. Abundant, up to an elevation of 3000 feet. Nom. vulg. "Amba-" and "Attamba-gass."

### 2. ANACARDIUM, Rottb.

1. **A. occidentale**, Linn. (W. et A. Prod. i. 168, cum syn.)

HAB. Naturalized in some of the warmer parts of the island. Nom. vulg. "Cadjoo-gass."

### 3. SEMECARPUS, Linn.

#### Sect. 1. *Paniculæ ramulos terminantes.*

1. **S. subpeltata**, Thw.; *foliis* magnis, ovato-lanceolatis, acuminatis, glabris, coriaceis, nitidis, subpeltatis; *drupis* transverse oblongis, parce compressis, dimidio inferiore *hypocarpio*\* magno carnosio subreniformi tectis.—C.P. 3004.

HAB. Singhe-rajah and other forests between Ratnapoora and Galle. Nom. vulg. "Maha-badoolla-gass."

*Arbor* magna, 30-40-pedalis. *Folia* 10-15 poll. longa, 4-6½ poll. lata; *petiolo*

\* The fleshy structure on which the drupe in this genus rests, and for which I propose the name of *hypocarpium*, can hardly be called a "torus incrassatus," as it is rather an enlargement of the base of the calyx.

crasso,  $1\frac{1}{2}$ –2 poll. longo. *Paniculæ* magnæ, ramis patentibus. *Drupæ* circiter  $\frac{3}{4}$  poll. in alt.,  $1\frac{1}{4}$  poll. in max. lat.

2. **S. Gardneri**, Thw.; *foliis* lanceolatis vel oblongo-lanceolatis, acuminatis, glabris; *petiolo* longiusculo; *drupis* oblique oblongis, parce compressis, dimidio inferiore *hypocarpio* magno carnosio cupuliformi tectis.—C.P. 1257, 2940.

HAB. Very common in the Central Province, up to an elevation of 3000 feet. Nom. vulg. "Badoolla-gass."

*Arbor* mediocris. *Folia*  $4\frac{1}{2}$ –13 poll. longa,  $1\frac{1}{2}$ –4 poll. lata; *petiolo* crassiusculo,  $\frac{3}{4}$ –2 poll. longo. *Paniculæ* folio subæquilongæ. *Drupæ* circiter 9 lin. longæ, *hypocarpio* circiter 7 lin. lato.

3. **S. obscura**, Thw.; *foliis* glabris, ovato-oblongis, obtusis, basi cuneatis, superne nitidis; *petiolo* brevi, vix incrassato; *drupis* (immaturis) oblongis, dimidio inferiore *hypocarpio* cupuliformi carnosio tectis.—C.P. 2556.

HAB. Deltotte, in the Central Province, at an elevation of 3000 feet.

*Arbor* mediocris. *Folia* 3–4 poll. longa,  $1\frac{1}{2}$ – $1\frac{3}{4}$  poll. lata, *petiolo* circiter 3 lin. longo. *Paniculæ* folio subæquilongæ.

4. **S. oblongifolia**, Thw.; *foliis* glabris, oblongis, apice rotundatis vel obscure obtuse acuminatis, basi cuneatis, *petiolo* basi incrassato; *drupis* oblongis, parce obliquis compressisque, apiculatis, *hypocarpio* (sicco) angustiore poculiformi suffultis.—? *S. ?Zeylanica*, Blume, Mus. Bot. Lugd.-Bat. p. 189.—C.P. 1258.

HAB. Hot, drier parts of the island, up to an elevation of 3000 feet; common. Nom. vulg. "Badoolla-gass."

*Arbor* mediocris. *Folia*  $3\frac{1}{2}$ –9 poll. longa, 1– $2\frac{1}{2}$  poll. lata, *petiolo* 4–6 lin. longo. *Drupæ* nuce avellana parum majores.

This species a good deal resembles *S. obscura*, but differs in its less coriaceous and paler leaves, and much smaller hypocarp.

5. **S. nigroviridis**, Thw.; *foliis* glabris, obovatis vel oblongis, obtuse brevi-acuminatis, basi angustatis, *petiolo* basi incrassato; *drupis* rotundatis vel obovatis, obliquis, compressis, uncinato-apiculatis, *hypocarpio* (sicco) angustiore poculiformi compresso suffultis.—C.P. 631, 2444 (641, 2586).

HAB. Central Province, at an elevation of 2000 to 4000 feet.

*Arbor* mediocris. *Folia* intense viridia,  $2\frac{1}{2}$ –6 poll. longa,  $1\frac{1}{2}$ – $2\frac{1}{2}$  poll. lata, *petiolo* 3–8 lin. longo. *Drupæ* 8–10 lin. in max. diam.

Distinguished from the preceding species by its differently-shaped and darker-coloured leaves and larger, more compressed drupes. From *S. obscura* it differs principally in the shape of the fruit and the smaller hypocarp.

6. **S. acuminata**, Thw.; *foliis* glabris, ovato-vel oblongo-lanceolatis, longe acuminatis, basi angustatis, *petiolo* parum incrassato; *drupis* obovatis, compressis, valde obliquis, uncinato-acuminatis, *hypocarpio* (sicco) multum angustiore poculiformi elongatulo suffultis.—C.P. 2676.

HAB. Forests in the Ratnapoora, Galle, and Ambagamowa Districts, at no great elevation.

*Arbor* mediocris. *Folia* 4–12 poll. longa,  $1\frac{1}{2}$ – $2\frac{1}{4}$  lata, *petiolo* 4–10 lin. longo. *Drupæ* 1 poll. longæ,  $\frac{3}{4}$  poll. in max. diam., *hypocarpio*  $\frac{1}{2}$  poll. longo.

Resembling *S. Gardneri* in many respects, but distinguished from it by its less coriaceous leaves, smaller panicles, and very differently-shaped fruit.

7. **S. coriacea**, Thw.; *foliis* glabris, obovatis vel obovato-oblongis,

valde coriaceis, apice rotundatis vel obscure obtuse acuminatis, basi plus minus angustatis, *petiolo* costaque robustis; *paniculis* folio brevioribus, rache brevi, ramis interrupte spicæformibus; *drupis* obcordatis, obliquis, compressis, mucronulatis; *hypocarpio* (sicco) mediocri, poculiformi.—C.P. 313, 635.

HAB. Central Province, at an elevation of 5000 to 7000 feet.

*Arbor* mediocris. *Folia* 3½–7 poll. longa, 1½–3½ poll. lata, *petiolo* 2–5 lin. longo. *Drupæ* 8 lin. in alt., 11 lin. in max. lat., *hypocarpio* 4½ lin. longo.

8. **S. parvifolia**, Thw.; *foliis* glabris, ovato-oblongis, abrupte obtuse acuminatis vel apice rotundatis, basi angustatis, coriaceis; *petiolo* superne sulcato, non incrassato; *paniculæ* ramis racemiformibus, paucifloris; *drupis* (immaturis) elongato-conicis, uncinato-acuminatis, parce obliquis, *hypocarpio* subæquilato poculiformi suffultis.—C.P. 3444.

HAB. Hinidoon Corle, in the Galle District. Nom. vulg. "Heenbadoolla-gass."

*Arbor* mediocris. *Folia* 1½–3½ poll. longa, ¾–1½ poll. lata, *petiolo* 3–6 lin. longo.

9. **S. Moonii**, Thw.; *foliis* magnis, glabris, coriaceis, lanceolatis, brevi-acuminatis, basi angustatis, brevi-petiolatis; *drupis* (immaturis) obliquis, oblongis, compressis, apiculatis, *hypocarpio* mediocri.—C.P. 3338.

HAB. South of the island, at no great elevation.

*Arbor* mediocris. *Folia* 6–12 poll. longa, 2–5 poll. lata, *petiolo* 1–3 lin. longo. *Paniculæ* elongatæ, multifloræ.

10. **S. obovata**, Moon, Cat. p. 22; *foliis* glabris, superne nitidis, obovatis, apice rotundatis vel abrupte obtuse acuminatis, ad basin angustatam rotundatis vel subcordatis, brevi-petiolatis; *drupis* (immaturis) oblongis, parce obliquis, apiculatis; *hypocarpio* (sicco) angustiore, elongatulo.—C.P. 3339.

HAB. Caltura, Moon; near Ratnapoora.

*Arbor* mediocris. *Folia* 3–7 poll. longa, 1¼–5 poll. lata; *petiolo* 1–2 lin. longo, incrassato.

11. **S. pubescens**, Thw.; *foliis* lanceolatis, acuminatis, basi subrotundatis, superne (costa excepta) glabris, subtus cum *costa* *petioloque* incrassato pubescentibus; *paniculis* elongatis; *drupis* parvis.—C.P. 163.

HAB. Ratnapoora District, at no great elevation.

*Arbor* parva. *Ramuli* elongati, teretes, pubescentes. *Folia* 5–13 poll. longa, 1½–3 poll. lata, *petiolo* ½–1 poll. longo. *Drupæ* oblongæ, parce obliquæ, circiter 6 lin. longæ, 3 lin. latæ, *hypocarpio* brevi carnosio cupuliformi subæquilato suffultæ.

## Sect. 2. *Paniculæ caulinae*.

12. **S. marginata**, Thw.; *foliis* magnis, lanceolatis, conspicue membranaceo-marginatis, acuminatis, ad basin angustatis cordatisque, *petiolo* valde incrassato; *paniculis* caulinis, ramosis, delicatulis, pilosiusculis.—C.P. 2677.

Var. *a. glabra*; *foliis* glabris.

Var. *β. hirsuta*; *foliis* subtus pilis patentibus, rigidis hirsutis.

HAB. Damp forests in the Galle, Ratnapoora, and Ambagamowa Districts.

*Frutex* erectus, 6–8-pedalis, parce ramosus. *Folia* 6–14 poll. longa, 2½–5 poll. lata, *petiolo* ¼–1½ poll. longo. *Drupæ* ovoideæ, obliquæ, parce compressæ, circiter

4 lin. longæ, rufescentes, *hypocarpio* æquilato dimidio breviorè poculiformi coccineo suffultæ.

#### 4. GLYCYCARPUS, Dalzell.

Hook. Journ. of Bot. vol. ii. (1850) p. 38.

1. **G. racemosa**, Dalz. l. c.; Hook. Icon. Pl. ix. t. 842.—c.p. 1260.

HAB. Very common, up to an elevation of 3000 feet. Nom. vulg. "Bala-gass."

*Drupæ* maturæ nigro-purpureæ. *Cotyledones* sæpissime purpureo-tinctæ.

#### 5. BUCHANANIA, Roxb.

1. **B. angustifolia**, Roxb. (W. et A. Prod. i. 169, cum syn.; Wight, Icon. t. 101.)—c.p. 471, 1255.

HAB. North of the island, *Gardner*; Galagama.

*Buchanania*? *Zeylanica*, Bl. Mus. Bot. Lugd.-Bat. p. 185, is probably a state of *Mangifera Indica*, Linn., judging from the native name he has given of the plant.

#### 6. CAMPNOSPERMA, Thw.

Hook. Journ. of Bot. vol. vi. p. 65. t. 1 A. Walpers, Ann. vol. iv. p. 444.

1. **C. Zeylanicum**, Thw., l. c.—c.p. 246.

HAB. Saffragam and Ambagamowa Districts, generally on the banks of streams, up to an elevation of 1000 feet.

#### 7. ODINA, Roxb.

1. **O. Wodier**, Roxb. (W. et A. Prod. i. 171, cum syn.; Wight, Icones, t. 60.)—c.p. 1161.

HAB. Warmer parts of the island, up to an elevation of 1500 feet. Nom. vulg. "Hig- or Hik-gass."

A considerable quantity of gum exudes from the trunk of this tree, which the natives use as a medicinal plaster.

#### 8. EVIA, Comm.

Blume, Mus. Bot. Lugd.-Bat. p. 234. *Spondias*, Auct. (*partim*).

1. **E. amara**, Comm. (Blume, l. c. cum syn.)—c.p. 1263.

HAB. Common, up to an elevation of 2000 feet. Nom. vulg. "Ambaralla-gass."

#### 9. PROTIUM, Burm.

1. **P. caudatum**, W. et A. Prod. i. 176.—c.p. 1261.

HAB. Hot, drier parts of the island, at no great elevation.

#### 10. SCUTINANTHE, Thw.

Hook. Journ. of Bot. vol. viii. (1856) p. 266. t. 8 B.

*Flores* abortu dioici. *Calyx* cupuliformis, carnosus-coriaceus, in seg-

mentis 5 erectis valvatis ad medium divisus, persistens. *Corolla* cum calyce æquilonga; *petalis* 5, erectis, valvatis, carnosocoriaceis, persistentibus. *Stamina* 10; *filamentis* in anulum parte inferiore cum *disco* glanduloso interno et basi calycis corollæque consolidata cohærentibus, apice liberis; *antheris* oblongis, introrsis, dorso affixis. *Ovarium* biloculare, loculis biovulatis; *ovulis* collateralibus, axi affixis, amphitropis. *Stylus* brevissimus. *Stigma* disciforme. *Nuculania* carnosae; *putamine* mediocri, osseo, abortu monospermo. *Semen* pendulum; *testa* membranacea; *cotyledonibus* magnis, foliaceis, integris, plicato-rugosis; *radicula* parva.

1. **S. brunnea**, Thw., l. c.—c.p. 1149.

HAB. Central Province, at an elevation of 2000 to 3000 feet. Nom. vulg. "Maha-booloomora-gass."

*Arbor* 50–60-pedalis. *Ramuli*, cum foliis junioribus, rufo-tomentosi. *Folia* imparipinnata, 10–20 poll. longa, exstipulata; *petiolo* superne sulcato, basi tumido, striato; *foliolis* 5–11, oblongis, parce obliquis, acuminatis, integris, 4–8 poll. longis, 2–3 poll. latis, rufescentibus, *petiolulo* 4 lin. longo. *Paniculæ* axillares, multifloræ, tomentosæ. *Flores* 3 lin. lati. *Nuculania* oblonga, rufo-tomentosa, utrinque attenuata, pollicaria.

## II. CANARIUM, Linn.

1. **C. Zeylanicum**, Blume, Mus. Bot. Lugd.-Bat. p. 218, cum syn.—*C. balsamiferum*, Moon, Cat. p. 68.—c.p. 2649.

HAB. Warm, moister parts of the island, up to an elevation of 1500 feet. Nom. vulg. "Kaakoona-gass."

*Stipulæ* majusculæ, petiolares, a ramulo parum distantes, brevi-stipitatae, obliquæ, rotundato-sagittatae, cito deciduæ.

The resinous balsam, which exudes copiously from the trunk of this tree, is used by the natives for burning, after being mixed with paddy-chaff; and the smoke given off is reputed to drive away snakes from the domicile.

## LVII. CHAILLETIACEÆ.

### 1. MOACURRA, Roxb.

1. **M. gelonioides**, Roxb. Fl. Ind. ii. 69.—c.p. 1245.

Var.  $\beta$ ; *foliis* subtus plus minus tomentosis.—c.p. 1231.

HAB. Common in the moister parts of the island, and up to an elevation of 3000 feet.

I can find nothing to distinguish var.  $\beta$  from the other form of this species but its usually greater luxuriance of growth, and the tomentum, which varies exceedingly in amount, upon the ramuli and the under surface of its leaves.

## LVIII. HOMALINEÆ.

### 1. BLACKWELLIA, Comm.

1. **B. Ceylanica**, Gardner, Calcutta Journ. of Nat. Hist. vol. vii. p. 452.—*B. tetrandra*, Wight, Icones, t. 1851.—c.p. 388.

HAB. Moister parts of the island, up to an elevation of 3000 feet. Nom. vulg. "Leeyang-gass."

*Arbor* 30–40-pedalis; *ramulis foliisque junioribus rubris. Florum spicæ sæpissime valde elongatæ, pendulæ.*

The wood of this tree is strong, and useful for building purposes.

## LIX. CONNARACEÆ.

### 1. ROUREA, Aub.

1. **R. santaloides**, W. et A. (Blume, Mus. Bot. Lugd.-Bat. p. 262, cum syn.)—C.P. 749.

HAB. Common, up to an elevation of 3000 feet. Nom. vulg. "Kerindi-wel."

### 2. CONNARUS, Linn.

1. **C. monocarpus**, Linn. (Blume, Mus. Bot. Lugd.-Bat. p. 265, cum syn.)—C.P. 323.

HAB. Very abundant in the hot, drier parts of the island. Nom. vulg. "Radaleya-gass."

2. **C. Championii**, Thw.; *ramulis foliisque novellis cum paniculis rufo-nigro-furfuraceis; foliis 5–7-foliolatis; foliolis ellipticis, acuminatis; folliculis transverse rugulosis, glabris, subtruncato-obovatis, versus basin angustatis.*—C.P. 2399.

HAB. Central Province, up to an elevation of 4000 feet.

*Frutex scandens. Foliola*  $2\frac{1}{2}$ –4 poll. longa,  $1\frac{1}{2}$ – $2\frac{1}{2}$  poll. lata; *petiolulo* incrassato,  $2\frac{1}{2}$  lin. longo. *Panicularum* rami ultimi racemiformes. *Pedicelli* breves. *Calycis lacinia* oblongæ, externe furfuraceæ. *Petala* longiora, albida, rufo-punctata, parce pilosa. *Stamina* 10, basi coalita; 5 antherifera, cum totidem plus quam dimidio brevioribus, antheris effœtis alternantia. *Ovarium* rufo-pilosum.

3. **C. unifoliolatus**, Thw.; *foliis glabris, petiolatis, 1-foliolatis; foliolo elliptico vel ovato, abrupte acuminato, reticulato; petiolulo brevi, tumido, rugulosulo; paniculis parvis, axillaribus, 1-, 2-, vel 3-nis, racemiformibus, paucifloris.*—C.P. 2443.

HAB. Central Province, at an elevation of 3000 to 4000 feet; rather rare.

*Arbor* mediocris, 20–30-pedalis; *ramulis petiolisque nigricantibus; novellis fusco-pilosis. Foliola*  $1\frac{3}{4}$ – $3\frac{1}{2}$  poll. longa,  $1$ – $1\frac{3}{4}$  poll. lata; *petiolulo* 1 lin. longo, cum *petiolo* 3 lin. longo basi tumidulo articulato. *Paniculæ* 4–7-floræ, pallide fusco-hirsutæ; *bracteolis* linearibus, deciduis. *Flores* abortu dioici, pallide virides, circiter 3 lin. in exp., brevi-pedicellati, cum *paniculæ* ramorum apicibus articulati. *Calycis lacinia* acutæ, *petalis* externe pilosis plus quam dimidio breviores. *Stamina* basi coalita; 5 antherifera, cum totidem brevibus anatheris dentiformibus alternantia. *Ovarium* strigosum, in *stylum* brevem attenuatum; *stigmatibus* dilatato. *Folliculi* (immaturi) fulvo-tomentosi, arcuato-falcati, acuti, basi in stipitem longum attenuati.

## LX. LEGUMINOSÆ.

## SUBORD. 1. PAPILIONACEÆ.

## 1. HEYLANDIA, DC.

1. *H. latebrosa*, DC. (W. et A. Prod. i. 180, cum syn.)—c.p. 1275.  
HAB. Abundant in the Badulla District.

## 2. CROTALARIA, Linn.

(Nom. vernac. "Andanahiriya.")

1. *C. biflora*, Linn. (W. et A. Prod. i. 190, cum syn.)—c.p. 3325.  
HAB. Hot, drier parts of the island, *Gardner*.
2. *C. prostrata*, Roxb. (W. et A., l. c. p. 189, cum syn.)—c.p. 1266  
(*partim*).  
HAB. Batticaloa District.
3. *C. ferruginea*, Grah. (Benth. Lond. Journ. of Bot. ii. 476, cum  
syn.)—c.p. 1265.  
HAB. Central Province, up to an elevation of 4000 feet.
4. *C. evolvuloides*, W. et A. Prod. i. 188.—c.p. 1266 (*partim*).  
HAB. Ceylon.
5. *C. bifaria*, Linn. (W. et A., l. c. p. 188, cum syn.; Wight, Icones,  
t. 30.)—c.p. 3593.  
HAB. Central Province.
6. *C. multiflora*, Benth. l. c. p. 478; Walp. Rep. v. 437.—c.p. 1268.  
HAB. Central Province, at an elevation of 3000 to 5000 feet.
7. *C. Wightiana*, Grah. (W. et A., l. c. p. 181.—? *C. scabrella*, W.  
et A., l. c.)—c.p. 2772.  
HAB. Central Province, at an elevation of 1000 to 3000 feet.  
A very variable species, and I suspect that *C. scabrella*, W. et A., is only a form  
of it.
8. *C. retusa*, Linn. (W. et A., l. c. p. 187, cum syn.)—c.p. 1274.  
HAB. Very common, up to an elevation of 2000 feet. Nom. vulg.  
"Kaha-andanahiriya-gass."
9. *C. verrucosa*, Linn. (W. et A., l. c. cum syn.; Wight, Icones,  
t. 200.)—c.p. 1273.  
HAB. Abundant, up to an elevation of 3000 feet. Nom. vulg. "Nil-  
andanahiriya," "Yak-bairiye."
10. *C. semperflorens*, Ventn. (Walp. Rep. v. 439, cum syn.)—c.p. 12  
(280).  
HAB. Central Province, at an elevation of 4000 to 7000 feet.
11. *C. juncea*, Linn. (W. et A., l. c. p. 185, cum syn.)—c.p. 1264.  
HAB. Not uncommon, up to an elevation of 3000 feet. Nom. vulg.  
"Hanna."  
The Cinghalese make cord of the strong fibre of this plant.
12. *C. lunulata*, Heyne, (W. et A., l. c. p. 183, cum syn.; Wight,  
Icones, t. 480.)—c.p. 1270 (1271).  
HAB. Hot, drier parts of the island.



13. *C. calycina*, Schr. (Benth. l. c. p. 564, cum syn.; Walp. Rep. l. c. 441.)—C.P. 526.

HAB. Common on open, grassy places in the Central Province, at an elevation of 3000 to 6000 feet.

14. *C. Mysorensis*, Roth, (Benth. l. c. p. 566, cum syn.; Walp. l. c.)—C.P. 3594.

HAB. Batticaloa District.

15. *C. albida*, Heyne, (Benth. l. c. p. 567, cum syn.; Walp. l. c. 442.)—C.P. 269 (2482).

HAB. Central Province, in the Badulla and Maturatte Districts, at an elevation of 2000 to 4000 feet.

16. *C. linifolia*, Linn. (Walp. l. c. cum syn.)—C.P. 1277.

HAB. Near Galle, *Gardner*.

17. *C. nana*, Burm. (W. et A., l. c. p. 191, cum syn.)—C.P. 3301.

HAB. Hotter parts of the island, at no great elevation.

18. *C. umbellata*, Wight, (W. et A., l. c.)—C.P. 1276.

HAB. Central Province, at an elevation of 3000 to 4000 feet.

If my specimens are correctly named, they approach almost too near to *C. nana* to be considered other than a variety of that species; and *C. umbellata* is perhaps only the mountain form of *C. nana*. I find the seeds of the former fewer in number, as well as rather larger and more shining, than those of *C. nana*.

19. *C. medicaginea*, Lam. (W. et A., l. c. p. 192, cum syn.)—C.P. 1278.

HAB. Hot, drier parts of the island, up to an elevation of 1500 feet.

This species varies extremely, sometimes occurring quite erect, and sparingly branched, with more numerous flowers, and with leaves a good deal larger than are seen on the procumbent seaside form. Can *C. hernarioides*, W. et A., and *C. neglecta*, W. et A., be forms of it?

20. *C. laburnifolia*, Linn. (W. et A., l. c. p. 193, cum syn.)—C.P. 367.

HAB. Very common, up to an elevation of 3000 feet. Nom. vulg. "Yak-baireya."

21. *C. quinquefolia*, Linn. (W. et A., l. c. p. 194, cum syn.; var. *trifolia*.)—C.P. 1272.

HAB. Bintenne, *Gardner*.

22. *C. striata*, DC. (Walp. l. c. p. 449, cum syn.)—C.P. 3608.

HAB. Common on roadsides, in the Nandian District.

CROTALARIA INCANA, Linn. (C.P. 1269), is a common weed in the Botanic Garden, but, no doubt, introduced.

### 3. PAROCHÆTUS, Ham.

1. *P. communis*, Ham. (Walp. Ann. iv. 478, cum syn.)—C.P. 2409.

HAB. Abundant in the Central Province, at an elevation of 4000 to 7000 feet.

TRIFOLIUM REPENS, Linn., is becoming naturalized in the neighbourhood of Nera Ellia.

### 4. ROTHIA, Pers.

1. *R. trifoliata*, Pers. (W. et A. Prod. i. 195, cum syn.)—C.P. 1452.

HAB. Abundant in the hot, drier parts of the island.

## 5. INDIGOFERA, Linn.

(Nom. vernac. "Awaree.")

1. *I. linifolia*, Retz, (W. et A. Prod. i. 198, cum syn.; Wight, Icones, t. 313.)—C.P. 3514.

HAB. Banks of the Mahawelliganga, on the Lower Badulla Road from Kandy.

2. *I. cuneaphylla*, Linn. (W. et A., l. c. p. 199, cum syn.; Wight, Icones, t. 403.)—C.P. 2775.

HAB. Hotter parts of the island; not very common.

3. *I. aspalathoides*, Vahl, (W. et A., l. c. cum syn.; Wight, Icones, t. 332.)—C.P. 1455.

HAB. Hot, drier parts of the island.

4. *I. viscosa*, Lam. (W. et A., l. c. p. 200, cum syn.; Wight, Icones, t. 404.)—C.P. 1459.

HAB. Hot, drier parts of the island; very common.

5. *I. tenuifolia*, Rottb. (W. et A., l. c.)—C.P. 1462.

HAB. Bintenne and Trincomalee, *Gardner*.

6. *I. trita*, Linn. f. (W. et A., l. c. p. 203, cum syn.; Wight, Icones, t. 315, 386.)—C.P. 1463.

HAB. Common in the hot, drier parts of the island.

7. *I. trifoliata*, Linn. (W. et A., l. c. p. 201, cum syn.; Wight, Icon. t. 314.)—C.P. 3592.

HAB. Near Minnery, on the borders of the tank.

8. *I. paucifolia*, Delile, (W. et A., l. c. p. 201, cum syn.; Wight, Icones, t. 331.)—C.P. 1454.

HAB. North of the island, *Gardner*.

9. *I. flaccida*, Koenig, (W. et A., l. c. p. 204, cum syn.)—C.P. 1460.

HAB. Haragam and other places on the Lower Badulla Road from Kandy.

10. *I. hirsuta*, Linn. (W. et A., l. c.)—C.P. 1456 (1457, 1458).

HAB. Very abundant in the hot, drier parts of the island.

11. *I. galegoides*, DC. Prod. ii. 225.—C.P. 1461.

HAB. Matele, *Gardner*.

12. *I. inamœna*, Thw.; *caule* erecto, suffruticoso, incano-strigoso, anguloso; *foliis* petiolatis, impari-pinnatis; *foliolis* 6-9-jugis, oblongis, recurvato-mucronulatis, basi angustatis, brevi-petiolulatis; *racemis* folio dimidio brevioribus, multifloris; *leguminibus* erectis, rostrato-apiculatis, teretibus, subcarinatis.—C.P. 3513.

HAB. Caloopahane, between Hapootelle and Balangodde, at an elevation of about 2000 feet.

*Folia* 1½-2½ poll. longa, *foliolis* 3-5 lin. longis, circiter 1½ lin. latis, stipulis stipellis subulatis. *Legumina* 1-1¼ poll. longa, circiter 10-sperma. *Flores* parvi, purpurei: *Calycis* laciniae subulatae.

## 6. ACANTHONOTUS, Benth.

Hook. Niger Flora, p. 293. Walpers, Ann. vol. ii. p. 362.

1. *A. echinatus*, Benth. l. c.—*Indigofera echinata*, Willd.; W. et A. Prod. i. 198, cum syn.; Wight, Icones, t. 316.—C.P. 1453.

HAB. Common in the hot, drier parts of the island.

## 7. PSORALEA, Linn.

1. **P. corylifolia**, Linn. (W. et A. Prod. i. 198, cum syn.)—c.p. 1443.  
 HAB. Not uncommon in the hotter parts of the island.

## 8. SESBANIA, Pers.

1. **S. Ægyptiaca**, Linn. (W. et A. Prod. i. 214, cum syn.; Wight, Icones, t. 32.)—c.p. 1512.  
 HAB. Habarane, *Gardner*.  
 2. **S. aculeata**, Pers. (W. et A., l. c. cum syn.)—c.p. 1513.  
 HAB. Not uncommon, up to an elevation of 3000 feet.

## 9. TEPHROSIA, Pers.

1. **T. suberosa**, DC. (W. et A. Prod. i. 210, cum syn.; Wight, Illust. i. 198. t. 82.)—*Mundulea suberosa*, Benth.; Walp. Ann. iv. 491.—c.p. 1486.  
 HAB. On rocks, near the Temple at Damboul.

2. **T. tinctoria**, Pers. (W. et A., l. c. 211, cum syn.; Wight, Icones, t. 388.)—c.p. 1449.

HAB. Var. *a*. W. et A., l. c., Central Province, up to elev. 5000 feet.  
 Var. *γ*. W. et A., l. c., near Matele, *Gardner*. Nom. vulg. "Aloo-pilla."

Although the extreme forms of this species vary so exceedingly in habit, degree of pubescence, and size of the leaflets, stipules, bracts, and calyx, it seems impossible, as Wight and Arnott remark, to separate them.

3. **T. maxima**, Pers. (W. et A., l. c. p. 213, cum syn.)—c.p. 1444.

HAB. Not uncommon in the warmer parts of the island.

4. **T. purpurea**, Pers. (W. et A., l. c. cum syn.)—c.p. 1445.

HAB. Very abundant in the warmer parts of the island. Nom. vulg. "Pilla."

The root of this plant is used as an anthelmintic for children.

5. **T. hirta**, Ham. in Linn. Trans. xiii. 546.

HAB. Hantain, *Gardner* (in Herb. Hook.).

6. **T. villosa**, Pers. (W. et A., l. c. p. 212, cum syn.)—c.p. 1447.

Var. *β*. *argentea*.—*T. argentea*, Pers.; W. et A., l. c. cum syn.—c.p. 1446.

HAB. Hot, drier parts of the island. Nom. vulg. "Boo-pilla."

*Flores purpurei vel albid.*

I have never met with any plant in the island agreeing in character with Persoon's *T. senticosa*.

## 10. STYLOSANTHES, Sw.

1. **S. mucronata**, Willd. (W. et A. Prod. i. 218, cum syn.)—c.p. 1451.

HAB. Common towards the north of the island.

## 11. ZORNIA,\* Gmel.

1. **Z. angustifolia**, Sm. (W. et A. Prod. i. 217, cum syn.)—c.p. 3598 (524 *partim*).

\* Bentham (Martius, Fl. Bras. Leg. p. 82) includes under *Z. diphylla*, Pers., all the Ceylon species, viz.—No. 1, under var. *λ*, *vulgaris punctata*, and No. 2 and 3, under var. *μ*, *Zeylonensis*.

HAB. Not uncommon in the hot, drier parts of the island.

2. **Z. conjugata**, Sm.—*Z. Zeylonensis*, Pers.; W. et A., l. c. cum syn.—C.P. 3600.

HAB. Colombo.

The joints of the legume of this species are twice as long and broad as those of the preceding and following species.

3. **Z. Walkeri**, Arn. Pug. p. 12 (330); Walp. Rep. i. 726.—C.P. 3599 (524 *partim*).

HAB. Central Province, not uncommon, up to an elevation of 3000 feet.

*Leguminis articuli parce inconspicue echinulati.*

The few echinulations generally present upon the legume of this species do not project beyond its border, and are therefore easily overlooked.

## 12. SMITHIA, Ait.

1. **S. conferta**, Sm. (DC. Prod. ii. 323.)—*S. sensitiva*, var.  $\beta$ , W. et A. Prod. i. 220, cum syn.—C.P. 2777.

HAB. Common in damp, grassy places, in the warmer parts of the island.

2. **S. blanda**, Wall. (W. et A., l. c. 221; Wight, Icones, t. 986.)—*S. paniculata*, Arn. Pug. 12 (330.)—C.P. 58.

HAB. Central Province, at an elevation of 4000 to 7000 feet.

A very variable species as regards the size of the leaves and flowers, and the amount of pubescence upon the stems and calyces.

## 13. ÆSCHYNOMENE, Linn.

(Nom. vernac. "Deya-seyembala.")

1. **Æ. aspera**, Linn. (W. et A. Prod. i. 219, cum syn.; Wight, Icon. t. 299.)—C.P. 1514.

HAB. Common in the hotter parts of the island.

2. **Æ. Indica**, Linn. (W. et A., l. c. cum syn.; Wight, Icones, t. 405.)—C.P. 1515.

HAB. Very abundant in the warmer parts of the island.

## 14. ORMOCARPUM, Beauv.

1. **O. sennoides**, DC. (W. et A. Prod. i. 216, cum syn.; Wight, Icones, t. 297.)—C.P. 1438.

HAB. Hot, drier parts of the island; not uncommon.

## 15. URARIA, Desv.

1. **U. picta**, Desv. (W. et A. Prod. i. 221, cum syn.; Wight, Icones, t. 411.)—C.P. 1437.

HAB. Not rare in the hotter parts of the island.

2. **U. hamosa**, Wall. (W. et A., l. c. 222, cum syn.; Wight, Icones, t. 284.)—*D. rufescens*, DC. (fid. Benth.)—C.P. 3590.

HAB. Nilgalla, in the Badulla District.

3. **U. crinita**, DC.

HAB. Ceylon, *Walker* (*Hb. Hook.*).

16. **DENDROLOBIUM**, Benth.*Desmodium*, Subgenus *Dendrolobium*, W. et A.

1. **D. umbellatum**, W. et A. Prod. i. 224, cum syn.; Walp. Ann. iv. 543.—C.P. 1436.

HAB. North of the island, *Gardner*.

2. **D. cephalotes**, W. et A., l. c. cum syn.; Wight, Icones, t. 373.—C.P. 2780.

HAB. Very common, up to an elevation of 3000 feet.

17. **PHYLLODIUM**, Desv.

Walpers, Ann. vol. iv. p. 544.

1. **P. pulchellum**, Desv.—*Dicerma pulchellum*, DC.; W. et A. Prod. i. 230, cum syn.; Wight, Icones, t. 418.—C.P. 1279.

HAB. Hot, drier parts of the island. Nom. vulg. "Hampinne."

18. **DICERMA**, DC.

1. **D. biarticulatum**, DC. (W. et A. Prod. i. 230, cum syn.; Wight, Icones, t. 419.)—C.P. 1450.

HAB. Hotter parts of the island.

19. **PTEROLOMA**, Desv.

1. **P. triquetrum**, Benth. (Walp. Ann. iv. 543.)—*Desmodium triquetrum*, DC.; W. et A. Prod. i. 224, cum syn.—C.P. 1431.

HAB. Central Province; abundant. Nom. vulg. "Bāloleyā."

20. **CATENARIA**, Benth. in Pl. Jungh. ii. 220.

1. **C. laburnifolia**, Benth. l. c. cum syn.; Walp. Ann. iv. 544.—C.P. 1435.

HAB. Central Province, at an elevation of 1500 to 3000 feet.

21. **DESMODIUM**, DC.

1. **D. triflorum**, DC. Prod. ii. 334, cum syn.; Wight, Icones, t. 292.—C.P. 2779.

HAB. Abundant, up to an elevation of 2000 feet. Nom. vulg. "Heen-oondoopeyelli."

This is valued as a medicine for the cure of dysentery.

2. **D. heterophyllum**, DC., l. c. cum syn.—*D. triflorum*, var. *a*, W. et A. Prod. i. 229; Wight, Icones, t. 291.—C.P. 2778.

HAB. Equally abundant with the last, from which it is certainly distinct. Nom. vulg. "Maha-oondoopeyelli."

3. **D. parvifolium**, DC. (W. et A., l. c. cum syn.)—C.P. 1429.

HAB. Central Province, at an elevation of 3000 to 5000 feet.

4. **D. polycarpon**, W. et A., l. c. 227, cum syn.

Var. *a*. *caulibus adpresse pilosis; bracteis longe acuminatis; leguminibus hirsutis vel sæpe glaberrimis*.—C.P. 1425.

Var.  $\beta$ . *caulibus* superne villosis; *bracteis* lanceolatis, nec longe acuminatis.—C.P. 3512.

HAB. Central Province, up to an elevation of 5000 feet.

5. **D. ferrugineum**, Wall. Cat. 5732 A.—*D. rufescens*, W. et A., l. c. 228 (non De Candolle); Wight, Icones, t. 984.—C.P. 1426.

HAB. Central Province, up to an elevation of 4000 feet.

This, like the preceding species, varies much in the amount of pubescence upon the several parts, as well as in the relative length of the acumen of the bracteæ.

6. **D. gyrans**, DC. (W. et A., l. c. 227, cum syn.; Wight, Icones, t. 294.)—C.P. 1430.

HAB. Central Province, up to an elevation of 3000 feet.

7. **D. gyroides**, DC. (W. et A., l. c.)—C.P. 3326.

HAB. Pusalawa, in the Central Province, according to the statement of the Plant Collector, but I have never seen it truly wild in the island.

8. **D. latifolium**, DC. (W. et A., l. c. 225, cum syn.; Wight, Icones, t. 270, 272.)—C.P. 3588.

HAB. Near Bibili, in the Badulla District.

9. **D. Wightii**, Gr. (W. et A., l. c. 226; Arn. Pug. p. 13, 331.)—C.P. 3129.

HAB. Abundant in the warmer parts of the island, up to an elevation of 1500 feet.

10. **D. ormocarpoides**, DC.

HAB. Ceylon, Gardner (*Hb. Hook.*).

11. **D. podocarpum**, DC. Prod. i. 336 (non Hook. et Arn.)—C.P. 1434.

HAB. Central Province, up to an elevation of 4000 feet.

*Flores* parvi, albid.

12. **D. strangulatum**, W. et A., l. c. 228; Wight, Icones, t. 985.—C.P. 1433.

Var.  $\beta$ . *minor*; *caule* villosa; *foliis* rotundato-rhomboideis, obtusis, minute apiculatis, lateralibus vix obliquis; *stipulis* villosis; *floribus* multo minoribus; *leguminibus* articulis minus constrictis.—C.P. 3329.

An species distincta?

*Flores* (var.  $\alpha$ ) aurantiaco-rubri.

*Desmodium Walkeri*, Arn. Pug. p. 13, 331, C.P. 2973, appears to be identical with De Candolle's *Desm. cajanifolium*, and can scarcely have any claim to be considered indigenous to Ceylon.

## 22. PSEUDARTHRIA, W. et A.

1. **P. viscida**, W. et A. Prod. i. 209, cum syn.; Wight, Icones, t. 286.—C.P. 1432.

HAB. Common up to an elevation of 3000 feet.

## 23. ALYSICARPUS, Neck.

(Nom. vernac. "Aswenna.")

1. **A. vaginalis**, DC. (Walp. Ann. iv. 548, cum syn.)—C.P. 1428.

HAB. Abundant up to an elevation of 3000 feet.

2. **A. bupleurifolius**, DC. (W. et A. Prod. i. 233, cum syn.)—C.P. 1427.

HAB. Hot, drier parts of the island, *Gardner*.

3. **A. scariosus**, Grah.—*A. styracifolius*, W. et A., l. c. 234 (non DC.)—C.P. 3589.

HAB. Minnery tank.

4. **A. Heyneanus**, Wight.

HAB. Ceylon, *Walker* (*Hb. Hook.*).

## 24. DUMASIA, DC.

1. **D. villosa**, DC. Prod. ii. 241.—Var.  $\beta$ , *D. pubescens*, DC., l. c. Arn. Pug. p. 12, 330. *D. congesta*, Grah.; W. et A. Prod. i. 206; Wight, Icones, t. 445.—C.P. 663.

HAB. Common in the Central Province, at an elevation of 5000 to 7000 feet.

## 25. CLITOREA, Linn.

1. **C. Ternatea**, Linn. (W. et A. Prod. i. 205, cum syn.)—C.P. 1485.

HAB. Not uncommon in the Batticaloa District. Nom. vulg. "Nil-Kattarodoo."

## 26. SHUTERIA, W. et A.

1. **S. vestita**, W. et A. Prod. i. 207; Wight, Icones, t. 165.—C.P. 661.

HAB. Central Province, at an elevation of 4000 to 7000 feet.

## 27. SOJA, Moench.

1. **S. Wightii**, Grah. (Walp. Ann. iv. 552, cum syn.)—C.P. 1477.

HAB. Common up to an elevation of 2000 feet.

## 28. GALACTIA, P. Browne.

1. **G. tenuifolia**, W. et A. Prod. i. 206, cum syn.—C.P. 1468.

HAB. Central Province, up to an elevation of 3000 feet.

## 29. GLYCINE, Linn.

1. **G. labialis**, Linn. (W. et A. Prod. i. 208, cum syn.; Wight, Icones, t. 168.)—? *G. mollis*, W. et A., l. c.; Arn. Pug. p. 12, 330.—C.P. 1471.

HAB. Hotter parts of the island, up to an elevation of 2000 feet.

This is such a variable plant that I cannot help suspecting, judging from the description, that *G. mollis* of W. et A. is only a form of it.

## 30. CANAVALIA, DC.

(Nom. vernac. "Wal-awara.")

1. **C. obtusifolia**, DC. (W. et A. Prod. i. 253, cum syn.)—C.P. 1484.

HAB. Common near the sea-shore.

2. **C. gladiata**, DC. (W. et A., l. c. cum syn.; Wight, Icones, t. 753.)—C.P. 1481.

HAB. North of the island, *Gardner*.

3. **C. virosa**, W. et A., l. c. cum syn.—c.p. 2782.

HAB. Central Province, abundant up to an elevation of 3000 feet.

### 31. MUCUNA, Adans.

1. **M. monosperma**, DC. (W. et A. Prod. i. 254, cum syn.; Wight, Icones, t. 35.)—c.p. 2662.

HAB. Central Province, up to an elevation of 3000 feet.

2. **M. atropurpurea**, DC. (W. et A., l. c.)—c.p. 1483.

HAB. Not uncommon, up to an elevation of 2000 feet.

3. **M. gigantea**, DC. (W. et A., l. c.)—c.p. 3587.

HAB. Batticaloa? *Gardner*.

4. **M. prurita**, Hook. (W. et A., l. c. 255.)—Var.  $\beta$ . *pedunculis brevibus*, sæpissime 2-floris.—c.p. 3150.

HAB. Hot, drier parts of the island, not uncommon. Var. *a*. In exposed situations. Var.  $\beta$ . In shady places. Nom. vulg. "Aachāriya-paloo."

### 32. ERYTHRINA, Linn.

1. **E. Indica**, Lam. (W. et A. Prod. i. 260, cum syn.; Wight, Icones, t. 58.)—c.p. 1464.

HAB. Hot, drier parts of the island. Nom. vulg. "Errabadoo-gass."

This is used medicinally for men and cattle, and the young, tender leaves are eaten in curries.

2. **E. ovalifolia**, Roxb. Fl. Ind. iii. 254; Wight, Icones, t. 247.—c.p. 3499.

HAB. Hot, drier parts of the island. Nom. vulg. "Yak-errabadoo-gass."

### 33. BUTEA, Roxb.

1. **B. frondosa**, Roxb. (W. et A. Prod. i. 261.)—c.p. 1465.

HAB. Hot, drier parts of the island. Nom. vulg. "Gass-kaala."

### 34. STRONGYLODON, Vogel.

1. **S. ruber**, Vog. (Walp. Ann. iv. 559, cum syn.)—c.p. 1479.

HAB. Near Kornegalle, apparently quite wild.

*Legumen* oblongum, inflatum, membranaceum, marginatum, indehiscens, rostratum, stipitatum,  $2\frac{1}{2}$ -3 poll. longum,  $1\frac{1}{4}$ - $1\frac{1}{2}$  latum, 1-2-spermum, *stipite*  $\frac{1}{2}$  poll. longo. *Semina* ovoidea, lævia, nigra, 8-9 lin. longa, 7 lin. lata, *carunculo* rubro 1 lin. alto semicineta, *testa* coriacea, *cotyledonibus* duris crassis consolidatis.

### 35. PHASEOLUS, Linn.

(Nom. vernac. "Wal-maa.")

1. **P. Truxillensis**, H.B.K.—*P. rostratus*, Wall. (W. et A. Prod. i. 244, cum syn.; Wight, Icones, t. 34.)—c.p. 1480.

HAB. Warmer parts of the island.

2. **P. radiatus**, Linn. (W. et A., l. c. 246, cum syn.)—c.p. 1473.

HAB. Very abundant up to an elevation of 2000 feet.



3. **P. trinervius**, Heyne, (W. et A., l. c. 245, cum syn.)—C.P. 1476.  
 HAB. Central Province, up to an elevation of 4000 feet.
4. **P. trilobus**, Ait. (W. et A., l. c. 246, cum syn.; Wight, Icones, t. 94.)—C.P. 1474.  
 HAB. Common amongst grass in the hotter parts of the island, varying greatly in the shape of the leaflets.
5. **P. Grahamianus**, W. et A., l. c. 244.—C.P. 3586.  
 HAB. Batticaloa.  
*Bracteolæ* minutæ, rotundatæ, membranacæ, deciduæ. *Calyx* ciliatus,  $1\frac{1}{2}$  lin. longus, *labio* superiore truncato emarginato, *laciniis* inferioribus brevibus subacutis. *Corolla* purpurea, circiter  $\frac{1}{2}$  poll. longa.
6. **P. semierectus**, Linn.—*Phas. psoralioides*, W. et A. Prod. i. 244; Wight, Icones, t. 249.—C.P. 3609.  
 HAB. Central Province.

### 36. VIGNA, Savi.

1. **V. vexillata**, Benth.—*V. carinalis*, Benth.; Walp. Rep. v. 538.—C.P. 2781.  
 HAB. Abundant in the Central Province, up to an elevation of 4000 feet.
2. **V. luteola**, Benth. MSS. An *Dolichos luteus*, Sw.?—C.P. 1482.  
 HAB. Not uncommon near the seacoast. Galle, *Gardner*; Colombo, *Ferguson*.

### 37. DOLICHOS, Linn.

1. **D. uniflorus**, Lam. (W. et A. Prod. i. 248, cum syn.)—C.P. 1475.  
 HAB. Central Province, *Gardner*.
2. **D. ciliatus**, Klein, (W. et A., l. c. 249, cum syn.)—C.P. 1469.  
 HAB. Hot, drier parts of the island.
3. **D. falcatus**, Klein, (W. et A., l. c. cum syn.)—C.P. 1472.  
 HAB. Hantani, *Gardner*; Batticaloa District.

### 38. LABLAB, Adans.

1. **L. vulgaris**, Savi, (W. et A. Prod. i. 250, cum syn.; Wight, Icones, t. 57, 203.)—C.P. 1470.  
 HAB. Common in the warmer parts of the island.

### 39. CAJANUS, DC.

1. **C. Indicus**, Spr. (W. et A. Prod. i. 256, cum syn.)—C.P. 720.  
 HAB. Ambagamowa, *Gardner*. Nom. vulg. "Rata-tora."  
 This is said to be not truly wild in the island, and the native name would also seem to indicate it.

### 40. DUNBARIA, W. et A.

1. **D. Heynei**, W. et A. Prod. i. 258.—*D. oblonga*, Arn. Pug. 15, (333.)—C.P. 1478.  
 HAB. Central Province; common, up to an elevation of 3000 feet.

41. **ATYLOSIA**, W. et A.

1. **A. Candollei**, W. et A. Prod. i. 257, cum syn.; Wight, Icones, t. 754.—? *A. major*, W. et A., l. c.—c.p. 55 (209, 224).

HAB. Very abundant amongst grass, in the Central Province, up to an elevation of 4000 feet. Nom. vulg. "Atta-tora."

2. **A. albicans**, Benth.—*Cantharospermum albicans*, W. et A., l. c. p. 256, cum syn.—c.p. 2783.

HAB. Hantani, *Gardner*.

3. **A. rugosa**, W. et A., l. c. p. 257.—c.p. 1441.

HAB. Central Province, up to an elevation of 5000 feet. Nom. vulg. "Wal-Cölooo."

4. **A. scarabæoides**, Benth.—*Cantharospermum pauciflorum*, W. et A. 255, cum syn.—c.p. 1442.

HAB. Hot, drier parts of the island.

42. **RHYNCHOSIA**, Lour.

1. **R. cana**, DC. (W. et A. Prod. i. 240, cum syn.)—c.p. 1280.

HAB. Abundant in the warmer parts of the island.

2. **R. minima**, DC. (Prod. ii. 385, cum syn.)—*R. medicaginea*, DC., l. c. 386; W. et A., l. c. 238, cum syn.—c.p. 1440.

HAB. Jaffna, *Gardner*.

3. **R. rufescens**, DC. (W. et A., l. c. 239, cum syn.; Arn. Pug. p. 14, 332, n. 42.)—c.p. 1424.

HAB. Batticaloa District; common.

43. **ABRUS**, Linn.

1. **A. precatorius**, Linn. (W. et A. Prod. i. 236, cum syn.)—c.p. 1466.

HAB. Not uncommon in the hotter parts of the island. Nom. vulg. "Olinda-wel."

2. **A. pulchellus**, Wall.; *foliolis* lineari-oblongis, 7-10-jugis; *paniculis* terminalibus, longe pedunculatis, racemiformibus, *ramis* earundem subsecundis incrassatis curvatis, *pedicellos* serie continua diu gerentibus; *leguminibus* compressis, subfalcatis, brevi-rostratis, circiter 9-spermis.—c.p. 1467.

HAB. Not uncommon in the hot, drier parts of the island.

*Frutex* scandens, volubilis, plus minus adpresse pilosus. *Stipulae* stipellæque parvæ, lanceolato-subulatae. *Calyx* campanulatus, minute 5-dentatus, pedicello subæquilongus, 1 lin. longus. *Corolla* calyce 3-4-plo longior, pallide purpurea. *Legumina* 1½-2¼ poll. longa, 3½-4 lin. lata. *Semina* subrotundata, compressa, minute tuberculata, nigra, 2 lin. longa.

44. **NOMISMIA**, W. et A.

1. **N. nummularia**, W. et A. Prod. i. 237, cum syn.; Wight, Icones, t. 283.—c.p. 1439.

HAB. Jaffna, *Gardner*.

45. **CYANOSPERMUM**, W. et A.

1. **C. tomentosum**, W. et A. Prod. i. 260, cum syn.; Wight, *Illust.* i. p. 198. t. 81, 84.—C.P. 2471.

HAB. Central Province, at an elevation of 2000 to 3000 feet.

46. **PYCNOSPORA**, Brown, W. et A.

1. **P. nervosa**, W. et A. Prod. i. 197, cum syn.—C.P. 279.

HAB. Not uncommon up to an elevation of 2000 feet.

47. **FLEMINGIA**, Roxb.

1. **F. strobilifera**, Brown, (W. et A. Prod. i. 243, cum syn.; Wight, *Icones*, t. 267.)—C.P. 697.

HAB. Warmer parts of the island. Nom. vulg. "Hampilla."

2. **F. lineata**, Roxb. (W. et A., l. c. 242, cum syn.; Wight, *Icones*, t. 327.)—C.P. 1423.

HAB. Hot, drier parts of the island.

3. **F. congesta**, Roxb. (W. et A., l. c. 241, cum syn.; Wight, *Icones*, t. 390.)—C.P. 2663.

HAB. Ambagamowa District.

4. **F. semialata**, Roxb. (*Fl. Ind.* iii. p. 340; Wight, *Icones*, t. 326.)—C.P. 256, 3585.

HAB. Not uncommon in the warmer parts of the island.

48. **PTEROCARPUS**, Linn.

1. **P. Marsupium**, Roxb. (W. et A. Prod. i. 266, cum syn.)—C.P. 1495.

HAB. Central Province, up to an elevation of 3000 feet. Nom. vulg. "Gammāloo-gass."

The gum kino which exudes from the bark of this tree is used medicinally by the natives.

49. **PONGAMIA**, Linn.

1. **P. glabra**, Ventn. (W. et A. Prod. i. 262, cum syn.; Wight, *Icones*, t. 59.)—C.P. 1489 (1490).

HAB. Banks of rivers, common up to an elevation of 2000 feet. Nom. vulg. "Magool-karanda-gass."

50. **DERRIS**, Lour.

(Nom. vernac. "Calla-wel.")

The tough bark of the species of this and of some of the allied genera is used by the Cinghalese for making ropes.

1. **D. uliginosa**, Benth. (Walp. *Ann.* iv. p. 581.)—*Pongamia uliginosa*, DC.; W. et A. Prod. i. 262, cum syn.—C.P. 1494.

HAB. Hotter parts of the island.

2. **D. ovalifolia**, Benth., var.—*Pongamia ovalifolia*, W. et A. Prod. i. 262; Wight, *Icones*, t. 328.—C.P. 1493.

HAB. Hantani, Saffragam, and Ambagamowa Districts.

Scandens. *Foliola* ovata vel obovato-oblonga, apice rotundata vel abrupte brev-acuminata, juniora rufescentia.

3. **D. sinuata**, Benth.; *foliolis* 3-jugis cum impari, ovatis vel ovato-lanceolatis, glabris, apice emarginatis, basi rotundatis; *paniculis* sæpissime terminalibus, elongatis, adpresse pilosis; *leguminibus* compressis, nitidis, reticulatis, rostrato-acuminatis, sæpe seminum abortu varie sinuatis, *marginè* seminifera alata.—C.P. 1491.

HAB. Batticaloa, *Gardner*; Cattura District.

Scandens. *Foliola* 1-4 poll. longa,  $\frac{3}{4}$ -2 poll. lata; *petiolulis* incrassatis, 2 $\frac{1}{2}$  lin. longis. *Pedicelli* calyce breviores. *Bracteolæ* minutæ, acutæ, cito deciduæ. *Corolla* pallide purpurea; *verillo* orbiculari, emarginato; *alis* anguste lanceolatis, subfalcatis, carinam apice fissam arcte amplectentibus. *Stamina* diadelpa, 9 et 1; *antheris* oblongis. *Semina* oblonga, parce curvata.

## 51. BRACHYPTERUM, Benth.

1. **B. scandens**, W. et A. (Walp. Ann. iv. 579, cum syn.; Wight, Icones, t. 275.)—C.P. 1492.

HAB. Abundant in the hotter parts of the island.

2. **B. Benthamii**, Thw.; *foliolis* 2-3-jugis cum impari, oblongis, obtuse brevi-acuminatis, glabris; *paniculis* terminalibus, elongatis, adpresse pilosis, *ramis* racemiformibus; *pedicellis* solitariis; *leguminibus* oblongis, adpresse pilosis.—C.P. 2925.

HAB. Ooma Oya, on the Lower Badulla Road from Kandy.

Scandens. *Foliola* 1 $\frac{1}{2}$ -4 $\frac{1}{2}$  poll. longa,  $\frac{1}{2}$ -2 poll. lata, *petiolulo* 1 $\frac{1}{2}$ -2 lin. longo. *Pedicelli* calyce æquilongi. *Calyx* rubiginosus, antice brevi-3-dentatus. *Petala* pallide rosea. *Ovarium* circiter 4-ovulatum. *Legumina* 1-1 $\frac{3}{4}$  poll. longa, 6-7 lin. lata.

This species, in general appearance, closely resembles the preceding, but is sufficiently distinguished by the different character of the inflorescence and the width of the legumes.

3. **B. elegans**, Thw.; *foliolis* 3-4-jugis cum impari, ellipticis vel oblongis, abrupte brevi-acuminatis, glabris, subtus minutissime reticulatis; *paniculis* axillaribus terminalibusque, glabris, foliis subæquilongis, racemiformibus; *pedicellis* fasciculatis; *leguminibus* linearibus, apice rotundatis, oblique apiculatis, 1-2-spermis, glabris.—C.P. 2508.

HAB. Central Province, at no great elevation.

Scandens. *Foliola* 1 $\frac{1}{4}$ -2 poll. longa,  $\frac{1}{2}$ -1 poll. lata, membranacea; *petiolulo* tenui, 2 lin. longo. *Pedicelli* 2 lin. longi. *Calyx* minute ciliatis, 1 lin. longus, antice brevi-3-dentatus, basi *bracteolis* 2 minutis lanceolatis ciliatis suffultus. *Ovarium* 2-ovulatum. *Legumina* 1 $\frac{3}{4}$ -2 $\frac{3}{4}$  poll. longa,  $\frac{1}{2}$  poll. lata.

## 52. DALBERGIA, Linn.

1. **D. Mooniana**, Thw.—*D. Lanceolaria*, Moon's Cat. p. 51; *foliolis* 5-8, alternis, majusculis, ovatis vel oblongis, acuminatis, subtus pallidioribus minute reticulatis; *paniculis* terminalibus; *leguminibus* linearibus, marginatis, stipitatis, rostratis, 1-6-spermis, sæpe seminum abortu varie sinuatis.—C.P. 439.

HAB. Southern and central parts of the island, at no great elevation. Nom. vulg. "Nadoong-gass."

*Arbor* ingens. *Ramuli* albo-punctati. *Gemmæ* pilosæ. *Folia* glabra, rachi petiolisque pilosis, denique glabris. *Foliola* 2 $\frac{1}{2}$ -7 poll. longa, 1 $\frac{1}{2}$ -3 poll. lata,

*petiolulo*  $1\frac{1}{2}$  lin. longo. *Stipulæ* lanceolatae, pilosæ, deciduæ, 2 lin. longæ. *Legumina* coriacea, lævia, obscure reticulata, 11 lin. lata, marginibus 1 lin. latis superiore crassiore.

This tree produces a very good timber, well adapted for furniture and for buildings.

2. **D. Championii**, Thw. — *Endespermum Zeylanicum*, Champion, MSS.; *foliolis* 2–4, alternis vel suboppositis, ellipticis acuminatis reticulatis, magnitudine valde variis, superne glabris, subtus fugaci pilosulis, *paniculis* terminalibus pilosis foliosis, leguminibus membranaceo-foliaceis, monospermis, lineari-lanceolatis, subacutis, stipitatis, adpresse pilosulis, medio reticulatis.—C.P. 761 (772).

HAB. Central Province, at an elevation of 1000 to 3000 feet.

Scandens. *Foliola* 2–6 poll. longa, 1–3 poll. lata; *petiolulo* tumidulo,  $\frac{1}{4}$  poll. longo. *Flores* 3 lin. longi. *Calyx* albidus, ferrugineo-pilosulus, dentibus æquilongis, tubo duplo brevioribus, 2 sup. rotundatis, reliquis acutis. *Petala* albida, pallide rufo-venosa, longe unguiculata. *Stamina* 9, monadelphæ. *Ovarium* longe stipitatum, 1-ovulatum.

This species is evidently very closely allied to Blume's *Endespermum scandens*.

3. **D. frondosa**, Roxb. (W. et A. Prod. i. 266, cum syn.)—C.P. 1496.

HAB. Not uncommon in forests, in the Central Province and elsewhere, up to an elevation of 3000 feet.

I have not met with any plants in Ceylon corresponding with the descriptions of *Dalbergia Lanceolaria*, Linn. f., and *Dalbergia Zeylanica*, Roxb.

4. **D. monosperma**, Dalzell.—C.P. 243.

HAB. Ceylon, *Walker* (in Herb. Hook.).

### 53. SOPHORA, Linn., R. Br.

1. **S. tomentosa**, Linn. (W. et A. Prod. i. 179, cum syn.; Walp. Ann. iv. 586, cum syn.)—C.P. 1487.

HAB. Seacoast, in the Galle District. Nom. vulg. "Moodoo-moroonga."

2. **S. heptaphylla**, Linn. (Walp. Ann. l. c.; Wight, Icones, t. 1155.)—C.P. 570.

HAB. Central Province, at an elevation of 4000 to 7000 feet.

3. **S. violacea**, Thw.; *foliolis* 7–12-jugis cum impari, ellipticis, obtusis, brevissime mucronulatis, pilosulis, subtus pallidioribus; *racemis* terminalibus, multifloris, adpresse pilosis, folio subæquilongis; *leguminibus* pilosulis, valde constrictis, nodis ovoideis.—C.P. 3546.

HAB. Galle District, between Dickwelle and Matura.

*Frutex* mediocris. *Ramuli* petiolique adpresse pilosi. *Stipulæ* bracteæque minutæ, subulatae, deciduæ. *Foliola* 6–15 lin. longa, 3–7 lin. lata, *petiolulo*  $\frac{1}{2}$ –1 lin. longo. *Calyx* tubulosus, margine 5-lobus, pedicello æquilongo oblique affixus. *Petala* violacea, longe unguiculata, *vexillo* distante recurvato, *alis* carina angustioribus longioribusque. *Ovarium* circiter 7-ovulatum. *Semina* ovoidea, rubra.

### SUBORD. 2. CÆSALPINIÆ, R. Br.

### 54. GUILANDINA, Linn.

1. **G. Bonduc**, Linn. (W. et A. Prod. i. 280, cum syn.)—C.P. 1524.

HAB. Common up to an elevation of 3000 feet. Nom. vulg. "Koombooroo-wel."

Every part of this plant is used medicinally by the natives.

### 55. CÆSALPINIA, Linn.

1. *C. paniculata*, Roxb. (W. et A. Prod. i. 281; Wight, Icones, t. 36.)—C.P. 1525.

HAB. Common up to an elevation of 3000 feet. Nom. vulg. "Deyawawool-atteya."

2. *C. digyna*, Rottl. (W. et A., l. c. cum syn.)—C.P. 1527.

HAB. Kornegalle, *Gardner*.

3. *C. mimosoides*, Lam. (W. et A., l. c. cum syn.; Wight, Icones, t. 392.)—C.P. 3601 (2784 *partim*).

HAB. Abundant in the warmer parts of the island. Nom. vulg. "Goda-wawool-atteya."

4. *C. sepiaria*, Roxb. (W. et A., l. c. 282, cum syn.; Wight, Icones, t. 37.)—C.P. 3602 (2784 *partim*).

HAB. Abundant up to an elevation of 3000 feet.

### 56. TAMARINDUS, Linn.

1. *T. officinalis*, Hook. (Walp. Ann. iv. 595, cum syn.)—Var. *a*, *F. Indica*, DC.—C.P. 1501.

HAB. Hot, drier districts of the island, but probably not truly indigenous to Ceylon. Nom. vulg. "Seyembala."

The ripe fruit is used as a condiment for curries: a decoction of the fresh leaves is employed as a detergent for foul wounds and sores.

### 57. CASSIA, Linn.

1. *C. Fistula*, Linn. (W. et A. Prod. i. 285, cum syn.; Walp. Ann. iv. 595, cum syn.)—*C. rhombifolia*, Roxb. Fl. Ind. ii. 334; Wight, Icones, t. 269.—C.P. 1502.

HAB. Very abundant in the hot, drier parts of the island. Nom. vulg. "Ahalla-gass."

Every part is used medicinally by the Cinghalese as a purgative. The heart of the tree yields a good timber.

2. *C. Roxburghii*, DC. (W. et A., l. c. 286, cum syn.; Wight, *Illustr.* i. 198. t. 83.)—C.P. 1505.

HAB. Frequently growing with the preceding, and is equally abundant. Nom. vulg. "Ratoo-waa."

3. *C. tomentosa*, Linn. f. (W. et A., l. c. cum syn.; Walp. Ann. iv. 596, cum syn.)—C.P. 2410.

HAB. Newera Ellia, at an elevation of 7000 feet.

4. *C. Sophera*, Linn. (W. et A., l. c. 287, cum syn.)—C.P. 1507.

HAB. Hotter parts of the island; not uncommon. Nom. vulg. "Ooroo-tora."

5. *C. occidentalis*, Linn. (W. et A., l. c. 290, cum syn.)—C.P. 1262.

HAB. Very abundant up to an elevation of 2000 feet. Nom. vulg. "Penni-tora."

The tender leaves of this and the next species are eaten by the Cinghalese in their curries.

6. **C. Tora**, Linn. (W. et A., l. c. cum syn.; Walp. Ann. l. c. cum syn.)—C.P. 2785.

HAB. Very abundant on roadsides, etc.

7. **C. glauca**, Linn. (W. et A., l. c. p. 289, cum syn.; Walp. l. c. p. 597, cum syn.)—C.P. 1526.

HAB. Colombo, *Moon*. Nom. vulg. "Wal-ahalla-gass."

8. **C. auriculata**, Linn. (W. et A., l. c. p. 290, cum syn.)—C.P. 1503.

HAB. Abundant in many parts of the island near the seacoast. Nom. vulg. "Ranawarā."

A decoction or infusion of the leaves of this plant is much esteemed as a cooling medicine by the Cinghalese.

9. **C. florida**, Vahl, (W. et A., l. c. p. 288, cum syn.; Walp. Ann. iv. 597, cum syn.)—C.P. 3485.

HAB. Banks of rivers, in the Central Province, up to an elevation of 2000 feet. Nom. vulg. "Waa-gass."

The wood of this tree is extremely hard and durable.

10. **C. Timoriensis**, DC. Prod. ii. 499; Walp. l. c. cum syn.—C.P. 1504.

HAB. Not uncommon in the warmer parts of the island. Nom. vulg. "Arramana."

11. **C. Absus**, Linn. (W. et A., l. c. p. 291, cum syn.; Walp. l. c. p. 598, cum syn.)—C.P. 1506.

HAB. Very common amongst high grass, up to an elevation of 3000 feet. Nom. vulg. "Boo-tora."

12. **C. mimosoides**, Linn. (DC. Prod. ii. 503; Walp. Rep. i. 837; Walp. Ann. iv. 598.)—C.P. 1510.

Var.  $\gamma$ . *villosula*; tota planta pilis longis subpatentibus flavis sparse tecta; *foliis* patello majoribus, aristato-mucronatis.—C.P. 3603. An species distincta?

HAB. Abundant amongst high grass, in exposed situations, up to an elevation of 4000 feet. Var.  $\gamma$ . Maturatte District.

Var.  $\gamma$  is probably a distinct species; the bracteolæ near the apex of the pedicel are nearer to the base of the flower than in the typical form of *C. mimosoides*, and the petiolar gland is a little below instead of between the last pair of leaflets.

13. **C. Wallichiana**, DC.; *caule* erecto, rigido, parce ramoso; *foliolis* 20-25-jugis, oblongo-linearibus, aristato-mucronatis; *glandula* paulo supra medium petioli subsessili; *pedicellis* supra-axillaribus, fasciculatis; *staminibus* 10, valde inæqualibus.—C.P. 2786.

HAB. Open, grassy places in the Central Province. Hantani, *Macrae*. Haragam, near Kandy. Nom. vulg. "Bin-seyembala."

*Caulis* teres, 1-4-pedalis, pilosulus. *Foliola* glabra, margine ciliolata, 5-9 lin. longa, 1-1½ lin. lata. *Fasciculi* circiter 4-flori, *pedicellis* 2-4 lin. longis. *Calyx* pilosulus. *Petala* basi interne, cum staminibus partim, rufo-tincta. *Legumina* pilosula.

Sufficiently distinguished from the preceding species by the darker colour of the stems and foliage, the much larger leaves, shorter pedicel, and different shape and position of the petiolar gland.

14. **C. Kleinii**, W. et A., l. c. p. 293, cum syn.; Walp. Rep. i. 837.—C.P. 1508 (1509).

Var.  $\beta$ . *pilosa*; planta tota patentim dense pilosa.

HAB. Abundant in similar situations to the last. Var.  $\beta$ . Hantani District. Nom. vulg. "Bin-seyembala."

CASSIA ALATA, Linn., has become naturalized in various parts of the island.

### 58. JONESIA, Roxb.

1. **J. Asoka**, Roxb. (W. et A. Prod. i. 284, cum syn.; Wight, Icon. t. 206; Walp. Ann. iv. 609.)—C.P. 653.

HAB. Sides of streams, under the shade of larger trees, up to an elevation of 3000 feet. Very abundant in the Bintenne District. Nom. vulg. "Deya-ratmal."

### 59. HUMBOLDTIA, Vahl.

1. **H. laurifolia**, Vahl, (W. et A. Prod. i. 285, cum syn.; Wight, Icones, t. 1605.)—C.P. 328.

HAB. Very abundant in the Ambagamowa and Ratnapoora Districts, on the shady banks of streams. Nom. vulg. "Gal-karanda."

### 60. DIALIUM, Linn.

1. **D. ovoideum**, Thw.; *foliolis* bijugis, suboppositis, cum impari, lanceolatis, apice retusis, superne glabris, subtus sparse minutissime puberulis; *paniculis* terminalibus adpresse incano-pilosis, *folio* longioribus; *bracteolis* lanceolatis, cito deciduis; *sepalis* lanceolatis; *legumine* ovoideo, parce compresso, subsessili, 1-2-spermo, velutino.—C.P. 3149.

HAB. Districts north of Kandy, at no great elevation. Nom. vulg. "Gal-seyembala-gass."

*Arbor* ingens, ramulis rugosulis. *Foliola* 2-3 poll. longa,  $\frac{3}{4}$ -1 $\frac{1}{4}$  lata, *petiolulo* tumidulo 2 lin. longo. *Flores* albidi, circiter 3 lin. longi, *pedicellis* parum longiores. *Antheræ* oblongæ, extrorsæ, in anthesi filamentis subæquilongæ. *Ovarium* in disco annulari viridi sessile, 2-ovulatum; *stylo* subulato, apice obtusiusculo. *Legumen* atrobrunneum, 7-8 lin. longum, 5-6 lin. latum. *Semina* rotundata, compressa, striata, nitida, ochraceo-vel rufo-brunnea, 4 lin. longa.

The fruit of this species is of an agreeable acid flavour, and is sold in the bazaars. The wood is strong and handsome, and well adapted for ornamental furniture.

### 61. CYNOMETRA, Linn.

1. **C. ramiflora**, Linn. (W. et A. Prod. i. 293; Walp. Ann. iv. 601.)—C.P. 1500.

Var.  $\alpha$ ; *foliolis* omnibus 1-jugis.—C.P. 3604.

Var.  $\beta$ . *heterophylla*; *foliolis* versus basin innovationum 2-jugis, versus apicem 1-jugis.

HAB. Var.  $\alpha$ . Batticaloa District. Var.  $\beta$ . Trincomalee, Gardner. Cultura District. Nom. vulg. "Gal-mendora-gass."

Not having been so fortunate as to obtain the flowers or fruit of var.  $\alpha$ , I am unable to determine whether var.  $\beta$  is really distinct, as a species, from it, but I almost suspect it will prove so, the leaflets being much less acuminate, and slightly different in other respects.



62. **BAUHINIA**, Linn.

1. **B. tomentosa**, Linn. (W. et A. Prod. i. 295, cum syn.; Walp. Ann. iv. 602, cum syn.)—c.p. 1498.

HAB. Hot, drier parts of the island. Nom. vulg. "Kaha-pettang."

63. **PILIOSTIGMA**, Hochstett.

1. **P. racemosum**, Benth. (Walp. Ann. iv. 607, cum syn.)—*Bauhinia racemosa*, Lam.; W. et A. Prod. i. 295, cum syn.—c.p. 1497.

HAB. Very abundant in the hot, drier parts of the island. Nom. vulg. "Myla-gass."

The bark of this tree is very tough, and is made into ropes by the Cinghalese. The leaves are a very favourite food of the elephant.

SUBORD. 3. *MIMOSEÆ*.64. **ENTADA**, Adans.

1. **E. scandens**, Benth. (Hook. Journ. of Bot. iv. 332.)—*E. Pursætha*, DC.; W. et A. Prod. i. 267, cum syn.—c.p. 2661.

HAB. Very abundant, up to an elevation of 2000 feet. Nom. vulg. "Poos-wel."

The tough bark of this plant is used for making cord and ropes: the juice of the leaves is employed for stupefying fish.

65. **ADENANTHERA**, Linn.

1. **A. pavonina**, Linn. (W. et A. Prod. i. 271, cum syn.)—c.p. 3605.

HAB. Warmer parts of the island; very common, but generally in or near native gardens. Nom. vulg. "Madateya-gass."

2. **A. bicolor**, Moon's Cat. p. 34 (sine syn.); *foliis* bipinnatis, glabris; *pinnis* distantibus, 1-3-jugis, suboppositis, apiculo parvo terminatis, 6-8-foliolatis; *foliolis* alternis, ovato-lanceolatis, minute apiculatis; *leguminibus* linearibus, varie contortis; *seminibus* rotundatis, compressis, nitidis, dimidiato-bicoloribus.—c.p. 324.

HAB. Not uncommon in the south of the island, at no great elevation. Nom. vulg. "Mas-moroo-gass."

*Arbor* mediocris, 15-20-pedalis. *Ramuli* teretes, glabri, nitidi, infra nodos bicallosi, cum racemis foliorumque rachi, atro-rubri. *Petioli* basi tumidi, superne minute glandulosi, subsulcati. *Stipulæ* minutissimæ, subulatæ. *Foliola* 1-1½ poll. longa, 6-9 lin. lata, superne nitida, *subtus* opaca, pallidiora, rufescentia, costa nervisque rubris, *petiolulo* tumidulo 1 lin. longo. *Racemi* axillares, multiflori, folio multum breviores, parce fugaci-pilosi. *Pedunculi* 1½ lin. longi, infra medium articulati, basi 1-bracteolati. *Flores* glabri, rubri, 1 lin. longi. *Calyx* minutus, acute 5-dentatus. *Petala* lanceolata. *Stamina* glandula terminali alba, polline flavo. *Ovarium* stipitatum, circiter 10-ovulatum. *Legumina* 5-6 lin. lata. *Semina* 4 lin. in diam., coccinea, dimidio apicali nigro, *funiculo* 2 lin. longo.

66. **DESMANTHUS**, Benth.

Hook. Journ. of Bot. vol. iv. p. 356.

1. **D. virgatus**, Willd. (W. et A. Prod. i. 270, cum syn.; Walp. Ann. iv. 614, cum syn.)—c.p. 3606.

HAB. Peradenia, *Alwis*. Probably introduced.

## 67. NEPTUNIA, Lour.

1. **N. oleracea**, Lour. (Benth. in Hook. Journ. of Bot. iv. 354; Walp. Rep. i. 863, cum syn.)—c.p. 1516.

HAB. Very common in shallow water, and on the borders of tanks in the hotter parts of the island. Nom. vulg. "Gass-nidi-koomba."

MIMOSA PUDICA, *Linn.*, is become quite naturalized, and a troublesome weed in some parts of the island.

## 68. DICHROSTACHYS, DC.

1. **D. cinerea**, W. et A. Prod. i. 271, cum syn.; Benth. in Hook. Journ. of Bot. iv. 353; Wight, Icones, t. 357.—c.p. 1523.

HAB. Abundant in the hot, drier parts of the island. Nom. vulg. "Andara-gass."

## 69. ACACIA, Willd.

1. **A. eburnea**, Willd. (W. et A. Prod. i. 276, cum syn.; Benth. in Lond. Journ. of Bot. i. 501; Walp. Rep. i. 912.)—c.p. 1521.

HAB. North of the island, *Gardner*.

2. **A. tomentosa**, Willd. (W. et A., l. c.; Benth. l. c.; Walp. Rep. l. c.)—c.p. 744.

HAB. North of the island, *Gardner*.

3. **A. leucophloea**, Willd. (W. et A., l. c. p. 277, cum syn.; Benth. l. c. 503; Walp. l. c.)—c.p. 1520.

HAB. Hot, drier parts of the island. Nom. vulg. "Katoo-andara-gass."

4. **A. Catechu**, Willd. (W. et A., l. c. p. 272, cum syn.)—c.p. 3547.

HAB. Hot, drier parts of the island. Nom. vulg. "Rat-kihiri-gass."

An infusion of the wood of this tree is much esteemed by the Cinghalese as a purifier of the blood; and drinking-cups are often made of it.

5. **A. concinna**, DC. (Benth. l. c. p. 514; Walp. l. c. p. 916; W. et A., l. c. p. 277, cum syn.)—c.p. 1519.

HAB. Very common, up to an elevation of 3000 feet. Nom. vulg. "Hingooroo-wel."

6. **A. Intsia**, Willd. (Benth. l. c. p. 515; Walp. l. c.; W. et A., l. c. p. 278.)—c.p. 3607 (1519 *partim*).

HAB. Hot, drier parts of the island.

7. **A. pennata**, Willd. (Benth. l. c. p. 516; Walp. l. c. p. 917; W. et A., l. c. p. 277, cum syn.)—c.p. 3300.

HAB. Haragam, near Kandy.

My specimens of this plant are nearly free from prickles.

ACACIA FARNESIANA, *Willd.*, has become quite naturalized, and very abundant in some of the hotter parts of the island.

## 70. ALBIZZIA, Durazzini, Benth.

1. **A. Lebbek**, Benth. Hook. Journ. of Bot. iii. 87; Walp. Rep. v. 596.—*Acacia speciosa*, Willd.; W. et A. Prod. i. 275, cum syn.—c.p. 3130.

HAB. Anooradhapoorā, *Mr. Brodie*; banks of the Ooma Oya, in the Badulla District.

2. **A. odoratissima**, Benth. l. c. p. 88; Walp. l. c.—*Acacia odoratissima*, Willd.; W. et A., l. c. cum syn.—c.P. 1529.

HAB. Warmer parts of the island, up to an elevation of 2000 feet. Nom. vulg. "Hooree-mārā-gass."

3. **A. amara**, Boivin, (Benth. l. c. p. 90; Walp. l. c. p. 597.)—*Acacia amara*, Willd.; W. et A., l. c. p. 274, cum syn.—c.P. 1518.

HAB. North of the island, *Gardner*.

4. **A. stipulata**, Benth. l. c. p. 92; Walp. l. c. p. 598, cum syn.—*Acacia stipulata*, DC.; W. et A., l. c.—c.P. 1517.

HAB. Warmer parts of the island, up to an elevation of 2000 feet. Nom. vulg. "Cabal-mārā-gass."

## 71. PITHECOLOBIUM, Martius.

Sectio *Concordia*, Benth. in Hook. Journ. of Bot. vol. iii. p. 202.

1. **P. geminatum**, Benth. l. c.; Walp. Rep. v. 611.—*Inga geminata*, W. et A. Prod. i. 269, cum syn.—c.P. 1531.

HAB. Jaffna, *Gardner*; Bintenne District.

2. **P. umbellatum**, Benth. l. c.; Walp. l. c.—*Inga umbellata*, Willd.; W. et A., l. c. p. 270, cum syn.—c.P. 1530.

HAB. North of the island, *Gardner*; Teldenia.

Sectio *Clypearia*, Benth. l. c. p. 207.

3. **P. bigeminum**, Mart. (Benth. l. c.; Walp. l. c.)—*Inga bigemina*, Willd.; W. et A., l. c. p. 269, cum syn.—c.P. 644.

HAB. Very abundant in the warmer parts of the island, and up to an elevation of 4000 feet. Nom. vulg. "Calateya-gass."

*Legumina* interne flavo-rufa. *Semina* oblonga, 7 lin. longa, 5 lin. lata; *testa* membranacea, nigra, subnitida, interne fusca; *funiculo* tenui, nigro, 2 lin. longo. *Embryo* mollis, graveolens; *cotyledonibus* crassis, *radiculam* includentibus, basi fissis.

4. **P. subcoriaceum**, Thw.; *ramulis*, *paniculis* *petiolisque* fusco-puberulis; *pinnis* 3-jugis; *foliolis* 15-20-jugis, subcoriaceis, lineari-oblongis, basi apiceque valde obliquis, subtus fusco-puberulis; *glandulis* clypeiformibus; *petiolaribus* sæpissime 2, inter se et petioli basi apiceque distantibus; *jugalibus* solitariis, inter pinnarum paria mediis.—c.P. 337.

HAB. Central Province, at an elevation of 4000 to 6000 feet; not uncommon. Nom. vulg. "Meemini-mārā."

*Arbor* ingens, 30-40-pedalis. *Ramuli* subangulati. *Foliola* 2-7 lin. longa,  $1\frac{1}{4}$ -2 lin. lata, subtus pallidiora. *Capitula* 3-12-flora. *Flores* puberuli, pallide virides, 1-bracteolati. *Corolla* calyce tubuloso 5-dentato plus duplo longior. *Stamina* numerosa, longe exserta. *Ovarium* longe stipitatum, pubescens. *Legumen* puberulum, cochleatum, circiter 6-spermum.

## LXI. ROSACEÆ.

### 1. PHOTINIA, Lindl.

1. **P. Notoniana**, W. et A. Prod. i. 302; Wight, Illust. t. 86; Icones, t. 991.—c.P. 136.

HAB. Central Province, at an elevation of 6000 to 7000 feet.

The usually bilocular ovary of this plant would seem to distinguish it principally from *Photinia integrifolia* of Lindley, in which the ovary is said to be "semper triloculare."

## 2. RUBUS, Linn.

### Sect. 1. *Foliis pinnatis.*

1. **R. lasiocarpus**, Smith, (W. et A. Prod. i. 299, cum syn.; Wight, Icones, t. 232.)—C.P. 1533, 1534.

Var.  $\beta$ . *subglaber*.—*R. leucocarpus*, Arn. Pug. p. 16 (334); Walp. Rep. ii. p. 19.

HAB. Central Province, up to an elevation of 6000 feet.

This species, like many other plants previously referred to, has, when growing in the more elevated districts of the island, somewhat larger flowers than when it occurs nearer the level of the sea.

### Sect. 2. *Foliis 3-foliolatis.*

2. **R. flavus**, Ham. (Don, Prod. Fl. Nep. 234; DC. Prod. ii. 559.)—*R. Goureephul*, Roxb.; W. et A., l. c. 298, cum syn.; Wight, Icones, t. 230.—C.P. 2596.

HAB. Central Province, at an elevation of 4000 to 7000 feet.

### Sect. 3. *Foliis simplicibus.*

3. **R. rugosus**, Smith, (W. et A. l. c.)

Var.  $\alpha$ ; *bracteis* ovatis, integris vel ad apicem plus minus incisodentatis; *fructibus* parvis, rubris. Wight, Icones, t. 225.—C.P. 2765.

Var.  $\beta$ ; *foliis* minoribus acutioribusque; *stipulis bracteisque* profunde capillaceo-incisis; *petalis* parvis; *fructibus* mediocribus, atorubentibus.—*R. micropetalus*, Gardn. in Calc. Journ. of Nat. Hist. vol. viii. p. 6.—C.P. 1536, 1537.

Var.  $\gamma$ ; *tomento* pallido; *foliis* valde coriaceis rugosisque; *stipulis bracteisque* plus minus profunde incisis; *paniculis* densis; *calycis* lobis ovato-lanceolatis, conspicue dentatis; *fructibus* mediocribus, atro-sanguineis.—*R. Fairholmianus*, Gardner, l. c. p. 5.—C.P. 134.

Var.  $\delta$ ; *aculeis* brevibus; *stipulis bracteisque* plus minus profunde incisis vel inciso-dentatis; *paniculis* paucifloris; *floribus* majoribus; *petalis* calyce subæquilongis; *fructibus* majoribus nigris.—*R. macrocarpus*, Gardner, l. c. p. 7.—C.P. 1535.

HAB. Var.  $\alpha$ . Not uncommon at an elevation of 1000 to 5000 feet. Var.  $\beta$ . Rambodde, at 5000 feet. Var.  $\gamma$ . Bopatalawe Plains (Gardner), and Oova District, at 6000 feet. Var.  $\delta$ . Newera Ellia, etc., at 6000 to 7000 feet.

The most characteristic forms of the above varieties might, in a less variable genus, pass very well for species; but, after an examination of numerous specimens, I find it quite impossible to draw up such distinctive definitions of them as could with propriety be considered specific, and therefore think it safer to describe them as forms of one variable species, which I believe them to be.

## 3. POTENTILLA, Linn.

1. **P. Mooniana**, Wight, Icones, t. 233.—? *P. polyphylla*, Wall. Cat. 1026; Lehm.; Walp. Rep. ii. p. 30.—C.P. 2767.

HAB. Horton Plains, at an elevation of 7000 feet.

*Herba* perennis, stolonifera. *Caules* suberecti, elongati, patentim pilosi. *Folia* interrupte pinnata, pilosa. *Foliola* numerosa, (majora oblonga, minora orbicularia,) subsessilia, acute dentata, subtus pallidiora, costa venisque longe adpresse pilosis. *Achenia* lævia.

In my specimens of this plant the bracteolæ are entire and smaller than the calyx-lobes, and not, as in Dr. Wight's figure, larger and toothed. *P. polyphylla*, Wall., would seem to differ principally in being quite glabrous and in the leaflets being crenately and not acutely toothed. *P. splendens*, Don, Prodr. Fl. Nep. p. 230, is evidently closely allied to these.

2. **P. Kleiniana**, W. et A. Prod. i. 300; Wight, Illust. t. 85.—C.P. 2768.

HAB. Central Province, at an elevation of 6000 to 7000 feet.

#### 4. AGRIMONIA, Tournef.

1. **A. Eupatoria**, Linn. (Wight, Icon. t. 224.)—*A. Zeylanica*, Moon's Cat. p. 37.—C.P. 2769.

HAB. Central Province, at an elevation of 5000 to 7000 feet.

#### 5. ALCHEMILLA, Tournef.

1. **A. vulgaris**, Linn. (Wight, Icones, t. 229.)

Var.  $\beta$ . *sarmentosa*.—*A. Indica*, Gardner, in Calc. Journ. of Nat. Hist. vol. viii. p. 8.—C.P. 2770.

HAB. Central Province, at an elevation of 5000 to 7000 feet; not uncommon.

#### 6. POTERIUM, Linn.

1. **P. Indicum**, Gardner, in Calc. Journ. of Nat. Hist. vol. viii. p. 10; "*caulibus* angulatis, basi ferrugineo-villosis; *foliolis* ovatis, obtusis, grosse dentato-serratis, basi subcordatis; *inferioribus* multo minoribus; *capitulis* polygamis; *bracteis* ciliatis; *calycibus fructiferis* osseo-induratis, reticulato-rugulosis, 4-angularibus; *angulis* alatis." Gardner.—C.P. 2771.

HAB. "Adam's Peak, 1835. Mr. Alwis," Gardner.

I have searched in vain for this plant in my visits to Adam's Peak, and Mr. Alwis has now no vivid recollection of gathering it there. The specimens in this Herbarium are in a very imperfect condition, and bear a considerable resemblance to *P. Sanguisorba*, Linn.

#### 7. PYGEUM, Gærtner.

1. **P. Ceylanicum**, Gærtner. Fruct. i. 218. t. 46. — *P. Walkerii*, Blume; Walp. Ann. iv. 642.—*Polyodontia? Walkerii*, Wight, Illust. i. 203.—? *Pygeum acuminatum*, Colebr. Linn. Tr. xii. 360. t. 18.—C.P. 1532.

HAB. Warm, moister parts of the island, up to an elevation of 3000 feet. Nom. vulg. "Kankoombalakatteya-gass," "Gal-mora-gass."

*Arbor* majuscula. *Folia* 4-7 poll. longa, 2-3 poll. lata, *petiolo* circiter  $\frac{1}{2}$  poll. longo. *Racemi* foliis breviores. *Drupæ* brevi-tomentosæ, denique subglabræ.

2. **P. Wightianum**, Blume, Walp. l. c.—*Polyodontia? Ceylanica*, Wight, l. c.; Icones, t. 256.—C.P. 638.

Var.  $\beta$ . *parvifolium*.—*P. parvifolium*, Gardn. MSS.—c.p. 1596.

HAB. Central Province, at an elevation of 4000 to 8000 feet. Nom. vulg. "Oonoonoogass."

*Arbor mediocris. Folia ovata vel ovato-lanceolata, brevi-acuminata, glabra, 1½–5 poll. longa, ¾–2½ poll. lata, petiolo 2½–4 lin. longo. Racemi pilosi, foliis breviores. Drupæ glabræ, transverse oblongæ.*

The ripe seeds of this and of the preceding species have, when bruised, a very powerful odour of prussic acid.

## LXII. COMBRETACEÆ.

### 1. COMBRETUM, Löffl.

1. **C. Wightianum**, Wall. (W. et A. Prod. i. 317; Wight, Icones, t. 227.)—c.p. 1601, 1602.

HAB. Not uncommon up to an elevation of 3000 feet. Nom. vulg. "Kadoorookettya-wel."

When growing on the hills, this plant has somewhat larger flowers than when it occurs near the level of the sea.

### 2. LUMNITZERA, Willd.

1. **L. racemosa**, Willd. (W. et A. Prod. i. 316, cum syn.)—c.p. 1594.

HAB. Seacoast; not uncommon. Nom. vulg. "Baireya-gass."

### 3. CONOCARPUS, Gærtner.

1. **C. latifolia**, Roxb. (W. et A. Prod. i. 316, cum syn.; Wight, Icones, t. 994.)—*Anogeissus*, Wall.—c.p. 1220.

HAB. Open, grassy places, in the district north of Kandy, up to an elevation of 1500 feet. Nom. vulg. "Daawoo-gass."

### 4. TERMINALIA, Linn.

#### Sect. 1. *Myrobalanus*.

1. **T. Belerica**, Roxb. (W. et A. Prod. i. 313, cum syn.)—c.p. 1605.

HAB. Open, grassy places, up to an elevation of 2000 feet. Nom. vulg. "Booloo-gass."

2. **T. Chebula**, Retz, (W. et A., l. c. cum syn.)—c.p. 1604.

HAB. Similar places with the last. Nom. vulg. "Araloo-gass."

The drupes of this and of the preceding species are much valued as a medicine, and are sold in the bazaars.

3. **T. parviflora**, Thw.; *foliis suboppositis, oblongis, abrupte acuminatis, integris, glabris, subtus minutissime reticulatis petioloque eglandulosis; spicis pilosulis, terminalibus, paniculatis, foliis subæquilongis; drupis ovoideis, carnosis, subpentagonis; putamine profunde lateque 5-sulcato*.—c.p. 618, 646.

HAB. Margins of woods in the Central Province, up to an elevation of 4000 feet; abundant in the Ambagamowa District. Nom. vulg. "Hampalanda-gass."

*Arbor* magna. *Ramuli* petiolique juniores fugaci-ferrugineo-pilosi. *Folia* 3-5 poll. longa,  $1\frac{1}{2}$ - $2\frac{1}{2}$  poll. lata, *petiolo* 3-5 lin. longo. *Flores* parvi, albidi. *Calyx* campanulatus, brevi-dentatus, externe pilosulus, interne villosus. *Drupæ* glabræ, virides, pollicares.

Sect. 2. *Pentaptera*.

4. **T. glabra**, W. et A., l. c. p. 314, cum syn.—c.p. 1603.

HAB. Very abundant on the banks of rivers, up to an elevation of 2000 feet. Nom. vulg. "Koombook-gass."

In my specimens the ramuli and the under side of the young leaves are puberulous, furnishing an additional reason for believing *Pent. tomentosa*, Roxb., to be merely a variety of the present plant, as suspected by Dr. Wallich.

LXIII. MELASTOMACEÆ.

Sect. *Melastomeæ*.

1. **OSBECKIA**, Linn.

(Nom. vernac. "Bowitteya," "Heen-bowitteya.")

† *Floribus tetrameris*.

1. **O. Leschenaultiana**, DC. Prod. iii. 142; Wight, Icon. t. 996; Spic. Nilgh. i. t. 68; non Naudin, Ann. des Sci. Nat. ser. 3. vol. xiv. p. 65 (quoad descript.).—*O. truncata*, Don; W. et A. Prod. i. 322; Arnott in Comp. to Bot. Mag. ii. 308; Walp. Rep. ii. pp. 135, 136. *O. parvifolia*, Arn. l. c.; Walp. l. c.—*O. erythrocephala*, Naud. l. c. p. 58.—c.p. 1576 (P c.p. 284).

HAB. Abundant in the Central Province, up to an elevation of 7000 feet.

A most variable species as regards the amount of pubescence and the size of the leaves, and I am unable to divide it into even distinct varieties, so completely do the several forms merge one into the other when a large series is examined. A specimen in the Herbarium, numbered c.p. 284, is, I suspect, a hybrid between the present species and *O. aspera*, some of its flowers having eight and others ten stamens, with anthers much longer than those of *O. Leschenaultiana*. In habit and pubescence it resembles the last-named species, but in the shape of the leaves and in other respects it is just intermediate between the two.

2. **O. Zeylanica**, Linn. f. (W. et A., l. c. cum syn.; non Naudin, l. c. p. 71).—*O. serialis*, Naud. l. c. p. 67.—c.p. 1575.

HAB. Amongst grass, in sandy soil: in the hot, drier parts of the island.

3. **O. Rheedii**, Gardn. MSS.; *foliis* rigidis, 3-nerviis, sparse adpresse setulosis, ovato-lanceolatis, basi plus minus angustatis, subtus pallidioribus; *floribus* ternis, subsessilibus; *calycis* strigosi dentibus triangularibus, tubo subæquilongis; *antheris* oblongis, obtusis, 1-porosis; *connectivo* infra loculos conspicue producto.—c.p. 129 (441).

HAB. Adam's Peak, near the summit.

*Frutex* parvus. *Ramuli* tetragoni, scabri. *Folia*  $\frac{3}{4}$ - $1\frac{1}{2}$  poll. longa, 4-7 lin. lata, subtus ad nervos scabra, *petiolo* scabro  $1\frac{1}{2}$ -2 lin. longo. *Petala* rotundata, acuminata, pallide purpurea, circiter 4 lin. longa. *Ovarium* 4-loculare, loculis circiter 8-ovulatis.

†† *Floribus pentameris.*

4. *O. Walkeri*, Arnott, l. c. p. 309; Walp. Rep. ii. 137.—c.p. 1571.

HAB. Central Province, at an elevation of 6000 to 8000 feet.

5. *O. buxifolia*, Arnott, l. c.; Walp. l. c. 136; Naud. l. c. p. 63.—c.p. 1572.

Var. *β. minor*; *foliis floribusque* minoribus; *calyce* dense sericeo-villoso, nec stellatim piloso.—c.p. 2618 (forsan species distincta).

HAB. Central Province, at an elevation of 6000 to 8000 feet.

If some of the species of this genus did not individually vary so much in the size of the leaves and flowers, and in the clothing of the calyx, I should without hesitation have made a separate species of var. *β*, but it bears so great a resemblance to var. *α* in all other than the above respects, that I cannot venture to divide them.

6. *O. aspera*, Blume.—*Melastoma aspera*, Linn. (*partim*).

Var. *α*; *calyce* setuloso, apicem versus parce squamuloso.—*O. aspera*, Bl.; W. et A., l. c. 323; Naudin, l. c. p. 74.—c.p. 1568.

Var. *β*; *calyce* setuloso, apicem versus appendiculis linearibus stellato-capitatis squamisque penicillatis parce ornato.—*O. Kleinii*, W. et A., l. c.; Arnott, l. c. p. 309.—c.p. 1569.

Var. *γ*; *floribus* majoribus; *calyce* sericeo-crinito; cæteroquin ut in var. *β*.—c.p. 506 (600, 1573).

Var. *δ*; *foliis* ellipticis, obtusiusculis, subcoriaceis; *paniculis* sæpe abbreviatis; *calyce* copiosius appendiculato; cæteroquin ut in var. *γ*.—*O. Kotigueda*, Naud. l. c. 64. ? *O. Wightiana*, Benth.; Wight, Icones, t. 998; Arnott, l. c.—c.p. 1570.

HAB. Var. *α* and *β*. Warmer parts of the island, up to no great elevation. Var. *γ*. Central Province, from 2000 to 4000 feet. Var. *δ*. From 4000 to 5000 feet.

The several varieties enumerated above, so run one into another, that I do not feel justified in constituting distinct species of them. Of the plants referred to by Linnæus as representing his *Melastoma aspera*, Herman's would appear to be the only one corresponding with any form of the present species, for the others are mentioned as having succulent fruit, and the one described by Burman was probably a state of *Melastoma Malabathricum*.

7. *O. rubicunda*, Arn. l. c.; Walp. l. c.—*O. elliptica*, Naud. l. c. 62.—c.p. 52 (208).

Var. *β*; *ramis* patentim pilosis, hispida; *calyce* (cum appendiculis) hispidissimo.—*O. rubicunda*, Naud. l. c.

HAB. Central Province, at an elevation of 5000 to 7000 feet.

Closely allied to the previous species, and much resembling its var. *δ*, and is principally distinguished by its more rigid pubescence, more coriaceous leaves, and fewer, larger, sessile capitate flowers.

8. *O. Moonii*, Thw.; pilosula, repens, ad nodos radicans; *foliis* membranaceis, ovatis, acutis, longiuscule petiolatis; *paniculis* paucifloris, ramulos terminantibus; *bracteis* rotundatis, ciliatis; *calyce* hemisphærico, appendiculis linearibus capitatis setosis, hispido; *dentibus* lanceolatis, tubo æquilongis, apice appendiculatis; *antheris* subulatis; *connectivo* ad basin parum producto.—*Melastoma repens*, Moon's Cat. 35, non Lam.—c.p. 2800.

HAB. Colombo, Moon. Saffragam District, rather rare.

*Folia* 3-nervia, 1–2¼ poll. longa, ½–1¼ poll. lata, *petiolo* 2–4 lin. longo. *Bractea* 1 lin. longæ. *Flores* mediocres, purpurei, *pedicellis* 1½ lin. longis.

9. *O. virgata*, Don, (W. et A., l. c. 323; Naud. l. c. 61).—*O. polyce-*



*phala*, Naud. l. c. 67. ? *O. cupularis*, Don; W. et A., l. c.; Naud. l. c. 61. ? *O. octandra*, DC. Prodr. iii. 142, cum syn.—c.p. 507, 2619.

HAB. Common up to an elevation of 3000 feet.

A very variable plant in the amount of pubescence upon the stems and leaves, and in the number of setose appendages to the calyx.

## 2. MELASTOMA, Linn.

1. **M. Malabathricum**, Linn. (W. et A. Prod. i. 324, cum syn.)—*M. ellipticum*, Naud. Ann. des Sci. Nat. ser. 3. vol. xiii. p. 291; Walp. Ann. ii. 567. *M. Royneii*, Blume; Walp. Repert. ii. 133.—c.p. 1574.

Var.  $\beta$ ; *ramulis* scaberrimis; *foliis floribusque* majoribus; *calycis* dentibus lanceolatis, tubo longioribus.—c.p. 1574 (2692).

HAB. Var. *a*. Common in the hotter parts of the island. Var.  $\beta$ . Up to an elevation of 3000 feet. Nom. vulg. "Bowitteya," "Mahabowitteya," "Katakaloowa."

The flat scales which clothe the calyx of var.  $\beta$  are shorter and less shining than in var. *a*: both sometimes occur with pale purple flowers. The fruit, when ripe, bursts irregularly, and exposes a dark-coloured pulp, covered with the small seeds: this pulp is eaten by children, and stains the inside of their mouths of a black colour.

## 3. MEDINILLA, Gaud.

? *Triplectrum*, Don, MSS., W. et A.

1. **M. maculata**, Gardner in Calc. Journ. of Bot. vol. viii. p. 13; *foliis* ovatis vel ovato-lanceolatis, subcordatis, crenatis, apice retusis; brevissime petiolatis, 3-nerviis; *pedunculis* axillaribus, 1-3-floris; *floribus* parvis, 4-meris; *antheris* basi postice calcaratis, antice bi-uncinatis.—c.p. 1578.

Var.  $\beta$ . *cuneata*; *foliis* majoribus, rotundato-cuneatis, abrupte breviacuminatis vel retusis.—c.p. 3443. (An species distincta?)

HAB. Var. *a*. Central Province, at an elevation of 3000 to 5000 feet. Var.  $\beta$ . Hellessee, Pasdoon Corle.

*Frutex* epiphytus. *Ramuli* 4-goni, angulis undulatis. *Folia* subcarnosa, superne glabra, subtus pilis minutissimis glandulosis conspersa,  $\frac{3}{4}$ -1 $\frac{3}{4}$  (var.  $\beta$ , 1 $\frac{1}{2}$ -2 $\frac{1}{2}$ ) poll. longa, 4-12 lin. (var.  $\beta$ ,  $\frac{3}{4}$ -2 poll.) lata. *Pedunculi* folio multo breviores. *Calyx* urceolatus, minutissime 4-dentatus, carnosus. *Petala* pallide rosea, subrotundata, reflexa, circiter 2 $\frac{1}{2}$  lin. longa. *Antheræ* subulatæ. *Baccæ* rubræ.

2. **M. fuchsioides**, Gardner, l. c.; *foliis* 3-5-nerviis, rhomboideo-oblongis, basi truncatulis, superne glabris, subtus sparse minutissime glandulosis, brevissime petiolatis; *pedunculis* in axillis foliorum delapsorum solitariis, 1-3-floris, folio multo brevioribus; *floribus* pendulis, 4-meris, petalis rotundatis, conniventibus; *antheris* subulatis, basi 3-tuberculatis.—c.p. 138.

HAB. Newera Ellia and other most elevated parts of the Central Province.

*Frutex* epiphytus, dichotome ramosus. *Ramuli* teretes, cinerei. *Folia* 2-3 poll. longa, 1-1 $\frac{1}{2}$  poll. lata, subtus pallidiora, costa rubra. *Inflorescentia* rubra. *Pedicelli* apicem versus incrassati, basi bibracteolati. *Flores* pollicares. *Calyx* urceolatus, 4-dentatus. *Stylus* filiformis. *Stigma* minutum. *Baccæ* subglobosæ, atro-rubræ, circiter 4 lin. in diam.

## 4. PACHYCENTRIA, Blume.

1. **P. Walkeri**.—*Medinilla?* *Walkeri*, Wight, *Illust.* i. 217.—*M. Walkerii*, Gardner in *Calc. Journ. of Nat. Hist.* vol. viii. p. 11; *foliis* 5-nerviis, oblongis vel ovato-oblongis, utrinque angustatis, obtusis; *junioribus* fugaci-ferrugineo-furfuraccis; *floribus* majusculis, speciosis, terminalibus, fasciculatis vel umbellatis, longe pedicellatis; *pedicellis* basi bracteatis.—C.P. 385.

HAB. Central Province, at an elevation of 3000 to 5000 feet.

*Frutex* epiphytus, scandens, copiose radicans. *Folia* 1½–3 poll. longa, ¾–1½ poll. lata, subtus pallidiora, subintegra, in plantis junioribus ciliato-dentata. *Petioli* 2–3 lin. longi. *Pedicelli* circiter 1½ poll. longi, cum bracteis calycibusque sparse ferrugineo-furfuracei. *Bracteæ* oblongæ, obtusæ, 2–3 lin. longæ. *Calyx* ruber, margine expanso 4-lobo carnosus. *Petala* carnosæ, ovato-rotundata, circiter 1½ poll. longa, interne sanguineo-rosea, externe pallidiora. *Stamina* æqualia, filamentis parce compressis; *antheris* sigmoideo-subulatis, compressis, postice acute calcaratis, 4 lin. longis. *Ovarium* calyce consolidatum, apice liberum, 4-loculare; *placentis* axilibus pluri-ovulatis. *Stylus* filiformis. *Stigma* minutum. *Capsula* unilocularis, apice 4-valvis, polysperma. *Semina* minuta, prismatica.

This is one of the most beautiful plants of the island, covering the trunks and branches of forest trees with a mantle of the most lovely rose-colour, when it is in flower.

## 5. SONERILA, Roxb.

† *Foliis* æqualibus, brevi- vel mediocri-petiolatis.

1. **S. robusta**, Arn. *Comp. to Bot. Mag.* ii. 308; *Walp. Rep.* ii. 122; *Naudin, Ann. des Sc. Nat.* ser. 3. vol. xv. p. 327.—C.P. 426.

HAB. Adam's Peak, at an elevation of 6000 to 7000 feet.

2. **S. Gardneri**, Thw.; erecta, sublignosa, hirsuta; *foliis* 5–7-nerviis, rotundatis vel ovatis, acutis, subcordatis, serrulatis, brevi-petiolatis; *pedicellis* 1-bracteatis; *bracteis* foliaceis; *floribus* majusculis; *antheris* cordato-lanceolatis, subacutis.—C.P. 63 (486).

HAB. Below Horton Plains, towards Galagama, on wet rocks, at an elevation of about 5000 feet.

*Caulis* teres, subsimplex, rufo-niger, glanduloso-hirsutus, 1–1½-pedalis. *Folia* ½–1¼ poll. longa, 4–8 lin. lata, utrinque hirsuta, subtus pallidiora; *petiolo* 1–3 lin. longo, glanduloso-hirsuto. *Pedicelli* calyce breviores et cum eo glanduloso-hirsuti. *Petala* sanguineo-rubra. *Capsula* ovoidæa, gibba, obscure 6-costata, muricato-hirsuta.

3. **S. Harveyi**, Thw.; erecta, sublignosa, superne ramosa, subglabra; *foliis* rotundatis vel ovatis, subacutis, 7-nerviis, setuloso-denticulatis, supra versus apicem prope marginem setulosis, cæteroquin glabris, basi subcordatis, brevi-petiolatis; *racemis* parce foliaceo-bracteatis; *antheris* breviusculis, cordato-acuminatis.—C.P. 2974.

HAB. Central Province, between Maturatte and Rambodde, at an elevation of 6000 feet, Oct. 1853.

1–2½-pedalis. *Rami* 4-goni, rubro-nigri, glabri vel parcissime glanduloso-pilosi, ad nodos setosi. *Folia* 3–10 lin. longa, subtus pallidiora; *petiolo* ½–1½ lin. longo. *Pedicelli* circiter 6, glabri. *Calyx* 6-costatus, glanduloso-pilosus; *dentibus* triangularibus, setosis. *Petala* purpurea, rotundato-acuminata.

4. **S. lanceolata**, Thw.; suberecta, parce ramosa, glaberrima; *foliis* 3-nerviis, lineari-lanceolatis, supra medium serrulatis, basi cordatis,

brevissime petiolatis; *racemis* simplicibus vel ramosis; *antheris* subulatis; *capsulis* longiusculis et cum pedicellis incrassatis profunde 6-sulcatis.—C.P. 3007.

HAB. Carawitte Kanda, near Ratnapoora, at an elevation of 1000 feet.

*Folia*  $1\frac{1}{2}$ –3 poll. longa, 3–8 lin. lata. *Petala* rubra, rotundata, apiculata.

5. **S. rhombifolia**, Thw.; suberecta, ramosa, glaberrima; *foliis* 3-nerviis, ovato-vel lanceolato-rhomboides, utrinque acutis, incurvo-denticulatis, petiolatis; *calycis* 6-costati dentibus longiusculis, acutis; *antheris* subulatis.—C.P. 2799.

HAB. Hinidoon and Kokool Corles, at no great elevation.

1–2-pedalis. *Rami* obscure 4-goni. *Folia*  $\frac{3}{4}$ – $1\frac{1}{2}$  poll. longa, 2–7 lin. lata, subtus rufescentia; *petiolo* 1–3 lin. longo. *Petala* rubra.

I have named this "*S. glaberrima*, Arn.," in some distributed collections, but it can scarcely be that species.

†† *Foliis plus minus inæqualibus basi que obliquis, longiuscule petiolatis.*

6. **S. Arnottiana**, Thw.; suberecta, ramosa, sublignosa; *foliis* 3–5-nerviis, lanceolatis vel ovato-lanceolatis, ciliato-denticulatis, parum obliquis inæqualibusque, superne glabris, subtus parce hirsutis; *petiolo*, cum nervis subtus, plus minus glanduloso-piloso; *racemis* simplicibus vel sæpius bifurcis, glanduloso-pilosis; *antheris* cordato-oblongis, brev-acuminatis.—C.P. 2615.

HAB. Central Province, at an elevation of 4000 to 7000 feet.

1–2-pedalis. *Rami* plus minus glanduloso-pilosi. *Folia*  $\frac{3}{4}$ –3 poll. longa, 4–10 lin. lata, subtus pallidiora, utrinque acuta vel basi parce rotundata. *Petala* ovato-acuminata, purpurea,  $3\frac{1}{2}$  lin. longa. *Capsulæ* hypocrateriformes, 6-costatæ, muricato-setosæ, *pedicello* subæquilongæ.

7. **S. Hookeriana**, Arn. l. c. p. 308; Walp. l. c.; Naud. l. c. p. 321.—C.P. 173.

HAB. Adam's Peak, on the Saffragam side, at an elevation of 5000 to 6000 feet.

*Racemi* simplices vel bifurci. *Petala* purpurea. *Antheræ* aurantiacæ.

8. **S. hirsutula**, Arn. l. c. p. 307; Walp. l. c.—? *S. Wightiana*, Arn. l. c.—C.P. 276.

HAB. Central Province, at an elevation of 6000 to 7000 feet.

*Petala* pallide purpurea.

9. **S. pilosula**, Thw.; suberecta, ramosa; *foliis* majusculis, penniveniis, ovato-lanceolatis, acuminatis, serrulatis, utrinque brevi-pilosulis, basi sæpissime valde obliquis, inæqualibus; *racemis* pilosulis; *antheris* longiusculis, cordato-rostratis; *capsulis* pilosis, hypocrateriformibus, 6-nervosis; *pedicellis* subæquilongis.—C.P. 2687.

HAB. Saffragam District, at an elevation of 1000 to 2000 feet.

$\frac{1}{2}$ – $1\frac{1}{2}$ -pedalis. *Rami* subteretes, pilosuli. *Folia*  $\frac{3}{4}$ –4 poll. longa,  $\frac{1}{2}$ – $1\frac{1}{2}$  poll. lata, subtus sæpe purpurascens. *Petala* rubra.

This species bears considerable resemblance to *S. versicolor*, Wight, Icones, t. 1057, but in that species the leaves are represented as scarcely oblique, the racemes glabrous, and the pedicels relatively shorter.

10. **S. rostrata**, Thw.; herbacea, erecta, ramosa; *foliis* 3–5-nerviis, ovato-lanceolatis, dentatis vel crenato-denticulatis, glabris vel sparse pilosis; *antheris* longis, rostrato-subulatis; *capsulis* brevibus, hypo-

crateriformibus, 6-nerviis, *pedicello* subæquilongis.—? *S. glaberrima*, Arn. l. c.—C.P. 2644.

HAB. Hinidoon and Reigam Corles, and Ambagamowa District, at an elevation of 1000 to 2000 feet.

$\frac{1}{2}$ –1-pedalis. *Rami* glabri, acute 4-goni. *Folia* 4–12 lin. longa, subtus rufescentia. *Petala* rubra, rotundato-acuminata, 4 lin. longa.

11. *S. affinis*, Arn. l. c.; Walp. l. c. 122.—C.P. 353.

HAB. Central Province, at an elevation of 4000 feet.

*Folia* 3–5-nervia, 2–10 lin. longa. *Petala* pallide rosea, rotundato-acuminata, 3 lin. longa. *Antheræ* cordato-subulatæ. *Capsulæ* *pedicellis* breviores, costulis parum prominentibus.

Closely allied to *S. rostrata* and *S. Ceylanica*, but distinguished from the former by its more sharply denticulated leaves, smaller, paler flowers, and less acuminate anthers; and from the latter by its subulate anthers and differently-shaped, less prominently costate capsules.

12. *S. Ceylanica*, W. et A. (Arn. l. c.; Walp. l. c.; Naud. l. c. 321.)—C.P. 349, 3345 (1577).

HAB. Central Province, at an elevation of 3000 to 5000 feet.

*Petala* purpurea vel albida. *Antheræ* cordatæ, subacuminatæ. *Capsulæ* longe hypocrateriformes, prominente 6-costatæ.

13. *S. tomentella*, Thw.; erecta, ramosa; *caule, ramis, petiolis racemisque* brevi-tomentosis; *foliis* ovato-lanceolatis, ciliato-denticulatis, utrinque sparse pilosis, costa venisque subtus tomentosis; *antheris* brevibus, oblongis, subapiculatis; *capsulis* hypocrateriformibus, setosis; *pedicellis* æquilongis.—C.P. 2616.

HAB. Ambagamowa District, at an elevation of 3000 to 4000 feet.

$\frac{1}{2}$ –1 $\frac{1}{2}$ -pedalis. *Folia* 3–12 lin. longa, basi rotundata, subpennivenia. *Petala* alba vel pallide rosea, rotundata, acuta, 3 $\frac{1}{2}$  lin. longa.

Very similar, in general appearance, to *S. Ceylanica*, but easily distinguishable by its tomentose stems and differently shaped anthers and capsules.

14. *S. pumila*, Thw.; erecta, sæpe ramosissima, multiflora; *foliis* 3–5-nerviis, rotundatis, ovatis vel ovato-lanceolatis, ciliato-denticulatis, superne sparse setosis et sæpissime albo-maculatis, subtus glabris; *racemis* paucifloris, sessilibus, glabris; *floribus* parvis; *antheris* brevibus, oblongis, obtusis, majuscule 2-porosis; *capsulis* longiusculis, hypocrateriformibus, 6-costatis.—C.P. 2617.

HAB. Newera Ellia and other more elevated parts of the Central Province.

1–10-pollicaris. *Rami* 4-goni, glabri. *Folia* 2–14 lin. longa; *petiolo* glabro. *Petala* alba. *Antherarum* loculi ab apice fere ad medium fissi.

The smaller, obtuse, large-pored anthers and less prominently costate capsules distinguish this from *S. Ceylanica* and other closely allied species.

15. *S. pedunculosa*, Thw.; herbacea, repens, radicans; *foliis* rotundatis vel ovatis, obtusis, crenatis, superne sparse villosis, subtus subglabris; *pedunculis* elongatis, paucifloris, sparse glanduloso-pilosis; *floribus* parvis; *antheris* cordato-acutis; *capsulis* turbinatis, 6-costatis.—C.P. 3091.

HAB. Hinidoon Corle, at an elevation of 1000 feet.

*Folia* viridi-rufescentia,  $\frac{1}{2}$ –1 $\frac{1}{4}$  poll. longa. *Petala* sanguinea, ovato-acuminata, 2 $\frac{1}{2}$  lin. longa.

This is a very distinct species, but my materials are not sufficient to enable me to give so complete a description of it as could be wished.

MARUMIA ZEYLANICA, *Blume*, has, I fear, no claim to be considered a native of Ceylon.

Sect. *Memecyleæ*.

6. MEMECYLON, Linn.

1. *Foliis 1-nerviis vel basi sub-3-nerviis*.

† *Disco calycino radiatim alato*.

1. **M. capitellatum**, Linn. (*Blume*, Mus. Bot. Lugd.-Bat. i. p. 354, cum syn., sed "Burm. Thes. Zeyl. p. 76. t. 30," vix excludend.)—C.P. 1564 (1565).

HAB. Not uncommon, up to an elevation of 1000 feet. Nom. vulg. "Welli-kaha."

2. **M. edule**, Roxb. Cor. Pl. i. t. 82.—? *M. Royneii*, *Blume*, l. c. 360.—C.P. 1566.

Var.  $\beta$ ; *foliis* majoribus, oblique acuminatis; *paniculis* elongatulis.—C.P. 3442.

Var.  $\gamma$ ; *foliis* elliptico-lanceolatis, basi subrotundatis; *paniculis* petiolo longiusculo subæquilongis.—? *M. Heyneanum*, Benth. Wight, Icon. t. 278.—C.P. 1563.

*Ramuli* teretes. *Folia* 2–6½ poll. longa, 1–3½ poll. lata, *petiolo* 2–4 lin. longo. *Pedicelli* calyce subæquilongi.

3. **M. sylvaticum**, Thw.; *ramulis* teretibus; *foliis* ovatis vel ellipticis, subabrupte acuminatis, basi in *petiolum* brevissimum vel subnullum angustatis, coriaceis, obscure penniveniis; *paniculis* brevibus, plus minus ramosis; *pedicellis* brevissimis, crassiusculis.—C.P. 1558, 1567, 3302 (2929).

HAB. Common in forests, at an elevation of 2000 to 4000 feet.

*Arbor* parva. *Folia* 1½–5½ poll. longa, ¾–2½ poll. lata. *Flores* cæruleo-purpurascens. *Baccæ* sphæricæ, atro-purpureæ, 4–5 lin. in diam.

Nearly allied to *M. edule*, from which it differs in its more abruptly acuminated leaves, shorter petioles and flowerstalks, and somewhat larger flowers.

4. **M. leucanthum**, Thw.; *ramulis* obscure 4-gonis; *foliis* subnitidis, rotundatis, ovatis vel ellipticis, abrupte breveque acuminatis, obtusis, basi acutis, brevi-petiolatis, venis primariis prominentibus; *paniculis* parum elongatis; *pedicellis* calyce dimidio longioribus.—C.P. 1559.

HAB. Central Province, at an elevation of 3000 to 5000 feet.

*Arbor* parva. *Folia* 1½–2½ poll. longa, ¾–1½ poll. lata, *petiolo* 1 lin. longo. *Flores* albi, petalis reflexis. *Baccæ* sphæricæ.

5. **M. rhinophyllum**, Thw.; *ramulis* obscure 4-gonis; *foliis* opacis, eveniis, ovatis vel ellipticis, valde acuminatis, basi acutis, subsessilibus; *paniculis* corymbosis vel umbellato-corymbosis, compactis, multifloris; *pedicellis* tenuibus, calyce subduplo longioribus; *floribus* parvis.—C.P. 2928.

HAB. Central Province, up to an elevation of 3000 feet.

*Arbor* parva. *Folia* (acumine ½-poll.) 2–3 poll. longa, ¾–1½ poll. lata. *Flores* pallide cyanæ. *Baccæ* sphæricæ, atro-purpureæ, 3 lin. in diam.

6. **M. macrocarpum**, Thw.; *ramulis* teretibus; *foliis* majusculis, valde coriaceis, rotundis, obtusis, basi plus minus angustatis, petiolatis; *pedunculis* brevibus; *pedicellis* calyce subæquilongis; *baccis* majusculis, sphæricis.—C.P. 2798.

HAB. Ambagamowa, at an elevation of about 3000 feet.

*Arbor parva. Folia* 3-4 poll. longa,  $2\frac{1}{2}$ - $3\frac{1}{2}$  poll. lata, *petiolo* 2 lin. longo. *Baccæ* (immaturæ) 8 lin. in diam.

†† *Disco calycino lævi vel obscure radiatim costulato.*

\* *Ramulis teretibus.*

7. **M. revolutum**, Thw.; *foliis* coriaceis, oblongo-rotundatis, obtusis, subcordatis, marginibus revolutis, brevi-petiolatis, venis primariis utrinque parum prominentibus; *pedunculis* multifloris, plus minus elongatulis ramosisque; *pedicellis* calyce dimidio longioribus; *floribus* parvis.—C.P. 2975.

HAB. Rambodde, at an elevation of 5000 feet.

*Arbor parva. Folia* 2-3 poll. longa, fere æquilata, *petiolo* 1 lin. longo. *Petala* alba. *Baccæ* nigro-purpureæ, 4-5 lin. in diam.

8. **M. umbellatum**, Burm. Fl. Ind. p. 87.—*M. ramiflorum*, Lam. DC. Prod. iii. p. 6; Wight, Illust. i. 215. t. 93. *M. umbellatum* et *M. ramiflorum*, Blume, l. c. pp. 863, 864, (quoad descript.)—C.P. 387, 764, 1561, 1562.

HAB. Very abundant, up to an elevation of 2000 feet. Nom. vulg. "Cora-caha."

The leaves of this species, mixed with the wood of *Morinda citrifolia* and *Cæsalpinia Sappan*, are employed by the Cinghalese for producing a permanent red dye.

9. **M. angustifolium**, Wight, Icones, t. 276.—C.P. 1556, 3346.

HAB. Common on the banks of rivers, up to an elevation of 2000 feet.

*Frutex* 6-8-pedalis. *Folia* evenia, superne nitidula, ad costam canaliculata, subtus opaca, pallidiora. *Pedunculi* ramosi, simplices vel subnulli. *Petala* cyanea. *Baccæ* pisi magnitudine, nigro-purpureæ.

10. **M. rostratum**, Thw.; *foliis* ellipticis, abrupte rostrato-acuminatis, obtusiusculis, basi acutis, inconspicue venosis, petiolatis; *pedunculis* multifloris, brevibus vel subnullis; *pedicellis* calyce 2-3-plo longioribus; *floribus* parvis.—C.P. 1560.

HAB. Central Province, at an elevation of 3000 to 4000 feet.

*Arbor parva. Folia* (acumine  $\frac{1}{2}$ -poll.)  $1\frac{1}{2}$ - $2\frac{1}{2}$  poll. longa,  $\frac{1}{2}$ - $1\frac{1}{4}$  poll. lata, *petiolo*  $1\frac{1}{2}$  lin. longo. *Petala* alba. *Antheræ* cyaneæ. *Baccæ* sphaericæ.

11. **M. fuscescens**, Thw.; *foliis* membranaceis, nitidulis, obscure venosis, ovatis vel ellipticis, basi plus minus angustatis, brevissime petiolatis; *paniculis* brevibus, parce ramosis, paucifloris; *pedicellis* subnullis.—C.P. 287.

HAB. Kokool Corle, at no great elevation.

*Folia* (acumine 4-6-lin.)  $1\frac{1}{2}$ -3 poll. longa, 6-11 lin. lata. *Baccæ* sphaericæ, 4 lin. in diam.

12. **M. macrophyllum**, Thw.; *foliis* magnis, coriaceis, cordato-vel subcordato-lanceolatis, brevissime petiolatis; *pedunculis* brevissimis vel subnullis; *pedicellis* brevibus.—C.P. 2685.

HAB. Ambagamowa District and Pasdoon Corle, at an elevation of 1000 to 3000 feet.

*Folia* 6-12 poll. longa,  $2-3\frac{1}{4}$  poll. lata. *Petala* pallide cyanea. *Baccæ* sphaericæ, 4 lin. in diam.

Closely allied to, and possibly a state of, the next species, from which it is principally distinguished by its larger leaves cordate at the base, and shorter petioles.

13. **M. jambosoides**, Wight, *Illust.* i. p. 215; *Icones*, t. 277.—C.P. 1555, 2468, 3347.

HAB. Damp, shady places, up to an elevation of 3000 feet.

*Petala* pallide cyanea. *Baccæ* sphaericæ, 4 lin. in diam.

14. **M. cuneatum**, Thw.; *foliis* coriaceis, opacis, eveniis, rotundatis vel oblongis, obtusis, basi cuneatis, brevi-petiolatis; *pedunculis* nullis; *floribus* confertis, brevissime pedicellatis; *calycis* margine subintegro, alabastris acutis.—C.P. 2538.

HAB. Central Province, at an elevation of 3000 feet.

*Arbor* parva. *Folia* 1-2½ poll. longa, ½-1½ poll. lata, *petiolo* 1 lin. longo. *Petala* pallide cyanea. *Baccæ* sphaericæ.

15. **M. orbiculare**, Thw.; *foliis* coriaceis, subnitidis, orbicularibus, obscure obtuseque acuminatis vel retusis, brevissime petiolatis, venis primariis vix conspicuis; *pedunculis* brevibus vel subnullis; *pedicellis* calyce subæquilongis.—C.P. 3441.

HAB. Hinidoon Corle, at an elevation of 1000 feet.

*Arbor* parva. *Folia* 1½-2 poll. longa. *Petala* cyanea.

16. **M. ovoideum**, Thw.; *foliis* majusculis, valde coriaceis, opacis, obovato-rotundatis, retusis, subsessilibus; *pedunculis* subnullis; *pedicellis* fasciculatis, confertis, calyce subæquilongis; *baccis* ovoideis.—C.P. 2621.

HAB. Ambagamowa District, near Adam's Peak, at an elevation of 4000 to 5000 feet.

*Arbor* parva. *Folia* 2-4 poll. longa, 1½-3 poll. lata. *Baccæ* 6 lin. longæ, 4 lin. latæ, nigro-purpureæ.

\*\* *Ramulis tetragonis*.

17. **M. ellipticum**, Thw.; *foliis* nitidulis, ellipticis, abrupte rostrato-acuminatis, obtusis, basi rotundatis vel parce angustatis, inconspicue venosis, brevissime petiolatis; *pedunculis* brevissimis vel subnullis; *pedicellis* confertis, calyce subæquilongis.—C.P. 3479.

HAB. Singhe Rajah Forest, between Galle and Ratnapoora, at no great elevation.

*Arbor* parva. *Folia* (acumine ½-poll.) 2-3 poll. longa, 1¼-1½ poll. lata; *petiolo* ½ lin. longo. *Petala* albida.

18. **M. elegantulum**, Thw.; *foliis* lanceolatis, longe acuminatis, obtusiusculis, basi plus minus angustatis, brevissime petiolatis, obscure venosis; *paniculis* filiformibus, subelongatis, parce ramosis, paucifloris; *pedicellis* subsolitariis, calyce dimidio longioribus.—C.P. 2684.

HAB. Saffragam District, at no great elevation.

*Folia* (acumine ½-¾-poll.) 2-3 poll. longa, 6-8 lin. lata. *Flores* albi. *Baccæ* sphaericæ, pisi magnitudine.

19. **M. varians**, Thw.; *foliis* opacis, brevi-petiolatis, obscure venosis; *pedunculis* brevibus vel parum elongatis; *floribus* capitellatis; brevissime pedicellatis.

Var. *a.* *acuminatum*; *foliis* ovatis vel ellipticis, acuminatis, basi acutis.—C.P. 2927.

Var. *β.* *rotundatum*; *foliis* rotundatis vel oblongis, utrinque obtusis.—C.P. 2959 (170).

HAB. Central Province. Var. *a*. At an elevation of 2000 to 4000 feet.  
Var. *β*. At 4000 to 5000 feet.

*Arbor parva. Folia* 1-3 poll. longa,  $\frac{1}{2}$ -1 $\frac{1}{2}$  poll. lata, *petiolo* 1-1 $\frac{1}{2}$  lin. longo. *Petala* cyanea vel raro albida. *Baccæ* sphaericæ, 3-4 lin. in diam.

20. **M. parvifolium**, Thw.; *foliis* coriaceis, opacis, eveniis, rotundatis vel oblongis, obtusis retusisve, basi cuneatis, brevi-petiolatis; *pedunculis* brevissimis vel subnullis; *floribus* capitellatis, subsessilibus.—C.P. 2955.

HAB. Central Province, at an elevation of 7000 feet.

*Arbor parva. Folia*  $\frac{1}{2}$ -1 poll. longa, 3-8 lin. lata, *petiolo*  $\frac{1}{2}$ -1 lin. longo. *Petala* albida. *Baccæ* sphaericæ.

Nearly allied to, and possibly a variety of, the preceding, from var. *β* of which it differs principally in its leaves being cuneate at the base and a good deal smaller,

\*\*\* *Ramulis 4-alatis.*

21. **M. Hookeri**, Thw.; *foliis* majusculis, cordato-lanceolatis, acuminatis, sessilibus, subtus costa et venis primariis intramarginalique valde prominentibus; *pedunculis* brevibus vel subnullis; *pedicellis* longiusculis.—C.P. 2686.

HAB. Ambagamowa District, in shady forests, on the banks of streams, at an elevation of 1000 feet.

*Frutex* erectus, parce ramosus, 8-12-pedalis. *Folia* 6-12 poll. longa, 1 $\frac{1}{2}$ -3 poll. lata. *Petala* cyanea. *Baccæ* sphaericæ, 4 lin. in diam., rufescentes, denique nigro-purpureæ.

22. **M. Wightii**, Thw.—? *M. amplexicaule*, Wight, Icon. t. 279 (vix Roxb. Fl. Ind. ii. 260.)—C.P. 2570.

HAB. Central Province, at an elevation of 3000 to 4000 feet.

*Arbor parva. Folia* subnitida, cordato-lanceolata, plus minus acuminata, obscure venosa, 2-5 poll. longa,  $\frac{3}{4}$ -1 $\frac{1}{2}$  poll. lata, *petiolo* 1 lin. longo. *Pedunculi* brevissimi vel subnulli. *Pedicelli* conferti, longiusculi. *Petala* cyanea. *Baccæ* sphaericæ, 3-4 lin. in diam.

The shape of the leaves and smaller size of the fruit seem to preclude the possibility of the present being Roxburgh's plant; and the winged ramuli and petioled leaves of our plant do not agree with Dr. Wight's description of his species, though his figure quoted above bears a very considerable resemblance to it.

2. *Foliis trinerviis.*

23. **M. Arnottianum**, Wight, MSS.—*Eugenia?* *capitellata*, Arn. Pug. 17 (335); Walp. Rep. ii. 182.—C.P. 1589 (1868).

HAB. Hinidoon and Pasdoon Corles, and Saffragam District, at no great elevation.

*Frutex* vel *arbor parva*, 10-12-pedalis. *Ramuli* teretes. *Folia* 1-4 poll. longa,  $\frac{1}{2}$ -2 poll. lata, juniora rufescentia. *Capitella* 3-5-flora, longe pedunculata. *Petala* albida. *Baccæ* sphaericæ, pisi magnitudine, nigro-purpureæ.

24. **M. Gardneri**, Thw.; *ramulis junioribus* anguste 4-alatis; *foliis* opacis, pellucido-punctulatis, rotundatis, ellipticis vel rhomboideo-oblongis, abrupte breveque acuminatis, obtusis retusisve, basi in *petiolum* brevem angustatis; *capitellis* 3-plurifloris, brevi-pedunculatis vel sessilibus.—C.P. 2387.

HAB. Central Province, at an elevation of 2000 to 5000 feet.

*Arbor parva. Folia* 1-3 poll. longa,  $\frac{1}{2}$ -1 $\frac{1}{2}$  poll. lata, *petiolo* 1 lin. longo. *Flores* albi. *Baccæ* sphaericæ, pisi magnitudine, nigro-purpureæ.



## LXIV. MYRTACEÆ.

## 1. MYRTUS, Tournef.

Sect. *Rhodomyrtus*, DC.

1. **M. tomentosa**, Ait. (W. et A. Prod. i. 328, cum syn.; Wight, Icones, t. 522; Illust. ii. 18. t. 97\* or 122. f. 3.)—C.P. 1591.

HAB. Common in the Central Province, at an elevation of 6000 to 8000 feet.

## 2. EUGENIA, Linn.

1. **E. Willdenovii**, DC. Pr. iii. 265, cum syn.; Wight, Icones, t. 545.—C.P. 1586.

HAB. Common in the hotter parts of the island. Nom. vulg. "Tambaleya-gass."

2. **E. bracteata**, Roxb. (W. et A. Prod. i. 331, cum syn.)—C.P. 1590.

HAB. Jaffna, *Gardner*.

This is, as remarked by Dr. Wight, very closely allied to the preceding species, from which it differs principally in its smaller flowers and wider leaves, with shorter petioles. They may possibly prove to be forms only of one species.

3. **E. terpnophylla**, Thw.; *foliis lanceolatis vel oblongo-lanceolatis, longe acuminatis, basi plus minus angustatis; junioribus cum ramulis inflorescentiaque ferrugineo-pilosis*.—C.P. 2623.

Var.  $\beta$ ; *foliis subtus plus minus breviter rufo-pilosis*.

HAB. Ambagamowa and Ratnapoora Districts, and Reigam Corle, up to no great elevation.

*Arbor mediocris. Ramuli teretes. Folia 2½–5 poll. longa, ½–2 poll. lata, petiolo ¼–½ poll. longo. Peduncululi axillares terminalesque, breves, pauciflori. Pedicelli petiolo subæquilongi. Petala alba, lobis calycinis rotundatis duplo longiora. Bacca sphaericæ, puberulæ, cerasi parvi magnitudine.*

4. **E. amœna**, Thw.; *foliis sessilibus, cordato-vel ovato-lanceolatis, acuminatis, glabris, subtus reticulatis; floribus sæpissime terminalibus, subfasciculatis; pedicellis folio multo brevioribus*.—C.P. 3439.

HAB. Kokool Corle and Dolosbage District, up to an elevation of 1500 feet.

*Arbor parva. Ramuli teretes. Folia 2–6 poll. longa, 1¼–2¼ poll. lata. Pedicelli 7–10 lin. longi, raro breviores. Petala 4, alba, oblonga, circiter 3 lin. longa, lobis calycinis rotundatis longiora.*

5. **E. Mooniana**, Wight, Illust. ii. 13; Icones, t. 551.—C.P. 11, 365 (77, 733, 1581, 2451).

Var.  $\beta$ ; *foliis subsessilibus, rotundatis, acuminatis; floribus sessilibus*.—C.P. 2803. (An species distincta?)

HAB. Abundant in the Central Province, up to an elevation of 4000 feet.

A most variable species in the shape of the leaves and the length of the petioles and flower-stalks. Var.  $\beta$  may possibly prove to be a distinct species, but I am disposed at present to consider it only a variety.

6. **E. mabœoides**, Wight, Illust. ii. 13.—C.P. 445, 1588 (2693).

HAB. Central Province, at an elevation of 4000 to 7000 feet.

*Folia*  $\frac{1}{2}$ –2 poll. longa,  $\frac{1}{4}$ – $1\frac{1}{4}$  poll. lata, dense glanduloso-punctata. *Baccæ* sphaericæ, cerasi parvi magnitudine.

7. **E. floccifera**, Thw.; *foliis* oblongo-lanceolatis, acuminatis; *junioribus* cum *ramulis inflorescentiaque* dense fulvo-tomentosis; *pedicellis* axillaribus, brevibus, fasciculatis; *calycis* lobis 4, lanceolatis, tubo multum longioribus.—C.P. 463.

HAB. Reigam Corle, at no great elevation.

*Arbor* parva. *Ramuli* teretes. *Folia* 4–8 poll. longa, 1–2 poll. lata, *petiolo* 3–4 lin. longo. *Petala* alba, 4 lin. longa, lobis calycinis æquilonga.

8. **E. fulva**, Thw.; *foliis* ellipticis vel oblongis, abrupte brevi-acuminatis, basi rotundatis, superne lævibus, subtus venosis; *junioribus* cum *ramulis inflorescentiaque* fulvo-tomentosis; *floribus* axillaribus caulinisque parvis, sessilibus, confertis.—C.P. 3008.

HAB. Pasdoon Corle, at no great elevation.

*Arbor* parva, *ramulis* teretibus. *Folia* 4–6 poll. longa,  $2\frac{1}{4}$ –3 poll. lata, *petiolo* 4 lin. longo.

9. **E. rivulorum**, Thw.; *foliis* majusculis, lanceolatis, acuminatis, basi angustatis, subtus minutissime fulvo-pilosis, denique glabris *floribus* majusculis, sessilibus, axillaribus caulinisque; *calycis* externe pilosi lobis 4, rotundatis, tubo æquilongis.—C.P. 3440.

HAB. Banks of streams, in the Singhe-rajah forest, between Galle and Ratnapoora, at no great elevation.

*Arbor* parva. *Ramuli* teretes. *Folia* 10–14 poll. longa,  $3-4\frac{1}{2}$  poll. lata, *petiolo* circiter  $\frac{1}{2}$  poll. longo. *Petala* oblonga, pallide purpurea, 5 lin. longa.

10. **E. decora**, Thw.; *foliis* oblongis, acuminatis, basi rotundatis, glabris, superne nitidis, subtus pallidioribus, minute reticulatis; *floribus* sessilibus, confertis, axillaribus caulinisque; *baccis* transverse oblongis, dispermis.—C.P. 3545.

HAB. Near Galle, at no great elevation.

*Arbor* parva, *ramulis* teretibus. *Folia* 4–5 poll. longa, circiter 2 poll. lata, *petiolo* 4 lin. longo. *Flores* albi. *Baccæ* rubro-nigrescentes, 6 lin. in alt., 8 lin. in lat. max.

When one seed only is produced, the berry would be no doubt spherical, and perhaps this will prove to be the most usual form.

### 3. JAMBOSA, Rumph.

1. **J. aquea**, DC. (W. et A. Prod. i. 332, cum syn.; Wight, Icones, t. 216.)—*Eugenia (J.) aquea*, Wight, Illust. ii. p. 14. *Eug. sylvestris*, Moon's Cat. p. 38.—C.P. 418 (609).

HAB. Abundant in the Central Province, up to an elevation of 5000 feet. Nom. vulg. "Wal-jambo-gass."

This species varies a good deal in the length of the calyx-tube and in the colour of the flowers, which latter are either of a pale-purple or more commonly of a deep-crimson colour.

2. **J. cylindrica**.—*Eug. (J.) cylindrica*, Wight, Icones, t. 527, et forsan *E. (J.) pauciflora*, Wight, Ic. t. 526.—C.P. 601.

HAB. Ambagamowa District, up to an elevation of 3000 feet.

*Arbor* mediocris. *Petala* alba, purpurascens. *Baccæ* subsphaericæ, urceolatæ, 6–8 lin. in diam.

4. **STRONGYLOCALYX**, Blume.

Mus. Bot. Lugd.-Bat. i. p. 89.

1. **S. hemisphæricus**, Bl. l. c. 90.—*Eug. (J.) hemisphærica*, Wight, Illust. ii. p. 14. t. 97\* or 122. fig. 8; Icones, t. 525.—C.P. 2450, 2540, 3438.

HAB. Central Province and Kokool Corle, up to an elevation of 4000 feet.

*Arbor mediocris. Folia* 2–6 poll. longa,  $\frac{3}{4}$ –2 poll. lata. *Petala* alba. *Baccæ* sphæricæ, circiter 1 poll. in diam.

5. **SYZYGIUM**, Gærtn.† *Cymis ad foliorum delapsorum axillas.*

1. **S. caryophyllifolium**, DC.—*Eug. (S.) caryophyllifolia* [*E. (S.) Jambolana?* var. *microcarpa*], Wight, Icones, t. 553. *Eug. caryophyllifolia*, Lam.; Roxb. Fl. Ind. ii. p. 486. *Calyptranthes Cumini*, Moon's Cat. p. 39.—C.P. 1584.

HAB. Common up to an elevation of 3000 feet. Nom. vulg. "Mā-dang-gass."

The fruit, which is of about the size of a pea, is eaten by the Çinghalese.

2. **S. polyanthum**.—*Eug. (S.) polyantha*, Wight, Illust. ii. p. 17; Icones, t. 543. An *Syz. balsameum*, Wall. Wight, Ill. ii. p. 16? *Calyp. caryophyllifolia*, Moon's Cat. p. 39.—C.P. 2801.

HAB. Common up to an elevation of 3000 feet. Nom. vulg. "Batta-domba-gass."

*Arbor magna. Baccæ* sphæricæ, nigrescentes, cerasi parvi magnitudine.

The leaves of this species are larger than those of the preceding species, with far fewer and more distant primary veins, and shorter petioles; the flowers also are larger. The fruit is not eaten.

†† *Cymis terminalibus axillaribusque.*\* *Ramulis junioribus teretibus.*

3. **S. cordifolium**.—*Eug. (S.) cordifolia*, Wight, Illust. ii. p. 16; Icon. t. 544.—C.P. 350, 2622.

HAB. Near Cattura, Moon; Ambagamowa and Ratnapoora Districts, up to an elevation of 3000 feet.

*Baccæ* subsphæricæ, pollicares.

4. **S. montanum**.—*Eug. (S.) montana*, Wight, Icon. t. 1060?—C.P. 2694.

HAB. Ambagamowa District, at an elevation of 4000 feet.

My specimens bear considerable resemblance to Dr. Wight's figure, but they are in too imperfect a state to enable me to feel certain they are identical with his species.

5. **S. sylvestre**.—*Eug. (S.) sylvestris*, Wight, Icon. t. 532. *Calyptranthes Jambolana*, Moon's Cat. p. 39.—C.P. 2862.

HAB. Common up to an elevation of 3000 feet. Nom. vulg. "Aloo-bo-gass."

*Arbor magna. Baccæ* sphæricæ, nigro-purpureæ, circiter  $\frac{1}{2}$  poll. in diam.

6. **S. assimile**, Thw.; *foliis* ovato-lanceolatis vel obovatis, plus minus

abrupte acuminatis, basi angustatis vel rarius rotundatis, petiolatis; cymis ad apices ramulorum confertis, folio dimidio brevioribus; baccis sphaericis.—An *Eug. (S.) Arnottiana*, Wight, Icon. t. 999, *varietas?*—C.P. 46, 634, 2449 (291, 1582, 2495).

HAB. Central Province, up to an elevation of 5000 feet, in open places.

*Arbor mediocris. Folia* 2–5 poll. longa,  $1\frac{1}{4}$ –2 poll. lata, *petiolo* 2–5 lin. longo. *Baccæ* atro-purpureæ, circiter  $\frac{3}{4}$  poll. in diam.

This closely resembles in general appearance Dr. Wight's figure of *Eug. (S.) Arnottiana*, quoted above, but the petals in our plant seem generally to fall off without expanding, and the fruit is spherical instead of being oblong.

7. **S. Neesianum**, Arn. Pug. (335) 17.—*Eug. (S.) Neesiana*, Wight, Icon. t. 533; Illust. ii. p. 15. *Calyptranthes cordifolia*, Moon's Cat. p. 39. An *Myrtus androsæmoides*, Vahl, Fl. Zeyl. n. 184?—C.P. 735 (763).

HAB. Common up to an elevation of 3000 feet. Nom. vulg. "Panookaara-gass."

8. **S. caryophyllæum**, Gært. (W. et A. Prod. i. 329.)—*Eug. (S.) caryophyllæa*, Wight, Illust. ii. p. 15; Icones, t. 540.—C.P. 1583.

HAB. Common up to an elevation of 3000 feet. Nom. vulg. "Dang-gass."

The berries, which are of about the size of a pea, are eaten by the Cinghalese.

9. **S. revolutum**.—*Eug. (S.) revoluta*, Wight, Illust. ii. p. 17; Icones, t. 534.—C.P. 2521.

HAB. Central Province, at an elevation of 3000 to 6000 feet.

*Arbor mediocris. Ramuli* aliquando obscure tetragoni. *Baccæ* sphaericæ, rubiginosæ, cerasi parvi magnitudine.

10. **S. micranthum**, Thw.; *foliis* ovatis vel ellipticis, obtuse rostrato-acuminatis, venosis, obscure punctulatis, basi angustatis, petiolatis; *cymis* folio subæquilongis; *floribus* parvulis; *calyce* acute 4-dentato.—C.P. 1580.

HAB. Central Province, at an elevation of 3000 to 5000 feet.

*Arbor mediocris. Folia* (acumine  $\frac{1}{2}$ – $\frac{3}{4}$  poll. longo) 2–4 poll. longa,  $\frac{3}{4}$ –2 poll. lata, *petiolo* 2–5 lin. longo. *Baccæ* sphaericæ, rubiginosæ, pisi magnitudine.

11. **S. Gardneri**, Thw.; *foliis* ovato-lanceolatis, rostrato-acuminatis, brevissimis, subtus delicatule parallelo-venosis, pellucido-punctatis; *cymis* folio brevioribus; *calyce* obscure 4-dentato.—C.P. 2496.

HAB. Central Province, at an elevation of 2000 to 4000 feet.

*Arbor mediocris. Folia*  $2\frac{1}{2}$ –4 poll. longa, 1– $1\frac{1}{2}$  poll. lata, *petiolo* tenui 4–5 lin. longo.

\*\* *Ramulis junioribus tetragonis.*

12. **S. lissophyllum**, Thw.; *foliis* ovato-lanceolatis, acuminatis, lævissimis, subtus delicatule venosis, punctulatis; *cymis* folio sæpissime longioribus; *dentibus calycinis* brevibus, rotundatis.—An "*Eug. (S.) cymosa*, Lam." Wight, Icon. t. 555?—C.P. 2452, 2453.

HAB. Forests in the Central Province, at an elevation of 3000 to 5000 feet.

*Arbor mediocris. Folia*  $1\frac{1}{2}$ –3 poll. longa, 6–13 lin. lata, *petiolo* 2–3 lin. longo.

Closely resembling the preceding species in the shape and venation of the leaves, but these are smaller and more shortly petioled in the present plant, which is moreover at once distinguished from the other by its quadrangular ramuli. *Eug. (S.)*

*toddalioides*, Wight, Icon. t. 542, must be very closely allied to this, but the longer-pedicelled flowers would seem, judging from the figure, sufficient to prove it distinct.

13. **S. umbrosum**, Thw.; *foliis* ovatis, breve et obtuse acuminatis vel retusis, basi angustatis, utrinque lævissimis, subeveniis; *cymis* folio longioribus.—C.P. 2539.

HAB. Open, grassy places in the Central Province, at an elevation of 5000 to 6000 feet.

*Arbor* mediocris, ramosissima. *Folia* 1-2 poll. longa, 6-10 lin. lata, *petiolo* 1½-2 lin. longo.

Differs from the last in the shape and venation of the leaves: the inflorescence of the two species is very similar.

14. **S. spathulatum**, Thw.; *foliis* parvis, spathulato-obovatis vel lanceolatis, obtusis, superne nitidis, subtus opacis, parce parallelo-venosis, dense glanduloso-punctatis; *cymis* folio subæquilongis vel brevioribus; *floribus* parvis, brevi-pedicellatis.—C.P. 2493 (2494).

HAB. Central Province, at an elevation of 3000 to 5000 feet.

*Arbor* parva. *Folia* 1-1¾ poll. longa, 4-8 lin. lata, *petiolo* 1-1½ lin. longo. *Baccæ* sphaericæ, rubro-nigrescentes, pisi magnitudine.

Apparently closely allied to *S. rubiginosum*, W. et A. Prod. i. 330.

15. **S. calophyllifolium**.—*Eug. (S.) calophyllifolia*, Wight, Icones, t. 1000.—C.P. 160.

HAB. Adam's Peak, at an elevation of 6000 to 7000 feet.

16. **S. sclerophyllum**, Thw.; *foliis* parvis, valde coriaceis, obovatis vel subrotundatis, sessilibus, cordatis, superne glabris, subtus venosis; *costa* ad basin incrassata; *cymis* folio subæquilongis.—C.P. 274.

HAB. Central Province, at an elevation of 6000 to 8000 feet.

*Arbor* mediocris. *Folia* 5-12 lin. longa, 4-10 lin. lata. *Baccæ* sphaericæ, nigrescentes, cerasi parvi magnitudine.

17. **S. rotundifolium**, Arn. Pug. (335) 17.—*Eug. (S.) rotundifolia*, Wight, Illust. ii. p. 17.—C.P. 1587.

HAB. Central Province, at an elevation of 6000 to 8000 feet.

*Arbor* parva. *Folia* orbicularia, obtusissima vel obscure acuminata, basi subcordata vel parum angustata, 4-8 lin. longa, *petiolo* ½-1½ lin. longo. *Baccæ* sphaericæ, rubro-nigrescentes, cerasi parvi magnitudine.

Nearly allied to the preceding, but I think certainly distinct.

18. **S. oliganthum**, Thw.; *foliis* parvis, obovatis, obtuse brevi-acuminatis vel retusis, basi angustatis, leviter venosis, punctatis; *capitulis* paucifloris, brevi-pedunculatis, terminalibus.—C.P. 452 (588).

HAB. Ambagamowa District, at an elevation of 3000 to 5000 feet.

*Arbor* parva. *Folia* 4-10 lin. longa, 3-6 lin. lata, *petiolo* 1 lin. longo. *Baccæ* sphaericæ, rubro-nigrescentes, pisi magnitudine.

## 6. ACMENA, DC.

1. **A. Zeylanica**.—*Eug. (A.) Zeylanica*, Wight, Illust. ii. p. 15, cum syn.; Icones, t. 73.—C.P. 56, 380 (470, 1579).

HAB. Common in the hot, drier parts of the island, up to no great elevation. Nom. vulg. "Marang-gass."

*Arbor* mediocris. *Folia* lineari-vel ovato-lanceolata, acuminata, basi angustata vel rotundata, dense glanduloso-punctata, 1-4 poll. longa, 1½-10 lin. lata, *petiolo* 1-3 lin. longo. *Calyx* valde glandulosus.

2. **A. lanceolata**.—*Eug. (A.) lanceolata*, Wight, *Illust.* ii. p. 15; *Icones*, t. 530. *Syzygium lanceolatum*, W. et A. *Prod.* i. p. 330, cum syn. An *Eugenia (A.) Wightiana*, Wight, *Illust.* ii. p. 15; *Icones*, t. 529, cum syn. ?—C.P. 2863.

HAB. Ambagamowa and Kornegalle Districts, up to an elevation of 2000 feet.

## LXV. BARRINGTONIACEÆ.

### 1. BARRINGTONIA, Forst.

† *Calyce primum indiviso vel apice minute aperto, denique in lobis 2-3 longitudinaliter fisso.*

1. **B. speciosa**, Linn. (W. et A. *Prod.* i. 333, cum syn.; Wight, *Icones*, t. 547.)—C.P. 3610.

HAB. On the sea-shore, between Galle and Matura.

2. **B. racemosa**, Roxb. (W. et A., l. c. cum syn.; Wight, *Icones*, t. 152; Thomson in *Linn. Soc. Journ.* ii. p. 47. t. 1. fig. 3.)—C.P. 2682.

Var.  $\alpha$ ; *calyce juniore indiviso, apiculato.*

Var.  $\beta$ ; *foliis floribusque minoribus, pedicellis longioribus, calyce juniore ad apicem minute aperto.*—*B. Ceylanica*, Gardn. MSS.

HAB. Warm, moister parts of the island, up to an elevation of 1500 feet. Nom. vulg. "Deya-midella-gass."

†† *Calyce 4-lobo, lobis imbricatis.* Stravadium, *Juss.*

3. **B. acutangulum**, Gært. (W. et A., l. c. cum syn.)—C.P. 1592 (1593).

HAB. Hotter parts of the island, up to no great elevation. Nom. vulg. "Ella-midella-gass."

### 2. CAREYA, Roxb.

1. **C. arborea**, Roxb. (W. et A. *Prod.* i. 334, cum syn.; Wight, *Illust.* ii. p. 20, tt. 99, 100.)—C.P. 3169.

HAB. Central Province, in exposed, grassy places, up to an elevation of 5000 feet. Nom. vulg. "Cahaata-gass."

*Ovula in loculis biserialia, singula dimidio inferiore membrana placentali vestita.*

The bark of this tree is very astringent, and is used medicinally and for tanning.

In *Barringtonia* and *Careya* I find the young leaves furnished with very minute stipules.

### 3. ANISOPHYLLEA, Br.

*Anisophyllum*, Don. *Tetracrypta*, Gardner and Champion.

1. **A. Zeylanica**, Benth. in *Niger Flora*; Hook. f. and Th. in *Linn. Soc. Journ.* ii. p. 86.—*Tetracrypta cinnamomoides*, Gard. and Ch. in *Hook. Journ. of Bot.* i. p. 314, and v. p. 378. t. 5.—C.P. 2205.

HAB. Southern and central parts of the island, up to an elevation of 1500 feet. Nom. vulg. "Wellapeyenna-gass."

*Calycis lobi persistentes. Fructus indehiscens. Embryo exalbuminosus, axialis, cotyledonibus nullis vel subnullis.*

The structure and germination of the embryo are precisely as in *Careya* and *Barringtonia*, so well described by Dr. Thomson in the Linnæan Society's Journal, vol. ii, p. 47.

## LXVI. RHIZOPHORACEÆ.

### SECT. 1. RHIZOPHOREÆ.

#### 1. RHIZOPHORA, Linn.

Nom. vulg. "Cadol-gass."

1. **R. mucronata**, Lam. (Blume, Mus. Bot. Lugd.-Bat. i. p. 132, cum syn.; Wight, Icones, t. 238.)—c.p. 1969.

HAB. Trincomalee, Calpentyu, *Gardner*. Negumbo and other parts of the coast.

This and some of the other species of the family are used for tanning.

2. **R. conjugata**, Linn. (Blume, l. c. cum syn.)—c.p. 1968.

HAB. With the preceding, and equally common.

#### 2. BRUGUIERA, Lam.

1. **B. gymnorhiza**, Lam. (Blume, l. c. p. 136, cum syn.)—An quoque *B. cylindrica*, Bl. l. c. p. 137, cum syn. ?—c.p. 3611 (1967, *partim*).

HAB. Southern coast of the island.

2. **B. Rheedii**, Blume, l. c. p. 138, cum syn.—*B. Rumphii*, Bl., and *B. Wightii*, Bl. l. c. cum syn.; Wight, Icones, t. 189 A.—c.p. 3612 (1967, *partim*).

HAB. Common near the sea-coast, *Gardner*, etc.

Blume seems to have divided this genus into an unnecessary number of species, and it may be presumed, from the differences being so trivial, that they will all prove to be varieties of one variable species.

#### 3. KANILIA, Blume, l. c. p. 140.

Walp. Ann. ii. p. 528.

1. **K. caryophylloides**, Bl. l. c., cum syn.; Walp. l. c. p. 529.—c.p. 1965.

HAB. Trincomalee and Calpentyu, *Gardner*.

#### 4. CERIOPS, Arn.

Blume, l. c. p. 142.

1. **C. Candolleana**, Arn. (Bl. l. c. cum syn.; Wight, Icones, t. 240.)—c.p. 1966.

HAB. Trincomalee, *Gardner*. Negombo, etc.

### SECT. 2. LEGNOTIDEÆ.

#### 5. CARALLIA, Roxb.

Blume, l. c. p. 128.

† *Calyce* 6-7-loba.

1. **C. integerrima**, DC. Prod. iii. p. 33; Blume, l. c. p. 129; Benth.

in Linn. Journ. Bot. iii. p. 74.—*C. Ceylanica*, Arn. Pug. 16 (334). *C. Sinensis*, Arn. l. c. 17 (335).—c.p. 1763, 1764.

HAB. Not uncommon, up to an elevation of 3000 feet. Nom. vulg. "Dawata-gass."

The wood is strong, and being rather ornamental, is adapted for furniture.

†† *Calyce 4-lobo.*

2. *C. calycina*, Benth. in Linn. Journ. Bot. iii. p. 75; *foliis* glabris, integris, ovatis, brevi-acuminatis, basi angustatis, parallelo-venosis, subtus punctulatis, petiolatis; *cymis* axillaribus, solitariis, paucifloris, folio brevioribus; *calyce* parum costato.—c.p. 3458.

Var.  $\beta$ ; *foliis* brevioribus, plus rotundatis coriaceisque.—c.p. 2588.

HAB. Var. *a*. Singhe Rajah Forest, between Galle and Ratnapoora, at no great elevation. Var.  $\beta$ . Central Province, at an elevation of 4000 to 5000 feet.

*Arbor* magna. *Folia*  $1\frac{1}{2}$ –3 poll. longa,  $1-1\frac{3}{4}$  poll. lata, *petiolo* 1–3 lin. longo. *Stipulae* interpetiolares circiter 8 lin. longæ, cito deciduæ. *Cymæ* supra medium parce ramosæ. *Bracteolæ* rotundatæ, retusæ, sæpe mucronulatæ, circiter 1 lin. longæ. *Flores* subsessiles, 3 lin. longi. *Calycis* lobi tubo subæquilongi. *Petala* albida, unguiculata, margine inciso-sinuata.

## 6. ANSTRUTHERIA, Gardner.

Calcutta Journ. of Nat. Hist. vol. vi. p. 344. t. 4. Walp. Ann. ii. p. 173.

1. *A. Zeylanica*, Gardn. l. c.—c.p. 1116.

HAB. Galle District, Gardner. Trincomalee, Dr. W. H. Harvey. Dambool, etc., up to no great elevation.

Apparently very closely allied to *Cassipourea*, Aubl.

## LXVII. LYTHRARIACEÆ.

### SECT. 1. LYTHREÆ.

#### 1. AMMANNIA, Houst.

Subgenus 1. DIPLOSTEMON, DC., Blume.

1. *A. octandra*, Linn. (Blume, Mus. Bot. Lugd.-Bat. ii. p. 132, cum syn.; Walp. Ann. iv. p. 686.)—c.p. 2797.

HAB. Hotter parts of the island.

Subgenus 2. CORNELIA, Blume.

2. *A. salicifolia*, Monti, (Bl. l. c. p. 132, cum syn.; Walp. l. c. p. 687.)—c.p. 2796.

Var.  $\beta$ ; *foliis* basi rotundatis vel cordatis.—c.p. 3296.

HAB. Common in the hotter parts of the island.

My specimens of this species have all apetalous flowers.

Subgenus 3. HAPALOCARPUM, W. et A.

3. *A. Indica*, DC. (Bl. l. c. p. 133. t. 46 A, cum syn.; Walp. l. c. p. 687.)—c.p. 1542 (1544).

HAB. Very abundant in the hotter parts of the island.



## Subgenus 4. SELLOWIA, Blume.

4. **A. pentandra**, Roxb. (Bl. l. c. p. 134. t. 46 *B*, cum syn.; Walp. l. c.; Wight, Icones, t. 260 *B*.)—c.P. 1548.  
 HAB. Very common.

## 2. AMELETIA, DC.

1. **A. Indica**, DC. (W. et A. Prod. i. p. 303; Wight, Icon. t. 257.)—*A. Indica*, Bl., and *A. elongata*, Bl. l. c. 135. t. 47, cum syn.; Walp. l. c. p. 686.—c.P. 1546.  
 HAB. Very abundant.

Judging from the descriptions only, the two forms, constituted distinct species by Blume, occur occasionally on the same plant.

## 3. ROTALA, Linn.

1. **R. verticillaris**, Linn. (Bl. l. c. p. 136, cum syn.; Wight, Icones, t. 260 *A*.)—c.P. 1547.  
 HAB. Very abundant.

## 4. PEMPHIS, Forst.

1. **P. acidula**, Forst. (Bl. l. c. p. 128. t. 43, cum syn.; Walp. l. c. p. 688.)—c.P. 1553.  
 HAB. North of the island, *Gardner*.

## 5. GRISLEA, Læffl.

1. **G. tomentosa**, Roxb. (Bl. l. c. p. 127. t. 45, cum syn.)—c.P. 1552.  
 HAB. Common in exposed places, in the Maturatte and Oova Districts, up to an elevation of 4000 feet.

## 6. LAWSONIA, Linn.

1. **L. alba**, Lam. (W. et A. Prod. i. p. 307, cum syn.; Wight, Illust. i. 207. t. 87.)—c.P. 1551.  
 HAB. Hot, drier parts of the island: Manaar, *Gardner*. Batticaloe.

## SECT. 2. LAGERSTRÆMIÆ.

## 7. LAGERSTRÆMIA, Linn.

1. **L. Reginæ**, Roxb. (Bl. l. c. p. 126. t. 41; Wight, Icones, t. 413.)—c.P. 1554.

HAB. Not uncommon in the warm, moister parts of the island, up to an elevation of 1500 feet. Nom. vulg. "Mooroota-gass."

## 8. AXINANDRA, Thw.

Hook. Journ. of Bot. vol. vi. p. 66. Walp. Ann. iv. p. 690.

1. **A. Zeylanica**, Thw. l. c. t. 1 *E*; Walp. l. c.—c.P. 2668.

HAB. Ambagamowa District, up to an elevation of 1500 feet.

## SECT. 3. SONNERATIÆ.

## 9. SONNERATIA, Linn.

1. **S. acida**, Linn. (Blume, l. c. p. 336, cum syn.; Wight, Icones, t. 340.)—c.p. 1595.

HAB. Caltura, Negombo, and other places on the coast. Nom. vulg. "Gedde-killala-gass."

*Ramuli* glandulis 2, parvis, disciformibus, ovalibus, paulo infra foliorum singulorum insertionem immersis, stipulas abortivas simulantibus muniti.

## LXVIII. ONAGRACEÆ.

## 1. JUSSIÆA, Linn.

1. **J. repens**, Linn. (W. et A. Prod. i. 335, cum syn.)—c.p. 2793.

HAB. Common in the warmer parts of the island. Nom. vulg. "Bairoo-deyanella."

Very variable in the degree of pubescence on the foliage and capsules.

2. **J. villosa**, Lam. (W. et A., l. c. 336, cum syn.)—c.p. 2794, 3297.

Var.  $\beta$ . *subglabra*.—c.p. 1540.

HAB. Very abundant, up to an elevation of 3000 feet.

## 2. LUDWIGIA, Roxb.

1. **L. parviflora**, Roxb. (W. et A., l. c. p. 336, cum syn.)—c.p. 1539, 3298, 3510.

Var.  $\beta$ ; plus minus pubescens.—c.p. 3299.

HAB. Very abundant in the warmer parts of the island, up to no great elevation.

2. **L. prostrata**, Roxb. (Fl. Ind. i. p. 420; Wight, Icon. t. 762.)—c.p. 2795.

HAB. Caltura, Moon.

## LXIX. HALORAGÆ.

## 1. MYRIOPHYLLUM, Willd.

1. **M. Indicum**, Willd. (W. et A. Prod. i. 339, cum syn.)—c.p. 1549.

HAB. Warmer parts of the island. Jaffna, etc., Gardner.

## 2. SERPICULA, Linn.

1. **S. Indica**, Thw.—*S. hirsuta*, W. et A., l. c. 338; Wight, Icones, t. 1001; Arn. Pug. 18 (336). *S. brevipes*, W. et A., l. c. (An *Haloragis oligantha*, Arn. l. c.; non W. et A.?)—c.p. 146, 447, 451, 1545, 2811.

HAB. Abundant in the Central Province, at an elevation of 5000 to 8000 feet.

*Suffrutex* procumbens, glaber vel plus minus hirsutus. *Folia* alterna, opposita vel subverticillata, rotundata, oblonga vel linearia, cuneata, inciso-dentata. *Stamina* 8, antheris linearibus. *Fructus* 8-costatus, costissæpissime 3-tuberculatis.

## 3. TRAPA, Linn.

1. **T. bispinosa**, Roxb. (W. et A., l. c. 337, cum syn.)—c.p. 2932.  
 HAB. Tanks, in the hotter parts of the island.

## LXX. CUCURBITACEÆ.

## TRIBE NHANDIROBEÆ.

## 1. ZANONIA, Linn.

1. **Z. Indica**, Linn. (W. et A. Prod. i. 340, cum syn.; Wight, *Illust.* ii. p. 31. t. 103.)—c.p. 1628.

HAB. Not uncommon, up to an elevation of 2000 feet. Nom. vulg. "Wal-rasakinda."

*Embryonis* oblongi exalbuminosi *cotyledones* planæ, carnosæ, parum inæquales, *radicula* versus hilum spectante.

I have never been so fortunate as to be able to get the female flowers of this species, so cannot give an analysis of them. The Cinghalese value this plant as a febrifuge.

## 2. PESTALOZZIA, Moritzi.

Walp. *Ann.* i. p. 316.

1. **P. laxa**, Thw.—*Zanonia laxa*, Wall. *Pl. As. Rar.* ii. p. 28; Walp. *Rep.* ii. p. 194; Arnott in *Hook. Journ. of Bot.* iii. p. 272, in adn. *Z. Wightiana*, Arn. *Pug.* 38 B (356); Walp. l. c. (An *Pest. pedata*, Zoll. and Mor.; Walp. l. c. ?)—c.p. 2582.

HAB. Central Province, at an elevation of 3000 to 5000 feet.

FL. MASC. *Antherarum* loculi longitudinaliter dehiscentes. FL. FÆM. *Ovarium* sphaericum, 3-loculare, *placentis* axilibus. *Ovula* in loculis solitaria, ab apice pendula. *Styli* 3, basi coaliti, apice divergentes, emarginati, bicuspidati. *Baccæ* sphaericæ, pisi magnitudine, 1-3-spermæ. *Semina* exalata.

## TRIBE CUCURBITEÆ.

## 3. MELOTHRIA, Linn., Endl.

*Stamina* 3, æqualia, bilocularia; *filamentis* medio tubi insertis.

1. **M. deltoidea**, Thw.—*Æchmandra deltoidea*, Arn. l. c. p. 274. *Bryonia deltoidea*, Arn. *Pug.* 19 (337).—c.p. 1610, 2581, 3128.

HAB. Common up to an elevation of 5000 feet.

*Folia* deltoidea, hastata vel digitata, parce denticulata, superne calloso-scabra, subtus glabra vel ad nervos pilosa, 1½-3 poll. longa, *petiolo* ½-1½ poll. longo. *Cirrho* simplices. *Flores* albidæ, limbo expanso, 4 lin. in diam., ad faucem hirsutissimi. MASC.: *Filamenta* brevia. *Antheræ* latæ, dorso rufo-hirsutæ, *loculis* lateralibus. *Glandula* centralis, parva, rotundata, lævis. FL. FÆM. Axillares, solitarii vel bini, sæpe cum masculis intermixti, longiuscule pedicellati. *Stamina abortiva* 3, filiformia. *Ovarium* fusiforme. *Stylus* filiformis, basi annulo glandulari cinctus. *Stigma* magnum, 3-lobatum. *Fructus* cylindricus, utrinque attenuatus vel fusiformis, teres vel plus minus acute 6-costatus. *Semina* lævia, complanata, ovalia, albida, 1½ lin. longa.

## 4. ZEHNERIA, Endl.

*Stamina 3, æqualia, bilocularia; filamentis prope basin tubi insertis.*

1. **Z. Hookeriana**, Arn. in Hook. Journ. of Bot. iii. p. 275.—*Bryonia Hookeriana*, W. et A., l. c. 345; Wight, Icones, t. 758.—c.p. 1613.

HAB. Very abundant in the Central Province, at an elevation of 2000 to 4000 feet.

2. **Z. Garcini**, Stocks in Hook. Journ. of Bot. iv. (1852) p. 149.—*Bryonia Garcini*, Willd.; W. et A., l. c. cum syn.—c.p. 1611.

HAB. Jaffna, Gardner.

3. **Z. umbellata**, Thw.—*Karivia umbellata*, Arn. l. c. *Bryonia umbellata*, Herb. Mad.; W. et A., l. c. cum syn.—c.p. 1619, 3506.

HAB. Common up to an elevation of 3000 feet.

*Flores* albidæ, ore contracti, glandula triloba, fulvo-hirsuta. MASC. *Bracteolæ* carnosæ, apiculatæ. *Connectivum* rotundatum, ad apicem fulvo-hirsutum. FÆM. *Stamina abortiva* 3, filiformia, truncata. *Ovarium* subcylindricum, versus apicem parum contractum.

## 5. ÆCHMANDRA, Arn., Walp.

*Stamina 5, unilocularia, fauce inserta.*

† *Filamentis antherisque 3-adelphis.*

1. **Æ. epigæa**, Arn. l. c. p. 274.—*Bryonia epigæa*, Rottl.; W. et A., l. c. p. 346, cum syn.; Wight, Icones, t. 503.—c.p. 1622.

HAB. Hot, drier parts of the island.

†† *Filamentis 3-adelphis, antheris distinctis.*

2. **Æ. Zeylanica**, Thw.; *foliis* oblongis vel hastato-oblongis, cordatis, acuminatis vel obtusis, subintegris vel sinuatis, parce denticulatis, superne plus minus calloso-scabris, subtus subglabris; *floribus* monoicis; *dentibus calycinis* brevibus; *corollæ* lobis rotundatis, acuminatis. MASC. Racemosis, pedunculatis. FÆM. Brevi-pedunculatis; *ovario* sphaerico, 4-6-ovulato.—c.p. 3002, 3500.

HAB. Central Province, at an elevation of 3000 to 5000 feet.

*Folia* 3-6 poll. longa, *petiolo*  $\frac{1}{2}$ -1 $\frac{1}{2}$  poll. longo. *Flores* flavi, 3-4 lin. in exp. *Antheræ* oblique patelliformes, dorso affixæ. *Filamenta* brevissima. *Baccæ* transverse oblongæ, 2-6-spermæ, coccineæ, circiter 6 lin. in max. diam. *Semina* ampuliformia, parum compressa, 3 lin. in diam., nigra, zona pallida incomplete marginata.

## 6. MUKIA, Arn., Walp.

*Stamina 5, unilocularia, prope basin tubi inserta; filamentis antherisque 3-4-adelphis.*

1. **M. scabrella**, Arn. l. c. p. 276.—*Bryonia scabrella*, Linn.; W. et A., l. c. p. 345, cum syn.—c.p. 1616.

HAB. Very abundant up to an elevation of 3000 feet.

2. **M. leiosperma**, Thw.—*Bryonia leiosperma*, W. et A., l. c. cum syn.—c.p. 2700.

HAB. Central Province, at an elevation of 4000 to 5000 feet.

Closely allied, as observed by Wight and Arnott, to the preceding, but I find in the male flowers of the present plant a rudimentary pistil, surrounded at the base, as in the female flowers, with a glandular annulus, whilst in *M. scabrella* the male flowers have merely a small rounded gland at the base of the tube.

7. **BRYONIA**, Linn.

1. **B. laciniosa**, Linn. (W. et A., l. c. 344, cum syn.; Wight, Icones, t. 500.)—c.p. 1606.

HAB. Common up to an elevation of 5000 feet.

8. **CITRULLUS**, Neck.

1. **C. Colocynthis**, Arn. l. c. p. 276; Wight, Icones, t. 498.—*Cucumis Colocynthis*, Linn.; W. et A., l. c. p. 342, cum syn.—c.p. 1607.

HAB. Jaffna and Batticaloa, *Gardner*. Nom. vulg. "Yak-komadoo."

9. **MOMORDICA**, Linn.

1. **M. dioica**, Roxb. (W. et A., l. c. p. 348, cum syn.; Wight, Icon. tt. 505, 506.)—c.p. 197 (1617).

Var.  $\beta$ . *denudata*; *floribus* masc. et fœm. parum minoribus; *bracteis* parvis, inconspicuis.—c.p. 1615. (An species distincta?)

HAB. Var.  $\alpha$ . Very abundant in the warmer parts of the island. Var.  $\beta$ . Central Province, at an elevation of 3000 to 4000 feet. Nom. vulg. "Toomba-karawilla."

Although the var.  $\beta$ , owing to the absence of the large bractæ, has a very different appearance to var.  $\alpha$ , I have searched in vain for any other character of the least importance to distinguish them, and therefore think it safer to describe them as forms of one species.

2. **M. Charantia**, Linn. (W. et A., l. c. cum syn.; Wight, Icones, t. 504.)—c.p. 1614.

HAB. Common in bushy places, near native gardens, up to an elevation of 3000 feet. Nom. vulg. "Karawilla," "Battoo-karawilla."

The several cultivated varieties of this species are much used as a vegetable by the Cinghalese.

10. **LUFFA**, Tournef.

† *Antheris distinctis*.

1. **L. pentandra**, Roxb. (W. et A., l. c. p. 343, cum syn.; Wight, Icon. t. 499.)—c.p. 2805.

HAB. Common up to an elevation of 3000 feet. Nom. vulg. "Neyang-ratta-coloo."

†† *Antheris 3-adelphis*.

2. **L. acutangula**, Roxb. (W. et A., l. c. cum syn.)—c.p. 1624.

HAB. Common in and near native gardens. Nom. vulg. "Vatta-coloo."

I have found the anthers triadelphous in all the specimens of this species I have examined.

The young fruit of this and of the preceding are used as vegetables by the Cinghalese, but the present one is most esteemed.

11. **TRICHOSANTHES**, Linn.

1. **T. cucumerina**, Linn. (W. et A., l. c. p. 350, cum syn.)—c.p. 2806.

Var.  $\beta$ . *laciniosa*.—c.p. 1620.

HAB. Very common in the warmer parts of the island. Nom. vulg. "Doommaala."

This plant is much esteemed as a febrifuge.

2. **T. nervifolia**, Linn. (W. et A., l. c. p. 349, cum syn.)—c.p. 1608.

HAB. Not uncommon up to an elevation of 5000 feet.

*Flores* albi. *Fructus* ovatus vel subrotundus, acuminatus, coccineus. *Semina* pauca, compressa, margine incrassato angulatoque, fusca, longe funiculata, epidermide coccinea vestita.

3. **T. integrifolia**, Thw.; dioica, *foliis* glabris, subcoriaceis, integris, venosis, ellipticis ovatisve, acuminatis vel rarius lanceolatis, trinerviis, basi rotundatis vel cordatis; *cirrhis* simplicibus; *floribus* majusculis, solitariis, axillaribus; *fructibus* majusculis, carnosus, sphaericis, obtuse brevi-apiculatis; *seminibus* numerosis, compactis, horizontalibus, in quoque loculo 2-serialibus.—c.p. 1629.

HAB. Central Province, at an elevation of 2000 to 4000 feet.

*Folia* 2½-6 poll. longa, 1¼-2 poll. lata, *petiolo* 3-5 lin. longo. *Flores* albid. *Fructus* coccineus, 2 poll. in diam. *Semina* lævia, oblonga, obliqua, compressa, ad hilum truncata, apice bidentata, 6 lin. longa, epidermide nigro-viridi vestita.

4. **T. palmata**, Roxb. (W. et A., l. c. cum syn.)—c.p. 1626.

HAB. Common up to an elevation of 2000 feet. Nom. vulg. "Tittahondala."

## 12. GYMNOPETALUM, Arn., Endl.

1. **G. Wightii**, Arn. l. c. p. 278; Walp. Rep. ii. p. 203.—c.p. 1625.

HAB. Central Province, up to an elevation of 5000 feet.

*Flores* dioici, albid, 6-7 lin. in exp., *tubo* piloso 5 lin. longo. *Calycis* lobi lineares. *Petala* oblonga, subacuta. MASC. *Filamenta* 3-adelpha, supra medium tubi inserta. FÆM. *Ovarium* fusiforme, pilosum. *Stylus* cylindricus. *Stigmata* 3, linearia. *Fructus* ovalis, rostratus, coccineus, 1½-2 poll. longus. *Semina* fusco-nigra, 3 lin. longa, pulpa pallide coccinea immersa.

2. **G. Zeylanicum**, Arn. l. c.—*Bryonia tubiflora*, W. et A., l. c. p. 347.

HAB. Trincomalee, Klein, Wight.

I have never met with this plant, so cannot give a decided opinion as to whether it may or not be specifically different from the preceding. The differences described hardly seem, in this family, sufficient to indicate more than forms of one species. The leaves on some of my specimens of *G. Wightii* exhibit an approach in form to those of the present one. I find the calyx always hairy in the former, and Wight and Arnott describe the fruit of the latter as slightly pilose. The seeds appear to be the same in both.

## 13. CUCUMIS, Linn.

1. **C. pubescens**, Willd. (W. et A., l. c. p. 342, cum syn.)—c.p. 3534.

HAB. Common on rubbish and on roadsides. Nom. vulg. "Gongkakiree."

2. **C. trigonus**, Roxb. (W. et A., l. c. cum syn. ? Wight, Icon. t. 497 ?)—c.p. 1618.

HAB. Aripo and Batticaloa, Gardner.

My specimens closely resemble Wight's figure, but are not sufficiently perfect to enable me to feel quite certain as to their identity.

## 14. COCCINEA, W. et A.

1. **C. Indica**, W. et A., l. c. p. 347, cum syn.—c.p. 1609.  
 HAB. Common in waste places. Nom. vulg. "Kowakka."

## LXXI. PASSIFLORACEÆ.

## 1. MODECCA, Linn.

1. **M. Wightiana**, Wahl. (W. et A. Prod. i. p. 353, cum syn.; Wight, Icon. t. 179.)—c.p. 1621.

HAB. Putlam, *Gardner*.

2. **M. palmata**, Lam. (W. et A., l. c. cum syn.; Wight, Icon. t. 201.)—c.p. 1627.

HAB. Common up to an elevation of 3000 feet. Nom. vulg. "Hondala."

The root is said to be poisonous; it is used by the Cinghalese as a medicine.

*Passiflora minima*, Jacq. (*P. Walkeria*, Wight, Illust. ii. p. 39. t. 108), is not uncommon in cultivated ground, but can scarcely be considered indigenous. The same may be said of *P. foetida*, L.

## LXXII. BEGONIACEÆ.

## 1. BEGONIA, Linn.

(Nom. vulg. "Hakambala.")

1. **B. Malabarica**, Dryand. in Linn. Soc. Trans. i. p. 171; ? Roxb. Fl. Ind. iii. p. 648; Rheede, Hort. Mal. ix. p. 167. t. 86.

Var.  $\beta$ . *dipetala*; *floribus* dipetalis vel sæpe petalis interioribus 1 vel 2 angustis ornatis.—*B. dipetala*, Grah. in Hook. Bot. Mag. t. 2849; Wight, Icon. t. 1813. *Haagea dipetala*, Klotzsch, Begon. Gatt. und Art. p. 104; Walp. Ann. iv. p. 921.—c.p. 2807.

*Flores* rosacei vel albid.

There can scarcely be a doubt, I think, of this being the plant figured by Rheede. The one described by Roxburgh would seem to have larger and differently shaped inner petals.

Sect. *Reichenheimia*.

2. **B. Thwaitesii**, Hook. Bot. Mag. t. 4692. — *Reichenheimia Thwaitesii*, Klotzsch, l. c. p. 55. t. 4 A; Walp. l. c. p. 894.—c.p. 2808.

HAB. Hunasgria and Matele Districts, in the Central Province, at an elevation of 3000 to 4000 feet.

3. **B. subpeltata**, Wight, Icon. t. 1812.—*Reichenheimia subpeltata*, Klotzsch, l. c.; Walp. l. c.—c.p. 2597.

HAB. Ambagamowa District, at an elevation of from 3000 to 4000 feet.

This is closely allied to the last species, but its leaves are more angular, its peduncles longer and less hairy, flowers larger.

Sect. *Diploclinium*.

4. **B. cordifolia**,—*Diploclinium cordifolium*, Wight, Icon. t. 1816; Klotz. l. c. p. 71; Walp. l. c. p. 904.—c.p. 3584.

HAB. On shady banks, in the Batticaloa and Bintenne Districts, at no great elevation.

Ceylon specimens differ a little from Dr. Wight's figure in having the petioles more or less pilose, and the flowers rather larger, but they closely resemble it in other respects.

Another species of *Begonia* (c.p. 3460) occurs in the south of the island, but my specimens are too imperfect to describe from. The stem is a few inches in length, closely pitted with the scars of the fallen leaves, and the whole plant is covered with a dense red pubescence.

## LXXIII. CRASSULACEÆ.

1. **KALANCHOE**, Adams.

1. **K. floribunda**, W. et A. Prod. i. p. 359?—c.p. 540.

HAB. Hantani, at an elevation of 3000 feet.

Glaberrima,  $1\frac{1}{2}$ –3-pedalis. *Folia* ovalia, crenato-serrata, longiuscule petiolata, *floralia* acuta, *radicalia* valde obtusa. *Cymæ* laxæ, paniculatæ. *Flores* lutei, rubro-tincti.

This differs from the described form of *K. floribunda* in being perfectly smooth. The branches of the cyme, too, of the present plant can hardly be called racemiform.

2. **K. laciniata**, DC. (W. et A., l. c. cum syn.; Wight, Icon. t. 1158.)

HAB. Central Province, Moon, H. de Alwis.

I have not found this species, nor are there specimens of it in this Herbarium, but Mr. Alwis, the draughtsman of this establishment, assures me he has seen it growing wild on the hills near Kandy.

2. **BRYOPHYLLUM**, Salisb.

1. **B. calycinum**, Salisb. (W. et A., l. c. cum syn.)—c.p. 541.

HAB. Not very uncommon; up to an elevation of 5000 feet. Nom. vulg. "Akka-paana," "Rata-gowa." Said to have been introduced.

## LXXIV. CACTACEÆ.

1. **RHIPSALIS**, Gærtn.

1. **R. Cassytha**, Gærtn. (DC. Prod. iii. p. 476.)—c.p. 2976.

HAB. Not uncommon; upon rocks and trees throughout the Central Province, and certainly indigenous.

*Rami* teretes. *Flores* albidi. *Calycis* segmenta 4–5, acuta, valde inæqualia, apice rubro-tincta. *Petala* 5–6, oblonga, obtusa. *Stamina* circiter 20. *Ovarium* incomplete 3–4-loculare, pluri-ovulatum. *Stylus* simplex. *Stigma* 3–4-lobatum. *Bacca* ovalis, albida, subtranslucens, pisi parvi magnitudine. *Semina* oblonga, nigra, nitida, inconspicue striata.



## LXXV. SAXIFRAGACEÆ.

## 1. VAHLIA, Thunb.

1. *V. oldenlandioides*, Roxb. (W. et A. Prod. i. p. 364, cum syn.; Wight, Icon. t. 562.)—C.P. 1686.

HAB. Hot drier parts of the island.

2. *V. viscosa*, Roxb. (W. et A., l. c. cum syn.; Wight, Icon. t. 563.)—C.P. 1687.

HAB. Jaffna, *Gardner*.

## LXXVI. UMBELLIFERÆ.

## 1. HYDROCOTYLE, Tournef.

1. *H. Asiatica*, Linn. (W. et A. Prod. i. p. 366, cum syn.; Wight, Icon. t. 565.)—? *H. lurida*, Hance in Walp. Ann. ii. 690.—C.P. 2810.

HAB. Very abundant up to the most elevated parts of the island. Nom. vulg. "Heen-gotoocola." Used by the Cinghalese as an anthelmintic.

2. *H. Javanica*, Thunb. (DC. Prod. iv. p. 67.)—*H. Nepalensis*, Hook. Ex. Fl. i. t. 30. *H. Zeylanica*, et var.  $\beta$  *Heyneana*, DC. l. c. *H. polycephala*, W. et A., l. c.; Wight, Icon. t. 1003.—C.P. 1634.

Var.  $\beta$ ; *foliis, petiolis pedunculisque* plus minus villosis, *pedicellis fructu maturo æquilongis*.—An *H. globata*, Blume, Bijd. p. 883?—C.P. 2812.

HAB. Var. *a*. Very abundant up to an elevation of 3000 feet. Var.  $\beta$ . In the more elevated parts of the island, up to 7000 feet. Nom. vulg. "Maha-gotookola."

3. *H. nitidula*, Rich. (Hook. l. c. t. 29; DC. l. c. p. 66).—*H. rotundifolia*, Roxb. Fl. Ind. ii. p. 88; Wight, Icon. t. 564. An *H. ranunculoides*, L., var. *incisa*, Blume, l. c. p. 884?—C.P. 1635.

HAB. Central Province, up to an elevation of 7000 feet.

*Folia* subtus villosula. *Fructus* brevissime pedicellati, mericarpiis glabris ad latera 1-costata.

## 2. SANICULA, Tournef.

1. *S. Europæa*, L., var. *elata*.—*S. elata*, Ham.; W. et A., l. c. p. 367, cum syn.; Wight, Icon. tt. 334 et 1004.—C.P. 2813.

HAB. Newera Ellia and Maturatte.

## 3. HELOSCIADIUM, Koch.

1. *H. Heyneanum*, DC. (W. et A., l. c. p. 368, cum syn.)—*Sium triternatum*, Moon, Cat. p. 22.—C.P. 182.

HAB. Four Corles, *Moon*. Very abundant in the Oova District, up to an elevation of 4000 feet. Nom. vulg. "Wal-asamodagan."

## 4. PIMPINELLA, Linn.

1. **P. Leschenaultii**, DC. Pr. iv. p. 122; W. et A., l. c. p. 3; Wight, Icon. t. 1005.—C.P. 515 (530).

HAB. Between Horton Plains and Galagama, *Gardner*. Bopatalawa Plains, at an elevation of 6000 feet.

*Caulis* 1-4-pedalis, pilis deflexis hirsutus. *Folia* radicalia caulinaque cordato-orbicularia vel subdeltoidea, grosse dentata, superne glabra, subtus pilosula. *Umbellæ* radii sæpissime 12-20, *involucro* nullo. *Umbellularum* radii 15-20, *involucellis* circiter 6-phyllis. *Flores* albi.

My specimens of what I believe to be this species, are somewhat intermediate between those described by De Candolle and Wight and Arnott, and the *P. Candolleana* of the latter authors, but agree most essentially with the former in the character of the fruit, and in the leaves being glabrous above. Are they all forms of one species?

## 5. BUPLEURUM, Tournef.

1. **B. falcatum**, Linn., var.—*B. ramosissimum*, W. et A., l. c. p. 370; Wight, Icones, t. 1007. *B. virgatum*, W. et A., l. c. *B. nervosum*, Moon, Cat. p. 22.—C.P. 225 (508).

HAB. Oova District, *Moon*. Newera Ellia, Maturatte, and other elevated parts of the Central Province.

## 6. PALIMBIA, Bess., DC.

1. **P. ramosissima**, DC.—*Peucedanum Zeylanicum*, Gardn. MSS.—C.P. 135.

HAB. Above Galagama, *Gardner*.

In one of the specimens collected by Dr. Gardner there are the remains of a few-leaved involucre at the base of one of the umbels.

## 7. HERACLEUM, Linn.

1. **H. Sprengelianum**, W. et A., l. c. p. 372; Wight, Icon. t. 1008.—C.P. 145.

HAB. Central Province, at an elevation of 6000 to 8000 feet.

*Folia* sparse pilosa vel subtus præcipue dense villosa, *segmentis* acutis vel rotundatis. *Mericarpium* vittæ commissurales aliquando 6, 2 exteriora multo breviora.

## LXXVII. ARALIACEÆ.

## 1. HEDERA, Linn.

† *Stylis distinctis, denique divergentibus.* (ARALIA.)

1. **H. acuminata**, Wight, Icon. t. 1062; Walp. Ann. i. p. 358.—*H. parasitica*, Don, Pr. Fl. Nep. p. 188; DC. Prod. iv. p. 265.—C.P. 4.

HAB. Banks of streams, near Adam's Peak, at an elevation of 4000 to 5000 feet.

My specimens are 5-7-foliolate, but agree so well in every other respect with Dr. Wight's figure, that there can be but little doubt of the two plants being identical.

I adopt Wight's name in preference to Don's, as from what I observed of the small trees of this species, it had no appearance of being parasitical.

†† *Stylis brevibus, coadunatis.* (EUHEDERA.)

2. **H. racemosa**, Wight, Icon. t. 1015; Walp. l. c.—c. p. 549 (559).

HAB. Central Province, at an elevation of 3000 to 5000 feet.

*Arbor* magna, partibus novellis fugaci-ferrugineo-furfuraceis. *Paniculæ* sæpe valde elongatæ ramosæque. *Pedicelli* fasciculato-racemosi. *Flores* 5-7-meri.

This agrees in so many respects with Dr. Wight's figure and description, that it is most probably identical with his plant, though he makes no mention of the ferruginous tomentum upon the young parts, which however speedily disappears.

††† *Stylis nullis.* (PARATROPIA.)

\* *Foliis digitatis.*

3. **H. Vahlîi**, Thw. — *H. terebinthacea*, Vahl, Symb. iii. p. 42? *Paratropia terebinthacea*, Arn. Pug. 20 (338); Walp. Rep. ii. p. 433.—c. p. 1632.

HAB. Very abundant up to an elevation of 3000 feet. Nom. vulg. "Itta."

*Arbor* scandens, epiphyta, balsamiflora, partibus novellis glaberrimis. *Paniculæ* juniores bracteatae; *bracteis* lanceolatis, 3-4 lin. longis, cito deciduis. *Umbellæ* 5-8-floræ. *Flores* 6-9-meri, sæpissime 8-meri. *Baccæ* flavæ.

Notwithstanding the discrepancy with Vahl's description of his plant, as having a single style of equal length with the stamens, there can be little doubt of the present being his species, as it is the common, and, I believe, the only one occurring in the lower part of the Island.

4. **H. exaltata**, Thw.; arborea; *foliolis* 7-9, majusculis, coriaceis, glabris, integris, oblongis, basi rotundatis vel subcordatis, petiolatis, venis non prominentibus; *umbellis* 10-15-floris, longiuscule pedunculatis, racemosis; *floribus* 8-9-meris.—An *Paratropia Wallichiana*, W. et A.?—c. p. 1633.

HAB. Central Province, at an elevation of 4000 to 6000 feet.

*Arbor* magna. *Partes novellæ* glabræ? *Foliola* 5-10 poll. longa, 2½-4 poll. lata; *petiolulis* ½-2½ poll. longis.

This would appear to be closely allied to *P. Wallichiana*, W. et A., if it be not identical with it, seeing how much the species vary in the number of parts of the flower.

\*\* *Foliis 1-3-foliolatis.*

5. **H. emarginata**, Moon's Cat. p. 18; scandens, fruticosa, glabra; *foliis* petiolatis; *foliolis* 1-3-nis, sæpissime solitariis, cuneatis, subsessilibus, apice truncatis, emarginatis; *paniculis* folio longioribus, parce ramosis; *umbellis* longiuscule pedunculatis, 5-8-floris; *floribus* 5-meris.—c. p. 655.

HAB. Saffragam, Moon. Central Province, up to an elevation of 4000 feet.

*Foliola* 1-2½ poll. longa, 4-12 lin. lata, utrinque (sicca) prominentim venosa, ad apicem incrassatum *petioli* 3-9 lin. longi articulata. *Flores* albidæ, rubro-tincti, circiter 1¼ lin. in diam., margine calycina obscure 5-dentata.

## LXXVIII. ALANGIACEÆ.

## 1. ALANGIUM, Linn.

1. **A. Lamarckii**, Thw.—*A. decapetalum*, Lam.; W. et A. Prod. i. p. 325, cum syn. *A. hexapetalum*, Lam.; W. et A., l. c. p. 326, cum syn. *A. tomentosum*, Lam.; DC. Prod. iii. p. 204.—c.p. 760 (1597, 1598, 1599, and 2498).

HAB. Hotter parts of the island, up to no great elevation. Nom. vulg. "Eepaatta."

The several forms of this species merge so completely into one another, that I find it quite impossible to separate them. The amount of pubescence and number of spines seem quite dependent upon the character of the climate, luxuriance of the plant, etc., specimens from the dry northern parts of the island being more strongly marked in that respect than those from the moister southern districts.

2. **A. glandulosum**, Thw.; *partibus novellis* rufo-furfuraceis; *foliis* ovatis vel oblongis, rostrato-acuminatis, basi 3-nerviis, superne glabris, subtus glandulis nitentibus minutissimis aspersis, ad axillas venarum primariarum glanduloso-porosis, petiolatis; *floribus* axillaribus, fasciculatis, externe glandulosis, parce hirsutis; *pedicellis* (petiolis subæquilingis) calycibusque rufo-glanduloso-furfuraceis.—c.p. 381.

HAB. Central Province, at an elevation of 2000 to 4000 feet.

*Arbor* parva, subscandens, plus minus spinosa. *Folia* 2½–6 poll. longa, 1–2½ poll. lata, *petiolo* circiter 3 lin. longo. *Calyx* parvus, 6–7-dentatus, basi 3–4-bracteolatus. *Petala* 5–6, ½ poll. longa. *Stamina* 15–20, interne prope basin pilosa. *Stylus* clavatus. *Stigma* discoideum, 4-radiatum. *Fructus* rufus, pulposus, sphericus, 6 lin. in diam.

Closely allied to the last, but well distinguished by the deep-red glandular furfurescence of the young shoots, and of the pedicels and calyces, by the glandular petals and under-surface of the leaves, and by the much smaller flowers and fruit.

## LXXIX. LORANTHACEÆ.

## 1. LORANTHUS, Linn.

Nom. vulg. "Pelilla."

Sect. 1. *Tibracteolati*.

1. **L. loniceroides**, Linn. (W. et A. Prod. i. p. 382, cum syn.; Wight, Icon. t. 303.)—c.p. 575, 2551 (1643).

HAB. Central Province, at an elevation of 4000 to 7000 feet.

2. **L. capitellatus**, W. et A., l. c.; Wight, Icon. t. 304.—c.p. 2814.

HAB. Central Province and Saffragam, up to an elevation of 3000 feet.

This, I suspect, is but a form of the preceding, which it very much resembles in many respects.

Sect. 2. *Involucrati*.

3. **L. Gardneri**, Thw.; *foliis* glabris, coriaceis, suboppositis, lanceolatis vel ovato-lanceolatis, basi rotundatis vel subcordatis, petiolatis; *involucris* subsessilibus, 5–6-phyllis, externe pilosulis, segmentis lanceolatis, flores 5–6 multo longiores cingentibus.—c.p. 1644.

HAB. Hantani, at an elevation of 3000 feet.

*Ramuli* teretes, glabri, rubro-nigrescentes. *Folia* 3-6 poll. longa,  $1\frac{1}{2}$ -3 poll. lata (juniora rufescentia), *petiolo* 3-5 lin. longo. *Involucra* rubra, segmentis 3-6 lin. longis. *Flores* rubri, versus apicem virides, pollicares. *Corolla* 5-fissa, fissura antica longiore, infra medium attingente.

Sect. 3. *Unibracteolati*.

† *Corolla* recta, æqualiter fissa.

4. **L. Hookerianus**, W. et A., l. c. p. 381.—c.p. 1645.

HAB. Batticaloa, *Gardner*; Hantani and Maturatte, in the Central Province.

*Spicæ* solitariæ vel binæ. *Flores* flavo-rufescentes, 4 lin. longi.

5. **L. nodiflorus**, Thw.; *foliis* glabris, coriaceis, ovato-lanceolatis, acuminatis, basi parum angustatis, petiolatis; *floribus* parvis, axillaribus, sessilibus, confertis; *bracteola* calyceque furfuraceis; *corolla* glabra, 4-fissa.—c.p. 2589.

HAB. Damp forests in the Ambagamowa District, at no great elevation.

*Ramuli* compressi, glabri, *gemma* parce furfuraceis. *Folia*  $2\frac{1}{2}$ -5 poll. longa, 1-2 $\frac{1}{2}$  poll. lata, *petiolo* 2-4 lin. longo. *Flores* rubri, 5 lin. longi.

6. **L. Neilgherrensis**, W. et A., l. c. p. 382; Wight, Icon. t. 1020.—c.p. 1647 (1648, 2483).

HAB. Central Province, at an elevation of 1000 to 7000 feet.

*Pedunculi* 2-9 lin. longi. *Corolla* rubro-sanguinea vel sæpius basi apiceque flavo-rubra, medio viridis.

May not this be a state of *L. elasticus*, Desv., from which it appears to differ but little except in its flowers being pedunculate?

†† *Corolla* plus minus curvata, inequaliter fissa.

‡ *Corollæ* segmentis 5.

7. **L. longiflorus**, Desv. (W. et A., l. c. p. 384, cum syn.; Wight, Icon. t. 302.)—c.p. 198 (1648).

Var.  $\beta$ .—*L. amplexifolius*, DC.; W. et A., l. c.—c.p. 2414, 2721.

HAB. Var.  $\alpha$ . Common in the warmer parts of the island. Var.  $\beta$ . Deltotte and Ambagamowa, up to an elevation of 3000 feet.

Var.  $\alpha$  and  $\beta$  merge so insensibly into one another, that I have not the least doubt of their being the same species. The flowers vary very much in length and in intensity of colour.

8. **L. ensifolius**, Thw.; *foliis* glabris, oppositis, membranaceis, anguste falcato-lanceolatis, 3-5-nerviis, petiolatis; *racemis* axillaribus terminalibusque, 1-2-nis, folio subæquilongis, 10-20-floris; *pedicellis* ovario longioribus; *bracteolis* minutis.—c.p. 3453.

HAB. Near Ratnapoora, at an elevation of 1000 feet.

*Ramuli* teretes, glabri. *Folia* 2-4 poll. longa,  $\frac{1}{2}$ -1 poll. lata, *petiolo* 2-4 lin. longo.

This species appears to be rare; I have not seen the flowers of it.

9. **L. suborbicularis**, Thw.; *partibus novellis* ferrugineo-furfuraceis; *foliis* coriaceis, rotundatis vel obovato-rotundatis, obtusis, 3-nerviis, basi plus minus acutis, petiolatis, (siccis) rugulosis; *racemis* axillaribus, 1-2-nis, tomentellis, folio brevioribus; *pedicellis* ovario subæquilongis; *calyce* subacutè 5-dentato; *bracteola* ovario dimidio breviorè; *corolla* 5-par-

tita, externe stellato-pilosula, segmentis linearibus; *baccis* oblongis.—C.P. 474.

HAB. Newera and other of the most elevated parts of the island.

*Ramuli* teretes, scabri, fusco-cinerei. *Folia* 1-2½ poll. longa, *petiolo* 2-4 lin. longo. *Flores* flavo-virides, apice rufescentes, pollicares.

10. **L. ligulatus**, Thw.; *foliis* glabris, carnosocoriaceis, eveniis, linearibus vel oblongis, apice rotundatis, basi acutis, petiolatis; *racemis* glabris, paucifloris, folio brevioribus vel subæquilongis; *pedicellis* ovario longioribus; *bracteola* minuta, rotundata; *calyce* obscure denticulato; *corolla* glabra, 5-fissa, segmentis spathulato-linearibus acutis.—C.P. 1640.

HAB. Jaffna and Calpentyne, *Gardner*.

*Ramuli* teretes. *Folia* ¾-2 poll. longa, 2½-5 lin. lata, *petiolo* 1-2 lin. longo. *Flores* pollicares.

11. **L. cuneatus**, Heyne (W. et A., l. c. p. 385, cum syn.)—C.P. 2554 (2552 partim).

Var.  $\beta$ ; *foliis* majoribus, plus minus stellato-puberulis, denique glabris.—*L. goodeniiflorus*, W. et A., l. c. *L. Candolleanus*, W. et A., l. c.; Wight, Icon. t. 305.

Var.  $\gamma$ ; *floribus* longioribus, cum *pedunculis pedicellisque* breve incano-tomentosis.—C.P. 1641.

HAB. Varr.  $\alpha$  and  $\beta$ . Very abundant in the warmer parts of the island, and very destructive to orange- and other fruit-trees. Var.  $\gamma$ . Jaffna, *Gardner*.

Varr.  $\alpha$  and  $\beta$  merge imperceptibly into one another, and the more pubescent state of the latter seems to differ in nothing from *L. Candolleanus*, W. et A., as figured by Dr. Wight. I have not seen var.  $\gamma$  in a fresh state.

12. **L. sclerophyllus**, Thw.; *foliis* valde coriaceis, obovato-spathulatis, obtusis, brevi-petiolatis, plerisque 3-nerviis; *junioribus* fugacissime stellato-pubescentibus; *inflorescentia* subglabra; *pedunculis* axillaribus, 1-3-nis, brevibus, 2-5-floris; *pedicellis* umbellatis, ovario subæquilongis; *bracteola* ovario dimidio brevior; *calyce* 5-dentato.—C.P. 2442 (2552, partim).

HAB. Central Province, up to an elevation of 7000 feet.

*Ramuli* teretes, scabri. *Folia* ¾-3 poll. longa, 3 lin., 1½ poll. lata, *costa marginique* sæpissime rubris. *Flores* pollicares, flavo-rufescentes, segmentis corollæ anguste lanceolatis.

Closely allied to the preceding species, from which its very coriaceous, 3-nerved leaves distinguish it. *L. memecylifolius*, W. et A., judging from Dr. Wight's figure in Icones, t. 142, would appear to resemble the present plant in many respects, but the shape and venation of the leaves are different.

• 13. **L. tomentosus**, Heyne, (W. et A., l. c. cum syn.; Wight, Icones, t. 378.)

Var. *normalis*; ferrugineo- vel griseo-tomentosus; *foliis* superne denique glabris; *bracteolis* ovario brevioribus.—*L. incanus*, Moon's Cat. p. 26.—C.P. 1642.

Var. *lanuginosus*; *floribus* brevioribus, longe et dense lanatis; *bracteolis* sessilibus et ovario parum longioribus vel majoribus, foliaceis, stipitatis; cæteroquin ut in var. præced.—C.P. 2492.

HAB. Var. *normalis*. Abundant up to an elevation of 5000 feet. Var. *lanuginosus*. In the Oova and Maturatte Districts, from 3000 to 5000 feet.

I have little doubt that the first of the above varieties is the typical form of the species, whilst in the latter variety, as well as in others described by previous authors, the bracteoles must be looked upon as in an abnormal condition, putting on, as they do, more or less of a leaf-like appearance.

‡‡ *Corollæ segmentis* 4.

14. **L. buddleioides**, Desv. (W. et A., l. c. p. 383, cum syn.)—C.P. 2391.

HAB. Drier parts of the island, up to an elevation of 4000 feet.

## 2. **VISCUM**, Tournef.

1. **V. orientale**, Willd. (W. et A., l. c. p. 379, cum syn.)—*V. verruculosum*, W. et A., l. c.—C.P. 412 (480, 481, 1639).

HAB. Central Province, up to an elevation of 7000 feet.

2. **V. capitellatum**, Smith, (W. et A., l. c. p. 380, cum syn.)—C.P. 1638.

HAB. Very common in the warmer parts of the island, often parasitic upon species of *Loranthus*.

3. **V. spathulifolium**, Thw.; *ramis* elongatis, teretibus, trichotomis, singulis ad basin involucri parvo cupuliformi cinctis; *foliis* oppositis, anguste spathulatis, obtusis, 3-nerviis; *floribus* laxè spicatis.—C.P. 336.

HAB. Adam's Peak, *Gardner*.

4. **V. attenuatum**, DC. (W. et A., l. c. cum syn.)—C.P. 479, 1637.

HAB. Jaffna, Galagama, *Gardner*.

5. **V. moniliforme**, Blume (W. et A., l. c. cum syn.; Wight, Icon. t. 1018, 1019.)—C.P. 295.

HAB. Common in the most elevated parts of the Central Province.

## LXXX. CAPRIFOLIACEÆ.

### SECT. LONICEREÆ.

#### 1. **DICHILANTHE**, Thw.

Hook. Kew Journ. of Bot. viii. pp. 270, 376; Hook. fil. et Thomson in Linn. Soc. Journ. (Bot.) ii. p. 164.

1. **D. Zeylanica**, Thw. l. c.; H.f. et T. l. c. p. 174.—C.P. 3422.

HAB. In forests between Galle and Ratnapoora, at an elevation of 1000 feet.

### SECT. SAMBUCEÆ.

#### 2. **VIBURNUM**, Linn.

1. **V. erubescens**, Wall. (H.f. et T. l. c. p. 177, cum syn.)—C.P. 1636.

HAB. Horton Plains, at an elevation of 7000 to 8000 feet.

2. **V. hebanthum**, W. et A. Prod. i. 388; Wight, Icon. t. 1023.—*V. coriaceum*, Bl. Var.  $\beta$ . H.f. et T. l. c. *V. Zeylanica*, Gardn. MS.—C.P. 39 (73).

HAB. Very abundant in the Central Province, at an elevation of 5000 to 7000 feet.

## LXXXI. RUBIACEÆ.

Sect. 1. *Corollæ segmentis æstivatione valvatis.*

† *Ovarii loculis bi-pluriovulatis.*

‡ *Placenta apicem prope loculi affixa, pendula.*

## 1. NAUCLEA, Linn.

† *Platanocarpum.*

1. **N. coadunata**, Roxb. (DC. Prod. iv. p. 344, cum syn.)—C.P. 1658.  
HAB. Common in the warmer parts of the island. Nom. vulg. "Bakmee-gass."

†† *Nauclearia.*

2. **N. Cadamba**, Roxb. (DC. l. c. cum syn.)—*N. orientalis*, L. (*partim.*)—C.P. 1659.

HAB. Common up to an elevation of 2000 feet. Nom. vulg. "Halamba-gass."

3. **N. parvifolia**, Roxb. (DC. l. c. cum syn.)—*N. orientalis*, Linn. (*partim.*)—C.P. 780.

HAB. Hot, drier parts of the island.

4. **N. tubulosa**, Arn.; *foliis* glabris (junioribus ad nervos minutissime puberulis), ovatis, obtusis vel parum acuminatis, basi rotundatis subcordatisve, petiolatis; *stipulis* magnis, membranaceis, obovatis; *pedunculis* terminalibus, sæpissime ternis, intermedio brevioribus vel sessilibus; *calyce* cylindrico, subintegro, bracteolis obtusis longioribus; *stigmatibus* longiusculis, utrinque truncatis; *capsulis* majusculis, oblongis, polyspermis.—C.P. 1657.

Var.  $\beta$ . *minor*; *foliis capitulisque* minoribus.—C.P. 1656.

HAB. Var.  $\alpha$ . Very abundant in the Kokool Corle. Var.  $\beta$ . Hot, drier parts of the island.

*Arbor* parva. *Folia* submembranacea, minutissime reticulata, 2–6 poll. longa, *petiolo* 4–6 lin. longo. *Capitula* in anthesi circiter  $1\frac{1}{2}$  poll. in diam. (var.  $\beta$  *pollicaria*), prope basin foliis stipulisque 2 deciduis bracteata. *Calyx* 1– $1\frac{1}{2}$  lin. longus. *Corolla* purpurea. *Antheræ* apiculatæ. *Capsula* 3–4 lin. longa.

††† *Pentacoryna.*

5. **N. cordifolia**, Roxb. (W. et A., l. c. cum syn.)—C.P. 1660.

HAB. Hot, drier parts of the island. Nom. vulg. "Colong-gass."

6. **N. peduncularis**, Wall. Cat. 6091; Walp. Rep. ii. p. 512.—*N. triflora*, Moon's Cat. p. 14. An *N. purpureæ*, Roxb., var. ?—C.P. 2820.

HAB. Saffragam and contiguous districts, up to an elevation of 1000 feet, not uncommon. Central Province, rare.

*Arbor* mediocris, *ramulis* parce compressis. *Pedunculi* 1–2 poll. longi, *intermedius* parum longior, paulo infra medium bracteolis 4 parvis munitus. *Capitula* in anthesi 8–9 lin. in diam. *Corolla* ochracea, purpurascens.



2. **UNCARIA**, Schreb.

1. **U. Gambier**, Hunt. (DC. Prod. iv. p. 347, cum syn.)—c.p. 1661.

HAB. Deltotte District, in the Central Province, at an elevation of 3000 feet. Near Colombo, *Mr. W. Ferguson*.

*Ramuli* subtetragoni. *Folia* juniora subtus ad nervos parce hispida.

This is a variable species in the shape of the leaves and length of the petiole. It agrees so well with Roxburgh's description that I cannot doubt its being his plant. The Cinghalese make no use, as far as I can learn, of this plant.

‡‡ *Placenta medio dissepimenti adnata.*

3. **LEUCOCODON**, Gard.

Calc. Journ. of Nat. Hist. vol. vii. p. 5.

*Florum capitula* terminalia, solitaria, involuero magno campanulato inclusa. *Calyx* tubulosus, subinteger. *Corolla* infundibuliformis, 5-loba. *Filamenta* brevia. *Antheræ* oblongæ. *Ovarium* biloculare. *Ovula* plurima, reniformia. *Stylus* simplex, exsertus. *Stigma* bilobum, lobis oblongis, obtusis. *Fructus* baccatus. *Semina* numerosa, sub-rhomboidea, compressa.

1. **L. reticulatum**, Gard. l. c.—c.p. 347 (695).

HAB. Central Province, at an elevation of 3000 to 5000 feet.

*Frutex* scandens, epiphyticus, radicans. *Folia* oblongo-lanceolata, utrinque acuta, 3-6 poll. longa, 1-2 poll. lata, *petiolo*  $\frac{1}{2}$ - $\frac{5}{4}$  poll. longo. *Stipulæ* membranaceæ, oblongæ, apice bidentatæ, basi connatæ, pollicares. *Involucrum* albidum, viridescens, 1 poll. in alt., margine dentatum, brevi-pedunculatum. *Corolla* albida, circiter 8 lin. longa, *lobis* lanceolatis acutis tubo interne minute pilosulo brevioribus. *Bacca* alba, oblonga, 4 lin. longa. *Semina* striata, brunnea, medio nigra,  $\frac{1}{2}$  lin. longa.—*A* *descript. cl. Gardner præcipue excerpt.*

4. **MUSSÆNDA**, Linn.

1. **M. frondosa**, Linn. (W. et A., l. c. p. 393, cum syn.; Wight, *Illust.* ii. t. 124.)—c.p. 1664.

HAB. Very abundant throughout the island. Nom. vulg. "Maassenda." The leaves are boiled and eaten as a vegetable by the Cinghalese.

5. **ACRANTHERA**, Arn.

Endl. *Suppl.* i. p. 1394.

1. **A. Zeylanica**, Arn. in *Ann. of Nat. Hist.* iii. p. 21.—c.p. 334.

HAB. On damp, shaded banks: in the Central Province, at an elevation of 3000 to 5000 feet.

6. **NEUROCALYX**, Hook.

1. **N. Zeylanicus**, Hook. *Icon. Pl.* ii. t. 174.—c.p. 286 (362).

HAB. Saffragam District, up to an elevation of 4000 feet.

The flowers of this and of all the following species are of a pale-purple colour.

2. **N. Wightii**, Arn. l. c. p. 20; Walp. *Rep.* ii. p. 515. An *N. Hookeriani*, Wight, *Icon.* t. 52, var. ?—c.p. 595.

HAB. Warm, moister parts of the island, up to an elevation of 2000 feet.

3. **N. Gardneri**, Thw. ; *foliis* majusculis, superne pilosulis, subtus ad nervos villosis, ovato-lanceolatis, acutis, obscure denticulatis, basi cuneatis ; *stipulis* bipartitis ; *segmentis* rotundatis, obliquis, longe acuminatis ; *racemis* villosis ; *bracteolis* oblongis, acuminatis, prope basin utrinque 1-dentatis ; *segmentis calycinis* rotundatis, subcordatis, acuminatis.—C.P. 1671.

HAB. Pasdoon Corle, *Gardner*.

Its wider, more hairy leaves, and villous racemes, distinguish this from the preceding, to which it is in other respects very closely allied, and the two may possibly prove to be forms of one species.

4. **N. Championii**, Benth. MSS. ; *foliis* oblongo-spathulatis, obtusis subacutisve, superne villosulis, denique subglabris, subtus ad nervos cum petiolo villosis ; *stipulis* bipartitis ; *segmentis* acutis, irregulariter 2-3-fissis ; *racemis* villosis ; *bracteolis* 3-partitis, segmentis lateralibus minoribus.—C.P. 3417, 3480.

HAB. Kokool Corle, up to no great elevation.

*Folia* 3-6 poll. longa, 1-2 poll. lata, *petiolo*  $\frac{1}{4}$ - $1\frac{1}{2}$  poll. longo.

## 7. SCHIZOSTIGMA, Arn.

Endl. Suppl. i. p. 1393.

1. **S. hirsutum**, Arn. Ann. of Nat. Hist. iii. p. 20.—C.P. 268.

HAB. Not uncommon in the south of the island, up to an elevation of 2000 feet.

*Flores* 1-3-ni. *Corolla* alba, 7-8 lin. in exp. *Bacca* cerasi parvi magnitudine.

## 8. AXANTHES, Blume.

1. **A. Zeylanica**, Wight, Icon. t. 1163.—C.P. 499 (707).

HAB. Central Province, at an elevation of 4000 to 7000 feet.

2. **A. elliptica**, Wight, Icon. t. 1164.—C.P. 359 (584).

HAB. In similar situations with the preceding.

The flowers and fruit of this and of the preceding species are of a dull orange-yellow colour.

††† *Placenta versus basin loculi affixa, plus minus erecta.*

## 9. OPHIORRHIZA, Linn.

(Nom. vulg. "Datkatteya.")

1. **O. Mungos**, Linn. (W. et A. Prod. i. 404, cum syn.)—C.P. 1704.

HAB. Very abundant in damp, shady places.

2. **O. nemorosa**, Thw. ; *caule* suberecto, parce ramoso ; *foliis* glabris, lanceolatis, acuminatis, basi cuneatis ; *stipulis* rotundatis, longe 1-2-setaceo-cuspidatis ; *corymbis* terminalibus, pedunculatis, tomentosis ; *bracteolis* minutis ; *corolla* longiuscula.—C.P. 205 (215, 484).

HAB. Central Province, at an elevation of 3000 to 5000 feet.

*Folia* 2-6 poll. longa,  $\frac{3}{4}$ - $1\frac{1}{2}$  poll. lata. *Corolla* alba, rubro-tincta, tubo semi-pollicari.

3. **O. radicans**, Gardn. MSS. ; *caule* procumbente ; *petiolis* pedun-

*culisque* tomentosis; *foliis* parvis, ovatis orbicularibusve, acutis vel obtusis, basi rotundatis cordatisve, superne glabris, subtus ad nervos tomentosis; *corymbis* terminalibus, parvis; *pedunculo* foliis longiore; *bracteolis* parvis.—C.P. 1706.

HAB. Not uncommon in the forests of the Central Province.

*Folia*  $\frac{1}{2}$ –1 poll. longa. *Corolla* alba, *tubo* 2 lin. longo, *limbo* 3 lin. in exp.

4. *O. angustifolia*, Thw.; *caule* erecto, glabro vel breve tomentoso; *foliis* anguste lanceolatis, utrinque angustatis, glabris vel subtus ad nervos tomentosis; *stipulis* setaceo-cuspidatis; *corymbis* laxis pedunculoque tomentosis; *bracteolis* linearibus, mediocribus; *alabastris* rugulosis.—C.P. 431.

HAB. Ambagamowa District.

*Folia* 1–4 poll. longa,  $1\frac{1}{2}$ –8 lin. lata, subtus albida. *Corolla* alba, *tubo* 1 lin. longo, *limbo* 3–4 lin. in exp.

Resembling the next species in many respects, but the smaller, narrower leaves, and the lax corymbs with their shorter bracteoles, of the present plant, seem sufficient to keep them distinct.

5. *O. pectinata*, Arn. Pug. 20 (338); Walp. Rep. vi. p. 57.—C.P. 712 (1707).

Var.  $\beta$ ; *corymbis* arcte compactis.—*O. imbricata*, Gard. MSS.—C.P. 750.

HAB. Abundant in the Central Province, up to an elevation of 5000 feet.

6. *O. pallida*, Thw.; *caule* tomentoso, plus minus procumbente; *foliis* hispidulis, lanceolatis vel ovato-lanceolatis, acuminatis, basi acutis vel rotundatis; *stipulis* setaceo-cuspidatis; *corymbis* compactis pedunculoque glabris vel rarius sparse pilosulis; *bracteolis* longiusculis, linearibus.—C.P. 711, 3344.

HAB. Forests in the Ambagamowa District.

Closely allied to the last, but distinguished by its more or less procumbent habit, its pubescent stems and leaves, and the latter, together with the corymbs, smaller and more delicate in texture.

7. *O. glechomæfolia*, Thw.; *caule* procumbente, villosulo; *foliis* parvis, rotundatis ovatisve, cordatis, subacutis, superne villosulis, subtus ad nervos petioloque villosis; *corymbis* terminalibus, pedunculatis; *bracteolis* linearibus, longiusculis, cum *lobis calycinis* ciliatis.—C.P. 1708.

HAB. Forests of the Central Province.

*Folia* 3–10 lin. longa, sæpe rugulosa.

This bears a close resemblance to *O. radicans*, differing from it however in its leaves being hairy on the upper surface, and in its relatively much longer-ciliated bracteoles and calyx-lobes.

## 10. HEDYOTIS, Lam.

1. *H. Lawsoniæ*, W. et A., l. c. p. 407, cum syn.; Wight, Icon. t. 1026.—C.P. 13, 1697 (102, 110).

HAB. Central Province, at an elevation of 5000 to 8000 feet.

*Folia* ovata vel plus minus anguste lanceolata,  $\frac{1}{2}$ –4 poll. longa, membranacea vel subcoriacea.

A very common species, varying very much in appearance according to the character of the locality in which it occurs.

2. *H. evenia*, Thw.; fruticosa; *ramulis* acute 4-gonis, *junioribus*

pilosulis; *foliis* parvis, glabris, nitidis, coriaceis, eveniis, lanceolatis, acutis, basi rotundatis, petiolatis; *stipulis* triangularibus, pilosulis, apice parce glanduloso-tuberculatis; *paniculis* terminalibus, pilosulis.

—C.P. 93.

HAB. Adam's Peak, *Gardner*.

Although this plant has a very dissimilar aspect to that of the preceding, yet I suspect it may prove to be but a variety of it.

3. **H. obscura**, Thw.; suffruticosa, erecta, (sicca) nigrescens; *ramulis* subteretibus; *foliis* glabris, lanceolatis, utrinque acutis, petiolatis; *venis primariis* parum conspicuis; *stipulis* triangularibus, plus minus connatis, pectinatim pinnatifidis, segmentis linearibus pilosis; *corymbis* axillaribus, pedunculatis, glabris, folio brevioribus, bracteolatis; *corolla* ad faucem pilosa.—C.P. 94 (111, 119).

Var.  $\beta$ ; *foliis* ovatis vel ovato-lanceolatis; *venis primariis* subtus prominentibus et sæpe pilosulis; *bracteolis* parvis; *corollæ lobis* interne, et externe ad apicem pilosis.—C.P. 91, 101.

HAB. Central Province, at an elevation of 6000 to 8000 feet. Var.  $\beta$ , in rather shady places.

Apparently closely allied to *H. stylosa*, Br., but distinguished from it, judging from Dr. Wight's figure of that species in *Icones*, t. 1027, by its inflorescence not being terminal.

4. **H. Lessertiana**, Arn. Pug. 22 (339); Walp. Rep. ii. p. 492.—C.P. 99, 115 (123, 1702).

Var.  $\beta$ . *major*; *foliis floribusque* majoribus.—C.P. 96.

Var.  $\gamma$ . *confertiflora*; *foliis* coriaceis; *floribus* majoribus confertisque.—C.P. 103 (109, 1701).

Var.  $\delta$ . *pilosa*; *ramulis foliisque* plus minus pilosis; *dentibus calycinis* brevibus, cæteroquin ut in var. *a*.—C.P. 2819.

HAB. Central Province. Varr. *a* and  $\delta$ . Abundant at an elevation of 3000 to 5000 feet. Varr.  $\beta$  and  $\gamma$ . From 5000 to 8000 feet.

The extreme forms of this plant look very different, but they merge so insensibly into one another that I find it impossible to separate them as distinct species.

5. **H. flavescens**, Thw.; *ramulis* teretibus; *foliis* glabris, coriaceis; subeveniis, ovato-vel oblongo-lanceolatis, acutis; *costa* prope basin *petioloque* dilatatis; *stipulis* scariosis, vaginantibus, persistentibus, prope marginem appendiculis linearibus, apice filamentosis denique caducis cinctis; *paniculis* terminalibus, compactis.—C.P. 86, 126.

HAB. Adam's Peak, and between Galagama and Horton Plains, at an elevation of 5000 to 7000 feet.

*Folia* 1-4 poll. longa,  $\frac{1}{2}$ -1 $\frac{1}{2}$  poll. lata, flavescentia, *petiolo* 1-6 lin. longo. *Corolla* ad faucem lanosa. *Fructus* oblongus, *dentibus calycinis* erectis, acutis, parce pilosis.

6. **H. quinquenervia**, Thw.; fruticosa, erecta; *ramulis* compressis; *foliis* parvis, coriaceis, subglabris, ovatis, obtuse brevi-acuminatis, 5-nerviis, subsessilibus, marginibus revolutis; *stipulis* connatis, cupuliformibus, externe pilosis plus minus setosisque, appendiculis linearibus, scariosis, pilosis, apice dilatatis fimbriatis, persistentibus; *capitulis* terminalibus, sessilibus.—C.P. 120.

HAB. Central Province, at an elevation of 7000 to 8000 feet.

*Folia* 3-8 lin. longa, flavescentia. *Calyx* 4-fidus, segmentis subacutis apice pilosis. *Corolla* alba, externe purpurascens, interne ad faucem pilosa. *Antheræ* oblongæ, purpureæ.

7. **H. verticillaris**, Wall. (W. et A., l. c. p. 409; Wight, Icon. t. 1029).  
—*H. plantaginifolia*, Arn. Pug. 22 (340).—c.p. 107.

HAB. Central Province, at an elevation of 7000 to 8000 feet.

8. **H. Gardneri**, Thw.; *foliis* confertis, glabris, nitidis, valde coriaceis, subeveniis, lanceolatis, acutis, basi ad petiolum dilatatum angustatis; *stipulis* oblongis, membranaceis, glabris, scariosis, persistentibus; *paniculis* terminalibus, elongatis, plurifloris, laxis, pluries trichotomis.—c.p. 116.

HAB. Adam's Peak, *Gardner*.

*Rami* circiter  $\frac{1}{2}$  poll. in diam., *stipulis* confertis squamosi. *Folia* 3–4 poll. longa. *Calyx* 4-dentatus, dentibus apice incrassatis parum reflexis.

9. **H. fruticosa**, Linn. (W. et A., l. c. p. 408, cum syn.)—c.p. 108, 2457.

HAB. Very abundant up to an elevation of 3000 feet. Nom. vulg. "Waranneya."

10. **H. cymosa**, Thw.; fruticosa, erecta, (sicca) nigrescens; *ramulis* lævibus, parum angulatis; *foliis* glabris, nitidis, anguste lanceolatis, utrinque acutis; *venis primariis* inconspicuis; *stipulis* triangularibus, longe acuminatis, carinatis, margine glanduloso-dentatis; *cymis* terminalibus, laxis, puberulis; *calyce* glabro, 4-partito; *segmentis* lanceolatis, acutis, demum recurvis; *corolla* externe glabra, fauce pilosa.—c.p. 3421.

HAB. Hinidoon Corle, at an elevation of 1000 feet.

*Folia* 2–3 poll. longa, 3–7 lin. lata.

Readily distinguished from *H. fruticosa*, to which it is very closely allied, by its regularly cymose lax inflorescence, and by the whole plant becoming black in drying.

11. **H. monosperma**, W. et A., l. c. p. 410, cum syn.; Wight, Icon. t. 1031.—c.p. 97 (112, 139).

Var.  $\beta$ . *subglabra*.—*H. Richardiana*, Arn. l. c. p. 22 (340); Walp. Rep. ii. p. 492.

HAB. Very abundant in the more elevated parts of the Central Province.

The whole plant is very fetid. It varies very much in the size and hairiness of the leaves.

12. **H. nummularia**, Arn. Pug. 23 (341); Walp. Rep. ii. p. 492.—c.p. 1692.

Var.  $\beta$ . *glabra*.—*H. nummulariformis*, Arn. l. c.; Walp. l. c. p. 493.—c.p. 113, 2817.

HAB. Abundant in wet, rocky places: in the Central Province, at an elevation of 5000 to 8000 feet.

*Caules* tetragoni. *Folia* 3–10 lin. longa, marginibus sæpe revolutis. *Flores* purpurei.

13. **H. auricularia**, Linn. (DC. Prod. iv. p. 420; W. et A., l. c. p. 412, cum syn.)—c.p. 1694.

HAB. Very abundant in the warmer parts of the island. Nom. vulg. "Gatta-cola."

The leaves are boiled, after being cut very small, and eaten by the Cinghalese with their rice.

14. **H. macrophylla**, Wall. (W. et A., l. c. p. 408, cum syn.)—c.p. 104, 466.

HAB. Forests about Adam's Peak.

*Stipulæ* membranaceæ, connatæ, setis dense pilosis limbo longioribus marginatæ. *Dentes calycini* lineari-lanceolati.

15. **H. nodulosa**, Arn. l. c. 22 (340); Walp. l. c. p. 493.—An *H. Walkeri*, Arn. l. c.; Walp. l. c.?

Var.  $\alpha$ ; *foliis* superne glabris vel obscure setulosis.—c.p. 87 (105, 1698).

Var.  $\beta$ ; plus minus pilosa.—*H. axillaris*, Gardn. MSS.—c.p. 88, 3095, 3542 (106, 117, 121, 1699).

HAB. Central Province and Saffragam, at an elevation of 3000 to 7000 feet.

*Folia* 2–6 poll. longa,  $\frac{1}{2}$ – $2\frac{1}{2}$  poll. lata, *petiolo* 2–10 lin. longo. *Stipulæ* membranaceæ, plus minus connatæ, dentibus subulatis brevioribus vel subæquilongis marginatæ. *Dentes calycini* subulati, recurvi.

A very variable species as regards the size and the amount of pubescence of the leaves, which are more or less tomentose or villous, or sometimes nearly altogether glabrous. From *H. macrophylla* it differs principally in the smaller, narrower teeth of the calyx.

16. **H. membranacea**, Thw.; suffruticosa, erecta, glabra; *foliis* oblongo-lanceolatis, acuminatis, basi acutis, petiolatis; *stipulis* appendiculatis linearibus, longiusculis, scariosis fimbriatis; *floribus* dense glomerato-verticillatis; *segmentis calycinis* lanceolatis, scariosis.—c.p. 3621.

HAB. Pasdoon Corle, at no great elevation.—A specimen in the herbarium, collected by Moon, has no locality affixed to it.

*Folia* 4–6 poll. longa, 1– $1\frac{1}{2}$  poll. lata. *Corolla* parva, flava.

An obscure plant, and possibly, as suspected by Dr. Gardner, an abnormal state of one of the two preceding species.

17. **H. cyanescens**, Thw.; suffruticosa, basi decumbens, radicans; *ramis* acute tetragonis, glabris vel pilosulis; *foliis* plus minus villosulis, ovato-lanceolatis, utrinque acutis, petiolatis; *stipulis* brevibus, membranaceis, connatis, dentibus subulatis æquilongis marginatis, parce pilosis; *floribus* dense verticillatis; *segmentis calycinis* lineari-lanceolatis, ovario 3–4-plo longioribus, villosulis.—c.p. 2384.

HAB. Muruta, Gardner. Ambagamowa District, not common.

*Folia*  $\frac{1}{2}$ – $2\frac{1}{2}$  poll. longa, *petiolo* 2–5 lin. longo. *Calycis* segmenta 2 lin. longa. *Corolla* albida, purpureo-tincta, externe glabra, tubo infundibuliformi fauce parum piloso.

The whole plant is often of a bluish-purple colour.

18. **H. inamœna**, Thw.; suffruticosa, erecta, subglabra; *ramis* subtetragonis; *foliis* lanceolatis, acuminatis, basi acutis, petiolatis, (siccis) fuscescentibus; *stipulis* triangularibus, pilosis, apice parum glandulosis, basi connatis, cito deciduis; *floribus* glomerato-verticillatis; *segmentis calycinis* lanceolatis, tubo corollæ brevioribus.—c.p. 95, 3543 (98).

Var.  $\beta$ ; *foliis* latioribus, (siccis griseo-viridibus); *stipulis* longiuscule rostrato-acuminatis.—c.p. 2639.

HAB. Ambagamowa and Saffragam Districts.

*Folia* 3–5 poll. longa,  $\frac{3}{4}$ –2 poll. lata, *petiolo* 2–5 lin. longo.

19. **H. nitida**, W. et A., l. c. p. 412.—*H. Neesiana*, Arn. Pug. p. 23 (341).—c.p. 122.

HAB. Common in the warmer parts of the island. Nom. vulg. "Pittasooddoopala."

The leaves of this plant are finely chopped up, boiled, and eaten with rice, by the Cinghalese.

20. **H. cærulea**, W. et A., l. c.—c.P. 1674.  
HAB. Sandy places, especially near the coast.

Div. *Oldenlandia*.

21. **H. umbellata**, Lam. (W. et A., l. c. p. 413, cum syn.)—*H. polygonoides*, Wall.—c.P. 125.

HAB. Abundant, especially near the sea. Nom. vulg. "Saayam-mool."  
Used as a dye by the Cinghalese.

22. **H. Burmanniana**, Br. (W. et A., l. c. p. 415, cum syn.)—*H. intermedia*, W. et A., l. c.—c.P. 1676, 2368.

HAB. Very abundant in the warmer parts of the island. Nom. vulg. "Wal-patpaadagam."

23. **H. racemosa**, Lam. (W. et A., l. c. p. 414, cum syn.; Wight, Icon. t. 312.)—*H. biflora*, Br.; W. et A., l. c. cum syn.—c.P. 1679.

HAB. Hotter parts of the island, common.

24. **H. quadrilocularis**, Thw.; herbacea, procumbens, ramosa, pilosula; ramis subteretibus; foliis ovato-lanceolatis, utrinque subacutis, petiolatis; stipulis membranaceis, rotundatis, ciliatis; floribus axillaribus terminalibusque, 1-2-nis, sessilibus; ovario 4-loculari, rarius 2-3-loculari.—c.P. 3132.

HAB. Badulla District, rare.

Folia  $\frac{1}{2}$ – $1\frac{1}{2}$  poll. longa,  $\frac{1}{4}$ – $\frac{3}{4}$  poll. lata, petiolo 1–5 lin. longo. Calyx turbinatus; lobis oblongis, ciliatis, interne setulosis, recurvis. Corolla tubulosa, interne glabra; lobis oblongis, acutis. Stylus filiformis, exsertus. Stigmata tot quot ovarii loculi, longiuscula, subclavata, reflexa. Capsula depressa, loculis ad basin dehiscentibus, circiter 3-spermis.

25. **H. trinervia**, (Roem. et Sch.); W. et A., l. c. cum syn.—c.P. 1681.  
HAB. Hotter parts of the island.

26. **H. Heynei**, Br. (W. et A., l. c. p. 416, cum syn.)—c.P. 1673.  
HAB. Very common in the warmer parts of the island.

27. **H. maritima**, Wall. (W. et A., l. c. p. 417, cum syn.)—c.P. 1678.  
HAB. Very common near the sea.

11. **DENTELLA**, Forst.

1. **D. repens**, Forst. (W. et A., l. c. p. 405, cum syn.)—c.P. 1683.  
HAB. Warmer parts of the island, not uncommon.

†† *Ovarii loculis 1-ovulatis.*

‡ *Ovulis erectis.*

12. **MORINDA**, Vaill.\*

1. **M. bracteata**, Roxb. (W. et A., l. c. p. 419, cum syn.; Wight, Illust. t. 126.)—An *M. citrifoliæ*, Linn., var. ?—c.P. 765.

\* The ovary in this genus appears to be normally two-celled, but these two cells are converted into four by the introverted carpellary leaves being so produced laterally as to reach the walls of the ovary. If this is the correct explanation of the ovarial structure, the genus would have to be placed in the previous division of the family, each cell in that case being really two-ovuled.

HAB. Not uncommon in the hotter parts of the island, in and near the gardens of the Cinghalese, who employ the roots as a dye. Nom. vulg. "Ahoogass."

Some fresh specimens of this plant, now before me, agree perfectly with Wight's figure of *M. bracteata*, whilst they so closely resemble too the figure in 'Hortus Malabaricus,' vol. i. t. 52, quoted for *M. citrifolia*, L., that I can scarcely believe our plant to be other than a form of the latter. Some of the capitula are without the bract-like appendage of the calyces, and the anthers are quite exerted in the flowers now under examination.

2. *M. exserta*, Roxb. (W. et A., l. c. cum syn. var.  $\beta$ .)—C.P. 1668.

HAB. Jaffna, Gardner. Badulla District, J. Bailey, Esq. Nom. vulg. "Ahoogass."

3. *M. umbellata*, Linn. (W. et A., l. c. p. 420, cum syn.)—*M. scandens*, Roxb. Fl. Ind. i. p. 548.—C.P. 1669.

HAB. Very common up to an elevation of 4000 feet. Nom. vulg. "Kirri-walla."

The tough stems of this plant are used, instead of ropes, by the Cinghalese for tying fences.

### 13. MEPHITIDIA, Reinw.

*Lasianthus*, Jack, Wight.

1. *M. Moonii*, Wight, in Calc. Journ. of Nat. Hist. vi. p. 504; Walp. Ann. ii. p. 760.—C.P. 92.

HAB. Abundant in the Saffragam District.

*Frutex* hispidus, parce ramosus. *Folia* superne glabra hispidulave. *Capitula* pedunculata vel sessilia. *Corolla* alba, 3 lin. in exp.; *tubo* brevi. *Bacca* hispidula, pulposa, rubra, subsphærica, cerasi magnitudine.

2. *M. Gardneri*, Thw.; fruticosa; *ramulis* compressis petiolisque strigosis vel hispidulis; *foliis* oblongo-lanceolatis, acuminatis, basi acutis, superne glabris, subtus pilosulis, venosis, venis secundariis numerosis parallelis; *stipulis* parvis, triangularibus, acutis, strigosis; *capitulis* sessilibus, pilosis; *segmentis calycinis* lanceolatis, acutis; *corolla* externe pilosula, interne villosa; *tubo* brevi.—C.P. 348 (1723).

HAB. Newera Ellia and Adam's Peak, Gardner.

*Folia* 3-8 poll. longa,  $1\frac{1}{2}$ - $2\frac{1}{2}$  poll. lata, *petiolo* 2-5 lin. longo.

3. *M. tomentosa*, Thw.; fruticosa; *ramulis* teretibus, vix compressis petiolisque tomentosis; *foliis* coriaceis, lanceolatis, longe acuminatis, basi acutis, superne glabris, subtus præcipue ad nervos tomentosis, venis secundariis paucis subparallelis; *capitulis* pilosis, paucifloris, sessilibus; *segmentis calycinis* lanceolatis, acutis; *corolla* externe pilosa; *tubo* brevi.—C.P. 2711.

HAB. Ambagamowa District, at an elevation of 3000 feet.

*Folia* 2-4 poll. longa,  $\frac{1}{2}$ -1 poll. lata, *petiolo* 3 lin. longo. *Corolla* albida. *Baccæ* pisi magnitudine, nigræ.

Closely allied to, and possibly a form of, the preceding species.

4. *M. rhizophylla*, Thw.; fruticosa; *ramulis* tenuibus, compressis, scabridis; *foliis* membranaceis, glabris, nitidis, delicatule venosis, lanceolatis, longe acuminatis, basi acutis, petiolatis; *capitulis* paucifloris, sessilibus; *calyce* piloso; *segmentis* lineari-lanceolatis, longiusculis.—C.P. 361.

HAB. Adam's Peak, Gardner.



*Folia* 2-3 poll. longa,  $\frac{1}{2}$ -1 poll. lata, *petiolo* 1-2 lin. longo.

The specimens I have are not sufficiently perfect to enable me to complete the description of this species.

5. **M. protracta**, Thw.; fruticosa; *ramulis* compressis petiolisque strigosulis; *foliis* elongato-lanceolatis, a prope basin gradatim angustatis, acuminatis, mucronatis, superne glabris, subtus ad nervos strigosulis, basi subacutis, brevi-petiolatis; *venis primariis* 7-8, folii versus apicem porrectis; *secundariis* numerosis, transverse parallelis; *stipulis* minutis, triangularibus, strigosis; *capitulis* paucifloris, sessilibus; *calyce* minute 4-dentato.—C.P. 354.

HAB. Central Province, Gardner.

*Folia* 4-8 poll. longa,  $1\frac{1}{4}$ - $2\frac{1}{4}$  poll. lata, *petiolo* 1-2 lin. longo.

I have not seen this species in a fresh state, and my specimens are too imperfect to enable me to describe the flowers and fruit.

6. **M. strigosa**, Wight, l. c. p. 512; Walp. l. c. p. 762.—C.P. 1725.

HAB. Common in the Central Province and the Saffragam District, at an elevation of 2000 to 5000 feet.

*Folia*  $2\frac{1}{2}$ -5 poll. longa, 1-2 poll. lata, *petiolo* 2-4 lin. longo, *venis secundariis* transverse parallelis. *Corolla* albida, externe pilosula; *tubo* 3 lin. longo; *limbo* circiter 4 lin. in exp., interne villosa. *Baccæ* subsphæricæ, opacæ, nigræ, circiter  $\frac{1}{2}$  poll. in diam.

7. **M. Walkeriana**, Wight, l. c. p. 513; Walp. l. c.—C.P. 289 (340, 1728).

HAB. Central Province, up to an elevation of 5000 feet.

*Folia* glabra vel subtus ad nervos parce strigosa, oblongo-vel lineari-lanceolata, basi acuta, 2-5 poll. longa, 4-14 lin. lata, *petiolo*  $1\frac{1}{2}$ -3 lin. longo. *Corolla* externe pilosula, *tubo* brevi, *limbo* interne villosa. *Baccæ* nigræ, pisi magnitudine.

The narrower, more acuminated, less pilose leaves and shorter flowers, principally distinguish this from the last species. In *M. strigosa* too the calyx-tube is much longer than in the present plant. I suspect, however, that they will prove to be only forms of one species; and *M. protracta*, Thw., will probably have to be united with them as a more rare variety.

8. **M. oligantha**, Thw.; fruticosa; *ramulis* compressis, scabridis; *foliis* oblongis, rostrato-acuminatis, basi rotundatis, utrinque venosis, glabris vel subtus ad nervos pilosulis, brevi-petiolatis; *stipulis* parvis, oblongis, pilosis; *capitulis* paucifloris, sessilibus; *floribus* parvis; *dentibus calycinis* brevibus, acutis.—C.P. 357 (1726, 1727).

Var.  $\beta$ . *obliqua*; *foliis* subsessilibus, basi cordatis, obliquis.—C.P. 3418.—An species distincta?

HAB. Var.  $\alpha$ . Hantani District. Var.  $\beta$ . Saffragam District, up to an elevation of 3000 feet.

Var.  $\alpha$  closely approaches some states of *M. strigosa*, Wight, and is not probably a form of it growing in poor or dry soil. The whole genus *Mephitidia* requires much investigation before the extent of variation to which the species are liable can be satisfactorily determined.

9. **M. varians**, Thw.; fruticosa, glabra; *ramulis* striatis, parum compressis; *foliis* coriaceis, ovatis vel oblongis, acutis retusisve, mucronatis, basi angustatis, petiolatis, utrinque reticulatim venosis; *stipulis* parvis, subacutis, apice pilosis; *capitulis* pedunculatis subsessilibusve, paucifloris; *dentibus calycinis* brevibus, triangularibus, acutis; *corolla* albida; *tubo* brevi; *limbo* interne villosa.—C.P. 57 (80, 290, 422, 423).

HAB. Central Province, at an elevation of 5000 to 8000 feet.

*Folia* 1- $2\frac{1}{2}$  poll. longa, 4 lin. ad  $1\frac{1}{2}$  poll. lata, *petiolo* 2-4 lin. longo. *Corolla* albida. *Baccæ* nigræ, pisi magnitudine.

14. **ALLÆOPHANIA**, Thw. (gen. nov.).

*Flores* polygamo-dioici. *Calyx* tubulosus, 4-dentatus. *Corolla* infundibuliformis, 4-loba. *Stamina* 4, fauce inserta, *filamentis* (in flor. ster.) valde exsertis, *antheris* oblongis dorso affixis. *Ovarium* 4-loculare. *Stylus* filiformis (in flor. fert.), valde exsertus. *Stigma* 4-lobum, lobis pilosis. *Bacca* sicca, parva, 4-sperma. — Frutex, Mephitidiæ valde affinis, sed facie Hedyotidis; ramulis subtetragonis; foliis membranaceis; stipulis connatis, tubulosis, scariosis, striatis, ad marginem subaristatis; capitulis multifloris, axillaribus, sessilibus; floribus parvis, albis, purpureo-tinctis.

1. **A. decipiens**, Thw.; foliis lanceolatis, acuminatis, basi acutis, pilosis vel glabris, subtus ad nervos adpresse pilosis.—c.p. 3093.

Var.  $\beta$ . *flavescens*, Thw.; foliis plus minus pilosis, subtus ad nervos villosis, ovatis, acutis, basi cordatis rotundatis vel subacutis, petiolatis sessilibusve.—c.p. 3094 (114).

HAB. Central Province, at an elevation of 3000 to 5000 feet. Var.  $\beta$ , up to an elevation of 7000 feet.

*Folia*  $1\frac{1}{2}$ –5 poll. longa,  $\frac{1}{4}$ – $1\frac{1}{2}$  poll. lata, *petiolo* 2–8 lin. longo.

In general appearance this species bears a very great resemblance to *Hedyotis nodulosa*, Arn.

15. **GRUMILEA**, Gærtn., W. et A.

1. **G. elongata**, Wight, Icon. t. 1036.—c.p. 1719.

HAB. Central Province, at an elevation of 4000 to 7000 feet.

2. **G. nudiflora**.—*Psychotria? nudiflora*, W. et A., l. c. p. 434. An *G. nigra*, Gærtn. De Fruct. i. p. 138. t. 28. f. 2?—c.p. 18 (2707 partim).

HAB. Very abundant in the Central Province, at an elevation of 3000 to 6000 feet.

*Corymbi* ad nodos ferrugineo-pilosi. *Flores* sæpissime 5-meri. *Baccæ* læves, nigrae, subsphaericae, pisi magnitudine.

3. **G. stenophylla**, Thw.; glabra; ramulis compressis, ad foliorum axillas glandulosis ferrugineo-villosisque; foliis lineari-lanceolatis, acuminatis, basi acutis, brevi-petiolatis; stipulis rotundato-triangularibus, brevi-cuspidatis, cito deciduis; corymbis longe-pedunculatis, 5-radiatis, ebracteolatis?, ad nodos ferrugineo-villosis; floribus ad radiorum apices pluribus, confertis; calyce sinuatim 5-dentato, corollæ tubo brevissimo, fauce piloso.—c.p. 2609.

HAB. Damp forests, at no great elevation, in the central and southern parts of the island.

*Folia* 2–6 poll. longa, 4–12 lin. lata, *petiolo* 2–3 lin. longo. *Baccæ* læves, sphaericae, nigrae, pisi magnitudine.

4. **G. Gardneri**, Thw.; glabra; ramulis compressis; foliis cuneato-oblongis, subito longiuscule cuspidatis, petiolatis; stipulis triangularibus, subacutis, cito deciduis; corymbis pedunculatis, ter quaterve umbellatim ramosis, multifloris, bracteolatis; bracteolis latis, brevibus, rotundatis; floribus 3–4-nis, sessilibus; calyce obscure 5-dentato; corollæ tubo brevissimo; baccis sphaericis, lævibus, parvis.—c.p. 1718.

HAB. Central Province, at an elevation of 3000 to 6000 feet, not very common.

*Folia* 5–12 poll. longa, 1– $3\frac{1}{2}$  poll. lata, *petiolo* 2–8 lin. longo. *Baccæ* nigrae.

5. **G. Wightiana**, Thw.—*Lasianthus? dubius*, Wight, in Calc. Journ. Nat. Hist. vi. p. 516.—c.p. 586.

HAB. Banks of streams, in the Ambagamowa and Saffragam Districts, at no great elevation.

*Frutex* parvus. *Ramuli* compressi, pilosi, denique glabri. *Folia* ovato-vel oblongo-lanceolata, acuminata, basi cuneata, petiolata, superne glabra, subtus pilosa, ad nervos tomentosa, 4-8 poll. longa, 1½-4 poll. lata, *petiolo* 4-8 lin. longo. *Stipulae* oblongae, connatae, pilosae, margine scariosae. *Corymbi* tomentosi, basi trichotomi, bracteolati, *divisionibus primariis* apice 3-5-tomis, lateralibus saepe reflexis. *Bracteolae* parvae, obtusae. *Flores* sessiles, capitellati. *Calyx* tubulosus, breve 5-dentatus, interne glaber. *Corolla* pilosula, tubo brevi. *Baccæ* pilosae, subsphaericæ, 3-4 lin. in diam. (siccae), subcarinatae.

6. **G. affinis**, Thw.; *ramulis* compressis, scabridis; *foliis* oblongo-vel lineari-lanceolatis, acuminatis, basi cuneatis, longiuscule petiolatis, glabris vel subtus ad nervos pilosulis; *stipulis* oblongis, pilosis; *corymbis* tomentosis, bracteolatis, basi trichotomis, divisione intermedia saepe ad apicem trichotoma; *floribus* sessilibus, capitatis; *calyce* tubuloso, obscure 5-dentato; *baccis* pilosis, pisi magnitudine.—c.p. 2706.

HAB. Similar situations to the last species.

Præcedenti valde affinis et ejusdem forsitan varietas.

7. **G. Moonii**, Thw.; *ramulis* glabris, compressis; *foliis* glabris, oblongis, abrupte acuminatis, basi cuneatis, petiolatis; *stipulis* oblongis, connatis, apice ciliatis, minute 1-2-dentatis; *corymbis* glabris, bracteolatis, basi trichotomis; *divisionibus lateralibus* reflexis; *centrali* apice trichotoma; *floribus* sessilibus, capitatis, bracteolis majusculis rotundatis ciliatis intermixtis; *calyce* tubuloso, 5-dentato, margine ciliato; *baccis* oblongis (siccis), utrinque obscure 4-sulcatis.—c.p. 585.

HAB. Reigam Corle, at no great elevation.

*Folia* 4-8 poll. longa, ¼-3½ poll. lata, *petiolo* 4-8 lin. longo.

This species, with *G. Wightiana* and *G. affinis*, forms an approach to the next genus, the albumen of their seeds being less evidently ruminant than in the other species of *Grumilea* herein described, and more or less sulcate by the intrusion of folds of the testa. It would perhaps be most desirable to form of *Grumilea* a section of *Psychotria*, seeing that it is quite impossible to determine, in the absence of tolerably mature fruit, to which of the two genera certain plants should be referred.

## 16. PSYCHOTRIA, Linn.

§ *Seminibus, cum albumine incluso, dorso 4-sulcatis.*

1. **P. sarmentosa**, Blume, Bijl. p. 964; DC. Prodr. iv. p. 522.—*P. scandens*, Hook. et Arn. Bot. Beech. Voy. p. 193; Walp. Rep. ii. p. 471.—c.p. 1712.

HAB. Not uncommon, up to an elevation of 4000 feet.

*Flores* pallide virides, fauce intus flavo-pilosa. *Baccæ* albæ.

§§ *Seminibus, cum albumine incluso, dorso 2-sulcatis.*

2. **P. bisulcata**, W. et A., l. c. p. 434; Wight, Icon. t. 1039.—c.p. 158 (669).

Var. β; *foliis* longioribus, minus coriaceis; *floribus* parum minoribus; *seminibus* longioribus, latius sulcatis.—c.p. 1720. (An species distincta?)

HAB. Var. *a*. Abundant in the Central Province, at an elevation of 5000 to 7000 feet. Var. *β*. From 2000 to 4000 feet.

3. **P. plurivenia**, Thw.; glabra; ramulis compressis; foliis lineari-lanceolatis, acuminatis, basi acutis, breve petiolatis, pluriveniis; stipulis oblongis, longe acuminatis, cito deciduis; corymbis terminalibus, basi trichotomis; floribus ternis, pedicellatis; calyce longiuscule 5-dentato; corollæ tubo brevi, fauce hirsuto; baccis (siccis) fusiformibus; seminibus late sulcatis.—C.P. 2652.

HAB. Hewahette, in the Central Province, at an elevation of 3000 to 4000 feet.

Folia  $2\frac{1}{2}$ – $4\frac{1}{2}$  poll. longa, 5–12 lin. lata, petiolo 2 lin. longo.

This species is apparently very closely allied to *P. polyneura*, DC.

4. **P. longipetiolata**, Thw.; ramulis compressis; foliis glabris, ellipticis vel oblongis, basi rotundatis vel parce acutis, longe petiolatis; stipulis oblongis, acutis; corymbis minutissime pilosulis, pedunculatis, 5-radiatis; radiis trichotomis, cymellis 7-floris terminatis; calyce brevi, 5-dentato; corollæ tubo brevissimo; baccis oblongis; seminibus latissime sulcatis.—C.P. 2707.

HAB. Saffragam District, at an elevation of 3000 to 4000 feet.

Folia flavescens, 3–7 poll. longa, 1–3 poll. lata, petiolo 1–2 poll. longo.

5. **P. sordida**, Thw.; ramulis compressis, glabris vel minutissime puberulis; foliis subcoriaceis, oblongis, plus minus acuminatis, basi acutis, petiolatis, venis secundariis inconspicuis; stipulis oblongis, sæpe brevicuspidatis; pedunculis 3-nis, terminalibus, glabris vel minutissime puberulis; floribus sessilibus, 3–4-nis; calyce brevi, 5-dentato; corolla brevi, infundibuliformi, interne glabra; baccis oblongis.—C.P. 333, 355, 1716 (390, 442, 1713).

HAB. Central Province, at an elevation of 4000 to 8000 feet.

Folia 1–4 poll. longa,  $\frac{1}{2}$ – $1\frac{1}{4}$  poll. lata, petiolo 2–5 lin. longo.

A very variable species as regards the shape of the leaves, which are oblong-lanceolate, with a long acumination, or more rarely ovate and shortly acuminate.

## 17. PRISTIDIA, Thw. (gen. nov.).

*Calyx* brevissimus, annuliformis. *Corolla* infundibuliformis, (ante anthesin oblonga, apice acute 4-costata); tubo brevi; limbo 4-lobo. *Stamina* fauce inserta; filamentis brevibus; antheris majusculis, oblongis, exsertis. *Stylus* simplex. *Stigma* oblongum, bilobum.—Frutex glaber, 5–6-pedalis; ramulis lævibus, parum compressis; foliis lævibus, parce venosis, oblongis, acuminatis, basi subacutis, petiolatis; stipulis oblongis, basi connatis, apice bidentatis; cymis longe pedunculatis, minute bracteolatis, supra-axillaribus vel rarius terminalibus, divisionibus ultimis spiciformibus; floribus parvis, late sessilibus, secundis.

1. **P. divaricata**, Thw.—*Grumilea?* *divaricata*, Thw. MSS. in Herb. Hook.—C.P. 2991.

HAB. Māpalagam, near Galle, at no great elevation.

Folia 3–7 poll. longa, 1– $2\frac{1}{2}$  poll. lata, petiolo 3–4 lin. longo. Cymæ foliis subæquilongæ. *Corolla* pallide viridis, externe minute puberula, interne parce pilosa.

I suspect that this species is polygamo-dioicous, and that I have only the barren flowers: the fruit I have not seen. *Psychotria tortilis*, Blume, is probably a congener of this plant.

## 18. CHASALIA, Comm., DC.

1. **C. curviflora**, Thw. — *Psychotria curviflora* et *P. ophioxylodes*, Wall. in Roxb. Fl. Ind. ii. pp. 167, 168; DC. Prod. iv. p. 520. *P. ambigua*, W. et A., l. c. p. 433.—c.p. 1714 (133, 391, partim).

HAB. Not uncommon, in shady places, up to an elevation of 4000 feet.

*Flores* polygamo-dioici, *fertiles* breviores, stigmatē exserto.

The structure of the fruit of this plant so precisely resembles that of *Chasalia*, as described by authors, that there can scarcely be a doubt of the propriety of referring it to that genus, as suggested by Wallich, at a time, too, it would appear, he was not acquainted with the fruit, since he has not described that organ.

## 19. GEOPHILA, Don.

1. **G. reniformis**, Don, (W. et A., l. c. p. 436, cum syn.; Wight, Icon. t. 54).—c.p. 1705.

HAB. Not uncommon in the Central Province.

## 20. SERISSA, Comm.

Bl. Bijl. p. 969. Endl.

*Dysodidendron*,\* Gardner, in Calc. Journ. of Nat. Hist. vol. vii. p. 2.

1. **S. Ceylanica**.—*Dysod. Ceylanicum*, Gard. l. c.—c.p. 261 (344, 433, 1710, 1711).

HAB. Not uncommon in forests of the Central Province.

*Frutex* 8–12-pedalis, glaber, *ramis* dichotomis, parum compressis, pallidis. *Folia* membranacea, oblonga vel oblongo-lanceolata, acuminata, basi plus minus acuta, brevi-petiolata, 2½–6 poll. longa, ¾–2½ poll. lata, *petiolo* 1–3 lin. longo. *Stipulæ* breves, plus minus connatæ, aristato-3-cuspidatæ, deciduæ. *Pedunculi* axillares terminalesque, 1–3-flori, folio multum breviores. *Pedicelli* basi apiceque 2-bracteolati, *bracteolis* connatis. *Flores* 4-meri. *Calyx* sinuato-4-dentatus. *Corolla* infundibuliformis, extus minute pubescens, fauce villosa, *limbo* 5–7 lin. in exp. *Baccæ* cyaneo-purpureæ, ovoideæ vel oblongæ, 4–9 lin. longæ, 1-loculares, 1–2-spermæ.—*A* *descript. cl. Gardner præcipue excerpt.*

2. **S. Gardneri**, Thw.; fruticosa, glabra, dichotomo-ramosa; *foliis* ovatis vel ellipticis, acuminatis, subsessilibus, basi rotundatis cordatisve; *stipulis* connatis, aristato-3-cuspidatis; *floribus* paucis, terminalibus, subsessilibus; *calyce* acute 4-fisso.—c.p. 82.

HAB. Forests about Galagama, not uncommon.

*Folia* 1¼–4 poll. longa, ½–2 poll. lata. *Corolla* alba, *limbo* 3 lin. in exp. *Baccæ* purpureo-cæruleæ, subsphæricæ, 4–5 lin. in diam.

This would seem to differ very slightly, except in having fewer flowers, from Dr.

---

\* I can find no structural character of sufficient importance to distinguish generically Dr. Gardner's species of *Dysodidendron* from *Serissa* of Commerson, as described by Blume; though the dichotomous ramification and much larger leaves of the former give them a different aspect to *Serissa foetida*, Comm., with its virgate habit, small leaves, and abbreviated ramuli. Dr. Wight, in describing his *Lasianthus? foetens* in Calc. Journ. vol. vi. p. 517 (*Dysod. Wightii*, Gard.), remarks upon its accordance in many respects with *Serissa*.

Gardner's *Dysod. glomeratum*, from the Peninsula of India; but I have no specimens of the latter for comparison.

3. **S. scabrida**, Thw.; fruticosa; ramulis scabridis; foliis ellipticis, acuminatis, basi acutis, glabris; costa petioloque scabridis; stipulis connatis, aristato-fimbriatis, scabridis; floribus terminalibus, sessilibus; calyce piloso, dentibus 4, lineari-lanceolatis.—C.P. 3306.

HAB. Near Ratnapoora.

The single specimen I possess of this plant has leaves 2 to 3 inches long and  $\frac{3}{4}$  to  $1\frac{1}{4}$  inch wide, with a petiole about a quarter of an inch long. The flowers and ripe fruit I have not seen.

## 21. HYDROPHYLAX, Linn.

1. **H. maritima**, Linn. (W. et A., l. c. p. 441, cum syn.; Wight, Icon. t. 760.)—C.P. 1672.

HAB. Common near the sea-shore. Nom. vulg. "Moohoodoo-gattacola.

## 22. RUBIA, Tourn.

1. **R. cordifolia**, Linn. (W. et A., l. c. p. 442, cum syn.; Wight, Icon. t. 187.)—C.P. 2705.

HAB. Abundant near Badulla, Mr. Ondaatze. Not uncommon throughout the Central Province.

## 23. GALIUM, Linn.

1. **G. asperifolium**, Wall. (W. et A., l. c. cum syn.)—C.P. 1682.

HAB. Central Province, at an elevation of 5000 to 7000 feet.

## 24. BORRERIA, Mey., DC.

1. **B. ocymoides**, DC. Prod. iv. p. 544; Burm. Fl. Ind. xxxiv. t. 13. fig. 1.—C.P. 1684.

HAB. South of the island, Moon, Gardner, Ferguson.

## 25. SPERMACOCE, Linn.

1. **S. hispida**, Linn. (W. et A., l. c. p. 438, cum syn.)—An *S. articularis*, Linn. fil.; W. et A., l. c. cum syn.?—C.P. 1695, 1696.

HAB. Very abundant.

I find it quite impossible to separate satisfactorily what I take, judging from the relative length of the corolla-tube, to be these two reputed species, and doubt their being really distinct.

‡‡ *Ovulis pendulis.*

## 26. KNOXIA, Linn.

1. **K. corymbosa**, Willd. (W. et A., l. c. p. 439, cum syn.; Wight, Illust. ii. t. 128.)—C.P. 3619 (1691 partim).

HAB. Central Province, up to an elevation of 5000 feet.

*Corymbi rami ultimi pauciflori. Fructus*  $1\frac{1}{2}$  lin. longi, 1 lin. lati.

2. **K. stricta**, Gært. De Fruct. i. p. 122. t. 25.—*K. mollis*, Br.; W. et A., l. c. cum syn.—C.P. 3620 (1691 *partim*).

HAB. Common in the Central Province, on the hills.

*Corymbi* rami ultimi pluriflori. *Fructus*  $\frac{3}{4}$  lin. longus.

Differs from the last in its longer, narrower leaves, its subumbellate and elongated branches of the corymb, and in its much smaller cocci.

3. **K. Zeylanica**, Linn. (W. et A., l. c. p. 440, cum syn.; excl. *K. stricta*, Gært.)—C.P. 1690.

HAB. Common in the southern parts of the island, up to no great elevation. Nom. vulg. "Ella-rat-mool."

4. **K. platycarpa**, Arn. Pug. 26 (344); Walp. Rep. ii. p. 468.—C.P. 1689 (2470).

Var.  $\beta$ . *hirsuta*.—*K. hirsuta*, Arn. l. c.; Walp. l. c. p. 469.—C.P. 1688.

Var.  $\gamma$ . *foliosa*; *foliis* majoribus, plus minus hirsutis, ovatis vel lanceolatis; *corymbis* foliosis; *lobis calycinis* longioribus (uno maximo sicut in varr. aliis).—C.P. 51 (118, 210, 422, 505).

HAB. Very abundant in the Central Province, at an elevation of 4000 to 7000 feet. Varr.  $\alpha$  and  $\beta$ , amongst grass; var.  $\gamma$ , in damp, rich ground.

An extremely variable species as regards the size of the leaves and amount of pubescence on the several parts of the plant, but the several forms merge insensibly into one another.

## 27. CANTHIUM, Lam.

1. **C. didymum**, Gært. (W. et A., l. c. p. 425, cum syn.)—*C. umbellatum*, Wight, Icon. t. 1034. *C. undulatum*, Champion, in Hook. Kew Journ. of Bot. iv. p. 198.—C.P. 1667, 3511.

Var.  $\beta$ . *lanceolatum*.—*C. lanceolatum*, Arn. Pug. 24 (342).—C.P. 2513.

Var.  $\gamma$ . *rostratum*; *foliis* ellipticis, rostrato-acuminatis; *floribus* paucioribus, subumbellatis, parum longioribus; *fructibus* parum majoribus.—C.P. 2607.

Var.  $\delta$ . *grandifolium*; *foliis* multo majoribus, superne nitidissimis.—C.P. 3419.

HAB. Varr.  $\alpha$  and  $\beta$ , abundant up to an elevation of 4000 feet; varr.  $\gamma$  and  $\delta$ , near Ratnapoora.

2. **C. montanum**, Thw.; arboreum, inerme; *foliis* valde coriaceis, ovatis vel orbicularibus, obtusis, pauciveniis, brevi-petiolatis; *pedunculis* axillaribus; *floribus* 5-meris, subumbellatis.—C.P. 2610, 2653.

Var.  $\beta$ . *minus*; *foliis* minoribus, utrinque parum angustatis; *pedicellis* post anthesin recurvis.—C.P. 1685.

HAB. Central Province, at an elevation of 4000 to 7000 feet.

Præcedenti valde affinis, et quoad inflorescentiam varietati  $\gamma$  ejusdem simillima, forsân varietas.

3. **C. parviflorum**, Lam. (W. et A., l. c. p. 426, cum syn.)—C.P. 1662 (1653).

HAB. Very abundant up to an elevation of 4000 feet.

4. **C. macrocarpum**, Thw.; fruticosum, subscandens, spinosum; *ramulis* teretibus, rufo-pilosulis; *foliis* lanceolatis, acuminatis, basi

rotundatis, utrinque rufo-pilosulis; *petiolo nervisque* tomentosus; *cymis* pilosis, axillaribus, brevibus, pedunculatis; *pedicellis* calyce subæquilongis; *floribus* 5-meris; *baccis* majusculis, rotundatis, parum compressis, emarginatis; *seminibus* tuberculatis.—C.P. 2816.

HAB. Hantani, at an elevation of 3000 feet.

*Folia* 1½–3 poll. longa, ¾–1½ poll. lata, *petiolo* 2–3 lin. longo. *Calyx* minute 5-dentatus. *Corolla* fusco-albida, *lobis* lanceolatis, *tubo* interne piloso parum longioribus. *Stamina* exserta. *Baccæ* pollicares.

5. **C. Rheedei**, DC. (W. et A., l. c. cum syn.; varr. β et γ.)—C.P. 1666.

Var. δ. *minus*; *foliis* minoribus; *spinis* rectis, adscendentibus; *alabastris* subito attenuatis, acutis.—C.P. 3420.—An species distincta?

HAB. Southern and central parts of the island, up to no great elevation. Var. δ. Near Ratnapoora.

6. **C. campanulatum**, Thw.; fruticosum, subscandens, spinosum; *ramulis* teretibus, pilosis; *foliis* parvis, glabris, rotundatis, ovatis vel lanceolatis, obtuse acuminatis, apiculatis, petiolatis; *fasciculis* 1–4-floris, axillaribus; *pedicellis* petiolo multum longioribus; *alabastris* acutis, medio constrictis; *corolla* campanulata; *limbo* 5-lobo; *baccis* obcordatis, compressis; *seminibus* tuberculatis.—C.P. 2472.

HAB. Deltotte, in the Central Province, at an elevation of 3000 feet; not common.

*Folia* 5–14 lin. longa, 3–8 lin. lata, *petiolo* 1–1½ lin. longo. *Pedicelli* glabri, tenues, 3–4 lin. longi. *Calyx* glaber, crateriformis, minute 5-dentatus. *Corollæ* *lobi* 2 lin. longi, *tubo* prope basin piloso æquilongi. *Antheræ* parum exsertæ. *Stylus* conicus. *Stigma* obconicum, utrinque truncatum, indivisum. *Baccæ* pisi magnitudine.

## 28. TIMONIUS, Rumph., DC.

*Nelitris*, Gærtner. *Eupyrena*, W. et A.

1. **T. Iambosella**, Thw.—*Nelitris Iambosella*, Gærtner. De Fruct. i. p. 134. t. 27 (excl. syn.). *Eupyrena glabra*, W. et A., l. c. p. 422.—C.P. 157 (664).

HAB. Very common in the southern and central parts of the island, up to an elevation of 5000 feet.

*Arbor* parva, 8–10-pedalis. *Folia* subtus ad costam glanduloso-porosa. *Inflorescentia* axillaris, polygamo-dioica. *Corolla* ochracea, *lobis* comosis, interne longitudinaliter 4–6-sulcatis, subimbricatis. FLOR. FÆM. Solitarii, cum apice pedunculi plus minus elongati articulati, bibracteolati. *Ovarium* sæpissime 12-loculare. *Stylus* simplex. *Stigmata* 12, in fasciculis 4 coalita. *Pedunculi* masc. bifurci, pauciflori. *Flores* laxè spicati, subsecundi. *Antheræ* lineari-oblongæ, subsessiles, inclusæ. *Stylus* apice minute bifidus.

### Sect. 2. *Corollæ segmentis contorto-imbricatis.*

† *Ovarii loculis* 1-ovulatis.

## 29. GUETTARDA, Linn.

1. **G. speciosa**, Linn. (W. et A., l. c. 422, cum syn.; Wight, Icon. t. 40.)—C.P. 2990.

HAB. Caltura, Moon. Near Galle. Nom. vulg. "Nil-pitcha."



30. *COFFEA*, Linn.

1. *C. Travancorensis*, Herb. Mad.; W. et A., l. c. p. 435.—*C. triflora*, Moon's Cat. p. 15 (non Forst.)—c.p. 2458.—An *C. Benghalensis*, Roxb. var.?

HAB. Warm, moister parts of the island, up to an elevation of 3000 feet. Nom. vulg. "Gas-pitcha."

*Frutex* 4-8-pedalis. *Ramuli* adscendentes, pilis brevibus scabri. *Folia* 1-4 poll. longa,  $\frac{1}{2}$ -2 poll. lata, *petiolo* circiter 1 lin. longo. *Baccæ* sphaericæ vel parum oblatae, nigrae, 4-5 lin. in diam. (siccae) subdidymæ. *Semina* hemisphaerica, ad hilum excavata.

2. *C. Wightiana*, Wall. (W. et A., l. c. p. 436; Wight, Icon. t. 1598.)—c.p. 1654.

HAB. Hot, drier parts of the island.

Differing from the last principally in its more horizontal ramification and smaller leaves, but, as remarked by Wight and Arnott, the two are probably forms of one species.

3. *C. ? elliptica*, Thw.; dioica, glaberrima; *foliis* subcoriaceis, integris, ellipticis vel oblongis, acuminatis, basi angustatis, petiolatis; *stipulis* majusculis, oblongis, acutis, convolutis; *pedunculis* terminalibus, ternis; *pedicellis* subæquilongis, ternis, folio multum brevioribus; *bracteolis* paucis, parvis, squamæformibus, versus basin pedicellorum sparsis; *calyce* integro vel minutissime denticulato; *corollæ tubo* intus ad apicem piloso, *lobis* oblongis obtusis duplo longiore. MASC. *Antheris* sessilibus, linearibus, apiculatis, inclusis, medio affixis; *stylo* tubo brevioris, subclavato; *stigmatis* lobis angustis, cohærentibus; *ovarii* rudimento subnullo.—c.p. 3482.

HAB. Hinidoon Corle, at no great elevation.

*Arbor* parva 10-12-pedalis, (sicca) nigrescens. *Folia* 4-10 poll. longa, 2-3 poll. lata, *petiolo*  $\frac{1}{2}$ -1 $\frac{1}{2}$  poll. longo. *Calycis tubus* 1 lin. longus. *Corolla* alba, *tubo* 12 lin. longo, *lobis* oblongis, obtusis, 8 lin. longis. *Fl. foem.* ignoti.

This species is placed provisionally only with *Coffea*, with which it agrees in many respects, until the structure of the female flower is known: it will then probably be found to be generically quite distinct.

*C. Arabica*, L., so extensively cultivated in the island, springs up in the forests from seeds carried by birds, monkeys, etc., but cannot be considered indigenous.

31. *PRISMATOMERIS*, Thw.

Hook. Kew Journ. Bot. viii. p. 268.

*Calyx* cupuliformis, subinteger. *Corollæ* segmenta carnosæ, prismatica, basi excavata. *Filamenta* brevia. *Antheræ* inclusæ, apicibus exsertis.

1. *P. albidiflora*, Thw. l. c. t. 7 A.—c.p. 728.

HAB. Not very uncommon in the southern and central parts of the island, up to an elevation of 4000 feet.

This plant would appear, at first sight, to offer an exception to the character of the section in which it is placed, but the convolute æstivation of the corolla is of necessity modified owing to the peculiar form and fleshy structure.

32. *IXORA*, Linn.

1. *I. coccinea*, Linn. (W. et A., l. c. p. 427; Wight, Icon. t. 153.)—c.p. 190.

HAB. Common in the warmer parts of the island, up to an elevation of 2000 feet. Nom. vulg. "Ratambala."

2. *I. calycina*, Thw.; glabra; *foliis* ovatis vel lanceolatis, acuminatis, basi cordatis rotundatis vel acutis, brevi-petiolatis; *venis primariis* subhorizontalibus; *corymbis* terminalibus, trichotomis; *pedicellis* brevibus, confertis; *bracteolis* longiusculis, linearibus; *segmentis calycinis* integris, lanceolatis, acutis, ovario 3-5-plo longioribus; *corollæ lobis* oblongo-rotundatis, acutis, *tubo* cylindrico multum brevioribus; *baccis* ovoideis vel sphaericis, calyce coronatis.—C.P. 448.

HAB. Central Province, up to an elevation of 7000 feet.

*Arbor* parva. *Folia*  $1\frac{1}{2}$ -4 poll. longa,  $\frac{1}{2}$ -2 poll. lata, *petiolo* 1-2 lin. longo. *Corolla* alba, rubro-tincta.

Allied apparently to *I. polyantha*, Wight, Icon. t. 1066, but distinguished by its much smaller leaves and flowers, and its glabrous corymbs.

3. *I. jucunda*, Thw.; *foliis* glabris, lanceolatis vel ovato-lanceolatis, acuminatis, basi angustatis, petiolatis; *corymbis* terminalibus, pilosulis, multifloris, basi trichotomis; *divisionibus primariis* elongatis; *bracteolis* parvis, acutis; *segmentis calycinis* truncatulis, ovario brevioribus.—C.P. 702 (1715, 2455, 2573).

Var.  $\beta$ ; *foliis* anguste lanceolatis; *segmentis calycinis* subacutis, quam *corollæ* lobi angustioribus.—C.P. 3424.

Var.  $\gamma$ ; *tubo corollæ* multo brevior.—C.P. 2 (406, 739).—An species distincta?

HAB. Not uncommon up to an elevation of 4000 feet.

*Arbor* parva, 10-20-pedalis. *Folia* 4-8 poll. longa, 1-4 poll. lata, *petiolo* 2-5 lin. longo. *Corymbi* sæpe coccinei. *Corolla* alba, rubro-tincta, *lobis* circiter 2 lin. longis, oblongis, acutiusculis; *tubo*, var.  $\alpha$  and  $\beta$ , 7-14 lin. longo; var.  $\gamma$ , 2-3 lin. longo. *Baccæ* subsphaericæ, atro-rubræ.

The short tube of the corolla in var.  $\gamma$  gives it an appearance very different from that of the other varieties, and further investigation may prove it to be specifically distinct; but I can at present discover no other character of importance sufficient to justify its separation as a species, and especially, too, as a similar variation is observable in *Pavetta Indica*, L.

4. *I. parviflora*, Vahl, (W. et A., l. c. p. 429, cum syn.; Wight, Icon. t. 711.)—C.P. 2575.

HAB. Hot drier parts of the island, up to no great elevation.

5. *I. acuminata*, Roxb. Fl. Ind. i. p. 383; Roxb. et Wall. Fl. Ind. p. 393; Wight, Icon. t. 706.—C.P. 748.

Var.  $\beta$ ; *ramulis foliisque* utrinque velutinis.

HAB. Not very uncommon up to an elevation of 3000 feet. Var.  $\beta$ . Hot drier parts of the island.

In my specimens the stipules are rounded and cuspidate, and the throat of the corolla is not hairy, but I cannot doubt their being specifically identical with Roxburgh's plant.

### 33. PAVETTA, Linn.

#### § *Floribus corymbosis.*

1. *P. Indica*, Linn. (W. et A., l. c. p. 431, cum syn.; Wight, Icon. t. 148.)—C.P. 1663 and 2456.

Var.  $\beta$ ; *corymbis* delicatulis, paucifloris; *pedicellis* elongatulis; *foliis* anguste lanceolatis, membranaceis.—C.P. 2611.

Var.  $\gamma$ ; *calyce* parum majore; *tubo corollæ* brevior quam in varr.  $\alpha$  et  $\beta$ .—C.P. 2512.—An *Ixora breviflora*, Roxb.?

Var.  $\delta$ ; *foliis* subtus *corymbisque* puberulis.—C.P. 3616.

HAB. Var.  $\alpha$ . Very abundant up to an elevation of 3000 feet. Var.  $\beta$ . Ambagamowa District. Var.  $\gamma$ . Common at an elevation of 3000 to 4000 feet. Var.  $\delta$ . Batticaloa. Nom. vulg. "Pawetta."

*Calyx* dentibus 4, acutiusculis, tubo 3-plo brevioribus, marginatus vel minute 4-denticulatus. *Corolla* externe glabra, *tubo* interne piloso.

2. **P. tomentosa**, Sm. (W. et A., l. c. cum syn.)—*Ixora tomentosa*, Roxb. Wight, Icon. t. 186.—C.P. 2815.

HAB. Abundant in the Batticaloa District.

*Dentes calycini* lanceolati, tubo longiores. *Corollæ lobi* externe parce pilosi.

This agrees so well in nearly every respect with Roxburgh's description and drawing, as copied in Wight's 'Icones,' that I cannot venture to separate them. The only difference of any importance appears to be the greater length of the calyx-lobes in the present plant.

3. **P. hispidula**, W. et A., l. c.—C.P. 742 (759, 771).

HAB. Central Province, at an elevation of 3000 to 5000 feet.

*Frutex* 2-4-pedalis. *Ramuli* teretes, tomentosi vel subglabri. *Folia* lanceolata vel obovato-lanceolata, acuminata, basi cuneata, membranacea, superne hispidula vel glabra, subtus tomentosa vel subglabra,  $2\frac{1}{2}$ -8 poll. longa, 1-3 poll. lata, *petiolo* 2-6 lin. longo. *Dentes calycini* parvi acutique vel *tubo* subæquilongi, obtusiusculi, vel rarius foliaceo-dilatati. *Corolla* albida. *Baccæ* nigre, nitidæ, subsphæricæ, semi-pollicares.

A variable species, distinguished from the preceding by its yellowish, rather large flowers, and its very long style, gradually attenuated from above the middle, and from the following by the very different shape of the leaves.

4. **P. angustifolia**, Thw.; *foliis* glabris (junioribus subtus ad nervos pilosulis), lineari-lanceolatis, utrinque acuminatis, petiolatis; *stipulis* membranaceis, apiculatis, externe hispidulis, interne glabris; *corymbis* brevibus, basi trichotomis, pilosulis; *pedicellis* longiusculis; *dentibus calycinis* acutis, apice recurvis; *tubo* brevioribus; *corollæ lobis* lineari-bus, acutis; *stylo* elongato, versus apicem attenuato.—C.P. 267.

HAB. Banks of streams, in the Central Province; not uncommon.

*Frutex* parvus. *Folia* 3-7 poll. longa,  $\frac{1}{2}$ -1 poll. lata, *petiolo* 1-5 lin. longo. *Corolla* albida, *tubo* 7-10 lin. longo, interne piloso, *lobis* 5 lin. longis. *Baccæ* pisi magnitudine.

### §§ *Floribus glomeratis.*

5. **P. involucrata**, Thw.; *foliis* glabris, obovatis vel obovato-lanceolatis, brevi-acuminatis, basi angustatis, rotundatis vel acutis, petiolatis; *capitulis* solitariis, axillaribus, arcte involucratis, pedunculatis.—C.P. 428 (458, 1717).

HAB. Central Province, at an elevation of 5000 to 7000 feet.

*Frutex* 10-12-pedalis, (siccus) nigrescens. *Folia*  $2\frac{1}{2}$ -5 poll. longa, 1-2 poll. lata, *petiolo*  $\frac{1}{4}$ - $\frac{1}{2}$  poll. longo. *Pedunculi* folio multum breviores. *Involucra* foliosa, *squamis* membranaceis, interne pilosis. *Calycis lobi* lineares, interne pilosi, *tubo* subæquilongi. *Corolla* albida, *lobis* acutis, *tubum* æquantibus. *Stylus* corolla duplo longior, apice pilosulus. *Baccæ* sphæricæ, nigre, pisi magnitudinæ.

†† *Ovarii loculis 2-pluriovulatis.*

### 34. EPITHINIA, Jack.

1. **E. Malayana**, Jack, (W. et A., l. c. p. 424, cum syn.)—c.p. 1665.  
HAB. Jaffna, *Gardner*.

As far as I can make out from dried specimens, the two ovules in each loculus appear to be attached to a central placenta, and one of them to be erect and the other pendulous.

### 35. HYPTIANTHERA, W. et A.

1. **H. macrocarpa**, Thw.; *foliis* glabris, oblongis, brevi-acuminatis, basi parum angustatis, petiolatis; *fasciculis* axillaribus vel paulo supra-axillaribus, paucifloris; *pedicellis* petiolo brevioribus; *calycis lobis* brevissimis, rotundatis, apiculatis; *baccis* oblongis, subsphæricis, paucispermis, 1-ocularibus; *seminibus* plus minus compressis; *albumine* ruminato.—c.p. 2709.

HAB. Ambagamowa District, and between Galle and Ratnapoora, at no great elevation.

*Frutex* 8-10-pedalis. *Folia* 5-8 poll. longa,  $1\frac{1}{2}$ - $2\frac{1}{2}$  poll. lata, *petiolo* circiter  $\frac{1}{2}$  poll. longo. *Corollæ tubus* cylindricus, 2 lin. longus, interne pilosus; *lobi* 5, breviores, rotundati, secus medium pilosi. *Antheræ* lineares. *Baccæ* subpollicares.

This species agrees in most of its characters with *Hyptianthera* of Wight and Arnott, but those authors make no mention of the seeds of their plant being ruminated.

### 36. SCYPHOSTACHYS, Thw.

*Calyx* brevis, tubulosus, 4-dentatus. *Corollæ tubus* brevis, *limbus* 4-lobus. *Antheræ* 4, oblongæ, subinclusæ. *Filamenta* brevia, fauce inserta. *Stylus* brevis. *Stigmata* 2, divergentia. *Orarium* 2-loculare; *loculis* circiter 4-ovulatis.—Frutices; *foliis lanceolatis*; *stipulis acuminatis, connatis*; *pedunculis axillaribus vel supra-axillaribus, ad apicem bracteolatis, paucifloris*; *bracteolis majusculis, connatis, vaginantibus*; *floribus sessilibus, parvis*. *Fructus baccatus*.

1. **S. pedunculatus**, Thw.; *ramulis* scabridis; *foliis* glabris vel subtus ad nervos pilosulis; *stipulis bracteolisque* striatis, cuspidatis, interne longe pilosis; *corolla* externe pilosa; *baccis* ovoideis.—c.p. 3423.

HAB. Pasdoon Corle, at no great elevation.

*Folia* 3-6 poll. longa,  $\frac{3}{4}$ - $2\frac{1}{4}$  lata, *petiolo* 2 lin. longo. *Pedunculi* 2-5 lin. longi, sæpissime deflexi. *Bracteolæ* sæpe 5-8, tanquam spicam formantes. *Baccæ* 7 lin. longæ, atro-purpureæ, sæpissime 1-spermæ. *Semina* lineari-oblonga, circiter 4 lin. longæ.

2. **S. coffeoides**, Thw.; *foliis* glabris, subcoriaceis; *pedunculis* brevibus vel subnullis; *bracteolis stipulisque* muticis, subscariosis, interne glabris; *corolla* externe glabra.—c.p. 2710.

HAB. Ratnapoora District, abundant, up to no great elevation. Nom. vulg. "Wal-copee."

*Folia*  $2\frac{1}{2}$ -5 poll. longa,  $\frac{3}{4}$ - $1\frac{3}{4}$  lata, *petiolo* 1-2 lin. longo. *Baccæ* rubræ. *Semina* oblonga, striata, circiter 2 lin. longæ.

## 37. DISCOSPERMUM, Dalzell, Walpers.

*Diplospora*, DC. (nomen prioritatem habens, sed quoad plantarum structuram decipiens).

1. **D. Dalzellii**, Thw.; *floribus* polygamo-dioicis. FÆM. *Ovarii loculis* 6-7-ovulatis.—An *D. sphaerocarpum*, Dalz. in Hook. Kew Journ. of Bot. ii. p. 257?—c.p. 561 (753, 767).

HAB. Central Province, at an elevation of 1000 to 3000 feet.

*Folia* 2-4 poll. longa,  $\frac{3}{4}$ - $1\frac{3}{4}$  lata, *petiolo* 3 lin. longo. *Flores* albidi, sessiles vel subsessiles. FÆM. *Corollæ tubus* brevior quam in flor. masc. *Ovula* prope marginem inferiorem placentæ subhemisphæricæ inserta. *Baccæ* sphaericæ, carnosæ, virides, pollicares.

The ovules of this plant are more numerous than described by Dalzell in his *D. sphaerocarpum*, and that author does not notice the dioicous inflorescence, but I suspect the two plants may prove identical.\*

2. **D. erythrosporum**, Thw.; *floribus* polygamo-dioicis; *calyce* minute 4-dentato; *staminibus* longiusculis. FLOR. FÆM. *Ovarii loculis* 3-4-ovulatis; *baccis* parvis, ovoideis vel subsphæricis; *seminibus* oblongis; *testa* rubro-glandulosa.—c.p. 2510, 3481.

HAB. Ambagamowa District, at an elevation of 3000 feet.

*Folia* 2-5 poll. longa, 1- $2\frac{1}{2}$  poll. lata, *petiolo*  $1\frac{1}{2}$ -3 lin. longo. *Flores* flavidi. *Baccæ* circiter 4 lin. in diam. *Semina* 3 lin. longa.

## 38. STYLOCORYNE, Cav.

1. **S. Webera**, A. Rich. (W. et A., l. c. p. 401, cum syn.; Wight, Icon. tt. 309, 584.)—c.p. 185.

Var.  $\beta$ . *montana*; *gemma* alabastrisque gummiferis; *floribus* majoribus.—*Webera cerifera*, Moon, Cat. p. 19.—c.p. 1649.

HAB. Very abundant throughout the island. Var.  $\beta$  in more elevated localities than var. *a*. Nom. vulg. "Tarrana-gass."

## 39. GRIFFITHIA, W. et A.

1. **G. fragrans**, W. et A., l. c. p. 400, cum syn.; Wight, Icon. t. 310.—c.p. 1234 (1651, 2522).

HAB. Not uncommon up to an elevation of 3000 feet.

*Ovarii loculi* circiter 8-ovulati, *ovulis* secus marginem placentæ oblongæ supra medium affixæ immersis.

2. **G. Gardneri**, Thw.; *inermis*; *foliis* glabris, nitidis, lanceolatis, acuminatis, basi angustatis, petiolatis; *corymbis* axillaribus, folio brevioribus, brevi-pedunculatis, parce pilosis, denique glabris; *pedicellis* elongatulis; *bracteolis* parvis; *calyce* minute 5-dentato, externe pilosulo; *corolla* externe subglabra, interne ad faucem pilosa; *lobis* lineari-

\* I cannot find any difference of importance. The calyx-limb differs much in the several species of this genus, and requires the generic character to be emended. It is 4-lobed in *D. sphaerocarpum* and *Dalzellii*; urceolate and entire in *D. apiocarpum*; subcampanulate and minutely 4-toothed in *D. erythrosporum*.—J. D. H.

lanceolatis; tubo duplo longioribus.—*Stylocoryne Ceylanica*, Gardn., MSS.—C.P. 657 (746).

HAB. Central Province, at an elevation of 2000 to 4000 feet.

*Arbor* parva, elegans, 15–20-pedalis. *Ramuli* teretes, parum compressi. *Folia* 2–4½ poll. longa, ¾–1½ poll. lata, *petiolo* 2–3 lin. longo. *Flores* flavidæ, *lobis* 4 lin. longis. *Placentæ* 8–12-ovulatae. *Baccæ* subsphæricæ, 4–5 lin. in diam.

3. *G. rugulosa*, Thw.; *foliis* rugulosis, ovatis vel ovato-lanceolatis, brevi-acuminatis, subtus ad nervos pilosis, denique glabris, petiolatis; *corymbis* adpresse pilosis, folio oppositis, ad axillam folii abortivi parvi spinæformis solitariis; *dentibus calycinis* brevibus, subulatis.—C.P. 245.

HAB. Ambagamowa, Saffragam, and Galle Districts; not common.

*Frutex*; *ramulis* striatulis, cinereis, *junioribus* pilosulis. *Folia* 2–5 poll. longa, 1–3 poll. lata, *petiolo* 2–3 lin. longo. *Stipulæ* rotundatae, cuspidatae, interne ad basin glandulosæ. *Baccæ* globosæ, pisi magnitudine.

#### 40. RANDIA, Houst.

1. *R. dumetorum*, Lam. (W. et A., l. c. p. 397, cum syn.; Wight, Icon. t. 580.)—*R. longispina*, DC., W. et A., l. c. p. 398; Wight, Icon. t. 582.—C.P. 1652.

HAB. Very abundant in the hotter parts of the island.

2. *R. uliginosa*, DC. (W. et A., l. c.; Wight, Icon. t. 397.)—C.P. 2936.

HAB. Trincomalee, *W. Morris, Esq.*

The unripe fruit of this plant, Mr. Morris informs me, is employed as a vegetable by the Cinghalese.

#### 41. GARDENIA, Ellis.

1. *G. carinata*, Wall. in Roxb. Fl. Ind. ii. p. 560; DC. Prod. iv. p. 380.—An *G. costata*, Roxb. var. ?—C.P. 3618 (1655 *partim*).

HAB. Batticaloa District, *Gardner*.

In the herbarium I have specimens with no locality indicated, and Mr. W. Ferguson has sent me from Colombo the leaves of another species of *Gardenia* ("Gallis-gass" of the Cinghalese), which seems to resemble in many respects *G. latifolia*, Ait., but Mr. F. is under the impression that the fruit is costate, so the matter must remain in doubt until flowers and fruit can be obtained for examination.

#### 42. WENDLANDIA, Bartl.

1. *W. Notoniana*, Wall. (W. et A., l. c. p. 403, cum syn.)—*W. bicuspidata*, W. et A. l. c.—C.P. 315.

HAB. Central Province, up to an elevation of 7000 feet.

The stipules are very variable in form; sometimes deeply bifid, but more frequently merely retuse at the apex, with the sinus mucronate, and often quite entire.

### LXXXII. VALERIANACEÆ.

#### 1. VALERIANA, Neck.

1. *V. Hardwickii*, Wall. (Roxb. Fl. Ind. i. p. 166.)—DC. Prod. iv.

p. 640.—*V. elata*, Don, Prod. Fl. Nep. p. 159. *V. Hookeriana*, W. et A. Prod. i. p. 444. *V. Arnottiana*, Wight, Ic. tt. 1045, 1046.—c.p. 150 (1729).

HAB. Newera Ellia, abundant.

The amount of pubescence upon the leaves varies very much. In my specimens the achenia are not at all hairy between the ribs.

### LXXXIII. DIPSACEÆ.

#### 1. DIPSACUS, Tourn.

1. **D. Leschenaultii**, Coult.—DC. Prod. iv. p. 647.—*D. Walkeri*, Arn. Pug. p. 26 (334); Wight, Ic. t. 1166.—c.p. 2404 (1730).

HAB. Newera Ellia, at an elevation of 7000 feet.

### LXXXIV. COMPOSITÆ.

#### 1. VERNONIA, Schreb.

1. **V. Javanica**, DC. Prod. v. p. 22, cum syn.—c.p. 1742.

HAB. Not uncommon, at the margins of damp woods, up to an elevation of 5000 feet: very abundant in the Ambagamowa District.

2. **V. Wightiana**, Arn. Pug. p. 27 (345); DC. Prod. vii. p. 263.—c.p. (1741).

Var.  $\beta$ ; *foliis* anguste lanceolatis, subtus parce tomentosis.—c.p. 164.—An species distincta?

HAB. Central Province, at an elevation of 5000 to 8000 feet. Var.  $\beta$ . In the Ambagamowa District.

The different form of the leaves, and their much smaller amount of pubescence, give a distinct appearance to var.  $\beta$ ; but it resembles var.  $\alpha$  so closely in the whole structure of the inflorescence that I cannot venture to separate them.

3. **V. Zeylanica**, Less. (DC. Prod. v. p. 26, cum syn.)—c.p. 1738.

HAB. Common, in the warmer parts of the island, up to an elevation of 3000 feet. Nom. vulg. "Poopoola."

4. **V. conyzoides**, DC. Prod. v. p. 25, cum syn.; Wight, Icon. t. 1076.—*V. Hookeriana*, Arn. Pug. p. 28 (346); DC. Prod. vii. p. 263.—c.p. 1737.

HAB. Central Province, at an elevation of 1000 to 4000 feet.

The outer series of the pappus in my specimens is very inconspicuous.

5. **V. cinerea**, Less. (DC. Prod. v. p. 24, cum syn.)—c.p. 1736.

HAB. Very abundant throughout the island. Nom. vulg. "Monera-koodimbeya."

6. **V. Neilgherryensis**, DC. l. c. p. 32; Wight, Icon. t. 1078.—*V. setigera*, Arn. Pug. p. 27 (345); DC. Prod. vii. p. 263.—c.p. 20 (169, 566).

HAB. Central Province, at an elevation of 6000 to 7000 feet.

7. **V. anthelmintica**, Willd. (DC. Prod. v. p. 61, cum syn.)—c.p. 487.

HAB. Galagama, Gardner. Common in native gardens, but I have never met with it truly wild.

8. **V. Gardneri**, Thw.; herbacea; *foliis* subsessilibus, lanceolatis vel anguste lanceolatis, acute serratis, subglabris, basi angustatis; *corymbo* oligocephalo; *capitulis* majusculis, multifloris, longe pedunculatis; *involucris squamis* lanceolatis, sæpe abrupte angustatis, squarrosis, *exterioribus* brevioribus angustioribusque; *achenis* cylindræis, pilosis, *pappo* *exteriore* brevissimo; *receptaculo* alveolo-fimbriato.—C.P. 27, 1745.

Var.  $\beta$ . *nervosa*; *foliis* firmis, ovatis vel rotundatis, acuminatis, venis subtus prominentibus, *invol. squamis* brevi-acuminatis.—C.P. 44.—An species distincta?

HAB. Central Province. Var.  $\alpha$ . In shady places, up to an elevation of 5000 feet. Var.  $\beta$ . On exposed, wet, rocky slopes, between Horton Plains and Galagama.

Var.  $\alpha$  of this species closely resembles *V. anthelmintica*, but the latter is distinguished by its smaller florets, obtuse involucre-scales, larger seeds, and naked receptacle. Var.  $\beta$  differs from var.  $\alpha$  in the shape of the leaves, and their veins being much more prominent beneath; but in other respects it seems to agree with undoubted forms of this very variable species.

9. **V. nemoralis**, Thw.; herbacea, *caule* minute adpresso piloso, *foliis* lanceolatis, acuminatis, distanter serratis, subglabris, basi angustatis, *capitulis* 20–30-floris, longiuscule pedunculatis, *involucris squamis* parce floccosis, lineari-oblongis, apice plus minus rotundatis, mucronulatis, *interioribus* subelongatis, *achenis* pilosis, *pappo* *exteriore* brevi, subpaleaceo, *receptaculo* alveolato-fimbriato.—C.P. 216.

HAB. Saffragam and contiguous districts, at no great elevation.

Closely allied to the preceding, from which it is principally distinguished by its fewer florets and the different shape of its involucre-scales.

10. **V. pectiniformis**, DC. Prod. v. p. 31; Wight, Ic. t. 1077.—*V. puncticulata*, DC. Prod. vii. p. 264.—C.P. 294 (567).

HAB. Central Province, to an elevation of 6000 to 7000 feet.

*Achænia* subglabra. *Pappus* plus minus rufescens, *serie* externa 3–4-plo brevior, decidua.

11. **V. scariosa**, Arn. Pug. p. 28 (346).—*Decaneurum?* *scariosum*, DC. Prod. vii. p. 264.—C.P. 389.

Var.  $\beta$ . *crassa*; *elatior*; *involucris squamis* villosis, latioribus, abrupte et longe subulato-acuminatis.—C.P. 2825.

HAB. Central Province, at an elevation of 4000 to 7000 feet. Var.  $\beta$ . In very damp shady places.

## 2. ELEPHANTOPUS, Cass.

1. **E. scaber**, Linn. (DC. Prod. v. p. 86; Wight, Icon. t. 1086).—C.P. 1772.

HAB. Very abundant in open places in the warmer parts of the island. Nom. vulg. "At-addeya."

## 3. AGERATUM, Linn.

1. **A. conyzoides**, Linn. (DC. Prod. v. p. 108, cum syn.)—C.P. 3633.

HAB. A most abundant weed throughout the island, from the sea-level to an elevation of 7000 feet.



4. **ADENOSTEMMA**, Forst.

1. **A. viscosum**, Forst. (DC. Prod. v. p. 111, cum syn.)—*A. latifolium*, Don; DC. l. c. p. 112; Wight, Ic. t. 1087. *A. erectum*, DC. l. c. p. 113. *A. reticulatum*, DC. l. c.; Wight, Icon. t. 1088. *A. angustifolium*, Arn. Pug. 29 (347); DC. Prod. vii. p. 266.—c.p. 594.

HAB. Very abundant throughout the island.

5. **LAGENOPHORA**, Cass.

1. **L. Harveyi**, Thw.; *foliis* spathulato-oblongis, sinuato-dentatis, plus minus hirsutis; *scapo* foliis 2-3-plo longiore; *involucris squamis* lineari-lanceolatis, subglabris; *achæniis* 3-serialibus, glabris, margine incrassato, rostro antice glanduloso; *receptaculo* conico.—c.p. 21 (513).

HAB. Central Province, at an elevation of 4000 to 6000 feet, not very uncommon.

The triserial arrangement of the achænia, pointed out to me by my friend Professor W. H. Harvey, when we were examining the plant together in the island, seems sufficient to separate this from the other described species of the genus.

6. **MYRIACTIS**, Less.

1. **M. Wightii**, DC. Prod. v. p. 308; Wight, Icon. t. 1091.—*M. Gmelini*, DC. l. c. p. 309.—An *M. Javanica* et *M. pilosa*, DC. l. c. p. 308?—c.p. 1740.

HAB. Very abundant in the Central Province, at an elevation of 5000 to 8000 feet.

7. **AMPHIRHAPIS**, DC.

1. **A. Zeylanica**, DC. Prod. vii. p. 279.—*Solidago Zeylanica*, Arn. Pug. 29 (347).—c.p. 152.

HAB. Abundant throughout the island.

8. **SPHÆRANTHUS**, Vaill.

(Nom. vulg. "Mooda-mahana.")

1. **S. microcephalus**, Willd. (DC. Prod. v. p. 369, cum syn.)—c.p. 1770.

HAB. Common in the warmer parts of the island.

2. **S. hirtus**, Willd. (DC. l. c. cum syn.; Wight, Icon. t. 1094.)—c.p. 1769.

HAB. Very abundant on the borders of paddy-fields.

3. **S. amaranthoides**, Burm. (DC. l. c. p. 370, cum syn.; Wight, Icon. t. 1149.)—c.p. 1768.

HAB. Not uncommon in the hotter parts of the island.

9. **DICHROCEPHALA**, DC.

1. **D. latifolia**, DC. Prod. v. p. 372, cum syn.; Wight, Icon. t. 1096.—c.p. 1766, 3303.

HAB. Very common in waste places.

## 10. GRANGEA, Adans.

1. **G. Maderaspatana**, Poir. (DC. Prod. v. p. 373, cum syn.; Wight, Icon. t. 1097.)—c.p. 1767.

HAB. Warmer parts of the island.

## 11. CONYZA, Less.

1. **C. Ægyptiaca**, DC. Prod. v. p. 382.—*C. semipinnatifida*, Wall.; DC. l. c. *Erigeron hispidum*, DC. l. c. p. 292, cum syn. *E. subliratum*, Roxb.; DC. l. c.—c.p. 1733.

HAB. North of the island, *Gardner*.

2. **C. viscidula**, Wall. (DC. l. c. p. 383.)—*C. Wallichii*, DC. l. c. p. 384, cum syn.—c.p. 1739.

HAB. Central Province, at an elevation of 1000 to 3000 feet.

## 12. BLUMEA, DC.

1. **B. amplectens**, DC. Prod. v. p. 433, cum syn.—c.p. 1730, 3523.

HAB. Hotter parts of the island.

2. **B. arenaria**, DC. l. c. cum syn.—c.p. 1732.

HAB. North of the island, *Gardner*.

3. **B. barbata**, DC. l. c. p. 434.—c.p. 1731.

HAB. Central Province, at an elevation of 3000 to 5000 feet.

4. **B. lacera**, DC. l. c. p. 436, cum syn.—*B. Wightiana*, DC. l. c. p. 435, cum syn.; Arn. Pug. 29 (347).—c.p. 1735.

HAB. Very abundant throughout the island, on the borders of paddy-fields, etc.

5. **B. hieraciifolia**, DC. l. c. p. 442; Wight, Icon. t. 1099.—c.p. 19, 403.

Var.  $\beta$ ; *caulibus foliisque subtus dense villosis*.

HAB. Central Province, at an elevation of 4000 to 7000 feet.

Dr. Wight's figure, quoted above, would seem to represent a stunted form or a very early stage of growth of the present plant, which here usually occurs from 3 to 5 feet or even more in height.

6. **B. crinita**, Arn. Pug. 30 (348); DC. Prod. vii. p. 283.—c.p. 147.

HAB. Adam's Peak and Newera Ellia, *Gardner*.

Closely allied to and very much resembling the preceding species, but apparently quite distinct. I have not seen the plant in a fresh state, my specimens having been collected by Dr. Gardner.

7. **B. alata**, DC. Prod. v. p. 448, cum syn.; Wight, Icon. t. 1101.—c.p. 551 (683).

HAB. Grassy places, in the Central Province, at an elevation of 4000 to 6000 feet.

8. **B. glandulosa**, DC. l. c. p. 438.—c.p. 1734, 2822.

Var.  $\beta$ ; *subglabra, eglandulosa*.—c.p. 560.

HAB. Common up to an elevation of 4000 feet. Var.  $\beta$ . In damp shady places.

9. **B. myriocephala**, DC. l. c. p. 445.—An *B. grandis*, DC. l. c. p. 447?—c.p. 1744 (2557).

HAB. Hantani, in the Central Province, at an elevation of 4000 feet.

Herbacea, 4-6-pedalis. *Folia* ad basin lobulis parvis paucis appendiculata. *Achænia* pubescentia.

10. **B. angustifolia**, Thw.; humilis; *foliis* glabris, integris, anguste lanceolatis, obtusiusculis, obscure et remote denticulatis, basi longe cuneatis; *corymbis* oligocephalis; *capitulis* plurifloris pedicellisque plus minus lanato-villosis; *involucris squamis* lineari-lanceolatis, apice fibrillatis, disco parum longioribus; *receptaculo* glabro; *achæniis* glabris.—C.P. 691.

HAB. Central Province, at an elevation of 1000 to 3000 feet.

Herbacea, subrecta, pedalis, basi procumbens, ramosa, radicans. *Folia* 1-3 poll. longa, 2-4 lin. lata. *Flosculi* flavidi. *Pappus* paucus, albus.

### 13. EPALTES, Cass.

1. **E. divaricata**, Cass. (DC. Prod. v. p. 461.)—*E. linearifolia*, DC. l. c. *E. pygmæa*, DC. l. c.—C.P. 234.

HAB. Very abundant in the warmer parts of the island. Nom. vulg. "Heen-moodoo-mahana."

### 14. VICOA, Cass.

1. **V. Indica**, DC. Prod. v. p. 474, cum syn.—C.P. 1757.

HAB. Abundant up to an elevation of 5000 feet.

### 15. ECLIPTA, Linn.

1. **E. erecta**, Linn. (DC. Prod. v. p. 490, cum syn.)—*E. prostrata*, Linn.; DC. l. c. cum syn. *E. parviflora*, Wall.; DC. l. c.—C.P. 1754.

HAB. Extremely abundant. Nom. vulg. "Keekerindi."

### 16. BLAINVILLEA, Cass.

1. **B. latifolia**, DC. Prod. v. p. 492, cum syn.—C.P. 1753.

HAB. Not uncommon in waste places.

### 17. SIEGESBECKIA, Linn.

1. **S. orientalis**, Linn. (DC. Prod. v. p. 495; Wight, Icon. t. 1103.)—C.P. 1752.

HAB. Common in waste places.

### 18. XANTHIUM, Tourn.

1. **X. strumarium**, Linn. (DC. Prod. v. p. 523, cum syn.)—*X. Indicum*, Roxb.; DC. l. c. cum syn.; Wight, Icon. t. 1104. *X. inæquilaterale*, DC. l. c. cum syn.—C.P. 1771.

HAB. Not uncommon in exposed sandy places.

### 19. MOONIA, Arn.

1. **M. heterophylla**, Arn. Pug. 31 (349); DC. Prod. vii. p. 289.

Var. *a*; *foliis* simplicibus.—C.P. 1758.

Var.  $\beta$ ; *foliis* 3-5-sectis.—*M. Arnottiana*, Wight, Icon. t. 1105.—c.p. 34.

HAB. Both varieties very abundant at an elevation of 6000 to 8000 feet.

Herbacea. *Achænia* ad apicem obscure bicornuta vel mutica.

## 20. WEDELIA, Jacq.

1. *M. calendulacea*, Less. (DC. Prod. v. p. 539, cum syn.; Wight, Icon. t. 1107.)—c.p. 1756.

HAB. Not uncommon in the hotter parts of the island, especially near the sea. Nom. vulg. "Ranwan-keekirindi."

## 21. WOLLASTONIA, DC.

1. *W. biflora*, DC. Prod. v. p. 546, cum syn.; Wight, Icon. t. 1108.—*W. asperrima*, DC. l. c. p. 547. *W. scabriuscula*, DC. l. c. cum syn. *W. glabrata*, DC. l. c. p. 548. *W. insularis*, DC. l. c.—c.p. 1760 (1759).

HAB. Common in the hotter parts of the island.

## 22. BIDENS, Linn.

1. *B. pilosa*, Linn. (DC. Prod. v. p. 597, cum syn.)—*B. Sundaica*, Bl.; DC. l. c. p. 598. *B. Wallichii*, DC. l. c. cum syn.—c.p. 3630.

HAB. A very common plant in cultivated ground. Nom. vulg. "Wal-teekola."

2. *B. decomposita*, Wall. (DC. l. c. p. 602, cum syn. P)—c.p. 3583.

HAB. Batticaloa District, not common.

*Involuceri squamæ* lineares, ciliatæ. *Ligulæ* flavæ, breves. *Achænia* elongata, 3-4-gona, subglabra vel versus apicem hirsuta, 3-4-aristata.

The achænia of *B. decomposita* are described as biaristate, whilst in my plant they are 3-4-aristate, but in other respects there would seem to be but slight difference, if any.

## 23. SPILANTHES, Jacq.

1. *S. Acmella*, Linn. (DC. Prod. v. p. 623, cum syn.)—*S. Pseudo-Acmella*, Linn.; DC. l. c. p. 625, cum syn. *S. calva*, DC. l. c. cum syn.—c.p. 684.

HAB. Very common in the Central Province, up to an elevation of 6000 feet. Nom. vulg. "Ac-malla."

## 24. ARTEMISIA, Linn.

1. *A. vulgaris*, Linn. (DC. Prod. vi. p. 112, cum syn.)—*A. Indica*, Willd.; DC. l. c. p. 114, cum syn.; Wight, Icon. t. 1111. *A. leptostachya*, DC. l. c. p. 113. *A. pauciflora*, Stechm.; DC. l. c. p. 102. *A. dubia*, Wall.; DC. l. c. p. 110. *A. Royleana*, Edgw.—c.p. 1755.

HAB. Not uncommon in or near native gardens. Nom. vulg. "Wal-kolondoo."

## 25. MYRIOGYNE, Less.

1. *M. minuta*, Less. (DC. Prod. vi. p. 139, cum syn.)—c.p. 2715.

HAB. Central Province, not uncommon.

## 26. GNAPHALIUM, Linn.

1. **G. Indicum**, Linn. (DC. Prod. vi. p. 231; DC. l. c. p. 274.)—C.P. 1762, 1764.

HAB. Very common in the Central Province, up to an elevation of 7000 feet.

*Caulis* simplex vel ramosus. *Capitula* in glomerulos paucos vel plurimos secus corymbi divisiones sparsos aggregata. *Involuceri squamæ* albidæ vel plus minus roseo-tinctæ.

2. **G. marcescens**, Wight, Icon. t. 1115.—C.P. 1765, 3522.

HAB. Central Province, at an elevation of 7000 to 8000 feet.

*Folia* superne glabra vel araneo-lanosa. *Achænia* (in meis specim.) glabra.

3. **G. Hookerianum**.—*Helichrysum Hookerianum*, W. et A.; DC. l. c. p. 201.—An *H. buddleioides*, DC. l. c. var. ?—C.P. 681.

HAB. Central Province, at an elevation of 6000 to 8000 feet.

Suffruticosum. *Receptaculum* luteo-glandulosum.

4. **G. Wightianum**, Wall. Cat. 2940.—*Anaphalis Wightiana*, DC. l. c. p. 273; Wight, Icon. t. 1117.—C.P. 528, 132, 568, 1763.

HAB. Central Province, at an elevation of 6000 to 8000 feet.

Suffruticosum. *Folia* lineari-oblonga, sæpe spathulata, late sessilia, superne subglabra vel plus minus lanuginosa, subtus dense cano-lanuginosa vel rarius subglabra.

Dr. Wight's figure, quoted above, so well represents a common form of this variable species, except that the capitula are somewhat larger in my specimens than shown in that figure, that there can be little doubt, I think, of the two plants being identical.

5. **G. brevifolium**.—*Anaphalis brevifolia*, DC. l. c. p. 273.—An *A. Neilgherriana*, DC. l. c. p. 272, et Wight, Icon. t. 478?—C.P. 1761.

HAB. Horton Plains and other of the most elevated parts of the Central Province.

## 27. GYNURA, Cass.

1. **G. lycopersicifolia**, DC. Prod. vi. p. 300, cum syn.—C.P. 420, 2826 (1750).

HAB. Very abundant up to an elevation of 5000 feet. Nom. vulg. "Kadoo-pāra."

2. **G. Nepalensis**, DC. l. c. cum syn.—C.P. 3539.

HAB. Galagama, at an elevation of 3000 feet.

3. **G. hispida**, Thw.; herbacea, subcarnosa, pilis flavis hispida; *caule* simplice vel parce ramoso, basi folioso; *foliis* obovato-lanceolatis, plus minus sinuato-dentatis, inferioribus in petiolum angustatis; *achæniis* glabris.—C.P. 3507.

HAB. On damp rocks, in the Central Province, at an elevation of 5000 to 7000 feet.

1-3-pedalis. *Folia* 2-6 poll. longa,  $\frac{3}{4}$ - $\frac{1}{2}$  poll. lata. *Capitula* 3-6, longe pedicellata, pluriflora, parce villosa. *Involucrum flosculis* flavo-aurantiacis parum brevius, *bracteolis* numerosis lineari-lanceolatis vix duplo longius.

Apparently very closely allied to *G. aurantiaca*, DC. (*Cacalia aurantiaca*, Blume).

## 28. EMILIA, Cass.

1. **E. sonchifolia**, DC. Prod. vi. p. 302, cum syn.—c.p. 3343.

HAB. Very abundant. Nom. vulg. "Kadoo-pāra."

An extremely variable plant as regards the shape of the leaves.

2. **E. prenanthoidea**, DC. l. c. p. 303, cum syn.—*E. angustifolia*, DC. l. c. cum syn.—An *E. scabra*, DC. l. c. ?—c.p. 430 (444, 581, 670).

HAB. Central Province, at an elevation of 6000 to 8000 feet.

*Folia* caulina cordato-sagittata, anguste lanceolata, vel rarius oblonga sublirataque, glabra vel (præcipue subtus) puberula. *Flosculi* purpurei.

## 29. DORONICUM, Linn.

1. **D. Walkeri**, Arn. Pug. 32 (350); DC. Prod. vi. p. 322.—c.p. 151 (688).

HAB. Central Province, at an elevation of 5000 to 7000 feet.

$\frac{1}{2}$ -3-pedale, plus minus pilosum. *Folia* oblonga, sessilia vel in petiolum longum basi sæpius appendiculatum angustata, dentata vel pinnatipartita, 1-6 poll. longa,  $\frac{1}{2}$ -2 poll. lata.

A most variable plant, the extreme forms of which, in the absence of the intermediate ones, it would be difficult to believe specifically identical.

2. **D. Gardneri**, Thw.; glabrum; *caule* adscendente, folioso; *foliis* lineari-lanceolatis, distanter grosse serratis, basi angustatis; *ligulis* 4-nerviis; *achæniis* pilosulis.—c.p. 435 (459).

HAB. Ambagamowa District, at an elevation of 3000 to 5000 feet.

$\frac{1}{2}$ -2-pedale. *Folia* 1-3 $\frac{1}{2}$  poll. longa, 2-4 lin. lata, subtus pallidiora. *Achænia* albida.

In the characters of the inflorescence, this very closely resembles the preceding species, but the very narrow smooth leaves seem sufficient to keep it distinct.

## 30. SENECIO, Less.

1. **S. corymbosus**, Wall. (DC. Prod. vi. p. 364; Wight, Icon. t. 1130.)—c.p. 565.

Var.  $\beta$ .—*S. Walkeri*, Arn.; DC. l. c.; Wight, Icon. t. 1131.—c.p. 272.

HAB. Central Province, not uncommon. Var. *a*. At an elevation of 7000 to 8000 feet.

An examination of numerous specimens has proved to me that the characters which have been employed to separate varr. *a* and  $\beta$  as distinct species, are by no means constant, so that I have no hesitation in uniting them.

2. **S. Walkeri**, Thw. (non Arn.; vide sp. prec. var.  $\beta$ .)—*Gynura Walkeri*, Wight, Icon. t. 1122.—c.p. 533.

HAB. Central Province, at an elevation of 7000 to 8000 feet.

This appears to be a true species of *Senecio*: it has not the styles of *Gynura*. The flowers are of a pale-yellow colour.

3. **S. Zeylanicus**, DC. l. c. p. 365.—*S. gracilis*, Arn. Pug. 31 (349).—An *S. Neilgherrianus*, DC. l. c. p. 368, var. ?—c.p. 1749.

HAB. Amongst high grass, at Newera Ellia and other of the most elevated parts of the island.

4. **S. Wightianus**, DC. l. c. p. 370, cum syn.; Wight, Icon. t. 1136.—c.p. 1751.

HAB. Common in the Maturatte and other districts of the Central Province.

*Folia* (in meis specim.) sæpissime ad basin pinnatim 2-4-lobata.

### 31. NOTONIA, DC.

1. *N. grandiflora*, DC. Prod. vi. p. 442, cum syn.—*N. corymbosa*, DC. l. c.; Wight, Icon. t. 484.—c.p. 1743.

HAB. Batticaloa District.

### 32. BRACHYRAMPHUS, DC.

1. *B. sonchifolius*, DC. Prod. vii. p. 177, cum syn.—*B. Heyneanus*, Wight, Icon. t. 1146. *Lactuca Heyneana*, DC. l. c. p. 140, cum syn.—c.p. 2827.

HAB. Galle, *Gardner*.

### 33. MICRORHYNCHUS, Less.

1. *M. sarmentosus*, DC. Prod. vii. p. 181, cum syn.; Wight, Illust. t. 133.—c.p. 1746.

HAB. Common near the seashore.

### 34. SONCHUS, Linn.

1. *S. oleraceus*, Linn.—c.p. 3631.

HAB. A very common weed in the Central Province, in newly-cleared ground.

### 35. YOUNGIA, Cass.

1. *Y. lyrata*, Cass. Opusc. iii. p. 86.—*Y. Mauritiana*, *Y. Thunbergiana*, *Y. runcinata*, *Y. napifolia*, *Y. Poosia*, *Y. striata*, et *Y. ambigua*, DC. Prod. vii. pp. 192, 193, cum syn. *Y. napifolia*, Wight, Icon. t. 1147.—c.p. 1747.

HAB. Very common in the Central Province, etc., up to an elevation of 5000 feet.

2. *Y. fuscipappa*, Thw.; glabra; *caulibus* sæpe e collo plurimis, erectis, parce foliosis; *foliis radicalibus* linearilanceolatis, acutis, runcinatis, in petiolum longum angustatis; *caulinis* summis linearibus; *corymbis* apertis.—c.p. 1748.

HAB. Central Province, at an elevation of 5000 feet.

1-2-pedalis. *Folia* 4-8 poll. longa,  $\frac{1}{2}$ - $\frac{3}{4}$  poll. lata. *Involucra* 3 lin. longa, minute bracteolata. *Achænia* fusiformia, parce compressa, puberula, striata. *Pappus* pallide fuscus.

## LXXXV. STYLIDIEÆ.

### 1. STYLIDIUM, Sw.

1. *S. uliginosum*, Sw. (DC. Prod. vii. p. 336.)—c.p. 2505.

HAB. South of the island.

## LXXXVI. GOODENOVIEÆ.

## 1. SCÆVOLA, Linn.

(Nom. vulg. "Taccāda.")

1. **S. Koenigii**, Vahl, (DC. Prod. vii. p. 505, cum syn.; Hook. fil. et T. Præcurs. ad Fl. Ind. in Linn. Soc. Journ. (Bot.) ii. p. 8, cum syn. seq.)—*S. Bela-Modagam*, R. & S.; DC. l. c. *S. Taccada*, Roxb. DC. l. c.; Wight, Illust. t. 137. *S. sericea*, Forst.; DC. l. c. 506. *S. Lambertiana*, De Vriese; Walp. Ann. ii. p. 1056. *S. chlorantha*, De Vriese; Walp. l. c.—C.P. 3632.

HAB. Very abundant between Colombo and Galle, and other parts of the coast.

2. **S. Plumieri**, Vahl, (DC. l. c. p. 506, cum syn.; Hook. fil. et T. l. c. cum syn. seq.)—*S. Thunbergii*, Eck. et Zeyh.; DC. l. c. *S. Senegalensis*, Presl; DC. l. c. *S. uvifera*, Stocks; Wight, Ic. t. 1613. *S. Macraei*, De Vriese (fide Auct.); Walp. l. c.—C.P. 1777.

HAB. Near Colombo, and other parts of the coast.

## LXXXVII. CAMPANULACEÆ.

## TRIBE I. CAMPANULEÆ.

## 1. WAHLENBERGIA, Schr.

1. **W. agrestis**, A. (DC. Prod. vii. p. 434; Wight, Icon. t. 1175; Hook. fil. et T. Linn. Soc. Journ. (Bot.) ii. p. 21, cum syn. seq.)—*W. dehiscens*, A. DC. l. c. *W. Indica*, A. DC. l. c.; Wight, Icon. t. 1176.—C.P. 1774.

HAB. Very abundant in grassy places in the Central Province, at an elevation of 4000 to 7000 feet.

## 2. CAMPANULA, Linn.

1. **C. fulgens**, Wall. in Roxb. Fl. Ind. ii. p. 99; DC. Prod. vii. p. 477; Wight, Icon. t. 1179; Illust. t. 136.—C.P. 1775.

HAB. Central Province, at an elevation of 5000 to 7000 feet, but not common.

## TRIBE II. LOBELIÆÆ.

## 3. LOBELIA, Linn.

1. **L. trigona**, Roxb. (DC. Prod. vii. p. 359; Wight, Icon. t. 1170.)—An *L. affinis*, Wall.; DC. l. c. p. 360? An *L. trialata*, Ham.; DC. l. c. ?—C.P. 1776, 2981.

HAB. Abundant throughout the island.

A most variable plant, and I have been unable to find sufficiently constant characters in any of its forms to justify my separating them as distinct species; nor have I ever met with specimens in which the stem, when in a fresh state, was not more or less triangular.



2. **L. excelsa**, Lesch. (DC. l. c. p. 381, cum syn.; Wight, Icon. tt. 1173, 1174.)—*L. nicotianæfolia*, Heyne; DC. l. c. cum syn.; Wight, Illust. t. 135. *L. aromatica*, Moon's Cat.; Wight, Icon. t. 1172.—c.p. 2592 (529, 531, 548).

HAB. Central Province, at an elevation of 4000 to 7000 feet; abundant at Newera Ellia and Maturatte. Nom. vulg. "Ras-nee."

The forms of this plant, which have been raised by some authors to the rank of species, merge so insensibly into one another, that I feel obliged to include them all under one specific name. The amount of pubescence upon the several parts, as well as the relative length of the calyx-lobes, are very variable, and I have found in the same raceme some pedicels with indications of the two minute bracteolæ, and others without the slightest trace of those appendages, and which latter seem to have their fullest development in plants of the most luxuriant growth.

## LXXXVIII. SPHENOCLEACEÆ.

### 1. SPHENOCLEA, Gærtn.

1. **S. Pongatium**, DC. Prod. vii. p. 548, cum syn.; Wight, Illust. t. 138.—c.p. 1773.

HAB. Warmer parts of the island; not uncommon.

## LXXXIX. VACCINIEÆ.

### 1. VACCINIUM, Linn.

1. **V. Leschenaultii**, Wight, Icon. t. 1188; Walp. Ann. i. p. 477.—*Agapetes? arborea*, DC. Prod. vii. p. 555, cum syn.—c.p. 270 (643).

Var.  $\beta$ ; *foliis lanceolatis vel ovato-lanceolatis; floribus pallidioribus.*—*Agapetes salicifolia*, Gardn. MSS.

HAB. Newera Ellia and other of the most elevated parts of the Central Province.

## XC. ERICACEÆ.

### 1. GAULTHERIA, Kalm.

1. **G. fragrantissima**, Wall. (DC. Prod. vii. p. 593, cum syn.; Wight, Icon. t. 1196.)—c.p. 42.

Var.  $\beta$ ; *ramulis foliisque subtus plus minus hispidis.*—*G. Leschenaultii*, DC. l. c.; Wight, Icon. t. 1195; Wight, Illust. t. 141 C.—c.p. 1790.

HAB. Central Province, at an elevation of 7000 to 8000 feet. Nom. vulg. "Kappooroo."

*Folia* elliptica vel oblongo-lanceolata, obtusa vel acuta, basi angustata, rotundata vel cordata. *Fructus* pulcherrime purpureo-cyanei.

### 2. RHODODENDRON, Linn.

1. **R. arboreum**, Sm. (DC. Prod. vii. p. 720, cum syn.; Wight, Icon. t. 1201; Wight, Illust. t. 140.)—c.p. 149 (521).

HAB. Central Province, at an elevation of 5000 to 8000 feet. Nom. vulg. "Ma-ratmal-gass."

*Folia subtus cano- vel pallide fulvo-lepidota.*

## XCI. LENTIBULARIÆ.

### 1. UTRICULARIA, Linn.

1. *U. flexuosa*, Vahl, En. Pl. i. 199; Oliver in Linn. Soc. Journ. (Bot.) iii. p. 175, cum syn. seq., scil.: *U. fasciculata*, Roxb. Fl. Ind. i. p. 143; Wall. Cat. 1499; Wight, Icon. t. 1568. *U. inæqualis*, Benjamin in Linnæa, xx. p. 304. *U. ramosa*, Vahl?—c.p. 2089.

HAB. Common in the warmer parts of the island.

2. *U. diantha*, Roem. et Sch. (non A. DC. Prod. viii. p. 21); Oliver, l. c. p. 176, cum syn. seq., scil.: Wight, Icon. t. 1569. *U. biflora*, Roxb. Fl. Ind. i. p. 43. *U. Roxburghii*, Spr. Syst. i. p. 52. *U. elegans*, Wall. Cat. 1502. *U. pterosperma*, Edgew. in Proc. Lin. Soc. i. 352.—c.p. 2828.

HAB. Not uncommon in wet paddy-fields.

3. *U. cærulea*, Linn. Herb. (non A. DC. Prod. viii. p. 19); Oliver, l. c. p. 179, cum syn., scil.: *U. pedicellata*, Wight, Icon. t. 1578. f. 2. *U. conferta*, Wight, Icon. t. 1575. *U. squamosa*, Wight, Icon. t. 1579. *U. uliginoides*, Wight, Icon. t. 1573.—c.p. 278 (2086).

Var.  $\beta$ . *affinis*; *scapo* robustiore, *floribus* minoribus, *calyce fructifero* latiore.—*U. affinis*, Wight, Icon. t. 1580. f. 1; Oliver, l. c. p. 178. *U. brachypoda*, Wight, Icon. t. 1578. f. 1.—c.p. 3308.

HAB. Central Province, at an elevation of 5000 to 8000 feet.

4. *U. reticulata*, Linn. (A. DC. Prod. viii. p. 19, cum syn.; Wight, Illust. ii. t. 143; Oliver, l. c. p. 180, cum syn. seq.)—*U. spiricaulis*, Miq. Pl. Hohen. 574.—c.p. 2090 (2091).

Var.  $\beta$ . *stricticaulis*; *scapo* firmo, brevior.—Oliver, l. c. cum syn. seq., scil.: *U. cærulea*, var. *stricticaulis*, Koen. in Hb. Brit. Mus. *U. uliginosa*, Vahl, En. Pl. i. 203. *U. humilis*, Heyne, Wall. Cat. 1495. *U. polygaloides*, Edgew. l. c. p. 351.—c.p. 2088.

HAB. Common in the hotter parts of the island. Nom. vulg. "Nilmonerasse."

5. *U. bifida*, Linn. Herb.; Oliver, l. c. p. 182, cum syn. seq., scil.: *U. biflora*, Wall. Cat. 1498. *U. antirrhinoides*, ejusd. 1498 b. *U. diantha*, A. DC. l. c. p. 21. *U. Wallichiana*, Benj. Bot. Zeit. 1845, 213. *U. humilis*, Wight, Icon. t. 1572. f. 2 (? Vahl.)—c.p. 2087.

Var.  $\beta$ . *Wallichiana*; *pedicellis* longioribus; *lobis calycinis* acutis.—*U. Wallichiana*, Wight, Icon. t. 1572. f. 1; Oliver, l. c.

HAB. Warmer parts of the island, common.

In my specimens of this plant, var.  $\alpha$  and  $\beta$  appear to merge insensibly into one another.

6. *U. capillacea*, Wight, Herb.; Wall. Cat. 6399; Oliver, l. c. p. 184.—c.p. 3307.

HAB. Near Colombo, Mr. W. Ferguson. Negumbo.

*Corolla* purpurea; *labio inferiore* 3-lobato, lobis lateralibus multo minoribus omnibus apice rotundatis; *calcare* lineari-conico, longiore, porrecto. *Lobi calycis fructiferi* subæquales, orbiculares.

7. **U. racemosa**, Wall. Cat. 1496; Wight, Icon. 1854, f. 1; Oliver, l. c. p. 186.—*U. cærulea*, A. DC. l. c. p. 19. An *U. filicaulis*, Wall. Cat. 1501; Oliver, l. c. p. 186.—c.p. 2085.

Var.  $\beta$ . *rosea*; *scapo* robustiore; *corollæ* labio inferiore majore et *calcar* obtusiusculum æquante vel superante.—*U. rosea*, Edgw. l. c. p. 352; Oliver, l. c. p. 184.—c.p. 277.

HAB. Var.  $\alpha$ . Warmer parts of the island. Var.  $\beta$ . At an elevation of 5000 to 6000 feet.

The points of difference between varr.  $\alpha$  and  $\beta$  are so unimportant that I feel quite satisfied that they are merely forms of one species. The colour of the corolla varies from a rather deep reddish-purple to nearly white: the flowers of var.  $\alpha$  are most frequently pale in colour.

8. **U. orbiculata**, Wall. Cat. 1500; Oliver, l. c. p. 187.—*U. glochidiata*, Wight, Icon. t. 1581.—c.p. 2085, 3350.

HAB. Common on wet rocks and mossy trunks of trees in the Central Province, at an elevation of 4000 to 6000 feet.

## XCII. PRIMULACEÆ.

### 1. LYSIMACHIA, Linn.

1. **L. ramosa**, Wall. Cat. 1490; DUBY in DC. Prod. viii. p. 65.—*L. alata*, Gardner, MS.—c.p. 1896.

HAB. Central Province, in shady places, at an elevation of 6000 to 8000 feet.

Glabra, 2-4-pedalis. *Caulis* erectus, alatus, ramosus. *Folia*  $1\frac{1}{2}$ -6 poll. longa,  $\frac{1}{2}$ - $1\frac{1}{2}$  lata, in *petiolum* brevem angustata. *Pedunculi* sesquipollicares. *Calycis* laciniæ (acumine 1 lin.) 3 lin. longæ. *Corolla* pallide lutea. *Capsula* 5-valvis, calyce subæquilonga. *Semina* triangularia, minute muricata.

2. **L. Japonica**, Thunb. Fl. Japon. 83. t. 16.—*L. deltoidea*, Wight, Illus. ii. p. 137. t. 144. *L. umbrosa*, Gardner, MS.—c.p. 1895.

HAB. Newera Ellia and Maturatte, at an elevation of 6000 to 7000 feet.

### 2. ANAGALLIS, Tourn.

1. **A. arvensis**, Linn.—Var.  $\beta$  *cærulea*.—c.p. 3131.

HAB. Oova District (possibly introduced).

## XCIII. MYRSINACEÆ.

### 1. MÆSA, Försk.

1. **M. Indica**, A. DC. Prod. viii. p. 80, cum syn.; Wight, Icon. t. 1206.—c.p. 1799.

HAB. Very abundant up to an elevation of 5000 feet. Nom. vulg. "Matabimbeya-gass."

### 2. EMBELIA, Burm.

1. **E. Ribes**, Burm. (A. DC. Prod. viii. p. 85.)—c.p. 1800.

HAB. Common up to an elevation of 4000 feet. Nom. vulg. "Wel-ambilla."

2. **E. robusta**, Roxb. Fl. Ind. ii. p. 287; A. DC. Prod. viii. p. 86.—*E. Tsjeriam Cottam*, Wight, Icon. t. 1209. *Basaal*, Rheede, Hort. Mal. v. p. 23. t. 12. An *Tsjeriam Cottam*, Rheede, l. c. p. 21. t. 11?—c.p. 1804.

HAB. Kornegalle and Batticaloa Districts.

### 3. SAMARA, Linn. Herb.

*Choripetalum*, A. DC.

1. **S. viridiflora**, Thw.—*S. Rheedei*, Wight, Icon. t. 1591. *Choripetalum viridiflorum*, A. DC. Prod. viii. p. 88. *Pu Walli*, Rheede, Hort. Mal. vii. p. 79. t. 42. An *C. aurantiacum*, A. DC. l. c. var.?—c.p. 610 (741).

HAB. Central Province, at an elevation of 3000 to 5000 feet.

*Flores* polygamo-dioici. MASC. *Filamenta* petalis multum longiora. *Ovarii rudimentum* parvum, *stylo* brevissimo. FÆM. *Ovarium* 3-ovulatum.

### 4. MYRSINE, Linn.

1. **M. capitellata**, Wall. in Roxb. Fl. Ind. ii. p. 295; A. DC. Prod. viii. p. 94. An *M. lucida*, Wall.; A. DC. l. c. p. 95?—c.p. 1806 (2599).

Var.  $\beta$ . *parvifolia*, A. DC. l. c., et var.  $\gamma$  *grandiflora*, A. DC. l. c.; Wight, Icon. t. 1211. An *M. umbellata*, A. DC. l. c.?—c.p. 623, 3009, 2598.

Var.  $\delta$ . *sessiliflora*; *floribus* sessilibus vel subsessilibus.—An *M. Porteriana*, Wall. List, 6525; A. DC. l. c.?—c.p. 285.

HAB. Central Province, very abundant up to an elevation of 8000 feet.

So variable is this plant as regards the shape and size of the leaves and length of the flower-pedicels, that some of its forms might well pass for distinct species, but that intermediate ones occur rendering it impossible to separate them. Some of my specimens have leaves varying from  $\frac{3}{4}$  to  $1\frac{1}{2}$  inch in length, whilst other specimens have them reaching to 7 inches, with a corresponding width; the former occurring upon the tops of the highest hills, and the latter in damp rich forests.

### 5. ARDISIA, Linn.

1. **A. paniculata**, Roxb. Fl. Ind. ii. p. 270; A. DC. Prod. viii. p. 126.—*A. Missionis*, var.  $\beta$  *verticillata*, A. DC. l. c. p. 130. *A. Courtalensis*, Wight, Icon. t. 1215.—c.p. 1802.

HAB. Central Province, not uncommon up to an elevation of 3000 feet.

2. **A. humilis**, Vahl, Symb. iii. p. 40; A. DC. l. c. p. 129 (in parte).—*Tinus humilis*, Burm. Fl. Zeyl. p. 222. t. 103.—c.p. 2829.

HAB. Near the seashore, Galee District, common. Nom. vulg. "Loonoo-dang-gass."

*Frutex* 1-6-pedalis. *Folia* ad apices ramulorum compressorum aggregata. *Panicule* terminales. *Drupæ* rubræ.

Well distinguished from the next species, with which it has been confounded, by its leaves clustered at the ends of the branchlets, its terminal panicles, and bright-red fruit.

3. **A. elliptica**, Thunb. nov. gen. pars 8. Upsal, 1795, ex Rœm. et Sch. Syst. iv. p. 517; A. DC. l. c. p. 138.—*A. umbellata*, Roxb. Fl. Ind. ii. p. 273. *A. polycephala*, Wall.; Wight, Illust. t. 145.—c.p. 1803.

Var.  $\beta$ . *solanacea*; omnino major.—*A. solanacea*, Roxb. Pl. Cor. i. t. 27. *A. humilis*, Wight, Icon. t. 1212; A. DC. l. c. p. 129 (in parte).

HAB. Var.  $\alpha$ . Not uncommon in the warmer parts of the island. Var.  $\beta$ . In gardens, and I have never found it in a wild state. Nom. vulg. "Balloo-dang."

*Racemi* axillares terminalesque, umbellæformes, *pedunculo* rarius ramoso.

4. **A. pauciflora**, Heyne, (Wall. in Roxb. Fl. Ind. ii. p. 279; A. DC. l. c. p. 127; Wight, Icon. t. 1214.)—c.p. 15 (28, 36).

HAB. Central Province, up to an elevation of 7000 feet.

In all my specimens the pedicels are smooth, or, when young, ferrugineo-squamulose, and not pilose, as represented in Dr. Wight's figure.

5. **A. divergens**, Roxb. Fl. Ind. ii. p. 275; A. DC. l. c. p. 130, cum syn.—c.p. 376 (740).

Var.  $\beta$ ; *foliis* sinuato-denticulatis.—An *A. crispa*, var.  $\beta$ , A. DC. l. c. 134, cum syn.?

HAB. Not uncommon in damp forests, up to an elevation of 5000 feet.

*Frutex* erectus, 3-6-pedalis. *Folia* membranacea, venosa, ad marginem creberrime tuberculato-glandulosa, 3-6 poll. longa, 1-3 poll. lata. *Flores* roseo-purpurei. *Drupæ* rubræ, 3-4 lin. in diam.

6. **A. Wallichii**, A. DC. l. c. p. 137, cum syn.—c.p. 2667.

HAB. Cultura, Moon. Damp shady forests in the Saffragam District.

*Suffrutex* humilis,  $\frac{1}{2}$ -2-pedalis, partibus junioribus ferrugineo-lepidotis. *Folia* oblonga vel obovata, abrupte obtuse acuminata, crenulata, venosa, basi angustata, nigro-viridia, subtus rufescentia, 4-8 poll. longa,  $1\frac{1}{2}$ -3 poll. lata. *Racemi* rubri, terminales axillaresque, sæpius multiflori simplicesque, folio breviores. *Pedunculi* angulares. *Pedicelli* sparsi, sursum incrassati, reflexi, flore subduplo longiores. *Corolla* rosea, laciniis 4 lin. longis. *Antheræ* flavæ. *Drupæ* rubræ, 4 lin. in diam.

## XCIV. ÆGICERACEÆ.

### 1. ÆGICERAS, Gærtn.

1. **Æ. majus**, Gærtn. (A. DC. Prod. viii. p. 142, cum syn.; Wight, Illust. ii. p. 141. t. 146.)—c.p. 1796 (1805).

HAB. Not uncommon on the coast.

## XCV. SAPOTACEÆ.

### 1. CHRYSOPHYLLUM, Linn.

1. **C. Roxburghii**, G. Don, (A. DC. Prod. viii. p. 162.)—*C. acuminatum*, Roxb. Fl. Ind. ii. p. 345.—c.p. 2689.

HAB. Not uncommon in the warmer parts of the island. Nom. vulg. "Lawooloo-gass."

*Folia* oblonga, abrupte rostrato-acuminata, 3-7 poll. longa,  $\frac{3}{4}$ -2 poll. lata; *petiolo* 3 lin. longo. *Nuculania* obscure pentagona, ferrugineo-tomentosa, denique glabra. *Semina* 8 lin. longa, 4 lin. lata.

The ripe fruit is eaten by the Cinghalese: the timber is used for building purposes.

## 2. SAPOTA, Plum.

1. **S. elengoides**, A. DC. Prod. viii. p. 176; Wight, Icon. t. 1218.—C.P. 1795.

HAB. Hot, drier parts of the island.

## 3. MIMUSOPS, Linn.

1. **M. Elengi**, Linn. (A. DC. Prod. viii. p. 202, cum syn.; Wight, Icon. t. 1586.)—C.P. 1794.

HAB. Warmer parts of the island. Nom. vulg. "Moonemal-gass."

The timber used for building, etc.

2. **M. Indica**, A. DC. l. c. p. 205, cum syn.; Wight, Icon. t. 1587.—C.P. 1793.

HAB. Very abundant in the hot, drier parts of the island. Nom. vulg. "Paloo-gass."

The timber of this valuable tree is extremely hard and strong, and very durable.

## 4. BASSIA, Kœnig.

*Calyx* 4-partitus. *Corolla* 6-12-loba. *Antheræ* subsessiles, 2-3-seriales.

1. **B. longifolia**, Linn. (A. DC. Prod. viii. p. 197, cum syn.; Wight, Illust. ii. t. 147.)—C.P. 1797.

HAB. Common in the hotter parts of the island, especially in native gardens. Nom. vulg. "Mee-gass."

The oil of the seeds, under the name of "Mee-tel," is used by the Cinghalese medicinally, and, when fresh, for cooking. The wood is valuable for building purposes.

## 5. DASYAULUS, Thw. (gen. nov.).

*Calyx* 4-partitus. *Corolla* 6-vel 8-loba. *Antheræ* corollæ lobis duplo numerosiores, subsessiles, 1-seriales, apice subulato-acuminatæ. Ceteroquin ut in *Bassia*.

1. **D. neriifolius**, Thw. (*Bassia neriifolia*, Moon's Cat. p. 36); *foliis* glabris, anguste oblongo-lanceolatis, obtusis, basi in petiolum brevem angustatis.—C.P. 235.

HAB. Common on the banks of rivers and streams in the warmer parts of the island. Nom. vulg. "Gang-mee."

*Arbor* mediocris. *Folia* 3-8 poll. longa, 6-12 lin. lata. *Fasciculi* 2-6-flori, versus apices ramulorum conferti. *Pedicelli* glabri, sesquipollicares. *Calyx* 4 lin. longus, lobis interioribus dorso sericeo-strigosis. *Corolla* ad medium 6-partita. *Ovarium* 6-loculare. *Nuculania* oblonga, acuta, 1-sperma,  $1\frac{1}{4}$  poll. longa. *Semina* albuminosa.

2. **D. microphyllus**, Thw.—*Bassia microphylla*, Hook. Ic. i. t. 74.

*B. parvifolia*, A. DC. l. c. p. 198.—c.P. 2648; foliis majoribus, c.P. 2830.

HAB. South of the island, not common.

*Corolla* 6-loba, rarius 5-loba.

My specimens, numbered c.P. 2830, were collected by Dr. Gardner near Galle, and have leaves a good deal larger than is usual in this species, but agree so well with it in other respects that I hesitate to separate them even as a distinct variety.

3. **D. Moonii**, Thw.; *ramulis junioribus* fulvo-tomentosis; *foliis* glabris, oblongis, obtuse acuminatis vel retusis, basi subacutis, petiolatis; *pedicellis* petiolo subæquilongis.—c.P. 3434.

HAB. Caltura, Moon; near Ratnapoora.

*Folia* 3-7 poll. longa,  $1\frac{1}{4}$ -3 poll. lata; *petiolo*  $\frac{1}{2}$ - $\frac{3}{4}$  poll. longo. *Fasciculi* 2-5-flori. *Corolla* 6-loba. *Nuculania* oblonga, acuta,  $1\frac{1}{2}$  poll. longa. *Semina* parce albuminosa.

4. **D. fulvus**, Thw.; *ramulis foliisque junioribus* dense fugaci-fulvo-tomentosis; *foliis* obovato-oblongis, abrupte brevi-acuminatis, basi angustatis, petiolatis; *pedicellis* petiolo subduplo longioribus.—c.P. 3446.

HAB. Hewessee, in the Pasdoon Corle.

*Arbor* mediocris. *Folia* 4-7 poll. longa,  $1\frac{1}{2}$ -3 poll. lata, *petiolo*  $\frac{3}{4}$  poll. longo. *Fasciculi* 6-8-flori. *Calyx* 3-4 lin. longus. *Ovarium* 6-loculare.

To the present genus is to be referred *Isonandra polyandra*, Wight, Ic. t. 1589.

## 6. DICHOPSIS, Thw. (gen. nov.).

*Calyx* 6-partitus, lobis 2-serialibus. *Corolla* 6-loba. *Stamina* 12, omnia fertilia. *Filamenta* brevia, 1-serialia, alterna longiora (corollæ lobis opposita) antheras subextrorsas, breviora (vix ulla) subintrorsas gerentia. *Antheræ* apice divaricato-bicornes. *Ovarium* 6-loculare.

Genus notabile, quoad stamina breviora *Dasyaulo*, longiora *Isonandræ* approximans.

1. **D. petiolaris**, Thw.—c.P. 3012.

HAB. Ambagamowa District, and the Hinidoon and Reigam Corles.

*Arbor* magna. *Folia* glabra, ovata, subabrupte brevi-acuminata, basi acuta, 3-5 poll. longa,  $1\frac{1}{2}$ -3 poll. lata; *petiolo* 1- $1\frac{1}{2}$  poll. longo. *Fasciculi* 2-5-flori, axillares, ad apices ramulorum conferti, fulvo-tomentosi. *Pedicelli* petiolo subæquilongi. *Calyx* 4 lin. longus. *Corolla* albida, 7 lin. longa, tubo interne pilosa. *Nuculania* subsphærica, 1-sperma,  $1\frac{1}{4}$  poll. in diam.

## 7. ISONANDRA, Wight.

*Calyx* corollæ lobis isomerus; *stamina* iisdem duplo numerosiora. *Filamenta* corollæ tubo subæquilonga. *Antheræ* extrorsæ, apice acuminatæ. *Nuculania* carnosæ.

### Sect. 1. *Calyx* 6-partitus.

1. **I. grandis**, Thw.; *foliis* glabris (junioribus cano-sericeis) obovatis, obtusis, abrupte brevi-acuminatis vel rotundatis, basi cuneatis, subtus prominentim venosis, petiolatis; *pedicellis* petiolo subæquilongis.—c.P. 619 (2507); foliis minoribus, c.P. 2402.

HAB. Central Province and Saffragam District, at an elevation of 3000 to 6000 feet. Nom. vulg. "Meeria-gass."

*Arbor* ingens. *Folia* 2-10 poll. longa, 1-6 poll. lata, *petiolo*  $\frac{1}{2}$ -1 poll. longo, superne plano. *Fasciculi* 4-6-flori, ad apices ramulorum conferti, adpresse fulvo-pilosi. *Calyx* 3-4 lin. longus. *Corolla* albida, calyce subduplo longior. *Ovarium* 6-loculare. *Nuculania* ovoidea, nigro-purpurea, sesquipollicaria. *Semina* exalbuminosa.

An oil is extracted from the seeds of this tree, which is used in the same way as that from the *Bassia longifolia*.

2. **I. rubiginosa**, Thw.; *foliis* coriaceis, ellipticis, obovatis vel obovato-oblongis, apice subacutis vel rotundatis, basi angustatis, subtus dense rubiginoso-pilosis, petiolatis; *pedicellis* petiolo æquilongis vel 2-3-plo longioribus.—c.p. 394, 3447.

HAB. Saffragam District and Hinidoon Corle, up to an elevation of 4000 feet.

*Arbor* ingens. *Folia*  $2\frac{1}{2}$ -7 poll. longa,  $1\frac{1}{4}$ -3 poll. lata, *petiolo*  $\frac{1}{2}$ - $\frac{3}{4}$  poll. longo, superne convexo. *Fasciculi* 6-10-flori, adpresse rubiginoso-pilosi. *Calyx* 4 lin. longus. *Ovarium* 6-loculare. *Nuculania* ovoidea.

3. **I. canaliculata**, Thw.; *foliis* glabris, oblongis, brevi-acuminatis, basi rotundatis vel subacutis, superne ad costam (cum *petiolo* longiusculo) canaliculatis; *pedicellis* petiolo æquilongis vel dimidio longioribus.—c.p. 3473 (3436).

HAB. Cultura District.

*Arbor* mediocris. *Folia* 3-11 poll. longa,  $1\frac{1}{4}$ - $4\frac{1}{2}$  poll. lata, *petiolo* 1- $1\frac{3}{4}$  poll. longo. *Fasciculi* 4-6-flori, adpresse fulvo-pilosi. *Calyx* 4 lin. longus. *Ovarium* 6-loculare. *Nuculania* ovoidea.

4. **I. pauciflora**, Thw.; *foliis* glabris, oblongis, brevi-acuminatis, basi acutis, superne ad costam (cum *petiolo* longiusculo) canaliculatis; *fasciculis* ad apices ramulorum paucis, 2-3-floris; *pedicellis* brevissimis, ovario 3-4-loculari; *nuculanis* oblongis.—c.p. 2680.

HAB. Near Ratnapoora, at no great elevation.

*Arbor* mediocris. *Folia* 3-8 poll. longa, 1-3 poll. lata, *petiolo*  $\frac{1}{2}$ -1 poll. longo. *Calyx*  $1\frac{1}{2}$  lin. longus.

5. **I. lævifolia**, Thw.; *ramulis* cum *petiolis* *fasciculisque* adpresse fusco-pilosis; *foliis* glabris, utrinque lævibus nitidisque, oblongo-lanceolatis, obtusis, basi in petiolum brevem, superne convexum, angustatis; *pedicellis* petiolo æquilongis, vel parum longioribus; *nuculanis* oblongis, acutis.—c.p. 2832.

HAB. Saffragam District and Reigam Corle, at no great elevation.

*Arbor* mediocris. *Folia* 2-6 poll. longa,  $\frac{3}{4}$ -2 poll. lata, *petiolo* 2-3 lin. longo. *Fasciculi* 4-6-flori, axillares, ad apicem ramulorum solitarii vel bini. *Calyx*  $1\frac{1}{2}$  lin. longus. *Antheræ* apice pilosæ. *Ovarium* 6-loculare.

### Sect. 2. *Calyx* 4-partitus.

6. **I. Wightiana**, A. DC. l. c. p. 187, cum syn.—*I. Perottetiana*, A. DC. l. c. p. 188; Wight, Ic. t. 1219. *I. Candolleana*, Wight, Ic. t. 1220. An *I. lanceolata*, Wight, Ic. t. 359?—c.p. 45 (47, 192, 2411).

Var.  $\beta$ . *angustata*; *foliis* lanceolatis, ab infra-medio versus apicem angustatis, subtus (ut in var.  $\alpha$ ) delicatule subparallelim venulosis.—c.p. 179 (191).

Var.  $\gamma$ . *montana*; *foliis* subcoriaceis, obovatis, apice rotundatis vel obscure brevi-acuminatis, subtus plus prominentim parceque venosis quam in varr.  $\alpha$  et  $\beta$ .—c.p. 3092.

HAB. Var.  $\alpha$ . Extremely abundant at an elevation of 2000 to 5000



feet. Var.  $\beta$ . Near Galagama, *Gardner*. Var.  $\gamma$ . Newera Ellia. Nom. vulg. "Kiriwarala-gass."

*Arbor* mediocris. *Folia* glabra (juniora plus minus pilosa), lanceolata, obovato-lanceolata, obovata vel suborbicularia, plus minus abrupte brevique acuminata, 1-5 poll. longa, 4 lin.  $2\frac{1}{4}$  poll. lata, *petiolo* 2-4 lin. longo. *Pedicelli* brevissimi vel subnulli. *Ovarium* sæpissime 4-loculare. *Nuculania* oblonga, 1-2-sperma, coccinea, 6 lin. longa. *Semina* castanea, nitida, albuminosa.

I was at one time disposed to regard some of the forms of this most variable plant as distinct, but further observation and the examination of a large number of specimens, satisfy me that they all belong to one species. I have little doubt, too, that Dr. Wight's *Is. lanceolata* is a form of the present plant, for I have examples of it extremely like his figure.

## XCVI. EBENACEÆ.

### 1. DIOSPYROS, Dal., Linn.

† *Flores pedunculati.*

‡ FLOB. MASC. *Corolla breviuscula, campanulata vel urceolata; staminibus inter se subæquilongis.*

1. **D. Embryopteris**, Pers. (A. DC. Prod. viii. p. 235, cum syn.; Wight, Icon. tt. 843, 844.)—C.P. 1915.

Var.  $\beta$ . *atrata; foliis membranaceis, gemmis, pedunculis calyceque nigro-pilosis.*—C.P. 2731.

Var.  $\gamma$ . *nervosa; foliis brevioribus, coriaceis, utrinque valde prominentim venosis, basi rotundatis; gemmis, pedunculis calyceque nigro-pilosis; lobis calycis fructiferi erectis.*—C.P. 1910.

HAB. Var.  $\alpha$ . Very abundant in the hotter parts of the island. Var.  $\beta$ . Less common, but generally distributed. Var.  $\gamma$ . In damp forests towards the south of the island. Nom. vulg. "Timberee-gass."

Though disposed at one time to regard the three well-marked varieties of this plant as distinct species, I now find it impossible, after a most careful examination, to discover any distinctive characters of sufficient importance to justify my separating them otherwise than as varieties. The black colour of the pubescence upon the gemmæ and inflorescence of var.  $\beta$  and  $\gamma$ , gives them a different aspect from that of var.  $\alpha$ , but black hairs are frequently intermixed with the pale pubescence of the latter, especially upon the gemmæ. In all the specimens I have examined, the ovary has been 8-celled.

The timber of this tree is used for buildings: the juice of the unripe fruit is employed for paying the seams of fishing-boats, and for putting upon their cordage and lines.

2. **D. cordifolia**, Roxb. (Pl. Cor. i. p. 38. t. 50; A. DC. l. c. p. 230, cum syn.)—*D. montana*, Wight, Ic. t. 1225 (non Roxb.).—C.P. 1909.

HAB. Jaffna, *Gardner*.

I have not seen the female flowers of this species, so am unable to ascertain whether they have four sterile stamens, as shown in Dr. Wight's figure, or a greater number, as represented in Roxburgh's; but it is probable, judging from other species, that the number of these organs is not very constant. The albumen of the seeds is not ruminant.

3. **D. sylvatica**, Roxb. (Roxb. Pl. Cor. i. p. 38. t. 47; A. DC. l. c. p. 231, cum syn.)—C.P. 2729.

HAB. Damp forests in the Hantani District and near Ratnapoora, up

to an elevation of 4000 feet. Nom. vulg. "Soodoo-Kadoombaireya-gass."

Tota planta siccitate nigrescens. Ovarium 6- vel 8-loculare. Nuculania sphaerica, 8 lin. in diam. Semina oblonga, 6 lin. longa, 4 lin. lata, albumine ruminato.

4. **D. Toposia**, Ham. (A. DC. l. c. p. 237, cum syn.)—*D. racemosa*, Roxb. Fl. Ind. ii. p. 536; Wight, Ic. t. 416.—C.P. 1911.

HAB. Not uncommon in damp forests, up to an elevation of 4000 feet. Nom. vulg. "Kahakaala-gass."

This agrees so well with Hamilton's description of his plant, except that he mentions the corolla as being five-lobed, that I cannot doubt their identity, and it certainly must be the one represented in Dr. Wight's figure, copied from Roxburgh's drawing. The inflorescence, however, is by no means racemose, but distributed over the lower leafless portion of the ramulus, and in the axils of a few of its lowest leaves. The female flowers are solitary, and the males in three-flowered cymes. In species of this genus, where the inflorescence is more apparently racemose, as in *D. Embryopteris*, it is not truly so, for the terminal flower being the first to expand, the inflorescence is really a modified cyme.

‡‡ FLOR. MASC. *Corolla longiuscula, hypocrateriformi; staminibus inter se valde inaequilongis.*

5. **D. crumenata**, Thw.; *foliis glabris, oblongis, abrupte acuminatis, obtusis, basi rotundatis vel parum angustatis.* MASC. *Cymis 3-5-floris, cernuis, pilosis, petiolo subaequilongis; calyce obscure 4-dentato.* FLOR. FÆM. *solitariis, majusculis, cernuis; calyce utrinque piloso, ad medium 4-lobo, inter lobos marsupio-dilatato; lobis rotundatis, apiculatis; ovario 8-loculari; nuculanis subsphaericis 6-8-spermis.*—C.P. 2438.

HAB. Central Province, at an elevation of 2000 to 4000 feet.

*Arbor ingens. Ramuli glabri. Folia 2-5 poll. longa, 1-2 poll. lata, siccitate fuscescentia, petiolo 3-4 lin. longo.* FLOR. MASC. *Calyx 2 lin. longus. Corolla 5 lin. longa. Stamina circiter 12.* FLOR. FÆM. *Calyx 4 lin. longus. Corolla calyce parum longior. Stamina ster. circiter 12. Nuculania 1½-2 poll. in diam., demum glabra, calyce aucto tetragono suffulta. Semina nigra, nitida, 1 poll. longa, ½ poll. lata, albumine ruminato.*

6. **D. affinis**, Thw.; *foliis glabris, subcoriaceis, prominentim reticulato-venosis, oblongis, plus minus acuminatis, obtusis, basi acutis vel subrotundatis.* MASC. *Cymis 3-7-floris, cernuis, pilosulis, petiolo parum longioribus; calyce breve 4-dentato.* FLOR. FÆM. *solitariis, majusculis, cernuis, pedunculo subaequilongis; calyce interne piloso, externe subglabro, ad medium 4-lobo, inter lobos marsupio-dilatato; lobis longiuscule acuminatis; ovario 6-loculari; nuculanis sphaericis, apiculatis, sæpissime 4-spermis.*—C.P. 2924.

HAB. Ooma Oya, on the lower road from Kandy to Badulla.

*Arbor mediocris. Ramuli subglabri. Gemmæ pilosæ. Folia 1¼-4½ poll. longa, 6-16 lin. lata, siccitate nigrescentia, petiolo 3-5 lin. longo.* FLOR. MASC. *Calyx 1½ lin. longus. Corolla 4 lin. longa. Stamina circiter 9.* FLOR. FÆM. *Calyx 5 lin. longus. Nuculania pollicaria, denique glabra. Semina 6-7 lin. longa, 4 lin. lata, albumine ruminato.*

Although very closely allied to the preceding species, I think it may be considered distinct. The timber of both species is suitable for building purposes.

7. **D. quæsitæ**, Thw.; *foliis glabris, oblongis, abrupte acuminatis, obtusis, basi parum angustatis, prominentim reticulato-venulosis, superne ad costam cum petiolo canaliculatis.* MASC. *Cymis 3-7-floris, cernuis, pilosis, petiolo subaequilongis; calyce breve 5-dentato.* FLOR.

FÆM. solitariis. calyce ad medium 5-lobo, inter lobos marsupio-dilatato; lobis acutiusculis; nuculanis subsphæricis.—c.p. 3010.

HAB. Singherajah and other forests between Ratnapoora and Galle. Nom. vulg. "Kaloomidereya-gass."

*Arbor* ingens. *Ramuli* glabri. *Folia* 3-7 poll. longa,  $1\frac{1}{2}$ -3 poll. lata, siccitate fusco-nigrescentia, *petiolo* 3-7 lin. longo. *Nuculania* (immatura) bipollicaria.

This species produces the most valuable of the timber known as Calamander-wood, so much esteemed for ornamental cabinet-work. I regret that my specimens are too imperfect to allow of my giving so full a description as would be desirable of this interesting plant, which is nearly allied to *D. crumenata*, but its larger leaves and fruit, and its pentamerous flowers, well distinguish it.

8. **D. Ebenum**, Retz, (A. DC. l. c. p. 234, cum syn.; Wight, Icon. t. 188.)—c.p. 1912 (1913, 2437, 2439).

HAB. Not uncommon up to an elevation of 5000 feet. Nom. vulg. "Kaloowara-gass."

*Arbor* ingens. *Folia* membranacea vel parum coriacea, brevi-acuminata, rarius retusa, 2-7 poll. longa,  $\frac{3}{4}$ - $2\frac{1}{2}$  poll. lata, *petiolo* 2-4 lin. longo. *Ovarium* 8-loculare. *Semina* albumine non ruminato.

This tree yields the best kind of Ebony-wood.

Dr. Wight's figure, copied from Roxburgh's drawing, is not characteristic of this species as it commonly occurs here, the lobes of the corolla being represented too large, and the leaves without the usual somewhat abrupt obtuse acumination.

9. **D. oocarpa**, Thw.; *foliis* glabris, membranaceis, ovatis vel ellipticis, obtuse acuminatis, basi rotundatis vel parum angustatis, inconspicue venosis. INFLO. MASC. racemoidea, pilosula; *pedunculo* 3-6-floro, *petiolum* æquante vel brevior; *pedicellis* brevissimis; *calyce* sub-integro vel obscure 3-4-dentato. FLOR. FÆM. 1-3-nis; *ovario* 6- vel 8-loculari; *nuculanis* ovoideis, rugulosis, sæpissime 6-spermis, calyce vix aucto suffultis.—c.p. 1914.

HAB. Kornegalle District, and at Haragam, near Kandy, at no great elevation. Nom. vulg. "Kalo-kadoombaireya-gass."

*Arbor* mediocris. *Ramuli* juniores minute pilosuli. *Folia* 2-4 poll. longa,  $1\frac{1}{2}$ - $2\frac{1}{2}$  poll. lata, *petiolo* 3-4 lin. longo. FLOR. MASC. *Calyx* 2 lin. longus. *Corolla* 5 lin. longa. *Stamina* circiter 12. FLOR. FÆM. quam masc. vix majores. *Nuculania* (immatura) pollicaria. *Semina* albumine non ruminato.

#### †† Flores sessiles vel subsessiles.

‡ FLOR. MASC. *Staminibus* indefinitis, inter se valde inæquilongis.

10. **D. insignis**, Thw.; *foliis* suboppositis vel alternis, majusculis, membranaceis, glabris, oblongis, acuminatis, parallelim delicatule venulosis, basi angustatis vel rotundatis. MASC. *Capitulis* multifloris; *calyce* ad medium 4-partito; lobis acutis. FLOR. FÆM. 1-2-nis; *ovario* 8-loculari; *nuculanis* subsphæricis, minute depresso-areolatis, calyce aucto, crasso, lignoso et tota margine quasi late reflexa suffultis.—c.p. 2730, 3477.

HAB. Damp forests, up to an elevation of 2000 feet. Nom. vulg. "Gona-gass."

*Arbor* ingens. *Ramuli* juniores pilosuli. *Folia* 4-14 poll. longa,  $1\frac{1}{2}$ -6 poll. lata, siccitate fuscescentia, *petiolo* 3-5 lin. longo. FLOR. MASC. *Calyx*  $1\frac{1}{2}$  lin. longus. *Corolla* 4 lin. longa. *Stamina* circiter 20. FLOR. FÆM. *Nuculania* sesquipollicaria. *Semina* 11 lin. longa, 6 lin. lata, albumine ruminato.

A valuable timber-tree.

11. **D. oppositifolia**, Thw.; *foliis* oppositis, glabris (junioribus pilosis), firmis, ovatis vel ellipticis, abrupte brevi-acuminatis, obtusis, basi rotundatis, obscure parallelim venulosis, brevi-petiolatis. — C.P. 3011.

HAB. Hinidoon Corle, up to an elevation of 1000 feet. Nom. locale “Kaloomidereya-gass.”

*Arbor* mediocris. *Folia* 2–6 poll. longa,  $1\frac{1}{2}$ –3 poll. lata, siccitate fusciscentia, *petiolo* 1–2 lin. longo. MASC. *Capitula* (in meis specim.) pauciflora. *Calyx* 1 lin. longus, fere ad medium 4-lobus, *lobis* acutis. *Corolla* tenuis, circiter 3 lin. longa. *Stamina* circiter 8. FLOR. FÆM. adhuc ignoti.

The timber of this tree very much resembles the true Calamander-wood.

12. **D. Gardneri**, Thw.; *foliis* membranaceis, glabris, nitidis, inconspicue venosis, oblongis, acuminatis, basi parum angustatis, petiolatis; *floribus* brevissime pedicellatis, pilosis; *calyce* ad medium 4-partito; *lobis* triangularibus, acutis; *corollæ* tubo parum inflato; *ovario* 8-loculari; *nuculanis* depresso-sphæricis, apiculatis, fulvo-pilosis, denique glabris, nitidis, ad dimidium inferius calyce valde aucto obtectis.—C.P. 1908.

HAB. Saffragam and Kornegalle Districts, and, less commonly, near Kandy. Nom. vulg. “Kadoombaireya-gass.”

*Arbor* mediocris. *Ramuli* teretes, glabri. *Gemmæ* fusco-pilosæ. *Folia* 3–7 poll. longa,  $1\frac{1}{2}$ –2 poll. lata, siccitate nigrescentia; *petiolo* 3–4 lin. longo. FLOR. MASC. 1–4-ni. *Calyx* 2 lin. longus. *Corolla* 5 lin. longa. *Stamina* circiter 12, filamentis pilosis. FLOR. FÆM. solitariis, erectis. *Calyx*  $2\frac{1}{2}$  lin. longus, *lobis* recurvatis, margine reflexis. *Nuculania* (immatura) pollicaria.

The timber of this tree is valuable for building and for cabinet-work.

‡‡ FLOR. MASC. *Staminibus* indefinitis, inter se subæquilongis.

13. **D. ovalifolia**, Wight, Ic. t. 1227.—C.P. 1815, 2533.

HAB. Jaffna, Gardner. Central Province, at Hewahette and below Hapootelle, at an elevation of 2000 to 4000 feet.

*Arbor* mediocris. *Folia*  $1\frac{1}{2}$ –6 poll. longa,  $\frac{1}{2}$ –2 poll. lata, siccitate fulvescentia, *petiolo* 2–3 lin. longo. *Flores* 4- vel 5-meri. MASC. *Stamina* 15–20. *Ovarii* rudimentum parvum, conicum. FÆM. *Capitula* 3–6-flora. *Stamina* sterilia 1–7. *Ovarium* 4- vel 6-loculare. *Stigma* 2–3-lobum, *lobis* reniformibus. *Nuculania* glabra, sphærica, glauca, 8 lin. in diam., sæpissime 1-sperma. *Semina* subsphærica, albumine non ruminato.

14. **D. Candolleana**, Wight, Ic. tt. 1221, 1222.—C.P. 3394.

HAB. Saffragam District and Hinidoon Corle. Nom. vulg. “Homedereya-gass.”

*Arbor* mediocris, *ramulis*, *foliis* junioribus inflorescentiaque ferrugineo-tomentosis. *Folia* sæpius longe acuminata, siccitate fulvescentia. MASC. *Stamina* 10–12. FÆM. *Nuculania* subsphærica, ferrugineo-pilosa, denique glabra, 7–8 lin. in diam., sæpissime 4-sperma. *Semina* oblonga, nigrescentia, transverse striata, 5 lin. longa,  $2\frac{1}{4}$  lin. lata, albumine ruminato.

‡‡‡ *Staminibus* tot quot lobi corollæ.

15. **D. hirsuta**, Linn. f. (Supp. p. 440; A. DC. l. c. p. 223.)—C.P. 382.

HAB. Forests in the Saffragam and Galle Districts.

*Arbor* mediocris, *ramulis*, *foliis* junioribus inflorescentiaque ferrugineo-tomen-

tos. *Folia* subcoriacea, elliptica vel oblonga, plus minus abrupte acuminata, basi parum angustata, subtus tomentosa, demum subglabra, siccitate rufescentia, 3–8 poll. longa, 1–3 poll. lata, *petiolo* tomentoso, 3–4 lin. longo. MASC. *Capitula* multiflora. *Calyx* 1 lin. longus, vix ad medium 5-partitus. *Corolla* circiter 2 lin. longa. *Ovar. rudim.* parvum. FLOR. FÆM. 1–3-ni. *Calyx* crassus, 3 lin. longus, ultra medium 5-partitus; *lobis* triangularibus, acutis, margine reflexis. *Corolla* calyce parum longior. *Ovarium* sæpissime 10-loculare. *Nuculania* ovoidea, apiculata, fulvo-tomentosa, 9 lin. longa, 6–10-sperma, calyce aucto suffulta. *Semina* nigra, transverse striata, 7 lin. longa, 3 lin. lata, *albumine* ruminato.

Female flowers are occasionally intermixed in the male capitula, and in that case are much smaller than when occurring alone. I have not yet been able to ascertain whether these smaller female flowers produce perfect fruit, but one which I dissected had the ovary only 6-celled.

16. **D. Moonii**, Thw.; *foliis* coriaceis, oblongis, acuminatis, basi rotundatis, superne glabris, costa petioloque canaliculatis, subtus sparse pilosulis; *calyce* fere ad medium 5-partito; *lobis* triangularibus, acutis; *ovario* 10-loculari; *nuculaniis* sphaericis, subglabris.—C.P. 2833.

HAB. Near Caltura, Moon; Pasdoon Corle.

*Arbor* mediocris. *Ramuli* teretes, pilosuli, cito glabri, cortice fisso, nigro-cinerascente. *Folia* 5–12 poll. longa, 1½–4 poll. lata, *petiolo* 3 lin. longo. *Inflorescentia* axillaris, et quoque in axillis foliorum delapsorum, fulvo-pilosa. MASC. *Capitula* 6–10-flora. *Calyx* 2 lin. longus. *Corolla* circiter 4 lin. longa. FLOR. FÆM. 1–2-ni. *Calyx* crassus, 3 lin. longus, margine revolutus. *Corolla* calyce subæquilonga. *Stigma* obscure 5-lobum. *Nuculania* sesquipollicaria, 6–10-sperma, calyce aucto suffulta. *Semina* oblonga, compressa, transverse rugosa, nigrescentia, 7 lin. longa, 4 lin. lata, *albumine* ruminato.

17. **D. acuta**, Thw.; *foliis* coriaceis, glabris vel subtus sparsissime pilosis, oblongo-lanceolatis, acutis, basi rotundatis, inconspicue venosis, superne costa petioloque canaliculatis; *calyce* ultra medium 4–5-partito; *lobis* lanceolatis, acutis; *ovario* 4- vel 6-loculari; *nuculaniis* acuminatis, sæpissime 2–3-spermis.—C.P. 3476.

HAB. Pasdoon Corle.

*Arbor* mediocris. *Ramuli* teretes, pilosuli, cito glabri. *Gemmæ* adpresse fulvo-pilosæ. *Folia* 5–12 poll. longa, 1–3 poll. lata, siccitate subtus rufescentia, *petiolo* ½–1 poll. longo. *Inflorescentia* adpresse fulvo-pilosa, aliquando monoica et tunc capitulis fœmineis versus apicem ramulorum, masculis infra. MASC. *Capitula* multiflora. *Calyx* 2 lin. longus. *Corolla* circiter 4 lin. longa. *Ovarii rudiment.* minutissimum. FLOR. FÆM. 1–4-ni. *Calyx* 4–5 lin. longus, *lobis* ad marginem plus minus reflexis. *Corolla* calyce subæquilonga. *Stigmata* 2–3, 1 lin. longa, spathuliformia. *Nuculania* sesquipollicaria, calyce parum aucto suffulta. *Semina* castanea, nitida, oblonga, acuminata, pollicaria, *albumine* non ruminato.

18. **D. attenuata**, Thw.; *foliis* membranaceis, glabris vel sparsissime pilosulis, oblongis, acuminatis, basi in petiolum brevem angustatis, creberrime venulosis; *calyce* ultra medium 4–5-partito; *lobis* lanceolatis, acutis; *ovario* 4-loculari; *nuculaniis* oblongis, acuminatis, sæpissime 2–3-spermis.—C.P. 3478.

HAB. Pasdoon Corle.

*Arbor* mediocris. *Folia* 1½–4 poll. longa, ½–1 poll. lata, siccitate pallide fusca, *petiolo* 1–2 lin. longo. *Inflorescentia* pilis testaceis, nigris intermixtis, strigosa. MASC. *Capitula* 3–10-flora. *Calyx* 1 lin. longus. *Corolla* circiter 2 lin. longa. FLOR. FÆM. solitarii. *Calyx* 3–4 lin. longus, *lobis* ad marginem plus minus reflexis. *Corolla* calyce parum longior. *Stigmata* 2, brevia, spathuliformia. *Nuculania* 1–1½ poll. longa. *Semina* castanea, nitida, oblonga, acuminata, *albumine* non ruminato.

2. **MACREIGHTIA**, A. DC.

1. **M. oblongifolia**, Thw.; *foliis* subcoriaceis, oblongis, acuminatis, basi rotundatis, subtus ad nervos cum petiolo brevi fulvo-hispidis, denique glabris. FLOR. FÆM. solitariis, brevi-pedunculatis; *calyce* hispido, ad basin 3-partito; *lobis* acutis; *ovario* piloso, 6-loculari; *nuculanis* subsphæricis, fulvo-tomentosis, 3-6-spermis.—C.P. 3396.

HAB. Near Ratnapoora, and in the Singherajah and other forests between that place and Galle.

*Arbor* parva. *Ramuli* juniores valde hispidi. *Folia* 3-7 poll. longa, 1-2½ poll. lata, *petiolo* 1-2 lin. longo. *Nuculanis* pollicaria, *calyce* non aucto suffulta. *Semina* 6 lin. longa, 4 lin. lata, nigrescentia, *radicula* cylindrica, *cotyledonibus* oblongis dimidio longiore, *albumine* non ruminato.

I have as yet only obtained specimens of this plant bearing ripe and unripe fruit, so am not quite certain that it is a true *Macreightia*, but it agrees with this genus in the characters my specimens exhibit.

3. **MABA**, Forst.

1. **M. buxifolia**, Pers. (A. DC. Prod. viii. p. 240, cum syn.; Wight, Ic. t. 763.)—C.P. 477.

Var.  $\beta$ . *microphylla*; *foliis* parvulis.—C.P. 1916.

Var.  $\gamma$ . *Ebenus*; *foliis* majoribus, membranaceis, parum acuminatis vel retusis, sæpe suborbiculatis.—*M. Ebenus*, Spr.; A. DC. l. c. p. 242, cum syn. *M. Neilgherrensis*, Wight, Ic. tt. 1228, 1229.—C.P. 3395.

Var.  $\delta$ . *angustifolia*; *foliis* lanceolatis vel lineari-lanceolatis, obtusis.—*M. angustifolia*, Miq.—C.P. 1917.

HAB. Varr.  $\alpha$ . and  $\beta$ . Hot, drier parts of the island. Var.  $\gamma$ . Ambagamowa District, and near Ratnapoora. Var.  $\delta$ . On the banks of rivers. Nom. vulg. "Kaloc-habaraleya-gass."

I have devoted much time to the examination of the several very different-looking varieties of this plant, expecting to discover some sufficiently important constant characters to enable me to separate them specifically, but I find them so completely connected together by intermediate forms, that I have no hesitation in considering them all as representing only one very variable species: variable, it may truly be called, since the leaves in var.  $\beta$  are sometimes not a quarter of an inch in length, whilst in var.  $\delta$  they reach to five inches in length.

## XCVII. AQUIFOLIACEÆ.

1. **ILEX**, Linn.

† *Flores tetrameri.*

1. **I. denticulata**, Wall. (Wight, Illust. t. 142.)—C.P. 573.

HAB. Bopatalawa Plains, at an elevation of 6000 feet.

*Arbor* magna. *Ramuli* angulosi, minutissime puberuli, cito glabri, cinerascetes. *Folia* subcoriacea, glabra, oblonga, abrupte rostrato-acuminata, acute serrata, 1½-3½ poll. longa, ¾-1½ poll. lata, siccitate fusciscentia, *petiolo* 2-3 lin. longo. *Pedunculi* subnulli vel plus minus elongati. *Pedicelli* 3-4 lin. longi, umbellati vel sparsi. *Corolla* 3½-4 lin. in exp.

†† *Flores pentameri.*

2. **I. Wightiana**, Wall. (Wight, Ic. t. 1216.)—C.P. 2656, 3484.

HAB. Southern and central parts of the island, up to an elevation of 4000 feet.

*Folia* 2-5 poll. longa, 1-2½ poll. lata, *petiolo* 3-4 lin. longo. *Pedunculi* sæpe elongati ramosique. *Pedicelli* umbellati, minutissime puberuli.

3. **I. Walkeri**, Wight et Gardn. MSS.; *foliis* parvis, coriaceis, glabris, nitidis, integris vel rarius parce denticulatis, oblongis vel rotundatis, acutis vel retusis mucronulatisque, basi plus minus angustatis; *umbellis* sessilibus vel brevi-pedunculatis. — An *I. integra*, Thunb. Fl. Jap. p. 77?—c.p. 298 (622, 647).

HAB. Central Province, at an elevation of 5000 to 8000 feet.

*Arbor* parva. *Ramuli* angulosi, dense foliosi. *Folia* 4-12 lin. longa, 2-7 lin. lata, *petiolo* 1½-2 lin. longo, parum alato. *Pedicelli* circiter 1½ lin. longi. *Flores* parvuli, albid.

## XCVIII. SYMPLOCEÆ.

### 1. SYMPLOCOS, Jacq.

Sect. *Ciponima*, A. DC.

*Ovarium* 2-loculare.

1. **S. pendula**, Wight, Ic. t. 1237; Walp. Ann. i. p. 500.—c.p. 504 (555, 665).

HAB. Central Province, at an elevation of 5000 to 8000 feet; abundant.

Sect. *Hopea*, A. DC.

† *Folia* secus ramulum fere ad basin disposita.

2. **S. spicata**, Roxb. (A. DC. Prod. viii. p. 254, cum syn.; Wight, Illust. ii. t. 150, et var.  $\beta$  *Zeylanica*, A. DC. l. c. cum syn.)—c.p. 473.

HAB. Very abundant. Var. *a* in more elevated situations, up to 7000 feet. Var.  $\beta$  in the hotter, lower parts of the island. Nom. vulg. "Bõmboo-gass."

3. **S. rufescens**, Thw.; arborea, glaberrima; *foliis* oblongo-lanceolatis, brevi-acuminatis, obtusis, inconspicue venosis, a medio ad apicem obscure crenulatis denticulatisque, basi acutis; *racemis* simplicibus, folio multo brevioribus; *floribus* inferioribus pedicellatis; *bracteis* *bracteolis*que parvis, glabris, margine ciliatis.—c.p. 3639 (473 partim).

HAB. Ambagamowa District.

*Ramuli* angulosi. *Folia* 1½-4 poll. longa, ¾-1½ poll. lata, siccitate rufescentia; *petiolo* 2-3 lin. longo. *Bracteæ* rotundatæ, cito deciduæ. *Bracteolæ* oblongæ, calyce dimidio breviores, subpersistentes. *Calyx* glaber, lobis rotundatis, parum acutis, margine ciliatis.

Closely resembling var.  $\beta$  of the preceding species, but apparently quite distinct. I have not seen the fruit of the present plant.

4. **S. læta**, Thw.; arborea, glaberrima; *foliis* ovato-lanceolatis, acuminatis, obscure denticulatis, basi acutis, subtus parce prominentim venosis; *racemis* folio subæquilongis vel brevioribus, basin versus nudis; *floribus* sessilibus vel inferioribus subsessilibus; *bracteis* glabris, parvis, oblongis; *bracteolis* nullis; *baccis* oblongis.—c.p. 165 (582, 2436).

HAB. Not uncommon in the more elevated parts of the Central Province.

*Ramuli* teretes, striati. *Folia*  $1\frac{1}{2}$ -4 poll. longa, 6-16 lin. lata, juniora cyaneo-viridia, denique flavescentia; *petiolo* 3-8 lin. longo. *Calyx* glaber, tubo obconico; *lobis* rotundatis, subacutis, ciliolatis.

5. **S. bractealis**, Thw.; arborea, glaberrima; *foliis* ovatis vel ovato-lanceolatis, acutis, glanduloso-serratis, creberrime venosis, basi obtusis vel subacutis, brevi-petiolatis; *racemis* folio brevioribus; *floribus* subsessilibus; *bracteis* *bracteolisque* majusculis, glabris, margine ciliatis, deciduis; *baccis* oblongis.—C.P. 174 (676).

HAB. Forests in the more elevated parts of the Central Province.

*Ramuli* teretes, striati. *Folia*  $1\frac{1}{2}$ -4 poll. longa, 1-2 poll. lata; *petiolo* 1-2 $\frac{1}{2}$  lin. longo. *Bracteæ* rotundatæ, 2-3 $\frac{1}{2}$  lin. longæ, *bracteolis* oblongis majores. *Calyx* glaber, *lobis* rotundatis.

6. **S. obtusa**, Wall. (A. DC. l. c. p. 255; Wight, Ic. t. 1233).—C.P. 626 (673).

Var.  $\beta$ . *major*; *foliis* majoribus, brevissime acuminatis, rotundatis vel retusis; *floribus* parum minoribus.—C.P. 1820.

Var.  $\gamma$ . *obovata*; *foliis* minoribus, obovatis vel suborbicularibus, basi acutis; *racemis* paucis sæpe 1-floris.—*S. obovata*, Wight & Gardn. MSS. C.P. 1819.

Var.  $\delta$ . *cucullata*; *foliis* majoribus, valde coriaceis, oblongis vel obovatis, utrinque obtusis, ad marginem cucullato-revolutis, superne prominentim, subtus inconspicue venosis; *petiolo* brevi, crasso; *racemis* ut in var.  $\alpha$ .—C.P. 2835.

HAB. Central Province, at an elevation of 5000 to 8000 feet.

A very variable plant, and perhaps var.  $\beta$ , in which the leaves frequently reach to seven inches in length and three inches in width, is to be considered the most typical form of the species. In var.  $\alpha$  the racemes sometimes become branchlike, the bracts being converted into leaves, and thus it approaches var.  $\gamma$ , in which the leaves of a branch often bear each a single stalked flower in its axil. The leaves in var.  $\gamma$  show not uncommonly a disposition to a verticillate arrangement. Var.  $\delta$  has, at first sight, a very distinct appearance, but the peculiar form of its leaves proves it to be in an abnormal state, arising probably from certain conditions of moisture and exposure.

7. **S. hirsuta**, Wight and Gardn. MSS.; arborea; *ramulis* plus minus hispidis; *foliis* membranaceis, creberrime venosis, oblongis, brevi-acuminatis, basi rotundatis, serrulatis, superne glabris, subtus ad costam *petioloque* hispidulis; *racemis* folio brevioribus, pilosis; *floribus* inferioribus brevi-pedicellatis, superioribus sessilibus; *bracteis* *bracteolisque* sæpius externe pilosis; *calyce* glabro vel rarius lobis externis pilosis; *baccis* oblongis.—C.P. 368.

Var.  $\beta$ . *minor*; *foliis* brevioribus, ovatis, acutis; *petiolo* longiore quam in var.  $\alpha$ .—C.P. 273, 2204.

HAB. Central Province, at an elevation of 5000 to 8000 feet.

*Folia* 1-3 $\frac{1}{2}$  poll. longa,  $\frac{1}{2}$ -1 $\frac{1}{2}$  poll. lata; *petiolo* 2 lin. (var.  $\beta$  3 lin.) longo. *Baccæ*  $\frac{1}{2}$  poll. longæ.

This species would seem to be very nearly allied to *S. Hamiltoniana*, Wall., A. DC. l. c. p. 254, and to *S. foliosa*, Wight, Ic. t. 1234.

8. **S. elegans**, Thw.; arborea; *ramulis* teretibus, plus minus rufo-vel testaceo-tomentosis; *foliis* membranaceis, creberrime venosis, oblongis vel ovatis, acute acuminatis, subintegris vel obscure serrulatis,



superne glabris vel basi ad costam pilosis, subtus *petioloque* plus minus pilosis, basi rotundatis vel subacutis; *racemis* folio brevioribus, pilosis; *bracteis bracteolisque* externe pilosis; *calyce* glabro, lobis ciliolatis; *baccis* oblongis, subampulliformibus.—C.P. 23.

Var.  $\beta$ . *angustata*; *ramulis* tomento fulvo parcius tectis; *foliis* angustioribus, longius acuminatis.—C.P. 78, 155.

HAB. Central Province, at an elevation of 5000 to 7000 feet.

*Folia* 1-3½ poll. longa, ½-1½ poll. lata; *petiolo* 1½-2 lin. longo. *Baccæ* 5 lin. longæ.

Closely allied to and resembling the preceding species, and I have sometimes felt disposed to regard them as varieties, but the different character of the pubescence, and the more acuminated leaves of the present plant, are perhaps sufficient to warrant their being kept distinct.

9. **S. hispidula**, Thw.; arborea; *ramulis* subteretibus, parum angulosis, plus minus hispidis; *foliis* membranaceis, creberrime venosis, integris vel obscure denticulatis, obovatis vel oblongis, abrupte brev-acuminatis, basi rotundatis, subcordatis vel rarius subacutis, superne pilosulis vel glabris, subtus *petioloque* pilosis; *racemis* folio brevioribus, hispidis; *floribus* brevi-pedicellatis; *bracteis, bracteolis, calyceque* externe pilosis; *baccis* subcylindricis, pilosis.—C.P. 407.

HAB. Ambagamowa and Saffragam Districts, at an elevation of 3000 to 5000 feet.

*Folia* 2½-7 poll. longa, 1½-3½ poll. lata; *petiolo* 2-4 lin. longo. *Baccæ* pilosæ, 5-6 lin. longæ.

The entire leaves, of a larger size and different shape, and the hairy calyx of the present plant, are the principal characters to distinguish it from *S. hirsuta*, W. & G.

10. **S. jucunda**, Thw.; arborea; *ramulis* teretibus, fulvo-tomentosis; *foliis* membranaceis vel subcoriaceis, integris vel obscure distanter denticulatis, longiuscule acuminatis, basi rotundatis, superne pilosulis vel glabris, subtus creberrime venosis et, cum *petiolo*, tomentosus; *racemis* folio brevioribus, cum *bracteis, bracteolis calyceque* externe, tomentosus; *floribus* brevi-pedicellatis; *baccis* subcylindricis, pilosis.—C.P. 2435, 3454.

HAB. Deltotte and Saffragam, at an elevation of 3000 to 4000 feet.

*Folia* 2½-6 poll. longa, 1-2 poll. lata; *petiolo* 2-3 lin. longo. *Baccæ* 6-7 lin. longæ.

Differs from the preceding in its more acuminated leaves and in the character of the pubescence. From *S. pulchra*, Wight, Ic. t. 1230, to which it bears a close resemblance, the entire leaves of the present plant seem sufficient to distinguish it.

11. **S. acuta**, Thw.; arborea; *foliis* subcoriaceis, oblongo-lanceolatis, acuminatis, utrinque acutis, superne glabris, lævibus, ad costam canaliculatis, subtus venosis, costa venisque primariis, cum *petiolo*, adpresse pilosulis; *racemis* *petiolo* subæquilongis vel 2-3-plo longioribus, puberulis; *pedicellis* *calyce* brevioribus; *calyce* tubo puberulo, lobis rotundatis, subglabris; *baccis* longiuscule sub-ampulliformibus, subglabris.—C.P. 2681.

HAB. Saffragam District, at an elevation of 2000 feet.

*Ramuli* teretes, juniores adpresse pilosi. *Folia* 2-5 poll. longa, ¾-1¼ poll. lata; *petiolo* 3-5 lin. longo. *Baccæ* 5-6 lin. longæ.

†† *Folia versus apicem ramuli disposita.*

12. **S. cuneata**, Thw.; arborea; *foliis* subcoriaceis, integris, oblongo-

lanceolatis, acutis, basi in petiolum brevem gradatim angustatis, superne glabris, lævibus, ad costam venasque primarias canaliculatis, subtus prominentim venosis, costa venisque, cum petiolo, adpresse pilosis; *racemis* brevibus, tomentosus; *floribus* subsessilibus; *baccis* elongatis; subcylindricis; cum *lobis calycinis*, pilosis.—C.P. 2920.

HAB. Pasdoon Corle, at no great elevation.

*Ramuli* teretes, juniores dense adpresse pilosi. *Folia* 3–6 poll. longa,  $\frac{3}{4}$ – $1\frac{1}{2}$  poll. lata; *petiolo* 2–3 lin. longo. *Baccæ* 6–7 lin. longæ.

The different arrangement of the leaves, as well as the subsessile flowers of the present species, sufficiently distinguish it from the preceding, to which it bears very considerable general resemblance.

13. **S. cordifolia**, Thw.; arborea; *foliis* glabris, subcoriaceis obovatis, obtusis vel brevissime acuminatis, versus basin angustatis, auriculato-cordatis, creberrime venosis, glanduloso-denticulatis, brevi-petiolatis; *racemis* tomentosus, folio brevioribus; *bracteis* *bracteolisque* majusculis, externe tomentosus; *calyce* tubo glabro, lobis externe tomentosus.—C.P. 271.

HAB. Central Province, at an elevation of 6000 to 7000 feet.

*Folia* 3–6 poll. longa,  $1\frac{3}{4}$ – $3\frac{1}{2}$  poll. lata; *petiolo* 1–2 lin. longo. *Baccæ* oblongæ, glabræ, calycis lobis tomentosus coronatæ.

14. **S. marginalis**, Thw.; arborea; *ramulis* teretibus et, cum *petiolis* *racemisque*, dense ferrugineo-tomentosis; *foliis* coriaceis, integris, oblongis, abrupte et brevissime acuminatis, basi subacutis, superne glabris, ad marginem parum incrassatum tomentosus, ad costam basi tomentosam canaliculatis, subtus creberrime prominenti-venosis tomentosisque; *calyce* tomentoso.—C.P. 2688.

HAB. Ambagamowa District.

*Folia*  $3\frac{1}{2}$ –7 poll. longa,  $1\frac{3}{4}$ – $3\frac{1}{2}$  lata; *petiolo* 3–5 lin. longo.

+++ *Folia ad summum apicem ramuli subverticillata.*

15. **S. apicalis**, Thw.; subarborea, erecta, parce ramosa; *ramulis* teretibus; *foliis* majusculis, membranaceis, subintegris, minute denticulatis, creberrime venosis, oblongo-lanceolatis, acutis; *racemis* folio multum brevioribus, axillaribus et sæpe in axillis squamarum ramuli delapsarum, hirsutis; *floribus* subsessilibus; *bracteis*, *bracteolis* *calyceque* externe pilosis; *baccis* oblongis, pilosis.

Var. *a. strigosa*; *ramulis* strigosis; *foliis* basi acutis vel subacutis, costa petioloque strigosis.—C.P. 438.

Var. *β. glabrifolia*; *ramulis* glabris; *foliis* majoribus, glabris, a supra-medio versus basin gradatim angustatis, basi abrupte obtusis.—C.P. 2933.

HAB. Var. *a.* Reigam Corle. Var. *β.* In the Singhe-rajah Forest, between Ratnapoora and Galle, in wet places, at no great elevation.

*Folia* 5–15 poll. longa,  $1\frac{1}{2}$ – $5\frac{1}{2}$  poll. lata; *petiolo* 2–3 lin. longo. *Baccæ* 6–7 lin. longæ.

16. **S. coronata**, Thw.; arborea; *ramulis* teretibus, ferrugineo-tomentosis; *foliis* coriaceis, oblongis, utrinque acutis, serrulatis, creberrime venosis, superne glabris, subtus petioloque tomentosus; *racemis* in axillis squamarum ramuli delapsarum, tomentosus, folio multum brevioribus; *floribus* subconfertis, sessilibus; *bracteis*, *bracteolis* *calyceque* externe pilosis; *baccis* ovoideis, pilosis.—C.P. 127.

HAB. Central Province, at an elevation of 2000 to 4000 feet.  
*Folia* 5-9 poll. longa, 2-4 poll. lata; *petiolo* 2-4 lin. longo. *Baccæ* 4-5 lin. longæ.

### XCIX. ANCISTROCLADEÆ.

(Planchon, Ann. des Sc. Nat. (Ser. 3) xiii. p. 316.—Walp. Ann. ii. p. 175.)

#### 1. ANCISTROCLADUS, Wall.

(Thw. in Linn. Soc. Trans. xxi. p. 225.)

*Semen* copiose albuminosum, cerebriformi-ruminatum. *Embryonis radícula* cylindrica, *cotyledones* breves, subfoliacei, divergentes.

1. **A. Vahlîi**, Arn. Pug. 7 (325); Planch. l. c. p. 317; Walp. l. c.; Thw. l. c. t. 24.—C.P. 1600.

HAB. Central and southern parts of the island, up to an elevation of 2000 feet. Nom. vulg. "Gona-wel," "Gonapittan-wel."

### C. OLEACEÆ.

#### 1. OLEA, Tourn.

1. **O. glandulifera**, Wall. (A. DC. Prod. viii. p. 285; Wight, Ic. t. 1238.)—C.P. 2481.

HAB. Central Province, at an elevation of 2000 to 4000 feet.

2. **O. Gardneri**, Thw.; glabra; *foliis* oppositis, versus apicem ramuli aggregatis, coriaceis, integris, ovatis vel oblongis, plus minus acuminatis vel rotundatis, parce venosis, basi acutis; *cymis* pedunculatis, 3-9-floris, rarius plurifloris, axillaribus et infra folia secus ramulum sparsis; *pedicellis* flore subæquilongis; *corolla* campanulata, ad medium 4-partita, lobis rotundatis.—*Notelæa Zeylanica*, Gardn. MSS.—C.P. 275, (627, 1821.)

HAB. Central Province, at an elevation of 5000 to 8000 feet.

*Arbor* 10-12-pedalis. *Folia* 1-2½ poll. longa, ½-1½ poll. lata; *petiolo* 2-3 lin. longo. *Calyx* acute 4-lobus. *Corolla* albida, 1 lin. longa. *Stamina* inclusa. *Nuculaniam* rotundata, pisi magnitudine, putamine tenui.

#### 2. LIGUSTRUM, Tourn.

1. **L. robustum**, Bl. Mus. Bot. Lugd.-Bat. i. p. 313.—*Visania robusta*, A. DC. Prod. viii. p. 289, cum syn.—C.P. 475.

HAB. Central Province, up to an elevation of 5000 feet; common in the Hapootelle District.

#### 3. CHIONANTHUS, Linn.

(Blume, Mus. Bot. Lugd.-Bat. i. p. 317.)

1. **C. Zeylanica**, Willd. i. p. 47. *Linociera purpurea*, Vahl; A. DC. Prod. viii. p. 297, cum syn. *L. dichotoma*, Wall. A. DC. l. c.—C.P. 1818.

HAB. Abundant in the hotter parts of the island. Nom. vulg. "Gherriatta-gass."

*Corolla* alba, siccitate subpurpurascens.

As remarked by Blume, there seems to be scarcely sufficient ground for separating *Linociera* as a genus distinct from *Chionanthus*. Of *Chion. cotinifolia*, Willd., founded on Plukenet's description, quoted by Burmann in *Thes. Zeyl.* p. 31, I know nothing, nor am I acquainted with any species of this genus, indigenous to Ceylon, which has the under side of its leaves villous.

2. **C. albidiflora**, Thw.; *foliis* subcoriaceis, oblongis, abrupte rostrato-acuminatis, paucissime venosis, basi acutis; *cymis* axillaribus, subracemoideis, paucifloris, folio multum brevioribus, glabris; *bracteis* minutissime pilosulis; *floribus* brevi-pedicellatis; *calyce* glabro, lobis rotundatis, subacutis; *petalis* elongatulis, canaliculato-linearibus; *nuculanis* obovoideis, apiculatis, glabris, putamine crustaceo.—C.P. 472.

HAB. Not uncommon at an elevation of 1000 to 3000 feet. Nom. vulg. "Taccada-gass."

*Arbor* parva. *Folia*  $2\frac{1}{2}$ –5 poll. longa,  $\frac{3}{4}$ – $1\frac{3}{4}$  poll. lata; *petiolo* 2 lin. longo. *Petala* 3 lin. longa, pallide flava. *Nuculania* 6 lin. longa, purpurea.

This may possibly be a variety of *Linociera Malabarica*, Wall., Wight, *Ic. t.* 1246, which it very much resembles, but the leaves of that species are represented with a shorter acumination, and the flowers are described as sessile, with the pedicels and calyces hairy, in all which respects it differs from the present plant.

3. **C. leprocarpa**, Thw.; *foliis* subcoriaceis, glabris, oblongis, acuminatis, subeveniis, basi acutis; *cymis* axillaribus, racemoideis, paucifloris, folio multum brevioribus, minutissime glanduloso-pilosulis; *floribus* brevi-pedicellatis; *calyce* lobis latis, subtruncatis, ciliatis; *petalis* (immaturis) margine involutis; *nuculanis* ovoideis, parce leprosis, putamine lignoso-chartaceo.—C.P. 2475.

HAB. Central Province, at an elevation of 3000 to 4000 feet; rather rare.

*Arbor* mediocris. *Folia*  $2\frac{1}{2}$ – $5\frac{1}{2}$  poll. longa,  $1$ – $1\frac{3}{4}$  poll. lata; *petiolo* 2–3 lin. longo. *Nuculania* purpurea, 1 poll. longa, 6–7 lin. lata.

4. **C. rostrata**, Thw.; *foliis* subcoriaceis, glabris, oblongis, longiuscule rostrato-acuminatis, basi acutis; *cymis* axillaribus, racemoideis, brevibus, paucifloris, minutissime pilosulis; *floribus* brevi-pedicellatis; *calyce* lobis rotundatis, ciliatis; *nuculanis* glabris, subfusiformibus, putamine lignoso-chartaceo.—C.P. 3472.

HAB. Hinidoon Corle.

*Folia*  $2\frac{1}{2}$ –4 poll. longa,  $\frac{3}{4}$ – $1\frac{1}{4}$  poll. lata; *petiolo* 3– $3\frac{1}{2}$  lin. longo. *Nuculania* 14 lin. longa, 6 lin. lata.

## CI. JASMINACEÆ.

### 1. JASMINUM, Tourn.

(Nom. vulg. "Pitcha.")

Sect. 1. *Unifoliata*, A. DC. *Prod.* viii. p. 301.

1. **J. Sambac**, Ait. (A. DC. l. c. var. *a*, cum syn.; Wight, *Ic. t.* 704.)—C.P. 1812.

HAB. Galle, *Gardner*; Batticaloa.

2. **J. angustifolium**, Vahl, (A. DC. l. c. p. 303, cum syn.; Wight, Ic. t. 698.)—c.p. 1807.

HAB. Very abundant in the hot, drier parts of the island, and very variable in the shape of its leaves.

3. **J. rigidum**, Zenk. (A. DC. l. c. p. 305; Wight, Ic. t. 1247, cum syn. seq.) *J. myrtophyllum*, Zenk.; A. DC. l. c. p. 304. *J. tetraphis*, Wight & Gardn. in Calc. Journ. of Nat. Hist. v. 27. p. 55.—c.p. 1809.

HAB. Hot, drier parts of the island.

*Ramuli puberuli.*

4. **J. auriculatum**, Vahl, (A. DC. l. c. p. 309, cum syn.; Wight, Ic. t. 700.)—c.p. 1810.

HAB. Very abundant in the warmer parts of the island.

Obs. *Jasm. laurifolium*, Roxb., occurs in gardens, but I have never found it in a wild state.

### Sect. 2. *Trifoliolata*, A. DC.

5. **J. flexile**, Vahl, (A. DC. l. c. p. 310; Wight, Ic. t. 1253; Burm. Thes. Zeyl. p. 127. t. 58. fig. 1.)—c.p. 724 (1814).

HAB. Not uncommon, up to an elevation of 3000 feet.

*Foliola oblonga vel lanceolata, acuminata, basi rotundata vel acuta.*

### Sect. 3. *Alternifolia*, A. DC.

6. **J. humile**, Linn. (A. DC. l. c. p. 313.) *J. revoluti*, Sims, var.  $\beta$ ; A. DC. l. c. p. 312; Wight, Ic. t. 1258, cum syn. seq. *J. bignoniaceum*, Wall., et *J. aureum*, Don.—c.p. 1811.

HAB. Elephant Plains, *Gardner*; below Horton Plains, at an elevation of 6000 feet.

## 2. NYCTANTHES, Juss.

1. **N. arbor-tristis**, Linn. (A. DC. Prod. viii. p. 314, cum syn.)—c.p. 1808.

HAB. Jaffna, *Gardner*. Not uncommon in native gardens and near Buddhist temples, but hardly truly wild. Nom. vulg. "Saipaala-gass."

A colouring matter is obtained from the corolla-tube of this plant, which is employed by the native priests for dyeing cloth yellow.

## CII. SALVADORACEÆ.

(Planchon in Ann. des Sc. Nat. (Ser. 3) x. p. 189).

### 1. SALVADORA, Linn.

1. **S. Wightiana**, Planch. MSS. in Herb. Hook. *S. Indica*, Wight, Illust. ii. p. 229. t. 181 (? Royle). *S. Persica*, Roxb. Fl. Ind. i. p. 389; Decaisne in Jacq. Voy. vol. iv. p. 140, adn.; Walp. Ann. i. p. 547 (non *S. Persica*, Linn.).—c.p. 2267.

HAB. Near the seacoast, towards the north of the island.

2. **AZIMA**, Lam.(Wight, *Illust.* ii. p. 156.)

1. **A. tetracantha**, Lam. (*Walp. Ann.* iii. p. 16, cum syn.; Wight, l. c. p. 157. t. 153.)—C.P. 2216.

HAB. Hot, drier parts of the island.

## CIII. APOCYNACEÆ.

1. **WILLUGHBEIA**, Roxb.

1. **W. Ceylanica**. — *Chilocarpus Ceylanicus*, Wight, *Ic.* t. 1288. *Winchia cirrhifera*, Gardner, MSS.—C.P. 1829.

HAB. Not uncommon in forests, up to an elevation of 4000 feet.

*Fructus* carnosus, subsphæricus vel parum pyriformis, flavo-rubescens, 4–5 poll. in diam. *Semina* oblonga, compressa,  $\frac{3}{4}$  poll. longa,  $\frac{1}{2}$  poll. lata, *testa* tenui, membranacea, *cotyledonibus* carnosis, rufescentibus, *radicula* parva.

The large, handsome fruit of this species is a favourite food of monkeys.

2. **CARISSA**, Linn.

(Nom. vulg. "Karanda-gass.")

1. **C. Carandas**, Linn. (*A. DC. Prod.* viii. p. 332, cum syn.; Wight, *Ic.* t. 426.)—C.P. 1823.

HAB. Jaffna, Gardner; Kornegalle.

2. **C. diffusa**, Roxb. (*A. DC. l. c.*; Wight, *Ic.* t. 427.)—C.P. 1822.

HAB. Very abundant in the hotter parts of the island.

3. **OPHIOXYLON**, Linn.

1. **O. serpentinum**, Willd. (*A. DC. Prod.* viii. p. 342, cum syn.; Wight, *Ic.* t. 849.)—C.P. 1834.

HAB. Common up to an elevation of 2000 feet. Nom. vulg. "Aikawaireya."

2. **O. densiflorum**, Thw.—*O. Ceylanicum*, Wight, *Ic.* t. 1291. *O. Neilgherrense*, Wight, *Ic.* t. 1292. *Tabernæmontana densiflora*, Wall. *Edw. Bot. Reg.* v. 15. t. 1273; *A. DC. l. c.* p. 373.—C.P. 1836.

HAB. Central Province, up to an elevation of 6000 feet.

4. **ALYXIA**, R. Br.

1. **A. Ceylanica**, Wight, *Ic.* t. 1293.—C.P. 1835.

HAB. Ambagamowa District, at an elevation of 1000 to 2000 feet.

*Flores* flavidi. *Carpella* matura, coccinea, 2–7-sperma.

5. **HUNTERIA**, Roxb.

(Nom. vulg. "Maddeya-gass.")

1. **H. Zeylanica**, Gardn. MSS. — *Cameraria Zeylanica*, Retz ;

Willd.; A. DC. Prod. viii. p. 389, cum syn.—An *Hunteria corymbosa*, Roxb. var. ?—c.p. 1827.

HAB. Warmer parts of the island; common.

*Flores* flavidi. *Carpella* oblonga, rostrato-acuminata, stipitata, pollicaria.

The wood of this species has a very fine, close grain, resembling that of the Box-tree.

2. ? *H. Roxburghiana*, Wight, Ic. t. 1294.—c.p. 2518.

HAB. Central Province, in the Maturatte and Badulla Districts.

*Folia* lineari-lanceolata, obtuse acuminata, basi acuta, superne nitida, subtus opaca pallidioraque, venis primariis parallelis parum conspicuis, 2–6 poll. longa,  $\frac{1}{2}$ –1 poll. lata, *petiolo* 3–5 lin. longo. *Flores* flavidi. *Carpella* (immatura) parce acuminata.

This would seem to have the leaves narrower and less numerous and strongly veined than Dr. Wight's plant, as represented in his figure, but in other respects there appears to be great similarity.

## 6. CERBERA, Linn.

1. *C. Odollam*, Gært. (A. DC. Prod. viii. p. 353, cum syn.; Wight, Ic. t. 441.)—c.p. 1832.

HAB. Hotter parts of the island, especially near the sea. Nom. vulg. "Gōng-kadooroo-gass."

## 7. OCHROSIA, Juss.

1. *O. Borbonica*, Gm. (A. DC. Prod. viii. p. 356, cum syn.)—*O.?* *platyspermos*, A. DC. l. c. cum syn. *Cerbera parviflora*, Moon's Cat. p. 19. An Willd. i. p. 1222?—c.p. 1833.

HAB. Caltura, and other places near the sea. Nom. vulg. "Moodoo-kadooroo-gass."

## 8. TABERNÆMONTANA, Plum.

1. *T. dichotoma*, Roxb. (A. DC. Prod. viii. p. 366, cum syn.; Wight, Ic. t. 433.) *Cerbera Manghas*, Linn. (*partim.*) *Manghas lactescens*, etc., Burm. Thes. Zeyl. 151. t. 70. f. 1.—c.p. 2834.

HAB. Very common in the warmer parts of the island. Nom. vulg. "Diwi-kadooroo-gass."

*Semina* matura in pulpa sanguinea nidulantia.

This is no doubt the plant intended by Burmann, judging from his description of the fruit, though he has given the native name of another species.

## 9. VALLARIS, Burm.

1. *V. Pergulana*, Burm. (A. DC. Prod. viii. p. 399, cum syn.; Wight, Ic. t. 429.)—c.p. 2515.

HAB. Colombo, Mr. W. Ferguson.

2. *V. dichotoma*, Wall. (A. DC. l. c. p. 400; Wight, Ic. t. 438, cum syn.)—c.p. 2519.

HAB. Hot, drier parts of the island, at no great elevation.

The smaller foliage, and a slight difference in the shape of the corolla of the present plant, give it a somewhat distinct appearance from the preceding, but the two are so closely allied that it would perhaps be quite safe to consider them as mere

varieties of one species. The flowers of *V. Pergulana* have a disagreeable odour, but I do not remember whether the same character applies to those of *V. dichotoma*. I can discover no difference in the nectaries of the two plants.

## 10. HELIGME, Blume, Endl.

*Parsonsiæ* Sect. 2, A. DC.

1. **H. spiralis**.—*H. Rheedii*, Wight, Ic. t. 1303, cum syn. *Parsonsia spiralis*, Wallich; A. DC. Prod. viii. p. 402.—c.p. 1862.

HAB. Very abundant, up to an elevation of 3000 feet.

*Flores virides, externe rufescentes.*

## 11. WRIGHTIA, R. Br.

1. **W. tomentosa**, R. et Sch. (A. DC. Prod. viii. p. 404, cum syn.; Wight, Ic. t. 443.)—c.p. 2691.

HAB. Central Province; not very uncommon.

*Corolla* flavida, rubro-tincta. *Corona* coccinea, breve et obtuse 5-loba, lobis subcrenatis. *Flores* siccitate cyaneo-nigrescentes.

2. ? **W. Rothii**, G. Don, (A. DC. l. c. p. 406, cum syn.; Wight, Ic. t. 1319.) Var.  $\beta$  *puberula*.—c.p. 1837.

HAB. Damboul, Gardner.

*Ramuli* cum *foliis junioribus* puberuli. *Antheræ* dorso pilosæ.

This will probably prove to be distinct from *W. Rothii*, but my specimen is not sufficiently perfect to justify a new species being established upon it.

3. **W. angustifolia**, Thw.; *foliis* glabris, membranaceis, lineari-lanceolatis, obtusis, basi acutis, brevi-petiolatis; *cymis* delicatulis; *floribus* minute pubescentibus; *corona* glabra, 5-partita; *lobis* linearibus, antheris oppositis æquilongisque, fere ad medium bifurcatis, segmentis simplicibus vel ad apicem parce laciniatis.—c.p. 1839.

HAB. Damboul and Trincomale, Gardner.

*Folia*  $1\frac{1}{2}$ –4 poll. longa,  $\frac{1}{2}$ – $\frac{3}{4}$  poll. lata; *petiolo* circiter 1 lin. longo. *Pedicelli* tenues, semipollicares. *Calyx*  $\frac{3}{4}$  lin. longus, lobis rotundatis, subacutis, ciliatis, *corollæ* tubo brevioribus. *Corollæ* lobi lineari-oblongi, obtusi, 3 lin. longi. *Antheræ* dorso parum pilosæ.

This much resembles *W. tinctoria*, R. Br. (c.p. 1838, 2385), but its much narrower and more membranaceous leaves, and very differently shaped corona, seem sufficiently to distinguish it from that species. Of the latter I have only garden specimens in the herbarium, and I have no evidence of its having been detected wild in the island, though it may be expected to occur so in the northern districts.

4. **W. Zeylanica**, R. Br. (A. DC. l. c. p. 407, cum syn.)—*W. anti-dysenterica*, R. Br.; A. DC. l. c. cum syn.—c.p. 1825.

HAB. Common in the south of the island. Nom. vulg. "Sooddoo-idda."

*Corolla* alba, tubo glabro, lobis puberulis. *Corona* alba, antheris longior, pluri-laciniata.

## 12. ALSTONIA, R. Br.

1. **A. scholaris**, R. Br. (A. DC. Prod. viii. p. 408, cum syn.; Wight, Ic. t. 422.)—c.p. 1840.



HAB. Common up to an elevation of 3000 feet. Nom. vulg. "Roo-kattana-gass."

The light wood of this tree is employed for making coffins.

### 13. HOLARRHENA, R. Br.

1. *H. mitis*, R. Br. (A. DC. Prod. viii. p. 413.)—c.p. 756.

HAB. Not uncommon, up to an elevation of 1500 feet. Nom. vulg. "Kirri-walla-gass."

*Arbor mediocris. Cymæ folio breviores, 5-12-floræ. Corolla alba.*

The wood of this tree is light in weight, of a pale colour, and fine, close grain, and is used for inlaying cabinet-work.

### 14. CHONEMORPHA, G. Don.

1. *C. macrophylla*, G. Don, (A. DC. Prod. viii. p. 430, cum syn.; Wight, Ic. t. 432.)—c.p. 2467.

HAB. Not very uncommon, up to an elevation of 2000 feet.

### 15. AGANOSMA, G. Don.

1. *A. elegans*, G. Don, (A. DC. Prod. viii. p. 433, cum syn.; Wight, Ic. t. 1304.)—c.p. 1850.

HAB. Hot, drier parts of the island, not uncommon.

*Folliculi teretes, 5-9 poll. longi, 2½ lin. lati, testaceo-pilosi, denique subglabri. Semina glabra, lineari-oblonga, nigra, 7 lin. longa, 1½ lin. lata, brevi-rostrata, longe comosa.*

### 16. CLEGHORNIA, Wight, Walpers.

1. *C. acuminata*, Wight, Ic. t. 1310.—*C. cymosa*, Wight, Ic. t. 1312.—c.p. 1861.

HAB. Common in the Central Province, at an elevation of 2000 to 5000 feet.

The two forms of this plant, described by Dr. Wight under two specific names, are connected by intermediate ones, and have, therefore, no claim to be kept distinct.

### 17. ICHNOCARPUS, R. Br.

1. *I. frutescens*, R. Br. (A. DC. Prod. viii. p. 435, cum syn.; Wight, Ic. t. 430.)—c.p. 1863.

HAB. Common up to an elevation of 3000 feet. Nom. vulg. "Kirri-wel."

### 18. ANODENDRON, A. DC.

1. *A. paniculatum*, A. DC. Prod. viii. p. 444, cum syn.; Wight, Ic. t. 396.—c.p. 1843.

HAB. Common up to an elevation of 2000 feet. Nom. vulg. "Dool."

The fibre of the stem is very fine and strong, and is much used by the Cinghalese.

2. *A. rhinosporum*, Thw.; glabra; foliis subcoriaceis, oblongis, utrinque plus minus abrupte angustatis, obtusis, subtus pallidioribus,

inconspicue venosis; *cymis* terminalibus, delicatulis, valde divaricato-ramosis; *pedicellis* flore longioribus; *seminibus* rostro gracili æquilongo ornatis.—C.P. 2579.

HAB. Ambagamowa and Maturatte Districts; not common.

*Folia* 2-4 poll. longa,  $\frac{1}{2}$ -1 $\frac{1}{2}$  poll. lata; *petiolo* 3-5 lin. longo. *Calyx*  $\frac{1}{2}$  lin. longus, lobis ovatis, subacutis. *Corolla* flava, 3 lin. in exp. *Antheræ* sessiles. *Folliculi* 3-5 poll. longi. *Semina* glabra, minute rugulosa (rostr. excl.), 6 lin. longa ad apicem copiose comosa.

*Vinca rosea*, Linn., is become quite naturalized in the neighbourhood of Colombo and other places. *Plumeria acutifolia*, Poir., is also widely distributed, especially about Buddhist temples.

## CIV. ASCLEPIADACEÆ.

### 1. HEMIDESMUS, R. Br.

1. **H. Indicus**, R. Br. (Decaisne in DC. Prod. viii. p. 494, cum syn.; Wight, Ic. t. 594.)—C.P. 183 (1865).

HAB. Very abundant in the warmer parts of the island. Nom. vulg. "Eremoosoo."

The root of this plant (Indian Sarsaparilla) is much used medicinally by the Cinghalese.

### 2. CRYPTOLEPIS, R. Br.

(Blume, Mus. Bot. Lugd.-Bat. i. p. 146.)

1. **C. Buchanani**, Roem. et Sch. (Walp. Rep. vi. p. 474, cum syn.; Wight, Ic. t. 494.)—C.P. 2548.

HAB. Common, up to an elevation of 3000 feet.

### 3. SECAMONE, R. Br.

1. **S. emetica**, R. Br. (Decaisne in DC. Prod. viii. p. 501, cum syn.; Wight, Ic. t. 1283.)—C.P. 1851.

HAB. Not uncommon in the warmer parts of the island.

### 4. TOXOCARPUS, W. et A.

1. **T. Kleinii**, W. et A. (Decaisne in DC. Prod. viii. p. 505; Wight, Ic. t. 886.)—C.P. 2578.

HAB. Maturatte District, at an elevation of 3000 feet; not common.

### 5. CYNOCYTONUM, Er. Mey., Decaisne.

1. **C. pauciflorum**, Decaisne in DC. Prod. viii. p. 528, cum syn.—*Cynanchum pauciflorum*, R. Br.; Wight, Ic. t. 354. An *C. pedunculatum*, Thunb. DC. Prod. viii. p. 548.—C.P. 2466.

HAB. Very common in the Central Province. Nom. vulg. "Kangkoombala."

The Cinghalese eat the young leaves of this and of many other plants of this Natural Family, in their curries.

## 6. CALOTROPIS, R. Br.

1. **C. gigantea**, R. Br. (Decaisne in DC. Prod. viii. p. 535, cum syn.; Wight, Illust. ii. t. 155.)—C.P. 1831.

HAB. Hot, drier parts of the island. Nom. vulg. "Warra."

The fibre of this plant is very fine and strong. The milky juice is employed medicinally.

## 7. PENTATROPIS, R. Br.

1. **P. microphylla**, W. et A. (Decaisne in DC. Prod. viii. p. 536, cum syn.; Wight, Ic. t. 352.)—An *Cynanchum acuminatum*, Thunb. DC. l. c. p. 548?—C.P. 1853.

HAB. Not uncommon in the warmer parts of the island.

## 8. SARCOSTEMMA, R. Br.

1. **S. Brunoniana**, W. et A. (Decaisne in DC. Prod. viii. p. 538; Wight, Ic. t. 1282.)—C.P. 1830.

HAB. Hot, drier parts of the island; common. Nom. vulg. "Moowa-keeriya."

The stems are used medicinally.

## 9. OXYSTELMA, R. Br.

1. **O. esculentum**, R. Br. (Decaisne in DC. Prod. viii. p. 543, cum syn.)—C.P. 2837.

HAB. Hot, drier parts of the island.

## 10. DÆMIA, R. Br.

1. **D. extensa**, R. Br. (Decaisne in DC. Prod. viii. p. 544, cum syn.; Wight, Ic. t. 596.)—*Cynanchum echinatum*, Thunb. DC. l. c. p. 459.—C.P. 1841.

HAB. Not uncommon in the warmer parts of the island.

## 11. TYLOPHORA, R. Br.

1. **T. carnosia**, Wall. (Decaisne in DC. Prod. viii. p. 607; Wight, Ic. t. 351.)—C.P. 1852.

HAB. Hotter parts of the island.

2. **T. cordifolia**, Thw.; volubilis; ramulis pilosulis; foliis glabris, cordatis, mucronatis, basi laminae parce glandulosis; petiolo superne pilosulo; pedunculis 1-2-floris, folio subæquilongis, cum floribus glabris; coronæ stamineæ radiatæ foliolis oblongis, truncatis, carnosis, postice superneque parum productis et ibi lateraliter compressis; gynostegium convexum umbilicatum subæquantibus.—C.P. 717.

HAB. Common in the Central Province.

Folia 3-13 lin. longa, 2-9 lin. lata; petiolo 1-1½ lin. longo. Calyx parvus, lobis ovatis, subacutis. Corolla fusco-purpurea, 4 lin. in exp., lobis ovatis, subacutis. Corona staminea 1 lin. lata, flavida. Folliculi sæpius abortu solitarii, attenuati, glabri, bipollicares.

3. **T. micrantha**, Thw.—*T. Zeylanica*, Decaisne, l. c. p. 608. *Cynanchum micranthum*, Thunb. DC. l. c. p. 548.—c.p. 2517.

HAB. Central Province, at an elevation of 2000 to 4000 feet.

*Folia* 2½–5 poll. longa, 1¼–3 poll. lata; *petiolo* ½–1 poll. longo. *Folliculi* elegantuli, glabri, 5 poll. longi.

4. **T. fasciculata**, Ham. (DC. l. c. p. 608; Wight, Ic. t. 848.)—c.p. 369.

HAB. Not uncommon in the Central Province, at an elevation of 4000 to 7000 feet.

*Flores* flavidi, purpureo-tincti, vel omnino fusco-purpurei.

5. **T. asthmatica**, W. et A. (DC. l. c. p. 611, cum syn.; Wight, Ic. t. 1277.)

Var. *a. pubescens*, DC. l. c.—c.p. 1857 (1858).

Var. *β. glabra*, DC. l. c.—An *Cynanchum flavens*, Thunb. et *C. bracteatum*, Thunb. DC. l. c. p. 549?—c.p. 1849 (1860).

HAB. Very abundant in the warmer parts of the island. Nom. vulg. "Binnooga."

*Folia* ad apicem petioli sæpe parce glandulosa. *Flores* flavidi, plus minus purpureo-tincti.

A very variable plant as regards the shape of the leaves and the amount of pubescence. The natives employ the roots medicinally.

## 12. COSMOSTIGMA, Wight.

1. **C. racemosum**, Wight (DC. in DC. Prod. viii. p. 613, cum syn.; Wight, Ic. t. 591.)—*C. acuminatum*, Wight, Ic. t. 1270.—c.p. 1854.

HAB. Warmer parts of the island; common.

## 13. MARSDENIA, R. Br.

1. **M. tenacissima**, W. et A. (DC. in DC. Prod. viii. p. 616, cum syn.; Wight, Ic. t. 590.)—c.p. 2860.

HAB. Hotter parts of the island; not very common.

## 14. GYMNEMA, R. Br.

1. **G. sylvestre**, R. Br. (DC. in DC. Prod. viii. p. 621, cum syn.; Wight, Ic. t. 349.)—c.p. 2549.

Var. *β. Decaisneanum*.—*G. Decaisneanum*, Wight, Ic. t. 1271. *G. hirsutum*, DC. l. c. p. 622 ("non W. et A.," Wight, l. c.)—c.p. 1844.

HAB. Not uncommon, up to an elevation of 3000 feet.

2. **G. rotundatum**, Thw.; volubilis; *ramulis junioribus* tomentosus; *foliis* ovato-rotundatis, obtuse acuminatis, superne pilosulis, ad apicem petioli pluri-glandulosus, subtus cum *petiolo* longiusculo brevi-tomentosis; *pedunculis* petiolo duplo brevioribus; *umbellis* geminatis, confertifloris; *stigmatibus* conico staminibus longiore; *folliculis* attenuatis, glabris.—c.p. 3086.

HAB. Central Province, at an elevation of 3000 feet.

*Folia* 2–4 poll. longa, 1½–3 poll. lata; *petiolo* ¾–1½ poll. longo. *Pedicelli* flore duplo longiores. *Calycis lobi* oblongo-rotundati, pilosi. *Corolla* flava, calyce duplo longior.

3. *G. lactiferum*, R. Br. (DC. l. c. p. 622, cum syn.)—*G. Zeylanicum*, Decaisne, l. c.—C.P. 2580.

Var.  $\beta$ ; *foliis* subtus molliter parce pubescentibus.—C.P. 1847.

HAB. Not uncommon, up to an elevation of 3000 feet.

*Folliculi* (var.  $\alpha$ ) lanceolati,  $3\frac{1}{2}$  poll. longi,  $\frac{3}{4}$  poll. in lat. max., obtusi, sæpius abortu solitarii, juniores ovati vel oblongi.

## 15. BIDARIA, Endl.

1. *B. pergularioides*, Thw.; volubilis; *ramulis* cum *petiolis*, *pedunculis* *pedicellis*que pubescentibus; *foliis* subglabris, ellipticis vel oblongis, plus minus acuminatis, basi acutis vel rotundatis, ad apicem petioli glandulosis; *pedunculis* petiolo subæquilongis; *pedicellis* brevibus; *folliculis* lineari-lanceolatis, valde attenuatis, puberulis, denique subglabris.—*Gymnema pergularioides*, Wight et Gardner, MSS.—An *Cynanchum cuspidatum*, Thunb. DC. l. c. p. 548?

Var.  $\alpha$ ; *corollæ* tubo lobis calycinis 2-3-plo longiore; *folliculis* longiusculis.—C.P. 2671, 2672.

Var.  $\beta$ ; *foliis* minoribus, firmioribus; *corollæ* tubo lobis calycinis vix longiore; *folliculis* brevioribus.—C.P. 1848.

*Folia*  $1\frac{1}{2}$ -6 (var.  $\beta$ , 1-2) poll. longa,  $\frac{1}{2}$ -2 poll. (var.  $\beta$ , 3-10 lin.) lata; *petiolo* 2-7 lin. longo. *Pedicelli* *corollæ* tubo subæquilongi. *Calycis* lobi ovati, subacuti. *Corolla* flava, sæpius plus minus rubro-tincta, 3-4 (var.  $\beta$ , 2-3) lin. in exp., lobis ovato-lanceolatis. *Folliculi* 5-6 (var.  $\beta$ , circiter 3) poll. longi, 3 lin. lati.

This variable species is evidently very closely allied to *B. elegans*, Decaisne.

## 16. LEPTADENIA, R. Br.

1. *L. reticulata*, W. et A. (Decaisne in DC. Prod. viii. p. 628, cum syn.; Wight, Ic. t. 350.)—An *Cynanchum ovatum*, Thunb. DC. l. c. p. 548?—C.P. 1846.

HAB. Anooradhapoorra, Gardner.

## 17. HETEROSTEMMA, W. et A.

1. *H. Tanjorensis*, W. et A. (DC. in DC. Prod. viii. p. 630; Wight, Ic. t. 348.)—C.P. 1856 (1859).

HAB. Central Province, up to an elevation of 4000 feet.

## 18. DISCHIDIA, R. Br.

1. ? *D. nummularia*, R. Br. (DC. in DC. Prod. viii. p. 632, cum syn.)

I have a drawing, made some years ago, from a plant collected in the island, which is either *D. nummularia* or a species very closely allied to it, but unfortunately the specimen was not preserved, nor is the particular locality in which it was found recorded.

## 19. HOYA, R. Br.

1. *H. ovalifolia*, W. et A. (Wight, Ic. t. 847.)—C.P. 2670.

HAB. Not very uncommon in the Ambagamowa District, at no great elevation.

*Folia* ad apicem petioli glandulosa. *Flores* albidi, plus minus rubro-tincti.

2. **H. Wightiana**, Thw.; volubilis; *foliis* glabris, carnosis, lineari-lanceolatis, acuminatis, obtusis, basi angustatis; *pedunculis* brevibus, multifloris; *corollæ lobis* ovatis, parce acuminatis, obtusis, minute papillois; *corona staminea* parum depressa, *foliolis* obovatis, obtusis, interne acutis.—C.P. 2762.

HAB. Central Province, at an elevation of 3000 to 5000 feet.

*Folia* 1-3 poll. longa, 2-5 lin. lata; *petiolo* purpurascente, 1-3 lin. longo. *Pedicelli* glabri, 6 lin. longi. *Calyx* minutus, glaber. *Corolla* alba, 6-7 lin. in exp. *Corona staminea* purpurea, 2 lin. in exp.

A very beautiful species, bearing considerable resemblance to the figure of Dr. Wight's *H. pauciflora* (*H. parviflora* in desc.) Ic. t. 1269; and its more numerous flowers and obtusely-pointed leaves seem to be the principal points of distinction.

3. **H. viridiflora**, R. Br. (DC. l. c. p. 639; Wight, Ic. t. 586.)—C.P. 1855.

HAB. Common, up to an elevation of 3000 feet. Nom. vulg. "Kirriangoona."

The leaves of this species are eaten by the Cinghalese.

## 20. CEROPEGIA, Linn.

1. **C. elegans**, Wall. (Bot. Mag. t. 3015; DC. in DC. Prod. viii. p. 642; Wight, Ic. t. 1265.)—*C. sphenanthera*, W. et A.; DC. l. c. p. 643. *C. Mysorensis*, Wight, Ic. t. 846. *C. Walkeri*, Wight, Ic. t. 1266.—C.P. 738.

HAB. Not uncommon, up to an elevation of 3000 feet.

This and the other species of the genus are very variable as regards the shape and size of the leaves and colour of the flowers, and far from constant in the shape and size of the latter.

2. **C. Gardneri**, Thw.; volubilis, glabra; *foliis* lanceolatis, acuminatis; *pedunculis* petiolo subæquilongis, paucifloris; *calycis lobis* linearibus, acutis; *corolla* ad tubi basim parum inflata, supra medium subito valde ampliata, apice aperte 5-crumenata, margine ciliata; *coronæ staminæ lobis exterioribus* linearibus, acutis; *interioribus* multo longioribus latioribusque et apice reflexis; *folliculis* longiusculis, terebibus.—C.P. 2838.

HAB. Rambodde, at an elevation of 4000 to 5000 feet, Gardner.

*Pedicelli* pedunculo subæquilongi. *Corolla* albida, purpureo-maculata, circiter 2 poll. longa, 1½ poll. in exp. *Folliculi* (immaturi) 7 poll. longi, 2 lin. lati.

A very pretty species, allied to the preceding, but it can scarcely be a variety of it.

3. **C. Thwaitesii**, Hook. Bot. Mag. t. 4758.—C.P. 1842.

HAB. Central Province; not very common.

This will possibly prove to be a small variety of *C. Decaisneana*, Wight, Ic. t. 1259, to which it is evidently very closely allied.

4. **C. candelabrum**, Linn. (DC. l. c. p. 643, cum syn.)—*C. intermedia*, Wight, Ic. t. 1263.—C.P. 774.

HAB. Hotter parts of the island; not uncommon.

An extremely variable species, and I suspect that several Indian plants, described as distinct species, will be discovered to be mere varieties of it.

21. **CARALLUMA**, R. Br.

1. **C. attenuata**, Wight, Ic. t. 1268.—An *C. adscendens*, R. Br.; DC. in DC. Prod. viii. p. 647, cum syn. P.—c.p. 3304.

HAB. On rocky ground at Ooma Oya, on the lower road from Kandy to Badulla, at no great elevation.

My specimens of this plant so much resemble Roxburgh's figure of *S. adscendens* in Pl. Cor. t. 30, that it seems not improbable that the cilia were accidentally absent, or overlooked in the corolla of his specimen.

22. **BOUCEROSIA**, W. et A.

1. **B. umbellata**, W. et A. (DC. in DC. Prod. viii. p. 648, cum syn.; Wight, Ic. t. 495.)—*B. campanulata*, Wight, Ic. t. 1287.—c.p. 2861.

HAB. Rocks near Kornegalle.

*Asclepias curassavica*, Linn., is become quite naturalized and distributed throughout the island.

CV. **LOGANIACEÆ**.1. **MITRASACME**, Labill.

1. **M. Indica**, Wight, Ic. t. 1601; Bentham in Linn. Soc. Proc. i. p. 92, cum syn.—c.p. 1677.

HAB. Common in the warmer parts of the island.

2. **FAGRÆA**, Thunb.

1. **F. Zeylanica**, Thunb. (DC. Prod. ix. p. 29, cum syn.; Blume, Rumphia, t. 78. fig. 2.)—*Solandra oppositifolia*, Moon's Cat. p. 15.

HAB. Central Province, not common. Nom. vulg. "Atamboorogass."

2. **F. obovata**, Wall. in Roxb. Fl. Ind. ii. p. 33; DC. l. c.—*F. Malabarica*, Wight, Ic. t. 1317.—c.p. 757.

HAB. Central Province, at an elevation of 2000 to 5000 feet.

This differs from *F. Zeylanica* in its more scandent habit, much shorter flowers, and more oblong fruit; but in other respects they are very closely allied. In both, the *placentæ* are distinct, not, as in the following species, cohering in the centre of the ovary.

3. **F. Gardneri**, Thw.; scandens; *foliis* majusculis, coriaceis, glabris, subeveniis, obovato-cuneatis, apice rotundatis, basi truncatis, brevipetiolatis; *petiolo* parum stipulaceo; *corolla* late infundibuliformi; *tubo* calyce circiter 3-plo longiore; *placentis* in medio ovarii coherentibus.—c.p. 1826.

HAB. Mantani, in the Central Province, at an elevation of 3000 to 4000 feet.

The present species has weaker, more scandent branches, than the preceding. Its leaves have a more crowded appearance, and the calyx is larger, with its pair of bracteoles at its immediate base, instead of halfway down the pedicel, as in the preceding. The two plants are, however, very closely allied, and their flowers are very similar in general appearance. The somewhat coriaceous flowers of all three species are of a yellowish-white colour; the fruit is shining and glaucous.

## 3. STRYCHNOS, Linn.

## Sect. 1. Scandentes.

1. **S. minor**, Blume, (A. DC. Prod. ix. p. 14; Benth. in Journ. Linn. Soc. Bot. i. p. 100.)—c.p. 187.

Var.  $\delta$ . *parvifolia*, Benth. l. c.—c.p. 341.

HAB. Var. *a*. Warmer parts of the island. Var.  $\delta$ . Central Province, up to an elevation of 6000 feet.

2. **S. colubrina**, Linn. (A. DC. l. c.; Benth. l. c. p. 101.)—c.p. 2516.

HAB. Hot, drier parts of the island, not uncommon.

*Rami juniores parce puberuli, cito glabrati. Folia sæpe ad basin sub-5-nervia. Corollæ lobi interne versus apicem pilis paucis rigidis erectis ornati.*

Distinguished from the preceding by its pentamerous flowers and larger fruit, as indicated by Mr. Bentham. I have never seen the fruit nearly so large as it is represented by Rheede, but have never met with it quite mature. The leaves of the present species are most frequently somewhat 5-nerved at the base, owing to the distinctness of the submarginal nervure.

3. **S. laurina**, Wall. (A. DC. l. c. p. 13; Benth. l. c. p. 102.)—c.p. 3540.

HAB. Galle, Kornegalle, and other of the warmer parts of the island.

*Nuculania (immatura) pollicaria.*

4. **S. cinnamomifolia**, Thw.; glabra; *cirrhis simplicibus; foliis oblongis, rostrato-acuminatis, obtusis, basi acutis, triplinerviis; cymis terminalibus, pedunculatis, trichotomis, folio brevioribus; corollæ lobis glabris; nuculanis majusculis, sphaericis, polyspermis.*—c.p. 1867.

HAB. Hantani District, at an elevation of 3000 feet. Nom. vulg. "Atta-kirindi-wel."

*Folia 2½-4 poll. longa, 1¼-1¾ poll. lata; petiolo 2-3 lin. longo. Cymæ tomentellæ, ramulos breves terminantes. Corolla albida, tubo interne infra medium sparse piloso, 4-5 lin. longo, lobis 1½ lin. longis. Nuculania circiter 3 poll. in diam. Semina pollicaria.*

This large climbing plant much resembles Blume's figure of *S. Tiente*, Lesch., Rumphia, t. 24, and would seem also to have many characters in common with *S. ovalifolia*, Wall.; but in these latter the inflorescence is axillary, whilst in the present species it terminates the small ramuli, as in *S. nux-vomica*. From the last-named species the present differs in, besides its scandent habit, having narrower, more acuminate leaves, and larger flowers.

## Sect. 2. Arborescentes.

5. **S. nux-vomica**, Linn. (A. DC. l. c. p. 15, cum syn.)—c.p. 2839.

HAB. Hotter parts of the island. Nom. vulg. "Goda-kadooroo-gass."

*Corolla interne infra medium pilosa. Nuculania sæpe 1-2-sperma.*

6. **S. potatorum**, Linn. f. (A. DC. l. c. cum syn.; Wight, Illust. ii. t. 156.)—c.p. 1866, 3367.

HAB. Hot, drier, and especially the northern parts of the island. Nom. vulg. "Ingini-gass."

## 4. GÆRTNERA, Lam.

1. **G. rosea**, Thw. (Benth. in Linn. Soc. Proc. i. p. 111.)—c.p. 2673.



HAB. Ambagamowa and Saffragam Districts, up to an elevation of 2000 feet.

*Frutex* glaber, 8-10-pedalis. *Ramuli* bicostati. *Folia* elliptico-vel ovato-lanceolata, longe acuteque acuminata, basi in petiolum brevem angustata,  $1\frac{1}{4}$ -3 poll. longa,  $\frac{1}{4}$ - $1\frac{1}{4}$  poll. lata. *Stipulæ* carinatae, biaristatae, in tubum coalitæ. *Flores* terminales, terni, sessiles ad apicem *pedunculi* brevis vel subnulli. *Calyx* tubulosus, 1 lin. longus, irregulariter 4-6-denticulatus. *Corolla* 8-10 lin. longa, externe roseo-purpurea, lobis 4, linearibus, tubo multum brevioribus. *Antheræ* 4, oblongæ, subsessiles. *Nuculania* subsphærica, pallide cyanea, 4-5 lin. in diam.

2. **G. Walkeri**, Wight, *Illust.* ii. t. 156 *b*; Bentham, l. c. (*partim*).—*Sykesia Walkeri*, Arn. *Pug.* 36, 354; DC. *Prod.* ix. p. 35.—C.P. 288.

HAB. Central Province, at an elevation of 3000 to 6000 feet.

*Frutex* 8-10-pedalis. *Ramuli* teretes, glabri. *Folia*  $1\frac{1}{2}$ -4 poll. longa,  $\frac{1}{2}$ - $1\frac{1}{2}$  poll. lata; *petiolo* 2-5 lin. longo. *Stipulæ* coalitæ, truncatæ vel brevi-acuminatæ et apice bifidæ. *Cymæ* terminales, glabræ, sæpius 5-floræ, folio subæquilongæ. *Nuculania* subsphærica, cyanea.

3. **G. Gardneri**, Thw.; *ramulis* teretibus, minute puberulis; *foliis* glabris, anguste lanceolatis, acute acuminatis, basi acutis, brevi-petiolatis; *stipulis* longe coalitis, parum acuminatis bifidisque; *cymis* puberulis, sæpius 3-floris et folio brevioribus, floribus erectis.—C.P. 346 (363).

HAB. Central Province, at an elevation of 5000 to 7000 feet.

*Frutex* 8-10-pedalis. *Folia* 1- $2\frac{1}{2}$  poll. longa, 2-5 lin. lata. *Flores* fructusque fere ut in præcedente sed parum majores.

The puberulous ramuli and inflorescence, differently-shaped leaves, fewer-flowered cymes, and rather larger flowers and fruit, distinguish this from the preceding species, to which it is very closely allied.

4. **G. ternifolia**, Thw.; *foliis* ternis, parvis, glabris, rigidulis, linearibus, mucronatis, utrinque acutis, eveniis, brevi-petiolatis; *floribus* terminalibus, solitariis, pedicellatis, cernuis.—*G. Walkeri*, var. *angustifolia*, Benth. l. c. (*partim*).—C.P. 440 (457).

HAB. Ambagamowa District, near Adam's Peak, at an elevation of 4000 to 5000 feet.

*Frutex* 4-6-pedalis. *Ramuli* puberuli, 3-costati. *Folia* 4-7 lin. longa,  $\frac{1}{2}$ -1 lin. lata, costa subtus prominenti. *Stipulæ* breves, acuminatæ, basi coalitæ. *Pedicelli* folio breviores, puberuli. *Calyx* brevi-campanulatus, acute 5-dentatus, puberulus, basi bracteolatus. *Corolla* alba, 5-6 lin. longa, parum campanulata, externe glabra, tubo intus superne villosulo, lobis longiore. *Filamenta* corollæ tubi medio affixa, *antheras* oblongas, exsertas æquantia. *Nuculania* subsphærica, cyanea, 4 lin. in diam.

A very beautiful and very distinct species.

5. **G. Koenigii**, Wight, *Ic.* t. 1318; Benth. l. c. p. 112.—*G. acuminata*, Benth. l. c.; *Sykesia Koenigii*, Arn. *Pug.* 35 (353); DC. l. c.—C.P. 544.

Var.  $\beta$ . *thyrsiflora*.—*G. thyrsiflora*, Blume, *Mus. Bot.* i. p. 174; Benth. l. c. *G. oryphylla*, Benth. l. c. *Sykesia thyrsiflora*, Arn. l. c.; DC. l. c.—C.P. 3342.

HAB. Var. *a*. Very abundant up to an elevation of 3000 feet. Var.  $\beta$  south of the island, at no great elevation.

The somewhat smaller leaves and flowers, and the elongated and often depauperated cymes of var.  $\beta$ , appear to me to be the only characters to separate it from var. *a*.

## CVI. GENTIANACEÆ.

## TRIBE I. GENTIANEÆ.

## 1. EXACUM, Linn.

Sect. 1. *Pseudochironia*, Griseb.

1. **E. Zeylanicum**, Roxb. (Griseb. in DC. Prod. ix. p. 45, cum syn.; Hook. Bot. Mag. t. 4423.)—*Chironia trinervia*, Linn.; Moon's Cat. p. 16.—c.p. 512.

Var.  $\beta$ ; *floribus* minoribus pallidioribusque; *antheris* brevioribus.—*E. Ramboddeanum*, Gardn. MS.—c.p. 38.

HAB. Common in damp, grassy places, up to an elevation of 5000 feet. Nom. vulg. "Bindara."

*Caulis* plus minus tetragonus vel subteres. *Folia* ovata vel anguste lanceolata, utrinque plus minus acuminata. *Antheræ* var.  $\alpha$  circiter 3 lin. (var.  $\beta$   $1\frac{1}{2}$  lin.) longæ.

At one time I considered, as did Dr. Gardner, var.  $\alpha$  and  $\beta$  distinct species, but after the examination of very numerous specimens, I am unable to find any essential permanent character to separate them.

2. **E. macranthum**, Arn. (Griseb. l. c. p. 46; Hook. Bot. Mag. t. 4771.)—*E. ovale*, Griseb. l. c.—c.p. 1877.

HAB. Central Province, at an elevation of 5000 to 7000 feet.

*Caulis* teres vel parum tetragonus. *Corollæ* purpureæ, *lobis* rotundatis, plus minus acuminatis. *Antheræ* 4–5 lin. longæ.

This beautiful plant differs from the preceding only in the deep purple colour of its larger flowers and the greater size of their anthers, and it might safely perhaps be considered a mere mountain variety of it.

3. **E. Walkeri**, Arn. (Griseb. l. c. p. 45.)—c.p. 1878.

HAB. Central Province, at an elevation of 5000 to 6000 feet.

*Caulis* tetragonus, ad nodos minute glandulosus. *Corolla* alba.

4. **E. axillare**, Thw.; *caule* parum procumbente, 4-alato; *foliis* 3-nerviis, ovato-lanceolatis, utrinque acuminatis; *pedicellis* axillaribus, solitariis, calyce brevioribus; *floribus* 4-meris; *calyce* late alato; *corollæ lobis* lanceolatis, acuminatis.—c.p. 2998.

HAB. Near Hewessee, in the Pasdoon Corle, at no great elevation.

*Folia* membranacea,  $1\frac{1}{2}$ –3 poll. longa. *Calyx* semipollicaris; *alis* membranaceis, laxè venosis. *Corolla* cyanea, *lobis*  $\frac{1}{2}$  poll. longis.

Sect. 2. *Pseudosebæa*, Griseb.

5. **E. pedunculatum**, Linn. (Griseb. l. c. p. 46, cum syn.; Wight, Ic. t. 336.)—c.p. 1876.

HAB. Hotter parts of the island, not uncommon.

*Corolla* pallide cyanea vel albida.

6. **E. sessile**, Linn. (Griseb. l. c. p. 47, cum syn.; Wight, Ic. t. 1324, f. 1.)—c.p. 250.

HAB. Common in grassy places, up to an elevation of 2000 feet.

*Corolla* cyanea vel albida.

2. **PLADERA**, Roxb.

1. **P. pusilla**, Roxb. (Griseb. in DC. Prod. ix. p. 63, cum syn.)—*Cicendia fastigiata*, Griseb. l. c. p. 62.—c.p. 2840.

HAB. Common in damp, grassy places, in the hotter parts of the island.

3. **CANSCORA**, Lam.

1. **C. diffusa**, R. Br. (Griseb. in DC. Prod. ix. p. 64, cum syn.)—c.p. 2841.

HAB. Central Province, at an elevation of 2000 to 4000 feet.

2. **C. decussata**, Rœm. et Sch. (Griseb. l. c. cum syn.)—c.p. 2400.

HAB. Not uncommon in the Central Province, on the borders of paddy-fields, up to an elevation of 4000 feet.

3. **C. sessiliflora**, Rœm. et Sch. (Griseb. l. c. cum syn.)—c.p. 3581.

HAB. Batticaloa District; not common.

4. **C. perfoliata**, Lam. (Griseb. l. c. p. 65, cum syn.; Wight, Ic. t. 1327, f. 2.)—c.p. 1874.

HAB. North of the island, *Gardner*.

4. **SLEVOGTIA**, Reich.

1. **S. orientalis**, Griseb. in DC. Prod. ix. p. 65, cum syn.—c.p. 1871.

HAB. Hotter parts of the island, especially near the sea.

5. **GENTIANA**, Tournef.

1. **G. pedicellata**, Wall. (Griseb. in DC. Prod. ix. p. 107, cum syn.; Wight, Ic. t. 1328.)—*G. Zeylanica*, Griseb. l. c. p. 108.—c.p. 1873.

HAB. Abundant amongst grass, at Newera Ellia and other of the most elevated parts of the island.

6. **CRAWFURDIA**, Wall.

*Tripterospermum*, Blume.

1. **C. fasciculata**, Wall. (Griseb. in DC. Prod. ix. p. 120, cum syn.; Hook. Bot. Mag. t. 4838.)—Var.  $\beta$  *Championii*; *Tripterospermum Championii*, Gardn. MS.—c.p. 137.

HAB. Horton Plains, in the Central Province, at an elevation of 7000 feet.

*Corolla* cylindrico-campanulata, 1 poll. longa, 6 lin. in exp., omnino flavo-viridis vel margine plicisque purpureis. *Fructus* carnosus, ruber, stipite æquilongo concolore. *Placentæ* duæ. *Semina* triserialia, libera, acute trigona, brunnea.

The present plant corresponds in so many respects with the figure and description of *C. fasciculata*, that there can scarcely be a doubt of its being a variety of that species. It has, however, essentially the characters of the genus *Tripterospermum* of Blume, and I suspect that the fruit of *Crawfurdia* must have been inaccurately described from dried specimens.

## 7. OPHELIA, Don.

1. **O. Zeylanica**, Griseb. in DC. Prod. ix. p. 124.—C.P. 1875.

HAB. Central Province, in grassy places, at an elevation of 6000 to 7000 feet.

*Corolla* cyaneo-purpurea, *foveis* orbicularibus ad marginem longe radiato-fimbriatis.

## TRIBE II. MENYANTHÆ.

## 8. LIMNANTHEMUM, Gmelin.

(Nom. vulg. "Oloo," "Ambala.")

1. **L. Indicum**.—*Menyanthes Indica*, Linn. Sp. Pl. cum syn. (non Griseb.)—C.P. 3152.

Var. *α. stamineum*; *stylo* staminibus multo brevior.—*L. Kleinianum*, Griseb. in DC. Prod. ix. p. 139, cum syn.

Var. *β. stylosum*; *stylo* linearis, staminibus longior.—*L. Wightianum*, Griseb. l. c.

HAB. Common in the warmer parts of the island.

Between the forms *α* and *β* of the present plant I can find no difference, except in the relative length of the stamens and style; and, as there are instances of an exactly corresponding kind amongst the *Rubiaceæ* and *Erythroxyloæ*, and I believe also in the *Oxalidaceæ*, this difference is scarcely sufficient to found a specific character upon. The seeds in both forms are delicately muricated, or frequently nearly smooth. As regards the figure in Rumph. Amb. lib. xi. t. 72. f. 3, quoted by Linnæus, it would seem to me that, taken in connection with the description in page 173 of the same volume, there can be but little doubt of its being intended for the plant now under consideration, although Grisebach refers it to a species closely allied to, if it indeed be more than a state of *L. cristatum*.

2. **L. cristatum**, Griseb. l. c. cum syn.; sed Rheede, Hort. Mal. xi. p. 57. t. 29 (non p. 55. t. 28.)—An *L. Indicum*, Griseb. l. c. quoad descript. p.—C.P. 1870.

HAB. Abundant in the hotter parts of the island.

3. **L. Moonii**, Thw.; *foliis* parvis; *radicalibus* (*sterilibus*) deltoideis vel trapezoideis, *petiolo* plus minus elongato; *caulinis* (*floriferis*) cordato-orbiculatis vel obovoideis; *pedicellis* 2-5-nis; *floribus* parvis, 4-meris; *corollæ lobis* fimbriatis, disco versus apicem sparse pilosis; *glandulis* minutis, penicillato-pilosis; *seminibus* minute muricatis vel sublævibus.—*Menyanthes campestris*, Moon's Cat. p. 13. An *Villarsia parrifolia*, Wall. Cat. p.—C.P. 2842.

HAB. Not uncommon in the south of the island, growing in the water or on wet ground.

*Folia*  $\frac{1}{2}$ -1 poll. in diam. *Flores* albi, circiter 4 lin. in exp.

Nearly allied to the preceding species.

4. **L. biflorum**, Thw.; *caulibus* elongatis, simplicibus, v. parce ramosis plurinodiis; *nodis* monophyllis; *foliis* cordato-orbicularibus, subtus glanduloso-punctatis; *petiolo* longiuseculo, basi vaginante; *pedicellis* elongatulis, sæpius binis, in axilla bracteæ oblongæ, sub-acutæ, folio oppositæ; *calycis segmentis* ovatis, acutis; *corolla* glabra, lutea, calyce duplo longiore, ad faucem glandulis 5 minutis pilosis staminibus alternantibus munita; *lobis* margine fimbriatis; *capsula*

polysperma; *seminibus* subsphæricis, opacis, punctulatis.—*Menyanthes biflora*, Moon's Cat. p. 13. An *L. Forbesianum*, Griseb. l. c. ?—C.P. 1869.

HAB. South of the island.

This is probably *L. Forbesianum*, Griseb., in part; but it does not quite agree with the characters given of the flower of that species; nor does the author notice the very different habit the present plant exhibits to that of the other species of the section, the stem simply elongating, at each innovation, by the production of a single shoot at its apex, instead of producing, as in the other species (in addition to the fascicle of flower-pedicels), a sessile offset developing its own roots, and a succession of petioled, floriferous leaves, (perhaps more correctly termed petiole-like stems,) each terminated by a single leaf.

## CVII. BIGNONIACEÆ.

### 1. CALOSANTHES, Blume.

1. **C. Indica**, Blume, (DC. Prod. ix. p. 177, cum syn.; Wight, Ic. t. 1337, 1338.)—C.P. 3171.

HAB. Very common, up to an elevation of 2000 feet. Nom. vulg. "Totilla-gass."

### 2. SPATHODEA, Beauv.

1. **S. Rheedii**, Wall. (DC. Prod. ix. p. 206, cum syn.; Wight, Ic. t. 1339.)—C.P. 3170.

HAB. Hotter parts of the island. Nom. vulg. "Deya-danga-gass."

### 3. STEREOSPERMUM, Cham.

1. **S. chelonoides**, DC. Prod. ix. p. 210, cum syn.; Wight, Ic. t. 1341.—C.P. 1959.

HAB. Very abundant near the sea, and occurring up to an elevation of 2000 feet. Nom. vulg. "Loonoo-madala-gass."

*S. suaveolens*, DC. (C.P. 1960), is sometimes to be found in the neighbourhood of Buddhist temples, in the south of the island, but I cannot hear of its occurring truly wild. Its roots are much valued by the natives as a tonic medicine, and they attribute the same properties and give the same name ("Palol") to those of *Spathodea adenophylla*, which is occasionally found in gardens.

*Millingtonia hortensis* is very common, but in or near cultivated ground, and I have no evidence of its being really wild.

## CVIII. CYRTANDRACEÆ.

### 1. ÆSCHYNANTHUS, Jack.

1. **Æ. Zeylanica**, Gardn. in Calcutta Journ. vol. vi. p. 474; Wight, Ic. t. 1347.—An *Æsch. Perrottetii*, A. DC. Prod. ix. p. 261, var. ?—C.P. 1782.

HAB. Central Province, at an elevation of 4000 to 6000 feet.

My specimens differ from the description of *Æ. Perrottetii* principally in the leaves being narrower, and acute at the base, and in the corolla being externally sprinkled with glandular hairs.

## 2. DIDYMOCARPUS, Wall.

1. **D. Zeylanica**, R. Br. in Pl. Jav. Rar. p. 119; DC. Prod. ix. p. 266.—*D. longipetiolata*, Gardn. in Calc. Journ. vi. p. 475; Walp. Ann. iii. p. 96.—c.p. 352 (395, 1783).

HAB. Ambagamowa District, near Adam's Peak, at an elevation of 5000 to 6000 feet.

*Folia* basi cordata vel rarius subacuta; *petiolo* brevior, æquilongus vel 2-3-plo longior, basi sæpe rufescente. *Pedunculi pedicellique* rubri. *Calyx* flavo-rufescens, *lobis* lineari-lanceolatis, obtusis. *Corolla* alba, *tubo* cyaneo-purpureo. *Stamina sterilia* clavata. *Stigma* oblique orbiculare, album.

2. **D. Humboldtiana**, Gardn. l. c. p. 477; Walp. l. c.; Hook. Bot. Mag. t. 4757.—c.p. 1784.

Var.  $\beta$ . *D. primulaefolia*, Gardn. l. c. p. 478; Walp. l. c.; Hook. Bot. Mag. t. 5161.—c.p. 1785.

HAB. Common in damp, rocky places in the Central Province and elsewhere, up to an elevation of 5000 feet.

This is a most variable species, and there appears to me to be no permanent character to distinguish *D. primulaefolia* from *D. Humboldtiana*; and I think it is not improbable that *D. tomentosa*, *D. ovalifolia*, and *D. Rottleriana*, are some or all of them, varieties of the present plant.

3. **D. floccosa**, Thw.; subacaulis; *foliis* 3-4-nis, ovatis, crenulatis, superne tomentosis, subtus pallide fulvo-floccosis, apice rotundatis vel parum acutis, basi in *petiolum* alatum, plus minus elongatum angustatis; *cymis* compactis *pedunculoque* floccoso-tomentosis; *lobis calycinis* linearibus, subacutis; *corollæ tubo* valde ventricoso, ore contracto; *limbo* patulo, subæqualiter rotundatim 5-lobato; *ovario* glandulis sessilibus pilisque brevibus consperso.—c.p. 3368.

HAB. Dolosbage, in the Central Province, at an elevation of 2000 feet.

*Flores* pallide purpurei, 6-7 lin. in exp.

This is principally distinguished from the preceding by the different character of its pubescence, its larger foliage, more compact cymes, and the more contracted throat of the corolla.

## 3. CHIRITA, Ham.

1. **C. Moonii**, Gardn. in Calc. Journ. vi. p. 479; Hook. Bot. Mag. t. 4405; Walp. Ann. iii. p. 98.—*Martynia lanceolata*, Moon's Cat. p. 45.—c.p. 1789.

HAB. Four Corles, Moon; Hantani, at an elevation of 3000 to 4000 feet.

2. **C. Walkeri**, Gardn. l. c. p. 480; Hook. Bot. Mag. t. 4327; Walp. l. c.—c.p. 2843.

Var.  $\beta$ ; minus pubescens; *floribus* minoribus.—c.p. 542.

HAB. Var.  $\alpha$ . Raxawa and Newera Ellia, up to an elevation of 7000 feet. Var.  $\beta$ . Reigam Corle, at no great elevation.

Var.  $\beta$ . *Calyx* 3-4 lin. longus. *Corolla* pollicaris, pallide purpurea. *Antheræ* barbatae (sicut in var.  $\alpha$ ).

3. **C. Zeylanica**, Hook. Bot. Mag. t. 4182; Walp. Rep. vi. p. 522.—*C. communis*, Gardn. l. c. p. 481; Walp. Ann. iii. p. 98.—c.p. 1788.

Var.  $\beta$ ; *foliis lanceolatis, utrinque acutis, minus pubescentibus; venis primariis paucioribus; floribus parum minoribus pallidioribusque*.—c.p. 3437.

HAB. Var. *a*. Common in damp forests, up to 5000 feet. Var.  $\beta$ . At the edge of a rivulet in the Singherajah Forest between Galle and Ratnapoora.

#### 4. **KLUGIA**, Schlecht.

1. **K. Notoniana**, DC. Prod. ix. p. 276, cum syn.; Wight, Ic. t. 1353; Gardn. in Calc. Journ. vi. p. 487.—*K. glabra*, Gardn. l. c. p. 489.—c.p. 1787, 3369.

HAB. Common in the Central Province, on rocks and banks in running streams, up to an elevation of 6000 feet. Nom. vulg. "Deyanella."

2. **K. Zeylanica**, DC. l. c.; Gardn. l. c. p. 490; Hook. Bot. Mag. t. 4620.—c.p. 1786.

HAB. In similar situations with the last.

Præcedenti valde affinis, a qua *foliis floribusque minoribus, et lobis calycinis inter se æqualiter carinatis, solum differre videtur*. Varietates forsan species unice.

#### 5. **EPITHEMA**, Blume.

1. **E. carnosum**, Benth. (DC. Prod. ix. p. 288.)—*E. Zeylanica*, Gardn. in Calc. Journ. vi. p. 492; Wight, Ic. t. 1354.—c.p. 2844.

HAB. Common on damp rocks, in the Central Province, up to an elevation of 5000 feet.

*Corolla pallide cyaneo-purpurea.*

#### 6. **ISANTHERA**, Nees.

1. **I. permollis**, Nees in Trans. Linn. Soc. xvii. p. 82; DC. Prod. ix. p. 279; Wight, Ic. t. 1355.—*I. floribunda*, Gardn. in Calc. Journ. vi. p. 483; Walp. Ann. iii. p. 99.—c.p. 1670.

HAB. In damp forests, up to an elevation of 4000 feet.

*Corolla alba. Antheræ flavæ.*

#### 7. **CHAMPIONIA**, Gardn.

(Calc. Journ. vi. p. 485; Walp. Ann. iii. p. 96.)

1. **C. reticulata**, Gardn. l. c.; Walp. l. c.—c.p. 358.

HAB. Saffragam District, at an elevation of about 3000 feet; not uncommon.

*Caulis erectus, 6-12-pollicaris. Folia 2-7 poll. longa,  $\frac{3}{4}$ -1 $\frac{1}{2}$  poll. lata, petiolo  $\frac{1}{2}$ -1 $\frac{3}{4}$  poll. longo. Pedunculæ sæpe folio longiores, solitarii vel bini. Flores 4-5 lin. in exp.*

## CIX. SESAMEÆ.

## 1. SESAMUM, Linn.

1. **S. Indicum**, Linn. (DC. Prod. ix. p. 250, cum syn.; Wight, Illust. ii. t. 163.)—c.p. 2023.

HAB. Jaffna, *Gardner*; Trincomale, *Rev. S. O. Glenie*. Nom. vulg. "Tel-talla."

Cultivated in the hot, drier parts of the island, for the sake of the oil yielded by the seeds.

## 2. PEDALIUM, D. Roy.

1. **P. Murex**, Linn. (DC. Prod. ix. p. 256.)—c.p. 1779.

HAB. Common near the sea. Nom. vulg. "At-nairenchee."

MARTYNIA DIANDRA has become quite naturalized in the north of the island, on the coast.

## CX. HYDROLEACEÆ.

## 1. HYDROLEA, Linn.

1. **H. Zeylanica**, Vahl, (Choisy in DC. Prod. x. p. 180, cum syn.; Wight, Ic. t. 601.)—c.p. 1883.

Var.  $\beta$ . *ciliata*, Choisy, l. c. cum syn.—c.p. 1884.

HAB. Very common in damp places, in the hotter parts of the island. Nom. vulg. "Deya-kirilla."

## CXI. CONVOLVULACEÆ.

## 1. RIVEA, Choisy.

1. **R. tiliæfolia**, Choisy, in DC. Prod. ix. p. 325, cum syn.—*Argyreia tiliæfolia*, Wight, Ic. t. 1358.—c.p. 2848.

HAB. Very abundant up to an elevation of 3000 feet. Nom. vulg. "Mā-banda."

2. **R. Zeylanica**, Thw.—*Ipomæa Zeylanica*, Gært. de Fruct. ii. p. 482. t. 178; Moon's Cat. p. 14.

Var.  $\alpha$ . *populifolia*.—*Argyreia populifolia*, Choisy, l. c. p. 329.—c.p. 1943.

Var.  $\beta$ . *hirsuta*.—*R. hirsuta*, Wight, Ic. t. 891. *Argyreia hirsuta*, W. et A.; Choisy, l. c. p. 330.—c.p. 1944.

(Var.  $\gamma$ . *emarginata*.—*R. pomacea*, Wight, Ic. t. 888. *Argyreia pomacea*, Choisy, l. c.)

HAB. Varr.  $\alpha$  and  $\beta$ . Very abundant, up to an elevation of 5000 feet. Nom. vulg. "Girri-tilla."

I have no Ceylon specimens of var.  $\gamma$ , but it doubtless occurs in the island.

3. **R. ornata**, Choisy, l. c. p. 326, cum syn.; Wight, Ic. t. 1356.—c.p. 1946.

HAB. Hot, drier parts of the island.



## 2. ARGYREIA, Lour.

1. *A. speciosa*, Sweet, (Choisy in DC. Prod. ix. p. 328, cum syn.)—c.p. 2849.

HAB. Central Province, *Mr. H. de Alwis*. Nom. vulg. "Mahadoomooda."

2. *A. Leschenaultii*, Choisy, l. c. p. 329, cum syn.—c.p. 1932.

HAB. Batticaloa, *Gardner*.

3. *A. splendens*, Sweet, (Choisy, l. c. cum syn.)—c.p. 2845.

HAB. Hantani District, *Gardner*.

4. *A. elliptica*, Choisy, l. c. p. 330, cum syn.—c.p. 1933.

HAB. Central Province; not uncommon.

5. *A. hancorniaefolia*, Gardn. MS.; *foliis* glabris, ellipticis vel ovatis, acuminatis, arcte parallelim venosis, basi rotundatis; *pedunculis* solitariis, unifloris, axillaribus terminalibusque, *petiolo* brevioribus, medio bibracteatis; *bracteis* valde caducis; *fructibus* carnosis, obovato-oblongis, apiculatis.—c.p. 1924.

HAB. Central Province, up to an elevation of 5000 feet.

*Ramuli juniores*, cum *petiolis pedunculisque*, adpresse fulvo-pilosi. *Folia* subcoriacea, 2-3½ poll. longa, 1¼-1¾ poll. lata, *petiolo* tenui, ¾-1 poll. longo. *Sepala* ovata, subacuta, 2-3 lin. longa. *Corolla* purpurea, tubo parum inflato, limbo 1½ poll. in exp.

6. *A. aggregata*, Choisy, l. c. p. 333, cum syn.—*Lettsomia aggregata*, Roxb. Wight, Ic. t. 1359.—c.p. 1945.

HAB. Abundant in the hot, drier parts of the island.

## BATATAS, Rumph.

1. *B. paniculata*, Choisy, in DC. Prod. ix. p. 339, cum syn.—c.p. 499.

HAB. Colombo, *Mr. W. Ferguson*. Nom. vulg. "Kirri-baddoo."

2. *B. Choisyana*, Wight, Ic. t. 491.—An *Convolvulus atropurpureus*, Wall. Fl. Ind. ii. p. 54; Plant. As. Rar. p. 38. t. 43?—c.p. 3647.

HAB. Batticaloa District.

The only specimen of this plant I possess agrees very well with Wight's figure, but the shape of the corolla would appear to be so very unlike that of Wallich's plant, that, taking the difference of locality also into consideration, I suspect they must be distinct, though Choisy, in DC. Prod. p. 366, describes them as one species, under the name of *Ipomœa atropurpurea*.

BATATAS EDULIS, Choisy, is abundantly cultivated as a vegetable. *B. pentaphylla* is common in gardens, and *Quamoclit coccinea* and *Q. vulgaris* are also common garden plants.

## 4. PHARBITIS, Choisy.

1. *P. Nil*, Choisy, in DC. Prod. ix. p. 343, cum syn.—c.p. 1938.

HAB. Hot, drier parts of the island.

PHARBITIS LEARII, Hook., has no claim to be considered indigenous to Ceylon.

## 5. CALONYCTION, Choisy.

1. *C. speciosum*, Choisy, in DC. Prod. ix. p. 345, cum syn.; Wight, Ic. t. 1361.—c.p. 3648.

HAB. Abundant in the warmer parts of the island. Nom. vulg. "Alanga."

A plant (c.p. 3580), closely allied to the present, though probably a distinct species, having longer, more fleshy pedicels, and smaller and purple flowers, is cultivated by the Cinghalese, who eat the pedicels in their curries. It is said to be not truly indigenous.

2. *C. ? comospermum*, Boj. (Choisy, l. c. p. 346.)—c.p. 3536.

HAB. Jaffna, Gardner; Dichwelle, near Matura, close to the sea.

*Planta* tota glabra. *Pedunculi* uniflori. *Sepala* obtusa, mutica vel mucronulata, demum pollicaria. *Corolla* magna, alba. *Semina* nigra, semipollicaria, margine longiuscule lanata.

Can this be the *Rivea hypocrateriformis*, Choisy, as regards the Trincomale plant? It appears to me to be a true species of *Ipomœa*.

## 6. IPOMŒA, Linn.

1. *I. reptans*, Poir. (Choisy, in DC. Prod. ix. p. 349, cum syn.)—c.p. 3535.

HAB. On the borders of tanks in the hot, drier parts of the island; also cultivated as a vegetable by the Cinghalese. Nom. vulg. "Kangkoong."

2. *I. Pes-Caprae*, Sw. (Choisy, l. c. cum syn.)—c.p. 1934.

HAB. Very abundant near the sea. Nom. vulg. "Moodoo-bin-tambooroo."

3. *I. rugosa*, Choisy, l. c. p. 350, cum syn.—c.p. 2854.

HAB. Very abundant in damp fields. Nom. vulg. "Bin-tambooroo."

4. *I. reniformis*, Choisy, l. c. p. 351, cum syn.—c.p. 3579.

HAB. Hot, drier parts of the island.

5. *I. littoralis*, Blume, (Choisy, l. c. p. 352.)—c.p. 1939.

HAB. Galle, near the sea.

6. *I. tridentata*, Roth, (Choisy, l. c. p. 353, cum syn.)—c.p. 1929.

Var.  $\beta$ . *I. filicaulis*, Blume, (Choisy, l. c. cum syn.)—c.p. 1930.

HAB. Very abundant in the warmer parts of the island. Nom. vulg. "Hawaree-maddoo."

The two varieties of this plant pass imperceptibly into one another, so that it is often impossible to determine to which of the two some specimens should be referred.

7. *I. campanulata*, Linn. (Choisy, l. c. p. 359, cum syn.; Wight, Ic. t. 1375.)—c.p. 2847.

HAB. Warmer parts of the island; not uncommon.

In general appearance this species closely resembles the more common *Rivea tiliifolia*, but the flowers are handsomer, the leaves less deeply cordate, and the fruit very different.

8. *I. jucunda*, Thw.; *foliis* rotundato-cordatis, longiuscule acuminatis, superne glabris, subtus tomentellis glanduloso-punctulatisque; *pedunculis* unifloris, petiolo subæquilongis; *sepalis* oblongis, subacutis,

externe plus minus pilosis; *seminibus* pilis sericeis longis olivaceis omnino dense tectis.—C.P. 3448.

HAB. Dolosbage District; rare.

Ingens, alte scandens. *Capsula* ovoidea,  $\frac{3}{4}$  poll. longa, *styli* base apiculata, *sepalis* pollicaribus parum reflexis suffulta. *Corollam* nondum vidi.

9. **I. Turpethum**, R. Br. (Choisy, l. c. p. 360, cum syn.)—*I. anceps*, R. et Sch.; Choisy, l. c.—C.P. 3649.

HAB. Hotter parts of the island. Nom. vulg. "Trasta-waaloo."

The roots of this plant are employed by the Cinghalese as a cathartic.

10. **I. Pes-Tigridis**, Linn. (Choisy, l. c. p. 363, cum syn.; Wight, Ic. t. 836.)—C.P. 1942.

Var.  $\beta$ . *I. hepaticifolia*, Linn.; Choisy, l. c. cum syn.

HAB. Very abundant in the hotter parts of the island. Var.  $\beta$ . Colombo, Mr. W. Ferguson. Nom. vulg. "Diwi-addeeya."

11. **I. Wightii**, Choisy, l. c. p. 364, cum syn.; Wight, Ic. t. 1364.—C.P. 2851.

HAB. Central Province, at an elevation of 2000 to 3000 feet.

12. **I. pileata**, Roxb. (Choisy, l. c. p. 365, cum syn.; Wight, Ic. t. 1363.)—C.P. 3501.

HAB. Peradenia; not common.

13. **I. sessiliflora**, Roth, (Choisy, l. c. p. 366, cum syn.; Wight, Ic. t. 169.)—C.P. 3107.

HAB. Central Province, up to an elevation of 4000 feet.

14. **I. obscura**, Bot. Reg. (Choisy, l. c. p. 370, cum syn.)—C.P. 1935.

HAB. Common up to an elevation of 3000 feet.

15. **I. sepiaria**, König, (Choisy, l. c. cum syn.; Wight, Ic. t. 838.)—C.P. 1937.

Var.  $\beta$ . *sagittata*.—C.P. 2859.

HAB. Common in the warmer parts of the island.

16. **I. cymosa**, R. et Sch. (Choisy, l. c. p. 371, cum syn.)—C.P. 1931.

HAB. Common throughout the island. Nom. vulg. "Maddoo."

17. **I. chryseides**, Bot. Reg. (Choisy, l. c. p. 382, cum syn.; Wight, Ic. t. 157.)—C.P. 1940.

HAB. Warmer parts of the island; not common.

18. **I. Coptica**, Roth, (Choisy, l. c. p. 384, cum syn.)—C.P. 1928.

HAB. North of the island, Gardner.

19. **I. palmata**, Försk. (Choisy, l. c. p. 386, cum syn.)—*I. pulchella*, Roth; Choisy, l. c.; Wight, Ic. t. 156.)—C.P. 3387.

HAB. Ambagamowa District.

## 7. ANISEIA, Choisy.

1. **A. uniflora**, Choisy, in DC. Prod. ix. p. 431, cum syn.; Wight, Ic. t. 850.—C.P. 3006.

HAB. Warmer parts of the island.

## 8. SHUTEREIA, Choisy.

1. **S. bicolor**, Choisy, in DC. Prod. ix. p. 435, cum syn.; Wight, Ic. t. 835.—C.P. 1970.

HAB. Common in the hot, drier parts of the island.

### 9. BREWERIA, R. Br.

1. **B. Roxburghii**, Choisy, in DC. Prod. ix. p. 438, cum syn.; Wight, Ic. t. 1370.—C.P. 2853.

HAB. South of the island; not uncommon.

### 10. CRESSA, Linn.

1. **C. Cretica**, Linn. ( $\beta$  *Indica*; Choisy, in DC. Prod. ix. p. 440.)—C.P. 1925.

HAB. North of the island, *Gardner*.

### 11. EVOLVULUS, Linn.

1. **E. alsinoides**, Linn. (Choisy in DC. Prod. ix. p. 447, cum syn.)—C.P. 1926.

HAB. Common throughout the island, in hot, dry places. Nom. vulg. "Visnoo-Kraanta."

#### TRIBE CUSCUTEÆ.

### 12. CUSCUTA, Tourn.

1. **C. reflexa**, Roxb. (Choisy in DC. Prod. ix. p. 454, cum syn.; Hook. Ex. Fl. t. 150.)—C.P. 1922.

HAB. Upon stems of, apparently, *Wendlandia Notoniana*, Newera Ellia, *Gardner*.

2. **C. Chinensis**, Lam. (Choisy, l. c. p. 457, cum syn.; Wight, Ic. t. 1373.)—C.P. 611.

HAB. Colombo, on stems of *Achyranthes aspera*.

## CXII. ERYCIBEÆ.

### 1. ERYCIBE, Roxb.

1. **E. paniculata**, Roxb. (DC. Prod. ix. p. 464, cum syn.)—C.P. 1881.

HAB. Common in the warmer parts of the island, up to an elevation of 3000 feet. Nom. vulg. "Atta-meeriya."

## CXIII. BORAGINACEÆ.

#### TRIBE CORDIÆÆ.

### 1. CORDIA, Plum.

#### Sectio *Myxa*, Endl.

1. **C. Myxa**, Linn. (DC. Prod. ix. p. 479, cum syn.; Wight, Illust. ii. t. 169.)—C.P. 3650.

Var.  $\beta$  *minor*; *foliis fructibusque minoribus*.—c.p. 1880. An species distincta?

HAB. Var.  $\alpha$ . Central Province, etc. Var.  $\beta$ . Hot, drier parts of the island. Nom. vulg. "Loloo."

Var.  $\beta$  bears a very great general resemblance to var.  $\alpha$ , but a more careful examination of fresh specimens may possibly prove it to be distinct. With the materials I have at present, I can discover no essential difference between the two plants.

2. **C. oblongifolia**, Thw.; *foliis glabris, oblongis, brevi-acuminatis, minutissime reticulato-venulosis, basi subacutis, longiuscule petiolatis; paniculis terminalibus, parce pilosis; calyce cylindrico, 3-5-dentato, interne villosa; corolla 5-7-loba; lobis oblongis, reflexis; tubo calyce æquilongo; fructibus majusculis, rotundatis, acuminatis*.—c.p. 2696.

HAB. Central Province, up to an elevation of 2000 feet.

Subscandens. *Ramuli teretes, glabri. Folia 2-5 poll. longa, 1-2½ poll. lata, petiolo pollicari, e basi tumida, subhamata oriente et cum eadem articulato. Calyx 4 lin. longus. Stylus profunde bis bifidus. Fructus subpollicaris, calyce aucto sub-integro suffultus.*

3. **C. diversa**, Thw.; *foliis ovatis, parum acuminatis, basi rotundatis, trinerviis, superne scaberrimis, subtus fusco-tomentosis; paniculis terminalibus, paucifloris, tomentosis; calyce obconico*.—c.p. 1954.

HAB. Hot, drier parts of the island.

*Ramuli juniores tomentosi. Folia 2-4 poll. longa, 1-2½ poll. lata, petiolo ½-1 poll. longo. Calyx 3 lin. longus, 3-4-dentatus, externe præcipue versus apicem tomentosus, interne glaber. Corolla glabra, calyce parum longior, 5-loba. Stylus profunde bis bifidus. Fructus rotundatus, semipollicaris, calyce aucto scabro irregulariter dentato suffultus.*

### TRIBE EHRETIEÆ.

#### 2. EHRETIA, Linn.

1. **E. lævis**, Roxb. (DC. Prod. ix. p. 505, cum syn.; Wight, Ic. t. 1382.)—c.p. 1879.

HAB. Common in the drier parts of the island.

2. **E. buxifolia**, Roxb. (DC. Prod. p. 509, cum syn.)—c.p. 1885.

HAB. Abundant in the hot, drier parts of the island. Nom. vulg. "Heen-tambala."

#### 3. RHABDIA, Martius.

1. **R. viminea**.—*Ehretia? viminea*, Wall.; DC. Prod. ix. p. 509. *E. (xeroderma) cuneata*, Wight, Ic. t. 1385.—c.p. 3491.

HAB. Ambagamowa District; not common.

*Flores roseo-purpurei. Stigma bilobum.*

#### 4. TOURNEFORTIA, Linn.

1. **T. argentea**, Linn. f. (DC. Prod. ix. p. 514, cum syn.)—c.p. 1882.

HAB. South of the island, close to the sea.

#### Sectio *Tetrandra*, DC.

2. **T. Wallichii**, DC. l. c. p. 527, cum syn.—c.p. 2697.

HAB. Southern and Central Provinces, up to an elevation of 2000 feet.

*Corolla* pallide flavo-viridis. *Fructus* carnosus, ovatus, glaber, circiter 2 lin. longus, albido-rufescens.

The sessile, tetramerous flowers distinguish this from the *T. reticosa*, Wight, which in general appearance it very much resembles.

### TRIBE HELIOTROPIEÆ.

#### 5. HELIOTROPIUM, Tourn.

1. *H. supinum*, Linn.— $\beta$  *Malabaricum*, Benth.; DC. Prod. ix. p. 533; Wight, Ic. t. 1387.—c.p. 2854.

HAB. North of the island, *Gardner*.

2. *H. marifolium*, Retz, (DC. l. c. p. 547; Wight, Ic. t. 1390.)—An *H. scabrum*, et *H. Rottleri*, Wight, Ic. tt. 1389, 1392?—c.p. 1889, 1891.

HAB. Common on the borders of paddy-fields, in the hotter parts of the island.

A very variable plant, and I suspect that Dr. Wight's three figures, quoted above, represent three forms only of it.

3. *H. linifolium*, Lehm. (DC. l. c. cum syn.; Wight, Ic. t. 1391.)—c.p. 1886.

HAB. Jaffna, *Gardner*.

I have not met with any plant in the island corresponding with the figure of *H. Zeylanicum*, Burm. Fl. Ind. t. 16. f. 2.

#### 6. HELIOPHYTUM, DC.

##### Sectio *Tiaridium*.

1. *H. Indicum*, DC. Prod. ix. p. 556, cum syn.—*Tiaridium Indicum*, Lehm.; Wight, Illust. ii. t. 171.—c.p. 1893.

Var.  $\beta$ . *foliis* minoribus; *corollæ tubo* longiore quam in var. *a*.—c.p. 1894.

HAB. Common in the warmer parts of the island. Var.  $\beta$ . Bintenne, *Gardner*.

#### 7. COLDENIA, Linn.

1. *C. procumbens*, Linn. (DC. Prod. ix. p. 558, cum syn.)—c.p. 1890.

HAB. Common in the hot, drier parts of the island.

### TRIBE BORAGEÆ.

#### 8. CYNOGLOSSUM, Tourn.

1. *C. micranthum*, Desf. (DC. Prod. x. p. 149, cum syn.)—*C. ovatum*, Moon's Cat.—c.p. 143.

HAB. Common in waste places throughout the island. Nom. vulg. "Boo-kattoo-handa."

2. *C. furcatum*, Wall. (DC. l. c. cum syn.; Wight, Ic. t. 1395.)—*C. decurrens*?, Moon's Cat. p. 12.—c.p. 1892.

HAB. Common in the most elevated parts of the island.

*Folia caulina* semiamplexicaulia vel plus minus decurrentia. *Corolla* cyaneo-purpurea.

As remarked by Dr. Wight, the present plant is very closely allied to *C. micranthum*, and the two may possibly prove to be varieties only of one species.

### 9. TRICHODESMA, R. Br.

1. **T. Indicum**, R. Br. (DC. Prod. x. p. 172, cum syn.; Wight, *Illust.* ii. t. 172.)—C.P. 1887.

HAB. Jaffna, *Gardner*; Trincomale, *Rev. S. O. Glenie*.

2. **T. Zeylanicum**, R. Br. (DC. l. c.)—C.P. 1888.

HAB. Hot, drier parts of the island.

## CXIV. SOLANACEÆ.

### 1. SOLANUM, Tourn.

1. **S. nigrum**, Linn.—*S. rubrum*, Roxb. Wight, *Ic.* t. 344.—C.P. 2865.

HAB. Common throughout the island. Nom. vulg. “Kaloo-kang-waireya.”

2. **S. verbascifolium**, Linn. (Duval in DC. Prod. xiii. p. 114, cum syn.; Wight, *Ic.* t. 1398.)—C.P. 1902.

HAB. Common in the Central Province. Nom. vulg. “Hakkarilla.”

3. **S. auriculatum**, Ait. (Dun. l. c. p. 115, cum syn.)—C.P. 3525.

HAB. Common in cultivated ground, and also called “Hakkarilla.”

4. **S. pubescens**, Willd. (Wight, *Ic.* t. 1402.)—C.P. 1901.

HAB. Hot, drier parts of the island.

5. **S. membranaceum**, Wall. (Dun. l. c. p. 143.)—C.P. 1900.

HAB. Central Province, up to an elevation of 4000 feet.

6. **S. denticulatum**, Blume, (Dun. l. c. p. 181; Wight, *Ic.* t. 1397.)—*Var. calyce* 5-dentato.—C.P. 2864.

HAB. Not uncommon in the Central Province, at an elevation of 4000 to 5000 feet.

This so closely resembles, except in the fewer teeth of its calyx, Wight's figure quoted above, that I cannot venture to describe it as a distinct species. It is very nearly allied, too, to *S. membranaceum*, but is much more hairy, and has larger flowers and fruit: they may possibly, however, be varieties of one species.

7. **S. ferox**, Linn. (Dun. l. c. p. 255, cum syn.; Wight, *Ic.* t. 1399.)—C.P. 2868.

HAB. Common up to an elevation of 4000 feet. Nom. vulg. “Malla-battoo.”

The ripe fruit is eaten by the Cinghalese.

8. **S. giganteum**, Jacq. (Dun. l. c. p. 258, cum syn.; Wight, *Ic.* t. 893.)—C.P. 1903.

HAB. Matturatte and Rambodde, in the Central Province, at an elevation of 3000 to 5000 feet.

9. **S. torvum**, Linn. (Dun. l. c. p. 260, cum syn.; Wight, *Ic.* t. 345.)—C.P. 2867.

HAB. Common on waste ground in the Central Province.

10. **S. trilobatum**, Linn. (Dun. l. c. p. 287, cum syn.; Wight, Ic. t. 854.)—C.P. 1906.

HAB. Hot, drier parts of the island. Nom. vulg. "Wal-tibbootoo."

11. **S. xanthocarpum**, Schrad. et Wendl. (Dun. l. c. p. 302, cum syn.)—*S. diffusum*, Roxb. Fl. Ind. ii. p. 250.—C.P. 1907.

Var.  $\beta$ . *Jacquini*.—*S. Jacquini*, Willd.; Wight, Ic. t. 1401.—C.P. 1905.

HAB. Common throughout the warmer parts of the island. Nom. vulg. Var.  $\alpha$ . "Ella-battoo." Var.  $\beta$ . "Kattoo-wel-battoo."

The fruit of var.  $\alpha$  and  $\beta$  is eaten by the Cinghalese. Every part of var.  $\beta$  is used medicinally.

12. **S. Indicum**, Linn. (Nees; Dun. l. c. p. 309, cum syn.; Wight, Ic. t. 346.)—C.P. 1904.

HAB. Very abundant in waste places throughout the island. Nom. vulg. "Tibbootoo."

The unripe fruit is eaten, after being boiled, by the Cinghalese.

*Solanum esculentum*, Dun., and several species and varieties of *Capsicum*, are commonly cultivated as vegetables and condiments.

## 2. PHYSALIS, Linn.

(Nom. vulg. "Mottoo.")

1. **P. minima**, Linn. (Dunal in DC. Prod. xiii. p. 445, cum syn.)—*P. Indica*, Lam.; Dun. l. c. p. 443. *P. Hermannii*, Dun. l. c. p. 444.—C.P. 2866.

HAB. Common in waste and cultivated ground in the warmer parts of the island.

2. **P. angulata**, Linn. (Dun. l. c. p. 449, cum syn.)—C.P. 1898.

HAB. Frequent in similar situations with the last.

## 3. WITHANIA, Duval.

1. **W. somnifera**, Dun. in DC. Prod. xiii. p. 453, cum syn.; Wight, Ic. t. 853.—C.P. 3651.

HAB. Common in cultivated ground. Nom. vulg. "Amookkarā."

The whole plant is used as a medicine by the Cinghalese.

## 4. DATURA, Linn.

(Nom. vulg. "Attana.")

1. **D. fastuosa**, Linn. (Dun. in DC. Prod. xiii. p. 542, cum syn.; Wight, Ic. t. 1396.)—C.P. 1897.

HAB. Very common in waste and cultivated ground.

The flowers of this variable species are frequently quite white. May not *D. Metel*, Linn., be a variety only of the present plant?

## CXV. SCROPHULARIACEÆ.

### 1. CELSIA, Linn.

1. **C. Coromandeliana**, Vahl, (Bentham in DC. Prod. x. p. 246, cum syn.; Wight, Ic. t. 1406.)—C.P. 2021.



HAB. North of the island, *Gardner*.

## 2. LINDENBERGIA, Lehm.

1. **L. urticæfolia**, Lehm. (Benth. in DC. Prod. x. p. 377, cum syn.)

HAB. Ceylon, *Champion* (*Hb. Hook.*).

## 3. PTEROSTIGMA, Benth.

1. **P. villosum**, Benth. l. c. p. 380, cum syn.—c.p. 2041 (723, 2054).

HAB. South of the island, common.

2. **P. capitatum**, Benth. l. c. cum syn.—c.p. 2880.

HAB. South of the island. Nom. vulg. "Gona Kola."

The Cinghalese frequently chew the leaves of this plant with their betel.

## 4. LIMNOPHILA, R. Br.

1. **L. serrata**, Gaud. (Benth. l. c. p. 387, cum syn.)—*L. conferta*, Benth. l. c. cum syn. *L. micrantha*, Benth. l. c. cum syn.—c.p. 2049.

HAB. Very abundant in paddy-fields and wet places.

2. **L. hirsuta**, Benth. l. c. p. 388, cum syn.—An *L. villosa*, Bl., Benth. l. c. ? An *L. diffusæ*, G. Don, Benth. l. c. var. ?—c.p. 2040.

HAB. Common in similar situations with the last.

3. **L. punctata**, Bl. (Benth. l. c. p. 388, cum syn.)—*L. erecta*, Benth. l. c. cum syn.—c.p. 2035, 535.

HAB. Very common in wet places.

4. **L. gratioloides**, R. Br. (Benth. l. c. p. 389, cum syn.)—*L. Roxburghii*, G. Don; Benth. l. c. p. 388.—c.p. 2038.

HAB. Common in the hotter parts of the island.

5. **L. sessiliflora**, Bl. (Benth. l. c.)—*L. heterophylla*, Benth. l. c. p. 390, cum syn.—c.p. 2037, 2389 (2039).

HAB. Very abundant.

6. **L. racemosa**, Benth. l. c. p. 390, cum syn.; Wight, Ic. t. 861.—c.p. 2036, 2996.

HAB. Abundant in the warmer parts of the island.

## 5. HERPESTIS, Gærtn. f.

1. **H. floribunda**, R. Br. (Benth. in DC. Prod. x. p. 400, cum syn.)—c.p. 2881.

HAB. Cultura, *Moon*.

2. **H. Monnieria**, H.B.K. (Benth. l. c. cum syn.)—c.p. 2031.

HAB. Very abundant in wet places. Nom. vulg. "Loonoo Weela."

Used as a medicine for children.

## 6. DOPATRIUM, Ham.

1. **D. nudicaule**, Benth. in DC. Prod. x. p. 407, cum syn.—c.p. 2046.

HAB. Not uncommon in the warmer parts of the island.

2. **D. lobelioides**, Benth. l. c. cum syn. ; Wight, Ic. t. 859.—c.P. 2042 (2043).

HAB. Common in the hotter parts of the island, in wet places.

3. **D. junceum**, Ham. (Benth. l. c.)—c.P. 2044.

HAB. Very abundant.

This plant, notwithstanding the great difference in the size of the flowers, is perhaps scarcely specifically distinct from the preceding.

### 7. ARTANEMA, Don.

1. **A. sesamoides**, Benth. in DC. Prod. x. p. 408, cum syn. ; Wight Ic. t. 1410.—c.P. 2024.

HAB. Very common, up to an elevation of 3000 feet.

### 8. TORENIA, Linn.

1. **T. Asiatica**, Linn. (Benth. in DC. Prod. x. p. 410, cum syn.)—*T. hirsuta*, Ham. ; Benth. l. c. cum syn. ; Wight, Ic. t. 862.—c.P. 2032.

HAB. Very abundant.

A very variable species, and I suspect that the following is but a form of it.

2. **T. rubens**, Benth. l. c.—c.P. 2033.

HAB. Central Province, up to an elevation of 4000 feet.

### 9. VANDELLIA, Linn.

1. **V. crustacea**, Benth. in DC. Prod. x. p. 413, cum syn. ; Wight, Ic. t. 863.—c.P. 2048 (2052).

HAB. Very abundant in paddy-fields and other wet places.

2. **V. multiflora**, G. Don, (Benth. l. c. cum syn.)—*V. hirsuta*, Ham. ; Benth. l. c. cum syn.—c.P. 3309.

HAB. Near Kornegalle.

3. **V. scabra**, Benth. l. c. cum syn.—c.P. 2885.

HAB. Very common in moist ground.

4. **V. pedunculata**, Benth. l. c. cum syn.—c.P. 652.\*

HAB. Abundant in paddy-fields, etc.

5. **V. angustifolia**, Benth. l. c. cum syn.—c.P. 2884.

HAB. Cultura, *Moon*.

### 10. ILYSANTHES, Rafin.

1. **I. hyssopioides**, Benth. in DC. Prod. x. p. 419, cum syn. ; Wight, Ic. t. 857.—*I. parviflora*, Benth. l. c. cum syn. *I. rotundifolia*, Benth. l. c. p. 420, cum syn.—c.P. 2047, 3111.

HAB. Common in paddy-fields.

The plants I have ventured to unite under this species merge so completely into one another, that I cannot separate them.

### 11. BONNAYA, Link and Otto.

(Nom. vulg. "Weela.")

1. **B. veronicæfolia**, Spr. (Benth. in DC. Prod. x. p. 421, cum syn.)

\* c.P. 652 in Herb. Hook. is *Ilysanthes hyssopioides*, Benth.—ED.

—*B. grandiflora*, Spr.; Benth. l. c. cum syn. *B. verbenæfolia*, Spr.; Benth. l. c. cum syn.—c.p. 3084, 2050, 2051.

HAB. Abundant in paddy-fields, etc.

An extremely variable plant; and further investigation will probably determine the following as well as the other described species of this genus to be forms only of one.

2. *B. tenuifolia*, Spr. (Benth. l. c. p. 422.)—c.p. 2883.

HAB. Central Province, *Macrae*.

## 12. MICROCARPÆA, R. Br.

1. *M. muscosa*, R. Br. (Benth. in DC. Prod. x. p. 433, cum syn.)—c.p. 2728.

HAB. Not uncommon in moist ground.

## 13. BUCHNERA, Linn.

1. *B. hispida*, Ham. (Benth. l. c. p. 496, cum syn.; Wight, Ic. t. 1413.)

HAB. Ceylon, *Walker* (*Hb. Hook.*).

## 14. STRIGA, Linn.

1. *S. hirsuta*, Benth. in DC. Pr. x. p. 502, cum syn.—c.p. 2028, 2886.

HAB. Common up to an elevation of 4000 feet.

This species has larger flowers when growing on the hills, than when it occurs in the low country.

2. *S. euphrasioides*, Benth. l. c. p. 503, cum syn.—*Buchnera euphrasioides*, Wight, Ic. t. 855.—c.p. 2887.

Var.  $\beta$ ; minus *hirsuta*, *caule* elongato, *foliis floralibus* brevioribus, *calyce* 15-costato, denique fisso.—c.p. 2029 (2030, 2031).

HAB. Var. *a*. Bintenne, *Gardner*. Var.  $\beta$ . Very common in swampy places in the hotter parts of the island.

*Flores albi.*

## 15. SOPUBIA, Ham.

1. *S. delphinifolia*, G. Don, (Benth. in DC. Prod. x. p. 522, cum syn.)—c.p. 2022.

HAB. Common up to an elevation of 4000 feet.

2. *S. trifida*, Ham. (Benth. l. c. cum syn.)—c.p. 525.

HAB. Abundant at Newera Ellia and other of the more elevated parts of the Central Province.

## 16. CENTRANTHERA, R. Br.

1. *C. Brunoniana*, Wall. (Benth. in DC. Prod. x. p. 525.)—*C. procumbens*, Benth. l. c.—c.p. 2026 (2027).

HAB. Common in grassy wet places, up to an elevation of 3000 feet. Nom. vulg. "Dootoo-satootoo."

2. *C. hispida*, R. Br. (Benth. l. c. cum syn.)—c.p. 2025.

HAB. In similar situations with the preceding species, but not so common.

3. *C. humifusa*, Wall. (Benth. l. c. cum syn.)—C.P. 2888.

HAB. South of the island.

### 17. PEDICULARIS, Linn.

1. *P. Zeylanica*, Benth. in DC. x. p. 580; Wight, Ic. t. 1419.—C.P. 229.

HAB. Abundant at Newera Ellia, Horton Plains, and other of the more elevated parts of the Central Province.

## CXVI. OROBANCHACEÆ.

### 1. ÆGINETIA, Linn.

*Calyx* spathaceus vel tubulosus irregulariter fissusque. *Antheræ* cohærentes; *superiores* uniloculares; *inferiores* biloculares, uno loculo sterili, oblongo, apiculato.

1. *Æ. Indica*, Roxb. (Reuter in DC. Prod. xi. p. 43, cum syn.; Wight, Ic. t. 895.)—C.P. 1778.

HAB. Hantani, *Macrae*.

2. *Æ. acaulis*, Walp. (Reut. l. c. cum syn.?)

Var. *a*; *calyce* spathaceo, *ramis placentæ* parce irregulariter lobatis.—C.P. 3635.

Var. *β*; *calyce* brevior, tubuloso, 2-5-dentato, *ramis placentæ* non lobatis.—C.P. 2972.

HAB. Var. *a*. Below Hapootelle, amongst *Andropogon Schænanthus*.  
Var. *β*. Rambodde District.

*Corolla* (var. *β*) albida, limbo ad marginem pulcherrime violaceo-purpureo, labio inferiore interne ad faucem aurantiaco.

Varr. *a* and *β* are probably distinct species, but I have not sufficient materials as regards var. *a* to enable me to determine this point satisfactorily. Var. *β* forms, with *Christisonia unicolor*, a connecting link between the present and the next genus.

### 2. CHRISTISONIA, Gardner.

(Calcutta Journ. of Nat. Hist. viii. p. 153. *Christisonia*, Wight; et forsan *Campbellia*, Wight, et *Oligopholis*, Wight, Ic.)

*Calyx* tubulosus, 5-dentatus. *Stamina* didynama. *Antheræ* cohærentes, biloculares, singularum uno loculo sterili, subulæformi, porrecto.

1. *C. grandiflora*, Gardn. l. c. p. 155.—C.P. 3085.

HAB. "Parasitic on the root of a large Acanthaceous shrub, on the ascent to Adam's Peak from Ratnapoorah, at an elevation of about 5000 feet, March," *Gardner*.

*Caulis* glaber, squamosus, parce ramosus, 2-3-pollicaris; *ramis* 1-3-floris. *Squamæ* ovatae, subacutæ, fuscae, circiter 4 lin. longæ. *Pedicelli* teretes, ebracteolati, pollicares. *Calyx* 5-angularis, 9 lin. longus, dentibus lanceolatis, æqualibus, 3 lin. longis. *Corolla* circiter 3 poll. longa; *limbo* roseo, 2 poll. in exp.; *lobis* crenatis, inferiore ad faucem croceo. *Stamina* inclusa.—*A* *descript. cl. Gardner excerpt.*

This is, I suspect, a form only of the following species, the only real difference appearing to be in the size and colour of the flowers; for the character derived from the position of the sterile loculus of the anthers is a fallacious one, this being, in all the species of the genus I have examined in a fresh state, porrected and nearly horizontal.

2. **C. tricolor**, Gardn. l. c. p. 156.—c.p. 2889.

HAB. Central Province, at an elevation of 3000 to 4000 feet; parasitic on roots of *Acanthaceæ*, Gardner.

*Caulis* glaber, squamosus, parce ramosus, 1-3-pollicaris; *ramis* 1-2-floris. *Pedicelli* teretes, ebracteolati, rufescentes, pollicares. *Calyx* 5-angularis, rufescens, 8 lin. longus; *dentibus* æqualibus, lanceolatis, acutis, 3 lin. longis. *Corolla* circiter 2½ poll. longa; *tubo* infundibuliformi, fauce lutea; *limbo* 1-1¼ poll. in exp.; *lobis* 2 superioribus denticulatis, sanguineis, 3 inferioribus subrenatis, pallide roseis. *Stamina* inclusa.—Gardner.

3. **C. bicolor**, Gardn. l. c. p. 160.—c.p. 3108.

Var.  $\beta$ . *pallidiflora*.—*C. pallida*, Gardn. l. c. p. 159.—c.p. 1781.

HAB. In similar situations with the last, Lear, Gardner.

*Caulis* simplex, 3-6 poll. longus, verrucoso-hispidus, apicem versus rufescens, sublævis squamosusque. *Squamæ* ovatæ, acutæ, circiter 4 lin. longæ, pilosulæ. *Flores* racemosi, sæpissime 5-6. *Pedicelli* 1-8 lin. longi, bibracteolati. *Calyx* tubulosus, pilosulus, rufescens, circiter 8 lin. longus; *dentibus* æqualibus, acutis, circiter 1 lin. longis. *Corolla* infundibuliformis, sesquipollicaris, pallide roseo, plus minus flavescens, vel omnino flava; *limbo* 1-1¼ poll. in exp., lobis undulatis. *Stamina* inclusa; *filamentis* glanduloso-pilosis.—*A descrip. cl. Gardn. præcipue excerpt.*

4. **C. unicolor**, Gardn. l. c. p. 161.—An *Campbellia cytinoides*, Wight, Ic. t. 1425, cum syn. ?—c.p. 2971, ? 1780.

HAB. Hunasgiria Range, Lear. Rambodde, at an elevation of 3000 to 4000 feet.

Tota planta flava. *Caulis* simplex, glaber, angularis, squamosus, 1-3-pollicaris, ad basin verrucoso-hispidus. *Squamæ* subrotundæ, 3-6 lin. longæ. *Flores* racemosi, sæpissime 5-6. *Pedicelli* 2-8 lin. longi, ebracteolati. *Calyx* pollicaris, angularis, irregulariter acute 5-dentatus. *Corolla* infundibuliformis, 1½-2 poll. longa; *lobis* integris, parum reflexis. *Stamina* exserta; *filamentis* glabris; *antherarum* superiorum loculo sterili parvo. *Placentæ* majusculæ.

## CXVII. PODOSTEMACEÆ.

### 1. DICRÆA, Pet. Th.

1. **D. elongata**, Tulasne in Ann. des Sci. Nat. (sér. 3), vol. ii. p. 102. Walp. Ann. iii. p. 437; Wight, Ic. t. 1917-1.—*Podostemon elongatus*, Gardner in Calc. Journ. of Nat. Hist. vol. vii. p. 188.—c.p. 2259.

HAB. Rocks in the bed of the Mahawelle Ganga, near Kandy, Gardner. Fl. January to March.

### 2. PODOSTEMON, Mich.

1. **P. subulatum**, Gardn. l. c. p. 184; Wight, Ic. t. 1918-1; Tulasne, l. c. p. 103.—*P. dendroides*, Thw. MS. in Hb. Hook.—c.p. 3088, 3366.

HAB. Rocks in the Mahawelle Ganga. Fl. January to March, Gardner.

2. **P. Gardneri**, Harvey, MS.; *caule* simplici, terete, glabro, foliis plurimis capillaceis coronato.—C.P. 2989.

HAB. On the membranaceous rhizomes of *Hydrobryum olivaceum*, Tul. In a rapid mountain stream at Rambodde, Oct. 1853, *Dr. W. H. Harvey*.

*Caulis*  $\frac{1}{2}$ –1-pollicaris. *Folia*  $1\frac{1}{2}$ –2 poll. longa, delicatula, nigro-viridia, enervosa. *Inflorescentia* ignota.

Until its inflorescence has been discovered, some doubt must remain to which genus of the present family this plant is to be referred, and my attempts to trace its further development from the state described above, have been hitherto unsuccessful. Can it possibly be an early stage of growth of *Hydrobryum olivaceum*?

### 3. HYDROBRYUM, Endl.

1. **H. olivaceum**, Tul. l. c. p. 104; Walp. Ann. iii. p. 439.—*Podostemon olivaceum*, Gardn. l. c. p. 181.—C.P. 3065.

HAB. On rocks in the Mahawelle Ganga, and large mountain streams in the Central Province, *Gardner*. Fl. January to March.

### 4. DALZELLIA, Wight, Ic. vol. v. par. ii. p. 34.

(*Lawia*, Tulasne. *Mnianthus*, Walp.)

1. **D. Ceylanica**, Wight, Ic. t. 1919–1.—*Lawia Zeylanica*, Tul. l. c. p. 112. *Mnianthus Zeylanicus*, Walp. l. c. p. 443. *Tristicha Ceylanica*, Gardn. l. c. p. 177.—C.P. 3089.

HAB. On rocks in the Mahawelle Ganga, near Kandy, *Gardner*.

## CXVIII. ACANTHACEÆ.\*

[The Indian *Acanthaceæ* being in process of examination and re-arrangement at Kew, by Dr. Thomas Anderson, F.L.S., of the Bengal Medical Service, who is preparing a new classification of the Order for the Linnæan Society of London, Mr. Thwaites has, at my suggestion, gladly availed himself of the opportunity of placing his enumeration of the Ceylon species (which was arranged according to Nees's method) in Dr. Anderson's hands. The latter gentleman has kindly devoted some weeks to the study of every species and its synonyms, and has classified the whole in accordance with his forthcoming 'Systema Novum Acanthacearum.' I have added the initials G. Thw. and T. A. to various paragraphs where doubts may arise as to the exact authority for the observations they contain."—*J. D. H.*]

SUBORD. 1. **THUNBERGIDEÆ**.—*Æstivatio corollæ contorta. Semina cupulâ suffulta. Plantæ scandentes.*

TRIB. **THUNBERGIEÆ.**

### 1. THUNBERGIA, Linn. fl.

1. **T. coccinea**, Wall. Tent. Fl. Nep.—*Hexacentris*, N. ab E. in Wall. Pl. As. Rar.; DC. Prod. xi. p. 61, cum. syn.

Macrae's specimens, referred to by Nees von Esenbeck, were collected in the Botanic Garden from a cultivated plant, as I am informed by Mr. Alwis, the draughtsman

\* The native name *Nelloo* is applied to the species of this Natural Family generally.—*G. Thw.*

of this establishment. Mr. Alwis is under the impression that he once saw the plant growing wild in a forest between Oova and Balangodde. I have however only seen it in a cultivated state, and suspect it is not truly indigenous.—*G. Thw.*

2. **T. Hawtynii**, Wall. Tent.—*Meyenia Hawtyniana*, N. ab E. in Wall. Pl. As. Rar.; DC. Prod. xi. p. 60, cum. syn.; Wight, Ic. t. 1487.  
HAB. Ceylon, *Walker, in Herb. Hook.*

Mr. Thwaites has never met with this species in the island.

3. **T. fragrans**, Roxb. Fl. Ind. iii. p. 33; DC. l. c. p. 57, cum. syn.—  
C.P. 2020.

HAB. Abundant up to an elevation of 7000 feet.

*Flores non fragrantæ. Corolla alba, tubo interne pallide viridi.—T. A.*

SUBORD. 2. *RUELLIDEÆ*.—*Æstivatio corollæ contorta. Semina papillâ v. retinaculo uncinato suffulta. Plantæ non scandentes.*

#### TRIB. NELSONIÆ.

### 2. ELYTRARIA, Vahl.

1. **E. crenata**, Vahl, (N. ab E. in DC. Prod. xi. p. 63, cum syn.)—*E. marginata*, Pal. de Beauv. *E. virgata*, N. ab E.; DC. l. c.—C.P. 240.  
HAB. Very common.

### 3. NELSONIA, R. Br.

1. **N. tomentosa**, Willd. (N. ab E. in DC. Prod. xi. p. 65, cum syn.)  
HAB. Ceylon, *Walker, in Herb. Hook.*

Mr. Thwaites has never met with this plant.

### 4. EBERMAIERA, N. ab E.

1. **E. glauca**, N. ab E. in DC. Prod. xi. p. 73, cum syn.; Wight, Ic. t. 1488.—C.P. 2717, 3575.

HAB. Banks of a stream near Batticaloa and Ambagamowa District, not uncommon.

Under *E. glauca* I include *E. Zeylanica, polybotrya, humilis*, and *thyrsoides* of Nees von Esenbeck, as well as *E. spathulata*, Hassk.—*T. A.*

### 5. ADENOSMA, N. ab E.

#### Sect. 1.—*Staminibus quatuor.*

1. **A. balsamea**, Spr. (DC. Prod. xi. p. 68; Wight, Ic. 446.)—C.P. 1996.

HAB. North of the island, *Gardner.*

2. **A. verticillata**, N. ab E. in Wall. Pl. As. Rar.; DC. l. c. p. 69; Wight, Ic. 1524.—C.P. 1995.

HAB. Not uncommon in the hotter parts of the island.

3. **A. uliginosa**, R. Br. (DC. l. c.)

HAB. In rice-fields, *Walker, in Herb. Hook.*

#### Sect. 2.—*Staminibus duobus.*

4. **A. Thwaitesii**, T. Anders.; *caule erecto, racemoso; foliis obovatis,*

subpetiolatis, superne tomentosis, subtus glaucis; *floribus* solitariis, in cymis laxis; *bracteis* parvis; *calyce* capsula dimidio brevior; *corolla* minuta, labio inferiore rugis inconspicuis; *staminibus* duobus (*T. A.*).—C.P. 1994.

HAB. Abundant in the Batticaloa District, at no great elevation.

### TRIB. RUELLIÆ.

#### SUBTRIB. 1. HYGROPHILÆ.

#### 6. HYGROPHILA, R. Br.

1. *H. quadrivalvis*, N. ab E. in Wall. Pl. As. Rar. p. 89. (DC. Prod. xi. p. 89, cum syn.)—C.P. 2993.

Var. *salicifolia*, T. Anders.—*H. salicifolia*, N. ab E. l. c. p. 92. (Wight, Ic. t. 1490.)—C.P. 591, 3109.

HAB. Common in the warmer parts of the island.

2. *H. spinosa*, T. Anders.—*Asteracantha longifolia*, N. ab E. in DC. l. c. p. 247, cum syn.; Wight, Ic. t. 449.—C.P. 2900.

HAB. Not uncommon in the warmer parts of the island.

*Asteracantha* is, in every respect, a true *Hygrophila*.—*T. A.*

#### SUBTRIB. 2. EU-RUELLIÆ.

#### 7. CALOPHANES, Don.

1. *C. depressa*, T. Anders.—*Dyschoriste*, N. ab E. in DC. Prod. xi. p. 106, cum syn.—C.P. 3537.

HAB. Hot, drier parts of the island.

2. *C. littoralis*, T. Anders.—*Dyschoriste*, N. ab E. in DC. l. c. cum syn.; Wight, Ic. t. 447.—C.P. 1997.

HAB. North of the island, *Gardner*.

#### 8. RUELLIA, Linn.

1. *R. prostrata*, Poir.—*Dipteracanthus prostratus*, N. ab E. in DC. Prod. xi. p. 124, cum syn. *D. dejectus*, N. ab E. in DC. l. c. p. 125.—C.P. 1987.

HAB. Very abundant up to an elevation of 3000 feet.

2. *R. patula*, Jacq.—*Dipteracanthus patulus*, N. ab E. in DC. l. c. p. 126, cum syn.; Wight, Icon. t. 1505.—C.P. 3628.

HAB. Trincomalee, *Rev. S. O. Glenie*.

#### 9. ÆTHEILEMA, R. Br.

1. *Æ. reniforme*, N. ab E. in Wall. Pl. As. Rar.; DC. Prod. xi. p. 261, cum syn.; Wight, Ic. t. 1533.—C.P. 398.

HAB. Not uncommon up to an elevation of 2000 feet.

#### SUBTRIB. 3. STROBILANTHÆ.

#### 10. STENOSIPHONIUM, N. ab E.

1. *S. Zeylanicum*, T. Anders.; scabrum, erectum; *foliis* petiolatis,



serratis; *floribus* confertis; *bracteis* ovatis, acutis, marginibus hispide ciliatis; *corolla* pallide cyanea, glabra; *staminibus* quatuor, didynamis, longe exsertis (*T. A.*).—C.P. 3577.

Var.  $\beta$ ; *foliis* marginibus integris.—C.P. 3661.

HAB. Bintenne and Batticaloa Districts, at no great elevation. Var. *a*. Central Province.

2. **S. Russellianum**, N. ab E. in Wall. Pl. As. Rar.; DC. Prod. xi. p. 105; Wight, Ic. t. 873, 1502.—*S. subsericeum*, N. ab E.—C.P. 2001.

HAB. Common in the hot, drier parts of the island. Varies greatly in the amount of pubescence.

## 11. STROBILANTHES, Blume.

### Sect. 1. ENDOPOGON.—*Floribus* spicatis.

† *Staminibus* duobus.

1. **S. consanguineus**, T. Anders.—*Endopogon consanguineus*. *E. amomum*, N. ab E. in DC. Prod. p. 104. *E. viscosum*, Wight, Ic. t. 1498. *Stenosiphonium diandrum*, N. ab E.—C.P. 2002, partim.

HAB. Maturatte District, at an elevation of 3000 feet.

Valde ramosus, 5-6-pedalis. *Corolla* pallide cyanea.

Very variable in the amount of pubescence.—*T. A.*

2. **S. hypoleucus**, T. Anders.—*Endopogon*, N. ab E. in DC. l. c.—C.P. 2897.

HAB. Central Province, *Gardner*.

3. **S. nigrescens**, T. Anders.; *cauli* erecto, tetragono; *foliis* glabris, ovatis, lanceolatis, utrinque acutis, petiolatis, venis plerumque nigrescentibus; *spicis* hispidis, *foliis* longioribus; *bracteis* lanceolatis (*T. A.*).—C.P. 3353.

HAB. Ambagamowa District, at an elevation of 3000 feet.

Suffrutex luridus, 4-5-pedalis, ad nodos rufescens. *Corolla* pallide pupurea. *Antherarum* loculi parum dislocati, in anthesin horizontales.—*T. A.*

4. **S. viscosus**, T. Anders.—*Endopogon viscosus*, *argutus*, et *digitalis*, N. ab E. in DC. l. c. *E. versicolor*, Wight, Ic. t. 1497.—C.P. 2002, partim, et 2898.

HAB. Central Province, up to an elevation of 7000 feet.

*Corolla* albida.—*G. Thw.*

A very variable species as regards the amount of the glandular pubescence, and in the size of the several parts of the inflorescence.—*T. A.*

5. **S. rhamnifolius**, T. Anders.—*Buterea rhamnifolia*, N. ab E. in D. C. l. c. p. 725. *Endopogon rhamnifolius*, Wight, Ic. t. 1521.—C.P. 2007.

HAB. Hantani District, at an elevation of 4000 feet.

*Corolla* pallide pupurea.—*G. Thw.*

†† *Staminibus* quatuor.

6. **S. Gardnerianus**, T. Anders.—*Endopogon*, N. ab E. in DC. l. c. p. 723.—C.P. 2006.

HAB. Hantani, at an elevation of 4000 feet.

1-2-pedalis. *Corolla* pupurea.—*G. Thw.*

7. **S. scaber**, T. Anders.—*Ruellia aspera*, N. ab E. in DC. l. c. p. 147, cum syn. *Strobilanthes scaber*, var.  $\beta$ , Wall. Pl. As. Rar., *fid.* N. ab E.

HAB. Ceylon, *Walker*, in *Herb. Hook.*

Sect. 2. EU-STROBILANTHES.—*Floribus in strobilis confertis.*

8. **S. Zeylanicus**, T. Anders.; *caule* fruticoso, erecto, quadrangulari, sulcato; *foliis* acutis, subtus scabris, marginibus serratis asperis; *strobilis* axillaribus, pedunculatis; *bracteis* fimbriatis, truncatis, reflexis vel acutis; *corolla* extus tomentosa (*T. A.*).—C.P. 3352.—*Strobilanthes callosus*, N. ab E., partim.

HAB. Ambagamowa District and Kokool Corle, at an elevation of 2000 feet.

3-5-pedalis. *Caulis* subglaber. *Folia* setulis stellatis scabra, elliptica vel ovata, 3-10 unc. longa, 1-4 unc. lata. *Bractea* trapezoidea vel acuta, foliacea, margine setulosa. *Corolla* albida, extus tomentosa.—*T. A.*

9. **S. Walkeri**, N. ab E. DC. l. c. p. 181.—C.P. 2008, 2009, partim, et 3517.

HAB. Central Province, at an elevation of 4000-7000 feet.

2-5-pedalis. *Corolla* albida, vel pallide purpurea. *Antheræ* exsertæ.—*T. A.*

It varies very much in the size of the flowers.

10. **S. Thwaitesii**, T. Anders.; *caule* fruticoso, tetragono, angulis scabris, setulis squamoso; *foliis* ovatis, ovato-lanceolatis, vel ellipticis, acuminatis, marginibus serratis, reflexis, supra asperis, subter hirsutis; *floribus* in capitulis strobiloideis; *pedunculis* axillaribus, tetragonis; *bracteis* lanceolatis, acuminatis, corrugatis, margine ciliatis; *staminibus* inclusis (*T. A.*).—C.P. 2009 partim, 3516.

HAB. Ambagamowa District.

*Pedunculi* foliis breviores, simplices vel ad apicem trifidi. *Capitula* circiter 3-flora, cernua. *Bracteæ* bullato-corrugatæ, ciliatæ, 7-10 lin. longæ. *Bracteolæ* multo minores, lineares, acutæ, ciliatæ. *Corolla* pollicaris, albida, externe glabra, interne pilosa.—*T. A.*

11. **S. deflexus**, T. Anders.; *caule* glabro; *foliis* ellipticis, caudatis; *pedunculis* axillaribus, solitariis vel binis, deflexis; *bracteis* foliaceis, acutis, glabris, calyce fere ad medium quinque-partito, segmentis lanceolatis; *staminibus* duobus (*T. A.*).—C.P. 1998.

HAB. Hantani, at an elevation of 4000 feet.

Viridis, 4-5-pedalis. *Pedunculi* versus apicem incrassati. *Capitula* 6-8 lin. longa. *Bracteæ* 6-8 interiores cum calycibus involventes. *Bracteolæ* angustissimæ. *Flores* terminales, 2-3. *Calyx* pilosulus. *Corolla* pallide purpurea, 8 lin. longa.—*T. A.*

12. **S. Hookeri**, N. ab E. DC. l. c. p. 185.—C.P. 1999.

HAB. Central Province, at an elevation of 7000 feet.

5-6-pedalis. *Corolla* albida, lobis rotundatis, retusis, ad faucem lineis 3 purpureis longitudinalibus notata. *Stamina* æquilonga, *antheræ* exsertæ.—*T. A.*

13. **S. calycinus**, N. ab E. DC. l. c. p. 186.—C.P. 309.—*S. coloratus*, N. ab E. l. c. p. 186.

HAB. Central Province, at an elevation of 7000 feet.

4-5-pedalis. *Bracteæ* glabræ vel glanduloso-pilosæ. *Corolla* straminea vel plus minus olivacea, lobis brevibus, rotundatis, retusis. *Stamina* æquilonga, *antheris* exsertis.—*G. Thw.*

Closely allied to *S. Hookeri*, but differing from it in having much shorter petioles, smaller hirsute leaves, longer and laxer heads of flowers, and oblong, more membranaceous bracts, as well as in the character of the corolla.—*T. A.*

14. ***S. trifidus***, N. ab E. DC. l. c. p. 185.—*S. rugosus*, Wight, Ic. t. 1619, et *S. campanulatus*, t. 1562.—C.P. 254, 2008.—*S. asperrimus*, *lupulinus*, *Heyneanus*, *tener*. (*S. involucratus*, Blume, *fid.* N. ab E.)

HAB. Very abundant up to an elevation of 6000 feet.

An extremely variable plant as regards amount of pubescence, size of the leaves, length of the heads of flowers, etc. In some specimens the bracts are nearly white, more or less tinged with red at the base and green at the tip of each, whilst in others they are of a uniform dull green colour, and harsher texture.—*T. A.*

15. ***S. laxus***, T. Anders.; *caule* tereti, glabro; *foliis* petiolatis, ovatis, acuminatis, cordatis, grosse serratis, utrinque subscabris, subtus glandulosis; *strobilis* axillaribus, multifloris, sæpe ramosis, laxis, inferioribus plerumque binis, superioribus solitariis; *bracteis* rotundatis, ovatis, margine ciliatis; *bracteolis* lanceolatis, acutis, ciliatis, calyce brevioribus; *calycis* segmentis lanceolatis, acutis, subæqualibus, inferiore angustiore, fere lineari; *corolla* extus glabra, ad medium quinque-partita, labio inferiore lineis duabus barbatis notato; *lobis* lanceolatis, acutis, reflexis; *staminibus* quatuor exsertis, filamentis æquilongis, linearibus, glabris (*T. A.*).—C.P. 2010.

HAB. Hantani, at an elevation of 4000 feet.

4-6-pedalis. *Folia* 1-2 unc. longa, 1-2 unc. lata. *Petiolus* 1½ unc. longus. *Strobili* ½-3 unc. longi. *Bracteæ* semipollicares. *Corolla* 9 lin. longa, flavo-rufescens, basi nigro-purpurascens. *Stamina* nigrescentia, *polline* flavo.—*T. A.*

16. ***S. vestitus***, N. ab E. in DC. l. c. p. 180.—C.P. 3660.

HAB. Maturatte, at an elevation of 5000-7000 feet.

17. ***S. Arnottianus***, N. ab E. in DC. l. c. p. 179.—C.P. 2005.

HAB. Newera Ellia, *Gardner*.

Closely allied to *S. vestitus*, but differing from it in the form of the leaves, and much smaller pilose flowers. Besides, the capsule and style of *S. vestitus* are quite glabrous, while they are hirsute and glandular in *S. Arnottianus*.—*T. A.*

18. ***S. caudatus***, T. Anders.; *caule* subterete, subglabro; *foliis* lanceolatis, caudato-acuminatis, lineolatis, subtus ad nervos tomentosus; *pedunculis* axillaribus, simplicibus vel trifidis; *capitulis* 3-6-floris, villosis; *bracteis* lanceolatis, tomentosus; *calycis* segmentis lineari-lanceolatis, æquilongis; *corollæ* lobis brevibus, rotundatis, retusis; *staminibus* inclusis; *antheris* subunilocularibus (*T. A.*).—C.P. 364.

HAB. Hantani, at an elevation of 2000 to 3000 feet.

4-5-pedalis. *Folia* 3-7 unc. longa, 1-1½ unc. lata. *Bracteæ* 4 lin. longæ, ciliatæ. *Corolla* pollicaris, flavide-rubro tincta.

19. ***S. adenophorus***, N. ab E. in DC. l. c. p. 182.—C.P. 162.—*Goldfussia myrtinia*, N. ab E. *Myrtinia lanceolata*, Macrae. *Stenosiphonium Moonianum*, Wight et Gardn. *Strobilanthes lanceolatus*, N. ab E.

HAB. Kokool Corle, and Saffragam District.

*Calyx* semipollicaris, ultra medium tubulosus, segmentis lanceolatis, acutis, ciliatis. *Corolla* pallide cyanea, 1½ unc. longa; *lobis* subacutis. *Antheræ* sagittatæ, exsertæ.—*T. A.*

Nees von Esenbeck has placed this plant once in *Goldfussia* and twice in *Strobilanthes*, and both in *Goldfussia* and *Strobilanthes* has quoted the same number of Macrae, 373.—*T. A.*

20. **S. anceps**, N. ab E. in DC. l. c. p. 189.—c.P. 2000.

Var. *punctatus* (*S. punctatus*, N. ab E. in DC. l. c. p. 182); *foliis* subtus punctatis; *bracteis* utrinque punctatis, ciliatis.—*T. A.*

HAB. Central Province, Saffragam District, at an elevation of 2000 to 6000 feet.

Var. *punctatus*, Walker, in Herb. Hook.

Sect. 3.—*Floribus in paniculis terminalibus caulibusve.*

21. **S. cerinthoides**, N. ab E. in Herb. Hook.—c.P. 66, 2593, 2594.—

*S. argutus*, N. ab E. in DC. l. c. p. 188. *S. sexennis*, N. ab E. in DC. l. c. p. 187.

Var. *a. hirsutissimus*, T. Anders.—c.P. 3626 (*S. hirsutissimus*, N. ab E. in DC. l. c.).

HAB. Very abundant at Newera Ellia, and other of the more elevated parts of the Central Province.

An extremely variable plant as regards amount of pubescence, shape of the leaves, length of the subpanicles of flowers, and relative size of the bracts and flowers.—*T. A.*

22. **S. helicoides**, T. Anders.—*Leptacanthus*, N. ab E. in DC. l. c. p. 170.—c.P. 1980.

HAB. Hantani and Ambagamowa Districts, up to an elevation of 3000 feet.

2-3-pedalis. *Corolla* purpurea; *tubo* parum inflato; *lobis* brevibus, rotundatis.—*G. Thw.*

23. **S. rubicundus**, T. Anders.—*Leptacanthus*, N. ab E. in DC. l. c. —c.P. 3351.

HAB. Saffragam and Ambagamowa Districts.

4-5-pedalis. *Corolla* violacea.—*G. Thw.*

24. **S. pulcherrimus**, T. Anders.—*Leptacanthus Walkeri*, N. ab E. in DC. l. c.; Wight, Ic. t. 1507.—c.P. 266 (301).

HAB. Central Province, at an elevation of 4000 to 7000 feet.

4-5-pedalis. *Paniculae* terminales caulinaeque, cum *bracteis* calycibusque plus minus rufescentes. *Corolla* violacea, parce curvata, *lobis* rotundatis retusis.—*G. Thw.*

12. **DÆDALACANTHUS**, T. Anders.

(*Eranthemum*, Wall. partim.)

*Calyx* minutus, scariosus, 5-dentatus. *Corolla* hypocraterimorpha; *tubo* gracili, elongato; *limbus* lobis in aestivatione valde contortis. *Stamina* quatuor, didynama, fauce inserta, sterilia brevissima. *Antheræ* biloculares, muticæ, loculis parallelis. *Stigma* subulatum, incurvum, dorso canaliculatum. *Capsula* oblonga, tetrasperma, retinaculis acutis, dissepimentis breviter adnatis. *Semina* subrotunda, compressa, lævia, margine sericea pilis adpressis.—Suffrutices, *bracteis pulcherrime variegatis*; *bracteolis paucis, scariosis*.—*T. A.*

1. **D. montanus**, (var. *a.*) T. Anders. *E. montanum*, Roxb. Var. *a.*

N. ab E. in DC. Prod. xi. p. 448 ; Wight, Ic. t. 466.—c.P. 2003.—*Spicis* laxis ; *bracteis* distantibus, lanceolatis, longissime attenuatis.

HAB. Common up to an elevation of 2000 feet.

A variable plant in the length and breadth of bracts, as well as in the amount of pubescence on the spikes.—*T. A.*

SUBORD. 3. *ACANTHIDEÆ*.—*Æstivatio corollæ imbricata vel imbricatobilabiata. Semina retinaculo uncinato suffulta.*

TRIB. *BARLERIÆ*.

13. *BARLERIA*, Linn.

1. *B. bispinosa*, Vahl, (DC. Prod. xi. p. 241.)—c.P. 2013.—*B. spina Ceylanica*, N. ab E. l. c. p. 242.

HAB. Hot, drier parts of the island.

2. *B. Arnottiana*, N. ab E. in DC. l. c. p. 232.—c.P. 682, 310, *ambipartim*.

HAB. Newera Ellia.

*Corolla* pallide cyanea.—*G. Thw.*

3. *B. cristata*, Linn. (N. ab E. in Prod. xi. p. 229, cum syn. ; Wight, Ic. t. 453.)—*B. dichotoma*, et *B. ciliata*, Roxb. Fl. Ind. *B. Nepalensis*, N. ab E. l. c. p. 228.

HAB. Common in gardens, but never met with truly wild in the island.

Repeated examinations of an extensive set of specimens in the Hookerian Herbarium have convinced me that *B. cristata*, *dichotoma*, *ciliata*, and *Nepalensis*, are forms of one variable species. The differences supposed to exist, in the leaves being stalked in *B. dichotoma* and sessile in *B. cristata*, are not at all constant. The inflorescence varies equally as much, the flowers being solitary in the axils of the leaves, or in short spikes of 4 or 5. The colour of the corolla varies from all shades of white to dark lilac or purple. Roxburgh describes and figures the corolla of his *B. dichotoma* as white.—*T. A.*

4. *B. nutans*, N. ab E. in DC. l. c. p. 227.—c.P. 2011, 2559.

HAB. Central Province, at an elevation of 1000 to 3000 feet.

*Corolla* cyaneo-purpurea.—*G. Thw.*

5. *B. Prionitis*, Linn. (N. ab E. in DC. l. c. p. 237 ; Wight, Ic. t. 452.)

HAB. Very abundant in the hotter parts of the island. Nom. vulg. "Kattoo Kurandoo."

This plant is employed as a medicine by the Cinghalese.—*G. Thw.*

6. *B. pentandra*, Arn. (N. ab E. in DC. l. c. p. 232 ; Wight, Ill. t. 164.)—c.P. 310, 682, *partim*, et 2558.—*B. involucrata*, N. ab E. in DC. l. c.

HAB. Ambagamowa District.

*Corolla* cyaneo-purpurea.—*G. Thw.*

*B. vestita*, T. Anders. ; herbacea, hirsuta pilis patentibus ; *foliis* ovatis, ellipticis, utrinque attenuatis, petiolatis vel subsessilibus ; *cymis* axillaribus terminalibusque, sparsifloris, laxis ; *bracteis* angustissime lanceolatis, acutis ; *calycis* segmentis inæqualibus, lanceolatis, strigosis, exteriorum superiore acuto, inferiore minute bifido, interioribus acutis,

brevioribus; *corolla* anguste infundibuliformi, 3-pollicari, lobis tubo æquilongis (*T. A.*).—C.P. 2004.—*B. pentandra*,  $\beta$ , N. ab E. in DC. l. c. 233.

HAB. Central Province, and in the Saffragam District, up to an elevation of 5000 feet.

6–8-pedalis, hispida. *Folia* integra, 6–12 unc. longa. *Cymæ* longe pedunculatæ, foliis paulo breviores. *Calyx* fulvus, strigosus. *Corolla* pallide purpurea vel albida.—*T. A.*

*B. noctiflora* and *B. buxifolia*, var.  $\beta$ . The recording of these two plants from Ceylon seems to be an error of Nees von Esenbeck, as no specimen from Ceylon occurs in any of the extensive collections I have consulted. For *B. noctiflora* no authority is quoted by Nees but “Herb. Hook.,” and for *B. buxifolia*, var.  $\beta$ , “Forsyth, in Herb. eod.,” and I am informed that this botanist made no collections in Ceylon.—*T. A.*

#### 14. CROSSANDRA, Salisb.

1. *C. infundibuliformis*, N. ab E. in Wall. Pl. As. Rar.; DC. Prod. xi. p. 280, cum syn.; Wight, Ic. t. 460, et t. 461.—C.P. 255 (2015), 2016.—*C. axillaris*, *C. oppositifolia*, N. ab E. l. c. (*T. A.*)

HAB. Common in the warmer parts of the island.

#### 15. LEPIDAGATHIS, Willd.

1. *L. Ceylanica*, N. ab E. Ill. Lepidag. (DC. Prod. xi. p. 259. *L. nervosa*, Wight, Ic. t. 1620.)—C.P. 1985. (*T. A.*)

HAB. Not uncommon up to an elevation of 4000 feet.

2. *L. fasciculata*, N. ab E. Ill. Lepidag. (DC. Prod. xi. p. 260.)—C.P. 1983.

HAB. Hot, drier parts of the island.

3. *L. Walkeriana*, N. ab E. in DC. l. c.; Wight, Ic. t. 1530.—C.P. 85, 1986.

HAB. Very common throughout the island.

4. *L. hyalina*, var. *a*, *lophostachyoides*, N. ab E. Ill. Lepidag. (DC. l. c. p. 252.)—C.P. 1971.—*Spicis* terminalibus, subsolitariis; *filamentis* lineis purpureis transversim notatis (*T. A.*)

HAB. Central Province, at an elevation of 3000 to 4000 feet.

This variety may ultimately prove to be distinct from *L. hyalina*. In the absence of a sufficient number of specimens to decide from, I do not feel justified in making a new species of it.—*T. A.*

#### TRIB. ACANTHÆ.

#### 16. BLEPHARIS, Juss.

1. *B. molluginifolia*, Juss. (DC. Prod. xi. p. 266, cum syn.)—C.P. 3576.

HAB. Batticaloa District.

2. *B. Boerhaaviaefolia*, Juss. (DC. l. c. cum syn.; Wight, Ic. t. 458.)—C.P. 2014.

HAB. Very abundant in the hot, drier parts of the island. Nom. vulg. “Laawaadaana.”

The Cinghalese use the leaves as a vulnerary.—*G. Thw.*

17. **ACANTHUS**, Linn.

1. **A. ilicifolius**, Blume, (DC. Prod. xi. p. 268, cum syn.; Wight, Ic. t. 459.)—C.P. 2018.—*Dilivaria ilicifolia*, Juss.  
 Var. *a. integrifolius*, T. Anders.—C.P. 2017.  
 HAB. Common near the sea.

TRIB. *ANDROGRAPHIDÆ*.18. **ANDROGRAPHIS**, Wall.

1. **A. paniculata**, N. ab E. in DC. Prod. p. 515, cum syn.; Wight, Ic. t. 518.)—C.P. 3664 (1981 partim).  
 HAB. South of the island, not uncommon.  
 Pedunculi glabri.—*T. A.*
2. **A. alata**, N. ab E. in DC. l. c. p. 516, cum syn.—C.P. 1977.  
 HAB. Common upon banks, in the hot, drier parts of the island.  
*Folia* linearia, lanceolata, vel oblonga, basi gradatim angustata. *Corolla* albida. *Antheræ* purpureæ.—*G. Thw.*
3. **A. macrobotrys**, N. ab E. in DC. l. c.—C.P. 3578, 1981 partim.  
 HAB. Near Kandy, not common.  
*Pedunculi* patentim glanduloso-pilosi. *Corolla* pallide purpurea, interne labio inferiore nigro-purpureo albo-marginato. *Antheræ* purpureæ, basi albo-barbatae.—*G. Thw.*
4. **A. Ceylanica**, N. ab E. in DC. l. c. p. 518; Wight, Ic. t. 1560.  
 HAB. Ceylon, *Walker, in Herb. Hook.*
5. **A. echioides**, N. ab E. in DC. l. c. cum syn.; Wight, Ic. t. 467.—C.P. 606.  
 HAB. Very common in the hotter parts of the island.

19. **GYMNOSTACHYUM**, N. ab E.

1. **G. sanguinolentum**, T. Anders.—*Cryptophragmium sanguinolentum*, cum syn. et *C. axillare*, N. ab E. in DC. Prod. xi. p. 96; Wight, Ic. t. 1496 (*T. A.*).—C.P. 186.  
 HAB. Very common in forests, up to an elevation of 4000 feet.  
 Planta tota plus minus purpurascens. *Corolla* violacea.—*G. Thw.*
2. **G. Ceylanicum**, W. Arn. et N. ab E.; DC. Prod. xi. p. 93.—C.P. 2019.  
 HAB. Shady places, in the hot, drier parts of the island; not uncommon.  
*Folia* superne secus costam venas primariasque albescentia. *Corollæ* tubo pallide purpureo; limbo flavo, externe viridescente. *Antheræ* albidæ.—*G. Thw.*
3. **G. Thwaitesii**, T. Anders.; tomentosum, caule radicante; *foliis* ovatis, obtusis, basi inæqualibus, petiolatis, hirsutis; *paniculis* terminalibus, elongatis, multifloris; *ramulis* oppositis, cymiformibus, laxis; *bracteis* sulcatis (*T. A.*).—C.P. 3356.  
 HAB. Ambagamowa District.
4. **G. paniculatum**, T. Anders.; *caule* subquadrangulare, sulcato-tomentoso; *foliis* ovatis, subacutis, basi in petiolo attenuatis, pilis ad-

pressis hirsutis; *paniculis* terminalibus, multifloris, ramulis multifloris; *floribus* confertis; *calyce* quinque-partito, segmentis brevibus lanceolatis, glandulosis, tomentosus; *corolla* extus subtomentosa, limbo breve, 5-fido, labio inferiore longiore paulo inflato (*T. A.*).—C.P. 2994.

Var. *lævius*; *foliis* caulibusque glabris.

HAB. Saffragam and Galle Districts, at an elevation of 1000 to 2000 feet.

Sæpe omnino purpurascens,  $1\frac{1}{2}$ –2-pedale. *Folia* 3–7 unc. longa,  $1\frac{1}{2}$ – $2\frac{1}{2}$  unc. lata. *Corolla* 7 lin. longa, violacea, limbo interne flavo-virescente. *Antheræ* albidæ.—*T. A.*

5. **G. hirsutum**, *T. Anders.*; *caule* erecto; *foliis* ovatis, acutis, basi attenuatis, petiolatis, subtus ad venos hirsutis, superne scabris, *paniculis* terminalibus confertis hirsutissimis; *bracteis* linearibus, ciliatis; *calyce* profunde 5-fido; *segmentis* longe setaceis, hirsute ciliatis; *corolla* calyce æquanti; *limbo* breviter 5-partito; *filamentis* basi paulo hirsutis; *capsula* 8–10-sperma (*T. A.*).

HAB. Central Province, at an elevation of 3000 to 5000 feet.

Purpurascens, 2–3-pedale. *Corolla* flavo-virescens. *Antheræ* albidæ.—*G. Thw.*

#### TRIB. JUSTICIEÆ.

### 19. JUSTICIA, Linn.

1. **J. Adhatoda**, Linn.—*Adhatoda Vasica*, *N. ab E.* in DC. Prod. xi. p. 387, cum syn.—C.P. 1991.

HAB. Common. Nom. vulg. “Paawetta.”

Used medicinally by the Cinghalese as an expectorant for children.—*G. Thw.*

2. **J. Betonica**, Linn.—*Adhatoda*, *N. ab E.* in DC. l. c. p. 385. *A. variegata*, *A. ramosissima*, *A. Nilgherrica*, *A. trinervia*, *A. arenaria*, *N. ab E.* in DC. l. c. p. 385, 386, 387.

HAB. Very abundant. Nom. vulg. “Soodoo-pooroo-gass.”

*A. Nilgherrica*, *Wight, Ic. t. 1544*, *A. trinervia*, and *A. arenaria*, are merely stunted states of *J. Betonica*; with a complete suite of specimens it is impossible to distinguish them.—*T. A.*

3. **J. Tranquebariensis**, Linn.—*Adhatoda*, *N. ab E.* in DC. l. c. p. 399, cum syn.; *Wight, Ic. t. 462.*—C.P. 1993.

HAB. Hot, drier parts of the island.

*Corolla* albida vel rubra, fauce rubro maculata.—*G. Thw.*

4. **J. Ceylanica**, *T. Anders.*—*Adhatoda*, *N. ab E.* in DC. l. c. p. 400, cum syn.—C.P. 2422, 2718, 3110 (713 partim).

HAB. Not uncommon up to an elevation of 3000 feet.

5. **J. Hookeriana**, *T. Anders.*—*Adhatoda*, *N. ab E.* in DC. l. c. p. 403.—C.P. 701. *Leptostachya Zeylanica*, *N. ab E.* partim.

HAB. Very abundant on the banks of streams in the Central Province.

*Bracteæ* lineares vel subspathulatæ.—*G. Thw.*

Macrae's plant in *Herb. Benth. n. 423*, quoted by *Nees von Esen.* as *Leptostachya Zeylanica*, is this species.—*T. A.*

6. **J. Gendarussa**, Linn.—*Gendurussa vulgaris*, *N. ab E.* in DC. l. c. p. 410, cum syn.—C.P. 3662.

HAB. Common. Nom. vulg. “Kalos-waaraneya.”



7. **J. glabra**, Koenig.—*Rhaphidospora*, N. ab E. in DC. l. c. p. 499, cum syn.; Wight, Ic. 1554.—c.p. 2718, 3663 (713 partim).

HAB. Common in the Central Province.

## 20. RHINACANTHUS, N. ab E.

1. **R. communis**, N. ab E. in Wall. Pl. As. Rar. iii. p. 109; DC. Prod. xi. p. 442; Wight, Ic. t. 464.—*R. Rottlerianus*, N. ab E. l. c.—c.p. 1982.

HAB. Very abundant in the hotter parts of the island.

## 21. ROSTELLULARIA, Reichen.

1. **R. Royeniana**, N. ab E. in DC. Prod. xi. p. 373, cum syn.—*R. rotundifolia*, N. ab E. in DC. l. c. p. 370. *R. procumbens*, N. ab E. l. c. p. 371, et Wight, Ic. t. 1539. *R. Abyssinica*, Brongn. *R. mollissima*, N. ab E. l. c. p. 373. *R. crinita*, N. ab E. l. c. *R. simplex*, Wight, Ic. t. 1542 (*T. A.*).—c.p. 142, 228, 404, 2904, 509.

HAB. Very abundant throughout the island.

I have failed in finding any character by which to separate these species even as varieties; I am therefore forced to consider them as states of one very variable species.—*T. A.*

2. **R. diffusa**, N. ab E. in Wall. Pl. As. Rar.; DC. l. c. 371, cum syn.—c.p. 1976, 1992.

HAB. Common in the hot, drier parts of the island.

## 22. PERISTROPHE, N. ab E.

1. **P. tinctoria**, N. ab E. in Wall. Pl. As. Rar.; DC. Prod. xi. p. 493, cum syn.

HAB. Ceylon, *Walker, in Herb. Hook.*

## 23. MONOTHECIUM, Hochst.

1. **M. aristatum**, T. Anders.—*Anthocometes aristatus*, N. ab E. in DC. Prod. xi. p. 312, cum syn.—c.p. 1984.

HAB. Hot, drier parts of the island; not very common.

*Corolla albida, labio inferiore fauce rufo maculato.*—*G. Thw.*

## 24. RUNGIA, N. ab E.

1. **R. parviflora**, N. ab E. in Wall. Pl. As. Rar. (DC. Prod. xi. p. 469, cum syn.)—*R. pectinata*, N. ab E. in DC. l. c. p. 470; Wight, Ic. t. 1547.

*R. muralis*, Royle. *R. polygonoides*, N. ab E. in Wall. Pl. As. Rar.; DC. l. c. p. 471. *R. organoides*, N. ab E. in Wall. Pl. As. Rar.; DC. l. c. (*T. A.*).—c.p. 257, 1975, 3354, 3357.

HAB. Very abundant up to an elevation of 7000 feet.

Like most of the species of this genus, this plant varies much in the size and shape of the leaves and the form of the spikes of flowers, as well as in the extent of the scarious margin of the bracts, and also in the extent of ciliæ on the calyx-segments.—*T. A.*

2. *R. repens*, N. ab E. in Wall. Pl. As. Rar.; DC. l. c. p. 472, cum syn.; Wight, Ic. t. 465.—c.P. 257, 1972, 1973, 1974, 2935.

HAB. Very common in the hot, drier parts of the island.

Like the preceding species, this is very variable. There is a state of it from the Central Province in which the bracts are acutely lanceolate, and their scarious margin nearly absent.—*T. A.*

3. *R. longifolia*, N. ab E. et W. Arn. in DC. l. c. p. 471.—*R. latior*, N. ab E.; DC. l. c. p. 472.—c.P. 89, 257, 301, 686, 708, 1974, 3355.

HAB. Ramboda and Adam's Peak.

## 25. DICLIPTERA, Juss.

1. *D. bivalvis*, Juss. in Ann. Mus. (DC. Prod. xi. p. 475.)—*D. Zeylanica*, N. ab E. in DC. l. c. 474.—c.P. 399, 2576.

HAB. Not uncommon on shaded banks, in the hot, drier parts of the island.

c.P. 2576 is the normal state of this species; c.P. 399 is probably only a stunted condition of it.—*T. A.*

## 26. ERANTHEMUM, Linn.

1. *E. crenulatum*, Wall. in Bot. Reg. t. 879 (DC. Prod. xi. p. 453.)—c.P. 1703, 1979.

HAB. Not uncommon.

2. *E. Ecbolium*, T. Anders.—*Justicia*, Linn.; N. ab E. in DC. l. c. p. 426, cum syn.; Wight, Ic. t. 463. *J. gymnostachya*, N. ab E. in DC. l. c. *J. lætevirens*, Vahl, En. i. p. 118? *J. strobilifera*, Lam. Ill. i. p. 36? *J. emarginata*, N. ab E. in DC. l. c. p. 427. *J. rotundifolia*, N. ab E. in Wall. Pl. As. Rar. iii. p. 108. *J. syringifolia*, Vahl, En. i. p. 117? *J. livida*, Wall. Cat. n. 2482 c. *J. dentata*, Klein. (*T. A.*)—c.P. 1978.

HAB. Common in the hot, drier parts of the island; a very variable and widely diffused plant.

## 27. PTYSSIGLOTTIS, T. Anders.

*Calyx* profunde 5-partitus, æqualis. *Corolla* infundibuliformis; *limbus* 5-partitus, *labio* inferiore longitudinaliter plicato, *lobis* lateralibus externis in æstivatione. *Stamina* duo. *Antheræ* biloculares, *loculis* parallelis, muticis, ovatis. *Stigma* capitatum, clavatum, obtusum. *Capsula* basi dissepimentis adnatis, apice tetrasperma. *Semina* discoidea, minute tuberculata (*T. A.*).

1. *P. radicata*, T. Anders.—*Rhytiglossa radicata*, N. ab E. in DC. Prod. xi. p. 344. *Rostellularia sarmentosa*, Zollinger, Pl. Jav.; DC. l. c. p. 370 (*T. A.*)—c.P. 3538.

HAB. Hot, drier parts of the island.

*Corolla* albida, 6 lin. longa, *labio* superiore bipartito; *segmentis* oblongis, subacutis, inferiore tripartito, *segmentis* rotundatis.—*G. Thw.*

## 28. ASYSTASIA, Blume.

1. *A. Gangetica*, T. Anders.—*A. Coromandeliana*, N. ab E. in Wall.

Pl. As. Rar.; DC. Prod. xi. p. 165, cum syn.; Wight, Ic. t. 1506. *A. plumbaginea*, N. ab E. in Wall. Pl. As. Rar. *A. quaterna*, N. ab E. in DC. l. c. p. 166. *A. intrusa*, N. ab E. in DC. l. c. *A. calycina*, Benth. Fl. Niger. p. 478. *A. Bojeriana*, N. ab E. in DC. l. c. *A. Comorensis*, Bojer; DC. l. c. (*T. A.*)—C.P. 1998.

HAB. Very abundant in the hotter parts of the island.

The extensive distribution of this species over tropical Asia and Africa renders the specific name adopted by Nees von Esenbeck quite inappropriate; and the same objection may be made to the original one given by Linnæus, and which, as being the oldest, I have revived. For so cosmopolitan a plant the specific term *communis* or *vulgaris* would perhaps be more suitable. The notes by Mr. Benthams to the *Asystasiæ* of the 'Flora Nigretiana' show that I have the best authority for the greater part of the identifications I have made of this variable species.—*T. A.*

2. *A. chelonoides*, N. ab E. in Wall. Pl. As. Rar.; DC. l. c. p. 164, cum syn.—*A. nemorum*, N. ab E. l. c. et DC. l. c. p. 167, cum syn. (*T. A.*)—C.P. 552, 1989.

HAB. Very abundant up to an elevation of 3000 feet.

Mr. Thwaites observes that the somewhat procumbent habit, smaller flowers and shorter pedicels, seem sufficiently to distinguish *A. nemorum* from *A. chelonoides*, and that when growing they look very different; but I have been unable to find any sufficient characters amongst the large series of specimens contained in the Hookerian Herbarium, and moreover Nees and others have partially united them.—*T. A.*

## CXVIII. LABIATÆ.

### TRIB. OCIMOIDEÆ.

#### 1. OCIMUM, Linn.

1. *O. canum*, Linn. (Benth. in DC. Prod. xii. p. 32, cum syn.)—C.P. 3624.

HAB. Very abundant in waste ground about native gardens. Nom. vulg. "Heen-talla."

*O. Basilicum*, Linn., is very common as a cultivated plant in native gardens. Is it really specifically distinct from *O. canum*?

2. *O. gratissimum*, Linn. (Benth. l. c. p. 34, cum syn.)—C.P. 2062.

HAB. Common in the warmer parts of the island.

3. *O. suave*, Willd. (Benth. l. c. p. 35, cum syn.)—C.P. 3623.

HAB. Ambagamowa; not common.

4. *O. sanctum*, Linn. (Benth. l. c. p. 38, cum syn.)—*O. tenuiflorum*, Linn.; Benth. l. c. p. 39.—C.P. 249.

HAB. Common in waste places. Nom. vulg. "Madooroo-tallu."

Much used as a medicine by the Cinghalese.

#### 2. GENIOSPORUM, Wall.

1. *G. elongatum*, Benth. Lab. p. 21 (DC. Prod. xii. p. 45.)—C.P. 2080.

HAB. Common in grassy places in the Central Province, at an elevation of 4000 to 6000 feet.

2. **G. prostratum**, Benth. in Wall. Pl. As. Rar. ii. p. 18 ; DC. l. c.—c.P. 2079.

Var.  $\beta$ . *gracile*.—*G. gracile*, Benth. Lab. p. 21 ; DC. l. c.—c.P. 2078 (2081).

HAB. Warmer parts of the island ; not uncommon.

I can find no constant character to distinguish *G. gracile* from *G. prostratum*, and they appear to pass insensibly into one another.

### 3. MOSCHOSMA, Reich.

1. **M. polystachyum**, Benth. in Wall. Pl. As. Rar. ii. p. 13 ; DC. Prod. xii. p. 48, cum syn.—c.P. 2056.

HAB. Hotter parts of the island, not uncommon.

### 4. ORTHOSIPHON, Benth.

1. **O. diffusus**, Benth. in DC. Prod. xii. p. 50.

HAB. Ceylon, *Walker, in Herb. Hook.*

2. **O. glabratus**, Benth. in Wall. Pl. As. Rar. ii. p. 14 ; DC. l. c. cum syn.—c.P. 2059.

HAB. Common in the hot, drier parts of the island.

This varies very much in the size of the flowers.

### 5. PLECTRANTHUS, L'Her.

1. **P. capillipes**, Benth. in DC. Prod. xii. p. 57.—c.P. 2064.

HAB. Central Province, at an elevation of 3000 to 5000 feet.

*Corolla* alba, labio inferiore rubro-maculato.

2. **P. nigrescens**, Benth. Lab. p. 710 ; DC. l. c. p. 58.—*P. Walkeri*, Benth. l. c. *P. scabrellus*, Benth. l. c. *P. hians*, Benth. l. c. p. 57.—c.P. 90, 425 (461, 583, 687, 2060, 2061).

HAB. Central Province, at an elevation of 4000 to 7000 feet.

*Corolla* alba, rubro-maculata.

A most variable plant, as regards the size and shape of the leaves, and the amount of pubescence.

3. **P. Coetsa**, Don, (Pr. Fl. Nep. p. 117 ; Benth. l. c. p. 58.)—*P. menthoides*, Benth. l. c. p. 59. *P. Macraei*, Benth. l. c. ; Wight, Ic. t. 1430.—c.P. 2063.

HAB. Central Province, at an elevation of 4000 to 6000 feet.

4. **P. subincisus**, Benth. in Wall. Pl. As. Rar. ii. p. 16 ; DC. Prod. xii. p. 66.—c.P. 2890.

HAB. Ceylon, *Macrae*.

I have never met with this species in a growing state, and there is no record of whence the very indifferent specimen in the Herbarium was obtained.

5. **P. Gardneri**, Thw. ; *caule* herbaceo, breve piloso, basi decumbente ; *foliis* ovatis vel rotundatis, grosse dentatis vel subincisis, crenatis, minute ciliatis, superne villosulis, subtus glanduloso-punctatis, basi subacutis, petiolo longiusculo nervisque tomentellis ; *racemis* elongatulis, glanduloso-tomentellis, parce ramosis ; *verticillastris* distantibus, 4–6-floris ; *bracteis* minutis, ciliatis ; *pedicellis* calyce fructifero nutante subæquilongis.—c.P. 16 (84, 264).

HAB. In damp, shady places, as Newera Ellia and other of the more elevated parts of the Central Province.

1-2-pedalis, plus minus rufescens. *Calyx fructifer* subglaber, prominentim parce reticulatus, 2 lin. longus, dente supremo rotundato, obtuso, inferioribus lanceolatis, infimis subulato-acuminatis. *Corolla* purpurea, calyce duplo longior, lacinia infima porrecta concava. (*P. Sieberi* affinis, *H. f.*)

6. **P. Zeylanicus**, Benth. Lab. p. 36; DC. l. c.—c.p. 2083.

HAB. Rambodde, in the Central Province, *Gardner*. Nom. vulg. "Ereweriya."

This is commonly cultivated in native gardens as a medicinal plant, but I have never found it truly wild.

*Plect. tuberosus*, Blume (*Coleus tuberosus* and *C. parviflorus*, Benth.), is cultivated by the Cinghalese for the sake of its tuberous roots, which are eaten as a vegetable.

## 6. COLEUS, Lour.

1. **C. barbatus**, Benth. in Wall. Pl. As. Rar. ii. p. 15; DC. Prod. xii. p. 71, cum syn.; Wight, Ic. t. 1432.—c.p. 2069.

HAB. Common on rocky places in the Oovah District, at an elevation of 2000 to 5000 feet.

2. **C. Malabaricus**, Benth. in Wall. Pl. As. Rar. ii. p. 16; DC. l. c. p. 76, cum syn.—*C. ovatus*, Benth. in DC. l. c. *C. Macraei*, Benth. l. c. p. 77. *C. Walkeri*, Benth. l. c. *C. leptostachys*, Benth. l. c. *C. mollis*, Benth. l. c.—c.p. 17, 2066, 2995, 3435 (2067).

HAB. Common in damp, bushy places, up to an elevation of 6000 feet.

A most variable species as respects amount of pubescence, size and form of the leaves and of the inflorescence; and the several forms run so insensibly into one another that I am quite unable to separate them satisfactorily, even as distinct varieties.

3. **C. inflatus**, Benth. Lab. p. 58; DC. Prod. xii. p. 78.—*C. Benthamianus*, Arn. in Nov. Act. Nat. Cur. xviii. p. 354; Benth. in DC. l. c.—c.p. 2065.

HAB. Abundant at Newera Ellia, Horton Plains, etc., at an elevation of 6000 to 8000 feet.

*Col. aromaticus*, Benth. (nom. vulg. "Kopprawalleya") is cultivated by the Cinghalese as a medicine for cattle. *C. Blumei*, Benth., with its variety *C. laciniatus*, Benth., is a common ornamental plant in the island.

## 7. ANISOCHILUS, Wall.

1. **A. carnosus**, Wall. Pl. As. Rar. ii. p. 18; Benth. in DC. Prod. xii. p. 81, cum syn.—c.p. 2055.

HAB. Warmer parts of the island, not very common. Nom. vulg. "Gal-kopprawalleya."

2. **A. paniculatus**, Benth. l. c. p. 82.

HAB. Ceylon, *Walker*, in *Herb. Hook.*

3. **A. suffruticosus**, Wight, Ic. t. 1437.—c.p. 3573.

HAB. Gunner's Quoin, in the Batticaloa District.

In the present plant the spikes of flowers are much shorter, the leaves are tomentose (not villous), and not so crowded together as in the plant figured by Dr. Wight, but from their general similarity I think they are no doubt forms of one species.

## TRIB. SATUREIÆ.

## 8. POGOSTEMON, Desf.

1. **P. Heyneanus**, Benth. in Wall. Pl. As. Rar. i. p. 31; DC. Prod. xii. p. 153, cum syn.; Wight, Ic. t. 1440.—c.p. 33 (2073, 2401).

HAB. Common throughout the island. Nom. vulg. "Gang-Kolangkola."

The leaves of this plant are used by the Cinghalese medicinally, as a sternutatory.

2. **P. rupestris**, Benth. Lab. p. 156; DC. l. c. p. 154.—c.p. 343.

Var.  $\beta$ . *hirsutus*.—*P. hirsutus*, Benth. in DC. l. c.—c.p. 283.

HAB. Central Province, at an elevation of 5000 to 8000 feet.

Although the extreme forms of var.  $\alpha$  and  $\beta$  of this species look very distinct, they are so completely connected by intermediate ones that I find it quite impossible to discover any constant character by which to separate them. The plant figured by Dr. Wight (Ic. t. 1442) under the name of *P. hirsutus* must certainly be, as he himself suspects, quite a distinct species.

3. **P. reflexus**, Benth. in DC. l. c. p. 155.—c.p. 154.

HAB. The more elevated parts of the Central Province, not very common.

## 9. DYSOPHYLLA, Blume.

1. **D. auricularia**, Bl. (Benth. in DC. Prod. xii. p. 156, cum syn.; Wight, Ic. t. 1445.)—c.p. 26.

HAB. Very common on the borders of paddy-fields. Nom. vulg. "Kamanilla."

2. **D. verticillata**, Benth. in Wall. Pl. As. Rar. i. p. 30; DC. l. c. p. 157, cum syn.—*D. crassicaulis*, Benth. l. c. p. 158. *D. tetraphylla*, Wight, Ic. t. 1444.—c.p. 2082.

HAB. Hotter parts of the island, not uncommon.

## 10. MENTHA, Linn.

1. **M. arvensis**, Linn. (Benth. in DC. Prod. xii. p. 171, cum syn.)—*M. Javanica*, Blume; Benth. l. c. cum syn.—c.p. 2077.

HAB. Borders of paddy-fields, in the Central Province. Nom. vulg. "Odo-talan."

## 11. CALAMINTHA, Benth.

1. **C. Clinopodium**, Benth.—Var. *umbrosa*, H. f. MS. *C. umbrosa*, Benth. in DC. Prod. xii. p. 232. *C. repens*, Benth. l. c. p. 233, cum syn. *Melissa umbrosa*, Wight, Ic. t. 1447.—c.p. 2058.

HAB. Newera Ellia.

## TRIB. STACHYDEÆ.

## 12. SCUTELLARIA, Linn.

1. **S. violacea**, Heyne, (Wall. Pl. As. Rar. i. p. 66; Benth. in DC. Prod. xii. p. 418, cum syn.; Wight, Ic. t. 1449.)—*S. floribunda*, Benth. l. c. *S. robusta*, Benth. l. c.—c.p. 217, 2716 (527).

Var.  $\beta$ ; *tota planta* glabra vel subglabra; *foliis* acutius dentatis.—  
An *S. oblonga*, var.  $\gamma$ , Benth. l. c. ?—c.p. 3370.

Var.  $\gamma$ ; *caule* repente, radicante, patentim piloso; *foliis* rotundatis, subacutis, crenatis, villosulis; *floralibus* majoribus quam in var. *a* et  $\beta$ .—c.p. 302.—An species distincta?

HAB. Very common in the Central Province, up to an elevation of 7000 feet. Var.  $\gamma$ . Horton Plains.

A very variable plant in the size of the flowers and in the amount and character of the pubescence, which latter upon the racemes varies from very glandular to being nearly or quite free from glands. Var.  $\gamma$  may possibly prove to be a distinct species, but I am disposed to believe it to be only a variety.

2. *S. oblonga*, Benth. l. c.—c.p. 2057.

HAB. Common on the rocky margins of streams in the Central Province, up to an elevation of 7000 feet.

All my specimens have the leaves with their base narrowed gradually into the petiole, and therefore belong to Bentham's var.  $\beta$  of this species, and which would seem to be the typical form of it.

### 13. ANISOMELES, R. Br.

1. *A. ovata*, R. Br. in Ait. Hort. Kew.; Benth. in DC. Prod. xii. p. 455, cum syn.; Wight, Ic. t. 865.—c.p. 2072.

HAB. Very common up to an elevation of 3000 feet. Nom. vulg. "Yakwanassa."

2. *A. Malabarica*, R. Br. in Bot. Mag. t. 2071; Benth. l. c. p. 456, cum syn.; Wight, Ic. t. 864.—c.p. 3572.

HAB. Hot, drier parts of the island.

### 14. LEUCAS, Benth.

1. *L. mollissima*, Wall. Pl. As. Rar. i. p. 62; Benth. in DC. Prod. xii. p. 525.—*L. pilosa*, Benth. in Wall. l. c.—c.p. 2075.

HAB. North of the island, *Gardner*.

I suspect that this is a mere state or variety of the following species, as the difference between the two plants is scarcely sufficient to be considered specific.

2. *L. marrubioides*, Desf. (Benth. in DC. Prod. xii. p. 528.)—c.p. 2071.

HAB. Very common in the warmer parts of the island. Nom. vulg. "Sooddoo-toomba."

3. *L. pubescens*, Benth. Lab. p. 610; DC. Prod. p. 528.—An *L. angularis*, Benth. in Wall. Pl. As. Rar. i. p. 62; DC. Prod. p. 526?

HAB. Ceylon, *Macrae* (Herb. Hook.).

The species of this genus being so variable, may not this be a form only of the preceding?

4. *L. procumbens*, Desf. (Benth. in DC. Prod. xii. p. 527.)—*L. biflora*, Br.; Benth. l. c.; Wight, Ic. t. 866.—c.p. 2053, 2074.

HAB. Very abundant throughout the island, from the level of the sea up to 8000 feet.

Tota planta molliter pubescens vel strigosula. *Calyx* omnino pubescens vel inter costas glaber, 3–4 lin. longus.

5. *L. Zeylanica*, R. Br. (Prod. p. 504; Benth. l. c. p. 531.)—*L. as-*

*pera*, Spr. Benth. l. c. p. 532. *L. diffusa*, Benth. l. c. cum syn. *L. hyssopifolia*, Benth. l. c. *L. Walkeri*, Benth. l. c.—C.P. 2076.

Cf. *L. dimidiata*, Spr., and *L. linifolia*, Spr.

HAB. Very common throughout the island. Nom. vulg. "Gatta-toomba."

A most variable species. The Cinghalese use the leaves as a medicine, and in some parts of the island, I am told, as a vegetable in their curries.

### 15. LEONOTIS, R. Br.

1. *L. nepetæfolia*, R. Br. (Prod. p. 504; Benth. in DC. Prod. xii. p. 535, cum syn.; Wight, Ic. t. 867.)—C.P. 2070.

HAB. Hot, drier parts of the island. Nom. vulg. "Maha-yakwan-assa."

#### TRIB. AJUGÆÆ.

### 16. TEUCRIUM, Linn.

1. *T. tomentosum*, Heyne, (Benth. in Wall. Pl. As. Rar. i. p. 58; DC. Prod. xii. p. 582; Wight, Ic. t. 1458.)—C.P. 212.

HAB. Central Province, at an elevation of 5000 to 7000 feet.

## CXIX. VERBENACEÆ.

#### TRIB. VERBENÆÆ.

### 1. BOUCHEA, Cham.

1. *B. Hyderabadensis*, Walp. Rep. iv. p. 12; Schauer in DC. Prod. xi. p. 559, cum syn.; Wight, Ic. t. 1462.—C.P. 3574.

HAB. On an open grassy spot between Madamahanewera and Aloo-newera, in the Central Province.

1-2-pedalis. Corolla roseo-purpurea, 8-9 lin. in exp., ad faucem minute pilosula et intensius colorata; lobis oblongis rotundatis, inferiore latiore, sæpe retuso. Antherarum 2 superiores parum majores. Stylus apicem prope abrupte deflexus. Stigma parvum. Cocci lineares, 4 lin. longi, membranaceo-marginati, externe striati, interne minute tuberculati.

### 2. STACHYTARPHA, St.

1. *S. Indica*, Vahl, Enum. i. p. 206; Sch. in DC. Prod. xi. p. 564.—C.P. 3153.

HAB. Very common throughout the island. Nom. vulg. "Balloona-koota."

*S. JAIMAICENSIS* is a very common weed in cultivated ground, having become quite naturalized in the island.

### 3. LIPPIA, Linn.

1. *L. nodiflora*, Rich. (Sch. in DC. Prod. xi. p. 585, cum syn.; Wight, Ic. t. 1463.)—C.P. 1947.

HAB. Very abundant near the sea. Nom. vulg. "Herimanadatta."



4. **LANTANA**, Linn.

1. **L. alba**, Mill. (Sch. in DC. Prod. xi. p. 606, cum syn.)—*Lantana Indica*, Roxb.; Wight, Ic. t. 1464.—c.p. 498.

HAB. Near Galagama, in the Central Province.

*Corolla* (in meis specimin.) pallide purpurea.

**L. MIXTA**, Linn. (*L. aculeata*, Hort.), introduced into the island about thirty-five years ago, has become not only quite naturalized, but covers many hundred acres of uncultivated land in many parts of the island. *Lant. trifolia*, Linn., is also not an uncommon weed in cultivated ground.

TRIB. *VITICEÆ*.5. **SYMPHOREMA**, Roxb.

1. **S. involucreatum**, Roxb. (Sch. in DC. Prod. xi. p. 621; Wight, Ic. t. 362.)—c.p. 3645.

HAB. Near Trincomalie, *Rev. S. O. Glenie*.

6. **PREMNA**, Linn.

1. **P. purpurascens**, Thw.; scandens; *ramulis* teretibus, striatis; *foliis* glabris, oblongis, abrupte breve et obtuse acuminatis, subcordatis, subtus minutissime reticulatis, petiolatis; *paniculis* tomentellis, terminalibus, divaricato-corymbosis, multifloris, laxis; *floribus* minutis; *calyce* subintegro; *corolla* subæqualiter 4-loba, lobis rotundatis, reflexis; *staminibus* exsertis, filamentis æqualibus, fauce insertis.—*Phænicanthus Zeylanicus*, Thw. in Herb. Hook.—c.p. 2698.

HAB. Ambagamowa District, at no great elevation.

*Folia* 3-6-poll. longa, 1½-3 poll. lata, læte viridia, venis cum petiolo ½-2 poll. longo purpurascens. *Paniculae* folio multum breviores, purpureæ. *Corolla* pallide purpurea.

2. **P. serratifolia**, Linn. (Sch. in DC. Prod. xi. p. 632, cum syn.)—*P. integrifolia*, Linn.; Wight, Ic. t. 1469.—c.p. 1953.

HAB. Very common in the hotter parts of the island. Nom. vulg. "Middee-gass."

The root of this plant is used medicinally by the Cinghalese

3. **P. latifolia**, Roxb. Fl. Ind. iii. p. 76; Sch. l. c. p. 635; Wight, Ic. t. 869.—An *P. viburnoides*, Wall. Sch. l. c. ?—c.p. 2893.

HAB. Trincomalie, *Rev. S. O. Glenie*. Cultura. Nom. vulg. "Middee-gass."

Like the preceding, of which this is possibly merely a variety, the present plant is employed as a medicine by the natives.

4. **P. micrantha**, Schauer, l. c. ?—An *P. mucronata*, Roxb. Sch. l. c. ?—c.p. 2541.

HAB. Maturatte District.

Arborea, 8-10-pedalis. *Ramuli juniores* pubescentes. *Folia* integra vel irregulariter serrata, basi angustata vel rotundata, superne puberula, subtus cum petiolo pubescentia. *Corolla* flavida, interne fulvo-fincta.

Though this does not quite agree with the description of either of the two species quoted above, it is not improbably one of them; and I hesitate, with the materials I have, to describe it under a new specific name.

5. **P. tomentosa**, Willd. (Sch. in DC. l. c. p. 634, cum syn.; Wight, Ic. t. 1468.)—C.P. 747.

HAB. Common up to an elevation of 3000 feet. Nom. vulg. "Boo-sairoo-gass."

6. **P. procumbens**, Wall. (Sch. l. c. p. 636.)—C.P. 325.

HAB. Hotter parts of the island, not uncommon.

7. **P. cordifolia**, Roxb. Fl. Ind. iii. p. 78; Sch. l. c. p. 632; Wight, Ic. t. 1483.—C.P. 193.

HAB. Hot, drier parts of the island.

*Folia* subglabra vel subtus tomentosa. *Corolla* flavida, plus minus rubro tincta.

### 7. CALLICARPA, Linn.

1. **C. Wallichiana**, Walp. Rep. iv. p. 125; Sch. in DC. Prod. xi. p. 641, cum syn.; Wight, Ic. t. 1480.—C.P. 722.

HAB. Very common up to an elevation of 4000 feet. Nom. vulg. "Eela-gass."

### 8. CLERODENDRON, Linn.

1. **C. inerme**, R. Br. in Ait. Hort. Kew.; Sch. in DC. Prod. xi. p. 660, cum syn.—C.P. 1949.

HAB. Very common near the sea. Nom. vulg. "Wal-gooranda."

2. **C. phlomoides**, Linn. (Sch. l. c. p. 663, cum syn.; Wight, Ic. t. 1473.)—C.P. 1950.

HAB. Hotter parts of the island, not very common.

3. **C. serratum**, Spr. (Sch. l. c. p. 664, cum syn.; Wight, Ic. t. 1472.)—C.P. 1951.

HAB. Not uncommon up to an elevation of 2000 feet. Nom. vulg. "Ken-handa."

4. **C. Linnæi**, Thw.—An *Volkameria scandens*, Linn. f. Supp. p. 292; Sch. l. c. p. 657?—C.P. 1948.

HAB. Hot, drier parts of the island.

Scandens. *Ramuli juniores* subteretes, pilosi. *Folia* obovato-rotundata, subcordata, apiculata, superne pilosula, subtus pallidiora, glanduloso-punctata et præcipue ad nervos pilosa, 2-3 poll. longa. *Petiolo* 2-3 lin. longo. *Paniculæ* terminales, foliosæ, corymbiformes, compactæ, incano-pilosæ. *Pedicelli* subnulli. *Calyx* hypocrateriformis, breve 5-dentatus, in anthesi  $1\frac{1}{2}$  lin. longus. *Corolla* alba, externe pilosa, *tubo* 3 lin. longo, *limbo* 5 lin. in exp., lobis oblongis obtusis, inferiore parum longiore. *Filamenta* valde elongata.

This is most probably, I think, *Volk. scandens*, Linn. f.; but, as the description of the latter does not quite accord with our plant, and as the name of *Cler. scandens* has been applied by Palisot de Beauvois to another species of the genus, I have thought it best to call the present *Cler. Linnæi*.

5. **C. infortpnatum**, Linn. (Sch. l. c. p. 667, cum syn.; Wight, Ic. t. 1471.)—C.P. 2894.

HAB. Not uncommon in damp forests, up to an elevation of 5000 feet. Nom. vulg. "Gas-pinna."

*C. SIPHONANTHUS*, R. Br., is common in gardens, but is not truly indigenous.

9. **GMELINA**, Linn.

1. **G. Asiatica**, Linn. (Sch. in DC. Prod. xi. p. 679, cum syn.)—*G. parvifolia*, Roxb. Pl. Cor. ii. t. 162; Sch. l. c. cum syn.—c.p. 1952.

HAB. Very common up to an elevation of 2000 feet. Nom. vulg. "Gatta-demmata."

2. **G. Rheedei**, Hook. Bot. Mag. t. 4395; Walp. Ann. iii. p. 239.—*G. arborea*, Wight, Ic. t. 1470.—An *G. arboreæ*, Roxb., var. ?—c.p. 128 (698).

HAB. Common up to an elevation of 5000 feet. Nom. vulg. "At-demmata."

Arbor ingens, 40-50-pedalis, patula.

The bark and roots of this and the preceding species are used medicinally by the Cinghalese.

10. **VITEX**, Linn.

1. **V. trifolia**, Linn. (Sch. in DC. Prod. xi. p. 683, cum syn.)—c.p. 1955.

HAB. Near the sea, not uncommon.

2. **V. Negundo**, Linn. (Sch. l. c. p. 684, cum syn.; Wight, Ic. t. 519.)—c.p. 1956.

HAB. Common on the banks of rivers and streams, up to an elevation of 3000 feet. Nom. vulg. "Sooddoo-nikka-gass."

3. **V. altissima**, Linn. f. (Sch. l. c. p. 685, cum syn.; Wight, Ic. t. 1466.)—An *V. alata*, Heyne, Sch. l. c. ?—c.p. 1958.

HAB. Common in forests, up to an elevation of 3000 feet. Nom. vulg. "Mililla-gass," "Meeyan-mililla-gass."

*Folia* subglabra vel subtus molliter pubescentia, *petiolo* sæpe plus minus alato.

This tree produces one of the most valuable timbers in the island for building and other purposes.

4. **V. pubescens**, Vahl, (Sch. l. c. cum syn.; Wight, Ic. t. 1465.)

HAB. Ceylon, *Walker*, (*Herb. Hook.*)

5. **V. leucoxylon**, Linn. f. (Sch. l. c. p. 692, cum syn.; Wight, Ic. t. 1467.)—*V. saligna*, Roxb. Sch. l. c.—c.p. 1957.

HAB. Hotter parts of the island, not uncommon.

TRIB. *AVICENNIEÆ*.11. **AVICENNIA**, Linn.

1. **A. officinalis**, Linn. (Sch. in DC. Prod. xi. p. 700, cum syn.; Wight, Ic. tt. 1481, 1482.)—c.p. 1961 (1962).

HAB. Not uncommon near the sea.

CXX. **PLUMBAGINACEÆ**.1. **PLUMBAGO**, Tourn.

1. **P. Zeylanica**, Linn. (Boissier in DC. Prod. xii. p. 692, cum syn.)—c.p. 2265.

HAB. Very common in the hotter parts of the island. Nom. vulg. "Ella-nettol."

*P. ROSEA*, Linn., called by the Cinghalese "Rat-nettol," is not uncommon in gardens, but I cannot hear of its occurring truly wild. It is much valued as a medicinal plant.

## CXXI. PLANTAGINACEÆ.

### 1. PLANTAGO, Linn.

1. *P. major*, Linn. (Var.  $\beta$  *Asiatica*, Dene. in DC. Prod. xiii.-i. p. 694.)—*P. Asiatica*, Linn.—c.p. 2246.

HAB. Common in the most elevated parts of the Central Province.

*P. LANCEOLATA*, Linn., is become a common weed at Newera Ellia, having been introduced with grass-seeds from Europe.

## CXXII. NYCTAGINACEÆ.

### 1. PISONIA, Plum.

1. *P. aculeata*, Linn.; Choisy in DC. Prod. xiii.-ii. p. 440, cum syn.—*P. villosa*, Poir. l. c. *P. Georgina*, Wall.—c.p. 2266.

HAB. Not uncommon up to an elevation of 3000 feet.

### 2. BOERHAAVIA, Linn.

3. *B. diffusa*, Linn.; Choisy in DC. Prod. xiii.-ii. p. 452, cum syn.—*B. procumbens*, Roxb.; Wight, Ic. t. 874.—c.p. 1099.

HAB. Very common on sandy or gravelly places. Nom. vulg. "Pittasooddoo-pala."

## CXXIII. POLYGONACEÆ.

### 1. POLYGONUM, Linn.

1. *P. barbatum*, Linn. (Meisner in DC. Prod. xiv. p. 104, cum syn.; Wight, Ic. t. 1798.)—c.p. 3000.

HAB. Warmer parts of the island, not uncommon.

2. *P. Posumbu*, Ham. in Don Pr. Fl. Nep. p. 71; Wall. Cat. N.; an Meisner?—*P. strictum*, Allioni; Wight, Ic. t. 1800.—c.p. 2591.

HAB. Dimboola, in the Central Province, at an elevation of 4000 feet.

3. *P. serrulatum*, Lag. (Meisn. l. c. p. 110, cum syn.)—c.p. 2999.

HAB. Galle.

This is considered by Dr. Hooker to be merely a variety of *P. mite*, Schr.

4. *P. glabrum*, Willd. (Meisn. l. c. p. 114, cum syn.; Wight, Ic. t. 1799.)—c.p. 2907.

HAB. Very common up to an elevation of 3000 feet.

Dr. Hooker believes this to be a variety of *P. Persicaria*, Linn.

5. **P. tomentosum**, Willd. (Meisn. l. c. p. 124, cum syn.)—C.P. 2248.  
HAB. Very common in the warmer parts of the island.
6. **P. perforatum**, Meisn. l. c. p. 128, cum syn.—*P. Nepalense*, Meisn. l. c. ; Wight, Ic. t. 1804.—C.P. 2251.  
HAB. Very abundant throughout the island.
7. **P. Chinense**, Linn. (Meisn. l. c. p. 130, cum syn. ; Wight, Ic. t. 186.)—C.P. 2249.  
HAB. A very common plant.
8. **P. strigosum**, R. Br. (Meisn. l. c. p. 134, cum syn.)—*P. muricatum*, Meisn. l. c. p. 133.—C.P. 2250.  
HAB. Most elevated parts of the Central Province.
9. **P. pedunculare**, Wall. (Meisn. l. c. p. 133, cum syn.)—C.P. 3348.  
HAB. South of the island.

## CXXIV. SALSOLACEÆ.

## 1. CHENOPODIUM, Tourn.

1. **C. murale**, Linn. (Moquin in DC. Prod. xiii.-ii. p. 69, cum syn.)—C.P. 3658.  
HAB. Trincomalie, *Rev. S. O. Glenie*.

## 2. ATRIPLEX, Gærtn.

1. **A. repens**, Roth, (Moq. l. c. p. 99.)—C.P. 2252.  
HAB. North of the island, *Gardner*.

## 3. ARTHROCNEMUM, Moquin.

1. **A. Indicum**, Moq. in DC. Prod. xiii.-ii. p. 151, cum syn.—*Salicornia Indica*, Willd. ; Wight, Ic. t. 737.—C.P. 2262.  
HAB. North of the island.

## 4. SUÆDA, Försk.

1. **S. nudiflora**, Moq. in DC. Prod. xiii.-ii. p. 155, cum syn.—C.P. 2261.  
HAB. Putlam, *Moon*. Trincomalie, *Rev. S. O. Glenie*. Batticoloa, etc.

In my specimens the styles vary in number from 2 to 5, but there are most commonly 4.

2. **S. Indica**, Moq. l. c. p. 156, cum syn. ; Wight, Ic. t. 1796.—C.P. 2260.

HAB. Putlam, *Moon*. Jaffna, and Calpentyne, *Gardner*.

I have not seen fresh specimens in flower of this species, so have not had an opportunity of comparing it so critically as I could wish with the preceding, to which it is evidently very closely allied.

BASELLA RUBRA, Linn., is commonly cultivated in gardens, but is certainly not indigenous.

## CXXV. AMARANTACEÆ.

## 1. CELOSIA, Linn.

1. *C. polygonoides*, Retz, (Moquin in DC. Prod. xiii.-ii. p. 238, cum syn.)—C.P. 2240.

HAB. North of the island, *Gardner*.

2. *C. pulchella*, Moq. l. c. cum syn.; Wight, Ic. t. 1768.—C.P. 2238.

HAB. Common in the hot, drier parts of the island.

3. *C. argentea*, Moq. l. c. p. 242, cum syn.; Wight, Ic. t. 1767.—An *C. cristata*, Moq. l. c., var. ?—C.P. 2245.

HAB. Warmer parts of the island. Nom. vulg. "Kirri-handa."

## 2. CHAMISSOA, Kunth.

1. *C. nodiflora*, Mart. (Moq. in DC. Prod. xiii.-ii. p. 249, cum syn.; Wight, Ic. t. 1770.)—*C. albida*, Moq. l. c. cum syn.; Wight, Ic. t. 1769. *C. dichotoma*, Moq. l. c. cum syn.; Wight, Ic. t. 1771. *C. esculenta*, Moq. l. c. cum syn. An *C. aspera*, Wight, Ic. t. 1772?—C.P. 2244, 2726, 3154.

HAB. Very common in the warmer parts of the island.

After examining a large series of specimens, I have found it quite impossible to arrive at any other conclusion but that the several plants I have enumerated are mere forms of one very variable species. The shape of the leaves and amount of pubescence are most inconstant characters; the seed varies in being more or less minutely punctate; and the whole plant is thin and flaccid, or of tolerably firm consistence.

## 3. AMARANTUS, Tourn.

(Nom. vulg. "Tampala.")

1. *A. oleraceus*, Roxb. Fl. Ind. iii. p. 605; Wight, Ic. t. 715 (vix *Euxolus oleraceus*, Moquin in DC. Prod. xiii.-ii. p. 273.)—*A. Gangesicus*, Linn.; Moq. l. c. p. 261, cum syn. *A. lanceolatus*, Roxb. l. c. p. 607.—An *A. melancholicus*, Linn.; Moq. l. c. p. 262, cum syn. ?—C.P. 3629.

HAB. Very common in native gardens, but hardly indigenous.

2. *A. polygamus*, Linn. (Roxb. Fl. Ind. iii. p. 603; Wight, Ic. t. 714.)—*Euxolus polygamus*, Moq. l. c. p. 272.—An *A. tristis*, Linn.; Moq. l. c. p. 260, cum syn.; Wight, Ic. tt. 514, 713?—C.P. 3642.

HAB. Common in cultivated ground.

The smaller size and procumbent habit of *A. polygamus* appear to me to be the principal points of distinction between it and *A. oleraceus*, to which it is extremely closely allied, if it be not indeed a form of it. In my specimens the *utriculi*, when perfectly ripe, have the circumscissal dehiscence of *Amarantus*.

3. *A. spinosus*, Linn. (Moq. l. c. p. 260, cum syn.; Wight, Ic. t. 513.—C.P. 2910.

HAB. Very abundant in waste places.

*A. FRUMENTACEUS*, Ham. (an *A. cruentus*, Linn.?), *A. hypochondriacus*, Linn., and *A. caudatus*, Linn., occur as cultivated plants. All the species of *Amarantus* and *Euxolus* are used as vegetables by the Cinghalese.

4. **EUXOLUS**, Rafin.

(Nom. vulg. "Koorá-tampala.")

1. **E. caudatus**, Moquin in DC. Prod. xiii.-ii. p. 274, cum syn.; Wight, Ic. t. 1773.—C.P. 2911.

HAB. Very abundant on waste ground.

2. **E. polygonoides**.—*Amblogyne polygonoides*, Rafin.; Moq. l. c. p. 270, cum syn. *Amarantus polygonoides*, L.; Willd.; Wight, Ic. t. 512.—C.P. 3643.

HAB. Very common in cultivated ground, etc.

This, as suspected by Moquin, does not differ generically from *Euxolus*. It is very closely allied indeed to *E. caudatus*, Moq., and might perhaps be safely considered a variety of it.

5. **PSILOTRICHUM**, Blume.

1. **P. scleranthum**, Thw.; suffruticosum; *ramulis* teretibus, tomentellis; *foliis* ovato-lanceolatis, acutis, basi in petiolum brevem angustatis, pilosulis; *spicis* terminalibus, parum ramosis, obtusis; *floribus* densis; *bracteolis* late ovatis, mucronatis, 2 interioribus glabris, exteriori minore, pubescente; *sepalis* subæqualibus, rigidis, lanceolatis, acutis, obscure 5-nerviis, externe pilosis, quam bracteolæ duplo longioribus.—C.P. 2239.

HAB. Anooradhapoorá, *Gardner*.

*Folia*  $\frac{3}{4}$ – $1\frac{3}{4}$  poll. longa, 4–7 lin. lata. *Flores* 2 lin. longi, albidí.

At first sight this might be taken for a small form of *Banalia thyrsiflora*, Moq., but its simple stigma brings it within the present genus.

6. **PTILOTUS**, R. Br.

1. **P. ovatus**, Moquin in DC. Prod. xiii.-ii. p. 281, cum syn.—C.P. 2235.

HAB. Hot, drier parts of the island.

7. **ÆERVA**, Försk.

1. **Æ. Javanica**, Juss. (Moquin in DC. Prod. xiii.-ii. p. 299, cum syn.; Wight, Ic. t. 876.)—C.P. 2255.

HAB. North of the island, *Gardner*.

2. **Æ. lanata**, Juss. (Moq. l. c. p. 303, cum syn.; Wight, Ic. t. 723.)—C.P. 2253.

HAB. Very abundant on waste ground. Nom. vulg. "Pol-hoonoo-pala."

3. **Æ. Monsonia**, Moq. l. c. p. 305, cum syn.; Wight, Ic. t. 725.—*Æ. setacea*, Moq. l. c.

HAB. North of the island, *Dyke*, (*Herb. Hook.*)

8. **NOTHOSÆERVA**, Wight, Ic. vi. p. 1.

*Pseudanthus*, Wight, Ic. t. 1776 (non Lieb.).

1. **N. brachiata**, Wight. l. c.—*Ærva brachiata*, Mart. Beitr. Amaran. p. 83; Moquin, l. c. (partim).—C.P. 2254.

HAB. Common in the hot, drier parts of the island.

## 9. ACHYRANTHES, Linn.

1. **A. bidentata**, Blume, (Moquin in DC. Prod. xiii.-ii. p. 312, cum syn.; Wight, Ic. t. 1779.)—C.P. 2236.

HAB. Very abundant on waste ground.

2. **A. aspera**, Linn. (Moq. l. c. p. 314, cum syn.; Wight, Ic. t. 1777.)—C.P. 2243.

HAB. Equally abundant with the preceding, and in similar places. Nom. vulg. "Gas-karral-habba."

This plant is employed medicinally in cases of dysentery.

3. **A. argentea**, Lam. (Moq. l. c. p. 315, cum syn.)—*A. porphyrostachya*, Wall.; Moq. l. c. p. 316. *A. rubro-fusca*, Wight, Ic. t. 1778. An *A. asperæ* var. ?—C.P. 2909.

HAB. Cultura, Moon.

I am quite disposed to think, with Dr. Hooker, that this is probably merely a variety of *A. aspera*.

## 10. CENTROSTACHYS, Wall.

1. **C. diandra**, Wall. (Moq. in DC. Prod. xiii.-ii. p. 322, cum syn.; Wight, Ic. t. 722.)—C.P. 2241.

HAB. Very common in the warmer parts of the island.

## 11. DIGERA, Forsk.

1. **D. arvensis**, Forsk. (Moq. in DC. Prod. xiii.-ii. p. 324, cum syn.)—*Desmochæte muricata*, Wight, Ic. t. 732.—C.P. 2912.

HAB. North of the island, Moon.

## 12. CYATHULA, Lour.

1. **C. prostata**, Blume, (Moquin in DC. Prod. xiii.-ii. p. 326, cum syn.; Wight, Ic. t. 733.)—C.P. 2242.

HAB. Very abundant. Nom. vulg. "Bin-karral-habba."

Employed as a medicine by the Cinghalese in cases of dysentery.

2. **C. capitata**, Moquin, l. c. p. 329; Wight, Ic. t. 1782.—C.P. 3112.

HAB. Ooma Oya, near Kandy.

In my specimens the *staminodia* are much wider and more copiously fimbriated than they are represented in Dr. Wight's figure.

## 13. PUPALIA, Juss.

1. **P. atropurpurea**, Moquin in DC. Prod. xiii.-ii. p. 331, cum syn.—*Desmochæte atropurpurea*, DC.; Wight, Ic. t. 731.—C.P. 2237.

HAB. Common in the hot, drier parts of the island.

2. **P. orbiculata**, Wight, Ic. t. 1783.—*Cyathula orbiculata*, Moquin, l. c. p. 330, cum syn.—C.P. 2234.

HAB. Hotter parts of the island; not uncommon.



14. **ALTERNANTHERA**, Mart.

1. **A. sessilis**, R. Br. (Moq. in DC. Prod. xiii.-ii. p. 357, cum syn.; Wight, Ic. t. 727.)—*A. nodiflora*, R. Br.; Moq. l. c. p. 356. *A. denticulata*, R. Br.; Moq. l. c. cum syn.—c.p. 2908.

HAB. Very abundant throughout the island. Nom. vulg. "Mokoo-noo-wanna."

## CXXVI. PHYTOLACCACEÆ.

1. **GIESECKIA**, Linn.

1. **G. pharnaceoides**, Linn. (Moq. in DC. Prod. xiii.-ii. p. 27, cum syn.; Wight, Ic. t. 1167.)—c.p. 1093.

HAB. Warmer parts of the island.

*Phytolacca octandra*, Moq., occurs occasionally as an escape from gardens; and *Rivina lævis*, Linn., is become naturalized in some places.

## CXXVII. PROTEACEÆ.

1. **HELICIA**, Lour.

1. **H. Ceylanica**, Gardner in Calcutta Journ. of Nat. Hist. vol. vii. p. 453; Meisner in DC. Prod. xiv. p. 438.—c.p. 690.

HAB. Central Province, at an elevation of 2000 to 4000 feet; not very common.

*Arbor mediocris. Pedunculique pedicellique purpurei. Flores pallide flavi. Fructus ovoidei, utrinque parum acuminati, nucis avellanæ magnitudine, atro-purpurei.*

I have examined a great many flowers of this species, and have found the ovary biovulate in all, in this respect agreeing with the other species of the genus. The loculus has at its upper part two shining slight prominences, which were probably mistaken by Dr. Gardner for another pair of ovules.

## CXXVIII. THYMELACEÆ.

1. **WIKSTRÆMIA**, Endl.

1. **W. virgata**, Meisner in DC. Prod. xiv. p. 547, cum syn.—*W. canescens*, Meisner, l. c. cum syn. *W. inamæna*, Meisn. l. c. *W. salicifolia*, Dcne.; Meisn. l. c. p. 546. *Daphne inamæna*, Gardn. in Calc. Journ. of Nat. Hist. vol. vii. p. 454.—c.p. 2206.

HAB. Central Province, at an elevation of 6000 to 8000 feet.

2. **LASIOSIPHON**, Fresen.

1. **L. eriocephalus**, Dcne. (Meisn. in DC. Prod. xiv. p. 597, cum syn.)—*L. speciosus*, Dcne.; Meisn. l. c. p. 598. *Gnidia eriocephala*, Wight, Ic. t. 1859. *Gnidia (Dingia) eriocephala*, Gardner in Calcutta Journ. of Nat. Hist. vol. vii. p. 456—c.p. 469.

Var.  $\beta$ . *Zeylanicus*, Meisn. l. c. ; *Gnidia* (*Dingia*) *insularis*, Gardner, l. c.—c.p. 2913.

HAB. Drier parts of the island, up to an elevation of 4000 feet. Nom. vulg. "Naha-gass."

The two varieties of this species merge insensibly into one another. The bark is used for poisoning fish ; a very excellent fibre is also obtained from it.

### 3. GYRINOPS, Gærtn.

1. **G. Walla**, Gærtn. (Meisn. in DC. Prod. xiv. p. 602, cum syn. ; Wight, Ic. t. 1850.)—c.p. 1828.

HAB. Common in the warmer parts of the island. Nom. vulg. "Walla-gass."

The bark of this plant yields a very strong fibre.

### 4. DRYMISPERMUM, Reinw.

1. **D. cauliflorum**, Thw. ; arborea, glabra ; *foliis* nitidis, oblongo-lanceolatis, longiuscule acuminatis, basi acutis, brevi-petiolatis ; *capitulis* caulinis, brevissime pedunculatis ; *calyce* glabro, *lobis* ab marginem villosulis ; *staminibus* biserialibus ; *filamentis* subæquilongis.—c.p. 3459.

HAB. Hewessee, in the Pasdoon Corle, on the banks of a stream.

*Arbor* parva, elegans ; *ramis* elongatis, subpendulis. *Folia* 6–9 poll. longa,  $1\frac{1}{2}$ – $3\frac{1}{2}$  poll. lata. *Bracteolæ* ovatæ, acutæ, circiter 3 lin. longæ. *Flores* albidi, 8 lin. longi ; *lobis calycinis* rotundatis, obtusis. *Fructus* ovoidei, parum acuminati, nucis avellanæ magnitudine, carnosus, sanguineus.

### 5. CANSJERA, Juss.

1. **C. Rheedeii**, Gmel. (Meisn. in DC. Prod. xiv. p. 519, cum syn. ; Wight, Ic. t. 1861.)—c.p. 189.

HAB. Very common in the warmer parts of the island.

## CXXIX. SANTALACEÆ.

### 1. PYRULARIA, Mich.

1. **P. Wallichiana**, A. DC. Prod. xiv. p. 629, cum syn.—*P. Ceylanica*, A. DC. l. c. *Sphærocarya Wallichiana*, Wight, Ic. t. 241.—c.p. 2555.

HAB. Central Province, at an elevation of 4000 to 6000 feet.

Partes juniores, cum inflorescentia, rubræ.

### 2. OSYRIS, Linn.

2. **O. arborea**, Wall. (A. DC. Prod. xiv. p. 633.)—*O. Wightiana*, Wall. ; A. DC. l. c. ; Wight, Ic. t. 1853.—c.p. 53 (206, 502.)

HAB. Oova, in the Central Province, at an elevation of 5000 to 6000 feet ; abundant.

## CXXX. ELÆAGNACEÆ.

## 1. ELÆAGNUS, Linn.

1. **E. latifolia**, Linn. (Schlecht. in DC. Prod. xiv. p. 610, cum syn.; Wight, Ic. t. 1856.)—*E. conferta*, Roxb. Fl. Ind.; Schl. l. c. p. 612. *E. arborea*, Roxb. l. c.; Schl. l. c. p. 611. *E. ferruginea*, Ach. Rich.; Schl. l. c. p. 610. *E. Wallichiana*, Schl. l. c. p. 612. *E. parvifolia*, Roxb.; Schl. l. c. *E. Thwaitesii*, Schl. l. c. p. 611. *E. Kologa*, Schl. l. c.—*Foliis* subtus argenteo-leprosis, squamis ferrugineis nullis vel paucissimis intermixtis.—c.p. 2263. *Foliis* firmioribus, sæpius majoribus, subtus squamis argenteis ferrugineisque intermixtis leprosis; *floribus* parum brevioribus.—c.p. 2264, 83. *Foliis* firmis, sæpius basi rotundatis, subtus ferrugineo-leprosis.—c.p. 2724.

HAB. Very common up to an elevation of 5000 feet.

I believe we have but one species of this genus indigenous to Ceylon, but it is an extremely variable plant. When occurring in the hotter parts of the island the leaves are more membranaceous, and have their leprous scales nearly all white; higher up, on the hills, the leaves are usually larger and firmer, and there is a greater intermixture of ferruginous scales, and sometimes the scales are all ferruginous.

## CXXXI. DATISCACEÆ.

## 1. TETRAMELES, R. Br.

1. **T. nudiflora**, R. Br. in Pl. Jav. Rar. p. 19. t. 17.—*S. Horsfieldii*, Steud. Nomenc. p. 171.—c.p. 3113.

HAB. Ambagamowa District, at no great elevation; not common.

## CXXXII. LAURACEÆ.

## 1. CINNAMOMUM, Burm.

1. **C. Zeylanicum**, Breyn. (N. ab E. System. Laurin. p. 45, cum syn.; Wight, Ic. tt. 128, 129, 134.)—*C. Capparum*, Coronde; Blume; N. ab E. l. c. p. 665, cum. syn.—c.p. 2283, 2284.

Var.  $\beta$ . *multiflorum*.—*C. multiflorum*, Wight, Ic. tt. 126, 131. *C. dubium*, N. ab E. l. c. p. 41; Wight, t. 135. *C. perpetuo-florens*, Wight, Ic. t. 141. *C. villosum*, Wight, Ic. t. 127.—c.p. 37, 2282.

Var.  $\gamma$ . *ovalifolium*.—*C. ovalifolium*, Wight, Ic. t. 125.—c.p. 263, (317, 628.)

HAB. Pretty generally distributed in forests up to an elevation of 3000 feet. Var.  $\beta$ . Common in the Central Province, up to 3000 feet. Var.  $\gamma$ . At an elevation of 4000 to 8000 feet.

Without the opportunity, which I have had, of seeing numerous examples of this very variable tree, I should scarcely have ventured to unite the above-enumerated varieties under one specific name, so unlike one another are the extreme forms of each; but in a large series of specimens it is frequently difficult, or even impossible, to determine to which variety to refer some of them, so intermediate are they in character.

The best cinnamon of commerce is the produce of what I consider a cultivated or selected form of var.  $\alpha$ , and the trees of this form have usually large leaves of some-

what irregular shape; but the bark of all the forms of this species possesses the odour of cinnamon in a greater or less degree, and it is not always possible to judge of the quality of the bark from the foliage, for I have observed the cinnamon-peelers, when collecting bark from uncultivated trees, taste a small portion before commencing their operations, and pass over some trees as unfit for their purpose. The bark of varr.  $\beta$  and  $\gamma$  is of very inferior quality, and I have been informed is never collected, except for the purpose of adulteration.

A fragrant oil, sold under the name of "Clove Oil," is obtained by distillation from the leaves of the cultivated Cinnamon-tree, and the roots yield, under a similar process, a certain quantity of camphor. The ripe seeds contain some amount of stearine, but I cannot learn that it is now extracted for any purposes.

I strongly suspect that *C. obtusifolium*, N. ab E. l. c. p. 33, cum syn.; Wight, Ic. t. 139; *C. Bazania*, N. ab E. l. c. p. 36, cum syn.; *C. Malabathrum*, Batka; N. ab E. l. c. p. 38, cum syn.; *C. iners*, Wight, Ic. tt. 122, 122 bis, 230; and *C. nitidum*, N. ab E. l. c. p. 43, cum syn.; Wight, Ic. t. 124, will prove to be mere forms of the present species.

2. **C. citriodorum**, Thw.; *foliis* glabris, nitidis, coriaceis, ovatis vel oblongis, plus minus obtuse acuminatis, penniveniis vel rarius ad basin breviter 3-nerviis, venis inconspicuis; *paniculis* multifloris, elongatulis, subterminalibus, incano-pilosulis; *lobis perianthii* ad basin deciduis.—C.P. 393.

HAB. Safragam District and near Galagama, at an elevation of 1000 to 2000 feet. Nom. vulg. "Pangaree-kooroondoo-gass."

*Arbor* 20-30-pedalis, dense ramosa, compacta. *Ramuli juniores* pilosuli. *Folia*  $1\frac{1}{2}$ -3 poll. longa,  $\frac{3}{4}$ - $1\frac{1}{2}$  poll. lata; *petiolo* semipollicari. *Pedicelli* floribus breviores. *Fructus* oblongus, basi perianthii truncata cyathiformi suffultus.

Well distinguished from every form of the preceding species by the venation of its leaves and by the truncated cup of the fruit. The bark of this species has much of the odour of citronella oil, intermixed with something of the fragrance of common cinnamon.

3. **C. litsææfolium**, Thw.; *foliis* glabris, coriaceis, ovato-lanceolatis, obtusiusculis, triplinerviis (nervis lateralibus infra medium folii evanescentibus), subtus pallidioribus; *paniculis* elongatis, apertis, multifloris, subterminalibus, incano-pilosulis; *lobis perianthii* ad basin deciduis.—C.P. 392.

HAB. Hapootelle, in the Central Province, at an elevation of 5000 feet. Nom. vulg. "Kooddoo-kooroondoo-gass."

*Arbor* 50-60-pedalis, versus apicem ramosum. *Ramuli juniores* pilosuli. *Folia*  $2\frac{1}{2}$ -6 poll. longa,  $1-2\frac{1}{2}$  poll. lata; *petiolo*  $\frac{1}{2}$ - $\frac{3}{4}$  poll. longo. *Pedicelli* floribus longiores.

Nearly allied to the preceding, but the larger size and different shape and venation of the leaves sufficiently distinguish it. The bark of this species is quite inodorous.

## 2. APOLLONIAS, N. ab E.

1. **A. Zeylanica**, Thw.; arborea, glabra; *foliis* oppositis, oblongis vel oblongo-lanceolatis, parum obtuse acuminatis, valde reticulatis, basi angustatis, petiolatis; *paniculis* multifloris; *perianthio* deciduo.—C.P. 2485.

HAB. Central Province, at an elevation of 3000 to 4000 feet.

*Arbor* 50-60-pedalis. *Folia* 4-8 poll. longa,  $1\frac{1}{2}$ -3 poll. lata, *petiolo* 4-8 lin. longo. *Flores* 1 lin. longi, pallide virides. *Fructus* cylindrico-oblongus, 8 lin. longus, 3 lin. latus, pedicello incrassato, truncato, 1-2 lin. longo.

This agrees so well in most respects with the other described species of *Apollonias*,

that the deciduous perianth of the present can scarcely be considered more than a specific character. It is evidently very closely allied to *Haasia Wightii*, Nees.

### 3. MACHILUS, Rumph. ; Nees.

1. **M. macrantha**, N. ab E. Syst. Laur. p. 174, cum syn. ; Wight, Ic. t. 1824.—*M. glaucescens*, Wight, Ic. t. 1825.—c.P. 2270, (2271.)

HAB. Central and South-western Provinces, at an elevation of 1500 to 4000 feet. Nom. vulg. "Oolooloo-gass."

The timber of this large tree is useful for building purposes.

### 4. ALSEODAPHNE, N. ab E.

1. **A. semicarpifolia**, N. ab E. Syst. Laur. p. 182, cum syn. ; Wight, Ic. tt. 1826, 1827.—c.P. 2269.

HAB. Hot, drier parts of the island ; not uncommon.

*Fructus* oblongus, 8 lin. longus, 5 lin. latus ; *pedicello* incrassato, tuberculato, 6-10 lin. longo.

A very variable plant as regards the size and shape of the leaves.

### 5. CRYPTO-CARYA, R. Br.

1. **C. Wightiana**, Thw.—*C. floribunda*, Wight, Ic. t. 1829, (non N. ab E. fid. Hook. f.)—c.P. 414, (2392.)

HAB. Common up to an elevation of 5000 feet. Nom. vulg. "Goloomora-gass."

*Folia* subtus ferrugineo-puberula. *Fructus* subsphæricus, nitidus, niger, semi-pollicaris.

A large tree, useful for building purposes.

2. **C. membranacea**, Thw. ; *foliis* oblongis vel rarius ovato-oblongis, abrupte acuminatis, subglabris, membranaceis, utrinque minute prominentim reticulatis, basi acutis et parum obliquis ; *paniculis* brevibus, paucifloris, ferrugineo-tomentellis ; *fructibus* oblongo-ovoideis, ecostatis.—c.P. 2701.

HAB. Saffragam District, up to an elevation of 2000 feet.

*Arbor* mediocris. *Folia* 2½-7 poll. longa, 1-2¾ poll. lata, *petiolo* 3-4 lin. longo. *Fructus* atro-purpureus, circiter 7 lin. longus.

This would seem, judging from Nees von Esenbeck's description of his *C. amygdalina*, in Syst. Laur. p. 208, to be closely allied to that species, and to differ from it principally in having much smaller panicles of flowers.

### 6. TETRANTHERA, N. ab E.

(Cum *Lepidadenia*, N. ab E.)

1. **T. tomentosa**, Roxb. (N. ab E. Syst. Laur. p. 511, cum syn. ; Wight, Ic. t. 1834.)—c.P. 2275.

HAB. Central Province, up to an elevation of 4000 feet ; common. Nom. vulg. "Kos-badda-gass."

2. **T. ligustrina**, N. ab E. l. c. p. 513, cum syn. ; Wight, Ic. t. 1835.—c.P. 2773.

HAB. Central Province, *Macrae*.

3. **T. Roxburghii**, N. ab E. l. c. p. 515, cum syn.—C.P. 2276.

HAB. Very common up to an elevation of 3000 feet. Nom. vulg. "Bo-mee-gass."

4. **T. nemoralis**, Thw.; *caule* erecto, simplici vel parce ramoso, terete, fusco-ferrugineo-velutino; *foliis* majusculis, lanceolatis, basi apiceque acutis, utrinque (junioribus) fusco-ferrugineo-villosis, denique superne subglabris, subtus reticulatis; *umbellulis* parvis, axillaribus caulinisque, fasciculatis, glabris vel sparsissime pilosis.—C.P. 2921.

HAB. Galle District, under the shade of large trees in forests.

10–12-pedalis. *Folia* 1–2-pedalia,  $2\frac{1}{2}$ –4 poll. longa, *petiolo*  $\frac{1}{2}$ – $2\frac{1}{2}$  poll. longo. *Pedunculi* 1–2 lin. longi. *Umbellulæ* parvæ, 5-floræ. *Perianthium* 6-partitum, lobis deciduis. *Fructus* subsphæricis, apiculatis, 4 lin. in diam., *pedicello* circiter 1 lin. longo.

5. **T. longifolia**, N. ab E. l. c. p. 528, et var.  $\beta$ .—C.P. 223, 79, (230, 417, 587.)

Var.  $\gamma$ ; *foliis* glabris vel subglabris.—C.P. 2274.

HAB. Common in the Central Province, at an elevation of 4000 to 7000 feet.

*Arbor* parva, parce ramosa. *Fructus* sphæricus, 3–4 lin. in diam.

6. **T. glaberrima**, Thw.; *foliis* ovatis vel ovato-lanceolatis, parum abrupte acuminatis, minute reticulatis, basi acutis, subtus glaucis: *venis primariis* paucis, multum arcuatis; *umbellulis* parvis, axillaribus, fasciculatis, 4-floris; *perianthio* 6-partito, lobis deciduis; *fructibus* oblongis.—C.P. 2606.

HAB. Allagalla, in the Central Province, at an elevation of 4000 feet.

*Arbor* parva. *Folia* 2–7 poll. longa, 1–3 poll. lata, *petiolo* 4–8 lin. longo. *Flores* dioici.

In general appearance closely resembling the last species, from which it is distinguished by being glabrous in every part, by its wider leaves with more arcuate primary veins, and its differently shaped fruit.

7. **T. Gardneri**, Thw.; arborea; *foliis* majusculis, trapezoideo-vel obovato-oblongis, obtusis, coriaceis, superne glabris, subtus albido-tomentellis, basi acutis; *inflorescentia* axillari, racemoidea, quam folium multo brevior, albido-tomentella; *fructibus* subsphæricis.—C.P. 40.

HAB. Central Province, at an elevation of 4000 to 6000 feet.

*Arbor* 40–50-pedalis. *Folia* 4–12 poll. longa, 2–6 poll. lata; *venis primariis* parum arcuatis, superne depressis, subtus valde prominentibus; *petiolo*  $\frac{1}{2}$ – $1\frac{1}{2}$  poll. longo. *Pedunculi* communes 1–3 poll. longi, *partiales* 2–5 lin. longi. *Umbellulæ* 5-floræ. *Flores* dioici; *perianthio* 6-rarius 4-partito, externe sericeo, interne glabro, lobis deciduis. *Stamina* sæpissime 9, 3 interiora biglandulosa. *Fructus* 4–5 lin. in diam.

This would seem to bear very considerable resemblance to *Lepidadenia Wightiana*, N. ab E., but the more numerous veined and oblique leaves of the latter, together with its hermaphrodite flowers and more numerous stamens, would appear sufficiently to distinguish it. In some of the male flowers, however, of the present plant there are, in addition to the nine fertile stamens, two or three central biglandular abortive ones, and thus is exhibited a character intermediate between that of *Tetranthera* and that of *Lepidadenia*, and justifying the suppression of the latter genus.

8. **T. iteodaphne**, N. ab E. l. c. p. 542.—C.P. 10, (351, 729, 2487.)

Var.  $\beta$ ; *foliis* minoribus, anguste lanceolatis, sæpe valde acuminatis.—C.P. 360.

Var.  $\gamma$ ; *foliis* ellipticis vel oblongis, parum acuminatis vel rarius apice rotundatis.—C.P. 2605.

HAB. Central Province, up to an elevation of 6000 feet.

*Arbor* mediocris, 20–30-pedalis. *Folia* subtus minute puberula, rarius glabra, 2–6 poll. longa,  $\frac{1}{2}$ –2 poll. lata, *petiolo* 2–4 lin. longo. *Perianthium* sæpissime 4-partitum, lobis deciduis. *Stamina* sæpissime 8, 4 interiora biglandulosa. *Fructus* oblongis, ruber.

A most variable plant, the extreme forms of which are so very dissimilar in appearance, that they might well be taken for distinct species; but they so merge one into the other that it is impossible to consider them other than varieties. Var.  $\gamma$  approaches very closely some forms of the next species.

9. **T. ovalifolia**.—*Lepidadenia ovalifolia*, Wight, Ic. t. 1839.—C.P. 167, (321, 2488, 2489, 2490.)

HAB. Central Province, at an elevation of 4000 to 7000 feet.

*Arbor* 30–40-pedalis, ramulis foliisque junioribus atro-violaceis, fugaci-puberulis. *Folia* ovata, oblonga vel ovato-lanceolata, obtusa, basi rotundata vel parum acuta, glabra, subtus pallidiora, 2–6 poll. longa, 1–3 poll. lata, *petiolo* 2–5 lin. longo. *Perianthium* 4–8-partitum, lobis deciduis. *Stamina* 8–16, biserialia, interioribus biglandulosis, sed sæpe numerosiora, valde irregularia confertaque. *Fructus* subsphæricus.

## 7. ACTINODAPHNE, N. ab E.

1. **A. glauca**, N. ab E. Syst. Laur. p. 599.—C.P. 3518, ? 2536.

HAB. lagalla, at an elevation of 7000 to 8000 feet.

*Arbor* 20–30-pedalis. *Flores* fœm. fasciculati,  $1\frac{1}{2}$  lin. longi, externe cum pedicellis 2 lin. longis ferrugineo-pilosi. *Stamina* sterilia spathulata, acuta, glabra. *Fructus* subsphæricus, basi perianthii aucta parum crenulata suffultus.

2. **A. stenophylla**, Thw.; arborea; *partibus junioribus inflorescentiaque* ferrugineo-tomentosis; *foliis* ad apicem ramuli subverticillatis, 5–7-nis, lineari-lanceolatis, obtusis, minutissime reticulatis, penniveniis, denique glabris, subtus albo-pruinosis, basi parum angustatis; *fasciculis* 4–8-floris, secus ramulum infra folia sparsis; *floribus* dioicis; *pedicellis* subæquilongis; *perianthii lobis* 6, deciduis; *fructibus* sphæricis.—C.P. 303, 2491.

HAB. Abundant in the Oovah and Badulla Districts, up to an elevation of 3000 feet.

*Arbor* 20–30-pedalis. *Folia* 2–6 poll. longa, 3–10 lin. lata, *petiolo* 2–5 lin. longo. *Flores*  $1\frac{1}{2}$  lin. longi. *Fructus* 3 lin. in diam.

3. **A. Moonii**, Thw.; arborea; *partibus junioribus inflorescentiaque* ferrugineo-tomentosis; *foliis* ad apicem ramuli subverticillatis, 4–5-nis, lanceolatis vel ovato-lanceolatis, acuminatis, obtusiusculis, penniveniis, superne minute reticulatis, subtus albo-pruinosis pilosulis denique glabris, basi rotundatis vel parum angustatis; *floribus* fasciculatis, dioicis, brevissime pedicellatis, axillaribus et infra folia secus ramulum sparsis; *fructibus* sphæricis, perianthio 6-lobo vix aucto suffultis.—C.P. 74.

HAB. Central Province, at an elevation of 5000 to 7000 feet.

*Arbor* 20-pedalis. *Ramuli* teretes, tomentosi. *Folia*  $1\frac{1}{2}$ –4 poll. longa,  $\frac{1}{2}$ – $1\frac{1}{2}$  poll. lata, *petiolo*  $1\frac{1}{2}$ –3 lin. longo. *Flores*  $1\frac{1}{2}$  lin. longi. *Fructus* 3 lin. in diam.

Nearly allied to *A. molochina*, N. ab E., but its penninerved acuminate leaves sufficiently distinguish it.

4. **A. elegans**, Thw.; arborea; *partibus junioribus inflorescentiaque* ferrugineo-tomentosis; *foliis* ad apicem ramuli 3–4-nis, lanceolatis, sæpe

longiuscule acuminatis, superne minute reticulatis, subtus albo-pruinosis, denique nudis, basi parum acutis; *floribus* dioicis, brevi-pedicellatis, fasciculatis, axillaribus et infra folia secus ramulum sparsis; *lobis perianthii* 6, deciduis; *fructibus* sphaericis.—C.P. 2277.

HAB. Central Province and in the Pasdoon Corle, up to an elevation of 3000 feet.

*Arbor* parva. *Folia*  $1\frac{1}{2}$ –4 poll. longa,  $\frac{1}{2}$ –1 poll. lata, *petiolo* tenui, 3–5 lin. longo. *Flores*  $1\frac{1}{2}$  lin. longi. *Fructus* 4 lin. in diam.

Differing from the last species, which it greatly resembles, in its narrower and more membranaceous leaves, with longer slenderer petioles, and its larger fruit, with truncated not lobed cup.

5. **A. speciosa**, N. ab E. l. c. p. 602; Wight, Ic. t. 1842.—C.P. 674.

Var.  $\beta$ ; *foliis* minus coriaceis, parcius hirsutis, ovato-vel oblongo-lanceolatis, acuminatis, basi acutis.—C.P. 3371.

HAB. Central Province, up to an elevation of 8000 feet. Var.  $\beta$ . At about 4000 feet.

*Arbor* 30–40-pedalis. *Fructus* subsphaericus, perianthii basi truncata aucta ore contracta suffultus.

6. **A. molochina**, N. ab E. l. c. p. 603; Wight, Ic. t. 1843.—C.P. 2272, v. 2373.

HAB. Pedrotalagalla, at an elevation of 6000 to 8000 feet.

*Arbor* parva. *Fructus* sphaericus, parum oblatus, 3– $3\frac{1}{2}$  lin. in diam., perianthio 6-lobo vix aucto suffultus.

## 8. DAPHNIDIUM, N. ab E.

1. **D. lancifolium**, Thw.; arboreum; *partibus junioribus inflorescentiaque* fulvo-sericeis; *foliis* ad apicem ramuli plus minus confertis, 5–7-nis, lanceolatis, gradatim acute acuminatis, superne laevibus vel minutissime rugulosis, subtus albo-pruinosis, basi acutis, triplinerviis, nervis lateralibus vix ad medium folii attingentibus; *gemmae* 3–4-floris, 4-squamosis, infra folia secus ramulum sessilibus.—C.P. 356.

HAB. Hapootelle District, in the Central Province.

*Folia* 2– $3\frac{1}{2}$  poll. longa, 6–10 lin. lata, *petiolo* 3–5 lin. longo. *Flores* subsessiles,  $1\frac{1}{2}$  lin. longi.

Apparently nearly allied to *D. melastomaceum*, N. ab E. l. c. p. 607. I have but a single specimen of the present plant, found mixed with some specimens of *Actinodaphne stenophylla*, a species it at first sight greatly resembles.

## 9. LITSÆA, Juss.

1. **L. Zeylanica**, N. ab E. Syst. Laur. p. 626, cum syn.; Wight, Ic. tt. 132, 1844. An *C. oblonga*, N. ab E. l. c. p. 672; Wight, Ic. t. 1845?—C.P. 2281, 3449 (413 partim).

Var.  $\beta$ ; *foliis* firmioribus, glaberrimis, ellipticis vel oblongis, parum acuminatis vel valde obtusis; *staminibus* plus minus pilosis. An *L. foliosa*, N. ab E. l. c. p. 622?—C.P. 2278, 2280, 72.

HAB. Very abundant up to an elevation of 4000 feet. Var.  $\beta$ . Up to 6000 feet. Nom. vulg. “Dawal-kooroondoo-gass.”

Plants of this species vary respectively very considerably as regards the size, shape, and texture of the leaves, but the character of the venation is very similar in all. The silky pubescence upon the young ramuli and leaves is in some persistent for a



considerable period, whilst in others it is exceedingly fugacious. The pruinose whiteness of the under surface of the leaves disappears sooner or later in different plants.

2. **L. orbicularis**, Thw. ; *ramulis junioribus* fusco-tomentellis ; *foliis* orbicularibus vel late obovatis, ad apicem ramuli confertis, 5-6-nis, triplinerviis, glabris, subtus albido-pruinosis ; *petiolo* costaque subtus ad basin tomentellis ; *gemmis florigeris* secus ramulum sessilibus, 4-floris ; *squamis* rotundatis, externis glabris ciliatis, internis ferrugineo-pilosis.—C.P. 2699.

HAB. Ambagamowa District, at an elevation of 4000 feet.

*Folia*  $\frac{3}{4}$ -1 $\frac{1}{2}$  poll. in diam., *petiolo* 2-3 $\frac{1}{2}$  lin. longo.

3. **L. fuscata**, Thw. ; arborea ; *ramulis, petiolis, foliis subtus inflorescentiaque* ferrugineo-tomentosis ; *foliis* versus apicem ramuli alternis, ovatis vel ellipticis, utrinque parum subabrupte angustatis, triplinerviis ; *nervis lateralibus* ad medium folii vel ultra attingentibus, externe venosis.—C.P. 413.

HAB. Central Province, at an elevation of 6000 to 8000 feet.

*Arbor* 20-pedalis. *Folia* 2-4 poll. longa,  $\frac{3}{4}$ -2 poll. lata, superne glabra, *petiolo* 5-9 lin. longo. *Gemmæ florigeræ* glomeratæ, axillares et infra folia et secus ramulum dispositæ. *Flores* 1 $\frac{1}{2}$  lin. longi, cum pedicellis subæquilongis, pallide fulvo-pilosi, *lobis perianthii* deciduis. *Fructus* subsphæricus, semipollicaris, pedicello 7-8 lin. longo.

## 10. CASSYTHA, Linn.

1. **C. filiformis**, Linn. (N. ab E. Syst. Laur. p. 642, cum syn. ; Wight, Ic. t. 1847.)—C.P. 1923.

HAB. Not uncommon in the hotter parts of the island, especially near the sea.

### *Lauraceis affines.*

#### 1. HERNANDIA, Plum., Blume, Endl.

1. **H. sonora**, Linn. (Willd. Sp. Pl. Linn. iv. p. 327, cum syn. ; Wight, Ic. t. 1855.)—C.P. 2914.

HAB. Not very uncommon near the seashore in the neighbourhood of Galle and Matura.

#### 2. GYROCARPUS, Jacq., Endl.

1. **G. Asiaticus**, Willd. (N. ab E. Syst. Laur. p. 701, cum syn.)—*G. Jacquini*, Roxb. Pl. Corom. i. t. 1.—C.P. 2202.

HAB. Hot, drier parts of the island ; not uncommon.

## CXXXIII. URTICACEÆ.

### TRIB. URTICEÆ.

#### 1. FLEURYA, Gaud.

1. **F. interrupta**, Gaud. (Weddell, Arch. du Mus. ix. p. 115 ; cum

syn. ; Wight, Ic. tt. 1975, 692.)—*Bæhmeria interrupta*, Willd. Sp. Pl. iv. p. 342, cum syn.—C.P. 2186.

HAB. Common in the warmer parts of the island. Nom. vulg. "Walahambillya."

## 2. LAPORTEA, Gaudich.

1. *L. terminalis*, Wight, Ic. t. 1972; Weddell, Arch. du Mus. ix. p. 125, cum syn.—C.P. 3336.

HAB. Central Province, at an elevation of 4000 to 6000 feet.

2. *L. crenulata*, Gaud. (Wedd. l. c. p. 113, cum syn.; Miquel, Fl. Ind. Bat. vol. i. pars altera, p. 230.)—*Urtica crenulata*, Roxb.; Wight, Ic. t. 686.—C.P. 2200.

HAB. Very common up to an elevation of 5000 feet, in damp forests. Nom. vulg. "Maoossa-gass."

## 3. GERARDINIA, Gaudich.

1. *G. Leschenaultiana*, Dcne. (Wedd. Arch. du Mus. ix. p. 165, cum syn.; Wight, Ic. t. 1976.)—C.P. 3520.

HAB. Central Province, at an elevation of 5000 to 6000 feet.

2. *G. Zeylanica*, Dcne. (Wedd. l. c. p. 166, cum syn.; Miquel, Fl. Ind. Bat. vol. i. pars altera, p. 233.)—*Urtica heterophylla*, Roxb.; Wight, Ic. t. 687.—C.P. 2193.

HAB. Not uncommon in the warmer parts of the island. Nom. vulg. "Gass-kahambillya."

## 4. PILEA, Lindley.

1. *P. Wightii*, Wedd. l. c. p. 205, cum syn.—*P. radicans*, Wight, Ic. t. 1974 (non Wedd.)—C.P. 2185.

HAB. Central Province, at an elevation of 5000 to 7000 feet.

2. *P. trinervia*, Wight, Ic. t. 1973; Wedd. l. c. p. 224, cum syn.—*P. oreophila*, Miquel, l. c. p. 235, t. 17, cum syn. *Urtica trinervia*, Roxb. Fl. Ind. iii. p. 582.—C.P. 2093.

HAB. Common in wet woods, up to an elevation of 5000 feet.

3. *P. angulata*, Blume, Mus. Lugd. Bat. ii. p. 55.—*P. stipulosa*, Miquel, l. c. p. 236; Wedd. l. c. p. 230, cum syn. *P. hygrophila*, Bl. l. c.; Wedd. l. c. p. 264, cum syn.—C.P. 2184.

HAB. Central Province, common at an elevation of 3000 to 5000 feet.

A very variable plant; when growing in very damp shady places it puts on the form described under the name of *P. stipulosa*.

## 5. PELLIONIA, Gaudich.

1. *P. Heyneana*, Wedd.; Arch. du Mus. ix. p. 287, cum syn.—C.P. 2179.

HAB. Hantani, in the Central Province, at an elevation of 4000 feet.

## 6. ELATOSTEMA, Forst.

1. *E. lineolatum*, Wight, Ic. t. 1984; Wedd. l. c. p. 312, a.—C.P. 456.

Var.  $\beta$ . *majus*.—C.P. 2183.

Var.  $\gamma$ . *falcigerum*.—C.P. 3427.

Var.  $\delta$ . *lineare*.—C.P. 3426.

HAB. Very abundant in damp forests, up to an elevation of 5000 feet.

2. **E. surculosum**, Wight, Ic. t. 2091, f. 4.; Wedd. l. c. p. 329, cum syn.—C.P. 2181, 2182.

HAB. Very common on the perpendicular surface of damp rocks, in forests of the central Province, up to an elevation of 6000 feet.

### 7. PROCRIS, Comm.

1. **P. lævigata**, Blume, Bijdr. p. 508; Miquel, Fl. Ind. Bat. vol. i. pars altera, p. 248, cum syn.—*P. Wightiana*, Wall.; Wedd. Arch. du Mus. ix. p. 336, cum syn.—C.P. 2180.

HAB. Common in forests, on damp rocks and trees, up to an elevation of 6000 feet.

*Flores masc. fasciculati. Fasciculi pluriflori, glabri. Pedicelli 1-1½ lin. longi. Perianthium 1 lin. longum; laciniis oblongis, obtusis. Pistilli rudimentum minutum. Flor. fœm. pedunculi pallidi, valde succosi.*

### 8. BOEHMERIA, Jacq.

1. **B. Malabarica**, Wedd. Arch. du Mus. ix. p. 355, cum syn.—*Urtica aquatica*, Moon's Cat. p. 62.—C.P. 2198.

HAB. Very common throughout the island. Nom. vulg. "Mahadeya-dool."

The bark of this plant is used by the Cinghalese for fishing-lines.

2. **B. platyphylla**, Don et Ham. (Wedd. l. c. p. 364, cum syn.) a.—C.P. 2952.

Var.  $\kappa$ . *macrostachya*, Wedd. l. c.—*Splizgerbera macrostachya*, Wight, Ic. t. 1977.—C.P. 2196.

Var.  $\mu$ . *Zeylanica*, Wedd. l. c.—C.P. 2197.

HAB. Common in the Central Province, up to an elevation of 6000 feet.

### 9. CHAMABAINIA, Wight.

1. **C. cuspidata**, Wight, Ic. t. 1981; Wedd. l. c. p. 387, cum syn.—C.P. 2882.

HAB. Newera Ellia, Mr. W. Ferguson.

### 10. POUZOLZIA, Gaudich.

1. **P. auriculata**, Wight, Ic. t. 1980, f. 2; Wedd. Arch. du Mus. ix. p. 393, cum syn.—*P. bicuspidata*, Wight, Ic. t. 2098, n. 32.—C.P. 2191.

HAB. Common in the hotter parts of the island.

*P. elliptica*, *P. scabra*, *P. Rheedei*, and *P. rostrata* of Wight's Icones are also quoted by M. Weddell as either forms or varieties of this species.

2. **P. Indica**, Gaud. (Wedd. l. c. p. 398, cum syn.)—*P. procumbens*, Wight, Ic. t. 2999, n. 35.—C.P. 2188.

Var.  $\beta$ . *alienata*, Wedd. l. c.—*P. Zeylanica*, Wight, Ic. t. 2101, n. 45.—C.P. 2189.

HAB. Very abundant up to an elevation of 4000 feet.

An extremely variable species; and Weddell includes amongst the synonyms of the present plant, under its several varieties or forms, *P. suffruticosa*, *P. tetraptera*, *P. Johnsoniana*, *P. pyramidata*, *P. pilosa*, and *P. minor* of Wight's Icones.

## 11. HYRTANANDRA, Miquel.

*Memorialis*, Ham. MSS., Weddell.

1. **H. Walkeriana**, Thw.—*Memorialis aquatica*, Wedd. Arch. du Mus. ix. p. 418 (nec *Urtica aquatica*, Moon, nec *U. verrucosa*, Moon). *Pouzolzia Walkeriana*, Wight, Ic. t. 2095, n. 16. *P. glabra*, Wight, l. c. n. 15.—C.P. 2192, 3372.

HAB. Common on moist banks, on the margins of woods, etc., up to an elevation of 2000 feet.

2. **H. hirta**, Miq. Fl. Ind. Bat. vol. i. pars altera, p. 261, cum syn.—*Memorialis hispida*, Ham. in Wall. Cat.; Wedd. l. c. p. 426, cum syn. Var. *Gardneri*, Wedd. l. c.—*Pouzolzia Gardneri*, Wight, Ic. t. 2092, n. 5.—C.P. 2195.

Var. *Bennettiana*, Wedd. l. c.—*P. Bennettiana*, Wight, Ic. t. 1978.—C.P. 2190.

HAB. Central Province, at an elevation of 4000 to 7000 feet.

Var. *Gardneri*, when growing upon dry banks, has frequently very small leaves, not exceeding 3 lines in length, when it closely resembles in appearance the following species, *H. triandra*; and I suspect may in some instances have been distributed by me in lieu of that species.

*P. ovalifolia*, *P. tomentosa*, *P. heterocarpa*, and *P. quadrialata*, of Wight's Icones, are quoted by Weddell as belonging to this species.

3. **H. triandra**, Miq. l. c. p. 262, cum syn.—*Memorialis parvifolia*, Wedd. l. c. p. 432. *Pouzolzia parvifolia*, Wight, Ic. t. 2092, n. 1. *Urtica triandra*, Blume, Bijdr. p. 496.—C.P. 2187.

HAB. Newera Ellia.

## 12. OREOCNIDE, Miquel.

*Villebrunia*, Gaud., Blume.

1. **O. sylvatica**, Miquel, Pl. Jungh. i. p. 40, cum syn.; Fl. Ind. Bat. vol. i. pars altera, p. 270, cum syn.—*Villebrunia sylvatica*, Blume, Mus. Bot. Lugd. Bat. ii. p. 167, t. 15; Wedd. Arch. du Mus. ix. p. 453, t. 15 c.—C.P. 2199.

HAB. Central Province, at an elevation of 2000 to 4000 feet.

## 13. MOROCARPUS, Sieb. et Zucc.

*Debregeasia*, Gaud., Weddell.

1. **M. longifolius**, Blume, l. c. p. 156; Miquel, Fl. Ind. Bat. vol. i. pars altera, p. 272, cum syn.—*Debregeasia velutina*, Gaud.; Wedd. Arch. du Mus. ix. p. 460, cum syn. *Conocephalus niveus*, Wight, Ic. t. 1959. *Urtica longifolia*, Burm. Fl. Ind. p. 197. *Urtica verrucosa*, Moon, Cat. p. 62.—C.P. 2194.

HAB. Common in the Central Province, at an elevation of 1000 to 3000 feet. Nom. vulg. "Gass-dool."

The Cinghalese make fishing-lines of the bark of this plant.

2. **M. Wallichianus**, Thw.—*Debregeasia Wallichiana*, Wedd. l. c. p. 464, t. 14. *Urtica leucophylla*, Wall. Cat. n. 4607.—C.P. 2201.  
 HAB. Hantani, in the Central Province.  
*Arbor* 15–20-pedalis.

TRIB. *ARTOCARPEÆ*.

14. **ARTOCARPUS**, Linn.

1. **A. nobilis**, Thw. (*A. pubescens*, Moon's Cat. p. 61 : non Willd.) ; monoica ; *foliis* magnis, subglabris, ovatis, brevi-acuminatis, basi obtusis, margine grosse crenatis ; *venis primariis* rectis, parallelis, apice arcuatis ; *lamina interposita* superne prominente et ad marginem folii parum saccata ; *amentis* squamis orbicularibus stipitatis peltatis puberulis persistentibus arcte tectis ; *masc.* lineari-cylindricis, *fœm.* oblongo-cylindricis.—C.P. 2818.

HAB. Not uncommon in the southern and central parts of the island, up to an elevation of 2000 feet. Nom. vulg. "Del-gass."

*Arbor* ingens, eximia, 40–50-pedalis. *Ramuli juniores* teretes, virides, albido-punctati, strigoso-scabri, denique brunnei, rugulosi. *Folia* 6–14 poll. longa, 4–10 poll. lata, superne nigro-viridia, costa venisque pallidis, subtus pallida, asperula ; *petiolo* cylindrico, minute strigoso,  $\frac{3}{4}$ –1½ poll. longo, 3 lin. in diam. *Stipulæ* nigro-virides, externe strigosulæ, 3 poll. longæ. *Pedunculi* 2–3½ poll. longi. *Amenta masc.* 3–6 poll. longa, 5–8 lin. in diam. ; *fœm.* breviora, 10–12 lin. in diam. ; *floribus* minutis, numerosissimis, papillæformibus, parum glandulosis. *Fructus* oblongus, parum nodulosus, papillosus, 6–8 poll. longus, 3½–4 poll. latus. *Semina* subsphærica, alba, semipollicaria.

*Folia* plantarum juniorum vel valde luxuriantium majora et pinnatifida.

This fine tree, which has, until now, been confounded with *Artocarpus pubescens* of Willdenow, is evidently quite distinct from that, and apparently from any other hitherto described species. The wood is of very good quality, but not considered of equal value with that of *Artocarpus integrifolia*. The seeds toasted are a favourite article of food with the Cinghalese.

2. **A. Lakoocha**, Roxb. Fl. Ind. iii. p. 524 ; Wight, Ic. t. 681 ; Trécul, Ann. des Sci. Nat. ser. iii. vol. iii. p. 117 ; Walpers, Annales, i. 660.—C.P. 2831.

Var.  $\beta$  ; *foliis* utrinque glabris vel subtus ad costam nervosque puberulis.—*A. Gomeziana*, Wall. MSS. Tréc. l. c. p. 118 ; Walp. l. c.—C.P. 2232 (2463).

HAB. Ratnapoora, Gardner. Var.  $\beta$ . Southern and central parts of the island, up to an elevation of 3000 feet. Nom. vulg. "Canna-gonagass."

*Fructus* minute velutinus.

The seeds of this, when toasted, are eaten by the Cinghalese.

*ARTOCARPUS INTEGRIFOLIA*, Linn., is a common tree in native gardens or where native gardens have been, but cannot be considered truly indigenous. The timber is very valuable for cabinet and other work ; and the fruit is much esteemed as an article of diet by the Cinghalese.

15. **CUDRANUS**, Rumphius.

*Cudranus* et *Maclura* (Sect. *Plecosperrum*), Miquel. *Cudrania*, Trécul.  
*Maclura* (Sect. *Leptosura*), Blume. *Trophis*, Willd., partim.

1. **C. Rumphii**, Thw.—*C. Bimanus* vel *Javanus*, *Javanensis*, et *Am-*

*boinensis*, Rumph. Amboin. v. p. 22, tt. 15, 16. *C. Amboinensis*, Miq. Fl. Ind. Bat. vol. i. pars altera, p. 290. *Cudrania Javanensis*, Trécul, Ann. des Sci. Nat. ser. iii. vol. viii. p. 123 (non Wight, Ic. t. 1961). *Maclura (Plecosperrum) Javanica*, Miq. l. c. p. 280. *Maclura (Leptosura) Javanica, Timorensis, et Amboinensis*, Blume, Mus. Bot. Lugd. Bat. ii. pp. 83, 84, t. 31. *Plecosperrum cuneifolium*, Thwaites, in Hook. Journ. of Bot. vi. (1854) p. 303. *Trophis spinosa*, Willd. Sp. Pl. iv. p. 734 (partim); Blume, Bijdr. p. 489 (non Roxb.); Wall. Cat. 4641 *B, D*. *Batis spinosa*, Wall. Cat. n. 4643 *B*.—C.P. 2526.

HAB. Hot drier parts of the island; not very common.

## 16. PLECOSPERMUM, Trécul.

1. **P. spinosum**, Tréc. (Ann. des Sci. Nat. ser. iii. vol. viii. p. 124; Wight, Ic. t. 1963.)—*Batis spinosa*, Roxb. Fl. Ind. iii. p. 763, quoad descr. *B. aurantiaca*, Wall. MSS. in Hort. Calc. *Trophis spinosa*, Heyne, Wall. Cat. 4641 *A*; an Willd. Sp. Pl. iv. p. 734, partim? *T. aculeata*, Roth, Sp. Nov. p. 368.—C.P. 2212.

HAB. Common up to an elevation of 2000 feet. Nom. vulg. "Katto-timbool."

## 17. ALLÆANTHUS, Thw.

(Hook. Journ. of Bot. vol. vi. (1854) p. 302.)

*Flores* dioici; *masc.* in spicis axillaribus unilateralibus pedunculatis dense aggregati. *Stamina* 4, perigonii segmentis opposita, æstivatione inflexa. *Ovarii rudimentum* parvum, conicum, subulatum. *Flores fœm.* pauci, cum pluribus abortivis squamæformibus intermixti, in capitulis globosis axillaribus congesti. *Perigonium* tubulosum, irregulariter 4-fissum. *Ovarium liberum*, sessile, 1-loculare. *Ovulum* pendulum, campylotropum. *Stylus* subterminalis, brevis. *Stigma* elongatum, simplex vel rarius secundo breviori comitatum.—*Broussonetiæ* valde affine.

1. **A. Zeylanicus**, Thw. l. c. t. 9 *B*.—C.P. 2215.

HAB. Central Province, at an elevation of 1000 to 2000 feet. Nom. vulg. "Allandoo-gass."

*Arbor* 30-40-pedalis. *Ramuli* teretes, pilosi, lactescentes. *Folia* alterna, disticha, cordato-lanceolata, acuminata, serrata, pennivenia, pilosula, subtus pallidiora, venosa, 3-4 poll. longa,  $1\frac{1}{4}$ - $1\frac{3}{4}$  poll. lata, ante anthesin simul decidua, *petiolo* piloso, 3 lin. longo. *Stipulæ* oblongæ, acuminatæ, membranacæ, sublæves, striatæ, obliquæ, deciduæ, 2 lin. longæ,  $1\frac{1}{2}$  lin. latæ. *Flor. masc.*: *Spicæ* 1- $2\frac{1}{4}$  poll. longæ, 2 lin. latæ, pilosæ, *pedunculo* 2-3 lin. longo. *Flores fœm.*: *Capitula* 4 lin. in diam., brevipedunculata, pilosa.

A very tough fibre is obtained from the inner bark of this tree, which is used by the Cinghalese for a variety of purposes.

## 18. ANTIARIS, Lesch.

*Lepurandra*, Nimmo.

1. **A. innoxia**, Blume, Rumphia, i. p. 172, t. 54.—*A. saccidora*, Dalz. Hook. Journ. of Bot. iii. (1851) p. 232; Wight, Ic. t. 1958. *A. toxicaria*, Hook. Comp. to Bot. Mag. i. p. 311, t. 17. *Lepurandra saccidora*, Nimmo, Plants of Bombay, p. 193. An *A. toxicariæ*, Lesch., *varietas*?—C.P. 2231.

HAB. Hot drier parts of the island; not uncommon. Nom. vulg. "Ritti-gass."

The liber, or inner bark, is carefully removed entire from portions of the trunk of the proper length, and used by the natives as sacks for containing rice and other articles.

## 19. STREBLUS, Lour., Blume.

*Epicarpurus*, Blume, olim.

1. **S. asper**, Lour. Fl. Cochin. ii. p. 615; Blume, Mus. Bot. Lugd. Bat. ii. p. 79, cum. syn.; Miq. Fl. Ind. Bat. i. pars altera, p. 278.—*Epicarpurus orientalis*, Bl. Bijdr. p. 488; Wight, Ic. t. 1961. *Trophis aspera*, Retz; Roxb. Fl. Ind. iii. p. 761.—C.P. 2214.

HAB. Very abundant up to an elevation of 2000 feet. Nom. vulg. "Gatta-nittool."

## 20. TAXOTROPHIS, Blume.

(Mus. Bot. Lugd. Bat. ii. p. 77.)

1. **T. Roxburghii**, Blume, l. c. p. 78, cum. syn.; Miq. Fl. Ind. Bat. vol. i. pars altera, p. 279.—*Epicarpurus spinosus*, Wight, Ic. t. 1962 (partim). *E. Timorensis*, Dcne. *Trophis spinosa*, Flor. Ind. iii. p. 762.—C.P. 1074.

HAB. Hot drier parts of the island; not uncommon.

*Cotyledones* carnosæ, valde inæquales, quarum major minorem et radiculam brevem amplectens.

2. **T. Zeylanica**, Thw.—*Epicarpurus Zeylanica*, Thw. Hook. Journ. of Bot. iv. (1852) p. 1, et vol. iii. t. 11.—C.P. 2213.

HAB. Haragam, near Kandy.

*Frutex* ramosus, parce spinosus, 8-10-pedalis. *Ramuli* puberuli. *Folia* glabra, membranacea, lanceolata vel trapezoideo-lanceolata, obtuse acuminata, mucronulata, serrata, minute glanduloso-punctulata. *Inflorescentia* racemosa; *masc.* brevis, oblonga, confertiflora; *fœm.* pauciflora, *pedicellis fructiferis* elongatis et apice incrassatis; *sepalis* parvis, demum refluxis, non auctis. *Cotyledones* subæquales, foliaceæ, plicato-convolutæ. *Radicula* longiuscula.

## 21. DORSTENIA, Plum.

1. **D. Indica**, Wight, Ic. t. 1964.—C.P. 29 (519).

HAB. Common in damp woods of the Central Province, at an elevation of 3000 to 6000 feet.

## 22. UROSTIGMA, Gasparr.

1. **U. religiosum**, Gasp. (Miquel, Fl. Ind. Bat. vol. i. pars altera, p. 333, cum syn.; Wight, Ic. t. 1967.)—*Ficus religiosa*, Linn.—C.P. 3672.

HAB. Common about native temples, being held in great veneration by the Cinghalese Buddhists. Nom. vulg. "Bo-gass."

2. **U. Arnottianum**, Miq. in Hook. Lond. Journ. of Bot. vi. p. 564; Walp. Ann. i. p. 688.—An *Ficus lucida*, Ait. Hort. Kew. v. p. 485?—C.P. 2856.

HAB. Hot drier parts of the island. Nom. vulg. "Kappootoo-bo-gass."

3. **U. infectorium**, Miq. l. c. p. 566, cum syn.; Walp. l. c.—*Ficus infectoria*, Willd.; Wight, Ic. t. 665. *U. ægeiophyllum*, Miq. l. c. *U. Tjakela*, Miq. l. c. cum syn. *U. Ceylonense*, Miq. l. c. p. 690.—c.p. 2931, 3083.

HAB. Common up to an elevation of 2000 feet. Nom. vulg. "Kirripalla-gass."

*Receptacula* brevi-pedunculata vel sessilia.

The Cinghalese sometimes use the bark of this tree, instead of the areca-nut, to chew with their betel.

4. **U. Wightianum**, Miq. l. c.; Walp. l. c.—An *U. perseæfolium*, Miq. Walp. l. c. p. 689?—c.p. 2223.

Var.  $\beta$ . *majus*; *foliis* majoribus, *costulis* utrinque 12-14; *pedunculis* *receptaculis* æquilongis vel multo brevioribus.—c.p. 2464.

HAB. Common in the Central Province, up to an elevation of 5000 feet.

*Receptacula* *matura* semipollicaria, flavida, plus minus rufescentia, punctata.

An extremely variable species, and the foliage of the larger variety very closely resembles that of the preceding species; the much larger size of the fruit, however, at once distinguishes the present plant.

5. **U. Benghalense**, Gasparr. (Miq. l. c. p. 571; Walp. l. c. p. 691, cum syn.)—*Ficus Benghalensis*, Linn. Amcen. *F. Indica*, Linn. Roxb.—c.p. 2225.

HAB. Hotter parts of the island, probably not indigenous. Nom. vulg. "Maha-nooga-gass."

6. **U. tomentosum**, Miq. l. c. p. 573.—*Ficus tomentosa*, Roxb. Fl. Ind. iii. p. 550; Walp. l. c.; Wight, Ic. t. 647.—c.p. 2226.

HAB. Trincomalie, Gardner. Bintenne.

7. **U. Mysorensis**, Miq. l. c. cum syn.; Walp. l. c. p. 691.—*Ficus Mysorensis*, Roth. *Ficus citrifolia*, Willd. Sp. Pl. iv. p. 1137.—c.p. 2222.

HAB. Not uncommon up to an elevation of 2000 feet. Nom. vulg. "Boonooga-gass."

8. **U. lacciferum**, Miq. l. c. p. 575; Walp. l. c. p. 692.—*Ficus laccifera*, Roxb. Fl. Ind. iii. p. 545; Wight, Ic. t. 656.—c.p. 2221.

HAB. Central Province, not uncommon. Nom. vulg. "Nooga-gass."

9. **U. Tjiela**, Miq. l. c. p. 580; Walp. l. c. p. 693.—*Ficus Tjiela*, Roxb. l. c. p. 549. *F. indica*, Linn. Syst. Veg. 922. *F. amplissima*, Linn.—c.p. 2537, 2220 (? 2218).

HAB. Central Province.

There is a noble specimen of this tree in the Royal Botanic Garden, whose branches spread over an area of ground 120 feet in diameter.

10. **U. retusum**, Miq. l. c. p. 581, cum syn.—*U. nitidum*, Miq. l. c. p. 582, cum syn. *U. ovoideum*, Miq. l. c. cum syn. *U. pisiferum*, Miq. l. c. p. 580, cum syn.; Walp. l. c. *Ficus retusa*, Linn. *F. nitida*, Thunb. *F. benjaminea*, Roxb. l. c. p. 550. *F. pallida*, Wall. Cat. n. 4567.—c.p. 14 (483).

HAB. Common in the Central Province, up to an elevation of 5000 feet.



11. **U. modestum**, Miq. l. c. p. 586; Walp. l. c. p. 696.—An *U. nervosi*, Miq. l. c., varietas?—c.p. 2219.

HAB. Central Province, at an elevation of 3000 to 5000 feet.

### 23. POGONOTROPHE, Miquel.

1. **P. dasyphylla**, Miq. in Hook. Lond. Journ. of Bot. vii. p. 74.—*P. Zeylanica*, Miq. l. c. p. 75; Walp. Ann. i. p. 701.—c.p. 233, 379.

HAB. Central Province, at an elevation of about 4000 feet.

### 24. FICUS, Linn., Miquel.

1. **F. asperrima**, Roxb. Fl. Ind. iii. p. 554; Wight, Ic. t. 633.—*F. politoria*, Moon's Cat. p. 74 (non Lam.)—c.p. 2229.

HAB. Common up to an elevation of 2000 feet. Nom. vulg. "See-wana-maddeya-gass."

The leaves are in general use amongst native cabinet-makers as a substitute for fine sand-paper.

2. **F. cinerascens**, Thw.; *foliis* coriaceis, ellipticis vel oblongis, obtuse brevissime acuminatis, 9- to 12-costulatis, reticulatis, superne glabris, nitidis, subtus pallidioribus, punctato-asperulis, basi subtrinerviis, rotundatis vel parum angustatis; *pedunculis* axillaribus, sæpius solitariis, petiolo brevioribus; *receptaculis* majusculis, sphericis, scabris, sæpe brevi-stipitatis, 3-bracteatis, pedunculo subæquilongis.—c.p. 2562.

HAB. Warmer parts of the island. Nom. vulg. "Wal-gona-gass."

*Arbor* ingens. *Ramuli* teretes (sicci sulcati), asperuli. *Folia* 3-6 poll. longa, 1½-3 poll. lata (sicca cinereo-viridia), *petiolo* asperulo, 6-12 lin. longo. *Receptacula* pollicaria. *Flores* monoici. *Perigonium* 5-phyllum. *Stigma* valde elongatum, bifidum.

3. **F. heterophylla**, Roxb. l. c. p. 532; Wight, Ic. t. 659; Miq. Hook. Lond. Journ. of Bot. vii. p. 231, cum syn.—*F. elongata*, Miq. l. c.; Walp. Ann. i. p. 712. *F. subpanduræformis*, Miq. l. c. p. 235; Walp. l. c. p. 714.—c.p. 2228.

HAB. Common in damp shady places. Nom. vulg. "Wal-ahattoo."

4. **F. parasitica**, Kœnig; Miq. l. c. p. 433, cum syn.; Walp. l. c. p. 718.—*F. Ampelos*, Kœn.; Roxb. Fl. Ind. iii. p. 553. *F. pervia*, Miq. l. c. *F. cuspidifera*, Miq. l. c. cum syn.; Walp. l. c. *F. uniglandulosa*, Wall. Cat. 4479. *F. sclerophylla*, Roxb. l. c. p. 546?—c.p. 2227.

HAB. Very abundant up to an elevation of 2000 feet. Nom. vulg. "Gass-nettool."

5. **F. disticha**, Blume, Bijdr. p. 458; Miq. l. c. p. 440; Walp. l. c. p. 721.—c.p. 2224, 3116.

HAB. Common in the Central Province, at an elevation of 3000 to 5000 feet.

6. **F. diversiformis**, Miq. l. c. p. 441; Walp. l. c.—*F. stipulata*, Moon's Cat. p. 74 (non Thunb.)—c.p. 2217.

HAB. Very common up to an elevation of 2000 feet.

### 25. COVELLIA, Gasparr.

1. **C. oppositifolia**, Gasparr.; Miq. in Hook. Lond. Journ. of Bot. vii. p. 461, cum syn.; Walp. Ann. i. p. 728.—*C. Dæmonum*, Miq. l. c.

cum syn.; Walp. l. c. *C. Assamica*, Miq. l. c. p. 464; Walp. l. c. *C. dasycarpa*, Miq. l. c. p. 465; Walp. l. c. p. 729. *Ficus oppositifolia*, Willd.; Wight, Ic. t. 638. *F. Dæmonum*, Vahl.—c.p. 2230.

HAB. Very abundant in the warmer parts of the island. Nom. vulg. "Kota-dimboola-gass."

2. *C. glomerata*, Miq. l. c.; Walp. l. c.—*Ficus glomerata*, Willd.; Roxb. Pl. Corom. ii. t. 123; Wight, Ic. t. 667.—c.p. 3305.

HAB. Common on the banks of rivers, up to an elevation of 2000 feet. Nom. vulg. "Atteekka-gass."

#### TRIB. CELTIDEÆ.

#### 26. ULMUS, Linn.

1. *U. integrifolia*, Roxb. Pl. Corom. i. t. 78; Wight, Ic. t. 1968.—*Holoptelæa integrifolia*, Planch. Ann. des Sci. Nat. ser. iii. vol. x. p. 266.—c.p. 2381.

HAB. Hot drier parts of the island.

#### 27. CELTIS, Tourn.

Sectio *Sponioceltis*, Planch.

1. *C. dysodoxylon*, Thw.; *partibus junioribus* ferrugineo-pilosis; *foliis* integris vel rarius obscure sinuato-dentatis, ovatis, plus minus abrupte acuminatis, firmis, trinerviis, glabris, basi parum obliquis angustatisque, costa venisque subtus puberulis; *cymis* brevibus, paucifloris, ferrugineo-pilosis; *drupis* parvis, subsphæricis, parum acuminatis.—c.p. 2563.

HAB. Central Province, up to an elevation of 5000 feet. Nom. vulg. "Gooranda-gass."

*Arbor* mediocris. *Folia* superne lævia, intense viridia, subtus pallidiora, inconspicue venosa (sicca fuscescencia),  $2\frac{1}{2}$ – $3\frac{1}{2}$  poll. longa,  $1\frac{1}{4}$ –2 poll. lata, *petiolo*  $1\frac{1}{2}$ –3 lin. longo. *Flores* 5-meri, interne ad basin dense pilosi. *Ovarium* versus apicem ferrugineo-strigosum. *Stigmata* elongata, acutata.

The freshly cut timber of this tree possesses a powerful and very disgusting odour.

Sectio *Solenostigma*, Planch.

2. *C. Wightii*, Planch. l. c. p. 307; Wight, Ic. t. 1969.—c.p. 50 (331).

HAB. Not uncommon in the hot drier parts of the island.

#### 28. SPONIA, Comm.

1. *S. orientalis*, Planch. Ann. des Sci. Nat. ser. iii. vol. x. p. 323.—*Celtis orientalis*, Linn. Fl. Zeyl. p. 176 (excl. syn. partim).

HAB. Common on the banks of rivers and on cleared ground. Nom. vulg. "Gadoomba-gass."

#### 29. GIRONNIERA, Gaudich.

*Nematostigma*, Planchon. *Helminthospermum*, Thw.

1. *G. subæqualis*, Planch. Ann. des Sci. Nat. ser. iii. vol. x. p. 339; Blume, Mus. Bot. Lugd. Bat. ii. p. 73, t. 25.

Var.  $\beta$ . *Ceylanica*, Planch.—*Helminthospermum scabridum*, Thw. in Hook. Journ. Bot. vi. (1854) p. 302, t. 9 C.—c.p. 716.

HAB. Central Province, up to an elevation of 4000 feet.

2. **G. reticulata**, Thw.; *partibus junioribus* fugaci-adpresso-pilosis; *foliis* ovatis vel oblongis, plus minus abrupte longiuscule acuminatis, minutissime prominentim reticulatis, penniveniis, superne glabris, subtus parum asperulis; *stipulis* lanceolatis, pilosis, fere ad apices in unam intra-axillarem connatis, deciduis; FLOR. MASC. in *paniculis* parvis axillaribus vel secus ramulum juniorem infra folia dispositis; FLOR. FÆM. solitariis, axillaribus, pedicellatis; *drupis* glabris, ovoideis, parum compressis, apice attenuatis.—c.p. 2565.

HAB. Badulla District, not uncommon up to an elevation of 3000 feet.

*Arbor* 30–40-pedalis. *Folia* 2–5 poll. longa,  $\frac{3}{4}$ –2 poll. lata (sicca rufescentia), *petiolo* 2–4 lin. longo. *Inflorescentia* subglabra. *Flores* parvi, 5-meri; *sepalis* rotundatis, obtusis. *Drupa* 6 lin. longa, 4 lin. lata, *stigmatibus* multo brevioribus coronata.

## CXXXIV. EUPHORBIAEÆ.

### TRIB. UNIOVULATÆ.

Sect. 1. EUPHORBIEÆ; *floribus involucratis, pedicellatis, nudis vel perianthii rudimento suffultis.*

### 1. EUPHORBIA, Linn.

1. **E. antiquorum**, Linn. (Willd. Sp. Pl. ii. p. 881; Roxb. Fl. Ind. ii. p. 468; Wight, Ic. t. 897; Rheede, Hort. Mal. ii. p. 81, t. 42.)—c.p. 2944.

HAB. Common in the hot drier parts of the island. Nom. vulg. "Dalook-gass."

2. **E. tortilis**, Rottler, (Wight, Ic. t. 898.)—An præcedentis varietas? —c.p. 2943.

HAB. Similar situations with the preceding, and perhaps only a variety of it. Nom. vulg. "Seenook-gass."

3. **E. Tirucalli**, Linn. (Willd. l. c. p. 890; Roxb. l. c. p. 470; Rheede, l. c. ii. t. 44.)—c.p. 2945.

HAB. Near the seashore. Nom. vulg. "Nawa-handee."

The milky juice of this plant is much employed as a medicine by the Malabars.

4. **E. nereifolia**, Linn. (Willd. l. c. p. 884, cum syn.; Roxb. l. c. p. 467; Rheede, l. c. p. 83, t. 43.)

HAB. Colombo, *Moon*. Nom. vulg. "Patook-gass."

I have never met with this plant truly wild in the island.

5. **E. hypericifolia**, Linn. (Willd. l. c. p. 895.)—*E. parviflora*, Linn.; Willd. l. c. p. 898; Roxb. l. c. p. 472. An *E. congenera*, Blume, Bijdr. p. 634? *Tithymalus erectus*, etc., Burm. Thes. Zeyl. p. 224, t. 105, f. 2.—c.p. 2125.

HAB. Very abundant. Nom. vulg. "Ella-dadakeeriya."

Tota planta, in meis speciminibus, plus minus hirsutula.

6. **E. rosea**, Retz, Obs. iv. p. 26; Willd. l. c. p. 895.—*E. saturoides*, Lam.—C.P. 2126, 2129.

HAB. Very common near the sea. Nom. vulg. "Moodoo-dadakeeriya."

7. **E. Burmanniana**, Gay.—*E. thymifolia*, Roxb. (Linn., Willd. partim). *Tithymalus humilis*, etc., Burm. l. c. t. 105, f. 3.—C.P. 2128.

HAB. Very abundant in cultivated ground. Nom. vulg. "Bin-dadakeeriya."

8. **E. Chamæsyce**, Linn. (Willd. l. c. p. 899.)—*E. uniflora*, Roxb. l. c. p. 473.—C.P. 3655.

HAB. Trincomalie, *Rev. S. O. Glenie*.

9. **E. bifida**, Hook. et Arn. Bot. Beech. p. 213.—*E. halophila*, Miquel.—C.P. 2131.

HAB. Not uncommon near the seashore.

*Folia integerrima, glauca. Involucris foliola glandulas vix excedentia*

10. **E. fimbriata**, Roth; annua, herbacea, prostrata, villosula; *foliis* oppositis, ovatis vel ovato-rotundatis, obtusis vel subacutis, serrulatis, oblique cordatis, brevi-petiolatis; *floribus* inter folia ad apices ramorum conferta nidulantibus; *pedunculis* brevibus, solitariis; *involucris foliolis* longiuscule laciniato-fimbriatis, laciniis pilosis; *glandulis* parvis, transverse oblongis.—C.P. 3568.

HAB. Gunner's Quoin, in the Batticaloa District.

*Folia* 2½–6 lin. longa, 2–3½ lin. lata. *Involucrum* brevi-campanulatum, foliolis radiantibus, 2 lin. in exp., albidis, rufescentibus. *Capsula* villosula.

11. **E. pilulifera**, Linn. (Willd. l. c. p. 897.)—*E. hirta*, Roxb. l. c. p. 472. (Linn., Willd., quoad plantas Zeylanicas.) *Tithymalus botryoides*, etc., Burm. l. c. pp. 223, 224, t. 104 et t. 105, f. 1.—C.P. 3337.

HAB. Very abundant in cultivated ground. Nom. vulg. "Boo-dadakeeriya."

12. **E. Rothiana**, Spr. (Wight, Ic. t. 1864.)—*E. læta*, Roth (non Ait.). *E. glauca*, Roxb. l. c. p. 473 (non Willd.). *E. oreophila*, Miq. *E. lanceolaria*, Herb. Heyne.—C.P. 2124.

HAB. Very common up to an elevation of 7000 feet.

Sect. 2. HIPPOMANEÆ; *floribus* spicatis vel subspicatis, arcte bracteatibus; *perianthio* uniseriali, segmentis in æstivatione imbricatis, vel plus minus imperfecto.

## 2. SAPIUM, Jacq.

1. **S. Indicum**, Willd. (Roxb. Fl. Ind. iii. p. 692; Wight, Ic. t. 1950.)—C.P. 2946.

HAB. Warmer parts of the island. Nom. vulg. "Kirri-makooloo-gass."

## 3. EXCŒCARIA, Linn.

1. **E. Agallocha**, Linn. (Willd. Sp. Pl. iv. 864; Roxb. Fl. Ind. iii. p. 756; Wight, Ic. t. 1865 B.)—C.P. 2169.

HAB. Very common near the sea. Nom. vulg. "Tellakeeriya-gass."

2. **E. oppositifolia**, Jack in Calc. Journ. of Nat. Hist. iv. p. 386; Walp. Ann. i. p. 621.—*E. crenulata*, Wight, Ic. t. 1865 A.—C.P. 2523.

HAB. Common in the Central Province, at an elevation of 4000 to 6000 feet.

#### 4. MICROSTACHYS, A. Juss.

1. **M. Chamælea**, A. Juss.—*Tragia Chamælea*, Linn.; Willd. Sp. Pl. iv. p. 326, cum syn.; Roxb. Fl. Ind. ii. p. 577.—c.p. 2132.

HAB. Very common amongst grass. Nom. vulg. "Rat-pittawaakka."

#### 5. FALCONERIA, Royle.

1. **F. insignis**, Royle, Him. Illust. p. 354, t. 98, f. 2.—Var. *ovario triloculari*.—*F. Malabarica*, Wight, Ic. t. 1866.—c.p. 654.

HAB. Hot, drier parts of the island.

Sect. 3. ACALYPHEÆ; *floribus masc. perianthio uniseriali, segmentis in æstivatione valvatis; flor. fœm. perianthio uniseriali vel sub-biseriali, segmentis in æstivatione sæpissime imbricatis.*

\* *Monoica*.

#### 6. DALECHAMPIA, Plum.

1. **D. bidentata**, Blume, Bijdr. p. 632; Miq. Fl. Ind. Bat. vol. i. pars altera, p. 417.—*D. Indica*, Wight, Ic. t. 1882. *D. velutina*, Wight, Ic. t. 1881.—c.p. 2170.

HAB. Gonagama, on the Lower Badulla road from Kandy; not common.

My specimens appear to be intermediate in character between Dr. Wight's *D. Indica* and *D. velutina*, and agree very well with Blume's description. The analysis of the male flower in both Dr. Wight's figures is incorrect, the filaments being represented as free to the base, instead of cohering into a central column.

#### 7. TRAGIA, Plum.

(Nom. vulg. "Wel-kahambilleya.")

1. **T. cannabina**, Willd. Sp. Pl. iv. p. 326; Roxb. Fl. Ind. iii. p. 575.—c.p. 3569.

HAB. Hot, drier parts of the island.

2. **T. involucrata**, Willd. l. c. p. 324; Roxb. l. c. p. 576, cum syn.—c.p. 2099, 2100, 2951.

Var.  $\beta$ . *montana*; *foliis ovato-lanceolatis, cordatis*.—c.p. 3519.

HAB. Very common. Var.  $\beta$ . Newera Ellia.

An extremely variable species, and I suspect that the preceding is but a form of it.

#### 8. ADENOCHLÆNA, Boivin:

In Baillon's Étude des Euph. p. 472. *Symphyllia*, Baill. l. c. p. 473.  
*Centrostylis*, Baill. l. c. p. 469.

1. **A. Zeylanica**, Thw.—*Centrostylis Zeylanica*, Baill. l. c. p. 470.—c.p. 396, 2111.

HAB. Rambodde and Happootelle, in the Central Province, at an elevation of 4000 to 5000 feet.

*Frutex suberectus, parce ramosus, partibus junioribus pubescentia stellata albida*

tectis. *Rami* elongati, debiles. *Folia* oblongo-lanceolata, acuminata, basi lata, truncata vel subcordata, trinervia, 3-7 poll. longa, 1-3 poll. lata, *petiolo*  $\frac{1}{2}$ -3 poll. longo. *Stipulæ* subreniformes. FLORES MASC. parvi, ad apicem pedunculi dense agglomerati. *Stamina* 4-5, sepalis alterna, *filamentis* ad basin pistilli rudiment. columnaris insertis, æstivatione incurvis, denique suberectis, *antheris* extrorsis. FLOR. FÆM. ad basin pedunculi, 1-3, singuli sæpe pedicello proprio, brevi, floribus masc. paucis cincto. *Sepala* 5-6, lanceolata. *Ovarium* pilosum. *Stylus* 3-lobus, majusculus, deciduus, lobis copiose profundeque laciniatis. *Calycis fructiferi* aucti segmenta pectinato-laciniata; *laciniis* longis, setaceis, apice incrassatis purpureisque. *Capsula* 3-loba, parum depressa, coccis rotundatis. *Semina* subsphærica, 2 lin. in diam., brunneo nigroque marmorata.

As M. Baillon's description of this plant, the type of his genus *Centrostylis*, is incorrect, having been derived probably from imperfect specimens, I have thought it best to describe it here pretty fully.

### 9. ACALYPHA, Linn.

1. **A. Indica**, Willd. Sp. Pl. iv. p. 523, cum syn.; Roxb. Fl. Ind. iii. p. 675; Wight, Ic. t. 877.—c.p. 2098.

HAB. Very common in cultivated ground. Nom. vulg. "Kooppa-maineya."

2. **A. betulina**, Willd. l. c. p. 522.—An *A. capitata* et *A. alnifolia*, Willd. l. c. p. 525?—c.p. 2094.

HAB. Hot, drier parts of the island.

3. **A. hispida**, Willd. l. c. p. 523.—c.p. 2092, 2097.

HAB. Common in the hotter parts of the island.

4. **A. ciliata**, Forsk. (Willd. l. c. p. 522; Roxb. l. c. p. 676.)—*A. lanceolata*, Willd. l. c. p. 524, cum syn.—c.p. 2095.

HAB. Not very uncommon in the warmer parts of the island.

5. **A. Wallichii**, Thw.; herbacea, ramosa, pilosula; *foliis* ovato-lanceolatis, dentatis, basi rotundatis, longe petiolatis; *inflorescentia* masc. axillari, spiciformi; *fæm.* terminali, paniculata, rarius axillari, spiciformi; FLOR. FÆM. *sepalis* 5.—*A. racemosa*, Wall. Cat. n. 7784. *Welia Capameni*, Rheede, Hort. Mal. x. p. 165, t. 83.—c.p. 2096 (2459).

HAB. Damp, shady places, up to an elevation of 3000 feet.

*Folia* 1-6 poll. longa,  $\frac{3}{4}$ -4 poll. lata, *petiolo*  $\frac{3}{4}$ -5 poll. longo. *Capsula* glandulosa. *Semina* subsphærica, minutissime reticulata.

\*\* *Dioicæ.*

### 10. CLAOXYLON, A. Juss. (non Wight).

*Erythrochilus*, Reinw., Blume. *Micrococca*, Benth.

1. **C. Mercurialis**, Thw.—*Micrococca Mercurialis*, Benth. in Hook. Fl. Nigrit. p. 503. *Tragia Mercurialis*, Willd. Sp. Pl. iv. p. 324; Roxb. Fl. Ind. iii. p. 576.—c.p. 3310.

HAB. Very common in cultivated ground.

2. **C. longifolium**, Baillon, Étude des Euphorb. p. 493.—*Erythrochilus longifolius*, Blume, Bijdr. p. 616.—c.p. 2102, 2499.

HAB. Central Province, up to an elevation of 5000 feet.

*Frutex* ramosus. *Folia* lanceolata, utrinque angustata, parce glanduloso-denticulata, 3-8 poll. longa,  $\frac{1}{4}$ -1 $\frac{3}{4}$  poll. lata, sæpe cyanescentia, *petiolo* apice superne glanduloso-

denticulata,  $\frac{1}{2}$ –3 poll. longo. *Pedunculi* axillares, solitarii. *Glomeruli* secus rachin sæpissime valde elongatam sparsi, hispidi; MASC. stipitato-glandulosi. FLOR. MASC. *Stamina* 12–15, glandulis stipitatis 5–6-radiatis intermixta; *filamentis* brevibus, *antheris* albidis. *Capsula* pilosa.

The present genus is at once distinguished from *Rottlera* by its stamens being intermixed with stalked glands; and by the linear squamulæ adpressed to the ovary, within and alternating with the calyx-segments.

### 11. CLEIDION, Blume.

1. **C. Javanicum**, Blume, Bijdr. p. 613?; Wall. Cat. n. 7718.—C.P. 656 (2104).

HAB. Central Province, up to an elevation of 2000 feet. Nom. vulg. “Okooroo-gass.”

*Antheræ* transverse dehiscentes. *Semina* spherica, 4–5 lin. in diam.; *testa* ossea.

### 12. SPATHIOSTEMON, Blume.

*Adelia*, Schreb., Miq., Roxb. (partim), Wight (non Linn.).

*Hæmatospermum*, Wall., Lindley.

1. **S. Javense**, Bl. Bijdr. p. 622.—Var.  $\beta$ ; *capsula* lævi vel sublævi.—*S. salicinum*, Zoll. et Moritz; Baill. Ét. des Euph. p. 293. *Adelia neriifolia*, Roxb. Fl. Ind. iii. p. 840; Wight, Ic. t. 1868; Miquel, Fl. Ind. Bat. vol. i. pars altera, p. 389. *Hæmatospermum neriifolium* et *H. riparium*, Wall. Cat. n. 7953, etc.—C.P. 651.

HAB. Very common on the banks of rivers, where it is frequently submerged; up to an elevation of 2000 feet.

I can find no sufficient character to separate, except as a mere variety, the present common Indian and Ceylon plant from Blume's species, for a specimen of which latter I am indebted to M. Binnendijk, of the Java Botanic Garden.

### 13. TREWIA, Linn.

1. **T. nudiflora**, Linn. (Roxb. Fl. Ind. iii. p. 837, cum syn.; Wight, Ic. tt. 1870, 1871.)—*Rottlera Indica*, Willd.; Blume, Bijdr. p. 607; Wall. Cat. n. 7817 B.—C.P. 2930.

HAB. Not uncommon in the hotter parts of the island.

### 14. ROTTLERA, Roxb.

1. **R. tetracocca**, Roxb. Fl. Ind. iii. p. 826.—C.P. 2118.

HAB. Common up to an elevation of 2000 feet. Nom. vulg. “Bookanda-gass.”

2. **R. rhombifolia**, Thw.—*R. dicocca*, Roxb. l. c. p. 829 (excl. syn.). *Croton rhombifolium*, Willd. Sp. Pl. iv. p. 555; Moon's Cat. p. 65.—C.P. 2115.

HAB. Not uncommon up to an elevation of 1500 feet.

*Frutex* vel *arbor* parva, subscandens, *partibus junioribus inflorescentiaque* pubescentia stellata ochraceis. *Folia*  $1\frac{1}{2}$ –4 poll. longa,  $1\frac{1}{4}$ –3 poll. lata, creberrime translucen-punctulata, subtus plus minus stellato-pilosa, disculis albidis nitentibus conspersa; *petiolo* cylindrico, piloso, 2 poll. longo. *Flores* numerosi. *Capsulæ* cocci 2, rotundati. *Semina* spherica, rufo-nigra, 2 lin. in diam.

3. **R. tinctoria**, Roxb. Cor. Pl. ii. t. 168; Willd. l. c. p. 832; Roxb. Fl. Ind. iii. p. 827, cum. syn.—*Croton coccineum*, Vahl; Willd. l. c. p. 544. An *C. punctatum*, Retz; Willd. l. c. p. 546?—c.p. 2103.

HAB. Not uncommon in the hotter parts of the island. Nom. vulg. "Hamparandella-gass."

4. **R. oppositifolia**, Blume, Bijdr. p. 608.—*Plagianthera oppositifolia*, R. et Zoll.; Baill. Étude des Euph. p. 424.—c.p. 196 (2106).

HAB. Common up to an elevation of 2000 feet.

*Frutex vel arbor parva, partibus junioribus pubescentia stellata ochraceo-fuscis. Folia ovato-lanceolata, utrinque angustata, acuminata, trinervia, 2-10 poll. longa, subtus disculis aureis nitentibus parce conspersa, venis venulisque prominentibus, petiolo ½-3 poll. longo. Capsula 3- vel 4-cocca, lævis, stellato-pubescentis, coccis subsphæricis.*

5. **R. fuscescens**, Thw.; glabra, *partibus junioribus, inflorescentia foliisque subtus disculis adpressis plus minus rufescentibus creberrime conspersis; foliis oppositis, ovato-lanceolatis, acuminatis, penniveniis vel aliquando sub-trinerviis, subtus venulis parum prominentibus; capsula 3- vel 4-cocca, sparse glandulosa, coccis subsphæricis.*—c.p. 2105.

HAB. Not uncommon up to an elevation of 2000 feet.

*Arbor parva. Folia 2-6 poll. longa, 1-3 poll. lata, basi parum angustata; petiolo ½-1½ poll. longo, apice obscure biglanduloso.*

6. **R. muricata**, Thw.—*Claoxylon muricatum*, Wight, Ic. t. 1886. An *Rottleræ aureo-punctatæ*, Dalz. in Hook. Lond. Journ. of Bot. iii. (1851) p. 122, varietas?—c.p. 2108.

HAB. Central Province, up to an elevation of 4000 feet.

*Arbor parva, subglabra. Folia apposita, cuneato-obovata, rostrato-acuminata, grosse sinuato-dentata vel rarius subintegra, subtus disculis aureis vel rufescentibus creberrime conspersa, 2-7 poll. longa, 1-3 poll. lata, petiolo ¼-1 poll. longo. Capsula 3-cocca, echinulis subglabris teretibus flexilibus prope basin articulatis caducisque munita.*

7. **R. digyna**, Thw.—*Claoxylon digynum*, Wight, Ic. t. 1884.—c.p. 2109.

HAB. Cultura.

*Arbor parva. Folia subtus disculis rufescentibus creberrime conspersa, prope basin biglandulosa.*

8. **R. eriocarpa**, Thw.; pubescens; *foliis oppositis, oblongo-lanceolatis, integris vel parum undulatis, trinerviis, subtus disculis aureis nitentibus parce conspersis, basi rotundatis vel subcordatis; capsula 3-cocca, echinulis plurimis patentim pilosis vestita, coccis subsphæricis. Croton montanum*, Herb. Wight; Wall. Cat. n. 7820 A.—c.p. 2112.

HAB. Hot, drier parts of the island, not very common.

*Frutex, pubescentia stellata ochraceo-fulva plus minus tomentosus. Folia 2-8 poll. longa, 1-4 poll. lata, petiolo ½-2 poll. longo. FLOR. MASC. Alabastra conica. Stamina numerosa, connectivo non producto. FLOR. FÆM. Ovarium staminibus sterilibus plurimis cinctum. Styli lobi 3, mediocres, interne valde plumosi.*

Wallich's plant, from Penang, has the staminal filaments hairy, whilst in the Ceylon plant they are glabrous; but in other respects the two plants appear to agree perfectly.

## 15. PODADENIA, Thw. (gen. nov.).

FLOR. MASC. *Stamina numerosa, glandulis parvis oblongis plurimis intermixta. FLOR. FÆM. Fructus indehiscens, carnosus, 3-spermus vel*



sæpius abortu 1-vel 2-spermus. *Semina* arillo copioso integro carnosio vestita.—Cæteroquin ut in *Rottlera*.

1. *P. sapida*, Thw.—*Rottlera (Stylanthus) Thwaitesii*, Baillon, Étude des Euphorb. p. 426.—c.p. 3428.

HAB. Ambagamowa and at Marai-Calai, near Ratnapoora, at an elevation of 1000 to 2000 feet.

*Arbor* ingens, *partibus junioribus* fulvo-tomentosis. *Folia* integra, obovata, breve et abrupte acuminata, pennivenia, eglandulosa, superne glabra, costa venisque pilosis, subtus pilosa, basi plus minus angustata, 5–10 poll. longa, 2½–5 poll. lata, *petiolo* cylindrico, tomentoso, ¾–2 poll. longo. *Inflorescentia* terminalis, paniculata, fulvo-tomentosa, glandulis stipitatis rubris creberrime conspersa. *Flores* rubri; MASC. secus racheos ramos dense fasciculati; alabastris acuminatis. FLOR. FÆM. *Styli lobi* 3, longiusculi, subulati, contorti, interne stigmatosi. *Fructus* subsphæricus, rufescens, glandulis majusculis stipitatis dense echinulatus, sesquipollicaris. *Semina* oblonga, 8–9 lin. longa, 4–5 lin. lata; *testa* ossea, brunnea. *Albumen* copiosum. *Arillus* albidus, sapidus.

The fleshy aril of the seed of this fine tree has an agreeable flavour, but I cannot learn that the Cinghalese eat it; the plant, however, being scarce and occurring in little-frequented localities, is probably known to but few of the natives.

## 16. MACARANGA, Dup. Th.

1. *M. tomentosa*, Wight, Ic. t. 1949–1.—c.p. 2171.

HAB. Very abundant up to an elevation of 3000 feet. Nom. vulg. “Kanda-gass.”

Sect. 4. CROTONEÆ; *floribus masc. et fœm. perianthio biseriali vel sub-biseriali, segmentis imbricatis.*

\* *Floribus apetalis.*

† *Perianthii lobis 5, quorum 2 minores, interni.*

## 17. GELONIUM, Roxb.

1. *G. lanceolatum*, Willd. Sp. Pl. iv. p. 832; Roxb. Fl. Ind. iii. p. 831; Wight, Ic. t. 1867.—*G. bifarium*, Willd. l. c. p. 831. An Roxb. l. c. p. 830? An *G. fasciculatum*, Roxb. l. c. p. 832? An *Erythrocarpus glomerulatus*, Blume, Bijdr. p. 604?—c.p. 2101.

Var. β; *foliis* minoribus et plus minus angustioribus, sæpe aculeato-dentatis; *floribus* minoribus, denique ad apicem ramorum brevium racheos bifarii dispositis.—An *Erythrocarpus spicatus*, Blume, Bijdr. p. 604?—c.p. 695, 252.

HAB. Common in rather open forests, up to an elevation of 4000 feet.

A very variable plant; and all the specimens I have seen of the genus, from different and distant parts of India, might with safety be referred, I think, to one species.

†† *Perianthii verticillis dimeris.*

## 18. CHÆTOCARPUS, Thw.

Hook. Lond. Journ. of Bot. vi. (1854) p. 300. Baill. Étude des Euphorb. p. 323.

(Nom. vulg. “Haddocka-gass.”)

*Stamina* sæpissime 8, biserialia, dimidio inferiore coalita et pistilli rudimentum cingentia.

1. **C. castanocarpus**, Thw.—*C. pungens*, Thw. l. c. p. 301 (excl. descrip. fruct.), t. 10 *A*, fig. 1–4. *Adelia castanocarpa*, Roxb. Fl. Ind. iii. p. 848; Wall. Cat. n. 7984. *Casearia ? coriacea*, Wall. Cat. n. 7196.—C.P. 2641.

Var.  $\beta$ ; *pubescens*; *ramulis foliisque subtus plus minus hispidis*.—C.P. 3013.

HAB. Common in the Ratnapoora and Ambagamowa Districts.—Var.  $\beta$ . Pasdun Corle.

*Flores* 2–2½ lin. longi. *Capsula* subsphærica, semipollicaris, aculeis rigidis fragilibus reflexe setulosis ochraceo-rufescentibus dense tecta; *putamine* mediocri.

2. **C. coriaceus**, Thw.; *foliis valde coriaceis, glabris, oblongis, brevium acuminatis, superne lævibus, subtus parce venosis, basi acutis; capsula muricata, sparse strigosa*.—*C. pungens*, Thw. l. c., quoad descrip. fruct. et fig. 6–9.—C.P. 1025.

HAB. With the preceding species, and equally common.

*Arbor* mediocris, *ramulis* subglabris. *Folia* 3–6 poll. longa, 1½–3 poll. lata, *petiolo* 3–5 lin. longo. *Fasciculi* multiflori, minute tomentosi. *Flores* 1–1½ lin. longi. *Filamenta* pilosa. *Capsula* ovoidea, obscure trigona, apiculata, 8 lin. longa; *putamine* osseo, 1 lin. in crass.

This very much resembles the preceding species; but the leaves of the present plant are much more coriaceous in texture, the flowers are smaller and on rather longer pedicels, and the capsule is very different. The timber of both is hard, and useful for building purposes.

††† *Perianthii verticillis trimeris.*

## 19. MISCHODON, Thw.

Hook. Journ. of Bot. vi. (1854) p. 299. Baillon, Étude des Euphorb. p. 335.

1. **M. Zeylanicus**, Thw. l. c. p. 300, t. 10 *B*.—C.P. 557.

HAB. Hantani, *Gardner*; Ooma Oya, on the Lower Badulla Road from Kandy; not common.

FLORE. FÆM. *Stylus* 3-partitus, *lobis* brevibus, reflexis, apice rotundato-dilatatis.

\*\* *Perianthii verticillis 4–5-meris, interiore petalino, aliquando in flor. fæm. minutissimo, vel nullo.*

† *Monoica.*

## 20. CROTON, Linn.

1. **C. lacciferum**, Linn.; Wight, Ic. t. 1915 (excl. syn. aliq.)—*Aleurites lacciferus*, Willd. Sp. Pl. iv. p. 590. *Croton foliis ovatis*, etc., Fl. Zeyl. n. 344. *Ricinoides aromatica arbor*, etc., Burm. Thes. Zeyl. p. 201, t. 91.—C.P. 2117.

HAB. Very common up to an elevation of 3000 feet. Nom. vulg. “Gass-kappiteya.”

The lac obtained from the old trees of this species is employed by the Cinghalese for medicinal and other purposes.

2. **C. aromaticum**, Linn. (Willd. l. c. p. 549, cum syn.)—An *C. Molluccanum*, Willd. l. c. p. 551 (excl. syn.)? *Croton foliis cordatis*, etc., Fl. Zeyl. n. 345.—C.P. 2116.

HAB. Abundant in the hotter parts of the island. Nom. vulg. “Wel-kappiteya.”

A præcedente, cui valde affinis, differt *ramulis* elongatis debilibus; *foliis* minus pilosis et sæpius aperte cordatis; *capsula* majore elongatioreque, pilis stellatis parce conspersa.

The present plant bears considerable resemblance to Roxburgh's *Croton drupaceum*; but, besides other differences, the fruit of the latter is much larger and more hairy, and the lateral nerves of its leaves form a much more acute angle with the costa.

3. **C. Klotzschianum**, Wight, Ic. t. 1914.—C.P. 2113.

HAB. Not uncommon in the hot, drier parts of the island.

*Frutex* vel *arbor* parva, *partibus junioribus* dense stellato-pubescentibus. *Folia* oblongo-lanceolata, acuminata, denique subglabra, incurvo-denticulata, creberrime translucenti-punctulata,  $1\frac{1}{2}$ –4 poll. longa,  $\frac{3}{4}$ –2 poll. lata, *petiolo* 3–15 lin. longo, glandulis disciformibus ad apicem subsessilibus. *Capsula* parum depressa, triloba, 4 lin. in diam., denique glabra.

I think this must be Dr. Wight's plant, as it agrees very well with his figure and short description. It is very closely allied to the two preceding species, from which it is best distinguished by its smaller, and at length glabrous, leaves and capsule; and by the latter being distinctly trilobed instead of subspherical.

4. **C. Moonii**, Thw.; *foliis* scabris, oblongis vel obovato-oblongis, acuminatis, obscure denticulatis, utrinque sparse stellato-pubescentibus, basi plus minus angustatis, rotundatis vel subcordatis, irregulariter trinerviis; *capsula* oblato-sphæroidea, tuberculata, furfuracea.—*C. punctatum*, Moon, Cat. p. 65 (non Retz).—C.P. 577.

HAB. Cultura.

*Arbor* parva, *ramulis*, *petiolis inflorescentiaque* pubescentia stellata scabris. *Folia* 3–6 poll. longa,  $1\frac{1}{2}$ – $2\frac{1}{2}$  poll. lata, siccitate superne nigrescentia, *petiolo*  $\frac{1}{2}$ – $1\frac{1}{4}$  poll. longo, glandulis disciformibus ad apicem subsessilibus. *Capsula*  $3\frac{1}{2}$  lin. in diam.

5. **C. nigro-viride**, Thw.; subglabrum, *partibus junioribus* sparse stellato-pubescentibus; *foliis* firmis, opacis, ovato-lanceolatis, obtuse acuminatis, serratis, irregulariter 3-nerviis, basi parum angustatis; *capsula* (immatura) stellato-pubescente furfuracea.—C.P. 3637.

HAB. Rambodde, in the Central Province, at an elevation of 5000 feet.

*Arbor* parva. *Folia* nigro-viridia, 3–5 poll. longa; *petiolo* semipollicari, glandulis disciformibus ad apicem brevi-stipitatis.

Allied to the preceding, but distinguished at once by its glabrous and distinctly serrated leaves.

6. **C. hypoleucum**, Dalz. in Hook. Journ. of Bot. iii. (1851) p. 123 (non Schlecht.).—*C. reticulatum*, Heyne, fid. Wall. Cat. 7724 A, B.—C.P. 2110.

HAB. Central Province, at an elevation of 2000 to 3000 feet.

*Frutex*, *ramulis*, *petiolis inflorescentiaque* albido-vel sæpius plus minus ferrugineo-leprosis. *Folia* membranacea, oblongo-lanceolata, acuminata, subintegra, superne glabra nitida (juniora fugaci-leprosa), subtus dense albido-leprosa, basi rotundata vel plus minus acuta, trinervia, 3–7 poll. longa,  $\frac{3}{4}$ – $2\frac{1}{2}$  poll. lata; *petiolo*  $\frac{1}{2}$ – $1\frac{1}{4}$  poll. longo, glandulis disciformibus ad apicem 2–4, stipitatis. *Capsula* ovoidea, trigona, apiculata, ferrugineo-leprosa, semipollicaris. *Semina* oblonga, nigro-fusca, albo-maculata.

7. **C. oblongifolium**, Roxb. Fl. Ind. p. 685.—*C. lævigatum*, Wall. Cat. 7735.—C.P. 2114.

HAB. Hot drier parts of the island, at no great elevation.

*Arbor* parva, *ramulis*, *petiolis inflorescentiaque* albido-leprosis. *Folia* glabra, oblongo-lanceolata, utrinque angustata, parum obtusata, serrata, glandulis disciformibus ad apicem petioli parvis immersis.

*Croton Tiglium*, Linn., occurs not uncommonly in native gardens, but can scarcely be considered truly wild.

## 21. BLACHIA, Baillon :

Étude des Euphorb. p. 385, t. 19, fig. 18-20.

1. **B. umbellata**, Baill. l. c.—*Croton umbellatum*, Willd. Sp. Pl. iv. p. 545 ; Wight, Ic. t. 1874.—C.P. 3014.

HAB. Very common in the south of the island, and especially near the sea.

Though it is difficult to point out any very decided character to separate this species generically from *Croton*, still the eglandulose leaves, the umbellate inflorescence, the anthers erect in æstivation, and other minor characters about the inflorescence of the present plant, are probably sufficient to justify its being constituted a distinct genus.

## 22. TRIGONOSTEMON, Blume.

1. **T. diplopetalus**, Thw. ; *foliis* subglabris, lanceolatis, mucronatis, distanter denticulatis, basi cuneatis ; *pedicellis* floribus subæquilongis ; *petalis* profunde bifidis, lobis rotundatis, laciniato-sinuatis, undulatis ; *capsula* lævi, pilosa.—C.P. 578.

HAB. Palahattoo, in the Reigam Corle, at no great elevation.—Sept. 1856.

*Frutex* vel *arbor* parva, *partibus junioribus inflorescentiaque* parce pilosis. *Folia* sublævia, pennivenia, siccitate nigro-fuscescentia, 6-9 poll. longa,  $1\frac{1}{2}$ - $2\frac{3}{4}$  poll. lata ; *petiolo* semipollicari, superne sulcato, ad apicem bidenticulato. *Stipulæ* minutæ, subulatae, caducæ. *Racemi* terminales, multiflori, foliis breviores ; *bracteis* minutis. *Petala* rubra. FLOR. MASC. *Pedicelli*  $1\frac{1}{2}$ -2 lin. longi. *Calycis lobi* rotundati, petalis dimidio breviores. *Antherarum loculi* ad apicem connectivi crassi divergentes. FL. FÆM. *Pedicelli* superne incrassati, 4-5 lin. longi ; *fructiferi* 6-7 lin. longi. *Calycis lobi* oblongi, plerumque dorso apiculati, petalis reflexis longiores. *Styli lobi* 3, breves, bis bifidi. *Capsula* 3-lobata, parum depressa, 2 poll. lata.

2. **T. nemoralis**, Thw. ; *foliis* subglabris, lanceolatis, acuminatis, mucronulatis, undulatis, obscure denticulatis, penniveniis, basi cuneatis, brevi-petiolatis ; *floribus* pedicellis subæquilongis ; *petalis* oblongis, quam lobi calycini 2- vel 3-plo longioribus ; *capsula* muricata, pilosa.—C.P. 3570.

HAB. Madamahanewera, in the Central Province, at an elevation of 2000 feet.

*Frutex* vel *arbor* parva, *partibus junioribus inflorescentiaque* adpresse pilosis. *Folia* sublævia, ad apicem ramuli parum conferta, 3-7 poll. longa, 1-2 poll. lata ; *petiolo* 2-4 lin. longo, apice bidenticulato. *Stipulæ* subulatae, 1- $1\frac{1}{2}$  lin. longæ. *Racemi* multiflori, foliis breviores. *Petala* rubra. FLOR. MASC. *Antherarum loculi* subparalleli, parum distantes, ad apicem connectivi rotundati. FL. FÆM. quam masc. parum majores, 6-7 lin. in exp. *Ovarium* hispidum. *Styli lobi* 3, breves, spathulati, emarginati, reflexi. *Semina* sphaerica, 3 lin. in diam., dorso parum carinata ; *testa* lævi, fusco-marmorata.

## 23. JATROPHA, Kth.

1. **J. glauca**, Vahl, (Willd. Sp. Pl. iv. p. 558.)—C.P. 2168.

HAB. Damboul, Gardner,

24. **DIMORPHOCALYX**, Thw. (gen. nov.)

**FLOR. MASC.** *Calyx* cupuliformis, margine undulatis vel breve 5-dentatus. *Petala* 5, erecta, apice reflexa. *Glandulæ* 5, squamæformes, petalis alternæ. *Stamina* sæpissime 10, biserialia, 5 externa breviora petalis opposita; *filamentis* coalitis, apice liberis; *antheris* introrsis, loculis parallelis, longitudinaliter dehiscentibus. *Ovarii rudimentum* nullum. **FLOR. FÆM.** *Calyx* profunde 5-lobus, segmentis plus minus inæqualibus. *Petala* ut in flor. masc. *Stamina abortiva* sæpius nulla. *Discus* annularis. *Ovarium* 3-loculare. *Styli lobi* 3, erecti, apicibus bifidis, reflexis. *Capsula* 3-cocca, calyce plus minus aucto suffulta. *Arbor* parva, ramosissima, glabra; *foliis* lanceolatis vel ovatis, margine parce glanduloso-punctatis, petiolatis; *stipulis* parvis, triangularibus, acutis; *cymis* paucifloris, terminalibus axillaribusque, sessilibus vel brevi-pedunculatis, sæpius unisexuis; *bracteis* parvis, imbricatis.

1. **D. glabellus**, Thw.—*Croton glabellum*, Herb. Wight. An *C. Lawianus*, Nimmo in App. to Graham's Bot. of Bombay?—c.p. 1046 (2167).

**HAB.** Common in the hot drier parts of the island. Nom. vulg. "Welle-wenne-gass."

*Folia* 1-6 poll. longa, *petiola* 2-6 lin. longo. *Pedicelli* 1-4 lin. longi. *Petala* alba, 3-4 lin. longa. *Capsula* 4-5 lin. in diam., pilosula.

†† *Dioicæ*.

25. **DESMOSTEMON**, Thw. (gen. nov.)

*Calyx* profunde 5-lobus, segmentis rotundatis, inæqualibus. *Petala* 5, erecta, apice reflexa, calyce longiora. **FLOR. MASC.** *Glandulæ* 5, majusculæ, carnosæ, contiguæ. *Stamina* circiter 15, 3-serialia; *filamentis* cohærentibus, apice liberis, 5 externis brevioribus et petalis oppositis; *antheris* dorso medio affixis, extrorsis, loculis parallelis et longitudinaliter dehiscentibus. **FLOR. FÆM.** *Ovarium* conico-ovoideum. *Discus* annularis. *Styli lobi* 3, bifidi. *Capsula* 3-cocca, subsphærica, parum 6-costata, externe subcarnosa. *Semina* oblonga, albumine copioso. *Arbor*, partibus junioribus fugaci-leprosis; *foliis* majusculis, oblongo-lanceolatis, acuminatis, serratis, glabris, penniveniis, longe petiolatis, *petiolo* ad apicem tumido et minute 2-glanduloso; *stipulis* minutis, interne glandula rotundata tectis; *paniculis* terminalibus, elongatis, pendulis; *floribus* plus minus glomeratis, brevi-pedicellatis.

1. **D. Zeylanicus**, Thw.—c.p. 2160.

Var.  $\beta$ . *minor*; subarboreus, partibus omnibus minoribus.—c.p. 2158.

**HAB.** Central Province, up to an elevation of 4000 feet. Var.  $\beta$ . Warmer and southern parts of the island. Nom. vulg. "Wal-kakoona-gass."

*Petala* albida, 3-4 lin. longa. *Capsula* 1 poll. (var.  $\beta$  6 lin.) longa, pubescentia stellata fulva vestita.

Between var.  $\alpha$  and  $\beta$  I can discover no distinctive character but that of size; and in that respect there is considerable difference: the former being a large forest tree; whilst var.  $\beta$  might in some instances be termed a mere shrub, and has flowers and fruit a good deal smaller than those of var.  $\alpha$ .

26. **GIVOTIA**, Griff.

1. **G. rottleriformis**, Griff.; Wight, Ic. t. 1889.—c.p. 1139.

HAB. Hot drier parts of the island, not common.

FLOB. MASC. *Stamina* sæpissime 15, basi coalita, 5 externa breviora.

## 27. SARCOCLINIUM, Wight.

1. **S. longifolium**, Wight. Ic. t. 1887, 1888.—C.P. 596.

HAB. Central Province, at an elevation of 4000 to 6000 feet. Nom. vulg. "Bairoo-gass."

The leaves of this plant, as they are of a firm consistence and do not rapidly decompose, are used by the Cinghalese for thatching.

2. **S. Hookeri**, Thw.; *foliis* magnis, glabris, nitidis, cuneato-lanceolatis, brevi-acuminatis, serrulatis, penniveniis, brevi-petiolatis; *stipulis* magnis, a basi lata lanceolatis, acutis. FLOB. FÆM. *Racemis* foliis subæquilongis, pendulis, glabris; *pedicellis* solitariis, infra medium articulatis; *sepalis* triangularibus, discum annularem magnum vix excedentibus.—C.P. 3429.

HAB. Near Eknalagodde, in the Ratnapoora District, at no great elevation. Nom. vulg. "Maha-bairoo-gass."

*Arbor* mediocris. *Folia* 2-3-pedalia, 5-7 poll. lata, *venis* primariis una ab altera 7-8 lin. distantibus, parallelis. *Stipulæ* 1½-2½ poll. longæ, striatæ, fuscæ. FLOB. FÆM. apetali (vel forsan petalis cito deciduis). *Pedicelli* 5-6 lin. longi. *Ovarium* obtuse trigonum, minute tomentosum. *Stylus* brevis, crassus, a medio 3-partitus; *segmentis* interne stigmatosis, apice emarginatis. *Capsulæ* cocci rotundati. *Semina* spherica, semipollicaria. *Embryo* in medio albuminis copiosi; *cotyledonibus* planis, orbicularibus, *radicula* cylindrica iisdem subæquilonga. *Flor. masc.* nondum vidi.

## 28. AGROSTISTACHYS, Dalz.

Hook. Journ. of Bot. ii. (1850) p. 41. Baillon, Étude des Euph. p. 310.

1. **A. Indica**, Dalz. l. c.—C.P. 2156.

HAB. Central Province, at an elevation of 1000 to 2000 feet.

FLOB. MASC. *Petala* 6-8, albida. *Stamina* 8-13, basi in tubum brevem, pistilli rudimentum cingentem coalita.

### TRIB. BIOVULATÆ.

Sect. 5. BRIEDELIEÆ.—*Perianthii biserialis verticillo interiore petalino.*

\* *Calycis segmentis valvatis.*

## 29. BRIEDELIA, Willd.

1. **B. retusa**, Spr.—*B. spinosa*, Willd. Sp. Pl. iv. p. 979. *Cluytia retusa*, Linn.; Willd. l. c. p. 883. *C. spinosa*, Roxb. Cor. Pl. ii. p. 38, t. 172; Rheed. Hort. Mal. ii. t. 16.—C.P. 2161.

HAB. Common up to an elevation of 2000 feet. Nom. vulg. "Kattakaala-gass."

*Folia* acuta vel obtusa, in arboribus vetustioribus retusa.

2. **B. Moonii**, Thw.; arborea; *foliis* majusculis, oblongis, parum acutis, obtusis vel retusis, minute venosis, subtus villosulis; *florum glomerulis* axillaribus; *fructibus* ovoideis, parum acutis.—*Cluytia retusa*, Moon's Cat. p. 71 (non Linn.).—C.P. 3475.

HAB. Equally common with the preceding, up to an elevation of 2000 feet. Nom. vulg. "Patkaala-gass."

The Cinghalese consider this quite distinct from the preceding, which it however very closely resembles, differing in its somewhat larger leaves, axillary, not spiked inflorescence, and ovoid fruit. They are probably mere varieties of one species. The timber of both is useful for building purposes.

### 30. AMANOA, Aubl.

1. *A. patula*, Thw.—*A. Indica*, Wight, Ic. t. 1911. *Cluytia patula*, Roxb. Cor. Pl. vol. ii. p. 37, t. 170; Willd. Sp. Pl. iv. p. 884.—C.P. 2151, 2164.

HAB. Not uncommon in the hotter parts of the island.

*Arbor* parva vel mediocris, glabra. *Folia* oblonga vel rotundata, longiuscule acuminata, 2–7 poll. longa. *Flores* pilosuli. *Capsula* estipitata vel stipite brevissimo, parce longiuscule pilosa.

2. *A. pallida*, Thw.; arborea, *partibus junioribus* pallide fulvo-pubescentibus; *foliis* oblongis, lanceolatis, acuminatis, mucronatis, basi acutis; *floribus* glomeratis, axillaribus, externe incano-pubescentibus; *capsula* subsessili, longiuscule pallide fulvo-pilosa.—C.P. 2166.

HAB. Hanguranketty, near Kandy.

*Arbor* parva. *Folia* 1½–5 poll. longa, *petiolo* 2–3 lin. longo. *Capsula* parum depressa, circiter 3 lin. in diam.

3. *A. ferruginea*, Thw.; arborea, *partibus junioribus* ferrugineo-pilosis; *foliis* subcoriaceis, oblongis, obtuse rostrato-acuminatis, superne glabris, nitidis, subtus opacis, pilosulis, venis venulisque prominentibus; *floribus* axillaribus, glomeratis, ferrugineo-pilosis; *capsula* stipitata, ferrugineo-tomentosa.—C.P. 2407.

HAB. Central Province, up to an elevation of 3000 feet.

*Arbor* parva. *Folia* 2–5 poll. longa, *petiolo* 2–3 lin. longo. *Capsula* parum depressa, 4 lin. in diam., *stipite* 1 lin. longo.

4. *A. collina*, Baill. Étude des Euph. p. 582.—*Cluytia collina*, Roxb. l. c. t. 169; Willd. l. c. p. 883.—C.P. 2163.

HAB. Colonna Corle, rare. Nom. vulg. "Madara-gass."

NANOPETALUM, Hassk., a specimen of which I have received from M. Binnendijk, in habit resembles the present genus, as well as in its valvate calyx and stipitate capsule; but in its exalbuminate seeds it resembles *Actephila*.

\*\* *Calycis segmentis imbricatis.*

### 31. ACTEPHILA, Blume.

*Anomospermum*, Dalz. An *Savia*, Willd.?

1. *A. Neilgherrensis*, Wight, Ic. t. 1910.—*A. Javanica*, Miq. Fl. Ind. Bat. vol. i. pars altera, p. 356. *Savia Actephila*, Hassk. Cat. Boy. p. 243. An *Anomospermum excelsum*, Dalz. in Hook. Journ. of Bot. iii. (1831) p. 228?—C.P. 2532, 3430, 3431.

HAB. Not very uncommon in the central and southern parts of the island, up to an elevation of 2000 feet.

*Arbor* parva. *Folia* ovato-vel elongato-lanceolata, plus minus acuminata, basi cuneata, 2–8 poll. longa, 1¼–3½ poll. lata, *petiolo* ½–1½ poll. longo. *Pedicelli* longitudine valde varii.

An extremely variable plant, and the different forms so merge into one another,

that I find it impossible to separate them into distinct varieties. In the ripe seeds I can discover no trace of albumen surrounding the embryo.

Sect. 6. PHYLLANTHÆ.—*Flores apetalæ; perianthio biseriali vel sub-biseriali; fructûs loculis dispermis.*

### 32. PROSORUS, Dalz., Thw.

*Cicca* (Sect. *ProSORUS*), Baill. *Zygospermum*, Thw. MSS. olim, Baill.

1. **P. Indica**, Dalz. in Hook. Journ. of Bot. iv. (1852) p. 346; Thw. ibid. vol. vi. (1854) p. 298, t. 10 C, figs. 3, 4, 5, 6; et vol. viii. (1856) p. 272.—C.P. 2155, 3099.

HAB. Not uncommon up to an elevation of 2000 feet. Nom. vulg. "Carron-gass."

FLOR. MASC. numerosissimi,  $1\frac{1}{2}$  lin. lati. *Discus* convexus, gibbus, rugulosus. *Pedicelli* 3-4 lin. longi. FLOR. FÆM. terni. *Pedicelli* 4-5 lin. longi.

2. **P. cyanosperma**, Thw.—*P. Gærtneri*, Thw. l. c. viii. p. 272, et vol. vi. t. 10 C, figs. 1, 2, 7, 8, 9, 10. *Croton? cyanospermum*, Gærtner. De Fruct. vol. ii. p. 120, t. 107. *Zygospermum zeylanicum*, Thw. MSS. olim; Baill. Étud. des Euph. p. 620.—C.P. 2601.

HAB. Ambagamowa and Ratnapoora Districts, up to an elevation of 1000 feet. Nom. vulg. "Sooddoo-leyang-gass."

FLOR. MASC. numerosi, 3 lin. lati. *Discus* concavus, lævis, cum calycis tubo adnatus. *Pedicelli* 4-5 lin. longi. FLOR. FÆM. solitarii. *Pedicelli* 5-8 lin. longi.

This and the preceding are entirely bare of leaves just before flowering. The timber of both is useful for buildings.

### 33. FLUEGGEA, Willd.

An *Chorisandra*, Wight, Ic. t. 1994?

1. **F. leucopyrus**, Willd. Sp. Pl. iv. p. 757; Wight, Ic. t. 1875.—C.P. 2154.

HAB. Very abundant in the hot, drier parts of the island. Nom. vulg. "Kattoo-pilla."

In my specimens the ovary is 3-locular, and the style is tripartite, with the apices bifid; in other respects the plant is well represented in Dr. Wight's figure quoted above. Two plants from the continent of India, given me by Dr. Thomson, and named respectively *F. retusa* and *F. virosa*, appear to be very closely allied to the present species, but have comparatively much shorter petioles.

### 34. PELTANDRA, Wight.

(Icones, tt. 1891, 1892.)

1. **P. flexuosa**, Thw.; suffruticosa, glabra; ramis parum angulatis, striatis, flexuosis; foliis teneris, membranaceis, integris, ovatis, utrinque plus minus acutis, longiuscule petiolatis.—C.P. 3571.

HAB. Minnery, not common.

Humilis, parce ramosa. *Folia* ejusdem stirpis magnitudine valde varia, glaucoviridia, superne secus costam venas primariasque albo-areolata, *petiolo* 1-8 lin. longo. *Inflorescentia* monoica, *rachi* brevissima, bracteata; *bracteis* lanceolatis, acuminatis. *Sepala* rotundata, membranacea. FLOR. MASC. numerosi, singulatim evoluti. *Pedicelli* circiter 2 lin. longi. FLOR. FÆM. sæpius solitarii ad basin racheos. *Pedicelli* circiter 7 lin. longi. *Stylus* tripartitus, lobis profunde bifidis, capitellatis.



35. **KIRGANELIA**, Juss., Baill.Sect. ovario 4- vel pluri-oculari.—*Anisonema*, A. Juss.

1. **K. multiflora**, Baill. Étud. des Euph. p. 614.—*Anisonema multiflora*, Wight, Ic. t. 1899. *Phyllanthus multiflorus*, Willd. Sp. Pl. iv. p. 581; Roxb. Fl. Ind. iii. p. 664. *P. pentandra*, Herb. Roxb. *P. puberula*, Miq. *Rhamnus Zeylanicus*, etc., Burm. Thes. Zeyl. p. 198, t. 88.—c.p. 2142.

Var.  $\beta$ ; *glabra*.—*P. scandens*, Herb. Roxb.—c.p. 327, 2947.

HAB. Hot, drier parts of the island, especially towards the north, very abundant. Nom. vulg. "Wel-kyla."

36. **PHYLLANTHUS**, Linn.

1. **P. Niruri**, Linn. (Roxb. Fl. Ind. iii. p. 659, cum syn.)—c.p. 2138.

Var.  $\beta$ ; *sepalis* 5.—Wight, Ic. t. 1894.—c.p. 2136.

HAB. A very abundant weed in gardens, etc. Var.  $\beta$ . Hot, drier parts of the island. Nom. vulg. "Pittawaaka."

2. **P. Urinaria**, Linn. (Roxb. l. c. p. 660, cum syn.)—*P. leprocarpus*, Wight, Ic. t. 1895-4.—c.p. 2137.

HAB. Equally common with the preceding. Nom. vulg. "Rat-pittawaaka."

3. **P. rotundifolius**, Klein. (Willd. Sp. Pl. iv. p. 584).—c.p. 2134.

HAB. Batticaloa, *Gardner*.

4. **P. Maderaspatensis**, Linn. (Willd. l. c. p. 575).—*P. obcordatus*, Roxb. l. c. p. 656.—c.p. 2133.

HAB. Hot, drier parts of the island, common.

5. **P. polyphyllus**, Willd. l. c. p. 586; Wight, Ic. t. 1895-2.—c.p. 2140, 2502 (2564).

HAB. A very common shrub or small tree in the northern parts of the island.

Sect. 2. *Emblica*, Gærtn.

6. **P. Emblica**, Linn. (Willd. l. c. p. 586, cum syn.; Roxb. l. c. p. 871).—*Emblica officinalis*, Gærtn.; Wight, Ic. t. 1896.—c.p. 2144.

HAB. Common on exposed grassy places, up to an elevation of 4000 feet. Nom. vulg. "Nelli-gass."

The fruit of this tree is much esteemed as a medicine by the Cinghalese. The timber is used for buildings.

Sect. 3. *Macræa*, Wight.

7. **P. simplex**, Willd. l. c. p. 573; Roxb. l. c. p. 654.—An *Macræa oblongifolia*, Wight, Ic. t. 1902-1?—c.p. 2139.

HAB. Common in the warmer parts of the island.

8. **P. Gardneri**, Wight, Ic. t. 1902-3.—*P. fruticosa*, Wall. MSS. An *Macræa ovalifolia*, Wight, Ic. t. 1902-4?—c.p. 5 (54, 296, 501).

Var.  $\beta$ . *pubescens*; tota planta plus minus pilosa.—c.p. 178.

HAB. Central Province, at an elevation of 3000 to 6000 feet.

Differs from the preceding, which it closely resembles in habit, in the firmer tex-

ture of its leaves; larger, more cordate and acuminate stipules, and larger calyx. In both, the capsule, when fresh, is slightly tuberculated. I suspect they will prove to be but forms of one species.

9. **P. myrtifolius**, Wight, Ic. t. 1902-2.—C.P. 650.

HAB. Banks of rivers, up to an elevation of 1500 feet.

*Frutex* majusculus, valde ramosus. *Folia* basi angustata, subauriculata. *Flores*, cum pedicellis, rufescentes.

A very distinct and elegant species, and becoming under cultivation a compact and extremely ornamental shrub.

10. **P. flaccidus**, Thw.; annua, erecta, plus minus ramosa, glabra; ramis striatis, parum angulatis; foliis teneris, ovatis, subacutis, mucronatis; stipulis anguste lanceolatis, petiolo longioribus; FLOR. MASC. 2-3-nis, brevi-pedicellatis; FLOR. FÆM. majoribus, in axillis versus apicem ramorum solitariis; pedicellis triangularibus, superne incrassatis, calyce subæquilongis.—C.P. 320 (510, 2147).

HAB. Central Province, at an elevation of 4000 to 7000 feet.

$\frac{1}{2}$ -3-pedalis. *Folia*  $\frac{1}{4}$ - $1\frac{1}{4}$  poll. longa, petiolo  $\frac{1}{2}$ -1 lin. longo. *Sepala* ovato-oblonga, subacuta. FLOR. MASC. glandulæ rotundatæ, tuberculatæ. FLOR. FÆM. discus margine minutissime crenulatus. *Stylus* 3-partitus, apicibus bifidis, divergentibus.

### 37. EPISTYLIUM, Sw., Baill.

Sect. *flor. fœm.* 6-sepalis.—*Reidia*, Wight, Miq. *Eriococcus*, Hassk.

1. **E. floribundum**.—*Reidia floribunda*, Wight, Ic. t. 1903. *Phyllanthus longiflorus*, Heyne.—C.P. 71, 2531.

HAB. Central Province, at an elevation of 3000 to 5000 feet.

2. **E. montanum**, Thw.; glabrum; foliis ellipticis vel oblongis, abrupte brevissime acuminatis, basi inæqualibus, brevi-petiolatis; pedicellis folio longioribus; sepalis ovatis, fimbriatis; FLOR. FÆM. disco obscure 6-lobo, sublævi; capsula glabra.—C.P. 2148.

HAB. Horton Plains, at an elevation of 7000 feet.

3-4-pedale. *Folia*  $2\frac{1}{2}$ -7 lin. longa,  $1\frac{1}{4}$ -4 lin. lata. *Flor. masc.* 1-3-ni; *fœm.* solitarii.

Allied to the preceding, which it much resembles, but distinguished from it by the absence of pubescence; smaller leaves; fimbriated sepals; and the nearly entire and smooth disk of the female flowers.

3. **E. latifolium**.—*Reidia latifolia*, Wight, Ic. t. 1904-2.—C.P. 2145.

HAB. Central Province, at an elevation of 2000 to 4000 feet.

2-3-pedale. *Folia* firma, subtus areolata, costa venisque dilatatis. *Flor. fœm.* ad apices ramorum racemosi.

4. **E. polyphyllum**.—*Reidia polyphylla*, Wight, Ic. t. 1904-4.—C.P. 432.

HAB. Banks of stream at Leeta-gangoola, near Adam's Peak, at an elevation of 5000 feet.

3-4-pedale. *Sepala* omnia fimbriata. FLOR. FÆM. discus subinteger vel obscure 6-lobus.

### 38. AGYNEIA, Linn.

1. **A. bacciformis**, A. Juss. (Wight, Ic. t. 1893).—*A. phyllanthoides*,

Spr. *Phyllanthus bacciformis*, Linn.; Roxb. Fl. Ind. iii. p. 661.—c.p. 2130.

HAB. Not uncommon near the sea.

### 39. SAUROPUS, Blume.

*Sauropus* et *Ceratogynum*, Wight.

1. **S. Gardneriana**, Wight, Ic. t. 1951-2.—*S. Zeylanica* et *S. Indica*, Wight, Ic. t. 1952; Wall. Cat. n. 7933 *B* et *E*. An *S. albicans*, Blume, Bijdr. p. 596. *Phyllanthus strictus*, Roxb. Fl. Ind. iii. p. 670? —c.p. 2146.

HAB. Central Province, not uncommon up to an elevation of 4000 feet.

I suspect that this will prove to be but a form of Blume's plant, which appears to differ from the present principally in having firmer leaves, with more prominent veins, judging from a specimen I have from M. Binnendijk.

2. **S. retroversa**, Wight, Ic. t. 1951-1.—c.p. 3134.

HAB. Oodoopussalawa, in the Central Province, at an elevation of about 4000 feet.

*Frutex* erectus, glaber. *Folia* 1-2½ poll. longa, ½-1¼ poll. lata, a basi lata vel sæpe parum cordata gradatim acuminata, *petiolo* 1-1½ lin. longo. FLOR. MASC. pallide virides, flavescentes, margine inferiore *perianthii* duplicati, pileiformis, circiter 12-dentata. FLOR. FÆM. *perianthium* 3-4 lin. in exp., lobis horizontalibus, rotundatis, 3 exter. minoribus, subacutis. *Capsula* albida, pyriformis, pollicaris.

3. **S. assimilis**, Thw.; fruticosa, glabra; *ramis* parum angulatis; *foliis* membranaceis, teneris, ovato-lanceolatis, acuminatis, basi plus minus acutis; FLOR. MASC. lobis *perianthii* duplicati, æqualibus, anguste cylindricis, obtusis, curvatis, suberectis.—c.p. 2855.

HAB. Allagalla, in the Central Province, at an elevation of 3000 feet.

*Folia* 1½-3 poll. longa, *petiolo* 1½ lin. longo. FLOR. MASC. pallide virides, flavescentes, lobis *perianthii* 2 lin. longis. *Antheræ* subsessiles. *Flor. fœm.* nondum vidi.

This in general appearance closely resembles *S. Gardneriana*, but the flowers are more like those of *Ceratogynum rhamnoides*, Wight, Ic. t. 1900, from which latter species, however, it would seem to be sufficiently distinguished by its much larger and differently shaped leaves, and the obtuse, cylindrical, narrower lobes of the perianth.

4. **S. rigida**, Thw.; fruticosa, erecta, glabra, rigida, ramosa; *ramis* angulatis; *foliis* ovalibus vel rotundatis, brevi-petiolatis; *floribus* paucis, axillaribus, brevi-pedicellatis; MASC. lobis *perianthii* duplicati horizontalibus, triangularibus, margine exteriori retusa vel parce sinuata; FÆM. *sepalis* obovoideis, obtusis, apiculatis; *ovario* apice concavo, *stigmata* bifida erecta includente.—c.p. 2135.

HAB. Hot, drier parts of the island, not very common.

1-2-pedalis. *Folia* 3-7 lin. longa, 1¼-5 lin. lata, superne nigro-viridia, medio albo-notata, subtus pallida. *Flores* pallide virides, flavescentes; MASC. 1 lin., FÆM. 3-4 lin. in exp. *Capsula* ovoidea, 3 lin. longa.

The ovary of this species bears a very considerable resemblance to that of *Agyneia bacciformis*. *Phyllanthus rhamnoides*, Roxb. (non Willd.), would appear, from the description, to be nearly allied to this; but the long-pedicelled male flowers of that plant are sufficient to prove it distinct.

## 40. MELANTHESA, Blume.

1. *M. rhamnoides*, Blume, Bijdr. p. 591 (non *Phyllanthus rhamnoides*, Willd. vel Roxb.); Wight, Ic. t. 1898-1.—*Phyllanthus Vitis-Idæa*, Kœn.; Roxb. Fl. Ind. iii. p. 665 (haud *Rhamnus Vitis-Idæa*, Burm. Fl. Ind. p. 61).—*P. cinerascens*, Wall. An *Melanthesa virgata* et *M. racemosa*, Bl. l. c. ?—c.p. 2141.

HAB. Very abundant up to an elevation of 3000 feet. Nom. vulg. "Gass-kyla."

Specimens from the Java Botanic Garden, named respectively *Melanthesa virgata* and *M. racemosa*, Blume, appear to me to differ in no essential respect from the present plant, but they are not in a sufficiently perfect state to enable me to determine satisfactorily.

2. *M. turbinata*, Wight, Ic. t. 1897.—*M. obliqua*, Wight, Ic. t. 1898-2. *Phyllanthus turbinatus*, Kœn.; Roxb. l. c. p. 666. *P. patens*, Roxb. l. c. p. 667. *P. rhamnoides*, Herb. Heyne. *P. glaucifolia*, Wall. Cat. n. 7911 B, E.—c.p. 2143.

HAB. Common up to an elevation of 3000 feet. Nom. vulg. "Walmooroonga."

## 41. GLOCHIDION, Forst., Blume.

*Bradleia*, Gært. *Gynoon*, A. Juss.

(Nom. vulg. "Hoonoo-kirilla-gass.")

1. *G. Zeylanicum*, Thw.—*Bradleia Zeylanica*, Gært. De Fr. ii. p. 12, t. 109. *B. obliqua*, Wall.

Var. *a*; tota planta glabra; FLOR. FÆM. *stylo* brevi, crassiusculo.—c.p. 2149.

Var. *β*; *ramulis, petiolis, foliis* subtus *inflorescentiaque* brevi-tomentosis; FLOR. FÆM. *stylo* subnullo.—"Falconer, n. 845, Amherst," in Hort. Calc.—c.p. 3432.—An species distincta?

HAB. Central and southern parts of the island, up to an elevation of 2000 feet.

*Arbor parva. Folia* firma, oblonga, acuminata, basi rotundata, obliqua, 3-6 poll. longa, circiter 2 poll. lata, *petiolo* 2 lin. longo. *Flores* umbellati. *Pedunculi* solitarii, axillares vel supra-axillares, *petiolo* 2-3-plo longiores. *Sepala* rotundata, subacuta. FLOR. MASC. *Antheræ* 5-6. FLOR. FÆM. *Ovarium* 5-6-loculare. *Capsula* oblato-sphærica, semipollicaris.

2. *G. coriaceum*, Thw.; glabrum; *foliis* majusculis, coriaceis, ovatis vel oblongis, obtuse acuminatis, mucronatis, basi obliquis, abrupte in *petiolum* angustatis; *sepalis* flaccidis, rotundatis vel late oblongis, subacutis; FLOR. MASC. *antheris* 3; FLOR. FÆM. sessilibus vel subsessilibus; *stylo* crasso, brevi, oblongo vel suborbiculari, apice 3-fisso; *capsula* subsphærica, depressa, parum triloba.—c.p. 3016, 342, 2529.

HAB. Central Province, at an elevation of 3000 to 4000 feet.

*Arbor parva. Folia* 2-6 poll. longa, 1-2½ poll. lata, *petiolo* circiter 2 lin. longo. FLOR. MASC. *Pedicelli* 3-4 lin. longi. FLOR. FÆM. *Capsula* 4-5 lin. in diam.

An extremely variable species.

3. *G. Jussieuium*.—*Gynoon Jussieuium*, Wight, Ic. t. 1908. *G. triandrum*, W. et A.?, Wight, l. c.—c.p. 2561 (? 2560).

HAB. Central Province, up to an elevation of 4000 feet.

I have received this species from Java, so that it would appear to be widely dis-

tributed. c.p. 2560 I believe to be an abnormal state of the present plant, in which the sepals have cohered into a tridentate tubular perianth.

4. **G. Moonii**, Thw.—*Gynoon hirsutum*, Wight, Ic. t. 1909 (haud *Bradleia hirsuta*, Roxb.). *Phyllanthus pubescens*, Moon's Cat. p. 65.—c.p. 68, 258, 2150.

HAB. Very common up to an elevation of 4000 feet.

*Frutex* vel *arbor* parva, *ramulis* hirsutis vel tomentosus. *Folia* ovato-vel oblongo-lanceolata, acuminata, plus minus tomentosa vel aliquando subglabra, 2–8 poll. longa, 1–3 poll. lata, basi parum acuta, brevi-petiolata. *Inflorescentia* hirsuta. *Sepala* lineari-oblonga. FLOR. FÆM. sessiles vel subsessiles. *Stylus* plus minus elongatus clavatusque, pilosus, apice trifidus. *Capsula* pilosa.

5. **G. montanum**, Thw.; arborea; *ramulis* parum angulatis et cum *petiolis inflorescentiaque* tomentosus; *foliis* ovato-vel oblongo-lanceolatis, obtusis, mucronatis, plus minus tomentosus vel subglabris, basi obliquis, rotundatis vel rarius parum acutis; *sepalis* oblongis; FLOR. FÆM. sessilibus; *stylo* cylindrico vel parum clavato, dense piloso vel subglabro, apice trifido; *capsula* glabra, depressa, triloba.—c.p. 3133.

HAB. Oova, in the Central Province, at an elevation of 4000 to 6000 feet.

15–20-pedalis. *Folia* 1½–3 poll. longa, ¾–1½ poll. lata, *petiolo* 1½ lin. longo.

Closely allied to the preceding species, from which it differs in its more arboreous habit, smaller, less acuminated leaves, and glabrous capsules.

6. **G. Gardneri**, Thw.; arborea; *ramulis* glabris; *foliis* glabris, acuminatis, mucronatis, basi obliquis, in *petiolum* brevem angustatis; *inflorescentia* plus minus pilosa; *sepalis* oblongis; FLOR. FÆM. sessilibus; *stylo* glabro, elongato, clavato, apice trifido; *capsula* glabra, depressa, triloba.—c.p. 3156.

Var.  $\alpha$ ; *foliis* ovatis, abrupte breveque acuminatis.

Var.  $\beta$ ; *foliis* oblongis, longiuscule acuminatis; *inflorescentia* minus pilosa.

HAB. Var.  $\alpha$ . Central Province, Gardner. Var.  $\beta$ . Reigam Corle.

*Arbor* parva. *Folia* (var.  $\alpha$ ) 1½–3 poll. longa, 1–1½ poll. lata; (var.  $\beta$ ) 2½–4 poll. longa, ¾–1¼ poll. lata.

7. **G. nemorale**, Thw.; arborea; *ramulis*, *petiolis inflorescentiaque* brevi-pilosis; *foliis* glabris, oblongo-lanceolatis, longiuscule acuminatis, subtus venis prominentibus, basi obliquis, acutis; *sepalis* oblongis; FLOR. FÆM. fasciculatis, *pedicellis* *petiolo* sæpius longioribus; *stylo* elongato, clavato, glabro; *capsula* glabra, depressa, triloba.—c.p. 3015.

HAB. Pasdun Corle, at no great elevation.

*Folia* 3½–7 poll. longa, 1½–2 poll. lata, *petiolo* circiter 2 lin. longo.

PHYLLANTHUS STELLATUS, Retz, Willd. Sp. Pl. iv. p. 577, is probably one of the foregoing species of *Glochidion*, but the description is too imperfect for the identification of the plant.

Sect. 7. PIERARDIÆ.—*Flores* apetalæ, *perianthio* biseriali vel sub-biseriali. *Ovula* bina in *loculo* sub *processu* magno cuculliformi *placentæ* affixa. *Fructus* monospermus vel *loculis* monospermis.

## 42. CYCLOSTEMON, Blume.

*Sphragidia*, Thw.

1. **C. Zeylanicum**, Baill. Étud. des Euphorb. p. 562.—*Sphragidia*

*Zeylanica*, Thw. in Hook. Journ. Bot. vii. (1855) p. 270, t. 10.—C.P. 2424.

HAB. Central Province, at an elevation of about 3000 feet.

*Arbor* 30-40-pedalis, *partibus junioribus* parce fugaci-pilosis. *Folia* 5-10 poll. longa,  $1\frac{1}{2}$ -3 poll. lata, *petiolo* 4-8 lin. longo. *Flores* pallide ochracei. *Sepala* 4-5, rotundata, obtusa, externe cum pedicellis sericeo-pilosa. *Fructus* subpollicaris.

#### 43. HEMICYCLIA, W. et A., Thw.

*Astylis*, Wight.

1. *H. sepiaria*, W. et A. (Wight, Ic. t. 1872; Thw. in Hook. Journ. of Bot. vii. (1855) p. 271).—C.P. 2120.

HAB. Abundant in the hot, drier parts of the island. Nom. vulg. "Weera-gass."

2. *H. Gardneri*, Thw. l. c.—C.P. 2121.

HAB. With the preceding species, but not so abundant.

*Arbor* mediocris; *partibus junioribus*, *petiolis inflorescentiaque* pubescentibus. *Folia* lanceolata, crenata, retusa, mucronata, basi obliqua, angustata et subtus secus costam pilis longis stipata, 1-3 poll. longa,  $\frac{1}{2}$ -1 poll. lata, *petiolo* 1-2 lin. longo. *Flores* pauci, pallide ochracei, 6 lin. in exp. MASC. *Stamina* circiter 24, *discum* medio excavatum cingentibus. *Ovarii rudimentum* nullum. FÆM. *Stylus* brevis. *Stigma* convexum, carnosum. *Drupa* oblonga, 4 lin. longa.

3. *H. lanceolata*, Thw.; arborea; *ramulis*, *petiolis inflorescentiaque* pilosis; *foliis* subglabris, lanceolatis, obtuse acuminatis, mucronatis, basi obliquis, subacutis et subtus secus costam longe pilosis; *floribus* paucis, parvis; *drupa* oblonga, ad apicem obliqua; *putamine* cristato, reticulato.—C.P. 2211.

HAB. Cultura.

*Arbor* mediocris. *Folia* 1-3 poll. longa,  $\frac{1}{2}$ -1 poll. lata, *petiolo* 1-2 lin. longo. *Flores* circiter 2 lin. in. exp. MASC. *Stamina* circiter 16. FÆM. *Stylus* brevis, in laminam bifidam dilatatus, reflexus. *Drupa* rubra, 6 lin. longa, 3 lin. lata.

#### 44. PALENGA, Thw.

Hook. Journ. of Bot. viii. (1856) p. 270. Baill. Étud. des Euph. p. 649.

1. *P. Zeylanica*, Thw. l. c. p. 271, t. 7 C.—C.P. 3349.

HAB. Ambagamowa District, at an elevation of about 2000 feet. Nom. vulg. "Palenga-gass."

*Arbor* 40-50-pedalis, glabra, *ramulis* teretibus. *Folia* lanceolata, acuminata, pennivenia, obliqua, nitida, nigro-viridia, subtus sparsim glanduloso-punctata, 3-5 poll. longa,  $1-1\frac{1}{4}$  poll. lata, *petiolo* ruguloso, 2 lin. longo. *Stipulae* minutæ. *Flores* albidæ,  $\frac{3}{4}$  lin. longi. *Pedicelli* 1 lin. longi. *Fructus* 8 lin. longus, 4 lin. latus, fusco-tomentosus.

#### 45. PUTRANJIVA, Wall.

*Nageia*, Roxb. (non Gærtn.)

1. *P. Roxburghii*, Wall. Tent. Fl. Nep. p. 61; Cat. n. 6814; Royle, Pl. Him. t. 100; Wight, Ic. t. 1876.—*Nageia Putranjiva*, Roxb. Fl. Ind. iii. p. 767.—C.P. 2122.

HAB. Hot, drier parts of the island.

46. **APOROSA**, Blume, Miq., Baill.*Scepa*, Lindley. *Lepidostachys*, Wall.

(Nom. vulg. "Kabella-gass.")

1. **A. Lindleyana**, Thw.—*Scepa Lindleyana*, Wight, Ic. t. 361.—C.P. 2153.

HAB. Very abundant up to an elevation of 2000 feet.

*Arbor* mediocris, valda ramosa. FLO. FÆM. *Pedicelli*  $\frac{1}{2}$ –1 lin. longi, fructiferi 2 lin. longi. *Ovarium* ampullæforme, fulvo-pilosum, 2-vel 3-loculare. *Stigma* 2–3-partitum, lobis bifidis, tuberculatis vel parum plumosis. *Fructus* subsphæricus, brevi-apiculatus, sessilis, semipollicaris. *Semina* tunica externa carnosâ, flavâ, sapida.Closely allied to and resembling in general appearance *Lepidostachys Roxburghii*, Wall. Cat. No. 6816, but differing from it in its smaller leaves, which are frequently rounded at the base, its pedicellate female flowers, and its fruit being not at all stipitate.2. **A. lanceolata**, Thw.—*Lepidostachys lanceolata*, Tul. in Ann. des Sci. Nat. (Bot.) sér. 3, v. 15, p. 254.—C.P. 2152.

HAB. Equally common with the last.

*Arbor* parva. FLO. MASC. sæpius diandri, ovarii rudimento minuto, inconspicuo. FLO. FÆM. *Spicæ* circiter 5-floræ. *Ovarium* ampullæforme, glabrum. *Stigma* breve, sessile, disciforme, 2-lobum, lobis bifidis sæpe denticulatis. *Fructus* subsphæricus, pisi magnitudine. *Semina* tunica externa carnosâ, flavâ, sapida.

The fruit of this, like that of the preceding species, is eaten by children.

3. **A. latifolia**, Thw.; *foliis* subintegris, majusculis, glabris, subcoriaceis, obovatis, obtusis, basi plus minus rotundatis vel subcordatis; *spicis* ferrugineo-tomentosis; *bracteis* rotundatis, ciliatis; FLO. MASC. sæpissime diandris; FLO. FÆM. sessilibus; *ovario* piloso; *stigmatibus* tripartito, lobis reflexis, apice bifidis; *fructibus* subsphæricis, parum conicis, pericarpio valde carnosâ firmo.—*Agyneia latifolia*, Moon, Cat. p. 65.—C.P. 3433.

HAB. South of the island, at no great elevation.

*Arbor* mediocris. *Folia* 4–8 poll. longa, 3–5 poll. lata, *venis* primariis utrinque circiter 7; *petiolo*  $\frac{1}{2}$ –1 poll. longo. *Fructus* pollicares.4. **A. fusiformis**, Thw.; *foliis* subintegris, coriaceis, glabris, obovatis vel rotundatis, breve obtuseque acuminatis vel retusis, basi acutis vel rotundatis; *spicis* ferrugineo-tomentosis; *bracteis* rotundatis, ciliatis; FLO. MASC. sæpius diandris; FLO. FÆM. confertis; *ovario* glabro; *stigmatibus* 2–3-partito, lobis revolutis, apice bifidis; *fructibus* fusiformibus, pericarpio parce carnosâ.—*Lepidostachys grandifolia*, Planch. MSS. in Herb. Hook.—C.P. 30.

HAB. Central Province, at an elevation of 4000 to 6000 feet.

*Arbor* mediocris. *Folia*  $2\frac{1}{2}$ –5 poll. longa,  $1\frac{3}{4}$ –3 poll. lata; *venis* primariis utrinque circiter 5; *petiolo*  $\frac{1}{2}$ – $\frac{3}{4}$  poll. longo. *Fructus* 6–8 lin. longi, 3–4 lin. lati.The very different form of the much less fleshy fruit of the present species abundantly distinguishes it from *A. latifolia*, to which it bears a very great general resemblance.5. **A. acuminata**, Thw.; *foliis* glabris, integris, membranaceis, ovato-vel oblongo-lanceolatis, acuminatis, basi acutis; *spicis* ochraceo-pilosis, brevibus, sæpius solitariis; FÆM. sæpissime unifloris; *ovario* ovoideo, piloso, 3-loculari; *stigmatibus* majusculo, 3-partito, lobis apice bifidis; *fructibus* subsphæricis, sessilibus, pericarpio valde carnosâ firmo.—C.P. 2530.

HAB. Ambagamowa District and near Balangodde, at an elevation of 2000 to 3000 feet.

*Arbor* parva, parce ramosa, *partibus junioribus* pilosis. *Folia*  $1\frac{1}{2}$ -4 poll. longa,  $\frac{1}{2}$ -2 poll. lata, *petiolo* 2-4 lin. longo. *Fructus* circiter  $\frac{3}{4}$  poll. in diam.; viridi-rufescentes, sæpius 3-spermi. *Semina* tunica externa carnosâ, flavo-rubra.

Sect. 4. ANTIDESMÆ; *perianthio uniseriali*; *'disco glanduloso maximo genitalia cingente.*—*Sectioni præcedenti valde affinis.*

#### 47. ANTIDESMA, Linn. Burm.

1. **A. paniculatum**, Roxb., Willd. Sp. Pl. iv. p. 764; Roxb. Fl. Ind. iii. p. 770; Tulasne in Ann. des Sci. Nat. (Bot.) sér. 3, vol. xv. p. 228.—*A. pubescens*, Roxb. Pl. Cor. ii. t. 167; Willd. l.c. p. 763; Roxb. Fl. Ind. iii. p. 770 (non Tulasne); Wight, Ic. tt. 820, 821.—c.p. 773.

HAB. Common up to an elevation of 2000 feet. Nom. vulg. "Boo-ambilla-gass."

2. **A. Bunius**, Spr., Syst. Veg. i. p. 826; Wight, Ic. t. 819; Tul. l.c. p. 186.—*A. Alexiteria*, Linn. (partim). *A. comptum*, Tul. l.c. p. 190. *A. floribundum*, Tul. l.c. p. 189. *Stilago Bunius*, Linn.; Roxb. Fl. Ind. iii. p. 758; Rheed. Hort. Mal. iv. t. 56.—c.p. 660 (766).

Var.  $\beta$ ; *foliis* firmioribus; *inflorescentia* delictiore et minus pubescente.—An *A. glabrum*, Tul. l.c. p. 188?—c.p. 2922.

HAB. Common up to an elevation of 3000 feet. Var.  $\beta$ ; Caltura. Nom. vulg. "Carawilla-cabilla-gass."

3. **A. lanceolarium**, Wight, Ic. t. 776.—*A. lanceolatum*, Tul. l.c. p. 195. *Stilago lanceolaria*, Roxb. l.c. p. 760.—c.p. 2602.

HAB. Ambagamowa, Ratnapoora, and Galle Districts, up to an elevation of 1500 feet.

A very variable plant as regards the size and shape of the leaves. I have a small form from the Singe Rajah forest, in which the leaves are linear-lanceolate, and some of them not more than 2 lines wide, but there is a gradual transition to the ordinary form of the plant.

4. **A. Zeylanicum**, Lam. Encyc. Bot. i. p. 207; Spr. Syst. Veg. i. p. 826; Tul. l.c. p. 209.—*A. Alexiteria*, Linn. (partim); Burm. Thes. Zeyl. p. 22, t. 10. (An *A. sylvestris*, Lam. l.c.; Spr. l.c.; Rheed. l.c. v. t. 26?)—c.p. 2210.

HAB. Common in the hotter parts of the island. Nom. vulg. "Heen-ambilla-gass."

5. **A. montanum**, Thw.; *arboreum*; *ramulis, petiolis, inflorescentia-que* cinereo-tomentellis; *foliis* integris, subglabris, ovatis vel oblongis, obtuse rostrato-acuminatis, superne ad costam canaliculatis, basi acutis: *racemis* paniculatis vel simplicibus, axillaribus terminalibusque; *bracteis* brevibus, latis, acutis; *floribus* sæpius 5-meris; *calyce* cupuliformi, *lobis* brevibus, rotundatis vel parum acutis.—c.p. 2208.

HAB. Central Province, at an elevation of 3000 to 6000 feet.

*Arbor* mediocris, ramosa. *Folia* 2-4 poll. longa,  $\frac{3}{4}$ -2 poll. lata, *petiolo* 2-5 lin. longo. *Pedicelli* calyce subæquilongi. FL. MASC. *Filamenta* in disco sublævi ovarium abortivum mamillæforme cingente inserta. FL. FÆM. *Fructus* subsphærici, parum compressi, pisi magnitudine, purpurei.

Closely allied to the preceding species, from which it differs in its larger leaves, canaliculate above, longer petioles, and pedicellate pentamerous flowers.



*Euphorbiaceis affines.*48. **DAPHNIPHYLLUM**, Blume, Baill.*Goughia*, Wight. *Gyandra*, Wall.1. **D. Neilgherrense**, Thw.—*Goughia Neilgherrensis*, Wight, Ic. tt. 1878, 1879.—c.p. 491.

HAB. Central Province, at an elevation of 4000 to 6000 feet.

49. **SARCOCOCCA**, Lindl.1. **S. pruniformis**, Lindl. Bot. Reg. xii. t. 1012.—*S. trinervia*, Wight, Ic. t. 1877.—c.p. 203.

HAB. Central Province, very abundant at an elevation of 5000 to 8000 feet.

Extremely variable in the shape of the leaves, which differ from nearly orbicular to narrow-lanceolate acuminate.

50. **PYRENACANTHA**, Hook.1. **P. volubilis**, Hook. Bot. Misc. ii. p. 108, tt. 9, 10.—c.p. 1024.

HAB. Hotter parts of the island.

CXXXV. **CALLITRICHACEÆ**.1. **CALLITRICHE**, Linn.1. **C. verna**, Linn.—*C. Wightiana*, Wall.; Wight, Ic. t. 1947.—c.p. 303, 3636.

HAB. Not uncommon in wet places, at Matturatte, Horton Plains, and other of the more elevated parts of the island.

CXXXVI. **CERATOPHYLLACEÆ**.1. **CERATOPHYLLUM**, Linn.1. **C. verticillatum**, Roxb. Fl. Ind. iii. p. 624.—*C. muricatum*, Cham. *C. tuberculatum*, Cham., et *C. Missionis*, Wall.; Wight, Ic. t. 1948.—c.p. 2311.

HAB. Hotter parts of the island.

There can be no doubt, I think, that Dr. Wight is correct in supposing that the three forms, figured by him in plate 1948 of the 'Icones,' are merely varieties of Roxburgh's plant.

CXXXVII. **NEPENTHACEÆ**.1. **NEPENTHES**, Linn.1. **N. distillatoria**, Linn. (Burm. Thes. Zeyl. p. 42, t. 17, cum syn.)—c.p. 262.

HAB. Very common in the south of the island, at no great elevation.  
Nom. vulg. "Bāndoorā-wel."

The long tough stems are used for tying fences, and for other purposes, by the Cinghalese.

## CXXXVIII. ARISTOLOCHIACEÆ.

### 1. ARISTOLOCHIA, Tourn.

1. **A. bracteata**, Retz, (Willd. Sp. Pl. iv. p. 160; Roxb. Fl. Ind. iii. p. 490.)—C.P. 2256.

HAB. North of the island, *Gardner*.

2. **A. Indica**, Linn.; Roxb. l. c. p. 489.—*A. lanceolata*, Wight, Ic. t. 1858.—C.P. 2258.

HAB. Very abundant up to an elevation of 3000 feet.

Most variable in the size and shape of its leaves; and these, in some of my specimens of what I take to be the present species, seem to make an approach in form to those of *A. acuminata*, Lamk.

### 2. BRAGANTIA, Lour.

1. **B. Wallichii**, R. Br. in Wall. Cat. n. 7415; Wight, Ic. t. 520.—*B. siliquosa*, Miq. *Trimeriza piperina*, Lind. in Bot. Reg. xviii. sub tab. 1543.—C.P. 2257, 3457, 3533.

HAB. Shady woods, up to an elevation of 4000 feet.

Varies very much in the amount of pubescence; in the size and shape of its leaves, which are sometimes somewhat obovate; and in the length of the seed-vessel.

### 3. TRICHOPIDIUM, Lindl., Endl.

*Trichopus*, Gærtn.

*Flores* hermaphroditi. *Perianthii lobi* 6, biseriales, persistentes. *Stamina* 6, libera, subsessilia, lobis perianthii opposita; *antheræ* loculis parum distantibus, parallelis, introrsis; *connectivo* in processum lanciformem terminalem anthera longiorem producto. *Ovarium* 3-loculare; *loculis* 2-ovulatis; *ovulis* pendulis superpositis. *Stylus* brevissimus. *Stigmata* 3, brevia, apicibus reflexis, bifidis.—Herba *glabra*, *subacaulis*; *foliis numerosis*, *petiolatis*, *forma valde variis*; *stipulis lanceolatis*, *acutis*, *persistentibus*; *petiolis in medio longitudinis florigeris*; *floribus fasciculatis*, *longiuscule pedicellatis*; *bracteis lanceolatis*, *acutis*.

1. **T. Zeylanicum**, Thw.—*T. cordatum*, *T. intermedium*, et *T. angustifolium*, Lindl. in Bot. Reg. xviii. sub tab. 1543. *Trichopus Zeylanicus*, Gærtner, De Fruct. p. 44, t. 14.—C.P. 467.

HAB. Common on the sandy margins of streams, at no great elevation. Nom. vulg. "Bin-tal."

*Folia* triangularia, lanceolata vel lineari-lanceolata, tri-vel tripli-nervia, 2-8 poll. longa, *petiolo* 1-8 poll. longo. *Flores* circiter 1 lin. longi, nigro-brunnei, lobis perianthii lanceolatis acutis. *Fructus* 6-7 lin. longi; *pedicello* longo, tenui, apicem versus incrassato.

## CXXXIX. PIPERACEÆ.

## 1. PEPEROMIA, Ruiz and Pav.

1. *P. reflexa*, A. Dietr. (Miq. Fl. Ind. Bat. vol. i., pars altera, p. 436, cum syn.; Wight, Ic. t. 1923-1.)—c.p. 2462.

HAB. Common in the Central Province, up to an elevation of 6000 feet.

2. *P. Heyneana*, Miq. (Wight, Ic. t. 1922-1.)—An *P. Dindygulensis*, Miq. Syst. p. 112; Illust. Piper. p. 19, t. 13; Wight, Ic. t. 1921?—c.p. 2956.

HAB. Not uncommon in the Central and Southern Provinces.

Varies greatly in the amount of pubescence and in the shape and size of the leaves.

3. *P. Courtallensis*, Miq. in Hook. Lond. Journ. Bot. v. p. 549; Wight, Ic. t. 1923-2.—An *P. portulacoides*, A. Dietr.; Wight, Ic. t. 1922-2? An *P. Zeylanica*, Miq. l. c. p. 550?—c.p. 2173.

Var.  $\beta$ ; *foliis majoribus acutioribusque, sæpius alternis*.—c.p. 2172. An species distincta?

HAB. Central Province, at an elevation of 3000 to 5000 feet.

## 2. POTHOMORPHE, Miq.

1. *P. subpeltata*, Miq. (Wight, Ic. t. 1925.)—c.p. 2714.

HAB. Common in damp shady places, up to an elevation of 4000 feet. Nom. vulg. "Mala-laboo."

## 3. CHAVICA, Miq.

1. *C. Roxburghii*, Miq. Syst. Piper. p. 239; Illust. Piper. p. 33, t. 30; Wight, Ic. t. 1928.—*Piper longum*, Linn.—c.p. 2953.

HAB. Hotter parts of the island, not very common; also in native gardens. Nom. vulg. "Tippili."

2. *C. Siriboa*, Miq. Fl. Ind. Bat. vol. i., pars altera, p. 438, cum syn.—*Piper Siriboa*, Linn.—c.p. 3690.

HAB. Cultivated in native gardens as a masticatory, and considered not indigenous. Nom. vulg. "Rata-boolat-wel," "Siribo."

A species (c.p. 3691), called by the Cinghalese "Seewiya-wel," evidently very closely allied to the present, if not, as I suspect, a mere form of it, occurs upon large trees and on rocks, near the banks of rivers and streams, in the Central Province; but I have not yet been so fortunate as to meet with specimens in flower. The leaves are rather firmer in texture and less suddenly acuminate than in the cultivated *C. Siriboa*, but in other respects are very similar in character to those of the latter.

3. *C. Betle*, Miq. l. c. p. 439, cum syn.; Wight, Ic. t. 1926.—*Piper Betle*, Linn.—c.p. 3692.

HAB. Cultivated very generally, as a masticatory, in native gardens in the warmer parts of the island, but not truly wild. Nom. vulg. "Boolat-wel."

## 4. PIPER, Linn.

1. *P. nigrum*, Linn. (Miq. Illust. Piper. p. 50, t. 50; Wight, Ic. t. 1934.)—c.p. 2176.

HAB. Ambagamowa District; also extensively cultivated in native gardens. Nom. vulg. "Gam-mirris-wel."

2. **P. trineuron**, Miq. in Hook. Lond. Journ. of Bot. v. p. 555; Walp. Ann. i. p. 601.—c.p. 2175.

HAB. Central Province, at an elevation of 4000 to 7000 feet.

A very distinct species, characterized by its narrow very coriaceous leaves, closely adnate bracts, and somewhat distant flowers. The fruit is slightly oblong, 3 lines in diameter.

3. **P. sylvestre**, Lam. (Wight, Ic. t. 1937.)—An *P. hymenophyllum*, Miq. l. c. p. 554; Wight, Ic. t. 1942?—c.p. 3688.

HAB. On forest trees, up to an elevation of 2000 feet. Nom. vulg. "Mala-mirris-wel," "Wal-merris-wel," "Wal-gam-mirris-wel."

4. **P. argyrophyllum**, Miq. (Illust. Piper. p. 55. t. 56; Wight, Ic. t. 1941.)—An *P. Walkeri*, Miq. in Hook. Lond. Journ. of Bot. iv. p. 436; Walp. Ann. iii. p. 344?—c.p. 3625.

HAB. Central Province.

Closely allied to the preceding, from which it seems to differ but little, except in the texture and colour of the foliage and in its more oblong fruit.

5. **P. arborescens**, Roxb. (Fl. Ind. i. p. 159; Miq. Fl. Ind. Bat. vol. i., pars altera, p. 452; Illust. Piper. t. 54; Wight, Ic. t. 1940.)—*P. Linkii*, Miq. l. c. p. 53, t. 53. An *P. Zeylanicum*, Miq. in Hook. Journ. of Bot. iv. p. 436?—c.p. 35, 2461.

HAB. Central Province, up to an elevation of 3000 feet.

6. **P. arcuatum**, Blume, Enum. Pl. Jav. i. p. 67; Miq. Illust. Piper. p. 57, t. 48.—c.p. 2177, 2178.

HAB. More elevated parts of the Central Province.

Very nearly allied to the preceding, from which it differs principally in its smaller leaves, shorter spikes of flowers, and the distinctly peltate bracts of its male flowers.

## CXL. CHLORANTHACEÆ.

### 1. CHLORANTHUS, Sw.

1. **C. brachystachys**, Blume, Fl. Jav. p. 13, t. 2; Miq. Fl. Ind. Bat. i. p. 801, cum syn.—*Ascarina serrata*, Bl. Enum. i. p. 80. *Sarcandra chloranthoides*, Gardner in Calc. Journ. of Nat. Hist. vi. p. 348; Walp. Ann. iii. p. 353; Wight, Ic. t. 1946.—c.p. 2207.

HAB. Not uncommon in forests of the Central Province, at an elevation of 3000 to 5000 feet.

## CXLI. BALANOPHOREÆ.

### 1. BALANOPHORA, Forst.

1. **B. Indica**, Wall. Cat. 7247; Weddell in Ann. Sc. Nat. ser. 3, vol. xiv. p. 167, t. 9, fig. 11-22; Hook. f. in Trans. Linn. Soc. vol. xxii. p. 46.—*Langsdorffia Indica*, Arn. in Hook. Ic. Plant. tt. 205, 206, et in Ann. Nat. Hist. ii. p. 36.—c.p. 200.

HAB. Forests in the Central Province, at an elevation of 3000 to 5000 feet.

## CXLII. TRIURIDEÆ.

## 1. SCIAPHILA, Blume, Endl.

† *Perianthio* 6-partito. *Staminibus in flore masc.* 3.—*Aphylleia*, Champion.

1. **S. secundiflora**, Thw. MS. (Bentham in Hook. Journ. of Bot. vii. (1855) p. 10.)—C.P. 2665.

HAB. Forest of Gattehatte, near Sittawake, at no great elevation.

*Rhizoma* repens, radicans. *Scapi* erecti, 6–14 poll. longi, albido-purpurascens, pauciflori. *Bracteæ* latæ, acutæ. *Flores* racemosi, monoici. *Pedicelli* secundi,  $1\frac{1}{2}$ – $2\frac{1}{2}$  lin. longi. *Perianthii segmenta* longe subulato-acuminata, imberbia, circiter  $1\frac{1}{2}$  lin. longa. FLOR. MASC. superiores. *Antheræ* late sessiles, transverse oblongæ, contiguæ, albæ. FLOR. FÆM. *Ovaria* parum tuberculata. *Stylus* ovarium vix superans, clavato-penicillatus.

2. **S. erubescens**, Miers in Trans. Linn. Soc. xxi. 48.—*Aphylleia erubescens*, Champion in Calc. Journ. of Nat. Hist. vii. p. 468.—C.P. 2666.

HAB. Narawalla, near Galle, *Champion*. Ponee, near Colombo, *Mr. W. Ferguson*.

†† *Perianthio* 8-partito. *Staminibus in flore masc.* 4.—*Hyalisma*, Champ. Miers.

3. **S. janthina**, Thw.—*Hyalisma janthina*, Champion, l. c. p. 466; Miers, l. c. p. 49.—C.P. 2664.

HAB. Damp forests in the south of the island.

*Obs.* In all the specimens of the genus *Sciaphila* which I have examined, the number of stamens in the male flower is half that of the lobes of the perianth, to each alternate lobe of which a stamen is opposed.

## CXLIII. CYCADEÆ.

## 1. CYCAS, Linn.

1. **C. circinalis**, Linn. Syst. Nat. quoad plant. Malab. et Zeylan. (syn. aliis excl.); Miq. Fl. Ind. Bat. ii. p. 1077, cum syn. (non *C. circinalis*, Roxb.).—*C. spherica*, Roxb.; Roxb. Fl. Ind. iii. p. 747; Todda-Panna, Rheed. Hort. Mal. iii. tt. 13–21.—C.P. 3689.

HAB. In scrubby hilly ground, up to an elevation of 1500 feet. Nom. vulg. “Maddoo-gass.”

The flour obtained from the seeds of this species is made into cakes and eaten by the Cinghalese, and is reputed a remedy for some disorders.

## MONOCOTYLEDONES.

## CXLIV. ORCHIDACEÆ.

Sect. 1. MALAXEÆ, Lindl.

Subsect. LIPARIDÆ, Lindl.

## 1. LIPARIS, Rich.

1. **L. atropurpurea**, Lindl. Gen. et Sp. Orch. p. 28.—*L. olivacea*, Wight, Ic. t. 903 (vix Lindl.). *L. Walkeriæ*, Wight, Ic. t. 905 (haud Grah.).—C.P. 3181.

HAB. More elevated parts of the Central Province.

2. **L. Walkeriæ**, Grah. in Bot. Mag. t. 3770 (haud Wight).—C.P. 2376.

HAB. Central Province, up to an elevation of 6000 feet.

Differs from the preceding in its smaller, more numerous flowers, and shorter pedicels; the labellum of the present plant is also more orbicular in outline and with deeper crenatures.

3. **L. Wightiana**, Thw.—*L. atropurpurea*, Wight, Ic. t. 904 (haud Lindl.).—C.P. 3179.

HAB. Central Province, at an elevation of 3000 to 5000 feet.

*Folia* 2-3, ovata, gradatim acuminata, basi cordata, parce obliqua. *Scapus* 3-12 poll., parum angulatus, pauciflorus. *Bracteæ* acutæ, suberectæ, pedicellis breviores. *Flores* viridi-purpurascens. *Sepala* oblonga, acuta. *Petala* linearia, marginibus revolutis. *Labellum* sepalis longius, planum, orbiculare vel suborbiculare, integrum vel minute retusum, 4 lin. in diam., basi callo lunato ornatum. *Columna* exalata.

4. **L. odorata**, Lindl. l. c. p. 26, cum syn.—*L. elata*, Lindl. l. c. p. 27, cum syn.—C.P. 3180, 3375.

HAB. Central Province, at an elevation of 2000 to 5000 feet.

An extremely protean plant, varying from 3 to 18 inches or more in height, with leaves from 2 to 5 in number, and ovate or sometimes narrow-lanceolate in form. The size of the flowers also varies, and these are yellow in colour, more or less tinged with purple. There cannot be a doubt, I think, of this being the plant figured by Rheede in Hort. Mal. vol. xii. t. 28; but I have no recollection as to whether the flowers are sweet-scented or otherwise.

5. **L. barbata**, Lindl. l. c.

HAB. Ceylon, *Macrae* (*Hb. Lindl.*).

I know nothing of this plant.

6. **L. longipes**, Lindl. l. c. p. 30; Wall. Pl. Asiat. Rar. t. 35; Wight, Ic. t. 906.—C.P. 3177.

HAB. Common up to an elevation of 3000 feet.

My specimens have much smaller flowers than those of the plant represented in Wallich's figure. The labellum is twice the length of the column, very much recurved, and of a pale green colour.

7. **L. viridiflora**, Lindl. l. c. p. 31, cum syn.—*L. elliptica*, Wight, Ic. t. 1735.—C.P. 3178.

HAB. Central Province, on rocks and trunks of trees, at an elevation of 3000 to 7000 feet.

Dr. Wight's figure well represents the present plant.

8. **L. angustifolia**, Lindl. l. c.—c.P. 2351.

HAB. Central Province, at an elevation of 2000 to 5000 feet.

*Folia* 2-4 poll. longa, 2-3 lin. lata, mucronata, in petiolum gradatim angustata. *Flores* minimi. *Capsula* ovata, 1½ lin. longa.

9. **L. gregaria**, Lindl. l. c. p. 33, cum syn.—c.P. 2359.

HAB. Hantani and elsewhere in the Central Province.

*Flores* sæpe rubro-tincti. *Labellum* basi bicallosum, longitudinaliter canaliculatum, transverse flexuosum, apice subacutum reflexum. *Anthera* 6-angula.

## 2. OBERONIA, Lindl.

1. **O. forcipata**, Lindl. Fol. Orch. No. 7.—c.P. 2511.

HAB. Hewahette District, on trees.

2. **O. longibracteata**, Lindl. Gen. Sp. Orch. p. 3; Fol. Orch. No. 8.—c.P. 543.

HAB. Common in the Hantani and Hewahette Districts.

*Flores* pallide rufescentes, *labello* rufo.

3. **O. truncata**, Lindl. Fol. Orch. No. 12.

HAB. Hantani, *Champion* (*Herb. Lindl.*).

I am not acquainted with this species.

4. **O. verticillata**, Wight, Ic. t. 1626.

Var.  $\beta$ ; *pubescens*, Lindl. Fol. Orch. No. 14.—c.P. 2516, 2572.

HAB. Kornegalle District.

5. **O. tenuis**, Lindl. l. c. No. 16.—c.P. 2654.

HAB. Hittawaka, on trees: March, 1853.

6. **O. Wightiana**, Lindl. in Bot. Reg. 1839, misc. 9; Fol. Orch. cum syn. No. 25; Wight, Ic. t. 1627.—*O. Arnottiana*, Wight, Ic. t. 1628.—c.P. 2506.

HAB. More elevated parts of the Central Province.

7. **O. Gardneriana**, Thw.; *bracteis* lanceolatis, acuminatis, integris, floribus subæquilongis; *sepalis* ovato-triangularibus, subacutis, integris; *petalis* oblongis, obtusis, denticulatis, parum reflexis.—c.P. 593.

HAB. Ambagamowa District, on trees, *Gardner*.

*Folia* lanceolata, mucronata, 1-3 poll. longa. *Spicæ* densifloræ, curvatæ, foliis longiores. *Flores* pallide rufescentes.

Closely allied to the preceding, with which I had confounded it; but the examination of recent specimens satisfies me that it is quite distinct.

8. **O. Scyllæ**, Lindl. Fol. Orch. No. 28.—c.P. 3124.

HAB. Matturatte, at an elevation of 5000 to 6000 feet: April, 1854.

## 3. MICROSTYLIS, Nutt.

1. **M. Rheedii**, Lindl. Gen. et Sp. Orch. p. 21, cum syn; Wight, Ic. t. 902.—*M. versicolor*, Lindl. l. c.; Wight, Ic. t. 901. An *M. luteola*, Wight, Ic. t. 1632?—c.P. 2375, 2743.

HAB. Common in shady forests in the Central Province.

Extremely variable in size and in the intensity of colour of the flowers and foliage.

I cannot doubt that the plant figured in Rheed. Hort. Mal. xii. t. 27 is the present species.

2. **M. discolor**, Lindl. p. 20; Wight, Ic. t. 1631.—C.P. 3698.

HAB. Damp forests in the Central Province.

This is very closely allied to the preceding, but the leaves are somewhat different in shape and more deeply tinged with purple; and, besides having the labellum edentulate, the sepals are relatively a good deal shorter than in *M. Rheedii*.

3. **M. lancifolia**, Thw.; cæspitosa; caule brevi; foliis lanceolatis vel lineari-lanceolatis, mucronatis; scapo elongato, pluri-bracteato; bracteis acuminatis, suberectis; floribus racemosis.—C.P. 2742.

HAB. On trunks of trees in the Saffragam District.

Folia 4-6,  $1\frac{1}{2}$ - $3\frac{1}{2}$  poll. longa, 2-4 lin. lata. Flores flavidi, quoad structuram iis præcedentis simillimi. Labellum concavum margine dentata, nec reflexa. Capsula oblonga, 6-costata, 3 lin. longa, pedicello 2 lin. longo.

4. **M. purpurea**, Lindl. l. c. p. 20.

HAB. Ceylon, *Macrae* (*Herb. Lindl.*).

I know nothing of this plant.

#### 4. DIENIA, Lindl.

1. **D. fusca**, Lindl. Gen. et Sp. Orch. p. 22.—C.P. 3182.

HAB. Central Province, *Macrae*.

Subsect. DENDROBIDÆ, Lindl.

#### 5. DENDROBIUM, Swartz.

1. **D. Macraei**, Lindl. Gen. et Sp. Orch. p. 75; Linn. Soc. Proc. iii. p. 6.—*D. nodosum*, Dalzell in Hook. Journ. of Bot. iv. (1852) p. 292.—C.P. 3695.

HAB. On trees, in the Central Province, not uncommon.

Flores sæpius bini, aliquando solitarii vel terni.

2. **D. crumenatum**, Swartz. (Lindl. Gen. et Sp. Orch. p. 88.)—C.P. 3696.

HAB. Common on the trunks of Cocoa-nut and other trees in the hotter parts of the island.

3. **D. aureum**, Lindl. l. c. p. 77; Wight, Ic. t. 1646.—*D. heterocarpum*, Wall. *D. rhombeum*, Lindl. in Bot. Reg. new ser. vol. xvii. t. 17.—C.P. 574.

HAB. More elevated parts of the Central Province.

4. **D. Maccarthiæ**, Thw. in Bot. Mag. t. 4886.—C.P. 3462.

HAB. Forest between Ratnapoora and Galle, flowering in May. Nom. vulg. "Wissak-mal."

Caules penduli, flagelliformes, teretes, striati, versus apicem foliosi,  $1\frac{1}{2}$ -2-pedales; nodis parce tumidis; internodis  $1\frac{1}{2}$ -2 poll. longis. Folia pauca, lanceolata,  $2\frac{1}{2}$ -3 poll. longa,  $\frac{3}{4}$ -1 poll. lata. Racemi laterales, 4-5-flori. Pedunculi basi bracteati. Pedicelli pallidi, 10-12 lin. longi. Flores pallide purpurei. Sepala lanceolata,  $2\frac{1}{2}$ -3 poll. lata. Petala subæquilonga, latiora. Labellum subtrilobum vel subtrapezi-forme, apice rotundatum, disco atro-purpureo. Columna albida, subquadrata. Mentum conicum, semiapertum, semipollicaris.

5. **D. macrostachyum**, Lindl. Gen. et Sp. Orch. p. 78; Wight, Ic. t. 1647.—C.P. 3183.



HAB. Common in the Central Province and Saffragam District.

6. **D. nutans**, Lindl. l. c. p. 90; Linn. Soc. Proc. iii. p. 17 in adnot.—*D. Jerdonianum*, Wight, Ic. t. 1644.—C.P. 2357.

HAB. Central Province, at an elevation of 4000 to 6000 feet.

A variable plant in the shape of the leaves and colour of the flowers, which latter are dark-ochraceous or sometimes nearly white.

7. **D. sanguinolentum**, Lindl. in Bot. Reg. new ser. vol. xvi. p. 62.

HAB. Ceylon, *Nightingale*.

I am not acquainted with this species.

8. **D. purpurascens**, Thw.; cæspitosum; *caulibus* brevissimis; *foliis* paucis, lineari-lanceolatis, apice angustatis, retusis, mucronatis; *racemis* foliis subæquilongis, 1-3-floris; *labello* sepalis petalisque latioribus lineari-lanceolatis æquilongo, indiviso, obtuso, versus apicem parum dilatato cymbiformique, minute crenulato; *mento* brevi, rotundato; *columna* sursum incrassata, margine utrinque breve 3-dentata.—C.P. 2352.

HAB. More elevated parts of the Central Province.

*Folia* 1-4 poll. longa, 2-6 lin. lata, sæpe purpurascens. *Flores* albidæ, semipollicares, *columna* atropurpurea. *Bracteæ* breves, rotundatæ.

9. **D. panduratum**, Lindl. in Linn. Soc. Proc. iii. p. 19.—An *D. graminifolium*, Wight, Ic. t. 1649?—C.P. 2353.

HAB. Central Province, up to an elevation of 6000 feet.

Cæspitosum. *Folia* lanceolata, 1-1½ poll. longa, *Flores* albidæ, purpureo-tincti, semipollicares.

## 6. BOLBOPHYLLUM, Thouars.

1. **B. elegans**, Gard. MS.; *scapo* unifloro; *sepalis* lateralibus in saccum oblongum, acuminatum, superne ad basin apertum cohærentibus, *dorsali* ovato, acuto, libero, duplo longioribus; *petalis* parvis, lanceolatis; *labello* valde carnosio, 3-lobo, *lobis* lateralibus brevibus, truncatis, erectis, *terminali* porrecto, linguæformi, apiculato, fere ad apicem fossulato.—C.P. 2350.

HAB. Hantani and Ambagamowa Districts, at an elevation of 3000 to 5000 feet.

*Rhizoma* repens. *Pseudobulbi* mediocres. *Folia* solitaria, lineari-lanceolata, utrinque gradatim angustata. *Flores* 1-1½ poll. longi, viridi-purpurascens, maculis purpureis plus minus conspersi, venis purpureis. *Labellum* flavum, purpureo-maculatum. *Columna* bicornis, flavida.

2. **B. petiolare**, Thw.; *pseudobulbis* parvis, confertis; *foliis* solitariis, ovatis, retusis, in petiolum longum angustatis; *floribus* capitatis; *sepalis* æqualibus, oblongis, acutis, quam petala plus quam duplo longioribus; *labello* obscure 3-lobo, late linguæformi, obtuso, convexo, medio profunde sulcato reflexoque.—C.P. 3184.

HAB. Ambagamowa, rare.

*Bracteæ* parvæ, rotundatæ, acutæ. *Capitula* circiter 4-flora. *Flores* 3 lin. longi, pallide ochracei, venis rubris.

3. **B. purpureum**, Thw.; *pseudobulbis* ovoideis, confertis; *foliis* solitariis, spathulato-ovatis, retusis; *floribus* capitatis; *sepalis* quam petala duplo longioribus; *lateralibus* in unum apice retusum cohærentibus,

*dorsali* libero; *labello* obscure 3-lobo, valde carnosus, medio profunde sulcato reflexoque, apice obtuso.—C.P. 3697.

HAB. Matolle, rare.

*Bracteæ* minutæ. *Capitula* 6-8-flora. *Flores* 1½-2 lin. longi, pallide purpurei, venis purpureis.

### 7. CIRRHOPETALUM, Lindl.

1. **C. Macraei**, Lindl. Gen. et Sp. Orch. p. 59 (haud Wight).—*C. Walkerianum*, Wight, Ic. t. 1657—C.P. 2362.

HAB. Hantani and Hewahette districts, at an elevation of 3000 to 4000 feet.

2. **C. grandiflorum**, Wight, Ic. t. 1656.—C.P. 3657.

HAB. Central Province, in the Maturatte and other districts, at an elevation of 3000 to 6000 feet.

3. **C. Wightii**, Thw.—*C. Macraei*, Wight, Ic. t. 1652 (haud Lindl.)—C.P. 3160, 2740 (632).

HAB. Central Province, at an elevation of 4000 to 7000 feet.

*Flores* albidii vel pallide flavi, plus minus rubro-maculati, vel omnino pallide virides. *Sepala* lateralibus linearibus vel lineari-lanceolata, obtusa, *dorsali* rotundato, apiculato 3-4-plo longiora. *Petala* parva, acuta, minute serrulata. *Labellum* carnosum, linguæforme, canaliculatum.

Very variable in size and in the colour of the flowers: or, possibly, further observation may show that I have confounded two distinct though closely allied species.

### 8. PHREATIA, Lindl.

1. **P. elegans**, Lindl. Gen. et Sp. Orch. p. 63.—C.P. 3186.

HAB. Central Province, *Macrae*.

### 9. ERIA, Lindl.

1. **E. bicolor**, Lindl. Gen. et Sp. Orch. p. 65 (haud in Linn. Soc. Proc. iii. p. 58).—C.P. 2982.

HAB. Central Province, at an elevation of 4000 to 7000 feet.

*Flores* albi, *columna* *labelloque* ad basin plus minus rubro-tinctis.

2. **E. Lindleyi**, Thw.—*E. bicolor*, Lindl. in Linn. Soc. Proc. iii. p. 58 (haud Gen. et Sp. Orch. p. 65).—*Dendrobium bicolor*, Lindl. l. c. p. 90.—C.P. 2761.

HAB. Central Province, common up to an elevation of 7000 feet.

3. **E. braccata**, Lindl. in Linn. Soc. Proc. iii. p. 46.—*E. reticosa*, Wight, Ic. t. 1637. *E. uniflora*, Dalzell in Hook. Journ. of Bot. iv. p. 111.—*Dendrobium braccatum*, Lindl. Gen. et Sp. Orch. p. 75.—C.P. 2356.

HAB. Central Province, not uncommon.

4. **E. muscicola**, Lindl. in Linn. Soc. Proc. iii. p. 47.—An *Dendrobium muscicola*, Lindl. Gen. et Sp. Orch. p. 75?—C.P. 2355.

HAB. Trunks of trees in the Central Province, at an elevation of 4000 to 7000 feet.

*Folia* acuta. *Pedunculi* 2-5-flori. *Labellum* acuminatum.

5. **E. velutina**, Thw.; molliter pubescens; *rhizomate* repente; cau-

libus brevibus, adscendentibus, 5-7-foliis; *foliis* anguste lanceolatis, coriaceis, subacutis; *pedunculis* terminalibus, solitariis vel 2-nis, paucifloris, folio longioribus; *sepalis* obtusis, *lateralibus* valde obliquis, *dorsali* lineari-lanceolato, cum petalis conformi: *labello* subcarnoso, linguæformi, recurvato, parum concavo, *lobis lateralibus* parvis, *terminali* rotundato, incrassato, papilloso.—C.P. 2349.

HAB. Dolosbage, in the Central Province, at an elevation of 3000 feet.

*Folia* 2-4 poll. longa. *Bracteæ* 3-4 lin. longæ. *Flores* parvi, flavi, siccitate rufescentes. *Columna* brevis, simplex. *Capsula* 6-costata.

## 10. ALWISIA, Lindl. Fol. Orchid.

1. **A. tenuis**, Lindl. l. c.—*Eria articulata*, Lindl. in Linn. Soc. Proc. iii. p. 47.—C.P. 2655.

HAB. Ambagamowa District, at no great elevation.

### Sect. 2. EPIDENDRÆ, Lindl.

#### Subsect. CÆLOGYNIDÆ, Lindl.

## 11. CÆLOGYNE, Lindl.

1. **C. breviscapa**, Lindl. Fol. Orch. No. 9.—*C. angustifolia*, Wight, Ic. t. 1641 (haud A. Rich.).—C.P. 3121.

HAB. Not uncommon in the more elevated parts of the Central Province.

2. **C. odoratissima**, Lindl. Gen. et Sp. Orch. p. 41; Fol. Orch. No. 10; Wight, Ic. t. 1640.—C.P. 518 (666, 667).

HAB. Newera Ellia and other elevated parts of the Central Province.

## 12. PHOLIDOTA, Lindl.

1. **P. imbricata**, Lindl. Gen. et Sp. Orch. p. 36; Wight, Ic. t. 907.—C.P. 3187.

HAB. Very common on the trunks of trees, up to an elevation of 3000 feet.

### Subsect. BLETIDÆ, Lindl.

## 13. PHAIUS, Lour.

1. **P. bicolor**, Lindl. Gen. et Sp. Orch. p. 128; Wight, Ic. tt. 1659, 1660.—C.P. 2368.

HAB. Central Province, at an elevation of 2000 to 4000 feet.

*Flores* 3-4 poll. in exp. *Sepala* petalisque flava, plus minus rubro-tincta.

2. **P. luridus**, Thw.; caulescens, *foliis* lanceolatis, acutis, glabris; *sepalis* cum *petalis* parum angustioribus lanceolatis, acutis; *labello* cucullato, rotundato, obscure trilobo, undulato, superne sparsim piloso et lineis duabus contiguis elevatis antice rugulosis ornato, subtus lævi, medio sulcato; *calcare* brevissimo; *columna* antice parum dilatata, parce pilosa, apice denticulata; *anthera* pilosa.—C.P. 613.

HAB. Rakwane, in the Saffragam District.

Terrestris. *Caulis* simplex, bipedalis. *Scapi* (in meis specim.) 2, foliis æquilongi, pluriflori. *Bracteæ* oblongæ, acutæ. *Flores* 2 poll. in exp. *Sepala petalæque* flavo-viridi-rufescentia. *Labellum* flavum, utrinque macula rubra irregulari-notatum.

#### 14. ARUNDINA, Blume.

1. **A. minor**, Lindl. Gen. et Sp. Orch. p. 125.—c.p. 485.

HAB. Not very uncommon amongst grass in swampy places in the Central Province, up to an elevation of 4000 feet.

#### 15. IPSEA, Lindl.

1. **I. speciosa**, Lindl. Gen. et Sp. Orch. p. 124 (Wight, Ic. t. 1663). c.p. 2364.

HAB. Not uncommon amongst long grass on exposed hillsides, at an elevation of 4000 to 5000 feet.

#### 16. APATURIA, Lindl.

1. **A. montana**, Lindl. Gen. et Sp. Orch. p. 131.—c.p. 3189.

HAB. Ambagamowa District, rare.

*Scapus* bipedalis, stramineus, plus minus purpureo-tinctus, basi squamis brevibus tectus. *Bracteæ* lineari-lanceolatæ, acutæ, reflexæ. *Flores* straminei, externe basi apiceque purpurei.

#### 17. ANIA, Lindl.

1. **A. bicornis**, Lindl. in Bot. Reg. xv. new ser. misc. p. 37.—*A. latifolia*, Wight, Ic. t. 914 (haud Lindl.); “*folio* oblongo-lanceolato, carnosio, *scapo* brevioris; *labelli* lobo medio emarginato, apiculato, basi lamellato, lamella tertia versus apicem interjecta; *anthera* bicorni,” Lindl. l. c.—c.p. 3190.

HAB. Hantani, in the Central Province, at an elevation of 3000 feet.

Tota planta plus minus purpureo-tincta. *Sepala petalæque* pallide flavo-viridia, purpurascentia.

2. **A. maculata**, Thw.; *folio* lanceolato, delicatule venuloso, longiuscule petiolato; *scapo* subæquilongo; *sepalis petalisque* subconformibus, lanceolatis, acutis; *labello* 3-lobo, disco trilamellato, *lobis lateralibus* parum brevioribus, erectis, obtusis, *intermedio* rotundato, subacuto, concavo; *calcare* brevi, cylindrico, truncato.—c.p. 3515.

HAB. A single specimen collected at Hapootelle, at an elevation of 5000 feet.

Glabra. *Pseudobulbus* conicus. *Scapus* bipedalis. *Bracteæ* anguste lanceolatæ, acutæ, flore breviores. *Sepala petalæque* pallide viridia, hæc seriebus linearibus 6, illa 4 macularum purpurearum longitudinaliter notata. *Labellum* album, interne versus basin purpureo-maculatum. *Columna* flava.

#### Sect. 3. VANDEE, Lindl.

#### Subsect. SARCANTHIDE, Lindl.

#### 18. EULOPHIA, R. Br.

1. **E. macrostachys**, Lindl. Gen. et Sp. Orch. p. 183 (Wight, Ic. tt. 1667, 1668).—c.p. 3188.

HAB. Not uncommon in the Central Province, up to an elevation of 4000 feet.

2. *E. virens*, R. Br. (Lindl. l. c. p. 182, cum syn. ; Wight, Ic. t. 913.)  
—C.P. 2369.

HAB. Central Province, not very common.

3. *E. grandiflora*, Lindl. l. c. p. 181.

HAB. Ceylon, *Macrae* (*Herb. Lindl.*).

I have not met with this species, nor is there a specimen of it in this herbarium.

4. *E. explanata*, Lindl. l. c. p. 180.

HAB. Ceylon, *Macrae* (*Herb. Lindl.*).

I am not acquainted with this species.

### 19. CYRTOPERA, Lindl.

1. *C. rufa*, Thw. ; *rhizomate* tuberoso ; *scapo* squamis distantibus circiter 3 vaginato ; bracteis anguste lanceolatis, ovario æquilongis ; *sepalis* æqualibus, oblongo-lanceolatis, acutis, quam *petala* obovata acuta parum longioribus ; *labello* cucullato, 3-lobo, interne venis pilosis, *lobis lateralibus* obtusis, *intermedio* rotundato, apiculato æquilongis ; *calcare* brevi, obtuso ; *columna* lineari, ad medium parum contracta.—C.P. 3566.

HAB. Hantani, at an elevation of 3000 feet.

*Scapus* bipedalis, rufescens. *Flores* rufi, 2 poll. in exp. *Capsula* oblonga, obtuse sexangula, 2½ poll. longa, 1 poll. in diam. *Folia* nondum vidi.

2. *C. Gardneri*, Thw. ; *acaulis* ; *rhizomate* tuberoso ; *foliis* anguste lanceolatis, plicatis ; *scapo* foliis æquilongis, 6-12-floro ; *sepalis* æqualibus, lanceolatis, acutis, quam *petala* oblonga longioribus ; *labello* parum saccato, 3-lobo, disco lineis pluribus elevatis ornato, *lobis lateralibus* latis, brevibus, rotundatis, erectis, *intermedio* reflexo, ovato, undulato, obtuso.—*C. laxiflora*, Gardn. MSS.—C.P. 2370.

HAB. Central Province, up to an elevation of 3000 feet.

*Scapus* bipedalis, rufescens. *Bractea* anguste lanceolata, acuta, ovario breviores. *Sepala* pollicaria, pallide viridia, plus minus rufescentia. *Petala* albida. *Labellum* albidum vel pallide flavum, disco rubro-tincto.

### 20. LUISIA, Gaud.

1. *L. tenuifolia*, Blume, Mus. Bot. Lugd. Bat. i. p. 64 ; Lindl. Fol. Orchid. No. 3, cum syn.—*Cymbidium tenuifolium*, Lindl. Gen. et Sp. Orch. p. 167.—An Wight, Ic. t. 1689 ?—C.P. 3530.

HAB. On trees in the Central Province, up to an elevation of 3000 feet.

*Flores* albidi, *sepalis* *petalis*que pallide purpurascens. *Labelli* *lobi* *laterales* parvi, interne purpurei ; *intermedius* oblongus, disco convexo marginibusque atropurpureis, postice lævis, antice convexus, 4-sulcatus, apice flavidus, plauus, truncatus, breve bilobus.

2. *L. Zeylanica*, Lindl. Fol. Orch. No. 7.—C.P. 2347.

HAB. Common on trees in the Central Province.

3. *L. bicaudata*, Thw. ; *caulescens* ; *foliis* membranaceis, lineari-lanceolatis, acutis, parum falcatis, apice breve bifidis, parce obliquis ; *racemis* oppositifoliis, folio longioribus ; *sepalis* ovatis, acutis, dorso ad apicem alato-carinatis ; *petalis* parum angustioribus, ecarinatis ; *labello*

inflato-cymbiformi, apice abrupte angustato, longe subulato-bifido.—  
C.P. 3494.

HAB. Ambagamowa and Saffragam Districts.

*Folia* 3-4 poll. longa,  $\frac{3}{4}$  poll. lata. *Flores*  $\frac{3}{4}$  poll. in exp., flavi, *labello* interne postice late carinato, antice versus marginem radiatim rubro-lineato.

## 21. COTTONIA, Wight, Ic. t. 1755.

1. **C. peduncularis**.—*C. macrostachya*, Wight, l. c.—*Vanda peduncularis*, Lindl. Gen. et Sp. Orch. p. 216.—C.P. 2361.

HAB. Warmer parts of the Central Province.

*Sepala* petalaeque reflexa, fusca, venis marginibusque flavis. *Labellum* atro-purpureum, margine pilosula fossulaque flavis, lobis lateralibus parvis adnatis auriculatum, oblongum, convexum, medio constrictum, prope basin rigide pilosum et tuberculo parvo stipitato piloso munitum, a medio versus apicem abrupte fossulatum. *Columna* erecta, apice triloba, lobis lateralibus minoribus gibbis, delicatule muricatis.

## 22. VANDA, R. Br.

1. **V. Roxburghii**, R. Br. in Bot. Reg. t. 506; Bot. Mag. t. 2245; Wight, Ic. t. 916.—C.P. 2346.

HAB. Hot, drier northern parts of the island.

2. **V. spathulata**, Spreng. (Lindl. Gen. et Sp. Orch. p. 216; Fol. Orchid.; Wight, Ic. t. 915.)—C.P. 2345.

## 23. ACAMPE, Lindl. Fol. Orchid.

1. **A. Wightiana**, Lindl. l. c.—*Vanda Wightiana?*, Wight, Ic. t. 1670. *V. fasciata*, Gardn. MSS.—C.P. 2342.

Var.  $\beta$ ; *foliis* latioribus; *pedunculis* elongatis, ramosis.—C.P. 3392.—An species distincta?

HAB. Jaffna, Gardner. Hantani.—Var.  $\beta$ . Ambagamowa and Hunasgiri Districts.

*Sepala* petalisque flava, interne transverse rubro-striata. *Labelli* lamina triangulari-ovata, parce transverse rubro-striata, convexa, subcarinata, basi tuberculata, (in meis specim.) edentata.

I suspect that var.  $\beta$  will prove to be a distinct species, but until I have an opportunity of examining fresh specimens of it I cannot determine satisfactorily.

2. **A. congesta**, Lindl. l. c.—*Vanda congesta*, Lindl. in Bot. Reg. 1839, Misc. n. 39.

HAB. Ceylon, Loddiges.

I know nothing of this species except from Dr. Lindley's description.

## 24. SACCOLABIUM, Blume.

1. **S. guttatum**, Lindl. Gen. et Sp. Orch. p. 220, cum syn.; Wight, Ic. tt. 1745, 1746.—C.P. 2344.

HAB. Hot, dry, northern parts of the island, on trees.

2. **S. curvifolium**, Lindl. l. c. p. 222.

HAB. Ceylon, Macrae (*Herb. Lindl.*).

With this species I am not acquainted, and there is neither specimen nor drawing of it in this collection.

3. **S. lineolatum**, Thw.; caulescens; *foliis* linearibus, canaliculatis, oblique retusis; *racemis* ramosis, multifloris; *sepalis petalisque* parum minoribus oblongis, obtusis; *labelli lamina* sagittato-triloba, lobis lateralibus acutis recurvis, intermedio majore, ovato, obtuso; *calcare* cylindrico.—C.P. 2741.

HAB. On trees in the Central Province, up to an elevation of 3000 feet.

*Folia* 6–8 poll. longa,  $\frac{3}{4}$  poll. lata. *Sepala petalaque* circiter 3 lin. longa, flava, interne delicatule transverse rubro-striata. *Labelli lamina* alba, sparsissime rubromaculata. *Calcar* flavum. *Capsula* cylindrica, 2 poll. longa, 3 lin. lata.

4. **S. niveum**, Lindl. l. c. p. 224.—C.P. 2340.

HAB. Central Province, not uncommon, at an elevation of 3000 to 4000 feet.

*Labelli lamina* carnososa, concava, obtusa. *Pollinia* aurantiaca. *Capsula* 2 lin. longa.

5. **S. acuminatum**, Thw.; caulescens; *foliis* carnosulis, lanceolatis, acuminatissimis; *racemis* lateralibus, brevibus; *sepalis petalisque* parum minoribus obovatis, obtusis; *labelli calcare* cylindrico, interne dente dorsali parvo bifido, *lamina* æquilonga, triloba, lobis lateralibus obtusis, intermedio ovato, concavo, obscure retuso.—C.P. 3376.

HAB. Saffragam District, at no great elevation.

*Folia* 4–5 poll. longa, 3 lin. lata. *Racemi* bipollicares. *Flores* 3 lin. in exp. *Sepala petalaque* pallide viridia, fulvo-marginata. *Labellum* albidum, lobis lateralibus rubris. *Capsula* reflexa, 8 lin. longa.

The dorsal tooth within the labellum of this species indicates an approach to *Cleisostoma*.

6. **S. gracile**, Lindl. l. c. p. 225.—C.P. 2528.

HAB. Hantani, in the Central Province, at an elevation of 3000 feet.

7. **S. roseum**, Lindl. l. c.—*Sarcanthus Walkerianus*, Wight, Ic. t. 1686.—C.P. 489.

HAB. Common upon trees in the Central Province, up to an elevation of 4000 feet.

8. **S. brevifolium**, Lindl. l. c.; Linn. Soc. Proc. iii. p. 35.—*S. virescens*, Gardn. MSS.—C.P. 488, 2341.

HAB. Common in forests in the Central Province. The variety with deep red flowers and more coriaceous leaves occurs at an elevation of 6000 to 8000 feet.

## 25. SCHŒNORCHIS, Blume.

1. **S. juncifolia**, Blume; Lindl. Gen. et Sp. Orch. p. 243.—*Sarcanthus filiformis*, Wight, Ic. t. 1684. *S. roseus*, Wight, Ic. t. 1685.—C.P. 633.

HAB. Trees in forests in the Central Province, at an elevation of 4000 to 6000 feet.

This would seem to be scarcely generically distinct from *Saccolabium*.

## 26. CLEISOSTOMA, Blume.

1. **C. maculosa**, Lindl. Gen. et Sp. Orch. p. 227.—C.P. 3193.

HAB. Central Province, not common.

*Folia* subtus præcipue versus basin, cum *pedunculis capsulisque*, creberrime purpureo-maculata. *Flores* flavidi, plus minus rubro-tincti.

2. **C. acaulis**, Lindl. l. c.—*Vanda fimbriata*, Gardn. MSS.—C.P. 3191.

HAB. Hantani, on trees, rare, Gardner.

*Labellum* saccatum, trilobum, lobis lateralibus brevibus, carnosus, erectis, parum acutis, antice convergentibus, terminali omnino externo, subtriangulari, fimbriato, reflexo.

3. **C. galeata**.—*Saccolabium galeatum*, Gardn. MSS.—Caulescens, *foliis* coriaceis, linearibus, obliquis, obtusis, emarginatis; *floribus* racemosis vel subpaniculatis, ad apicem pedunculi folio subæquilongi confertis; *sepalis petalisque* conformibus, obovato-spathulatis, obtusis; *labelli* saccati lobis lateralibus brevibus, truncatis, carnosus, terminali parvo, ovato, retuso, parce papilloso; *dente dorsali* triangulari, apice emarginato.—C.P. 2343.

HAB. Not uncommon on trees in the Central Province.

*Folia* 6 poll. longa,  $\frac{3}{4}$  poll. lata. *Flores* circiter 4 lin. in exp., flavi, rubro-maculati. *Capsulæ* 1 poll. longæ, 2 lin. latæ.

## 27. TÆNIOPHYLLUM, Blume.

1. **T. Alwisii**, Lindl. in Linn. Soc. Proc. iii. p. 42.—C.P. 3195.

HAB. Ambagamowa District. A single specimen detected upon the branch of a *Symplocos*, by Mr. H. De Alwis, the draftsman of the Peradenia Garden.

Minutum, acaule, aphyllum. *Radices* virides, carnosæ, repentes, complanatæ. *Scapus* erectus,  $\frac{1}{4}$  poll. longus. *Bracteæ* carinatæ. *Flores* spicati, pauci, minutissimi, pallide virides. *Sepala petalaque* conformia, cum labello in perianthium acute 6-dentatum connata. *Labellum* saccatum, lamina cymbiformi, marginibus liberis. *Columna* brevis. *Pollinia* 4, pyriformia, in glandulam sessilia. *Anthera* quadrata, subquadrilocularis.

## 28. OCTARRHENA, Thw. (gen. nov.)

*Sepala* æqualia, libera. *Petala* multo minora. *Labellum* petalis subconforme, parum cymbiforme. *Columna* brevissima, subhemisphærica. *Stigma* transverse oblongum. *Pollinia* 8 (4-4) libera, sphærica; *caudicula* lineari, medio bifurca; *glandula* parva, rotundata.

*Herba* parva, epiphyta, caulescens, cæspitosa; *foliis* distichis, subcylindricis, acutis; *racemis* axillaribus, paucifloris; *floribus* minutis, pallide viridibus.

1. **O. parvula**, Thw.—C.P. 3072.

HAB. Central Province, on forest trees, at an elevation of 3000 to 4000 feet.

## 29. AERIDES, Lour.

1. **A. Wightianum**, Lindl. Gen. et Sp. Orch. p. 238; Linn. Soc. Proc. iii. p. 40, cum syn; Bot. Mag. t. 5138.—*A. testaceum*, Lindl. Gen. et Sp. Orch. p. 238. *Vanda parviflora*, Lindl.; Wight. Ic. t. 1669.—C.P. 752.

HAB. Not uncommon upon trees, up to an elevation of 3000 feet.

2. **A. tessellatum**, Wight, (Lindl. l. c. p. 240?)—C.P. 3378.



HAB. Hunasgiria District; *Mr. H. De Alwis*.

*Sepala* pollicaria, lateralia obovata, dorsale *petalaeque* conformia angustiora, omnia pallide fusco-viridia, interne rubro-lineata maculataque. *Labelli* albidi calcar conicum; laminæ lobi laterales erecti, acuti; terminalis major, ovatus, retusus, flavo-tinctus, lineis rubris circiter 8 longitudinaliter notatus.

The above description is taken from a carefully executed drawing in this library: I have not seen the plant in a fresh state, nor are there specimens of it in the herbarium.

3. **A. cylindricum**, Lindl. l.c. (Wight, Ic. t. 1744.)—c.p. 2348.

HAB. On trees, not uncommon up to an elevation of 3000 feet.

*Flores* albidi. *Labelli* lobi laterales interne longitudinaliter delicatule purpureo-striati, intermedius disco luteus, 4-sulcatus, apice purpureo-maculatus.

### 30. DENDROCOLLA, Blume.

1. **D. serræformis**, Lindl. MSS. *Liparis serræformis*, Lindl. l. c. 33.—c.p. 3209.

HAB. Ambagamowa and Saffragam Districts, rare.

*Sepala petalaeque* lanceolata, acuta, hæc angustiora. *Columna* brevis, gibba. *Polinia* 4, 2 interiora multo minora. *Anthera* rotundata.

### 31. ŒCEOCLADES, Lindl.

1. **Œ. tenera**, Lindl. Gen. et Sp. Orch. p. 236 (Wight, Ic. t. 1683).—c.p. 2983.

HAB. Central Province, on trees, at an elevation of 3000 to 5000 feet.

*Sepala petalaeque* flava vel pallide viridia, sæpe venis rubris. *Labelli* laminæ lobi laterales truncati, parum rotundati crenulatique, petalis concolores, intermedius albus, calcar breve, cylindricum, dente interno dorsali truncato parum emarginato clausum.

### 32. DIPLOCENTRUM, Lindl.

1. **D. recurvum**, Lindl. Gen. et Sp. Orch. p. 218; Wight, Ic. t. 1680.—An *D. longifolium*, Wight, Ic. t. 1681.—c.p. 3192.

HAB. Central Province, *Gardner*.

### 33. ANGRÆCUM, Thouars.

1. **A. zeylanicum**, Lindl. in Linn. Soc. Proc. vol. iii. p. 40—c.p. 3693.

HAB. Narawelle, near Galle, *Champion*.

### 34. APPENDICULA, Blume.

1. **A. longifolia**, Blume, Bijdr. p. 304; Lindl. Gen. et Sp. Orch. p. 230<sup>p</sup>—c.p. 3208.

HAB. Ambagamowa District, rare.

*Folia* 3-5 poll. longa,  $\frac{1}{2}$  poll. lata, retusa, mucronata.

The present agrees very well with Blume's description, but in my specimens the flowers are not in a sufficiently perfect state for analysis.

### 35. PODOCHILUS, Blume.

1. **P. falcatus**, Lindl. Gen. et Sp. Orch. p. 234.—c.p. 2527.

HAB. Common in the Central Province, at an elevation of 3000 to 6000 feet.

2. *P. saxatilis*, Lindl. l.c. p. 235.—c.p. 3194.

HAB. Hantani and Ambagamowa Districts.

### 36. CYLINDROCHILUS, Thw. (gen. nov.)

*Sepala petalaeque subconformia, patentia. Labellum cylindrico-cucullatum, brevi-saccatum, erectum, columnæ basi producta articulatum, 3-lobum, lobis lateralibus oblongis, obliquis, intermedio brevissimo, truncato. Columna brevis, antice excavata. Pollinia 2, bipartibilia, rotundata, compressa, medio caudiculæ latæ brevis affixa; glandula parva, rotundata. Anthera rotundata, bilocularis.*—Herba epiphyta, caulescens, foliis coriaceis, lineari-oblongis, oblique retusis; pedunculis lateralibus; floribus paucis, capitatis; capsulis longiusculis, cylindricis.

1. *C. pulchellus*, Thw.—c.p. 2354.

HAB. Central Province and at Kornegalle, up to an elevation of 2000 feet.

*Flores albi, 7 lin. in exp. Labellum interne capitellato-pilosum, externe ad apicem maculis paucis fulvis cingulatum. Capsulæ 2½ poll. longæ, 1½ lin. latæ, 6-costatæ.*

### 37. JOSEPHIA, Wight.

1. *J. lanceolata*, Wight, Ic. t. 1742.—*Polystachya ramosa*, Gardn. MSS.—c.p. 2358.

HAB. Very abundant on trees in the Central Province.

Subsect. CRYPTOCHILIDÆ, Lindl.

### 38. ACANTHOPHIPPIUM, Blume.

1. *A. bicolor*, Lindl. in Bot. Reg. xx. t. 1730.—c.p. 2365.

HAB. Central Province, up to an elevation of 2000 feet.

*Pseudobulbi ovati. Folia sæpius 3, 9-18 poll. longa, 2½-4½ poll. lata, utrinque acuta. Scapus brevis, 2-7-florus. Bracteæ ovato-lanceolatæ, pollicares. Flores sesquipollicares, flavi, ad apicem plus minus rubro-tincti maculatique. Sepalorum apices liberi obtusi, reflexi. Petala angustiora. Labelli lobi laterales rotundati, incurvi; intermedius linguæformis, carnosus, obtusus, deflexus; discus 3-lamellatus; lamellis exterioribus tuberculatis, intermedia tenui, utrinque mucronata.*

Subsect. BRASSIDÆ, Lindl.

### 39. CYMBIDIUM, Swartz.

1. *C. ensifolium*, Swartz; Lindl. Gen. et Sp. Orch. p. 162, cum syn.—*C. hæmatodes*, Lindl. l.c.—c.p. 3694.

HAB. Hantani, in the Central Province.

The shape of the labellum of the Ceylon plant is so precisely similar to that of the variety of *C. ensifolium* figured in Bot. Reg. t. 1976, that I cannot doubt the plants being identical in species.

2. **C. bicolor**, Lindl. l.c. cum syn.—c.p. 3379.

HAB. Common on trees, up to an elevation of 3000 feet.

3. **C. aloifolium**, Swartz, (Lindl. l.c. cum syn.; Wight, Ic. tt. 1687, 1688).—c.p. 754.

HAB. Jaffna, *Gardner*.

Subsect. MAXILLARIDÆ, *Lindl.*

#### 40. POLYSTACHYA Hook.

1. **P. luteola**, Hook. Exot. Fl. t. 103; Lindl. Gen. et Sp. Orch. p. 73, cum syn.; Wight, Ic. t. 1678.—c.p. 2360.

HAB. Not uncommon in the warmer parts of the island, on trees.

Subsect. CALANTHIDÆ, *Lindl.*

#### 41. CALANTHE, R. Br.

1. **C. veratrifolia**, R. Br. (Lindl. Fol. Orchid. cum syn.)—c.p. 2367.  
Var.  $\beta$ . *discolor*.—c.p. 2371.

HAB. Not uncommon in the Central Province. Var.  $\beta$ . At an elevation of 6000 to 7000 feet.

2. **C. Masuca**, Lindl. Gen. et Sp. Orch. p. 249, cum syn.; Fol. Orchid. cum syn.—*C. purpurea*, Lindl. l.c. *C. emarginata*, Wight, Ic. t. 918.—c.p. 2366.

HAB. Ambagamowa District, at an elevation of 2000 to 4000 feet.

*Folia* utrinque molliter pubescentia. *Bractea* lanceolatae vel ovato-lanceolatae. *Labelli* discus tuberculatus; lobi laterales oblongi, obliqui; calcar pedicello æquilongum.

#### 42. GEODORUM, Jacks.

1. **G. dilatatum**, R. Br. (Lindl. Gen. et Sp. Orch. p. 175; Fol. Orchid. cum syn.; Wight, Ic. t. 912.)—c.p. 3196.

HAB. Central Province, amongst grass in shady places.

2. **G. fucatum**, Lindl. in Bot. Reg. t. 1687; Fol. Orch.

HAB. Ceylon, *Walker* (*Herb. Lindl.*).

I have never met with this species, nor is there a specimen or drawing of it in this collection.

Sect. 4. OPHREÆ, *Lindl.*

Subsect. SATYRIADÆ, *Lindl.*

#### 43. SATYRIUM, Swartz.

1. **S. Nepalense**, D. Don (Lindl. in Linn. Soc. Proc. iii. p. 44, cum syn. seq.; Wight, Ic. t. 929).—*S. Perottetianum*, A. Rich.; Wight, Ic. t. 1716. *S. albiflorum*, A. Rich.; Wight, Ic. t. 1717. *S. pallidum*, A. Rich.—c.p. 227 (522).

HAB. Abundant about Newera Ellia.

## Subsect. GYMNADENIDÆ, Lindl.

## 44. ATE, Lindl.

1. **A. virens**, Lindl. Gen. et Sp. Orch. p. 327 ; Wight, Ic. t. 928.—  
C.P. 3200.

HAB. Maturatte District, at an elevation of 4000 feet.

*Petala* oblata, obliqua, laciniis æqualibus.

2. **A. acuminata**, Thw. ; *foliis* anguste lanceolatis *bracteisque* rotundato-ovatis, setaceo-acuminatis ; *petalis* linearibus, obliquis lacinia anteriore longiore, anguste lineari ; *labello* trifido, lobis linearibus, obtusis, lateralibus divergentibus, longioribus ; *calcare* clavato, obtuso, ovario fere æquilongo.—C.P. 514.

HAB. Rambodde and above Galagama, at an elevation of 4000 to 5000 feet.

*Caulis* 1-3-pedalis, sparse foliosus. *Folia* 2-4 poll. longa. *Flores* secundi. *Sepala* rotundata *petalaque* pallide viridia. *Labellum* glabrum, fulvum.

This bears a considerable general resemblance to *Habenaria Heyneana*, Lindl., but is totally distinct from it.

## 45. HABENARIA, Willd.

1. **H. macrostachya**, Lindl. Gen. et Sp. Orch. p. 307.—C.P. 3197.

HAB. Hantani, Gardner.

*Sepala* longiuscule acuminata. *Petala* fere ad basin divaricato-bifida.

2. **H. dichopetala**, Thw. ; *foliis* ovatis vel ovato-lanceolatis, acutis ; *sepalis* lateralibus ovatis, acutis, dorsali rotundato, concavo ; *petalis* bipartitis, sinu lato, laciniis elongatis, setaceis, anteriore angustiore, parum longiore ; *labello* tripartito, laciniis elongatis, setaceis, æqualibus ; *calcare* cylindrico, parum clavato, subacuto, ovario æquilongo.—C.P. 3564.

HAB. Bintenne District, at no great elevation.

*Caulis* bipedalis, medio foliosus, dimidio inferiore squamatus. *Spica* pauciflora. *Sepala* circiter 3 lin. longa.

3. **H. dolichostachya**, Thw. ; *foliis* ovatis, gradatim acuminatis, acutis, basi rotundatis ; *spica* elongata ; *bracteis* ovato-acuminatis, floribus æquilongis ; *sepalis* lateralibus ovato-lanceolatis, acutis, dorsali parum brevior, rotundato, mucronato ; *petalis* subæquilongis, falcatis, bifidis, lacinia anteriore nana ; *labello* tripartito, laciniis linearibus angustissimis, lateralibus divergentibus, petalis æquilongis, *intermedia* parum longiore.—C.P. 3199.

HAB. Rambodde, Gardner.

*Caulis* 3-pedalis, medio foliosus, dimidio inferiore squamatus. *Folia* 2-3-poll. longa, superiora minora. *Spica* pedalis, laxiflora. *Sepala* circiter 3 lin. longa.

4. **H. viridiflora**, R. Br. (Lindl. Gen. et Sp. Orch. p. 319 ; Wight, Ic. t. 1705.)—C.P. 2372.

HAB. Common in the Central Province, at an elevation of 5000 to 7000 feet.

5. **H. pterocarpa**, Thw. ; *foliis* lanceolatis ; *floribus* majusculis ; *sepalis* lateralibus deflexis, oblique lanceolatis, acutis, dorsali parum brevior, rotundato, subacuto, *petala* æquilonga, falcato-linear, obtusa

includente; *labello* sepalis duplo longiore, tripartito, lobis linearibus, *lateralibus* acutis, parum longioribus, recurvis; *calcare* cylindrico, parum clavato, obtuso, quam ovarium 6-alatum fere triplo longiore.—C.P. 3201.

HAB. Rambodde, at an elevation of 4000 feet.

*Caulis* sesquipedalis, foliosus, *foliis* 2-6 poll. longis, superioribus gradatim minoribus. *Flores* albidi, *calcare* 3½ poll. longo, pallide viridi.

6. **H. latifolia**, Lindl. Gen. et Sp. Orch. p. 321.

HAB. Ceylon, *Macrae* (*Herb. Lindl.*).

I have not met with this species, nor are there specimens of it in this herbarium.

7. **H. plantaginea**, Lindl. l. c. p. 323, cum syn.; Wight, Ic. t. 1710.—C.P. 2374.

HAB. Not uncommon in the hot drier parts of the island.

8. **H. crinifera**, Lindl. l. c.; Wight, Ic. t. 926.—C.P. 207.

HAB. Central Province, up to an elevation of 3000 feet.

#### 46. PLATANThERA, Rich.

1. **P. cubitalis**, Lindl. Gen. et Sp. Orch. p. 292?—*Habenaria cubitalis*, R. Br. Prod. *Orchis cubitalis*, Linn.; Willd. iv. p. 17; Fl. Zeyl. n. 320.—C.P. 3202.

HAB. Pasdun Corle, at no great elevation.

*Petala* ovato-oblonga, obtusa.

There can be little doubt, I think, of this being the *Orchis cubitalis*, Linn., with the description of which in 'Flora Zeylanica' it agrees very well; but Lindley's plant would appear to be probably different, since he describes the petals as subulate.

2. **P. rhynchocarpa**, Thw.; *caule* dimidio inferiore, folioso; *foliis* lanceolatis; *spica* brevi, compacta; *bracteis* lanceolatis, acutissimis, ovario subæquilongis; *sepalis lateralibus* ovatis, subacutis, deflexis, *dorsali* parum brevioribus, rotundato, cucullato, *petala* lineari-spathulata, acuta, subæquilonga includente; *labello* sepalis lateralibus æquilongo, a medio tripartito, laciniis lineari-lanceolatis, acutis, lateralibus parum brevioribus; *calcare* cylindrico, obtuso, porrecto, quam ovarium longiuscule rostratum dimidio longiore.—C.P. 3058.

HAB. Above Galagama, at an elevation of 4000 feet.

*Caulis* 1-2-pedalis. *Folia* 2-6 poll. longa. *Flores* albi, *sepalis* 3-4 lin. longis, *calcare* 13 lin. longo, apice viridi.

#### 47. PERISTYLUS, Blume.

1. **P. goodyeroides**, Lindl. Gen. et Sp. Orch. p. 299.

HAB. Ceylon, *Walker* (*Herb. Hook.*).

2. **P. plantagineus**, Lindl. l. c. p. 3000; Wight, Ic. t. 921.

HAB. Ceylon, *Macrae* (*Herb. Lindl.*).

3. **P. spiralis**, Wight, Ic. t. 1696.—*P. Lawii*, Wight, Ic. t. 1695.—C.P. 226 (523).

HAB. Central Province, at an elevation of 4000 to 6000 feet.

4. **P. aristatus**, Lindl. l. c. p. 300.—*P. exilis*, Wight, Ic. t. 1698. *P. robustior*, Wight, Ic. t. 1699.—C.P. 3081, 2373.

HAB. More elevated parts of the Central Province, common.

5. **P. brevilobus**, Thw.; *caule* gracili, dimidio inferiore distanter folioso; *foliis* oblongo-lanceolatis, acutis; *spica* brevi, densiflora; *bracteis* ovato-lanceolatis, acutis, ovario subæquilongis; *sepalis* *petalisque* obtusis, his latioribus; *labello* obcordato, lobo intermedio minuto.—C.P. 3493.

HAB. Near Ratnapoora.

Subsect. CORYCIDÆ, Lindl.

#### 48. DISPERSIS, Swartz.

1. **D. tripetaloidea**, Lindl. Gen. et Sp. Orch. p. 371, cum syn.; Wight, Ic. t. 930.—C.P. 2363.

HAB. Not uncommon in damp forests of the Central Province, at an elevation of 3000 to 5000 feet.

*Flores* pallide purpurei.

Sect. 5. ABETHUSÆ, Lindl.

Subsect. GASTRODIDÆ, Lindl.

#### 49. GASTRODIA, R. Br.

1. **G. Javanica**, Lindl. Gen. et Sp. Orch. p. 385.—*Ephiphanes Javanica*, Blume, Bijdr. p. 421.—C.P. 3463.

HAB. Hinidoon Kanda, and Carawitta Kanda, near Ratnapoora.

This agrees so well with Blume's description, that I cannot doubt its being his plant.

#### 50. EPIPOGIUM, Gmel.

Lindl. in Linn. Soc. Proc. i. p. 176.

*Galera*, Blume. *Podanthera*, Wight. *Ceratopsis*, Lindl.

1. **E. nutans**, Lindl. l. c. p. 177.—*Galera nutans*, Blume, Bijdr. p. 415. *Podanthera pallida*, Wight, Ic. t. 1759.—C.P. 3205.

HAB. Not very uncommon, in damp shady places, up to an elevation of 2000 feet.

Tota planta albida, vel floribus bracteisque rubro-maculatis.

Subsect. VANILLIDÆ, Lindl.

#### 51. CYRTOSIA, Blume.

1. **C. Javanica**, Blume, Bijdr. p. 396; Rumphia, t. 69.—C.P. 3203.

HAB. Central Province, at the roots of Coffee-trees. Passbage, *Gardn.* Poondool Oya, *J. Nietner, Esq.*

#### 52. VANILLA, Plumier.

1. **V. Walkeriæ**, Wight, Ic. t. 932.—An *V. aphylla*, Wight, Ic. t. 931? (non Blume).—C.P. 2964.

HAB. Galle and Negumbo.

*Labellum* fauce rubescens.

I have unfortunately been unable to make a careful analysis of the flower of this beautiful plant, the only flower I have seen, and from which the draftsman made his drawing, having been inadvertently thrown away.

2. **V. Moonii**, Thw.; *foliis* carnosis, oblongis, acuminatis, subsessilibus; *spicis* brevibus, plurifloris; *bracteis* brevibus, acutis; *sepalis* *petalisque* conformibus, lanceolatis, conniventibus; *labello* cucullato, margine crenulato.—*V. aromatica?*, Moon's Cat. p. 60.—c.p. 3204.

HAB. Cultura, Moon. Negumbo and Kornegalle.

*Sepala* *petalaque* fulva, apice nigrescentia. *Labellum* pallide viride, disco albo, tribus seriebus apice convergentibus macularum fulvarum ornato. *Ovarium* albidum.

I have not seen the flowers of this plant in a fresh state. The description of their colour is taken from a drawing in this library.

### Sect. 6. NEOTTEÆ, Lindl.

#### Subsect. CRANICHIDÆ, Lindl.

### 53. ZOSTEROSTYLIS, Blume.

1. **Z. Zeylanica**, Lindl. Gen. et Sp. Orch. p. 446.—*Z. Walkerae*, Wight, Ic. t. 1748-4.—c.p. 383.

HAB. Not uncommon in shady forests of the Central Province, up to an elevation of 4000 feet.

*Folia* ovalia vel elliptica, subtrinervia, nitida, 3-7 poll. longa, longipetiolata. *Scapus* 1-3-pedalis. *Columna* brevissima, basi tuberculato-callosa. *Stigma* disciforme, emarginatum. *Clinandrium* latum, concavum, marginibus acutis nec membranaceis. *Anthera* disjunctim bilocularis. *Pollinia* basi attenuata curvataque; *glandula* communi parva, oblonga.

#### Subsect. SPIRANTHIDÆ, Lindl.

### 54. SPIRANTHES, L. C. Rich.

1. **S. australis**, Lindl. Linn. Soc. Proc. i. p. 178, cum syn.—*S. australis* et *densa*, Wight, Ic. t. 1724.—c.p. 550.

HAB. Common amongst grass, in the more elevated parts of the Central Province.

#### Subsect. PHYSURIDÆ, Lindl.

### 55. ZEUXINE, Lindl.

1. **Z. sulcata**, Lindl. Linn. Soc. Proc. i. p. 186, cum syn. seq.—*Z. emarginata*, Lindl. Gen. et Sp. Orch. p. 485. *Z. robusta*, Wight, Ic. t. 1726. *Z. brevifolia*, Wight, Ic. t. 1725.—c.p. 3017.

HAB. Amongst grass, in the warmer parts of the island.

56. **MONOCHILUS**, Wallich.

1. **M. longilabris**, Lindl. Linn. Soc. Proc. i. p. 186.—*M. affinis*, Wight, Ic. t. 1728.—c.P. 2377.

HAB. Hot drier parts of the island, not common.

2. **M. nervosus**, Wall. MSS. (Lindl. Gen. et Sp. Orch. p. 487.)

HAB. Ceylon, *Walker, Gardner (Herb. Hook.)*.

3. **M. flavus**, Wall. MSS.; Lindl. l. c.—c.P. 3120.

HAB. Raxawa and Ambagamowa, in the Central Province, at an elevation of 3000 feet.

4. **M. regius**, Lindl. l. c.—c.P. 2643.

HAB. Damp forests, in the Central Province, up to an elevation of 4000 feet. Nom. vulg. "Eeree-rajah."

57. **CHEIROSTYLIS**, Blume.

1. **C. parvifolia**, Lindl. Gen. et Sp. Orch. p. 488; Linn. Soc. Proc. i. p. 187.—c.P. 3071.

HAB. Ambagamowa District, rare.

2. **C. flabellata**, Wight, Ic. t. 1727; Lindl. l. c. cum syn. seq.—*Goodyera flabellata*, A. Rich.; Ann. des Sci. Nat. ser. 2, xv. p. 79, t. 12. *Zeuxine moniliformis*, Griff. Notulæ, iii. p. 397, t. 350.—c.P. 3377.

HAB. Ambagamowa District, rare.

Closely allied to the preceding: can it be merely a more highly developed form of it?

58. **RHAMPHIDIA**, Lindl.

Linn. Soc. Prod. i. p. 190.

1. **R. elongata**, Lindl.—*Goodyera* (sect. *Rhamphidia*) *elongata*, Lindl. Gen. et Sp. Orch. p. 494.—c.P. 2739.

HAB. Hewahette and Ambagamowa Districts, at an elevation of 4000 feet.

2. **R. Gardneri**, Thw.; subcaulescens; *foliis* oblongis, utrinque angustatis, glabris, scapo tomentello brevioribus; *spica* elongata, tenui, pluriflora; *bracteis* anguste lanceolatis, ovario subæquilongis; *sepalis* petalisque ovatis, obtusiusculis; *labello* parum brevioris, sacciformi, ovoideo, acuto, dimidio apicali clauso, interne parcissime brevi-piloso; *columna* labelli aperturæ apposita, subquadrata, apice spinis 2 uncinatis auriculata; *rostello* profunde bifido; *anthera* dorsali.—c.P. 3425.

HAB. Central Province, *Gardner*.

Bipedalis. *Folia* 3-5 poll. longa, 3-nervia, subtus purpurascens. *Sepala* petalaeque rufo-fuscescentia, 2 lin. longa. *Labellum* flavum.

59. **GOODYERA**, R. Br.

1. **G. procera**, Hook. Exot. Flor. t. 39; Lindl. Gen. et Sp. Orch. p. 493, cum syn.; Wight, Ic. t. 1729.—c.P. 597.

HAB. Common in damp forests, up to an elevation of 4000 feet.



2. **G. fumata**, Thw. ; subcaulescens, *foliis* majusculis, oblongis, utrinque angustatis, glabris, scapo brevioribus ; *spica* sparsiflora, elongata ; *rachi* glanduloso-pilosa ; *bracteis* lanceolatis, acutis, floribus subæquilongis ; *sepalis* oblongis, acutis ; *labello* concavo, abrupte in ligulam æquilongam arcte revolutam angustato.—c.p. 3668.

HAB. Central Province, rare.

*Folia* 5–10 poll. longa, concolora. *Rachis* rufo-fusca. *Bracteæ* rufæ. *Flores* viridi-rufescentes. *Labellum* albidum, fulvo-tinctum. *Stigma* obcordatum, rostratum, apice emarginatum.

## 60. GEORCHIS, Lindl.

1. **G. cordata**, Lindl. Gen. et Sp. Orch. p. 496.

HAB. Ceylon, *Macrae* (*Herb. Lindl.*).

I have not met with this species nor are there specimens of it in this herbarium.

## 61. ANÆCTOCHILUS, Blume.

1. **A. setaceus**, Blume, (Lindl. Gen. et Sp. Orch. p. 499 ; Wight, Ic. t. 1731.)—c.p. 384.

HAB. Damp forests, up to an elevation of 3000 feet. Nom. vulg. "Wanna-rajah."

## 62. PHYSURUS, L. C. Rich.

1. **P. Blumei**, Lindl. Gen. et Sp. Orch. p. 504 ; Linn. Soc. Proc. i. p. 181.—c.p. 598.

HAB. Not uncommon in the damp forests of the Central Province.

## 63. TROPIDIA, Lindl.

1. **T. curculigoides**, Lindl. Gen. et Sp. Orch. p. 497.—c.p. 3565.

HAB. Amongst high grass, in the hot, drier parts of the island.

## 64. CNEMIDIA, Lindl.

*Govindovia*, Wight.

1. **C. bambusifolia**, Thw. ; *foliis* oblongo-lanceolatis, acuminatis, basi acutis ; *ramis florigeris* brevibus, squamosis, rarius subfoliosis ; *spicis* brevissimis, confertifloris ; *bracteis* lanceolatis, acutissimis.—c.p. 3207.

HAB. Saffragam District.

*Caulis* 2–3-pedalis. *Folia* 4–9 poll. longa.

This would appear to be quite distinct from either of Lindley's species of this genus, as well as from the *Govindovia nervosa*, Wight, Ic. t. 2090, which latter has the leaves rounded at the base, and the spikes of flowers longer and less crowded.

## 65. CORYMBIS, Thouars.

1. **C. disticha**, Thouars, (Lindl. Fol. Orchid. cum syn.)—c.p. 3206.

HAB. Matturatte, *Moon*.

## CXLV. APOSTASIACEÆ.

## 1. APOSTASIA, Blume.

1. **A. Wallichii**, R. Br. in Wall. Pl. Asiat. Rar. i. p. 75, t. 84.—c.p. 2744.

HAB. Banks of streams, in the Saffragam District, at no great elevation.

## CLXVI. ZINGIBERACEÆ.

## 1. GLOBBA, Schieb.

1. **G. bulbifera**, Roxb. Fl. Ind. i. p. 78.—*G. marantinoides*, Wight, Ic. t. 2001 (*G. marantina*, Wight, non Willd.).—c.p. 3563.

HAB. Near Batticaloa.

My specimens agree extremely well with Dr. Wight's figure, except that in mine the superior wing-like appendages of the anther are twice as broad as the inferior ones. The flowers are of an orange-yellow colour.

## 2. ZINGIBER, Gærtn.

(Nom. vulg. "Wal-ingooroo.")

1. **Z. Zerumbet**, Rosc. (Roxb. Fl. Ind. i. p. 48, cum syn.; Wight, Ic. t. 2003.)—c.p. 3699.

HAB. Not uncommon up to an elevation of 2000 feet.

2. **Z. Cassumunar**, Roxb. l. c. p. 49.—Var. *subglabra*.—c.p. 3727.

HAB. Hantani, in the Central Province, rare.

This differs from the typical form of the species in being nearly smooth and in having smaller spikes of flowers.

3. **Z. Wightianum**, Thw.—*Z. squarrosum*, Roxb. ?; Wight, Ic. t. 2004 (vix Roxb.).—c.p. 2286.

HAB. Very common in forests, up to an elevation of 4000 feet.

1½–5-pedale. *Folia* 6–12 poll. longa, membranacea, superne subglabra, subtus villosula, vagina ad apicem tuberculis minutis conspersa, ligulæ bifidæ lobis 1½–3 lin. longis, rotundatis. *Flores* ochracei. *Corollæ lacinie exteriores* acuminato-lanceolatæ. *Labellum* brevius, purpureo-venosum maculatumque, 3-lobum, lobis lateralibus lanceolatis, acutis, intermedio majore, ovato, obtuso, bifido. *Capsulæ* pollicares.

In the present species the sheaths of the leaves have not the large conical processes at their apex, and the fruit is very much smaller than that of Roxburgh's *Z. squarrosum*, as described by him.

4. **Z. cylindricum**, Moon, Cat. p. 1.—*foliis* lanceolatis vel lineari-lanceolatis, utrinque acutis, subsessilibus, ad costam subtus aliquando pilosis; *vaginis* ad apicem tubulosis nec fissis; *ligula* brevi, truncata, externe pilosa; *spicis* radicalibus, pedunculatis, subcylindricis; *bracteis* imbricatis, rotundatis vel oblongis, subacutis; *labello* 3-lobo, lobis linearibus, truncatis, intermedio latiore.—c.p. 2287.

HAB. Very abundant in the forests of the Central Province.

3–6-pedale vel ultra. *Folia* 5–8 poll. longa, 1–2 poll. lata. *Pedunculi* 3–9

poll. longi, erecti, squamis pollicaribus obtusis ad medium tubulosis vaginati. *Bracteæ* pollicares, rufescentes vel pallidæ. *Flores* albidi. *Capsulæ* subsphæricæ, rubræ. *Semina* nigra, arillo albo.

This bears a very great external resemblance to *Z. officinale*, which is largely cultivated in native gardens; but, besides other differences, the tubular uncleft sheaths of the leaves at once distinguish the present plant.

### 3. CURCUMA, Linn.

1. **C. Zerumbet**, Roxb. Pl. Cor. iii. t. 201; Fl. Ind. i. p. 20, cum syn.—C.P. 3373.

HAB. Common in the Ambagamowa District. Nom. vulg. "Haran-kaha."

2. **C. aromatica**, Salisb. (Wight, Ic. t. 2005.)—*C. zedoaria*, Roxb. Fl. Ind. i. p. 23 (excl. syn.).—C.P. 3705.

HAB. Pusalawa and near Gampolle, in the Central Province. Nom. vulg. "Dada-kaha," "Wal-kaha."

The roots of this plant are used medicinally by the Cinghalese.

3. **C. albiflora**, Thw.; glabra, *foliis* ellipticis vel oblongis, utrinque acutis, petiolatis, concoloribus; *spica* radicali, pedunculo æquilonga; *bracteis* ovato-lanceolatis, parum reflexis, superioribus gradatim minoribus, omnibus florigeris; *perianthii* laciniis obtusis; *labello* rotundato, retuso.—C.P. 2737.

HAB. Maskellia, in the Ambagamowa District, at no great elevation.

*Folia* 6-8 poll. longa, 2½-3½ poll. lata. *Bracteæ* pallide virides. *Flores* albi, *labelli* disco luteo. *Antheræ* calcaria incurvata.

Mr. W. O. Brodie sent me some years ago, from Anooradhapoorra, a very small species of *Curcuma* (C.P. 3700), which appears to be undescribed, and the specimens are not sufficiently perfect to enable me to draw up a description.

*Curcuma longa*, Linn. (nom. vulg. "Kaha"), is cultivated in native gardens, and the roots are used in curries, as well as for medicine.

### 4. KÆMPFERIA, Linn.

1. **K. rotunda**, Linn. (Roxb. Fl. Ind. i. p. 16, cum syn.; Wight, Ic. t. 2029.)—C.P. 3175.

HAB. Warmer parts of the island. Nom. vulg. "Lonkenda."

The roots are employed medicinally by the Cinghalese.

2. **K. pandurata**, Roxb. l. c. p. 18, cum syn.—C.P. 3702.

HAB. Gangaroowa, near Peradenia. Nom. vulg. "Amba-kaha."

The Cinghalese use the roots of this medicinally. *K. Galanga*, Linn., is cultivated in native gardens as a medicine and a masticatory; but I have not heard of its occurring in a wild state in the island. Its native name is "Hingooroo-peyellee."

### 5. AMOMUM, Linn.

(Nom. vulg. "Neya-dandoo.")

#### Sect. 1.—*Floribus dense glomeratis.*

1. **A. echinatum**, Willd. Sp. Plant. i. p. 8, cum syn.—C.P. 3020.

HAB. Common in dense forests of the Central Province, up to an elevation of 4000 feet.

6-12-pedale. *Folia* firma, glabra, lineari-lanceolata, acuminata, sessilia, superne nigro-viridia, subtus pallidiora, 1-2-pedalia; *ligula* truncata vel parum rotundata, indivisa, 5-7 lin. longa. *Pedunculi* procumbentes, rubri, squamis obtusis muniti. *Capitulum* rotundatum vel ovoideum. *Bracteæ* oblongæ, obtusæ, pilosulæ, quam flores parum breviores. *Flores* flavi, rubro-tincti. *Labellum* parum carnosum, cucullatum, obscure trilobum, basi utrinque bidentatum, disco rubro-lineato maculatoque, lobo intermedio rotundato, aurantiaco. *Antheræ loculi* ciliati; *crista* transversim oblonga, breve triloba. *Capsulæ* rotundatæ, nitidæ, uncinato-aculeatæ, nigro-purpureæ, pollicares.

**2. *A. fulviceps*, Thw.;** *foliis* firmis, lanceolatis vel lineari-lanceolatis, utrinque acutis, acuminatis, sessilibus; *ligula* indivisa, rotundata, tam longa quam lata; *pedunculis* brevibus, procumbentibus; *capitulo* rotundato vel ovoideo, plurifloro; *bracteis* imbricatis, ovato-oblongis, retusis, adpressim dense fulvo-pilosis.—*a. foliis* glabris.—c.p. 3122. *β. foliis* subtus pilosis.—c.p. 3704.

**HAB.** Dense forests. **Var. *a.*** Raxawa in the Central Province. **Var. *β.*** Reigam Corle.

5-8-pedale vel ultra. *Folia* 1-2-pedalia. *Flores* pallide flavi, rubro-tincti. *Corollæ* laciniæ exteriores oblongæ, obtusæ, externe dense pilosæ, posterior subduplo latior. *Labellum* rotundatum, retusum, trilobum, basi bidentatum, dentibus longiusculis. *Antheræ loculi* pilosi, *crista* semilunaris, emarginata.

**3. *A. pterocarpum*, Thw.;** glabrum, *foliis* oblongo-lanceolatis, utrinque acutis acuminatis, sessilibus; *ligulæ* bifidæ lobis longis, lanceolatis, acutis, scariosis; *pedunculis* brevibus, procumbentibus; *capitulo* ovoideo vel globoso; *bracteis* oblongis, majusculis, cito deciduis; *capsulis* confertis, ovoideis, parum acutis, 9-costatis, costis alatis crenulatisque.—c.p. 3021.

**HAB.** Forests of the Central Province, up to an elevation of 4000 feet.

3-6-pedale vel ultra. *Folia* 1-2-pedalia. *Flores* albi. *Corollæ* laciniæ exteriores oblongæ, obtusæ, posterior latior. *Labellum* rotundatum, basi breve bidentatum, disco flavo, rubro-tincto. *Antheræ* *crista* brevis, triloba. *Capsulæ* sesquipollicares, virides.

**4. *A. masticatorium*, Thw.;** *foliis* firmis, glabris, lineari-lanceolatis; acuminatis, sessilibus, basi acutis vel parum rotundatis; *ligula* longiuscula, indivisa, lanceolata, superne scariosa; *pedunculis* procumbentibus; *capitulo* globoso; *bracteis* obovatis, obtusis, pilosulis, ciliatis, quam flores dimidio brevioribus; *capsulis* retundatis, echinatis.—c.p. 3701.

**HAB.** Common in forests of the Central Province, up to an elevation of 4000 feet.

2-8-pedale. *Folia* 5-15 poll. longa. *Bractæ* 8 lin. longæ. *Flores* flavi. *Corollæ* tubus externe villosulus, laciniæ exteriores oblongæ, obtusæ, posterior duplo latior. *Labellum* rotundatum, rubro-punctulatum, basi bidentatum, trilobum, lobo intermedio minore, bifido. *Anthera* glabra, loculis longiusculis, ad apicem valde divergentibus, *crista* semilunaria. *Capsulæ* 8 lin. in diam. nigro-virides.

The Cinghalese chew the aromatic rhizomes of this plant with their betel.

**5. *A. acuminatum*, Thw.;** *foliis* firmis, glabris, oblongo-lanceolatis, longiuscule acuminatis, basi parce acutis, petiolatis; *ligula* indivisa, rotundata, parum brevior quam lata; *pedunculis* procumbentibus; *capitulo* subgloboso, parvo; *bracteis* rotundatis vel obovatis, obtusis, puberulis, ciliatis, quam flores dimidio brevioribus; *labello* rotundato, trilobo, lobo intermedio emarginato; *antheræ* loculis ciliatis, *crista* semilunari crenulata.—c.p. 3466.

HAB. Ehualagodde, near Ratnapoora, at no great elevation.  
2-4-pedale. *Folia* (in meis specim.) circiter 6 poll. longa, 1-1½ poll. lata, petiolo 4 lin. longo. *Rhizomata* repentia, cum *pedunculis* brevibus, *squamis* rotundatis *bracteis*que rufescentia. *Flores* lutei, rubro-tincti.

Sect. 2.—*Floribus laxe spicatis vel paucis.*

6. **A. pulchellum**, Thw.; *foliis* oblongis, utrinque acutis, membranaceis, minute reticulatis, præter costam subtus pilosam glabris, petiolatis; *ligulæ* bifidæ lobis brevibus, rotundatis; *pedunculis* elongatis, procumbentibus, ramosis, *squamis* oblongis obtusis subglabris amplectentibus tectis; *floribus* quam bracteæ multo longioribus; *antheræ* crista majuscula, semicirculari; *capsulis* rotundatis, leviter 9-costatis.—C.P. 2736.

HAB. Forests of the Central Province, up to an elevation of 3000 feet.

1-2-pedale. *Folia* ½-2 ped. longa, 2½-5 poll. lata, *petiolo* ½-8 poll. longo. *Corollæ* laciniæ exteriores puberulæ, inter se subæquales, lineares, obtusæ, pallidæ, venis 3 fulvis. *Labellum* rotundatum, roseo-purpureum, disco flavo, 3-lobum, lobo intermedio retuso. *Antheræ* loculi glabri, ad apicem vix divergentes, *crista* roseo-purpurea. *Capsulæ* 6 lin. in diam., rubræ.

7. **A. hypoleucum**, Thw.; *foliis* oblongo-lanceolatis, utrinque acutis, acuminatis, superne glabris nitidis, subtus adpresse canescenti-pilosis, petiolatis; *ligula* brevi, obtusa, apice biloba; *scapis* secus rhizoma repens elongatum sparsis, brevibus, 3-floris; *capsulis* rotundatis, 9-costatis.—C.P. 3532.

HAB. Damp forests of the Central Province, up to an elevation of 4000 feet.

2-5-pedale. *Folia* ½-2-pedalia, petiolo 1-3 poll. longo. *Bracteæ* ovatæ, retusæ, mucronulatæ, pallidæ, glabræ, floribus multo breviores. *Corollæ* laciniæ exteriores albidæ, inter se subæquales, oblongæ, obtusæ, sæpe mucronatæ. *Labellum* rotundatum, retusum, disco flavo rubro-tincto. *Antheræ* loculi glabri, ad apicem vix divergentes, *crista* semilunari. *Capsulæ* rubræ, pollicares.

## 6. ELETARIA, Rheede.

### Sect. 1.—*Spicis laxifloris.*

1. **E. Cardamomum**, White, (Blume, Enum. i. p. 51; Miq. Flor. Ind. Bat. iii. p. 600, cum syn.)

Var. β; *foliis* parum latioribus firmioribusque; *capsulis* fusiformibus.—*E. major*, Smith.—An *Elettari*, Rheed. Hort. Mal. xi. t. 6?—C.P. 2431.

HAB. Var. β. Forests in the Central and Southern Provinces, up to an elevation of 3000 feet. Nom. vulg. "Ensal." Var. α occurs in Ceylon only as a cultivated plant; it is called by the Cinghalese Rata-ensal.

—A careful comparison of growing specimens satisfies me that the plants producing respectively the round and the long cardamoms of commerce are not distinct species, but merely varieties of the present species. In every essential particular the structure is similar in the two plants, the only difference being that var. α, which produces the round cardamom, is a little taller, with rather narrower and less firm leaves, and that its fruit is more aromatic as well as different in form from that of var. β. The seeds of both varieties are used by the Cinghalese to chew with their betel, and as medicines.

Sect. 2.—*Floribus dense glomeratis.**Geanthus*, Rot.

2. **E. involucrata**, Thw.; *foliis* majusculis, membranaceis, glabris, lineari-lanceolatis, utrinque acutis, acuminatis, sessilibus; *ligula* brevi, indivisa, rotundata; *pedunculis* erectis, longiusculis, squamosis; *capitulo* multifloro, ultra apicem truncatum explanatum squamis magnis involucrato.—C.P. 3019.

HAB. Central Province, in damp forests, at an elevation of 4000 to 6000 feet.

6-10-pedalis vel ultra. *Folia* 1½-3 ped. longa, 4-7 poll. lata. *Pedunculi* 1½-3 ped. alti, cum squamis oblongis vel ovato-oblongis obtusis 3-6 poll. longis bracteisque obovatis, rubri. *Flores* parvi, pallide ochracei. *Corollæ* laciniæ exteriores obovato-oblongæ, obtusæ. *Labellum* obovatum, 3-lobum, lobis subæqualibus rotundatis.

3. **E. floribunda**, Thw.; *foliis* glabris, firmis, lineari-lanceolatis, utrinque acutis, acuminatis, petiolatis; *ligula* brevi, indivisa, rotundata; *pedunculis* erectis, longiusculis, squamosis; *capitulo* multifloro, globoso; *bracteis* rotundatis vel ovato-rotundatis, retusis, mucronatis; *floribus* parvis, parum carnosis; *corollæ* laciniis exterioribus ovatis, obtusis, concavis; *labello* obovato, retuso, obscure trilobo.—C.P. 3374.

HAB. Forests in the Ambagamowa District.

3-6-pedalis vel ultra. *Folia* 1½-2 ped. longa, circiter 3 poll. lata. *Flores* pallide ochracei.

4. **E. nemoralis**, Thw.; *foliis* glabris, firmis, lanceolatis vel lineari-lanceolatis, utrinque acutis, acuminatis, brevi-petiolatis; *ligula* brevi, indivisa, rotundata; *pedunculis* procumbentibus, squamosis; *capitulo* parvo, subgloboso; *bracteis* glabris, valde membranaceis, exterioribus rotundatis vel obovatis, interioribus lanceolatis; *corollæ* laciniis exterioribus æqualibus, oblongis, obtusis, tubo longo; *labello* rotundato, 3-lobo, lobo intermedio minore, bifido; *ovario* piloso.—C.P. 3703.

HAB. Forests in the Reigam and Pasdoon Corles.

2-4-pedalis. *Folia* 8-15 poll. longa, circiter 2 poll. lata. *Flores* flavi, labello copiose purpureo-venoso. *Capsulæ* rotundatæ, breve rostratæ, semipollicares.

## 7. HEDYCHIUM, Kœnig.

1. **H. coronarium**, Kœn. (Wallich in Hook. Journ. of Bot. iii. (1853) p. 325, cum syn; Wight, Ic. t. 2010.)—C.P. 3018.

Var.  $\beta$ ; *floribus* lutescentibus.—Wall. l. c.; Wight, Ic. tt. 2008, 2009.—C.P. 3728.

HAB. Margins of woods, in the Central Province, up to an elevation of 6000 feet. Nom. vulg. "Ella-wal."

2. **H. angustifolium**, Roxb. Flor. Ind. i. p. 13; Wall. l. c. p. 371.—C.P. 3669.

HAB. In similar situations with the preceding.

## 8. ALPINIA, Linn.

1. **A. Galanga**, Linn. (Roxb. Fl. Ind. i. p. 59, cum syn.)—C.P. 3729.

HAB. Cultura, Moon. Nom. vulg. "Koloowala."

The Cinghalese employ the rhizomes medicinally.

2. **A. Allughas**, Roscoe, (Roxb. l. c. p. 61, cum syn.—C.P. 3157.)

HAB. Warmer parts of the island. Nom. vulg. "Aloo-gass."

The rhizomes of this species are also used medicinally by the Cinghalese.

3. **A. nutans**, Roscoe, (Roxb. l. c. p. 65, cum syn; Wight, Ic. t. 2027.)

Var.  $\beta$ ; *panicula* brevior; *floribus* minoribus.—*A. sericea*, Moon, Cat. p. 1.—C.P. 3312.

HAB. In a forest between Negombo and Kornegalle. Nom. vulg. "Rankeeriya."

Besides the distinguishing characters mentioned above, this differs from the common form of *A. nutans* in the flowers not being tipped with rose-colour.

4. **A. calcarata**, Roscoe, Roxb. l. c. p. 69; Wight, Ic. t. 2028.)—C.P. 3730.

HAB. Kandy, Moon, Walker. Nom. vulg. "Kattakeeriya."

I have never met with this truly wild. It grows abundantly in the Botanic Garden.

## 9. COSTUS, Linn.

1. **C. speciosus**, Smith, (Roxb. Fl. Ind. i. p. 58, cum syn.; Wight, Ic. t. 2014.)—C.P. 3082.

HAB. Margins of woods, in the Ambagamowa District. Nom. vulg. "Taiboo-gass."

The Cinghalese use the rhizomes as a medicine.

## CXLVII. MARANTACEÆ.

### 1. MARANTA, Plum.

1. **M. virgata**, Dietr. (Steud. Nom.; Wight, Ic. t. 2015.)—*Phrynium virgatum*, Roxb. Fl. Ind. ed. Carey, i. p. 4.—C.P. 3465.

HAB. Hotter parts of the island, not very uncommon. Nom. vulg. "Hoolankeeriya."

The Rev. S. O. Glenie, of Trincomalee, informs me that arrowroot of a very superior quality is obtainable from the rhizomes of this plant.

2. **M. spicata**.—*M. cæspitosa*, Dietr.; Steud. Nom. *Phrynium spicatum*, Roxb. l. c. p. 5.—C.P. 612.

HAB. Forests in the Central Province; rather rare.

This, after a careful analysis of the flower, appears to me to be a true species of *Maranta*.

### 2. PHRYNIUM, Willd.

1. **P. capitatum**, Linn. (Roxb. Fl. Ind. i. p. 8, cum syn.; Wight, Ic. t. 2016.)—C.P. 3123.

HAB. Forests of the Central Province. Nom. vulg. "Atbamikeeriya."

### 3. CANNA, Linn.

1. **C. Indica**, Linn. (Roxb. Fl. Ind. i. p. 1, cum syn.)—C.P. 3706.

HAB. Very abundant, both the red- and yellow-flowered varieties. Nom. vulg. "Bootsarana."

## CXLVIII. MUSACEÆ.

## 1. MUSA, Schreb.

1. **M. sapientum**, Linn. (Willd. Sp. Pl. iv. p. 894; Roxb. Fl. Ind. i. p. 663, cum syn.)—C.P. 3731.

HAB. Rocky woods in the Central Province, common. Nom. vulg. "Wal-kaihel-gass."

There cannot be a doubt that this is the species from which have originated the numerous varieties of sweet Plantains cultivated here.

## CXLIX. PONTEDERACEÆ.

## 1. MONOCHORIA, Presl.

(Nom. vulg. "Deya-habarella.")

1. **M. hastæfolia**, Presl, (Kunth, Enum. iv. p. 133, cum syn.)—*Pontederia hastata*, Linn.; Roxb. Fl. Ind. ii. p. 121.—C.P. 3213.

HAB. Common in swampy places in the hotter parts of the island.

This and the following species are used medicinally by the Cinghalese.

2. **M. vaginalis**, Presl, (Kunth, l. c. p. 134, cum syn.)—*Pontederia vaginalis*, Burm.; Roxb. Cor. Pl. ii. p. 110; Fl. Ind. l. c.—C.P. 2315.

HAB. Equally common with the preceding, and in similar places.

3. **M. plantaginea**, Kunth, l. c. p. 135.—*Pontederia plantaginea*, Roxb. Fl. Ind. ii. p. 123.—C.P. 3686.

HAB. Hotter parts of the island; less common than the two preceding species.

## CL. COMMELYNACEÆ.

## 1. COMMELYNA, Dill.

Sect. 1.—*Spatha complicata*.

1. **C. salicifolia**, Roxb. Fl. Ind. i. p. 172; Kth. Enum. iv. 39, cum syn.—An *C. communis*, Linn. forma?—C.P. 2979, 3358.

Var.  $\beta$ ; *angustata*; *foliis* longis, linearibus; *spathis* longe pedunculatis, subfalcatis, valde acuminatis.—C.P. 3215.

HAB. Very abundant. Var.  $\beta$ . Doombera District. Nom. vulg. "Girrapata."

Var.  $\beta$ , though differing a good deal in general appearance from the present plant as it ordinarily occurs, is notwithstanding, I suspect, merely an extreme form of it.

I do not remember having met with the typical *C. communis*, L., which has the smaller petal very narrow, whilst in the present plant I find the smaller petal varying from wide-cordate to oval.

Sect. 2.—*Spatha turbinato-cucullata*.

2. **C. Benghalensis**, Linn. (Kth. l. c. p. 50, cum syn.; Wight, Ic. t. 2065.)—C.P. 2314.



HAB. Very common in the warmer parts of the island. Nom. vulg. "Deya-mainaireya."

Used as a medicine by the Cinghalese.

3. *C. obliqua*, Don, (Prod. Fl. Nep. p. 45 ; an Vahl, Enum. ii. p. 173 ?) — *C. Donii*, Dietr. ; Kth. l. c. p. 58. *C. paludosa*, Blume, Enum. i. p. 2 ; Kth. l. c. p. 51 ; Miquel, Fl. Ind. Bat. iii. p. 533.—c.p. 2326.

HAB. North of Kandy, not very uncommon.

4. *C. longifolia*, Lam. (Vahl, Enum. ii. p. 165 ?)—c.p. 3224.

HAB. Hot drier parts of the island.

Subglabra vel plus minus pilosa. *Folia* anguste lanceolata, valde acuminata, parce obliqua, 2-6 poll. longa,  $\frac{1}{4}$ -1 poll. lata, vagina ad oram pilis rigidulis albis ciliata. *Spathæ* 1-3-næ, parum falcatae acuminatæque, externe pilosæ, singulæ pedunculum solitarium 3-florum includentes.

Nearly allied to the preceding species, from which it differs in its smaller size, and in its narrower, more acuminated, and less oblique leaves. It might perhaps be safe to consider it a variety only.

## 2. ANEILEMA, R. Br.

(Nom. vulg. "Tanna-patā.")

1. *A. ensifolium*, Wight, Ic. t. 2074.—*A. secunda*, Wight, Ic. t. 2075-1.—c.p. 2324.

HAB. Not uncommon in the south of the island.

2. *A. scapiflorum*, Wight, Ic. t. 2073.

Var.  $\beta$  ; minus ; foliis multo minoribus ; bracteis parum brevioribus latioribusque.—c.p. 3314.

HAB. Ceylon ; Mrs. Walker, in *Herb. Hook.* Var.  $\beta$  ; Oova District, at an elevation of 5000 feet.

Var.  $\beta$  may possibly prove to be a distinct species, but, without better materials than I at present possess, it is not in my power to determine satisfactorily. It appears to agree in many essential particulars with specimens of *A. scapiflorum*, from Khasia, collected by Drs. Hooker and Thomson.

3. *A. protensum*, Wall. Cat. No. 5218.—*Dictyospermum protensum*, Wight, Ic. t. 2071.—c.p. 3026.

HAB. Kokool-Corle.

4. *A. montanum*, Wight, in Wall. Cat.—*Dictyospermum montanum*, Wight, Ic. t. 2069.—c.p. 2325.

HAB. Not uncommon up to an elevation of 3000 feet.

5. *A. vaginatum*, R. Br. (Kth. Enum. p. 67, cum syn. ; Wight, Ic. t. 2076-1.)—c.p. 2328.

HAB. Common in the hotter parts of the island.

6. *A. nanum*, Kth. Enum. iv. p. 65, cum syn. ; Wight, Ic. t. 2077-2.—*Commelyna nana*, Roxb. Fl. Ind. i. p. 173.—c.p. 2329.

HAB. Very abundant in damp pastures.

7. *A. nudiflorum*, R. Br. (Kth. l. c. p. 66, cum syn. ; Miquel, Fl. Ind. Bat. iii. p. 537.)—c.p. 3027, 3313.

HAB. Common in the warmer parts of the island.

8. *A. dimorphum*, Dalz. in Hook. Journ. of Bot. iii. (1851) p. 138 ;

Bombay Flor. p. 254.—*A. paniculatum*, Wight, Ic. t. 2075-2.—C.P. 3025.

HAB. Common in the Central Province.

*Bracteæ* minutæ, rotundatæ, concavæ, apice incrassatæ, coloratæ, incurvæ. *Folia* glabra vel pilosula.

### 3. FLOSCOPA, Lour.

*Dithyrocarpus*, Kunth.

1. *F. paniculata*, Hassk. (Miq. Fl. Ind. Bat. iii. p. 542; Benth. Fl. Hongk. p. 377, cum syn.—*Dithyrocarpus petiolatus*, *D. Rothii*, et *D. undulatus*, Wight, Ic. tt. 2079, 2080.)—C.P. 553.

HAB. Very common up to an elevation of 4000 feet.

### 4. POLLIA, Thunb.

*Aclisia*, E. Meyer.

1. *P. Indica*.—*Aclisia Indica*, Wight, Ic. t. 2068.—C.P. 2327.

HAB. Not uncommon in the warmer parts of the island.

### 5. CYANOTIS, Don.

(Nom. vulg. "Goloo-bailoo.")

1. *C. axillaris*, R. et Sch. (Kth. Enum. iv. p. 105, cum syn.)—C.P. 2330.

HAB. Hotter parts of the island.

2. *C. cristata*, R. et Sch. (Kth. l. c. p. 102, cum syn.; Wight, Ic. t. 2082.)—C.P. 3217.

HAB. Common up to an elevation of 3000 feet.

3. *C. pilosa*, R. et Sch. (Kth. l. c. p. 105; Wight, Ic. t. 2083.)—C.P. 2331.

HAB. Newera Ellia and other elevated parts of the island.

4. *C. fasciculata*, R. et Sch. (Kth. l. c. p. 104, cum syn.; Wight, Ic. t. 2086-2.)—C.P. 2433 (2978).

HAB. Very abundant in the Central Province.

5. *C. lanceolata*, Wight, Ic. t. 2085.—C.P. 2332, 3223. Var. *subglabra*.—C.P. 3216.

HAB. Central Province, up to an elevation of 6000 feet.

Closely allied to the preceding species, but distinguished by its much larger size and by the different character of the pubescence, which is silky in this, but lanose in *C. fasciculata*.

## CLI. HYPOXIDACEÆ.

### 1. HYPOXIS, Linn.

1. *H. trichocarpa*, Wight, Ic. t. 2045.—*H. latifolia*, *H. leptostachya*, *H. pauciflora*, et *H. brachystachya*, Wight, Ic. tt. 2044, 2045, 2046.—C.P. 2288. Var. *linearifolia*.—C.P. 2290.

HAB. Central Province, up to an elevation of 5000 feet. Nom. vulg. "Mā-bintal."

A rather variable plant as regards the size and shape of the leaves and the amount of pubescence.

## 2. CURCULIGO, Gærtn.

1. *C. recurvata*, Dryand. (Roxb. Fl. Ind. ii. p. 145.)—c.p. 3210.

HAB. Saffragam, *Moon*.

I have never met with this plant in a wild state.

2. *C. orchoides*, Gærtn. (Roxb. Fl. Ind. ii. p. 144, cum syn.)—*C. brevifolia*, Ait.; Wight, Ic. t. 2043.—c.p. 2289.

HAB. Not uncommon in grassy places, up to an elevation of 3000 feet. Nom. vulg. "Heen-bintal."

Extremely variable in the amount of pubescence, in the shape and size of the leaves, and in the number of flowers; and I cannot help suspecting that *C. Malabarica*, Wight, will prove to be a form of it, and perhaps also *C. graminifolia*, Nimmo, of the Bombay Flora.

## CLII. AMARYLLIDACEÆ.

### 1. CRINUM, Linn.

1. *C. Asiaticum*, Linn. (Kth. Enum. v. p. 547, cum syn.)—*C. toxicarium*, Roxb. Fl. Ind. ii. p. 134.—c.p. 2969.

HAB. Very abundant on the sea-coast of the island. Nom. vulg. "Tolabo."

Frequently planted as a fence for native gardens near the sea.

2. *C. defixum*, Gawl. (Kth. l. c. p. 561, cum syn.)—*C. Asiaticum*, Roxb. l. c. p. 127.—c.p. 2338.

HAB. Hot dry northern parts of the island.

3. *C. ornatum*, Herb. (Kth. l. c. p. 573, varr. *a* et *β*, cum syn.)—*C. Zeylanicum*, Linn.; Roxb. l. c. p. 139. Var. *a*, c.p. 3734. Var. *β*, c.p. 3735.

HAB. Warmer parts of the island. Nom. vulg. "Goda-mānil."

### 2. PANCRATIUM, Linn.

1. *P. Malabaricum*, Herb. (Kth. Enum. v. p. 661, cum syn.)—c.p. 2339.

HAB. On the banks of rivers and streams, up to an elevation of 2000 feet. Nom. vulg. "Deya-mānil."

2. *P. Zeylanicum*, Linn. (Kth. l. c. p. 662, cum syn.; Roxb. Fl. Ind. ii. p. 124.)—c.p. 3211.

HAB. Amongst grass, in the warmer parts of the island. Nom. vulg. "Wal-Loonoo."

The Cinghalese employ the bulbs medicinally.

*P. verecundum*, Soland., stated to be a native of the island, I have never met with here, either wild or under cultivation.

## CLIII. BURMANNIACEÆ.

(Includ. *Taccaceas*; auct. J. D. Hooker.)

## 1. BURMANNIA, Linn.

1. **B. distachya**, R. Br. Prod. p. 52.—*B. disticha*, Linn.; Roxb. Fl. Ind. ii. p. 117, cum syn.—C.P. 2313.

HAB. Common in the hotter parts of the island. Nom. vulg. "Mā-deya-jāwāla."

2. **B. Championii**, Thw.; *caule* filiformi, simplici, aphylo, squamoso; *floribus* subcapitatis; *perigonii* tubo 3-costato, nec alato.—*B. pseudo-alata*, Champ. MSS.—C.P. 2735.

HAB. Saffragam District and Hinidoon Corle.

Tota planta albida, 4–8-pollicaris. *Squamæ* caulis sparsæ, lanceolatæ, 2 lin. longæ. *Flores* 9–12, circiter 3 lin. longi. *Perigonii* lobi majores lanceolati, marginibus involutis, *minores* lineares, truncati.

3. **B. pusilla**.—*Gonyanthes pusilla*, Miers, in Linn. Soc. Trans. xviii. p. 537, t. 38, f. 3.—C.P. 3023. Var.  $\beta$ ; parum elatior, alis tubi perigonii angustioribus.—C.P. 2312.

HAB. Var.  $\alpha$ ; North of the island, *Gardner*. Trincomale, *Rev. S. O. Glenie*. Var.  $\beta$ ; Caltura, *Gardner*; Kokool Corle.

There does not appear to me sufficient difference in the seed-vessels of the preceding three species to make it desirable to constitute two genera of them. I observe in my specimens of *B. distachya*, a disposition in the capsules to split transversely about their middle, just as takes place in the present species.

## 2. THISMIA, Griff.

*Tribrachys*, Champion.

1. **T. Gardneriana**, Hk. f. MSS.; *caule* unifloro, perianthii lobis latioribus rotundatis.—*Tribrachys Gardneriana*, Champ. MSS. et Ic. in Herb. Hook.

HAB. Galle, *Champion* (*Herb. Hook.*).

I have never met with this species. "Very nearly allied to *T. Brunonis*, Griff. Linn. Trans. xix. 341. t. 39, but differs in the solitary flower and broader lobes of the perianth."—*J. D. H.*

## 3. TACCA, Forst.

1. **T. pinnatifida**, Forst. (Roxb. Fl. Ind. ii. p. 172; Miquel, Fl. Ind. Bat. iii. p. 566, cum syn.)—C.P. 2320.

HAB. North of the island, *Gardner*. Trincomale, *Rev. S. O. Glenie*.

## CLIV. DIOSCOREACEÆ.

## 1. DIOSCOREA, Linn.

1. **D. pentaphylla**, Linn. (Kth. Enum. v. p. 396, cum syn.; Wight, Ic. t. 814.)—*D. triphylla*, Linn.; Kth. l. c. p. 392, cum syn.—C.P. 2869.

HAB. Very abundant in the warmer parts of the island. Nom. vulg. "Kattoo-wella."

The tuberous roots of this and of the following species, excepting *D. bulbifera*, are eaten by the Cinghalese.

2. *D. tomentosa*, Kœnig. (Roxb. Fl. Ind. iii. p. 805.)—*Helmia?* *tomentosa*, Kth. l. c. p. 439; Wight, Ic. t. 815.—c.p. 2304.

HAB. Common up to an elevation of 2000 feet. Nom. vulg. "Ooyala."

3. *D. bulbifera*, Linn. (Willd. Sp. Pl. iv. p. 793, cum syn.; Wight, Ic. t. 878.)—*Helmia bulbifera*, Kth. l. c. p. 435, cum syn.—c.p. 2977.

HAB. Very common on the banks of rivers and streams up to an elevation of 2000 feet. Nom. vulg. "Pannoo-Condöl."

The tubers of this species are employed by the Cinghalese for attracting fish to certain spots where they can be easily caught. The tubers are broken into pieces and thrown into the water daily for some time.

4. *D. oppositifolia*, Linn. (Roxb. l. c. p. 804; Wight, Ic. t. 813.)—c.p. 2303.

HAB. Abundant in the hotter parts of the island. Nom. vulg. "Hirri-talla."

Tota planta plus minus villosula, foliis denique subglabris.

5. *D. intermedia*, Thw.; glaberrima, inermis; caule tereti; foliis oppositis vel alternis, ellipticis oblongisve, acuminatis, mucronatis, 3-nerviis; nervis lateralibus margine subparallelis et exinde costa distantibus; spicis FÆM.: solitariis, simplicibus, folio subæquilongis; MASC.: folio brevioribus, fasciculatis vel longiuscule paniculatis.—c.p. 2870, 3022.

HAB. Hotter parts of the island, not very common.

Folia circiter 3 poll. longa,  $\frac{1}{2}$ – $1\frac{1}{2}$  poll. lata, petiolo  $\frac{1}{2}$ –1 poll. longo. Spicæ masculæ pollicares vel parum ultra. Flores 6-andri.

Allied to the preceding, but differs in being entirely glabrous, in the leaves having their lateral nervures more distant from the costa, and in the male flowers being less crowded and on somewhat longer spikes. The present is well distinguished from the following species, some forms of which it closely resembles, by its shorter and more numerous spikes of male flowers, and the more diverging lateral nervures of its leaves.

6. *D. spicata*, Roth. Pl. Nov. p. 571<sup>p</sup> a; foliis basi acutis.—c.p. 2871.  $\beta$ ; foliis basi rotundatis, cordatis vel etiam profunde cordatis.—c.p. 2872, 3119.

HAB. Very common in the Central Province, up to an elevation of 4000 feet.

Glabra, inermis. Folia alterna vel opposita, elliptica vel oblonga, acuminata, mucronata, 3–5 vel rarius 7-nervia, prominentim reticulata, petiolo  $\frac{1}{2}$ – $\frac{3}{4}$  poll. longo. Spicæ sparsifloræ, sæpius valde elongatæ; FÆM. solitariæ; MASC. solitariæ vel binæ, aliquando parce ramosæ. FLOR. MASC. 6-andri. FLOR. FÆM. stigmatibus subsessilibus, parvis, simplicibus, reflexis: staminibus sterilibus linearibus, apice emarginatis.

*Dioscorea sativa*, Linn. (nom. vulg. "Kattoo-kookoolala," "Jaa-waal,"), *D. alata*, Linn. (nom. vulg. "Kirri-koudöl," "Kahata-koudöl," "Lingoorella"), and *D. purpurea*, Roxb., are cultivated in native gardens.

## CLV. PANDANEÆ.

## I. PANDANUS, Linn.

1. **P. odoratissimus**, Linn. (Roxb. Plant. Cor. tt. 94, 95, 96; Flor. Ind. iii. p. 738, cum syn.; Miq. Fl. Ind. Bat. iii. p. 156, cum syn.)—  
An *P. fascicularis*, Willd. ?—C.P. 3739.

HAB. Very abundant near the sea. Nom. vulg. "Moodoo-kaiyeya."

*Stamina* in racemulis numerosis ramos spadiceis masculi dense investientibus.

The aerial roots are used medicinally by the Cinghalese.

2. **P. humilis**, Rumph. (Miq. l. c. p. 160, cum syn.)—C.P. 3740.

HAB. Used as a fence-plant for paddy-fields in the warmer parts of the island. Nom. vulg. "Doonoo-kaiyeya."

*Stamina* simplicia, densissime conferta in ramos spadiceis masculi, filamentis brevissimis. *Capitula fructus maturi* singula 4-6 poll. longa, 3-4 poll. lata.

The leaves of this plant are made into mats by the Cinghalese.

3. **P. furcatus**, Roxb. Fl. Ind. iii. p. 744; Miq. l. c. p. 162, cum syn. t. 37.—C.P. 2734.

HAB. Similar situations with the last. Nom. vulg. "O-kaiyeya."

*Stamina* penicillatim disposita, ad apices pedunculorum numerosorum ramos spadiceis masculi arcte tegentium.

## 2. FREYCINETIA, Gaud.

1. **F. radicans**, Gaud. in Freyc. It. Bot. p. 432. t. 43; Miq. Fl. Ind. Bat. iii. p. 171.—An *F. insignis*, Blume, Rumph. p. 158, t. 42, forma minor ?—C.P. 2333.

HAB. Southern and Central Provinces, up to an elevation of 4000 feet, climbing over the trunks and branches of large trees.

*Capitula fructus maturi* 2-3 poll. longa, pulchre rubra.

2. **F. angustifolia**, Blume, l. c. p. 159, t. 43; Miq. l. c. cum syn.—C.P. 366.

HAB. In similar situations with the preceding, and more abundant.

## 3. NIPA, Rumph.

1. **N. fruticans**, Thunb. (Miq. Fl. Ind. Bat. iii. p. 150, cum syn.)—C.P. 3333.

HAB. Mangrove swamps in the south of the island. Nom. vulg. "Gim-pöl."

## CLVI. PALMACEÆ.

## TRIB. I. ARECINEÆ, Mart.

## 1. ARECA, Linn.

1. **A. Catechu**, Linn. (Roxb. Pl. Cor. i. p. 54, t. 75; Fl. Ind. iii. p. 615; Miq. Fl. Ind. Bat. iii. p. 8, cum syn.)—C.P. 3471.

HAB. Abundant in native gardens throughout the warmer parts of the island. Nom. vulg. "Poowak-gass."

An apparently larger form of this (*Ar. alba*, Rumph., Miq., haud Willd.) occurs sparingly in native gardens, but it is said to have been introduced and not to be indigenous to the island. The seeds of both kinds are used by the natives as a masticatory with Betel.

2. **A. concinna**, Thw.; inermis, *caudice* tenui, simplici, annulato; *frondibus* paucis, pinnatisectis, subglabris; *segmentis* falcatis, valde acuminatis; *terminalibus* brevioribus, plus minus confluentibus, truncatis, apice dentatis; *spadice* ramoso; *floribus* ut in *A. Catechu* dispositis; MASC.: 6-andris; *petalis* ovato-lanceolatis, quam sepala rotundato-oblonga fere 3-plo longioribus; *ovarii rudimento* trigono-claviformi; FÆM.: *drupa* fibrosa, subfusiformi, umbonata; *albumine* ruminato.—“*Areca Dicksonii*, Roxb.?” Moon’s Cat. p. 66.—C.P. 620.

HAB. Forests in the Reigam and Pas-doon Corles. Nom. vulg. “Laina-terri-gass.”

*Caudex* 8–12-pedalis, viridis,  $1\frac{1}{2}$ – $1-\frac{3}{4}$  poll. in diam. *Fronde*s circiter 7, 3– $3\frac{1}{2}$  ped. longæ (excl. basi vaginante 16 poll. longa) *segmentis* mediis 2 ped. longis,  $2\frac{1}{2}$  poll. latis. *Drupa* sesquipollicaris, flavo-rufescens.

Apparently closely allied to *A. pumila*, Blume, Rumph. t. 99 (haud Mart.), but the male flowers of the present plant are biserial, not uniserial as described in Blume’s species. The Cinghalese chew the albumen of the seeds with their Betel.

## 2. PTYCHOSPERMA, Labill., Miq.

### Sect. *Spiranthæ*, Miq.

1. **P. rupicola**, Thw.; inermis; *caudice* elato, simplici, annulato; *frondibus* subglabris, pinnatisectis, *segmentis* lanceolatis, oblique acuminatis, *terminalibus* plus minus confluentibus, præmorso-dentatis, nervo medio superne prominenti, subtus squamoso; FLO. MASC.: 12-andris; *petalis* acutis, quam sepala rotundata obtusa 4–5-plo longioribus; *ovarii rudimento* parvo; FLO. FÆM.: *sepalis* *petalis*que rotundatis, obtusis, *ovario* rotundato, acuminato; *stigmatibus* minutissimis; *ovulo* pendulo.—“*Caryota mitis*, Willd.?” Moon’s Cat. p. 65.—C.P. 2732.

HAB. Rocky places in forests of the Southern and Central Provinces, at no great elevation. Nom. vulg. “Dotaloo-gass.”

*Caudex* 30–40-pedalis, 4–5 poll. in diam. *Fronde*s 6–8-pedales, basi vaginante 15 poll. longa. *Spathæ completæ* 2, acuminatæ, externe parce furfuraceæ, *exterior* crista acuta,  $\frac{3}{4}$  poll. lata marginata. *Spadix* valde ramosus, pallide ochraceus, plus minus rufescens, sesquipedalis, pedunculo brevi. *Drupa* fibrosa, parum carnosa, subsphærica,  $\frac{3}{4}$  poll. in diam., abrupte obliqueque acuminata. *Semen* pendulum, *albumine* ruminato, coccineo-variegato.

The seeds of this are used by the Cinghalese with their Betel, as a substitute for those of the *Areca Catechu*.

## 3. ONCOSPERMA, Blume.

1. **O. fasciculata**, Thw.; stolonifera; *caudice* elato, aculeato; *frondium* *segmentis* secus rachin fasciculatis, 4–7-nis, 4–7-stiche (vel subspiraliter) dispositis, apicibus dependentibus; *spathis* *spadice*que inermibus.—*Caryota horrida*, Moon’s Cat. p. 64 (haud Willd.)—C.P. 2337.

HAB. Forests in the Central Province, up to an elevation of 5000 feet. Nom. vulg. “Kattoo-kittool.”

*Caudex* 30–40-pedalis vel ultra, 5–6 poll. in diam., aculeis numerosis nigris compressis flexilibus patentibus armatus. *Fronde*s (excl. basi vaginante externe furfuracea

aculeataque  $2\frac{1}{2}$  ped. longa) 8-pedales; *rachide* fusco-furfuracea basin versus plus minus aculeata; *laminæ segmentis* lanceolatis, valde acuminatis, 12-18 poll. longis,  $1-1\frac{3}{4}$  poll. latis, basi parum induplicatis, nervo medio superne prominente subtus parce squamoso. *Spathæ completæ* 2, parcissime furfuracea. *Spadix* 2-pedalis, ramosus, densiflorus (2 masc. 1 fœm. in scrobic.). FLORE MASC. 9-andri, ovario sterili majusculo, fere ad basin trifido, segmentis cylindricis. *Drupa* sphaerica, semipollicaris, nigro-purpurea. *Albumen* ruminatum, albidum, ochraceo-variegatum.

Distinguished from *Oncosperma filamentosa*, Blume (*Areca Nibung*, Mart.), and *Areca horrida*, Griff., by its 9-androus male flowers and the fasciculate arrangement of the *laminæ* of its fronds.

#### 4. CARYOTA, Linn.

1. *C. urens*, Linn. (Roxb. Fl. Ind. iii. p. 625, cum syn.)—C.P. 3742.

HAB. Southern and central parts of the island, up to an elevation of 2000 feet; principally in native gardens. Nom. vulg. "Kittool-gass."

From the flowering stems of this plant toddy is drawn, from which a coarse kind of sugar is made by the Cinghalese. The wood is useful for building purposes.

#### TRIB. II. BORASSINÆ, Mart.

#### 5. BORASSUS, Linn.

1. *B. flabelliformis*, Linn. (Roxb. Cor. Pl. i. p. 50, tt. 71, 72; Fl. Ind. iii. p. 790.)—C.P. 3743.

HAB. Hot, drier, and especially the northern parts of the island. Nom. vulg. "Tal-gass."

Where this tree grows in abundance it is second only to the Cocoa nut in value to the native population, every part of the plant being made available for some useful purpose.

#### TRIB. III. CORYPHINÆ, Mart.

#### 6. CORYPHA, Linn.

1. *C. umbraculifera*, Linn. (Miq. Fl. Ind. Bat. iii. p. 49, cum syn.)—C.P. 2336.

HAB. Southern and central warm moister parts of the island. Nom. vulg. "Talla-gass."

Mats, useful for constructing temporary dwellings and for other purposes, are made from the large fronds of this fine tree.

#### 7. PHOENIX, Linn.

1. *P. sylvestris*, Roxb. Fl. Ind. iii. p. 787, cum syn.—*Palma dactylifera minor*, etc., Burm. Thes. Zeyl. p. 183, cum syn.—C.P. 3172.

HAB. Very common in the hotter parts of the island.

*Caudex* 8-10-pedalis. *Drupa* 6 lin. longa,  $3-3\frac{1}{2}$  lin. lata, nitida, rubro-nigrescente.

The Cinghalese eat the sweetish pulp of the ripe seeds, and construct mats and betel-boxes of the leaves.



TRIB. IV. *COCOINÆ*, Mart.8. *COCOS*, Linn.

1. *C. nucifera*, Linn. (Roxb. Pl. Cor. i. p. 52, t. 73; Fl. Ind. iii. p. 614; Miq. Fl. Ind. Bat. iii. p. 64, cum syn.)—c.p. 3744.

HAB. Commonly cultivated throughout the warmer parts of the island. Nom. vulg. "Pol-gass."

The value of this tree is so well known, that it is unnecessary here to make any observations on the subject.

TRIB. V. *LEPIDOCARYINÆ*, Mart.9. *CALAMUS*, Linn.

Nom. vulg. "Waiwel."

1. *C. longisetus*, Griff. Palms of British India, p. 42, tt. 189, A, B. (Martius, Gen. et Sp. Palm. p. 332; Miq. Fl. Ind. Bat. iii. p. 114.)—c.p. 2873.

HAB. Hantani District, in the Central Province, at an elevation of 2000 to 3000 feet.

*Planta fœm.*: Flores secus ramulos spadiceis per paria (1 masc., 1 hermaph.) dispositi. *Drupa* obovata (includ. rostro obtuso 2 lin. longo), pollicaris, squamis majusculis, in 5 vel 6 spiris, ochraceis, margine fuscis.

Closely allied to *C. erectus*, Roxb. (*C. schizospathus*, Griff. l. c. p. 41), of which I have specimens from Dr. Hooker, but besides the differences pointed out by Griffith in the work quoted, the rostrum of the fruit in the present species is longer and its scales larger and fewer than in *C. erectus*.

2. *C. rudentum*, Lour. (Willd. Sp. Pl. ii. p. 203; Roxb. Fl. Ind. iii. p. 776.)—*Palmajuncus albus*, Rumph. Amboin. v. p. 102, t. 53.—c.p. 2874.

HAB. Hotter parts of the island. Nom. vulg. "Mā-waiwel."

3. *C. Roxburghii*, Griff. l. c. p. 55, t. 112.—*C. Rotang*, Roxb. l. c. p. 777 (non Linn. neque Willd. nisi partim).—c.p. 3388.

HAB. Common in the hotter parts of the island.

4. *C. gracilis*, Roxb. l. c. p. 781; Griff. l. c. p. 64, t. 196, cum syn.—c.p. 2334.

HAB. South of the island.

*Frondis laminæ* latitudine valde varia, secus rachin fasciculatæ vel nonnunquam flabellatim apicales. *Spadices masculi* 3-5-pollicares. *Flores* plus minus conferti. *Spad. fœm.* nondum vidi.

5. *C. tenuis*, Roxb. l. c. p. 780; Griff. l. c. p. 57, tt. 93, A, B, C.—*C. monoicus*, Roxb. l. c. p. 789?—c.p. 2335.

HAB. Hotter parts of the island, not very uncommon.

Tota planta, frondium laminis exceptis, plus minus fugaci-ferrugineo-furfuracea. *Spadices masc.* supradecompositi, ramulis florigeris brevibus, recurvis, paucifloris. *Flores* distichi, conferti.

6. *C. delicatulus*, Thw.; *frondis laminis* numerosis, æquidistantibus, lineari-lanceolatis, valde acuminatis, margine et utrinque ad nervos longiuscule setigeris; *rachide* fugaci-ferrugineo-furfuracea, carina aculeis uncinatis armata; *spadicibus masculis* valde elongatis, filiformibus, de-

compositis vel parum supradecompositis; *spathis* inermibus, basi attenuatis, apice longiuscule ciliati; *floribus* alternatim distichis, parum distantibus.—C.P. 3159.

HAB. Galle District.

*Frondis laminæ* 8 poll. longæ,  $\frac{1}{2}$  poll. latæ. *Spad. masc. rami florigeri* circiter 18, singuli 1-5 poll. longi. *Bracteæ bracteolæque* minutæ. FLOR. MASC. *Calyx* urceolatus, 3-dentatus. *Corolla* duplo longior, fere ad basin tripartita, segmentis oblongis, acutis. *Stamina* filamentis longiusculis, subliberis. *Ovarii rudimentum* minutum.

I regret not being able at present to obtain sufficient materials to draw up so complete a description as could be desired of this apparently very distinct species.

## CLVII. TYPHACEÆ.

### 1. TYPHA, Linn.

1. **T. angustifolia**, Linn. (Kunth, Enum. iii. p. 91, cum syn.; Miq. Fl. Ind. Bat. iii. p. 173, cum syn.; Roxb. Fl. Ind. iii. p. 567.)—*T. elephantina*, Roxb. l. c. p. 566.—C.P. 3218.

HAB. Warmer parts of the island, not very common.

## CLVIII. PISTIACEÆ.

### 1. PISTIA, Linn.

1. **P. Stratiotes**, Linn. (Roxb. Fl. Ind. iii. p. 131, cum syn.; Miq. Fl. Ind. Bat. iii. p. 218, cum syn.)—C.P. 3334.

HAB. Warmer parts of the island. Nom. vulg. "Deya-parandella."

### 2. LEMNA, Linn.

1. **L. minor**, Linn.; Kunth, Enum. iii. p. 4, cum syn.—C.P. 2379.

HAB. Not uncommon.

2. **L. polyrhiza**, Linn.; Eng. Bot. t. 2458?—*L. major*, Griffith, Notulæ (Monocot.), p. 220, t. 264. An *L. orbiculata*, Roxb. Fl. Ind. iii. p. 565?—C.P. 2378.

HAB. North of the island, Gardner.

## CLIX. HYDROCHARIDEÆ.

### 1. HYDRILLA, Rich.

1. **H. verticillata**, Casp. in Pringsh. Jahrb. i. 494.—*H. dentata*, Casp. Bot. Zeit. (1854) p. 56. *H. Wightii*, Planch. Ann. des Sc. Nat. (Botanique) 3me ser. xi. p. 79. *H. ovalifolia*, Rich. *H. angustifolia*, Blume, Mus. Bot. Lugd. Bat. i. p. 82. *Vallisneria verticillata*, Roxb. Fl. Ind. iii. p. 751. *Serpicula verticillata*, Roxb. Pl. Cor. t. 164.—C.P. 2310 (2305, 2309).

HAB. Common in the warmer parts of the island.

*Fructus* (in meis specim.) demum dimidio superiore filamentis paucis longiusculis acutis squarrosis appendiculato.

## 2. NECHAMANDRA, Planch.

1. **N. alternifolia**.—*N. Roxburghii*, Planch. Ann. des Sc. Nat. (Bot.) 3me ser. xi. p. 78 (excl. syn. Roxb. Pl. Cor. t. 165). *Vallisneria alternifolia*, Roxb. Fl. Ind. iii. p. 750.—C.P. 3176.

HAB. Batticaloa District.

## 3. BLYXA, Thouars.

1. **B. octandra**, Planch. in Hb. Hook.—*B. Griffithii*, Planch. in Hb. Hook. *B. Roxburghii*, Rich.; Miq. Fl. Ind. Bat. iii. p. 237, cum syn. *Vallisneria octandra*, Roxb. Pl. Cor. t. 165; Fl. Ind. iii. p. 752. *Diplosiphon oryzetarum*, Dene., in Jacq. Voy. Bot. 167.—C.P. 237.

HAB. Common in the warmer parts of the island.

FLOR. HERMAPHR. : *Stamina* 3, cum perianthii laciniis interioribus alternantia, antheris linearibus polliniferis. *Pollen* sphaericum, echinulatum.

In all the female flowers I have examined of this plant I have found three stamens alternating with the inner lobes of the perianth.

## 4. ENHALUS, L. C. Rich.

1. **E. marinus**, Griff. Not. (Monocot.) p. 178, tt. 249, 250.—*E. acoroides*, Rich.; Zoll. Cat. p. 60. *E. Koenigii*, Rich.; Miq. Fl. Ind. Bat. iii. p. 237. *Stratiotes acoroides*, Linn. f.; Willd. Sp. Pl. iv. p. 820. *Acorus marinus*, Rumph. Herb. Amb. vi. p. 191, t. 75, f. 2—C.P. 2296.

HAB. Jaffna, Gardner.

## 5. OTTELIA, L. C. Rich.

1. **O. alismoides**, Pers. Syn. p. 400, cum syn.—*Damasonium Indicum*, Willd.; Roxb. Pl. Cor. t. 185. *Stratiotes alismoides*, Linn.—C.P. 2316.

HAB. Common in the warmer parts of the island.

## CLX. ALISMACEÆ.

## 1. ALISMA, Juss.

1. **A. glandulosum**, Thw.; acaule, *foliis* longe petiolatis, ovatis, subacutis, subtus sparsim glanduloso-punctatis, profunde cordatis, lobis posticis ovatis approximatis parum acutis; *scapo* foliis subæquilongis, apice paniculato, multifloro; *floribus* 3-nis, longe pedicellatis, 6-andris; *carpidiis* 4-5, dorso 4-costatis, costis margine tuberculatis.—C.P. 2318.

HAB. Tanks in the hotter northern parts of the island.

*Folia* 6-8 poll. longa, 4-5 poll. lata, *petiolo* 2-4-pedali vel ultra. *Bracteæ* majusculæ, lanceolatae, acuminatae. *Flores* parvi, sepalis petalisque rotundatis.

2. **A. obtusifolium**.—*Sagittaria obtusifolia*, Linn.; Kunth, Enum. iii. p. 158, cum nota ("potius *Alismatis* species?"); Roxb. Fl. Ind. iii. p. 646.—*Culi-tamara*, Rheed. Hort. Mal. xi. t. 45.—C.P. 2317.

HAB. In similar situations with the last species.

## CLXI. NAJADACEÆ.

## 1. CYMODOCEA, Kœnig.

1. *C. æquorea*, Kœn. (Kunth, Enum. iii. p. 118, cum syn.)—C.P. 2380.

HAB. Point de Galle, *Gardner*.

## 2. THALASSIA, Soland.

1. *T. stipulacea*, Kœn. (Kunth, Enum. iii. p. 120, cum syn.)—C.P. 3055.

HAB. Trincomalee, *Prof. W. H. Harvey*.

## 3. POSIDONIA, Kœn.

1. *P. serrulata*, Spr. (Kunth, Enum. iii. p. 122, cum syn.)—C.P. 3056.

HAB. Trincomalee, *Prof. W. H. Harvey, Rev. S. O. Glenie*.

*Folia* linearia (in meis specim.), 6 poll. longa,  $\frac{1}{4}$  poll. lata, 15-nervia, apice rotundata, argute serrulata, basi vaginante sesquipollicari, ligula brevissima truncata. Inflorescentiam nondum vidi.

## 4. RUPPIA, Linn.

1. *R. subsessilis*, Thw.; fasciculis versus pedunculi apicem per paria approximatis, hinc floribus 2-3-subverticillatis; nuculis maturis obtuse obovoideis, subsessilibus.—C.P. 3214.

HAB. Jaffna, *Moon*.

## 5. APONOGETON, Thunb.

Nom. vulg. "Kaikateya."

1. *A. monostachyon*, Linn. f. (Roxb. Pl. Cor. i. t. 81; Fl. Ind. ii. p. 210, cum syn.)—C.P. 2307.

HAB. North of the island, *Gardner*.

2. *A. crispus*, Thunb.; Spr. Syst. ii. 465.—*A. undulatus*. Roxb. Fl. Ind. ii. p. 211.—C.P. 2306, 2308, 3381.

HAB. Central Province, up to an elevation of 6000 feet.

*Folia* oblonga, lanceolata vel linearia, basi cordata vel acuta, breve vel longiuscule petiolata.

The larger, more translucent, and transversely-veined leaves of the present species at once distinguish it from the preceding. The roots are eaten by the Cinghalese.

## 6. POTAMOGETON, Linn.

1. *P. natans*, Linn. (Kunth, Enum. iii. p. 127, cum syn.)—C.P. 590.

HAB. Ambagamowa District.

2. *P. pectinatus*, Linn. (Kunth, l. c. p. 137, cum syn.)—C.P. 3531.

HAB. Caltura.

## CLXII. AROIDEÆ.

## TRIB. I. EUAROIDEÆ.

## 1. CRYPTOCORYNE, Fisch.

1. **C. spiralis**, Fisch.; Wight, Ic. t. 773.—*Ambrosinia spiralis*, Roxb. Fl. Ind. iii. p. 492.

HAB. Ceylon, *Walker (Herb. Hook.)*.

I have not met with this species, nor are there specimens of it in this Herbarium.

2. **C. Thwaitesii**, Schott, Prod. Syst. Aroid. p. 14.—c.p. 3464.

HAB. Singhe-rajah forest, and at Hewessee, in the Pas doon Corle.

*Folia* obovata, basi apiceque parum angustata, superne punctis elevatis densissime ornata, margine elegantule undulata, 1-3 poll. longa,  $\frac{3}{4}$ -1 $\frac{1}{2}$  poll. lata, basi cordata, lobis posticis rotundatis; *petiolo* subæquilongo. *Scapus* brevissimus. *Spatha* oblonga, in rostrum longiusculum gradatim angustata, nec torta.

## 2. LAGENANDRA, Dalzell.

1. **L. ovata**.—*L. toxicaria*, Dalz. in Hook. Journ. of Bot. iv. (1852) p. 289; Miq. Fl. Ind. Bat. iii. p. 192, cum syn. *Arum ovatum*, Linn. *Karin-pola*, Rheed. Hort. Mal. xi. p. 45, t. 23.—c.p. 3315.

HAB. Very abundant in the Central Province, etc. Nom. vulg. "Kettulla."

2. **L. lancifolia**.—*Cryptocoryne lancifolia*, Schott, l. c. p. 15.—c.p. 3173, 3174.

HAB. Banks of rivers and streams, in the southern and central parts of the island. Nom. vulg. "Atta-oodiyang."

The roots are used medicinally.

*Folia* ovato-lanceolata vel lanceolata, acuminata, superne sparse subtus dense albido punctulata, basi rotundata vel acuta, 2-4 poll. longa,  $\frac{3}{4}$ -1 $\frac{1}{2}$  poll. lata; *petiolo* subæquilongo. *Scapus* brevis. *Spatha* oblonga, plus minus rugosa, in rostrum abrupte vel gradatim angustata, 1-2-pollicaris, purpurascens, interne atro-purpurea. *Ovaria* 30-40, circa medium sub stigmate conico minute tuberculata.

3. **L. Koenigii**.—*Cryptocoryne Koenigii*, Schott, l. c. p. 16; *foliis* longiusculis, lineari-lanceolatis, acuminatis, sessilibus; *scapo* foliis brevioribus; *spatha* longe et gradatim acuminata.—c.p. 3496.

HAB. Singhe-rajah forest, between Galle and Katnapoora.

*Folia* 1-2-ped. longa, 3-8 lin. lata, subtus albido-punctulata. *Scapus* 2-8 poll. longus. *Ovaria* circiter 30. *Semina* 2 $\frac{1}{2}$  lin. longa, pluri-alato-costata.

## 3. ARUM, Linn.

Sect.—*Spadix apice subulato*. Typhonium, Schott.

1. **A. divaricatum**, Linn. (Benth. Fl. Hongkong. p. 342, cum syn.)—*Typhonium divaricatum*, Blume, Rumph. i. p. 130, t. 36; Miq. l. c. p. 192, cum syn.—c.p. 3666.

HAB. Common in the hotter parts of the island.

2. **A. trilobatum**, Linn. (an Roxb.?)—*Typhonium trilobatum*, Schott; Miq. Fl. Ind. Bat. iii. p. 194, cum syn. *A. Orixense*, Roxb. Fl. Ind. iii. p. 503; Wight, Ic. t. 801.—c.p. 2896.

HAB. Very common in damp grassy places in the Central Province.  
Nom. vulg. "Panoo-alla."

The leaf of Roxburgh's *A. Orizense*, as figured by Wight, is so similar in form to that of the present plant, that I cannot doubt their being the same species, and that the drawing of the inflorescence of the former must be incorrect. The Cinghalese employ the bruised roots to destroy maggots in the sores of cattle.

#### 4. ARISÆMA, Mart.

1. *A. curvatum*, Kunth, Enum. iii. p. 20, cum syn.—*Arum curvatum*, Roxb. Fl. Ind. iii. p. 506; Wight, Ic. t. 788.

HAB. Ceylon, *Walker. (Herb. Hook.)*

2. *A. filiforme*, Blume, Rumph. i. p. 102, t. 28.—An *A. curvati*, Kunth, var. ?—c.p. 3118.

HAB. South of the island.

In my specimens the five folioles are radiate, not pedate as in Blume's figure, but in other respects the resemblance appears to be perfect.

3. *A. papillosum*, Steud. (Schott, Synop. Aroid. p. 31.)—An *A. erubescens*, Sch. l. c. p. 30?—c.p. 546.

HAB. Central Province, at an elevation of 4000 to 6000 feet. Nom. vulg. "Wal-kidāran."

The roots are employed as a medicine by the Cinghalese.

#### 5. AMORPHOPHALLUS, Blume.

1. *A. campanulatus*, Bl. Rumph. i. p. 144, tt. 32, 33; Miq. Fl. Ind. Bat. iii. p. 201, cum syn.; Wight, Ic. t. 785.—c.p. 2823.

HAB. Hotter parts of the island, especially near the sea. Nom. vulg. "Kidāran."

The Cinghalese eat the roots of the cultivated variety (nom. vulg. "Kanna-kidāran") of this plant, and use those of the wild one medicinally.

2. *A. Zeylanicus*, Blume, l. c. p. 148.—*Brachyspatha Zeylanica*, Schott, Synop. Aroid. p. 36.

HAB. Between Tangulle and Matura, *Kœnig. (Blume, l. c.)*

I have not been so fortunate as to meet with this or with the next species, nor are there specimens of either in this Herbarium.

3. *A. dubius*, Blume, l. c. p. 142.—*Schena*, Rheed. Hort. Mal. xi. p. 35, t. 18.

HAB. Ceylon, *Rheede, l. c.*

#### 6. COLOCASIA, Ray.

1. *C. antiquorum*, Schott, Syn. Aroid. p. 40, cum syn. (Miq. Fl. Ind. Bat. iii. p. 202, cum syn.)—*Arum Colocasia*, Linn.; Roxb. Fl. Ind. iii. p. 494. *A. nymphæifolium*, Roxb. l. c. p. 495; Wight, Ic. t. 786.—c.p. 324.

HAB. Very abundant on the banks of streams and other damp places. Nom. vulg. "Gahala."

The very young leaves of this, and the tubers and young leaves of the cultivated variety (nom. vulg. "Kandalla") are eaten by the Cinghalese.

2. **C. macrorrhiza**, Schott, Mel. i. 18; Kunth, Enum. iii. p. 38.—*C. odora*, Brongn. Kunth, l. c. p. 39; Benth. Fl. Hongkong. p. 343, cum syn. *Alocasia macrorrhiza*, Schott, Miq. l. c. p. 205, cum syn. *Arum odorum*, Roxb. l. c. p. 499. *A. macrorrhizum*, Linn. Sp. 1369; Fl. Zeyl. no. 327.—c.p. 3725.

HAB. Warmer parts of the island. Nom. vulg. "Habarella."

The tubers and young leaves of the cultivated, and the very young leaves of the wild plant, are eaten by the Cinghalese.

3. **C. cucullata**, Schott, Mel. i. 18; Kunth, l. c. p. 38, cum syn.—*Alocasia cucullata*, Schott, Syn. Aroid. p. 48. *Arum cucullatum*, Lour.; Roxb. l. c. p. 501; Wight, Ic. t. 787.—c.p. 3746.

HAB. On the roadside between Cultura and Galle: possibly not truly indigenous.

4. **C. vivipara**.—*Remusatia vivipara*, Schott; Kunth, l. c. p. 36; Wight, Ic. tt. 798, 900. *Arum viviparum*, Roxb. l. c. p. 496.—c.p. 2323.

HAB. Common in forests of the Central Province.

*C. Indica* is cultivated in native gardens as a vegetable.

## TRIB. II. ORONTIÆ.

### 7: LASIA, Lour.

1. **L. spinosa**.—*L. heterophylla*, Schott; Miq. Fl. Ind. Bat. iii. p. 176, cum syn. *L. Hermannii*, Schott. *Pothos heterophylla*, Roxb. Fl. Ind. i. p. 437; Wight, Ic. t. 777. *Dracontium spinosum*, Linn. Fl. Zeyl. No. 328.—c.p. 2978 (3315 partim).

HAB. Warmer parts of the island, not uncommon. Nom. vulg. "Kohilla," "Mahakohilla."

A very variable species. The leaves and roots are used as a vegetable in curries, and are also employed medicinally by the Cinghalese.

### 8. SCINDAPSUS, Schott.

1. **S. Peepla**, Schott, Mel. i. 21; Kunth, Enum. iii. p. 62.—*Rhaphidophora Peepla*, Schott, Prod. Aroid.; Benth. Fl. Hongkong. p. 344, cum syn. *Pothos Peepla*, Roxb. Fl. Ind. i. p. 433; Wight, Ic. t. 780.—c.p. 3667.

HAB. Not uncommon in the Central Province, growing over the trunks of large trees.

2. **S. decursivus**, Schott, Mel. i. 21. (Kunth, l. c. cum syn.; Miq. Fl. Ind. Bat. iii. p. 186, cum syn.)—*Rhaphidophora decursiva*, Schott, Gen. Aroid. t. 77. *Pothos decursivus*, Roxb. l. c. p. 436; Wight, Ic. t. 779.—c.p. 2322.

HAB. Forests of the Central Province, at an elevation of 2000 to 4000 feet. Nom. vulg. "Dada-kaihel."

### 9. POTHOS, Linn.

1. **P. scandens**, Linn. (Roxb. Fl. Ind. i. p. 430, cum syn.; Wight, Ic. t. 776.)—*P. scandens* and *P. Hookeri*, Schott.—c.p. 2321.

HAB. Extremely abundant in the warmer parts of the island. Nom. vulg. "Pota-wel."

The seeds are eaten by the Cinghalese, after being a long time boiled.

2. **P. remotiflora**, Hook. Ic. Pl. ii. t. 133.—An *P. oxyphyllus*, Miq. Fl. Ind. Bat. iii. p. 181, t. 38?—C.P. 2432.

HAB. Central Province, up to an elevation of 4000 feet.

*Planta juvenis*: *Caulis* filiformis, elongatus, super truncos arborum repens et iisdem arcte adhærens. *Folia* parva, plus minus conferta, rotundata, apiculata, profunde cordata, 7-nervia, prominentim reticulata, brevi-petiolata.

It will be seen by the above description of it that the young state of this plant is so unlike the appearance it subsequently assumes, that without tracing the transition from one to the other it would be difficult to believe them to belong to one species.

### TRIB. III. ACOROIDEÆ.

#### 10. ACORUS, Linn.

1. **A. Calamus**, Linn. (Roxb. Fl. Ind. ii. p. 169; Kunth, Enum. iii. p. 87, cum syn.)—C.P. 3745.

HAB. Not very uncommon in native gardens. Nom. vulg. "Wadakahā."

The roots and leaves are used medicinally by the Cinghalese.

### CLXII. LILIACEÆ.

#### 1. ASPARAGUS, Linn.

(Nom. vulg. "Hātāwaareya.")

1. **A. falcatus**, Linn. (Willd. Sp. Pl. ii. p. 152; Burm. Thes. Zeyl. t. 13. f. 2.; Linn. Fl. Zeyl. No. 123.)—C.P. 2298.

HAB. Common in the warmer parts of the island.

*Rami* scandentes, angulati, aculeati. *Folia* 2½–4 poll. longa, 1–1½ lin. lata, acuta, sessilia, solitaria vel fasciculata. *Racemi* 5–7 lin. longi, 8–10 flori. *Pedicelli* 2 lin. longi, infra medium articulati. *Bracteæ* parvæ, ovatæ, acutæ. *Perianthii* segmenta oblonga, obtusa, 1 lin. longa.

2. **A. racemosus**, Willd. (Roxb. Flor. Ind. ii. p. 151; Wight, Ic. t. 2056.)—*A. floribunda*, Kunth, Enum. v. p. 98, cum syn. An *A. acerosus*, Roxb. ?—C.P. 2299 (737 partim).

HAB. Abundant in the warmer parts of the island.

*Folia* ½–1 poll. longa. *Racemi* sesquipollicares, ad apicem floriferi. *Pedicelli* 2 lin. longi, infra medium articulati. *Perianthii* segmenta 1½ lin. longa.

3. **A. sarmentosus**, Linn. (Kunth, l. c. p. 97, cum syn.)—C.P. 2300 (737 partim).

HAB. Not uncommon, up to an elevation of 5000 feet.

*Folia* 4–6 lin. longa, ¼–½ lin. lata, plus minus fasciculata. *Racemi* elongati, foliis terminati. *Pedicelli* 2 lin. longi, paullo supra medium articulati. *Perianthii* segmenta 1½–1¾ lin. longa.

This, like the preceding species, has wider leaves when growing in the hills than when occurring near the sea.

The species are all used medicinally by the Cinghalese.



## 2. DRACÆNA, Vand.

1. **D. elliptica**, Thunb. (Kunth, Enum. v. p. 14.)—*D. spicata*, Roxb. var. *floribus minoribus*. *Cordyline Sieboldii*, Planch. Miq. Fl. Ind. Bat. iii. p. 557, cum syn.—c.p. 2293.

HAB. Shady places, up to an elevation of 4000 feet.

*Folia* oblonga, oblongo-lanceolata vel lineari-lanceolata, longiuscule acuminata, basi acuta, 6-12 poll. longa,  $\frac{1}{4}$ -2 poll. lata, *petiolo* bipollicari. *Flores* pallide virides, 4-5 lin. longi.

## 3. SANSEVIERA, Thunb.

1. **S. Zeylanica**, Willd. (Kunth, Enum. v. p. 18. cum syn.)—c.p. 2297.

HAB. Hotter parts of the island. Nom. vulg. "Neyanda."

The leaves of this plant yield a fibre of very excellent quality.

## 4. DIANELLA, Lam.

1. **D. ensifolia**, Redouté, (Kunth, Enum. v. p. 50, cum syn.)—*D. nemorosa*, Lam. Roxb. Fl. Ind. ii. p. 164. (excl. syn. Rumph.)—c.p. 3212.

HAB. Common in the Central Province, up to an elevation of 4000 feet. Nom. vulg. "Monæra-pattang."

## 5. DISPORUM, Salisb.

1. **D. Leschenaultianum**, Don, (Kunth, Enum. iii. p. 207, cum syn.; Wight, Ic. t. 2048.)—*D. Mysorense* and *D. Ceylanicum*, Wight, Ic. t. 2049.—c.p. 2319.

HAB. Central Province, at an elevation of 4000 to 7000 feet, on the margins of woods.

## 6. SMILAX, Tourn.

1. **S. maculata**, Roxb. (Kunth, Enum. v. p. 218, cum syn.; Wight, Ic. t. 2059.)—*S. fulgens*, Wall. Kunth, l. c. p. 219. *S. Villandia*, Ham. in Wall. Cat. No. 5123. *S. Nilgherrensis*, St.—c.p. 75 (308).

HAB. Central Province, up to an elevation of 5000 feet.

2. **S. ovalifolia**, Roxb. Fl. Ind. iii. p. 794; Wight, Ic. t. 809; Kunth, l. c. p. 249.—*S. macrophylla*, Roxb. l. c. p. 793. *S. prolifera*, Roxb. l. c. p. 795. An *S. retusa*, Roxb. l. c. p. 793.—c.p. 2302.

HAB. Common in the warmer parts of the island. Nom. vulg. "Maha-kabarosa."

3. **S. Zeylanica**, Linn. (Kunth, l. c. p. 249, cum syn.; Wight, Ic. tt. 2057, 2058.)—*Kari Vilandi*, Rheed. Hort. Mal. vii. p. 59, t. 31.—c.p. 3670.

HAB. Central Province, up to an elevation of 5000 feet. Nom. vulg. "Heen-kabarosa."

## 7. CHLOROPHYTUM, Ker.

1. **C. parviflorum**, Dalz. in Hook. Journ. of Bot. ii. (1850) p. 141; Bombay Flora, p. 251.—*Phalangium? parviflorum*, Wight, Ic. t. 2039.—C.P. 2292.

HAB. Not uncommon in the southern parts of the island, at no great elevation.

2. **C. breviscapum**, Dalz. l. c. ?—C.P. 2291.

HAB. Kornegalle and Matale, *Gardner*; Central Province, common. Nom. vulg. "Bim-pöl."

*Folia* pedalia, spathulato-lanceolata, acuminata. *Scapi* foliis breviores, parum densiflori. *Pedicelli* breves, apice articulati. *Bractee* lineari-lanceolatae, acutae, inferiores pollicares. *Sepala* lanceolata, 3 lin. longa, apice plus minus cucullata apiculataque. *Ovarium* loculis circiter 15-ovulatis. *Capsula* triquetra, 4 lin. longa, loculis 8-10-spermis. *Semina* atra, rugosa, minutissime tuberculata.

This appears to agree in every respect with Mr. Dalzell's description, except in the more numerous ovules of the present plant.

Used medicinally by the Cinghalese.

## 8. ALLIUM, Linn.

1. **A. Hookeri**, Thw.; *scapo* elongato, subtriangulari, basi folioso; *foliis* membranaceis, linearibus, 9-11-nerviis, *scapo* brevioribus; *umbella* multiflora; *sepalis* linearibus, parum acutis, filamentis subaequilongis; *capsula* sæpius monosperma.—*Allium*, No. 7. in Herb. Ind. Or. Hook. f. and Thoms.—C.P. 3659.

HAB. Pedrotalagalla near Newera Ellia, at an elevation of 7000 feet.

*Scapus* 1½-2-pedalis. *Folia* 2½-3 lin. lata, acuminata, apice minutissime retusa. *Pedicelli* 6 lin. longi. *Semina* ovoidea, opaca, rugulosa, nigra, 1½ lin. longa.

This is closely allied to *A. Wallichii*, Kunth, but is smaller in all its parts, and its leaves have fewer nervures.

## 9. IPHIGENIA, Kunth.

1. **I. Indica**, Kunth, (Enum. iv. p. 213, cum syn.)—*Anquillaria Indica*, R. Br.; Wall. Pl. As. Rar. iii. t. 259. *Melanthium Indicum*, Linn.—C.P. 3680.

HAB. Trincomale, *Rev. S. O. Glenie*.

## 10. GLORIOSA, Linn.

1. **G. superba**, Linn. (Roxb. Fl. Ind. ii. p. 143, cum syn.)—*Methonica superba*, Lam.; Kunth, Enum. iv. p. 276, cum syn.; Bot. Reg. t. 77. *Mendoni*, Rheed. Hort. Mal. vii. t. 57.—C.P. 2295.

HAB. Common in bushy places, up to an elevation of 3000 feet. Nom. vulg. "Neyangalla."

## 11. OPHIOPOGON, Ker.

*Ophiopogon* and *Flüggea*, Kunth (haud *Flüggea*, Willd.).

1. **O. intermedius**, Don, Prod. Fl. Nep. p. 48.—*Flüggea intermedia*,

Schult. ; Kunth, Enum. v. p. 306. *F. dubia*, Kunth, l. c. p. 305. *F. Wallichiana*, Kunth, l. c. p. 303. *O. Indicus*, Wight, Ic. t. 2050 (an Royle?). An *F. Japonica*, Rich. ; Kunth, l. c. p. 302?—c.p. 2294.  
 HAB. Central Province, up to an elevation of 4000 feet.

## 12. FLAGELLARIA, Linn.

1. **F. Indica**, Linn. (Kunth, Enum. iii. p. 370, cum syn. ; Rheed. Hort. Mal. vii. t. 53.)—c.p. 2301.  
 HAB. Hotter parts of the island, not uncommon. Nom. vulg. "Goye-wel."

## CLXIII. JUNCACEÆ.

### 1. JUNCUS, DC.

1. **J. glaucus**, Ehrh. (Kth. Enum. iii. p. 316, cum syn.)—c.p. 1003.  
 HAB. More elevated parts of the Central Province.  
 2. **J. Leschenaultii**, J. Gay, (Kth. l. c. p. 336.)—*J. Sinensis*, J. Gay ; Kth. l. c. *J. Wallichianus*, Laharpe ; Kth. l. c. p. 338.—c.p. 844.  
 HAB. With the preceding species.

### 2. SUSUM, Blume.

1. **S. anthelminticum**, Blume, (Kth. Enum. iii. p. 371.)—c.p. 2738.  
 HAB. Marshy places in the south of the island, at no great elevation.  
 Nom. vulg. "Indooroo."

*Folia* 3-pedalia vel ultra, nigro-viridia, mucronata. *Flores* dioici. *Calycis* laciniæ orbiculatæ. FL. MASC.: *Stamina* 6, filamentis æquilongis, basi coalitis, calycis laciniis interioribus adnatis. *Ovarii rudimentum* parvum. FL. FEM.: *Stamina* effœta 6, glandulas 3 retusas calycis laciniis interioribus oppositas cingentia. *Stigmata* 3, approximata, discoidea. *Baccæ* atro-purpureæ, pisi parvi magnitudine.

## CLXIV. XYRIDACEÆ.

(Nom. vulg. "Ranmanissa.")

### 1. XYRIS, Linn.

1. **X. Indica**, Linn. (Kth. Enum. iv. p. 20, cum syn.)—c.p. 3028.  
 HAB. Marshy places in the hotter parts of the island.  
 2. **X. Walkeri**, Wight, (Kth. l. c. p. 19.)—c.p. 997.  
 HAB. Similar situations with the preceding.  
*Semina* nitidiora et minus elongata quam ea spec. præcedentis.  
 3. **X. schœnoides**, Mart. (Kth. l. c. p. 16, cum syn.)—c.p. 3380.  
 HAB. Paddy fields and other wet places in the Central Province, up to an elevation of 6000 feet.  
 4. **X. pauciflora**, Willd. (Kth. l. c. p. 17, cum syn.)—c.p. 999.  
 HAB. Hotter parts of the island, in wet places.

## CLXV. RESTIACEÆ.

## 1. ERIOCAULON, Linn.

(Nom. vulg. "Kok-mota.")

1. **E. setaceum**, Linn. (Kth. Enum. iii. p. 550, cum syn.)—*E. intermedium*, Körnicke, in Linnæa, xxvii.—c.p. 791 (794).

HAB. Hotter parts of the island, in marshy places.

2. **E. Wallichianum**, Mart. in Wall. Pl. Rar. iii. t. 249; Kth. l. c. p. 564, cum syn.—*E. longifolium*, N. ab Es., Kth. l. c. p. 567.—c.p. 220.

HAB. Not uncommon in the warmer parts of the island.

3. **E. sexangulare**, Linn. (Kth. l. c. p. 551, cum syn.)—*E. Thwaitesii*, Körnicke, l. c. (partim).—c.p. 795.

HAB. Central Province.

4. **E. truncatum**, Ham. (Kth. l. c. p. 555.)—*E. Thwaitesii*, Körn. l. c. (partim).—c.p. 790 (793).

HAB. Very common.

5. **E. Brownianum**, Mart. l. c. t. 248; Kth. l. c. p. 562.—c.p. 377, 378 (1001).

HAB. Central Province, up to an elevation of 7000 feet.

An extremely variable species, as regards the amount of pubescence, and the relative size of the leaves, peduncles, and capitula. The external sepals of the male flowers are more or less coherent.

6. **E. quinquangulare**, Linn. (Willd. Sp. Pl. i. p. 485; Kth. l. c. p. 556, cum syn.)—c.p. 792.

Var.  $\beta$ . *argenteum*.—*E. argenteum*, Mart. l. c.; Kth. l. c. p. 557.—c.p. 3562.

HAB. Very abundant on the borders of paddy-fields.

7. **E. luzulæfolium**, Mart. l. c. p. 28; Kth. l. c. p. 553.—An *E. trilobum*, Ham. in Herb. Wall.; Körn. l. c. ?—c.p. 796.

HAB. Central Province, up to an elevation of 4000 feet.

8. **E. Wightianum**, Mart. l. c.; Kth. l. c. p. 563; Steud. Synops. p. 271.

Var. *capitulis nigro-cinereis, parce pilosis*.—c.p. 3382.

HAB. Ambagamowa District.

9. **E. cristatum**, Mart. l. c.—Var. *bracteis floralibus denticulatis et longiuscule cuspidato-acuminatis*.—c.p. 789.

HAB. Rambodde, in the Central Province, Gardner.

This may possibly be a distinct species. The flowers are monœcious, and the flower bracts are different in shape from those of the ordinary form of *E. cristatum*; but the two plants are so conformable in other respects, that I cannot venture to separate them.

10. **E. atratum**, Körn. l. c.—c.p. 61.

Var.  $\beta$ . *major*.—*E. caulescens*, H. f. and T. MSS.—c.p. 131.

HAB. Central Province, at an elevation of 5000 to 7000 feet.

Parce caulescens. *Folia vaginis valde dilatatis. Sepala exteriora multum latiora quam in E. cristato, cui alioquin hæc species simillima est.*

## CLXVI. CYPERACEÆ.

## 1. CYPERUS, Linn.

\* *Stylo indiviso ; stigmatè simplici.*

1. **C. Hookerianus**, Thw.—*Cyperus*, No. 21 (bis), Hook. f. and T. Herb. Ind. Orient.—c.p. 3221.

HAB. Galle.

Sarmentosus. *Folia* firma, plana, ad apicem scabra, 4–12 poll. longa. *Culmi* foliis subæquilongi, trigoni, glabri. *Involucrum* 3–4-phyllum, *capitulo* subgloboso  $\frac{1}{2}$ -pollicari multoties longius. *Spiculæ* oblongæ, compressæ, sub-12-floræ. *Squamæ* (2 vel 3 inferiores majores vacuæ) arcte imbricatæ, oblongæ, mucronato-acutæ, nitidæ, rugulosæ, ochraceæ, nigro-castaneo plus minus tinctæ,  $1\frac{1}{4}$  lin. longæ, carina scabridæ. *Stamina* 3. *Caryopsis* oblonga, rostrato-acuminata, subtrigona, glabra, castanea, basi longiuscule 3-callosa, pallida.

The undivided style and stigma, and the remarkable callosities at the base of the nut, well distinguish this from every other described species.

\*\* *Stigmatibus* 2.

2. **C. pulvinatus**, N. and M. (N. ab Es. in Wight, Contr. p. 74, cum syn.)—*C. punctatus*, Roxb. Fl. Ind. i. p. 193. *C. patens*, Vahl, Enum. ii. p. 334; Kth. Enum. ii. p. 16.—c.p. 806.

HAB. Common in wet gravelly and sandy places.

3. **C. sanguinolentus**, Vahl, l. c. p. 351; N. ab Es. l. c. p. 75, cum syn.—c.p. 802.

HAB. Not uncommon in the Central Province.

4. **C. polystachyus**, Rottb. (haud Schlecht.) (Kth. l. c. p. 13, cum syn.; N. ab Es. l. c. cum syn.; Roxb. l. c.)—c.p. 800.

HAB. Very abundant.

5. **C. flavescens**, Linn. (Steud. Synop. p. 4.)—*C. capillaris*, Koen.; N. ab Es. l. c. p. 76, cum syn. *C. Junghuhnii*, Miq. Fl. Ind. Bat. iii. p. 13.—c.p. 801.

HAB. A very common species. Dr. Hooker observes that *C. tortuosus*, Roxb., is nearly a state of it.

**C. puncticulatus**, Vahl, l. c. p. 348; Kth. l. c. p. 15.—*C. inundatus*, Roxb. l. c. p. 201 (haud R. Br.)—N. ab Es. l. c. p. 76, cum syn.—c.p. 3751 (3316 partim).

HAB. Warmer parts of the island.

My specimens are somewhat smaller than the ordinary form of this plant, and their flower-scales are slightly mucronate just below the apex.

7. **C. alopecuroides**, Rottb. (N. ab Es. l. c. (haud Roxb.); Kth. l. c. p. 19, cum syn.)—c.p. 3560.

HAB. Batticaloa District.

\*\*\* *Stigmatibus* 3.

8. **C. compressus**, Linn. (N. ab Es. l. c. cum syn.; Roxb. l. c. p. 194 (excl. syn. Pluk.); Kth. l. c. p. 23, cum syn.)—c.p. 812.

Var. *spiculis* angustis.—c.p. 807.

HAB. Very common in the warmer parts of the island.

9. **C. arenarius**, Retz, (N. ab Es. l. c. p. 77, cum syn.; Kth. l. c. p. 46, cum syn.)—c.p. 798.

HAB. Batticaloa, Gardner.

10. **C. conglomeratus**, Rottb. (Steud. Synops. p. 15.)—*C. leucocephalus*, Retz, Steud. l. c. p. 51; Kth. l. c. p. 97.—c.p. 808.

HAB. North of the island, *Gardner*.

11. **C. aristatus**, Rottb. (N. ab Es. l. c. p. 78, cum syn.)—*C. uncinatus*, R. Br. Prod. 215. *C. intricatus*, Schrad.; Kth. l. c. p. 103. *C. inflexus*, Muehl, in Willd. Herb.; Kth. l. c. p. 22. *C. angustifolius*, Ham.; N. ab Es. l. c. p. 79. *C. versicolor*, N. ab Es. l. c. p. 78.—c.p. 966.

HAB. Very abundant.

12. **C. castaneus**, Willd. (N. ab Es. l. c. p. 79, cum syn.; Kth. l. c. p. 21, cum syn.)—c.p. 803.

HAB. Common in gravelly places.

13. **C. Silhetensis**, N. ab Es. l. c. p. 79; Kth. l. c. p. 33.—c.p. 3558.

HAB. Batticaloa District.

Closely allied to *C. Haspan*, Linn., from which it appears to be principally distinguished by its denser panicle and more incurved flower-scales.

14. **C. Haspan**, Linn. (N. ab Es. l. c. p. 80, cum syn.; Kth. l. c. p. 34 cum syn.)—*C. tenuispica*, Steud. Synops. p. 11.—c.p. 799, 805, 965.

HAB. Very abundant throughout the island.

15. **C. pallidus**, Heyne? (N. ab Es. l. c. p. 79; Kth. l. c. p. 40.)—c.p. 3559.

HAB. Batticaloa District.

16. **C. articulatus**, Linn. (N. ab Es. l. c. p. 80, cum syn.; Kth. l. c. p. 53.)—*C. nudus*, Roxb. l. c. p. 187 (haud p. 209).—c.p. 3561.

HAB. Hotter parts of the island.

17. **C. rotundus**, Linn. (Roxb. l. c. p. 197; Kth. l. c. p. 58, cum syn.)—*C. hexastachyus*, Rottb.; N. ab Es. l. c. p. 81, cum syn. *C. tenuiflorus*, Rottb.; N. ab Es. l. c. p. 82. *C. pertenuis*, Roxb. l. c. p. 198. *C. bulbosus*, Vahl; N. ab Es. p. 81, cum syn.—c.p. 804, 3045.

HAB. Extremely abundant in cultivated ground, and a most troublesome weed. Nom. vulg. "Kallandooroo."

The aromatic tubers of this very variable species are much used as a medicine by the Cinghalese.

18. **C. Retzii**, N. ab Es. l. c. p. 82, cum syn.; Kth. l. c. p. 73.—c.p. 3750.

HAB. Central Province.

19. **C. procerus**, Roth. (N. ab Es. l. c. p. 83, cum syn.)—c.p. 3752 (3316 partim).

HAB. Batticaloa District.

20. **C. pennatus**, Lam. (Kth. l. c. p. 80.)—*C. canescens*, Vahl; N. ab Es. l. c. p. 84, cum syn.—c.p. 678.

HAB. Hotter parts of the island.

21. **C. exaltatus**, Retz, (N. ab Es. l. c. cum syn.; Kth. l. c. p. 70.)—*C. altus*, N. ab Es. l. c. cum syn.—c.p. 3040.

HAB. Common in the warmer parts of the island.

22. **C. Roxburghii**, N. ab Es. l. c. cum syn.; Kth. l. c. p. 99.—*C. elatus*, Roxb. l. c. p. 204 (excl. syn.).—c.p. 3041.

HAB. Hotter parts of the island, not common.

*Antheræ* connectivo longo acuto piloso terminatæ.

23. **C. xanthopus**, Steud. Syn. p. 36.—*C. infraapicalis*, N. ab Es. in Hb. Wt.—c.p. 3044.

HAB. Doombera District.

*Culmus* 3-4-pedalis vel ultra, triqueter, glaber, basi foliosus. *Umbella* decomposita. *Spiculæ* sub-8-floræ, in spicas oblongas subcylindricas 7-10 lin. longas dense aggregatæ. *Squamæ* parum distantes, oblongæ, subacutæ, infra apicem mucronatæ, pallide ochraceo-cinereæ, dorso rufo-pluristriatæ. *Caryopsis* oblonga, trigona, brevirostrata, minutissime punctulata, pallide fusca.

24. **C. dubius**, Rottb. (N. ab Es. l.c. p. 85, cum syn.)—*C. kyllingioides*, Vahl; Kth. l. c. p. 94.—c.p. 855, 2942.

HAB. Central Province, not very uncommon up to an elevation of 4000 feet.

25. **C. pilosus**, Vahl, Enum. ii. p. 354.—*C. obliquus*, N. ab Es. l. c. p. 86. *C. piptolepis*, Steud. Syn. p. 40. An *C. marginellus*, N. ab Es. l. c. p. 83?—c.p. 797.

HAB. Very common in the Central Province.

26. **C. Iria**, Linn. (N. ab Es. l. c. p. 87, cum syn.; Kth. l. c. p. 38, cum syn.)—c.p. 811.

HAB. Very abundant.

27. **C. difformis**, Linn. (N. ab Es. l. c. p. 88, cum syn.; Kth. l. c. p. 38.)—c.p. 3042.

HAB. Common in the warmer parts of the island.

28. **C. nigroviridis**, Thw.—*C. elegans*, Wall. Cat. (haud Linn.)—c.p. 2879.

HAB. Kornegalle.

*Culmus* tenuis, glaber, triqueter, 1-2-pedalis, basi foliosus. *Folia* latiuscula, culmo subæquilonga, margine scabra. *Umbella* composita, longe involuocrata. *Spiculæ* laxifloræ, umbellatæ. *Squamæ* rotundatæ, mucronatæ, nervoso-striatæ, nigro-fuscae, margine membranacea fulvescentes. *Caryopsis* ovoidea, trigona, utrinque acuta, subglabra, nigra.

Resembling very much *C. elegans*, Linn., but distinguished at once by the very different shape of its fruit, which in *C. elegans* is much wider in proportion to its length, and is rounded at the apex.

29. **C. distans**, Linn. (N. ab Es. l. c. p. 88, cum syn.; Kth. l. c. p. 93.)—c.p. 810.

HAB. Very abundant.

30. **C. corymbosus**, Rottb. (Kth. l. c. p. 55, cum syn.)—*Papyrus corymbosus*, N. ab Es. l. c. p. 89.—c.p. 809.

HAB. Kornegalle.

31. **C. Paugorei**, Rottb. (Kth. l. c. p. 57, cum syn.)—*Papyrus Paugorei*, N. ab Es. l. c. p. 88.—c.p. 813.

HAB. Not uncommon in the warmer parts of the island.

32. **C. Neesii**, Kth. l. c. p. 101.—*C. elatus*, Linn. (partim). *Papyrus elatus*, N. ab Es. l. c.—c.p. 3043.

HAB. Caltura, Moon.

33. **C. dilutus**, Vahl, Enum. ii. p. 357; Kth. l. c. p. 92.—*Mariscus dilutus*, N. ab Es. l. c. p. 90.—c.p. 815.

HAB. Kornegalle, Gardner.

34. **C. umbellatus**, Benth. Flor. Hongk. p. 386.—*Mariscus umbellatus*, Vahl, l. c. p. 376. *M. cyperinus*, Vahl, l. c. p. 377. *M. paniceus*, Link? N. ab Es. l. c. p. 89, cum syn. *M. pictus*, N. ab Es. l. c. p. 90, cum syn.—C.P. 814, 816, 817, 2878.

HAB. Extremely abundant.

*Planta valde variabilis. Spiculæ 1-4-floræ.*

## 2. KYLLINGIA, Linn.

1. **K. monocephala**, Linn. (N. ab Es. in Wight, Cat. p. 91; Kth. Enum. ii. p. 129.)—C.P. 3753.

HAB. Very common amongst grass, in the warmer parts of the island.

Differs from the next species in its usually diandrous flowers, with their glumes whiter, and somewhat suddenly dilated above the middle.

2. **K. brevifolia**, Rottb. (N. ab Es. l. c. cum syn.; Kth. l. c. p. 130.)—C.P. 3755.

HAB. With the preceding species, but not so common.

3. **K. melanosperma**, N. ab Es. l. c.; Kth. l. c. p. 131.—C.P. 818, 2980.

HAB. Central Province, at an elevation of 4000 to 6000 feet.

4. **K. cylindrica**, N. ab Es. l. c. Var.  $\beta$ . *monostachya*, Kth. l. c. p. 133.—C.P. 3754.

5. **K. triceps**, Rottb. (N. ab Es. l. c.; Kth. l. c. cum syn.)—C.P. 3234.

HAB. Hot drier parts of the island.

## 3. REMIREA, Aubl.

1. **R. pedunculata**, R. Br. (Kth. Enum. ii. p. 139, cum syn.)—*R. Wightiana*, Wall. Cat. No. 3378; N. ab Es. in Wight, Contr. p. 92.—C.P. 3327.

HAB. Jaffna, *Gardner*.

## 4. PANDANOPHYLLUM, Hassk.

*Steudel, Synops. p. 134. Halostemma, Wall. MSS.*

*Char. emend. Squamæ unifloræ. Glumæ 6, biseriales, plicatæ; 3 exteriorum unaquæque stamen includens eidemque opposita; 3 interiorum una (posterior) pistillum amplectens, duæ vacuæ. FLOR. MASC.: Antheræ lineares, subsessiles, longitudinaliter dehiscentes. Ovarii rudimentum stylo longo clavato acuto terminatum. FLOR. FÆM.: Stamina sterilia paleæformia. Stylus basi incrassatus. Stigmata 2 (Hassk.) 3 (in meis specimin.). Caryopsis ossea.*

1. **P. Zeylanicum**, Thw.; *foliis glabris, longissime setaceo-acuminatis, margine scabris; caryopsi obovoidea, lævi, brevi-stipitata.*—C.P. 3029.

HAB. Banks of a stream, near Pittigalakanda in the Kokoole Corle, December, 1853.



*Folia* 3-4-pedalia, 6-7 lin. lata. *Culmi* plures, 3-9 poll. longi, basi vaginati. *Capitulum* pluriflorum, 6-7 lin. in diam. *Squamæ* ovatae, striatae, inferiores mucronatae, semipollicares. *Glumæ* squama æquilongæ, lineari-lanceolatae, acutæ, exteriores ad carinam scabrae. *Antheræ* 2 lin. longæ. *Caryopsis* albida, interne nigra, 2 lin. longa.

Hasskarl, in describing this genus as monandrous, must, I suspect, have taken the long clavate style of the male flower for an anther. The true anthers, 3 in number, which lie concealed, one within the fold of each of the exterior glumes of the flower, may easily escape observation, unless great care be taken in the dissection.

## 5. LEPIRONIA, Rich.

1. **L. mucronata**, Rich. in Pers. Syn. i. p. 70; Kth. Enum. p. 366.  
—*Chondrachne articulata*, R. Br.—C.P. 3228.

HAB. Caltura District.

*Pistillum* 18-24 paleis (6-8 exterioribus monandris, reliquis vacuis) lanceolatis acutis cinctum.

This genus is evidently closely allied to *Pandanophyllum*, the pistil in each being surrounded by a certain number of empty paleæ, and, outside these, by one or more series of monandrous paleæ. In *Pandanophyllum* the flower (or spikelet?) is unisexual, either the pistil or the stamens being sterile or rudimentary, whilst in the present genus the flower (or spikelet?) is hermaphrodite.

## 6. HYPOLYTRUM, Rich.

1. **H. latifolium**, Rich. in Pers. Syn. i. p. 70; Kth. Enum. ii. p. 271, cum syn.—*H. giganteum*, Wall. Cat. No. 3404; N. ab Es. in Wight, Contr. p. 93, cum syn. *H. trinervium*, Kth. l. c. p. 272.—C.P. 3, 219.

Var.  $\beta$ . *minor*.—*Spiculis squamisque* multo minoribus.

HAB. Common in dense forests in the southern and central parts of the island, up to an elevation of 3000 feet.

Var.  $\beta$ . Caltura.

*Glumæ* postice connatae, sed facile disruptæ. *Caryopsis* glabra vel parum rugulosa, rarius costulata, ochraceo- vel nigro-fusca, endocarpio nigro.

2. **H. longirostre**, Thw.; *spiculis* corymboso-paniculatis; *squamis* firmis, nitidis, plus minus acutatis, mucronatis; *glumis* postice connatis, oblongis, subacutis, carinis scabris; *staminibus* 2; *caryopsi* rotundata, rugosa, parum compressa, styli basi dilatata longiuscule rostrata.—C.P. 3468.

HAB. Hinidoon Corle.

*Caryopsis* fusca, sæpe obscure costulata,  $\frac{1}{2}$  lin. longa, rostro parum longiore; endocarpio nigro.

A præcedente, cui valde similis, foliis angustioribus, squamis acutis majoribus firmioribusque et caryopsi longe rostrata satis differt.

3. **H. costatum**, Thw.; *spicis* capitato-congestis; *caryopsi* majuscula, ovata vel rotundata, acuta, longitudinaliter grosse 6-12-costata, externe spongiosa.—C.P. 3222.

HAB. Central Province, but exact locality not recorded.

*Culmus* triqueter, lævis. *Folia floralia* longiuscula, 8 lin. lata, trinervia, margine scabra. *Panicula* oblonga, dense compacta. *Caryopsis* fusco-nigra, 3 lin. in diam., interne albida, endocarpio separabili.

The only specimen I possess is not sufficiently perfect to enable me to draw up so complete a description as could be desired, of this interesting plant.

## 7. LIPOCARPHA, R. Br.

1. *L. argentea*, R. Br. (Kth. Enum. ii. p. 266, cum syn.)—*L. lævigata*, N. ab Es. in Wight, Contr. p. 92, cum syn.—c.p. 819.

Var. *bracteis majoribus*.—c.p. 3557.

HAB. Common up to an elevation of 3000 feet.

2. *L. sphacelata*, Kth. l. c. p. 267.—*L. triceps*, N. ab Es. l. c.—c.p. 3756.

HAB. Doombera District.

*Capitulum* 3-6-stachyum.

## 8. FUIRENA, Linn.

1. *F. ciliaris*, Roxb. Fl. Ind. i. p. 180; N. ab Es. in Wight, Contr. p. 93, cum syn.—c.p. 3038.

HAB. Very abundant in paddy-fields and wet grassy places, in the warmer parts of the island.

2. *F. glomerata*, Lam. (Vahl, Enum. ii. p. 386; Kth. Enum. ii. p. 184.)—*F. Rottbællii*, N. ab Es. l. c. p. 94, cum syn.—c.p. 2748.

HAB. Similar places with the last.

3. *F. umbellata*, Rottb. (Kth. l. c. p. 185, cum syn.)—*F. pentagona*, W. et A.; N. ab Es. l. c. p. 93, cum syn.; Kth. l. c. cum syn.—c.p. 3229.

HAB. With the preceding and equally common.

4. *F. uncinata*, Wall. Cat. No. 3534 *d*; Kth. l. c. p. 184, cum syn.—c.p. 3230.

HAB. Similar places with the foregoing species.

## 9. ABILDGAARDIA, Vahl.

1. *A. monostachya*, Vahl, Enum. ii. p. 296; Kth. Enum. ii. p. 247.—*A. Rottbælliana*, N. ab Es. in Wight, Contr. p. 95, cum syn.—c.p. 3231.

HAB. Not uncommon in the warmer parts of the island.

2. *A. tristachya*, Vahl, l. c. p. 297; Kth. l. c. p. 248, cum syn.—c.p. 852.

HAB. North of the island, *Gardner*.

3. *A. fulvescens*, Thw.—*Umbella* 3-4-radiata; *involucro* brevi; *spicis* planis, compressis, acutis, plurifloris, solitariis vel binis in radiis; *squamis* ovatis, acutis, pilosulis; *floribus* triandris; *caryopsi* trigona, subpyriformi, tuberculata.—c.p. 679.

HAB. Reigam Corle.

Cæspitosa, 1-1½-pedalis. *Folia* culmo breviora vel subæquilonga, lineari, obtusiuseula, mucronata, margine scabra, ½-1½ lin. lata. *Culmus* filiformis, striatus. *Caryopsis* nigro-fusca, umbonata, ¼ lin. longa.

4. *A. cinnamometorum*, Thw.—*A. fusca*, N. ab Es. l. c.; Kth. l. c. p. 249.—*Fimbristylis Cinnamometorum*, Kth. l. c. p. 229. *Scirpus Cinnamometorum*, Vahl, l. c. p. 278.—c.p. 2752.

HAB. Not uncommon in wet grassy places.

## 10. FIMBRISTYLIS, Vahl.

\* *Stigmatibus* 2.

1. **F. nutans**, Vahl, Enum. ii. p. 285; N. ab Es. in Wight, Contr. p. 96; Kth. Enum. ii. p. 221, cum syn.—c.P. 832.

HAB. Very common in the warmer parts of the island.

2. **F. acuminata**, Vahl, l. c.; N. ab Es. l. c.; Kth. l. c. cum syn.—*F. setacea*, Benth. in Hook. Journ. of Bot.—c.P. 2747.

HAB. Hotter parts of the island.

Closely allied to the preceding, but certainly distinct.

3. **F. polytrichoides**, R. Br.? (N. ab Es. l. c. cum syn.)—*Scirpus polytrichoides*, Retz; Vahl, l. c. p. 249. *Scirpus scaber*, Roxb. Fl. Ind. i. p. 220.

HAB. Ceylon, *Kœnig*.

4. **F. schœnoides**, Vahl, l. c. p. 286; N. ab Es. l. c. p. 97, cum syn.; Kth. l. c. p. 222, cum syn.—c.P. 833.

HAB. Not uncommon in the hotter parts of the island.

5. **F. diphylla**, Vahl, l. c. p. 289; N. ab Es. l. c. p. 100, cum syn.—*F. tomentosa*, Vahl, l. c. p. 290; N. ab Es. l. c. *F. Royeniana*, varr.  $\alpha$  et  $\beta$ ; N. ab Es. l. c.—c.P. 839, 840 (841 partim).

HAB. A very abundant plant.

Varies greatly in the amount of pubescence: the flowers are commonly diandrous, more rarely monandrous; I have not met with it with triandrous flowers.

6. **F. ovalis**, N. ab Es. l. c. p. 98, cum syn.—c.P. 3737 (841 partim).

HAB. Kornegalle District.

Certainly distinct from the preceding species, though bearing considerable external resemblance to the tomentose form of it.

7. **F. rigidula**, N. ab Es. l. c. p. 99; Steud. Synops. p. 116.—c.P. 3232.

HAB. Warmer parts of the island, not common.

8. **F. ferruginea**, Vahl, l. c. p. 291; N. ab Es. l. c. p. 97, cum syn.; Kth. l. c. p. 236.—c.P. 848.

HAB. Kokoole Corle, *Gardner*.

9. **F. argentea**, Vahl, l. c. p. 294; N. ab Es. l. c. p. 100, cum syn.; Kth. l. c. p. 223.—c.P. 2877.

HAB. Hotter parts of the island.

10. **F. pallescens**, N. ab Es. l. c. p. 101, cum syn.; Steud. Synops. p. 111.—*Scirpus pallescens*, Roxb. Fl. Ind. i. p. 220.—c.P. 3758.

HAB. Warmer parts of the island, not very common.

\*\* *Stigmatibus* 3. *Trichelostylis*.

11. **F. Arnottii**, Thw.—*F. xyroides*, Arn. Cat. Hb. Wight, 2331 (haud *xyridis*, R. Br.)—c.P. 830.

HAB. Common on the borders of paddy-fields, etc.

Cæspitosa, aphylla. *Culmus* parum angularis, 6–18 poll. longus, basi vaginatus. *Spica* solitaria, erecta, rotundata. *Squamæ* arcte imbricatæ, obovatae, obtusæ, inferiores parum majores. *Caryopsis* oblonga, subclavata, delicatule cancellata, stipitata, albida, stylo subæquilonga.

12. **F. miliacea**, Vahl, l. c. p. 287; Kth. l. c. p. 230.—*Trichelostylis*

*miliacea*, N. ab Es. l. c. p. 103, cum syn. *F. tetragona*, N. ab Es. l. c. p. 104, cum syn. *Scirpus tetragonus*, Roxb. l. c. p. 228.—C.P. 836.

HAB. Very abundant in paddy-fields and other wet places.

13. *F. globulosa*, Wall. (Kth. l. c. p. 231.)—*Trichelostylis globulosa*, N. ab Es. l. c. p. 105, cum syn.—C.P. 842.

HAB. Warmer parts of the island.

14. *F. quinquangularis*, Kth. l. c. p. 229, cum syn.—*Trichelostylis quinquangularis*, N. ab Es. l. c. p. 104, cum syn.—C.P. 838.

HAB. Not uncommon in the hotter parts of the island.

15. *F. insignis*, Thw.—*Culmo* filiformi, striato, apice parum compresso, glabro, basi folioso; *foliis* brevibus, firmis, curvatis, obtusis, mucronatis, glabris, margine scabris; *umbellæ* apertæ radiis circiter 6; *involucro* brevissimo; *spicis* subcylindricis, acutis, 1-2- vel 3-nis in radiis, intermedia sessili majore; *squamis* ovato-oblongis, subacutis, mucronatis; *floribus* triandris; *caryopsi* brevi-obcordata, trigona, sublævi; *stylo* lineari, triquetro, glabro; *stigmatibus* breviusculis.—C.P. 3317.

HAB. Between Negombo and Kornegalle.

*Cæspitosa*, 2-pedalis. *Spicæ*  $\frac{1}{2}$ -1 poll. longæ,  $1\frac{1}{2}$  lin. latæ. *Squamæ* fusco-ferrugineæ, basi margineque pallidiores, 2 lin. longæ.

16. *F. complanata*, Link. (Kth. l. c. p. 228, cum syn.)—*Trichelostylis complanata*, N. ab Es. p. 103, cum syn.—C.P. 3220.

HAB. Ceylon, *Walker* (in *Herb. Hook.*).

17. *F. chætorhiza*, Kth. l. c. p. 240.—*Trichelostylis chætorhiza*, N. ab Es. l. c. p. 106. *T. torta*, N. ab Es. l. c. p. 105.—C.P. 837, 970.

HAB. Extremely abundant amongst grass.

A very variable plant. I find the flowers always triandrous, and the characters of the fruit, as described by Nees, very constant.

18. *F. retusa*, Thw.; *culmo* filiformi, acute angulari, apicem versus scabrido, basi foliato; *foliis* anguste linearibus, obtusis, mucronatis; *umbella* composita, compacta, involucro subæquilongo; *spicis* glomeratis, sessilibus; *squamis* ovatis vel oblongis, retusis, dense imbricatis, pluri-seriatis, muricato-punctulatis, ad apicem margine ciliatis; *floribus* monandris; *stylo* glabro; *caryopsi* late obovata, trigona, aspero-tuberculata.—C.P. 3760.

HAB. Near Galle.

*Cæspitosa*, 1-2-pedalis. *Squamæ* fusco-ferrugineæ. *Caryopsis* ochracea, ferrugineo tincta.

19. *F. Wightiana*, N. ab Es. l. c. p. 99.—C.P. 3759.

HAB. Trincomale, *Rev. S. O. Glenie*.

20. *F. salbundia*, Kth. l. c. p. 230, cum syn.—*Trichelostylis salbundia*, N. ab Es. l. c. p. 105, cum syn. *T. pentaptera*, N. ab Es. l. c. —C.P. 823, 843.

HAB. Newera Ellia, and other of the more elevated parts of the island.

21. *F. connectens*, Thw.—*Culmo* striato, versus apicem compresso scabroque, basi folioso; *foliis* anguste linearibus, culmo subæquilongis, glabris, margine scabris, abrupte acutis; *umbella* mediocri, decomposita, compacta, radiis scabris; *involucro* diphylo, foliolis linearibus, acutis, scabris, umbella subæquilongis; *spicis* oblongis, acutis; *squamis*

2-3-chis, ovatis, subacutis, mucronulatis, opacis; *floribus* triandris; *caryopsi* obovoidea vel subpyriformi, acute trigona, nitida, sublævi; *stylo* glabro.—C.P. 2967.

HAB. Maturatta District.

Cæspitosa, 1-2-pedalis. *Folia* glauco-viridia, circiter 1 lin. lata. *Squamæ* fusco-castaneæ, carina scabra margineque pallidiores, longitudinaliter delicatule striatæ. *Caryopsis* albida vel pallide fulva.

Allied to *F. salbundia*, and might perhaps with propriety be placed with it, in the genus *Abildgaardia*, close to *A. cinnamometorum*, with which they have great affinity.

## 11. ISOLEPIS, R. Br.

1. **I. squarrosa**, R. et S. (Kth. Enum. ii. p. 202, cum syn.)—*Scirpus squarrosus*, Linn.; Roxb. Fl. Ind. i. p. 222.—C.P. 854.

HAB. North of the island, *Gardner*.

2. **I. articulata**, N. ab Es. in Wight's Contr. p. 108, cum syn. (Kth. l. c. p. 199, cum syn.)—*I. incurvatus*, N. ab Es. l. c. cum syn. *I. prolongata*, N. ab Es. l. c. *Scirpus articulatus*, Linn.; Roxb. l. c. p. 214. *S. incurvatus*, Roxb. l. c. *S. subarticulatus*, Roxb. l. c. p. 215.—C.P. 846.

HAB. Common in the hotter parts of the island.

3. **I. supina**, R. Br. (N. ab Es. l. c. p. 107, cum syn.; Kth. l. c. p. 196.)—*Scirpus supinus*, Linn.; Roxb. l. c. p. 217.—C.P. 3233.

HAB. Hotter parts of the island.

4. **I. trifida**, N. ab Es. l. c. p. 108. (Kth. l. c. p. 213.)—C.P. 851.

HAB. More elevated parts of the Central Province.

5. **I. gracilis**, N. ab Es. l. c. p. 109. (Kunth, l. c. p. 217.)—C.P. 834.

HAB. Common in the warmer parts of the island.

6. **I. barbata**, R. Br. (N. ab Es. l. c. cum syn.; Kth. l. c. p. 208.)—*Scirpus monander* et *S. antarcticus*, Roxb. l. c. (excl. syn.)—C.P. 829.

HAB. Not uncommon in the hotter parts of the island.

7. **I. pulchella**, Thw.; *spicis* terminalibus, capitato-congestis; *culmo* lævi, filiformi, striato, basi foliato; *foliis* brevibus, setaceis, canaliculatis, basi in vaginam apice barbatam dilatatis; *involucro* quam capitulum 2-3-plo longiore, basi barbato; *squamis* late obtuse carinatis, pilosulis; *floribus* triandris; *caryopsi* rotundata, trigona, minute tuberculata, brevi-stipitata.—C.P. 3761.

HAB. Trincomale, *Rev. S. O. Glenie*.

Cæspitosa, 3-6-pollicaris. *Squamæ* ferrugineo-aureæ, versus apicem carinæ pallide virides. *Caryopsis* pallida, nigro-apiculata.

8. **I. fluitans**, R. Br. (Kth. l. c. p. 188, cum syn.)—*Eleogiton fluitans*, Link. Hort. i. 284. *E. curvulus*, N. ab Es. l. c. p. 110.—C.P. 835 (2634).

HAB. Common in the more elevated parts of the Central Province.

9. **I. dipsacea**, R. et S. (Kth. l. c. p. 205, cum syn.)—*Echinolytrum dipsaceum*, Desv.; N. ab Es. l. c. p. 96. *Scirpus dipsaceus*, Rottb.—C.P. 668.

HAB. Warmer parts of the island.

## 12. SCIRPUS, Linn.

1. **S. mucronatus**, Linn. (Kth. Enum. ii. p. 161, cum syn.)—*S. triangulatus*, Roxb. Fl. Ind. i. p. 217; N. ab Es. in Wight's Contr. p. 111. *S. javanus*, N. ab Es. l. c. p. 112.—c.p. 41.

HAB. Very common.

2. **S. juncoides**, Roxb. l. c. p. 216; Kth. l. c. p. 160.—*S. junciformis*, N. ab Es. l. c. p. 112.—c.p. 850.

HAB. Abundant throughout the island.

3. **S. subcapitatus**, Thw.; aphyllus; *culmis* confertis, rigidis, teretibus, vaginatis; *vaginis* subulato-apiculatis, superiore multum longiore; *involucro* brevi, ovato, 5-nervio, subulato-apiculato; *spicis* 3 vel 4, oblongis, acutis, subcapitatis; *squamis* sub-4-scriatis, ovatis, membranaceis, carinatis, mucronatis; *staminibus* 3; *caryopsi* obovato-oblonga, trigona, lævi; *setis* 6, tenuibus, denique caryopsi multiplo longioribus.—c.p. 306.

HAB. Newera Ellia.

6-18-pollicaris. *Capitula* ochracea, ferrugineo tincta. *Stylus*, cum stigmatibus et setis, nigro-rufus. *Caryopsis* ferruginea.

4. **S. grossus**, Linn. (Kth. l. c. p. 167, cum syn.)—*Hymenochæte grossa*, N. ab Es. l. c. p. 110, cum syn.—c.p. 847.

HAB. Hotter parts of the island.

Mats are made by the Cinghalese from the culms of this species.

5. **S. pectinatus**, Roxb. l. c. p. 218.—*Malacochæte pectinata*, N. et M.; N. ab Es. l. c. cum syn. An *Scirpus subulatus*, Vahl, Enum. ii. p. 268? (conf. *S. littoralis*, Schrad.)—c.p. 831.

HAB. Caltura.

## 13. CHÆTOCYPHERUS, N. ab Es.

1. **C. setaceus**, N. ab Es. (Steud. Synops. p. 73.)—*C. Limnocharis*, N. ab Es. in Wight's Contr. p. 96, cum syn. *Eleocharis chætaria*, Kth. Enum. ii. p. 140, cum syn. *Cyperus setaceus*, Retz.—c.p. 247.

HAB. Saffragam District.

## 14. ELEOCHARIS, R. Br.

1. **E. tetraquetra**, N. ab Es. in Wight's Contr. p. 113, cum syn.; Kth. Enum. ii. p. 150, cum syn.—c.p. 2397.

HAB. Rambodde in the Central Province, at an elevation of 4000 feet.

2. **E. ovata**, R. Br. (Kth. l. c. 151, cum syn.)—*E. congesta*, Don; Kth. l. c. p. 152, cum syn.—c.p. 2635.

HAB. More elevated parts of the Central Province.

*Spicæ* sæpe basi proliferæ. *Stigmata* 3. *Caryopsis* pyriformis, trigona, nitida, ochracea.

3. **E. capitata**, R. Br. (Kth. l. c. p. 150, cum syn.)—*Eleogenus capitatus*, N. ab Es. l. c. p. 112, cum syn.—c.p. 3039.

HAB. Saffragam District.

4. **E. fistulosa**, Schult. (Kth. l. c. p. 155, cum syn.)—*E. acutangula*,

N. ab Es. l. c. p. 114, cum syn. *Scirpus acutangulus*, Roxb. Fl. Ind. i. p. 213.—C.P. 3162 (3047).

HAB. Hotter parts of the island.

*Stigmata* (in meis specim.) 3.

5. **E. plantaginea**, R. Br. (Kth. l. c. p. 153, cum syn.)—*Limnocharis plantaginea*, N. ab Es. l. c. *Scirpus plantagineus*, Rottb.; Roxb. l. c. p. 212.—C.P. 3046.

HAB. South of the island.

6. **E. spiralis**, R. Br. (Kth. l. c. p. 155, cum syn.)—*Limnocharis spiralis*, P. de B.; N. ab Es. l. c. cum syn. *L. media*, N. ab Es. l. c. cum syn. *Scirpus spiralis*, Rottb., et *S. medius*, Roxb. l. c. pp. 212, 213.—C.P. 853.

HAB. Hotter parts of the island; Caltura, *Macrae*.

I can find no essential difference between *Macrae's* specimens, named *Limnocharis media* by Dr. Wight, and the ordinary *E. spiralis*, R. Br.

### 15. ARTHROSTYLIS, R. Br.

1. **A. filiformis**, Thw.; *culmis* confertis, filiformibus, longiusculis, glabris, sulcato-triquetris, basi vaginatis; *involucro* capitulo subæquilongis; *spicis* plurimis, 1-2-floris; *squamis* circiter 5, trinerviis (vel superioribus sub-5-7-nerviis) apice recurvis, aristato-subulatis; *floribus* 3-andris; *stylo* elongato; *stigmatibus* 2-3, elongatis; *caryopsi* obovata, trigona, obscure minute papillosa, apice vix umbonata.—C.P. 3169.

HAB. Carawitta Kanda, near Ratnapoora; and Dolosbage District.

*Rhizoma* repens. *Culmi* 2-3-pedales, *vaginis*  $\frac{1}{2}$ -2-poll. longis, cuspidatis. *Capitulum* semipollicare. *Squamæ* ochraceæ, fulvo tinctæ. *Stylus*, cum basi dilatata, nigro-rufus. *Caryopsis* albida.

### 16. RHYNCHOSPORA, Vahl.

1. **R. laxa**, R. Br. (Kth. Enum. ii. p. 298, cum syn.)—An *R. triflora*, Vahl, Enum. ii. p. 231?—C.P. 677.

Var.  $\beta$ . *minor*.—*R. Chinensis*, N. et M.; N. ab Es. in Wight's Contr. p. 115.—C.P. 2396.

HAB. Var. *a*, south of the island; var.  $\beta$ , very abundant amongst grass at an elevation of 4000 to 6000 feet.

2. **R. aurea**, Vahl, l. c. p. 291; Kth. l. c. p. 293, cum syn.—C.P. 199.

HAB. Common in swampy places in the hotter parts of the island.

3. **R. Zeylanica**, Kth. l. c. p. 294.—*Cephaloschænus Zeylanicus*, N. ab Es. in Wight's Contr. p. 115.—C.P. 3036.

HAB. Saffragam District.

Closely allied to and possibly a variety of *R. aurea*, but I am disposed to consider it distinct: the present plant is more glaucous, the leaves much narrower, the panicle fewer-flowered, and the rostrum of the fruit very different in shape. *R. articulata*, Schult., of which I have a specimen from Khasia given me by Dr. Hooker, is nearly allied to both the above-mentioned plants, but specifically distinct, I think.

4. **R. Wallichiana**, Kth. l. c. p. 289, cum syn.; Benth. Fl. Hongkong. p. 396, cum syn.—*R. Hænkei*, Pr.; Kth. l. c. p. 290. *Morisia Wallichii*, N. ab Es. l. c. p. 115. *Haplostylis Meyenii*, N. ab Es. l. c.—C.P. 2746.

HAB. Common amongst high grass, up to an elevation of 4000 feet.

17. **CLADIUM**, P. Browne.

1. **C. undulatum**, Thw. ; *foliis* rigidis, angustis, acutissimis, margine involutis, scabriusculis, *culmo* subtrigono, sulcato, glabro, tenui, subæquilongis ; *spicis* 1-floris, 2-3-nis, ad nodos ramorum undulato-flexuosorum paniculæ subsessilibus, bractea parva vaginante carinata aristato-acuminata suffultis ; *squamis* lanceolatis, acutis, glabris ; *caryopsi* rotundata, subtrigona, glabra, versus apicem rugulosa.—C.P. 3226.

HAB. South of the island.

1½-2½-pedalis. *Panicula* aperta, 4-8 poll. longa, ramulis interne sulcatis scabrisque. *Squamæ* ferrugineæ. *Flores* triandri. *Antheræ* mucronatæ. *Stylus* elongatus. *Caryopsis* ochracea, denique nigro-fusca.

18. **BAUMEA**, Gaud. ; Kunth.

1. **B. crassa**, Thw. ; *foliis* crassis, parum flaccidis, interne spongiosis, glabris, acutis, margine obtusiusculis, *culmo* complanato foliato subæquilongis ; *spicis* pilosulis, trifloris (tertio flore terminali imperfecto), ad apices ramulorum paniculæ majusculæ glomerato-capitatis.—C.P. 845.

HAB. Ambagamowa District.

4-6-pedalis. *Folia* 3-4 lin. lata. *Panicula* pedalis vel ultra, parum aperta. *Bracteæ squamæque* ovato-lanceolatæ, ferrugineæ. *Flores* triandri. *Stylus* (cum stigmatibus 3, rarius 2, longis) nigro-ferrugineus, *basi dilatata* albida, albido-villosa, *caryopsi* ovoideæ nitidæ ferrugineæ adnata.

19. **SCLERIA**, Linn.

1. **S. oryzoides**, Pr. (N. ab Es. in Wight's Contr. p. 116 ; Steud. Synops. p. 169.)—C.P. 828.

HAB. Very common in the south of the island.

2. **S. stipularis**, N. ab Es. (Steud. l. c. p. 171, cum syn.?)—"C.P. 3031," C.P. 3763.

HAB. Allagalla, in the Central Province, at an elevation of 3000 feet.

*Vaginæ foliorum* tri-alatæ, antice infra apicem pilis deflexis hispidæ, alis latiusculis superne ultra vaginam productis, dorsali longiore et ad apicem extremum libera. *Discus* trilobus, lobis lanceolatis acutissimis.

The present plant agrees very well with Nees's description, so far as it goes ; but the locality he gives for his species (Surinam and Guyana) being so distant from this, renders it not unlikely that the Ceylon plant will prove to be distinct.

3. **S. Sumatrensis**, Retz. (N. ab Es. in Wight's Cat. p. 116 ; Kth. Enum. ii. p. 357 ; Steud. Synops. Cyperac. p. 171.)—C.P. 3783.

HAB. Ambagamowa District.

4. **S. elata**, Thw. ; *culmis* elatis, triquetris ; *foliis* longiusculis, valde acuminatis, margine scaberrimis ; *vagina* exalata ; *ligula* brevi, rotundata, parum pilosa ; *paniculis* deltoideis, plurifloris ; *bracteis* subulatis. basin versus subito dilatatis pilosisque ; *caryopsi* ovoidea, apiculata ; *hypogynio* mediocri subtrilobo vel triangulari, angulis rotundatis.—*Scleria*, No. 15, in Herb. Ind. Orient. Hook. f. et Thoms.

Var. *a* ; *caryopsi* prominentim reticulato.—C.P. 3030, 825.

Var. *β* ; *caryopsi* sublævi.—C.P. 3032.

HAB. Not uncommon up to an elevation of 5000 feet.

5. **S. androgyna**, N. ab Es. l. c. p. 117 ; Kth. l. c. ; Steud. l. c. p. 168.—C.P. 3319.



HAB. Reigam Corle.

6. *S. lævis*, Retz. (N. ab Es. l. c.; Kth. l. c. p. 342; Steud. l. c. p. 169.)—C.P. 2745.

HAB. South of the island.

7. *S. tessellata*, Willd. (N. ab Es. l. c. p. 118, cum syn.; Kth. l. c. p. 343; Steud. l. c.)—C.P. 3034.

Var.  $\beta$ , *squamis* longioribus angustioribusque; *caryopsi* glabra et grossius cancellata quam in var. *a*.—*S. uliginosa*, Hochst. *S. parvula*, Steud. l. c. p. 174.—C.P. 3033.

HAB. Var. *a*, Ambagamowa and Saffragam Districts; var.  $\beta$ , near Galle.

*Flores masc.* 2-3-andri.

8. *S. hebecarpa*, N. ab Es. l. c.; Kth. l. c. p. 357; Steud. l. c.—C.P. 3318.

HAB. Warmer parts of the island.

*Spicæ fœm.* 1-floræ. *Flores masc.* 1-andri. *Caryopsis* hypogynio trilobo, lobis rotundatis, obtusis, subtus excavatis.

9. *S. lithosperma*, Willd. (haud N. ab Es.)—Kth. l. c. p. 349.—C.P. 826.

Var.  $\beta$ , *caryopsi* a basi fere ad apicem transversim rugoso-tuberculata, ferrugineo conspersa.—C.P. 2627.

HAB. Common in the hotter parts of the island.

*Flores masc.* monandri.

10. *S. pergracilis*, Kth. l. c. p. 354; Steud. l. c. p. 176.—*Hypoporum pergracile*, N. ab Es. l. c. p. 118.—C.P. 827.

HAB. South of the island, Gardner.

11. *S. Neesii*, Kth. l. c. p. 358; Steud. l. c. p. 175.—*Hypoporum capitatum*, N. ab Es. l. c.—C.P. 3037.

HAB. Common in the hotter parts of the island.

*Spicæ masc.*: *squamis* (in meis specim.) integris, lanceolatis, acutis, nec apice fimbriato-laceris; *floribus* triandris.

12. *S. junciformis*.—*Cylindropus junciformis*, N. ab Es.; Kth. l. c. p. 360; Steud. l. c. p. 180.—C.P. 3225.

HAB. Caltura, Moon.

There is only a single imperfect specimen of the present plant in this Herbarium, and I can discover nothing in its structure to justify its separation from *Scleria*.

## 20. DIPLACRUM, R. Br.

1. *D. caricinum*, R. Br. Prod. 241; Kth. Enum. ii. p. 360, cum syn.—*D. Zeylanicum*, N. ab Es. in Wight's Contr. p. 119.—C.P. 3035.

HAB. Common in the warmer parts of the island.

*Spicula fœm.* uniflora, sæpius brevi-stipitata; *squamis* oppositis, 2 (haud 4, ut N. ab Es. describ., cujus "squamæ inferiores" certe bracteæ sunt, una vel utraque spiculam masc. in axilla gerente). *Flores masc.* (in meis specim.) 1-2-andri.

## 21. CAREX, Linn.

\* *Spica solitaria, androgyna.*

1. *C. rara*, Boott in Linn. Trans. xx. p. 139; Illustrations of the genus *Carex*, p. 44, t. 109.—C.P. 3080.

HAB. Newera Ellia and Horton Plains, at an elevation of 6000 feet.

\*\* *Spicis 2 vel pluribus, androgynis.*

2. **C. nubigena**, Don, Fl. Nep. p. 42; N. ab Es. in Wight's Contr. p. 120, cum syn.; Kth. Enum. ii. p. 385; Boott, Illustr. Car. p. 1, t. 2.—C.P. 2395.

HAB. Newera Ellia and other of the more elevated parts of the Central Province.

3. **C. spicigera**, N. ab Es. l. c. p. 121; Kth. l. c. p. 512; Boott, l. c. p. 10, t. 24.—C.P. 822; et *forma minor*, C.P. 824; *forma perigyniis angustioribus, subrostratis*, C.P. 2629.

Var.  $\beta$ ; *spicis omnibus ad apicem culmi conglomeratis*.—*C. rubella*, Boott.—C.P. 2966.

HAB. Very abundant in the Central Province, up to an elevation of 6000 feet; *forma minor*, Pedrotallagalla, Gardner, M. Wichura.

4. **C. leucantha**, Arn. MSS. (Boott, l. c. p. 10, t. 28.)—C.P. 2631.

HAB. Not very uncommon up to an elevation of 2000 feet.

5. **C. gracilis**, R. Br. Prod. 242; Boott, l. c. p. 59, t. 154, cum syn.—*C. lachnosperma*, Wall. N. ab Es. l. c. p. 124. *C. flexilis*, Don. *C. lenta*, Don; Spreng. Syst. iii. p. 811.—C.P. 2632.

HAB. Newera Ellia and Horton Plains.

6. **C. longipes**, Don, Fl. Nep. p. 43; N. ab Es. l. c.; Kth. l. c. p. 418.—C.P. 2749.

HAB. Maturatta and Hewahette Districts, rare.

7. **C. Bengalensis**, Roxb. Fl. Ind. iii. p. 572; Boott, l. c. p. 85, tt. 240, 243.—*C. valida*, N. ab Es. l. c. p. 123. An *C. condensata*, N. ab Es. l. c. ?—C.P. 2628.

HAB. Common in the Central Province, up to an elevation of 4000 feet.

8. **C. cruciata**, Wahl. (Willd. Sp. Pl. iv. p. 248; N. ab Es. l. c.)—*C. filicina*, N. ab Es. l. c. *C. Lindleyana*, N. ab Es. l. c. *C. vacua*, Boott, Herb. Ind. Orient. Hook. f. and Thoms. (conf. *C. ramosa*, Schk., *C. meiogyna*, N. ab Es., *C. raphidocarpa*, N. ab Es., et *C. thyrsiflora*, Boott.)—C.P. 820, 3161.

HAB. Very abundant at an elevation of 3000 to 6000 feet.

An extremely variable plant, and, at a superficial glance, nothing can well look more distinct than the luxuriant form with spreading panicle, growing in damp shady places, and the smaller form with constricted panicle and larger flower-scales, growing on exposed downs; these forms, however, so merge into one another that I find it impossible to separate them into distinctly defined varieties.

9. **C. baccans**, N. ab Es. l. c. p. 122; Boott, l. c. p. 83, tt. 234–239; Kth. l. c. p. 513.—C.P. 821.

HAB. Common in the Central Province, up to an elevation of 6000 feet.

10. **C. Walkeri**, Arn. MSS. (Boott, l. c. p. 2, t. 4.)—C.P. 2751.

HAB. More elevated parts of the Central Province, not very common.

\*\*\* *Spicis unisexuis, vel intermediis androgynis, apicali mascula.*

11. **C. maculata**, Boott. (Linn. Trans. xx. p. 128; Illustr. Car. p. 9 t. 26.)—C.P. 2630.

HAB. Horton Plains, and in the more elevated parts of the Ambagamowa District.

12. *C. phacota*, Spr. (Syst. iii. p. 826; N. ab Es. l. c. p. 126; Boott, Illustr. Car. p. 63, t. 168, cum syn.)—C.P. 2965.

HAB. Not uncommon in the higher parts of the Central Province.

13. *C. Arnottiana*, Drejer, Symb. t. 5.—C.P. 3219.

HAB. Newera Ellia and other of the more elevated parts of the Central Province, common.

3-4-pedalis, foliis culmo subæquilongis. Spicæ cylindricæ, numerosæ, plerumque longiuscule pedunculatæ, intermediæ androgynæ. Squamæ oblongæ, subacutæ, fusco-purpureæ, carina cum apice cuspidato glabro viridi. Stigmata 2. Perigynium squama subæquilongum, ovoideum, compressum, fusco-ferrugineum, dimidio superiore scabriusculum, rostro brevi cylindrico truncato.

14. *C. lobulirostris*, Drejer, Symb. t. 14.—*C. Arnottiana*, Boott. Ann. Nat. Hist. xvii. p. 56.—C.P. 2633.

HAB. Horton Plains and Newera Ellia, at an elevation of 6000 to 7000 feet.

2-8-pedalis, foliis culmo longioribus. Bractæ longæ, basi vaginantes. Spicæ 4-5, cylindricæ, contiguæ, bipollicares, terminalis mascula, reliquæ fœmineæ, inferior pedunculata. Squamæ lanceolatæ, nigro-sanguineæ, medio trinerviæ fusco-virides, aristato-acuminatæ, acumine scabro. Stigmata 3, brevia. Perigynium ampulliforme, nervosum, fuscum, rostro truncato demum parum fisso.

15. *C. Jackiana*, Boott, Illustr. Car. p. 9, t. 25.—*C. instabilis*, Boott, in Herb. Ind. Or. Hook. f. and Thoms.—C.P. 3198. Var.  $\beta$ ; *breviculmis*.—C.P. 3781.

HAB. Horton Plains. Var.  $\beta$ . Ambagamowa District.

16. *C. breviculmis*, R. Br. (Prod. 242; Kth. l. c. p. 522.)—C.P. 2750.

HAB. Maturatta and Hewahette Districts, at an elevation of 4000 to 5000 feet.

1½-2-pedalis, bracteis longis, foliaceis. Spicæ fœm. 2-4, plus minus contiguæ, squamis late ovatis, acutis, brevicuspidatis, pallide ferrugineis carina viridi, quam perigynium tomentosum ovoideum rostratum multo brevioribus. Spica masc. squamis muticis.

The present plant agrees very well with Brown's description of his *C. breviculmis*, but not having an authentic specimen of the latter I cannot be quite certain of their identity. Can *C. Royleana*, N. ab Es., be a starved form of this species?

## CLXVII. GRAMINEÆ.

### 1. LEERSIA, Soland.

1. *L. hexandra*, Sw. Fl. Ind. Occ. i. p. 131; Kunth, Enum. i. p. 6, —C.P. 877.

HAB. Very common up to an elevation of 4000 feet. Nom. vulg. "Layoo."

### 2. HYGRORYZA, N. ab Es.

1. *H. aristata*, N. ab Es. in Edinb. Phil. Journ. (1833) p. 380; Steud. Synops. Gram. p. 5.—*Potamochoa Retzii*, Griff. Notulæ, pars

3, p. 3, t. 140. *Leersia aristata*, Roxb. Fl. Ind. ii. p. 207.—C.P. 878.

HAB. Hotter parts of the island. Nom. vulg. "Gojabbā."

### 3. ORYZA, Linn.

1. *O. sativa*, Linn. (Kth. Enum. i. p. 7, cum syn.)—C.P. 2876.

HAB. Not uncommon in wet places, and apparently truly wild. Nom. vulg. "Goyang."

Numerous varieties of this plant are extensively cultivated by the Cinghalese.

### 4. LEPTASPIS, R. Br.

1. *L. urceolata*, Br. et Benn. in Plant. Jav. Rar. p. 23, t. 6.—*L. Zeylanica*, N. ab Es. MSS.; Steud. Syn. Gram. p. 8.—C.P. 972.

HAB. Shady places in the warmer parts of the island, not very uncommon.

FLOR. MASC.: (in meis specim.) *Ovarii* rudimento minuto. FLOR. FÆM.: *Stamina* abortiva 6, minutissima.

2. *L. cochleata*, Thw.; *foliis* oblongis vel obovato-oblongis, acuminatis, glabris; *glumis* rotundatis, cuspidato-acuminatis, persistentibus. FLOR. FÆM.: *palea inferiore* incurvata, valde inflata, 5-costata, uncinato-pilosa.—C.P. 896.

HAB. Forests of the Central Province, up to an elevation of 4000 feet.

*Culmi* 2-3-pedales, basi procumbentes radicanterque. *Folia* 4-6 poll. longa, 1½-1¾ poll. lata, *vagina* 2-3 poll. longa, ad apicem, cum petiolo semipollicari interne, pilosa. *Paniculæ* rami verticillati, terui, ramulis alternis spiculam alteram masc. terminalem pedicellatam et alteram fœmin. sessilem minute bracteatum gerentibus. *Squamulæ* 2, breves, crassiusculæ, truncatæ. FLOR. MASC.: *palea inferior* rotundata, gradatim brevi-acuminata, *superior* linearis, apice bifida. *Ovarii rudimentum* stamin. abortiv. 6, minutissimis cinctum. *Stylus* linearis, basi inflexus. *Stigmata* 3, plumosa, stylo parum breviora.

This species differs from the genus, as characterized by Brown, in the flowers having squamulæ.

### 5. COIX, Linn.

1. *C. Lachryma*, Linn. (Roxb. Fl. Ind. iii. p. 568; Kth. Enum. i. p. 20.)—C.P. 942.

HAB. Very common in cultivated ground. Nom. vulg. "Keekirindee."

### 6. CHIONACHNE, R. Br.

1. *C. Koenigii*.—*C. barbata*, Br. et Benn. in Pl. Jav. Rar. p. 18. *Coix Koenigii*, Spr. Syst. i. p. 239. *C. barbata*, Roxb. Fl. Ind. iii. p. 569. *C. arundinacea*, Koen.; Willd. Sp. Pl. iv. p. 203 (haud Lam.)—C.P. 3137.

HAB. Central Province, not very uncommon.

### 7. PASPALUM, Linn.

1. *P. scrobiculatum*, Linn. (Kth. Enum. i. p. 53, cum syn.; Roxb. Fl. Ind. i. p. 278.)—*P. Kora*, Willd.; Roxb. l. c. p. 279. *P. orbiculare*, Forst. Prod. vii.

Var. *a*, Kth. l. c.; *pedicellis* simplicibus, *glumis* 7-8-nerviis.—C.P. 863.

Var. *β*, Kth. l. c.; *pedicellis* simplicibus, *glumis* 3-nerviis.—C.P. 865.

Var. *γ*, Kth. l. c.; *pedicellis* in media rachi bipartitis, *glumis* 3-nerviis.—C.P. 2434.

HAB. Very abundant; var. *a* is cultivated by the Cinghalese. Nom. vulg. "Ammoo."

2. **P. Royleanum**, N. ab Es. MSS.

Var. *β*; *gluma* inferiore nulla.—C.P. 859.

HAB. Very common on the borders of paddy-fields.

3. **P. filiculme**, N. ab Es. MSS.—An *P. bifarium*, Edgew. in Journ. As. Soc. (1852) p. 187; Steud. Syn. Gram. p. 32? *Milium filiforme*, Roxb. l. c. p. 314.—C.P. 860 (2573).

HAB. Not uncommon in the Central Province.

## 8. ERIOCHLOA, Humb. et Kth.

1. **E. annulata**, Kth. Enum. i. p. 73; Benth. Flor. Hongk. p. 409.—*Milium ramosum*, Roxb. Fl. Ind. i. p. 315.—C.P. 3323.

HAB. Hotter parts of the island.

## 9. PANICUM, Linn.

1. **P. sanguinale**, Linn. (Kth. Enum. i. p. 82, cum syn.) Var. *β*; australe.—*P. australe*, Spr.; Kth. l. c. p. 85, cum syn. *P. Ægyptiacum*, Retz; Kth. l. c. p. 83, cum syn.—C.P. 856, 857, 861.

HAB. Extremely abundant.

2. **P. ciliare**, Retz. (Kth. l. c. p. 82, cum syn.)—*P. Neesii*, Kth. l. c. p. 84.—C.P. 862.

HAB. Common in sandy places.

3. **P. Wallichianum**, W. U. (Steud. Synops. Gram. p. 41.)—C.P. 3249.

HAB. More elevated parts of the Central Province.

4. **P. Helopus**, Trin. (Steud. l. c. p. 57.)—*Urochloa pubescens*, Kth. l. c. p. 74, cum syn.—Var. *β*; spiculis glabris, Munro, MSS.—C.P. 3244.

HAB. South of the island.

5. **P. Javanicum**, Poir. (Steud. l. c. p. 61.)—*Urochloa panicoides*, Beauv.; Kth. l. c. cum syn.—C.P. 858.

HAB. Common in the hotter parts of the island.

6. **P. semialatum**, R. Br. i. 192.—*Urochloa semialata*, Kth. l. c.—C.P. 3239.

HAB. Oova District, up to an elevation of 5000 feet.

7. **P. cimicinum**, Retz, Obs. iii. p. 9.—*Urochloa cimicina*, Kth. l. c. cum syn.—C.P. 911.

HAB. Very abundant.

8. **P. Burmanni**, Retz, Obs. iii. p. 10; Steud. l. c. p. 44, cum syn.—*Oplismenus Burmanni*, Beauv.; Kth. l. c. p. 139. Var. spiculis glabris.—C.P. 3683.

HAB. Central Province, under the shade of trees.

9. **P. compositum**, Linn.—*Oplismenus compositus*, R. and S.; Kth. l. c. p. 141, cum syn.—c.P. 913.

HAB. Common throughout the island.

An extremely variable species, some examples being nearly glabrous, and others densely pubescent.

10. **P. Crus-galli**, Linn.—*Oplismenus Crus-galli*, Kth. l. c. p. 143, cum. syn. *P. stagninum*, Koen. in Willd. Sp. Pl. i. p. 337. *Oplism. stagninus*, Kth. l. c. p. 144, cum syn.—c.P. 900, 3237.

Var.  $\beta$ ; *minor*.—*P. Colonum*, Linn. *Oplism. Colonus*, H. and K.; Kth. l. c. p. 142, cum syn.—c.P. 901.

HAB. Very common in the hotter parts of the island.

11. **P. fuitans**, Retz. (Willd. Sp. Pl. i. p. 338; Roxb. Fl. Ind. i. p. 293; Kth. l. c. p. 78.)—*P. affine*, N. ab Es.—c.P. 864, 897 (898).

HAB. Abundant in wet places in the hotter parts of the island.

12. **P. brizoides**, Linn. (Jacq. Ecl. i. t. 2; Kth. l. c. cum syn.)—*P. flavidum*, Retz; Kth. l. c. cum syn.—c.P. 902.

HAB. Very common in the warmer parts of the island.

13. **P. cruciforme**, Sibth. (Kth. l. c. p. 78.)—*P. Caucasianum*, Trin. *P. Wightii*, N. ab Es.; Steud. l. c. p. 58.—c.P. 903.

HAB. Not uncommon in cultivated ground.

14. **P. distachyum**, Linn. (Kth. l. c. p. 91, cum syn.)—c.P. 904.

HAB. Very abundant.

15. **P. prostratum**, Lam. (Kth. l. c. p. 89, cum syn.; Steud. l. c. p. 62, cum syn.)—c.P. 905. Forma major, c.P. 3236.

HAB. Common in the hotter parts of the island.

16. **P. coccospermum**, Steud. l. c. p. 62.—*P. vestitum*, N. ab Es. (haud Kth.)—c.P. 3241.

HAB. Central Province, up to an elevation of 6000 feet.

17. **P. uncinatum**, Raddi. (Trin. Ic. t. 216; Steud. l. c. p. 60.)—*Echinolæna? polystachya*, H. and K.; Kth. l. c. p. 172, cum syn.—c.P. 710.

HAB. Common in the Central Province in damp, shady places.

18. **P. ovalifolium**, Poir. (Kth. l. c. p. 113, cum syn.; Benth. Flor. Hongk. p. 413.)—c.P. 889.

HAB. Common in the warmer parts of the island.

*Rhachis paniculæ* plus minus patentim longe pilosa, *floribus apicalibus* longe albido-villosi.

19. **P. trigonum**, Retz, Obs. iii. p. 9; Willd. Sp. Pl. i. p. 352.—*P. radicans*, Retz, iv. p. 18; Willd. l. c. p. 354. *P. pilipes*, N. ab Es.—c.P. 100, 886, 887, 888, 891 (709).

HAB. Very abundant throughout the island.

An extremely variable species as regards its size and hairiness. All the forms have the flat callous spot at the apex of the inferior palea of the fertile flower.

20. **P. Petiverii**, Trin. (N. ab Es. in Mart. Bras. ii. p. 111; Kth. l. c. p. 91, cum syn.)—c.P. 895.

HAB. Damboul, Gardner.

21. **P. Gardneri**, Thw.; culmis ramosis, basi repentibus, radicanti-

bus; *foliis* lanceolatis, acuminatis, basi rotundatis, subcordatis; *pani-  
cula* aperta; *spiculis* majusculis, solitariis, longe pedicellatis; *glumis*  
ovato-lanceolatis, acutis, *inferiore* 5-nervia, parum brevior, *superiore*  
7-nervia; *flor. ster.* bipaleaceo, *palea inferiore* gluma superiore con-  
formi; *flor. hermaphr.* obtuso, eadem parum brevior.—c.p. 894.

HAB. Forests of the Central Province, at an elevation of 4000 to 6000 feet.

Tota planta villosula vel subglabra, sæpius purpurascens. Culmi  $\frac{1}{2}$ -2-pedales. Folia  $1\frac{1}{2}$ -5 poll. longa, 4-10 lin. lata, vagina ad marginem superne villosa, ligula brevi, erosa. Panicula 3-6 poll. longa. Spicula  $1\frac{1}{2}$ -2 lin. longæ.

This species bears a very great general resemblance to *Isachne Walkeri*, W. et A., but is quite distinct from it. The much larger spikelets of the present distinguish it at once from *P. montanum*, Roxb.

22. **P. montanum**, Roxb. Fl. Ind. i. p. 313; Kth. l. c. p. 126; Benth. Flor. Hongk. p. 412.—*P. Courtallense*, N. ab Es.; Steud. l. c. p. 83. *P. euchroum*, Steud. l. c. p. 98, cum syn.—c.p. 892.

HAB. Central Province, not very common.

23. **P. antidotale**, Retz, Obs. iv. p. 17; Kth. l. c. p. 125; Steud. l. c. p. 77.—c.p. 971.

HAB. Hotter parts of the island.

Used medicinally by the Cinghalese.

24. **P. plicatum**, Lam. (Kth. l. c. p. 94.)—*P. Nepalense*, Spr.; Kth. l. c. *P. plicatum*, *P. nervosum*, et *P. costatum*, Roxb. Fl. Ind. pp. 311, 312. *P. neurodes*, Schultz.—c.p. 890.

HAB. Not uncommon up to an elevation of 2000 feet.

25. **P. miliaceum**, Linn. (Roxb. l. c. p. 310; Kth. l. c. p. 104.)—*P. miliare*, Lam. (Roxb. l. c. p. 309; Kth. l. c.)—c.p. 882, 893.

HAB. Very common. Nom. vulg. "Mainairee."

26. **P. psilopodium**, Trin. (Kth. l. c. p. 100, cum syn.)—c.p. 236. (An *P. miliacei* var.? Hook. f.)

HAB. Equally common with the preceding, and also cultivated by the Cinghalese under the same name.

27. **P. repens**, Linn. (Kth. l. c. p. 103, cum syn.)—*C. paludosum*, Roxb. l. c. p. 307. *P. ischæmoides*, Retz, Obs. 4, 17.—c.p. 883 (3049).

HAB. Very abundant. Nom. vulg. "Attora."

Much valued by the Cinghalese as a good fodder for cattle, but a troublesome weed in gardens.

28. **P. humile**, N. ab Es. (Steud. l. c. p. 84.)—c.p. 3243.

HAB. South of the island.

29. **P. Leptochloa**, N. ab Es. (Steud. l. c. p. 62.)—c.p. 918.

HAB. Not uncommon up to an elevation of 2000 feet.

30. **P. nodosum**, Kth. l. c. p. 97, cum syn.; Benth. Flor. Hongk. p. 412, cum syn.—*P. Arnottianum*, N. ab Es.; Steud. l. c. p. 59.—c.p. 964.

HAB. Common in damp shady places in the warmer parts of the island.

31. **P. curvatum**, Linn. (Steud. l. c. p. 67, vix Roxb. Fl. Ind. p. 286.)—c.p. 3240.

HAB. Hotter parts of the island.

*Gluma superior* basi saccata.

32. *P. auritum*, Presl, N. ab Es. (Kth. l. c. p. 113.)—c.p. 3242.  
HAB. Saffragam District, at no great elevation.

33. *P. Myrus*, Lam. (Kth. l. c. p. 86, cum syn.)—*P. serrulatum*, Roxb. l. c. p. 307.—c.p. 3238.

HAB. In hotter places in the hotter parts of the island.

I can find no sufficient specific difference between the Ceylon plant and a specimen of *P. Myrus*, from Guiana.

34. *P. interruptum*, Willd. (Kth. l. c. p. 87, cum syn.; Roxb. l. c. p. 289.)—c.p. 910.

HAB. Common in wet places in the warmer parts of the island.

35. *P. Indicum*, Linn. (Kth. l. c. p. 133; Benth. Flor. Hongk. p. 413, cum syn.)—*P. angustum*, Trin., et *P. contractum*, N. ab Es.; Steud. l. c. p. 84.—c.p. 885, 908, 909.

HAB. A very abundant species.

36. *P. glaucum*, Linn.—*Setaria glauca*, Beauv.; Kth. l. c. p. 149, cum syn.—c.p. 907.

HAB. Very common throughout the island. Nom. vulg. "Kāooloo."

37. *P. intermedium*, Rottb. Nov. Sp. p. 47.—*Setaria intermedia*, R. S.; Kth. l. c. p. 150.—c.p. 906.

HAB. Not uncommon in damp, shady places.

38. *P. verticillatum*, Linn.—*Setaria verticillata*, Beauv.; Kth. l. c. p. 152, cum syn.—c.p. 3246.

HAB. South of the island, apparently not very common.

*Panicum jumentorum*, Pers., and *P. barbinode*, Trin., are largely cultivated as fodder for horses and cattle. *P. italicum*, Linn., is cultivated by the natives as a food grain, as is also *Penicillaria cylindrica*.

## 10. ICHNANTHUS, Beauv.

1. *I. pallens*, Munro. (Benth. Fl. Hongk. p. 414, cum syn.)—*Panicum pallens*, Sw.; Kth. Enum. i. p. 89, cum syn.—c.p. 2755.

HAB. Deltotte, in the Central Province, at an elevation of 4000 feet.

## 11. STENOTAPHRUM, Trin.

1. *S. complanatum*, Schr. (Kth. Enum. i. p. 137, cum syn.)—*Rottbællia complanata*, Sw.—*Panicum dimidiatum*, Linn.; Roxb. Fl. Ind. i. p. 287.—c.p. 912.

HAB. Not uncommon in the hot, drier parts of the island.

## 12. ISACHNE, R. Br.

1. *I. Walkeri*, W. et A. (Steud. Synops. Gram. p. 97.)—c.p. 282.

HAB. Not very uncommon in forests of the Central Province.

2. *I. australis*, R. Br. (Prod. 196; Kth. Enum. i. p. 136; Benth. Fl. Hongk. p. 414, cum syn.)—*I. miliacea*, Rottb. Nov. Sp. p. 58. *Panicum atrovirens*, Trin.; Kth. l. c. p. 127. *P. Batavicum* et *P. Benjamini*, Steud. l. c. p. 96.—c.p. 880.

Var.  $\beta$ . *multiflora*; *elatior*; *panicula* majore, *multiflora*; *spiculis*



minoribus, glabris; *flore sterili* quam *hermaphr.* fere duplo longiore.—  
C.P. 3671.—Forsan species distincta.

HAB. Very abundant. Var.  $\beta$ . Oova District, at an elevation of 5000 feet.

3. **I. Kunthiana**, W. et A. (Steud. l. c. p. 96.)—*I. Neesiana*, Herb. Arn. *Panicum cuspidigluma*, Steud. l. c.—C.P. 881 (314).

HAB. More elevated parts of the Central Province.

*Glumæ* subrostrato-acuminatæ, glabræ vel versus apicem rigide pilosæ, floribus inter se æquilongis longiores.

A variable plant as regards size of the leaves and amount of pubescence. Possibly a variety only of the preceding species, which it greatly resembles.

4. **I. pulchella**, Roth, Nov. Sp. p. 58; Benth. Fl. Hongk. p. 414.—*Panicum Malaccense*, Trin.; Kth. l. c. p. 113. *Grayia elegans*, N. ab Es.; Steud. l. c. p. 119, cum syn.—C.P. 879.

HAB. Hantani, *Macrae*; Reigam Corle.

### 13. SPINIFEX, Linn.

1. **S. squarrosus**, Linn. (Kth. Enum. i. p. 175, cum syn.)—C.P. 947.

HAB. Common on the sand of the sea-coast.

### 14. THOUAREA, Pers.

1. **T. sarmentosa**, Pers. Syn. i. p. 110. (Kth. Enum. i. p. 174; Benth. Fl. Hongk. p. 415.)—C.P. 3260.

HAB. Sea-coast near Tangalle.

### 15. TRACHYS, Pers.

1. **T. mucronata**, Pers. Syn. i. p. 85. (Kth. Enum. i. p. 168, cum syn.)—*Panicum squarrosum*, Retz; Willd.; Roxb. Pl. Cor. t. 206; Roxb. Fl. Ind. i. p. 288.—C.P. 3262.

HAB. Jaffna, *Gardner*.

### 16. LAPPAGO, Schreb.

1. **L. racemosa**, Willd. (Kth. Enum. i. p. 170, cum syn.)—C.P. 3682.

HAB. Trincomale, *Rev. S. O. Glenie*.

### 17. ARUNDINELLA, Raddi.

1. **A. nervosa**, N. ab Es. (Steud. Synops. Gram. p. 115.)—*A. agrostoides*, Trin.; Steud. l. c. p. 116. *Holcus nervosus*, Roxb. Fl. Ind. i. p. 318.—C.P. 919.

HAB. Common in the Central Province, up to an elevation of 5000 feet.

2. **A. villosa**, W. et A. (Steud. l. c. p. 115.)—C.P. 960.

HAB. More elevated parts of the Central Province.

*Culmi* basi dense albido-tomentosi.

3. **A. avenacea**, Munro, MSS.—C.P. 3471.

HAB. Saffragam District, at no great elevation.

*Culmi* debiles, ramosi, radicanes. *Folia* parva, ovato-lanceolata, sparse pilosula,

dimidio inferiore distanter rigide ciliata, basi cordata, amplexicaulia. *Panicula* pollicaris, compacta. *Glumæ* lanceolatae, apice setaceae, inferior superne hispida. *Flos hermaphr.* pilosus, arista valida, complanata, torta, supra medium attenuata munitus.

### 18. GARNOTIA, Brongn.

*Miquelia*, N. ab Es.; *Berghausia*, Endl.

1. **G. stricta**, Brongn. (Benth. Fl. Hongk. p. 416, cum syn.)—*Berghausia mutica*, Munro, in Proc. Am. Acad. 4, p. 362.—c.p. 2968.

HAB. More elevated parts of the Central Province.

Much used for thatching.

2. **G. scoparia**.—*Berghausia scoparia*, Munro.—c.p. 943.

HAB. Hotter parts of the island, not uncommon.

*Culmi* 2-4-pedales, erecti. *Folia* angusta, *vaginis* latis, distichis, margine longe sericeo-pilosis. *Panicula* densa, 1-2-pedalis, ramis plurimis, longiusculis, suberectis. *Spiculæ* 1½ lin. longæ, basi pilis brevibus paucis cinctæ. *Gluma* anguste lanceolata, setaceo-acuminata, scabriuscula. *Palea inferior flor. hermaphr.* acuta, longe aristata.

3. **G. fuscata**, Thw.; *culmis* tenuibus, basi repentibus, ramosis, radicantibus, nodis glabris; *foliis* firmis, brevibus, lineari-lanceolatis, sparsissime longiuscule pilosis, vagina glabra apice pilosa; *panicula* parva, pauciflora; *pedicellis* apice piliferis; *gluma* subglabra, lanceolata, apice longiuscule setacea; *palea inferiore flor. hermaphr.* longe aristata.—c.p. 2756.

HAB. Saffragam District, at an elevation of 4000 feet.

*Culmi* 1-2-pedales. *Folia* 1-2 poll. longa, circiter 2 lin. lata.

4. **G. Courtallensis**.—*Miquelia Courtallensis*, Arn. and N. ab Es.; Steud. Synops. Gram. p. 119.—c.p. 454.

HAB. Central Province, at an elevation of 4000 feet.

Tota planta pallida vel plus minus nigrescens. *Culmi* ½-2-pedales. *Folia* pilosa vel glabra, 1-4 poll. longa, 1-3 lin. lata. *Spiculæ* 1½ lin. longæ. *Gluma* pilosa, acuta nec setacea.

5. **G. micrantha**, Thw.; *culmis* tenuibus, basi repentibus, radicantibus, nodis pilosis; *foliis* lineari-lanceolatis, valde acuminatis, breve pilosulis vel subglabris, vagina apice pilosa; *panicula* elongata, ramis capillaceis, scabris; *pedicellis* ad apicem parcissime piliferis; *spiculis* minutis, aliis aristatis, aliis muticis, vel omnino muticis; *gluma* scabra, acuta, apice setacea.—c.p. 944, 945.

HAB. Central Province, at an elevation of 2000 to 4000 feet.

*Culmi* 1-2½-pedales. *Folia* 3-8 poll. longa, 2-4 lin. lata. *Panicula* sæpe pedalis. *Spiculæ* 1 lin. longæ.

In general appearance very like *G. adscendens*, Munro, but the spikelets of the present plant are much smaller, and with scarcely a trace of the circle of hairs at their base.

### 19. OROPETIUM, Trin.

1. **O. Thomæum**, Trin. (Kth. Enum. i. p. 464, cum syn.)—*Rottbællia Thomæa*, Willd.; Roxb. Pl. Cor. ii. t. 133.—c.p. 872.

HAB. Bintenne Tank, Gardner.

### 20. HEMARTHRIA, R. Br.

1. **H. compressa**, R. Br. (Kth. Enum. i. p. 465, cum syn.)—*Rott-*

*bællia compressa*, Linn.; Roxb. Pl. Cor. ii. t. 156. *H. fasciculata*, Kth. l. c.—c.p. 3254.

HAB. Warmer parts of the island, not uncommon.

## 21. MNESITHEA, Kth.

1. *M. lævis*, Kth. Enum. i. p. 466, cum syn.—*Rottbællia lævis*, Retz; Willd. Sp. Pl. i. p. 465.—c.p. 873.

HAB. Hotter parts of the island.

## 22. ROTTBÆLLIA, Linn.; R. Br.

1. *R. exaltata*, Linn. (Kth. Enum. i. p. 466, cum syn.)—c.p. 2386.

HAB. Common in the hotter parts of the island.

2. *R. nigrescens*, Thw.; *culmis* suberectis, sæpe elongatis, nodis glabris; *foliis* lineari-lanceolatis, acuminatis, basi late subcordatis ciliatisque; *ligula* rotundata, lacerato-ciliata; *spicæ* axillares terminalesque, solitariae vel binæ; *gluma* exterior *spiculæ sessilis* ovata, rigida, tessellato-punctata, apice membranaceo-auriculata; *spicula pedunculata* parum brevior, epunctata, biflora, flore altero masculino, altero neutro.—c.p. 867.

HAB. Central Province, at an elevation of 4000 feet.

Siccitate nigrescens. *Folia* 4–8 poll. longa, 4–7 lin. lata, pilosa vel glabra, margine scabra. *Spicæ* 2–3-pollicares. *Spiculæ* 1½ lin. longæ.

Allied to *R. striata*, N. ab Es., but the pedicellate spikelet of the present species is much more developed, and the outer glume of the sessile one is wider and more entirely tessellated.

## 23. MANISURIS, Linn.

1. *M. granularis*, Sw. (Kth. Enum. i. p. 469, cum syn.; Roxb. Pl. Cor. ii. p. 118.)—c.p. 81.

HAB. Common in the hot, drier parts of the island.

## 24. APLUDA, Linn.

1. *A. aristata*, Linn. (Kth. Enum. i. p. 516, cum syn.; Steud. Syn. Gram. p. 403, cum syn.; Roxb. Fl. Ind. i. p. 324.)—*A. rostrata*, N. ab Es.; Steud. l. c.—c.p. 59, 3165, 3166.

HAB. Very abundant up to an elevation of 4000 feet.

## 25. ISCHÆMUM, Linn.

1. *I. muticum*, Linn. (Kth. Enum. i. p. 512, cum syn.)—c.p. 868, 3048.

HAB. Common in the hotter parts of the island.

2. *I. rugosum*, Salisb. (Willd. Sp. Pl. iv. p. 940; Gært. de Fruct. iii. p. 2, t. 181.)—*I. segetum*, Trin.; Steud. Synops. Gram. p. 376, cum syn. *Andropogon Tong-dong*, Steud. l. c. p. 375, cum syn.—c.p. 870.

HAB. Hotter parts of the island.

3. *I. barbatum*, Retz. (Willd. l. c.; Kth. l. c. p. 513, cum syn.; Benth. Fl. Hongk. p. 425.)—*Meoschium lodiculare*, N. ab Es. *M. Neesianum*, Arn., et *M. Meyenianum*, N. ab Es.; Steud. l. c. p. 375.—c.p. 700, 869, 994.

HAB. South of the island.

*Spiculæ* subglabræ vel plus minus pilosæ.

A coarser plant than the preceding, with the sessile *spiculæ* narrower in proportion to their length, and less regularly rugose. The two are, however, very closely allied.

4. **I. semisagittatum**, Roxb. Fl. Ind. i. p. 320; Steud. l. c. p. 376.—C.P. 2625.

HAB. Ambagamowa District.

*Folia* (in meis specimin.) tuberculata, pilosa. *Spicul. sess.* gluma inferior late 3-rugosa. *Spicul. pedicell.* pedicellus gluma inferiore plana striata parum brevior.

5. **I. conjugatum**, Roxb. Fl. Ind. i. p. 321; Kth. l. c. 513.—*Andropogon cordatifolius*, Steud. l. c. p. 376.—C.P. 3263.

HAB. Colombo, Mr. W. Ferguson.

6. **I. pectinatum**, Trin. (Steud. l. c. p. 369, cum syn.)—*Andropogon falcatus*, Steud. l. c.—C.P. 3322.

HAB. Hotter parts of the island, in wet ground.

7. **I. nervosum**.—*Hologamium nervosum*, N. ab Es. in Edinb. Phil. Journ. anno 1831, p. 185. *Andropogon nervosus*, Rottb.; Steud. l. c. p. 370. An *Andropogon striatus*, Klein, in Willd. Sp. Pl. iv. p. 903?—C.P. 3138. : *Setaria nervosum*

HAB. Oodoopusalawa in the Central Province, at an elevation of 3000 to 4000 feet.

## 26. SPODIOPOGON, Trin.

1. **S. obliquivalvis**, N. ab Es. in Pl. Meyen. 185; Benth. Fl. Hongk. p. 426, cum syn.—*Andropogon malacophyllus*, Hochs.; Steud. Synops. Gram. p. 372. *A. Macraei*, Steud. l. c. p. 377, cum syn. *A. Blumei* et *A. bifidus*, Steud. l. c. p. 373. An *Ischæmum geniculatum* et *I. tenellum*, Roxb. Fl. Ind. i. pp. 322, 323?—C.P. 875, 874, 3235, 3168, 3167.

HAB. Common throughout the island.

A very variable plant, and the extreme forms of it very different in appearance, but, from the examination of a large number of specimens, I feel satisfied they may be safely arranged under one specific name. The larger hairy form occurs at a considerable elevation on the hills.

## 27. APOCOPIS, N. ab Es.

1. **A. Wightii**, N. ab Es. MSS.—*Andropogon Courtallumensis*, Steud. Synops. Gram. p. 377.—C.P. 401.

HAB. Galagama, in the Central Province, at an elevation of 3000 to 4000 feet.

*Culmi* ramosi, 1-2-pedales. *Folia* angusta, acuminata, sparse tuberculata, longiuscule pilosa. *Spica* pollicaris, pallide fulva. *Rachis* uno latere pedicellis fulvo-pilosis. *Spicula sessilis* 1½ lin. longa, arista semipollicari. *Gluma inferior* incomplete 7-nervia, basi, cum margine rufescente, longiuscule fulvo-pilosa.

This would seem to be very closely allied to *A. Wrightii*, Munro, described in Bentham's Fl. Hongkongensis.

## 28. POGONATHERUM, Beauv.

1. **P. saccharoideum**, Beauv. (Benth. Flor. Hongk. p. 421.)—*P. cri-*

*nitum*, Trin.; Kth. Enum. i. p. 478, cum syn. *P. refractum*, N. ab Es.; Steud. Synops. Gram. p. 412.—C.P. 936.

HAB. Very common in the warmer parts of the island.

### 29. CHRYSOPOGON, Trin.

1. **C. aciculatus**, Trin.—*Andropogon acicularis*, Retz; Kth. Enum. i. p. 505, cum syn.—C.P. 917.

HAB. Extremely abundant in open ground, in the warmer parts of the island.

2. **C. Wightianus**.—*Andropogon Wightianus*, Steud. Synops. Gram. p. 395, cum syn.—C.P. 3248.

Var.  $\beta$ ; *leucantha*.—*Spiculis* minoribus, albidis, pilosioribus; *pedicellis* spicul. masc. brevibus, longe aureo-fulvo-pilosis.—C.P. 2954. An species distincta?

HAB. Doomberra District.

3. **C. Zeylanicus**.—*Raphis Zeylanica*, N. ab Es. MS. *Andropogon Zeylanicus*, Steud. l. c. p. 397.—C.P. 996.

HAB. Not uncommon up to an elevation of 6000 feet.

### 30. ANTHISTIRIA, Linn.

1. **A. ciliata**, Retz, Obs. iii. p. 11; Kth. Enum. i. p. 481, cum syn.—C.P. 3257.

Var.  $\beta$ ; *major*.—*A. arguens*, Willd.; Kth. l. c. p. 482, cum syn.—C.P. 962.

HAB. Common up to an elevation of 4000 feet.

2. **A. tremula**, N. ab Es. (Steud. Synops. Gram. p. 401.)—C.P. 961.

HAB. Very abundant throughout the island. Non. vulg. "Painee-bāroo."

3. **A. fasciculata**, Thw.; *culmis* suberectis, glabris; *foliis* linearibus, margine scabris, sæpius villosulis; *pedunculis* axillaribus, 2-5-nis, folio subæquilongis, singulis spicas duas 7-stachyas gerentibus: *spiculis masculis* 6, lanceolatis, pilosulis, pedicellatis, 2 superiorum gluma interiore apice setacea; *spicula hermaphr.* glumis parum truncatis, inferiore dense pilosa; *arista* longissima, valde robusta, aureo-pilosa.—C.P. 940.

HAB. Badulla District, up to an elevation of 3000 feet.

*Culmi* graciles, 2-pedales, uno latere sulcati. *Pedunculi* 3-pollicares, filiformes, infra medium articulati, bibracteati, bractea basali hyalina, superiore sæpe ad apicem pedunculi attingente. FLOR. MASC.: bipaleacei.

4. **A. heteroclita**, Roxb. Flor. Ind. i. p. 249.—*Andropogon heteroclitus*, N. ab Es.; Steud. l. c. p. 389.—C.P. 963.

HAB. Common in the more elevated parts of the Central Province.

### 31. ANDROPOGON, Linn.

1. **A. Halepensis**, Sibth. (Kth. Enum. i. p. 502, cum syn.—C.P. 2484.

HAB. Not uncommon in the Central Province.

2. **A. tropicus**, Spr. (Kth. l. c. p. 503, cum syn.; Benth. Flor. Hongk. p. 423.)—*Holcus fulvus*, R. Br., and var. *mutica*.—C.P. 419.

HAB. Badulla District.

3. **A. venustus**, Thw.; *culmis* erectis, altiusculis, basi albido-pilosis; *foliis* longis, linearibus, margine scabris, vagina ad apicem pilosa; *panicula* majuscula, decomposita, aperta, articulis pedicellisque ad apicem piliferis; *spiculis* anguste lanceolatis, acuminatis, subglabris, *sessili* hermaphr. longe aristata, *pedicellata* mascula, bipaleacea, mutica.—C.P. 2875.

HAB. Rambodde, in the Central Province, at an elevation of 4000 feet.

*Panicula* purpurascens, pedalis vel ultra, ramis capillaribus, distanter subverticillatis, plus minus pendulis. *Spiculæ*  $1\frac{1}{2}$ –2 lin. longæ, *mascula* brevi-pedicellata, *palea* inferiore aliquando apice setacea.

4. **A. Martini**, Roxb. Fl. Ind. i. p. 277; Kth. l. c. p. 494; Benth. Fl. Hongk. p. 424.—*A. flexuosus*, N. ab Es.; Steud. Synops. Gram. p. 388.—C.P. 2733.

HAB. Central Province, covering considerable areas at an elevation of 2000 to 4000 feet. Nom. vulg. "Maana."

From the leaves of a cultivated variety of this species (nom. vulg. "Pangaremaana") is distilled the "citronella oil" of commerce. "Lemon oil" is derived from *Andropogon Schoenanthus*, Linn., which is considered not to be indigenous to the island, and which rarely produces flowers.

5. **A. versicolor**, N. ab Es. (Steud. l. c. p. 388.)—C.P. 3135.

HAB. More elevated parts of the Central Province.

The inflorescence of this species has, when crushed, a rather agreeable aromatic odour. The essential oil appears to be situated principally at the base of the spikelets.

6. **A. distans**, N. ab Es. (Steud. Synops. Gram. p. 387.)—C.P. 3784.

HAB. Newera Ellia.

7. **A. lividus**, Thw.; *cæspitosus*, *culmis* erectis, rigidiusculis, nodis glabris; *foliis* lineari-lanceolatis, acuminatis, glabris, ligula majuscula oblonga; *pedunculis* terminalibus axillaribusque, 1–4; *spicis* binis, subsessilibus; *rachi* *pedicellis*que longe ciliatis; *spiculis* glabris, lanceolatis; *spiculæ sessilis gluma inferiore* supra medium ad nervos 2 laterales alata, *flore hermaphr.* aristato; *spiculæ pedicell. gluma inferiore* vix alata, *flore* masculo, unipaleaceo.—C.P. 953.

HAB. Newera Ellia and other of the more elevated parts of the Central Province.

Siccitate fuscescens. *Culmi* bipedales vel ultra. *Spicæ*  $1\frac{1}{2}$ –2 poll. longæ. *Spiculæ* 3 lin. longæ.

8. **A. polytychos**, Steud. l. c. p. 380, cum syn.—C.P. 32.

HAB. Pedrotalagalla, at an elevation of 7000 feet.

Dense *cæspitosus*. *Folia* longiuscula, pedalia vel ultra. *Culmi*  $1\frac{1}{2}$ –2-pedales. *Spicæ* sæpissime 3, valde fragiles, bipollicares. *Spiculæ*  $2\frac{1}{2}$  lin. longæ; *pedicellata flore masculo*, bipaleaceo, *paleis* pilosis.

9. **A. pertusus**, Willd. Sp. Pl. iv. p. 922; Roxb. Fl. Ind. i. p. 258.—*A. punctatus*, Roxb. l. c. p. 264. *A. fascicularis*, Roxb. l. c. p. 265.—C.P. 411, 951.

HAB. Very abundant in the hotter parts of the island.

*Spicæ* plus minus distanter subverticillatæ vel ad apicem racheos subfasciculatæ. *Gluma exterior* spicularum omnium, vel tantum sessilium, vel harum partim foveolata, vel aliquando omnino efoveolata. *Spicula pedicel.* mascula unipaleacea, vel neutra epaleacea.

10. **A. scandens**, Roxb. l. c. p. 258.—c.p. 3258.

HAB. Hot, drier parts of the island.

Closely allied to the preceding, and differing from it principally in the prominently nerved, hairy, and more acute inferior glume of the spikelets, and which I have in no instance observed foveolate.

11. **A. lancifolius**, Trin. (Steud. l. c. p. 382, cum syn.)—*Batratherum molle*, N. ab Es. MSS.—c.p. 3136.

HAB. Badulla District.

Basin versus spicarum spiculæ pedicellatæ sæpissime desunt.

12. **A. rudis**, Steud. l. c. p. 383.—*Batratherum rude*, N. ab Es. MS.—c.p. 866.

HAB. Not uncommon in the Central Province.

13. **A. muricatus**, Retz (Kth. Enum. i. p. 505, cum syn.)—*Anatherum muricatum*, Beauv.—c.p. 871.

HAB. Hotter parts of the island.

14. **A. Zeylanicus**, Arn. (haud N. ab Es.); Benth. Fl. Hongk. p. 423.—*A. pseudograya*, Steud. l. c. p. 395. *A. semiberbis*, Steud. l. c. *Schizachyrium semiberbe*, N. ab Es.—c.p. 3079.

HAB. Central Province, at an elevation of 3000 to 4000 feet.

### 32. HETEROPOGON, Pers.

1. **H. hirtus**, Pers. Syn. ii. p. 533; Benth. Fl. Hongk. p. 424.—*Andropogon contortus*, Linn.; Kth. Enum. i. p. 486.—c.p. 69.

HAB. Common throughout the island.

2. **H. concinnus**, Thw.; *culmis* tenuibus, ramosis; *foliis* linearibus, superne glanduloso-pilosis; *spicul. sessil. gluma inferiore* obovata, obtusa, ciliata, 9-nervia, vix convoluta, *superiore* lanceolata, 3-nervia, quam inferior parum longiore; *spicul. pedicell. gluma inferiore* obovata, villosula, plurinervia, *superiore* lanceolata, 5-nervia, *flore* masculo, uni-paleaceo, 3-andro.—c.p. 3556.

HAB. Bibili, in the Badulla District, at no great elevation.

*Spica* 1-1½-pollicaris, pallide viridis, rachi pedicellisque longe albo-pilosis. *Spiculæ* 1½ lin. longæ. *Arista* flor. hemaphr. 7 lin. longa, pilosula.

Without a careful examination, this might easily be taken for a species of *Apopis*.

### 33. POLLINIA, Trin. Benth. Fl. Hongk.

*Eulalia*, Kunth.

1. **P. tristachya**.—*Andropogon tristachyos*, Roxb. Fl. Ind. i. p. 256. *Saccarum tristachyum*, Steud. Synops. Gram. p. 408.—c.p. 995.

Forma spiculis majoribus.—c.p. 949.

HAB. Central Province, up to an elevation of 7000 feet.

Valde variabilis. *Spicæ* 2-10, 1½-6 poll. longæ. *Spiculæ* 1-2½ lin. longæ.

The form with larger spikelets occurs at the greatest elevations, and in wet peaty soil, and has most commonly two spikes to the panicle.

2. **P. Cumingii**, N. ab Es. MS.—*Andropogon aureo-fulvus*, Steud. l. c. p. 373, cum syn. An *A. aureo-villosus*, Steud. l. c. p. 380?—c.p. 959.

HAB. More elevated parts of the Central Province.

*Culmi* ad basin dense brunneo-tomentosi.

3. **P. Wallichiana**, N. ab Es. (Steud. l. c. p. 410.)—C.P. 950.  
HAB. Not uncommon in the Central Province.

### 34. DIMERIA, R. Br.

1. **D. ornithopoda**, Trin. (Kth. Enum. i. p. 472.)—*D. gracilis*, N. ab Es.; Steud. Synops. Gram. p. 413.—C.P. 952, 3321.

Var.  $\beta$ ; *spiculis* paucioribus, majoribus.—C.P. 3256.

Var.  $\gamma$ ; *spiculis* pubescentibus.—*D. fuscescens*, Trin.; Steud. l. c.; Benth. Fl. Hongk. p. 419.—C.P. 956.

Var.  $\delta$ ; *spiculis* muticis.—C.P. 955.

HAB. Not uncommon throughout the island. Var.  $\beta$ , at Newera Ellia, and Horton Plains, at an elevation of 6000 to 7000 feet.

*Folia* pilosa vel glabra.

An extremely variable plant, but all the forms enumerated above appear to me certainly to belong to one species.

2. **D. pilosissima**, Trin. (Steud. l. c.)—*Haplachne pilosissima*, Pr.; Kth. l. c. p. 473.—C.P. 24, 3261.

HAB. Southern and Central Provinces.

Differs, like the preceding, in the amount of pubescence, and in the size of the spikelets.

3. **D. pusilla**, Thw.; dense cæspitosa, *culmis* erectis, brevibus; *foliis* parvis, linearibus; *spica* simplici; *rachi* tenui, triangulari, glabra; *spiculis* aristatis, villosulis, basi attenuata dense albo-pilosa; *pedicello* brevi, glabro, uno latere ad apicem acute producto; *glumis* acutis, superiore ad apicem setacea.—C.P. 957.

HAB. Kokotodua, Gardner.

*Culmi* 4–8-pollicares. *Spica* pollicaris. *Spiculæ*  $1\frac{5}{8}$  lin. longæ, arista circiter 5 lin. longa.

### 35. IMPERATA, Cyr.

1. **I. arundinacea**, Cyr. (Kth. Enum. i. p. 477, cum syn.)—C.P. 968.

HAB. Common in the hotter parts of the island. Nom. vulg. "Illook."

The leaves make an excellent thatch.

### 36. SACCHARUM, Linn.

1. **S. spontaneum**, Linn. (Kth. Enum. i. p. 475, cum syn.)—*S. Egyptiacum*, Willd.; Kth. l. c. *S. semidecumbens*, Roxb. Fl. Ind. i. p. 236. *S. caudiculatum*, Roxb. l. c. p. 246; Kth. l. c.—C.P. 3681.

HAB. Hotter parts of the island.

### 37. PEROTIS, Ait.

1. **P. latifolia**, Ait. (Kth. Enum. i. p. 470, cum syn.)—C.P. 3164.

HAB. Abundant in the hotter parts of the island.



## 38. ZOYSIA, Willd.

1. **Z. pungens**, Willd. (Kth. Enum. i. p. 471, cum syn.; Benth. Fl. Hongk. p. 418, cum syn.)—c.p. 948.

HAB. Common in sandy ground near the sea.

## 39. SPOROBOLUS, R. Br.

1. **S. Indicus**, R. Br. (Kth. Enum. i. p. 211; Benth. Fl. Hongk. p. 426.)—c.p. 65.

HAB. Very common.

2. **S. diander**, Beauv. (Kth. l. c. p. 213, cum syn.)—*Agrostis diandra*, Retz; Willd. Sp. Pl. Ind. i. p. 371; Roxb. Fl. Ind. i. p. 317.—c.p. 3771.

HAB. Very abundant.

## 40. AGROSTIS, Linn.

1. **A. Royleana**, Trin. (Steud. Synops. Gram. p. 173, cum syn.)—*Calamagrostis Hookeriana*, and *C. Roylei*, Steud. l. c. pp. 192, 193.—c.p. 2394.

HAB. Central Province, at an elevation of 6000 to 8000 feet.

## 41. POLYPOGON, Desf.

1. **P. Monspeliensis**, Desf. (Kth. Enum. i. p. 232, cum syn.)—c.p. 144.

HAB. Oova District, rare, and possibly introduced.

## 42. ARISTIDA, Linn.

1. **A. cœrulescens**, Desf. (Kth. Enum. i. p. 190, cum syn.)—c.p. 915.

HAB. Common in the hot drier parts of the island.

2. **A. depressa**, Retz. (Kth. l. c.)—*A. vulgaris*, Trin. var.  $\delta$ ; Steud. Synops. Gram. p. 138.—c.p. 3684.

HAB. Trincomalee, *Rev. S. O. Glenie*.

## 43. PHRAGMITES, Trin.

1. **P. Roxburghii**, Kth. (Steud. Synops. Gram. p. 196, cum syn.)—*P. Nepalensis*, N. ab Es.; Steud. l. c. *Arundo Karka*, Roxb. Fl. Ind. i. p. 347.—c.p. 176.

HAB. Not uncommon.

## 44. AMPHIDONAX, N. ab Es.

1. **A. Heynei**, N. ab Es. (Steud. Synops. Gram. p. 197.)—c.p. 645.

HAB. Central and Southern Provinces, up to an elevation of 4000 feet.

2. **A. obtusiflora**, Thw.; *culmis* elatis, glabris; *foliis* ensiformibus, acuminatis, glabris, ligula brevissima dense pilosa; *panicula* aperta, ramis capillaribus; *pedicellis* basi glanduloso-callosis; *spiculis* bifloris;

*glumis* ovatis, glabris, apiculatis, quam flores fere dimidio brevioribus; *floribus* sessilibus, æquilongis, hermaphroditis, obtusiusculis, basi pilis subæquilongis cinctis.—C.P. 3470.

HAB. Ratnapoora, at no great elevation.

*Culmi* 3-4-pedales vel ultra. *Folia* bipedalia,  $\frac{3}{4}$ -1 poll. lata, firma. *Panicula* 6-8 poll. longa. *Spiculæ* 1 lin. longæ.

#### 45. CYNODON, Pers.

1. **C. Dactylon**, Pers. (Kth. Enum. i. p. 259, cum syn.)—C.P. 312.

HAB. Common in the warmer parts of the island.

2. **C. Neesii**, Thw.—*C. virgatus*, N. ab Es.; Steud. Synops. Gram. p. 213 (haud Willd. Enum.)—C.P. 3749.

HAB. Trincomalee, *Rev. S. O. Glenie*.

3. **C. gracilis**, N. ab Es. (Steud. l. c.)—*Leptochloa uniflora*, Hochst. Steud. l. c. p. 209.—C.P. 2948.

HAB. Common in the hot drier parts of the island.

#### 46. LEPTOCHLOA, Beauv.

1. **L. Chinensis**, N. ab Es. (Steud. Synops. Gram. p. 209; Benth. Fl. Hongk. p. 430, cum syn.)—*L. tenerrima*, R. and S.; Kth. Enum. i. p. 270.—C.P. 3748 (933 partim).

HAB. Atakalam Corle.

2. **L. filiformis**, R. and S. (Kth. l. c. p. 270, cum syn.)—C.P. 3245.

HAB. Colombo, *Mr. W. Ferguson*.

#### 47. ELEUSINE, Gærtn.

1. **E. Indica**, Gærtn. (Kth. Enum. i. p. 272, cum syn.)—C.P. 937.

HAB. Very abundant.

*E. Coracana*, Gærtn. (nom. vulg. "Koorakkan"), is extensively cultivated by the Cinghalese as a food-grain.

#### 48. DACTYLOCTENIUM, Willd.

1. **D. Ægyptiacum**, Willd. (Kth. Enum. i. p. 261, cum syn.)—C.P. 941.

HAB. Common in the warmer parts of the island.

#### 49. CHLORIS, Linn.

1. **C. barbata**, Sw. (Kth. Enum. i. p. 264, cum syn.)—C.P. 954.

HAB. Common in the hotter parts of the island, especially near the sea.

2. **C. decora**, N. ab Es. (Steud. Synops. Gram. p. 205.)—*C. Meccana*, Hochst.; Steud. l. c.—C.P. 958.

HAB. Hot drier parts of the island.

3. **C. digitata**, Steud. l. c. p. 207.—*Melica digitata*, Roxb. Fl. Ind. i. p. 326. *Gymnopogon digitatus*, N. ab Es. mss.—C.P. 939.

HAB. Hot drier parts of the island.

## 50. GYMNOPOGON, Beauv.

1. **G. rigidus**, Thw.; *culmis* erectis, glabris; *foliis* linearibus, acuminatis, glabris, lateribus inæquilatis, ligula brevi ciliata; *panicula* elongata, scabra; *spiculis* majusculis, brevi-pedicellatis, parum distantibus; *glumis* acutis, trinerviis; *inferiore* parum brevior, *superiore* aliquando aristata; *floribus* longe aristatis, *hermaphrodito* cylindrico *glumis* duplo longiore, *sterili* minutissimo—C.P. 914.

HAB. Hot drier parts of the island.

*Culmi* 2-3-pedales. *Folia* pedalia, 2-3 lin. lata. *Flos hermaphroditus* semipollicaris, *arista* æquilonga vel ultra.

## 51. AVENA, Linn.

1. **A. aspera**, Munro, mss.—C.P. 916.

HAB. Newera Ellia and other of the more elevated parts of the island.

*Culmus* erectus, 2-3-pedalis, nodis pilosulis. *Folia* longiuscule pilosula vel glabra, ligula truncata lacerato-ciliata. *Panicula* parce ramosa, aperta. *Spiculæ* 4-floræ, 6 lin. longæ, floribus parum distantibus. *Gluma inferior* 3-nervia, 2 lin. longa, apice brevi-setacea, *superior* 3½ lin. longa, acuta, 5-nervia. *Palea inferior* scabriuscula, 4 lin. longa, dimidio inferiore 9-nervia, a medio longe aristata fissaque, laciniis 3-nerviis acutis. *Palea superior* brevior, dorso ad carinas ciliata.

## 52. MEGALACHNE, Steud.

1. **M. Zeylanica**, Thw.; *culmis* gracilibus, erectis, glabris, rigidis, basi ramosis; *foliis* glabris, angustis, rigidis, superne prominentim trinerviis, marginum involutione cylindricis, basi tomentellis, ligula brevissima ciliata; *panicula* pauciflora; *spiculis* bifloris, rudimento nullo, flore superiore subsessili; *glumis* glabris, lanceolatis, apice brevi-aristatis; *palea inferiore* pilosa, longe aristata, *superiore* laciniis æqualiter longe aristatis.—C.P. 3247.

HAB. Cinnamon Gardens, Colombo.

*Culmi* 1½-2-pedales. *Panicula* 2-4 poll. longa, purpurascens. *Glumæ* 4 lin. longæ. *Paleæ* 3 lin. longæ, aristis 4-5 lin. longis. *Squamulæ* oblongæ, truncatæ, emarginatæ, ovario brevistipitato æquilongæ. *Stamina* 2. *Stylus* fere ad basin bifidus, stigmatibus plumosis.

## 53. URALEPIS, Nutt.

1. **U. fusca**, Steud. Synops. Gram. p. 247.—*Tridens Indicus*, N. ab Es. mss.—C.P. 934.

HAB. Elephant Pass, Gardner.

*Folia*, cum rachi, scabriuscula.

## 54. POA, Linn.

1. **P. annua**, Linn. (Kth. Enum. i. p. 349, cum syn.)—C.P. 2393.

HAB. Common on roadsides in the neighbourhood of Newera Ellia, possibly introduced.

## 55. ERAGROSTIS, Beauv.

1. **E. bifaria**, W. and A. (Steud. Synops. Gram. p. 264.)—*E. secunda*, N. ab Es.; Steud. l. c. *Poa bifaria*, Vahl; Kth. Enum. i. p. 327.—C.P. 60.  
HAB. Not uncommon up to an elevation of 5000 feet.
2. **E. unioloides**, R. and S. (Steud. l. c.; Benth. Flor. Hongk. p. 432.)—*E. amabilis*, W. and A. *Poa unioloides*, Retz; Kth. l. c. p. 335, cum syn.—C.P. 930.  
HAB. Common in the warmer parts of the island.
3. **E. Brownei**, N. ab Es. (Steud. l. c. p. 279; Benth. Flor. Hongk. p. 432, cum syn.)—*Poa polymorpha*, R. Br. *P. Brownei*, Kth. l. c. p. 333.—C.P. 931.  
HAB. Very abundant throughout the island.
4. **E. Zeylanica**, N. ab Es. (Steud. l. c. p. 265; Benth. l. c. p. 433.)—C.P. 3251. An præcedentis varietas?  
HAB. Hotter parts of the island.
5. **E. orientalis**, Trin. (Steud. l. c. p. 267; Benth. l. c.)—C.P. 925. An *E. Brownei*, N. ab Es., varietas?  
HAB. Central Province, not uncommon.
6. **E. pilosa**, Beauv. (Steud. l. c. p. 263; Benth. l. c. cum syn.)—C.P. *P. pilosa*, Linn., et *P. verticillata*, Cav.; Kth. l. c. p. 329. *P. Ægyptiaca*, Willd.; Kth. l. c. p. 336. *P. elegantula*, Kth. l. c. p. 346.—C.P. 929.  
HAB. Very abundant.
7. **E. megastachya**, Link.—*Poa megastachya*, Kœl.; Kth. l. c. p. 333, cum syn.—C.P. 3163.  
HAB. Hot drier parts of the island.
8. **E. paniculata**, Steud. l. c. p. 266.—*E. nigra*, N. ab Es.; Steud. l. c. p. 267. *Poa paniculata*, Roxb. Fl. Ind. i. p. 340.—C.P. 2626.  
HAB. Doombeera and Maturatte Districts.  
*Flosculi scabriusculi.*
9. **E. nutans**, Steud. l. c. p. 264, cum syn.—*Poa nutans*, Retz; Roxb. l. c. p. 335. *P. Kœnigii*, Kth. l. c. p. 346. *P. interrupta*, Kœn.; Roxb. l. c.—C.P. 3259 (933 partim).  
HAB. Hot, drier parts of the island.
10. **E. plumosa**, Link; Benth. l. c. p. 431.—*Poa plumosa*, Retz; Kth. l. c. p. 328, cum syn. *P. viscosa*, Kth. l. c. p. 336. An *P. ciliaris*, Linn. ?—C.P. 70, 926, 927 (928).  
HAB. Very common in the warmer parts of the island, and an extremely variable plant.

## 56. COELACHNE, R. Br.

1. **C. pulchella**, R. Br. (Steud. Synops. Gram. p. 284; Benth. Fl. Hongk. p. 430.)—*Isachne simpliciuscula*, W. A.; Steud. l. c. p. 96.—C.P. 884.  
HAB. Common in marshy places in the Central Province.
2. **C. perpusilla**, Thw.—*Isachne perpusilla*, Arn.; Steud. l. c.—C.P. 130 (450).

HAB. Central Province, in bogs, at an elevation of 5000 to 7000 feet.

Dense et late cæspitosa, culmis ramosis, 2-18 poll. longis. Folia  $\frac{1}{2}$ -1 poll. longa,  $\frac{1}{2}$ -1 lin. lata, vagina subæquilonga. Panicula sesquipollicaris, aperta, ramis alternis. Spiculæ 1 lin. longæ, flore superiore pedicellato.

### 57. ÆLUROPUS, Trin.

1. **Æ. lagopodioides**, Trin.—*Dactylis lagopodioides*, Linn.; Willd. Sp. Pl. i. p. 410; Burm. Ind. t. 12, fig. 2.—C.P. 876.

HAB. Not uncommon on sandy ground near the sea.

This must surely, I think, be the species figured by Burmann, as quoted above.

*Dactylis glomerata*, Linn., occurs at Newera Ellia, but has no doubt been introduced.

### 58. LOPHATHERUM, Brongn.

1. **L. gracile**, Brongn. (Kth. Enum. i. p. 391; Benth. Fl. Hongk. p. 433, cum syn.)—*L. Lehmanni*, N. ab Es.; Steud. Synops. Gram. p. 300.—C.P. 920, 921.

HAB. Not very uncommon up to an elevation of 4000 feet

### 59. ELYTROPHORUS, Beauv.

1. **E. articulatus**, Beauv. (Kth. Enum. i. p. 391, cum syn.)—C.P. 935.

HAB. Warmer parts of the island.

### 60. TRIPOGON, Roth.

1. **T. Zeylanicus**, N. ab Es.; Steud. Synops. Gram. p. 301.—C.P. 281.

HAB. More elevated parts of the Central Province.

Cæspitosus, glaucus, culmis gracilibus, 1-1 $\frac{1}{2}$  ped. longis, nodis glabris. Folia 4-8 poll. longa, rigidiuscula, pilosa, marginibus involutis. Spica 2-4 poll. longa. Gluma superior 2 lin. longa. Flosculi (setis except.) singuli 1 $\frac{1}{2}$  lin. longi.

### 61. CENTOTHECA, Desv.

1. **C. lappacea**, Desv.; Kth. Enum. i. p. 366, cum syn.—C.P. 923.

HAB. Common at the margins of woods in the warmer parts of the island.

### 62. BRACHYPODIUM, Beauv.

1. **B. sylvaticum**, R. and S. Syst. 2, p. 741.—*B. Nepalense*, N. ab Es.; Steud. Synops. Gram. p. 316 (sub *Festuca*).—C.P. 3253.

HAB. More elevated parts of the island, common.

### 63. STREPTOGYNE, Beauv.

1. **S. crinita**, Beauv.; Kth. Enum. i. p. 427.—C.P. 922.

HAB. Matele district, Gardner.

64. **ARUNDINARIA**, Rich.

1. **A. debilis**, Thw.; *culmis* valde elongatis, glabris vel cum vaginis, plus minus hispidis; *foliis* parvis, glaucis, lanceolatis, apice acutissimis, glabris vel superne pilosis, basi acutatis vel subtruncatis; *paniculis* simplicibus, ramos paucifloros breviusculos semiverticellatis terminantibus; *spiculis* 3-floris, floribus 2, inferioribus hermaphroditis; *glumis* subacutis, mucronatis, 6-7-nerviis, pilosulis, *inferiore* parum brevior; *palea inferiore* quam *superior* fere duplo longiore, gluma superiore subæquilonga.—C.P. 1.

HAB. More elevated parts of the Central Province.

*Culmi* 2-5 lin. in diam. *Stamina* 3.

The foliage of this plant is used at Newera Ellia as a fodder for horses.

2. **A. floribunda**, Thw.; *foliis* lanceolatis, acuminatissimis, nervosis, transversim venulosis, basi plus minus acutis, vagina ad apicem longe ciliatis; *paniculis* majusculis; *spiculis* laxis, 6-8-floris; *glumis* parvis, acutis, glabris, *inferiore* minore: *palea inferiore* pilosula, subulato-acuminata, *superiore* subæquilonga, quam gluma superior duplo longiore.—C.P. 2624.

HAB. Maturatte District, at an elevation of 5000 feet.

*Culmi* erecti, 2-5-pedales. *Spiculæ* pollicares vel ultra. *Flores* 3½ lin. longi. *Stamina* 3.

65. **BAMBUSA**, Schreb.

1. **B. Thouarsii**, Kth. Enum. i. p. 431; Steud. Synops. Gram. p. 329.—An *B. vulgaris*, Schrad.; Steud. l. c. ? *B. arundinacea*, Moon's Cat. p. 26. Et var.  $\beta$ . *picta*; *culmis* majoribus elatioribusque, stramineo et viridi variegatis.—C.P. 3252.

HAB. Southern and central parts of the island, up to an elevation of 2000 feet. Nom. vulg. "Oona-gass."

Much used by the Cinghalese for temporary buildings and other purposes. The flowers, which are very rarely produced, very much resemble those of the next species; but their outer paleæ are somewhat longer, and terminate in subulate points.

2. **B. arundinacea**, Willd. (Kth. l. c. cum syn.; Roxb. Pl. Cor. i. t. 79.)—*B. orientalis*, N. ab Es.; Steud. l. c. *B. spinosa*, Roxb.; Kth. l. c.—C.P. 3320.

HAB. Common on the margins of rivers and streams in the warmer parts of the island. Nom. vulg. "Kattoo-oona-gass."

The seeds are eaten by the Cinghalese.

3. **B. attenuata**, Thw.; *culmis* mediocribus, inermibus; *vaginis* sparsim pallide adpresso-pilosis; *foliis* glabris, lineari-lanceolatis, subabrupte acuminatis, basi rotundatis, petiolatis; *ramis florigeris* fasciculatis, breviusculis, foliatis; *panicula* simplici, spiciformi; *spiculis* glabris, subsessilibus, in fasciculos parvos congestis, angustis, attenuato-cylindricis, 5-floris, floribus 2 apicalibus plerumque sterilibus; *pedicellis* inclusis, quam flores dimidio brevioribus; *paleis inferioribus* 12-nerviis, acutis, mucronatis.—C.P. 3255.

HAB. Central Province, at an elevation of 4000 to 6000 feet.

*Culmi* 12-25-pedales, ½-1 poll. in diam. *Folia* 3-6 poll. longa, 4-9 lin. lata, arcte ramoso-striata. *Spiculæ* pollicares. *Paleæ inferiores* superioribus parum

longiores. *Stylus* elongatulus, infra medium incrassatus. *Stigmata* 3, plumosa, stylo multo breviora.

## 66. DENDROCALAMUS, N. ab Es.

1. **D. monadelphus**, Thw.; *foliis* lanceolatis, acuminatis, parum obliquis; *palea inferiore* apice subulato-acuta, quam superior parum longiore; *staminibus* 6, filamentis in tubum coalitis; *antheris* linearibus, angustis, subulato-apiculatis, apiculo piloso.—C.P. 3359.

HAB. Common in the Central Province, at an elevation of 4000 to 6000 feet.

*Culmi* inermes, teretes, fistulosi, 10–12-pedales, circiter 1 poll. in diam., ramosi, *vaginis* fugaci-albido-pilosis apice longe caduco-ciliatis. *Folia* 2–12 poll. longa,  $\frac{1}{2}$ – $1\frac{1}{2}$  poll. lata, plurinervia, interstitiis arctissime longitudinaliter venulosis. *Spiculæ* acutæ, 4–6 lin. longæ, sæpius 3-floræ, densissime glomeratæ. *Palea superior*, ciliata.

## Gen. nov. 67. OCHLANDRA, Thw.

*Flores* bipaleacei. *Squamulæ* 6–9 (vel forsan plura) lanceolata, 1–3-nervia. *Stamina* numerosa (7–30). *Filamenta* libera, inter se æqualia. *Antheræ* lineares, utrinque brevissime bifidæ, apice mucronulatæ. *Stylus* simplex, longiusculus, robustus, subprismaticus. *Stigmata* 4–5, brevissima, plumosa. *Caryopsis* majuscula, ovoidea, stylo persistente indurato acuminata, paleis squamisque spiculæ tecta.—Frutex, culmis suberectis, confertis; foliis majusculis; inflorescentia terminali, laxe spicata; spiculis in glomerulis parvis aggregatis, 1-floris; flore terminali.

1. **O. stridula**, Thw.—*Bambusa stridula*, Moon's Cat. p. 26.—C.P. 241 (429?).

HAB. Common in the warm, moister parts of the island, and very abundant in the Saffragam District. Nom. vulg. "Batta-gass."

The leaves make an excellent thatch.

## 68. TRITICUM, Linn.

1. **T. repens**, Linn.; Kth. Enum. i. p. 440, cum syn.—C.P. 924.

HAB. North of the island, Gardner.

## CLXVIII. LYCOPODIACEÆ.

## 1. LYCOPODIUM, Linn.

1. **L. Phlegmaria**, Linn. (Willd. Sp. Pl. v. p. 10.)—*L. mirabile*, Willd. l. c.—c.p. 987 (1195, 1277).

HAB. Common in the Central Province.

2. **L. Hookeri**, Wall. Cat. n. 146; Hook. et Grev. Ic. Fil. t. 185.—c.p. 986 (1193, 1194).—Forma *foliis* latioribus.—c.p. 3281.

HAB. Central Province, not uncommon.

Very variable in the form of its leaves, but the size and shape of the sporangia very constant. Spring refers this to *L. ulicifolium*, Vent.

3. **L. serratum**, Thunb. Fl. Jap. p. 341, t. 38; Willd. l. c. p. 21; Hook. et Grev. Ic. Fil. t. 37.—c.p. 1419 (1189).

Forma *foliis* minoribus, magis confertis, firmioribus, inconspicue serratis.—c.p. 1115 (1488, 1273).

HAB. More elevated parts of the Central Province.

Equally variable with the preceding.

4. **L. clavatum**, Linn. (Willd. l. c. p. 13.)—c.p. 3283 (1192, 1279).

HAB. Horton Plains, at an elevation of 7000 feet.

5. **L. complanatum**, Linn. (Willd. l. c. p. 14.)—c.p. 3489.

HAB. Horton Plains, with the preceding species.

6. **L. cernuum**, Linn. (Willd. l. c. p. 15.)—c.p. 984 (1187).

HAB. Very common in the Central Province.

7. **L. Carolinianum**, Linn. (Willd. l. c. p. 13.)—c.p. 1416 (1191, 1278).

HAB. Pedro talagalla, at an elevation of 7000 feet.

## 2. SELAGINELLA, Spring.

1. **S. rupestris**, Spring, Mon. Lycop. part ii. p. 55.—c.p. 1414.

HAB. Damboul, *Gardner*.

2. **S. stolonifera**, Spring, l. c. p. 209.—*Lycop. stoloniferum*, Sw.—c.p. 1412 (1185).

HAB. Common in the Central Province.

3. **S. caudata**, Spring, l. c. p. 139.—*Lycop. caudatum*, Desv.; Benth. Fl. Hongk. p. 437.—c.p. 3284 (1276).

HAB. Forests of the Central and Southern Provinces.

4. **S. caulescens**, Spring, l. c. p. 158.—c.p. 985 (1186).

HAB. Central Province, *Gardner*.

5. **S. atroviridis**, Spring, l. c. p. 124.—*S. monospora*, Spring, l. c. p. 135. *Lycop. atroviride*, Wall.; Hook. Gen. Fil. t. 117, B; Benth. l. c.—c.p. 1417, 1418 (1184, 1274).

HAB. Very abundant.

6. **S. concinna**, Spring, l. c. p. 199.—*Lycop. concinnum*, Sw.—c.p. 3280.

HAB. Raxawa, in the Central Province; and near Galle.

In my specimens the inner leaves scarcely differ in shape from the outer entire distichous ones.



3. **PSILOTUM**, Sw.

1. **P. triquetrum**, Sw. (Hook. Gen. Fil. t. 87.)—c.p. 1420 (1272).  
 HAB. Not uncommon in the Central Province.

## CLXIX. MARSILEACEÆ.

1. **MARSILEA**, Linn.

1. **M. quadrifolia**, Linn. (Willd. Sp. Pl. v. p. 538.)—c.p. 3051.  
 HAB. South of the island.

2. **M. erosa**, Willd. l. c. p. 540.—c.p. 1422.  
 HAB. Hotter parts of the island, north of Kandy.

2. **AZOLLA**, Lam.

1. **A. pinnata**, R. Br. Prodr. p. 167; Moore, Ind. Fil. p. 190, cum syn.—c.p. 1421.  
 HAB. Common in the warmer parts of the island.

## CLXX. FILICES.

1. **BOTRYCHIUM**, Sw.

1. **B. daucifolium**, Wall. (Hook. Bot. Mag. 3rd ser. xviii. t. 5340.)  
 —c.p. 1410 (1271).

HAB. Forests of the Central Province, up to an elevation of 4000 feet.

2. **B. Virginicum**, Willd.—Var.  $\beta$ . *Indicum*, Hook. Gard. Ferns, t. 29. Var.  $\beta$ . *lanuginosum*, Moore, Ind. Fil. p. 313, cum syn. Var.  $\beta$ . Wall. Cat. n. 48.—c.p. 3266 (1181).

HAB. Newera Ellia, on the turf under the shade of trees.

In general appearance very like the preceding species, but at once distinguished from it by the fertile branch of the rachis arising from the middle of the leafy portion of the frond, whilst in the preceding it originates below it.

2. **HELMINTHOSTACHYS**, Kaulf.

1. **H. Zeylanica**, Hook. Gen. Fil. t. 47, B, cum syn.; Gard. Ferns, t. 28.—*Botrychium Zeylanicum*, Sw. Syn. Fil. p. 172.—c.p. 1411.

HAB. South of the island, on the borders of paddy fields, etc., not uncommon.

3. **OPHIOGLOSSUM**, Linn.

1. **O. pendulum**, Linn. (Willd. Sp. Pl. v. p. 60; Sw. Syn. Fil. p. 170; Hook. Gard. Ferns, t. 53.)—c.p. 1409 (1180).

HAB. Common upon the trunks of trees in the Central Province.

2. **O. vulgatum**, Linn. (Willd. l. c. p. 58.)—*O. ovatum*, Willd. l. c.—c.p. 1408.

HAB. More elevated parts of the Central Province.

4. **ANGIOPTERIS**, Hoffm.

1. **A. erecta**, Hoffm. (Hook. Fil. Exot. t. 75, cum syn.; Benth. Fl. Hongk. p. 440.)—c.p. 1407 (967, 1177).

HAB. Common up to an elevation of 5000 feet.

5. **MARATTIA**, Sw.

1. **M. sorbifolia**, Sw. Syn. Fil. p. 168; Willd. Sp. Pl. v. p. 67.—c.p. 3147.

HAB. More elevated parts of the Central Province.

6. **OSMUNDA**, Linn.

1. **O. Javanica**, Blume, En. Fil. Jav. p. 252; Hook. Kew Journ. of Bot. ix. p. 360; Kze. in Schk. Fil. Suppl. t. 111; Benth. Fl. Hongk. p. 441, cum syn.—*O. Zeylanica*, Kze. in Schk. Fil. Suppl. t. 122 (not Linn.).—c.p. 2988 (961, 1178).

HAB. Common in the higher parts of the Central Province.

7. **LYGODIUM**, Sw.

1. **L. circinatum**, Sw. Syn. Fil. p. 153; Benth. Fl. Hongk. p. 441, cum syn.—*L. dichotomum*, Sw. l. c. p. 154. *Hydroglossum*, Willd.—c.p. 1406 (1269).

HAB. Not uncommon in the Central Province.

2. **L. scandens**, Sw. (Presl; Benth. l. c. cum syn.)—c.p. 1404 (1183, 1270).

HAB. Very common in the warmer parts of the island.

3. **L. Japonicum**, Sw. (Presl; Benth. l. c. p. 442.)—c.p. 1405 (1268).

HAB. Common in the Central Province.

8. **SCHIZÆA**, Sm.

1. **S. digitata**, Sw. (Willd. Sp. Pl. v. p. 86, cum syn.; Hook. Gard. Ferns, t. 49.)—*Actinostachys*, Wall.—c.p. 3105 (969, 1182).

HAB. Kittool Galle, and near Peradenia.

9. **GLEICHENIA**, Sw.

1. **G. dichotoma**, Willd. (Hook. Sp. Fil. i. p. 12.)—c.p. 3106 (970, 1053).

HAB. Extremely common throughout the island.

10. **CERATOPTERIS**, Brongn.

1. **C. thalictroides**, Brongn. (Hook. Sp. Fil. ii. p. 235; Gen. Fil. t. 12.)—c.p. 980 (1171).

HAB. Not uncommon on the borders of streams and of paddy fields.

## 11. ACROSTICHUM, Linn.

a. *Euacrostichum.*

1. **A. aureum**, Linn. (Moore, Ind. Fil. p. 5, cum syn.; Hook. Gen. Fil. t. 81; Sp. Fil. v. ined.—c.p. 1315 (1152)).

HAB. Common in wet places in the hotter parts of the island.

b. *Elaphoglossum.*

2. **A. laurifolium**, Thouars.—*Elaphoglossum laurifolium*, Moore, l. c. p. 359, cum syn.—c.p. 1310 (1165).

HAB. Common in forests of the Central Province.

[Probably a variety of *A. conforme*, Sw.—Hook.]

3. **A. marginatum**, Wall. Cat. n. 17.—*A. conforme*, Bl. Fil. Jav. t. 5, and Hook. Sp. Fil. v. ined. *Elaphog. marginatum*, Moore, l. c. p. 361, cum syn.—c.p. 1311.

HAB. Central Province; particular locality not recorded.

Distinguished from the preceding by the much broader paleæ of the rhizome.

4. **A. squamosum**, Sw. (Hook. l. c. v. ined.)—*Elaphog. squamosum*, J. Sm.; Moore, l. c. p. 367, cum syn. *A. paleaceum*, Hook. et Grev. Ic. Fil. t. 235.—c.p. 3292 (1164 partim).

HAB. Central Province, not very common.

5. **A. spathulatum**, Sw. (Bory, Itin. i. p. 363, t. 20, f. 1; Hook. l. c. v. ined.)—*Elaphog. piloselloides*,  $\delta$ , *spathulatum*, Moore, l. c. p. 363, cum syn.—c.p. 989 (1164, partim).

HAB. Newera Ellia, Gardner.

Mem.—I know nothing of *A. Lindenii*, Bory, and *A. Neitneri*, Kl. (*Elaphoglossum*, Moore, l. c. pp. 360, 362), said to occur in the island.

c. *Stenochlæna.*

6. **A. scandens**, Willd., not. ad *Lomaria scandens*, Sp. Pl. v. p. 294; Hook. l. c. v. ined.—*Stenochlæna scandens*, J. Sm. in Hook. Journ. of Bot. iii. (1841) p. 401.—c.p. 1312.

HAB. Common in the warmer parts of the island.

d. *Pæcilopteris.*

7. **A. repandum**, Blume, En. Pl. Jav. p. 104; Benth. Fl. Hongk. p. 444, cum syn.; Hook. l. c. v. ined.—*A. contaminans*, Wall. Cat.—c.p. 3075 (1161, 1232).

HAB. Forests of the Central Province, common.

8. **A. Hookerianum**.—*A. (Campium) proliferum*, Hook. Ic. Pl. vii. t. 681-2. *A. virens*, Wall. Cat. n. 1033; Hook. Ic. Pl. t. 681 and 682. Sp. Fil. v. ined. *Pæcilopteris Hookeriana*, Moore, Genera of Ferns, p. 20.—c.p. 1313.

HAB. Not uncommon in the forests of the Central Province.

e. *Polybotrya.*

9. **A. appendiculatum**, Willd. l. c. p. 114; Hook. l. c. v. ined.—*A. setosum*, Wall. Cat. n. 30. *Polybotrya appendiculata*, J. Sm.—c.p. 1314 (1160, 1234).

HAB. Very common.

f. *Gymnopteris*.

10. **A. decurrens**, Hook. Fil. Exot. t. 94; Hook. Sp. Fil. l. c. v. ined.; Benth. l. c. p. 443.

Forma *fronde* simplice.—C.P. 1316, 1317 (1157, 1135).

Forma *fronde* pinnatifida.—C.P. 1318.

HAB. Very abundant in damp forests.

11. **A. quercifolium**, Retz. (Willd. l. c. p. 112, cum syn.; Hook. l. c. v. ined.)—*Gymnopteris quercifolia*, Benth.; Hook. Ic. Pl. x. t. 905, cum syn.—C.P. 1319 (1170).

HAB. Shady banks in the warmer parts of the island.

g. *Hymenolepis*.

12. **A. spicatum**, Linn. f. (Hook. l. c. v. ined.)—*Hymenolepis spicata*, Presl. *H. ophioglossoides*, Kaulf. *Lomaria spicata*, Willd. l. c. p. 289.—C.P. 1303 (1135).

HAB. Forests of the Central Province, not uncommon.

## 12. DRYMOGLOSSUM, Presl.

1. **D. piloselloides**, Presl, Tent. 227, t. 10, ff. 5, 6; Hook. l. c. v. ined.; Moore, Ind. Fil. p. 343, cum syn.—C.P. 3076 (1156).

HAB. Common on the trunks of trees.

## 13. VITTARIA, Sm.

1. **V. scolopendrina**, Schk.—*Vittaria Zeylanica*, Fée, and Hook. Sp. Fil. v. ined. *Tæniopsis scolopendrina*, J. Sm. in Hook. Journ. of Bot. iv. (1842) p. 67. *Pteris scolopendrina*, Sw.; Bory; Willd. Sp. Pl. v. p. 356.—C.P. 1304 (1231).

HAB. Forests of the Central Province.

2. **V. lineata**, Sw. (Willd. l. c. p. 404; Hook. l. c. v. ined.)—*Tæniopsis lineata*, J. Sm. l. c.—C.P. 281 (1136).

HAB. Common in the Central Province.

## 14. LOMARIA, Willd.

1. **L. elongata**, Blume (Hook. Sp. Fil. iii. p. 3, cum syn.)—C.P. 1333 (1158).

HAB. More elevated parts of the Central Province.

## 15. BLECHNUM, Linn.

1. **B. orientale**, Linn. (Hook. Sp. Fil. iii. p. 52, cum syn.)—C.P. 977 (1085).

HAB. Very common on moist banks.

## 16. VAGINULARIA, Fée.

1. **V. Junghuhnii**, Mett.—*Monogramme*, Hook. Sp. Fil. v. p. 123, t. 288, B.—C.P. 1281.

HAB. Central Province, at an elevation of 3000 to 5000 feet, not very common.

## 17. DOODIA, R. Br.

1. **D. dives**, Kze.—Var.  $\beta$ . *Zeylanica*, Hook. Sp. Fil. iii. p. 73.—c.p. 976 (1086).

HAB. Woods of the Central Province.

## 18. MENISCIUM, Schreb.

1. **M. triphyllum**, Sw. Syn. Fil. pp. 19 and 206; Willd. Sp. Pl. v. 133; Hook. et Grev. Ic. Fil. t. 120; Hook. Sp. Fil. v. ined.—c.p. 1193.

HAB. Not uncommon in the Central and Southern Provinces.

2. **M. Thwaitesii**, Hook. Fil. Exot., not. ad t. 83.—c.p. 3145.

HAB. Oodoo pussalawa, in the Central Province, at an elevation of 4000 feet. [Traces of an involucre have been seen on the sori of this plant, which induce Sir Wm. Hooker to think it ought to be removed to the *Asplenium*-group. It will be further noticed in the Suppl. to 'Species Filicum.']

## 19. HEMIONITIS, Linn.

1. **H. cordata**, Roxb. (Hook. Sp. Fil. v. ined.; Moore.)—c.p. 1309 (1169).

HAB. Common in the warmer parts of the island.

2. **H. reticulata**, Forst.—*Antrophyum reticulatum*, Kaulf.; Hook. l. c. v. ined.; Spreng. Syst. iv. p. 67.—c.p. 1305 (1229).

HAB. Very common in the upper forests of the Central Province.

3. **H. plantaginea**, Cav.—*Antrophyum plantagineum*, Spreng. l. c.; Hook. l. c. v. ined.—c.p. 3290.

HAB. Central Province.

## 20. GRAMMITIS, Sw.

a. *Loxogramme*.

1. **G. involuta**, Don, Prod. Fl. Nep. p. 14; Hook. et Grev. Ic. Fil. t. 53.—*Gymnogramme*, Hook. Sp. Fil. v. p. 155. *Loxogramme involuta*, Pr.; Moore.—c.p. 1302 (1155).

HAB. Common upon the trunks of trees in the Central Province.

2. **G. lanceolata**, Sw. Syn. Fil. pp. 22 and 212, t. 1, f. 4; Willd. Sp. Pl. v. p. 139.—*Gymnogramme*, Hook. l. c. v. p. 156. *Loxogramme lanceolata*, Pr.; Moore.—c.p. 3146.

HAB. Pedro talagalla, at an elevation of 7000 feet.

b. *Leptogramme*.

3. **G. totta**, Pr.; Moore.—*Polypodium tottum*, Willd. l. c. p. 201. *Gymnogramme totta*, Schlecht.; Hook. l. c. v. p. 138. *G. Lowei*, Hook. et Grev. Ic. Fil. t. 89.—c.p. 1292.

HAB. Common in the more elevated parts of the Central Province.

c. *Gymnogramme*.

4. **G. serrulata**, Blume, En. p. 113; Fl. Jav. t. 42.—*Gymnogramme Javanica*, Blume, En. p. 112; Fl. Jav. t. 41; Hook. l. c. p. 145. *G.*

*falcata*, J. Sm. *Diplazium falcatum* and *D. fraxineum*, Don, l. c. pp. 12, 13. *Grammitis caudata*, Wall. Cat. n. 4.—c.p. 3264 (1225).

HAB. Horton Plains and Maturatte District, at an elevation of 5000 to 4000 feet.

d. *Stegnogramme*.

[5. **G. aspidioides**, Hook. Gen. Fil. t. 120, B, and Ic. Pl. x. t. 950; Hook. Sp. Fil. v. p. 150; Gardner, n. 1292.—I insert this on the authority of specimens in my herbarium from Mr. Gardner, the same number as Mr. Thwaites (*supra*) refers to *G. (Leptogramme) totta*. I possess both species from Ceylon.—Hook.]

21. **ASPLENIUM**, Linn.

a. *Thamnopteris*.

1. **A. Nidus**, Linn. (Hook. Sp. Fil. iii. p. 77, cum syn.)—c.p. 3267.

HAB. Very common.

b. *Euasplenium*.

2. **A. ensiforme**, Wall. (Hook. et Grev. Ic. Fil. t. 71; Hook. Sp. Fil. l. c. p. 89.)—c.p. 1334.

HAB. Rambodde, in the Central Province, Gardner.

3. **A. Wightianum**, Wall. (Hook. l. c. p. 105, t. 167, cum syn.)—*A. Walkeræ*, Hook. l. c. p. 108, t. 163. An *A. longipes*, Fée; Hook. l. c. p. 106<sup>p</sup>—c.p. 1010 (1070).

HAB. Common in the forests of the Central Province.

The two forms of *A. Wightianum* and *A. Walkeræ* merge insensibly into one another.

4. **A. persicifolium**, J. Sm.

Var. *latifolium*, Hook, l. c. p. 109.—c.p. 3461.

HAB. Newera Ellia, and other more elevated parts of the Central Province.

5. **A. elongatum**, Sw. Syn. Fil. p. 79; Hook. l. c. p. 117.—c.p. 1007 (1078).

HAB. Not uncommon in the Central Province.

6. **A. tenerum**, Forst.; Hook. l. c. p.—c.p. 1371.

HAB. Hantani, in the Central Province.

*Caudex brevis. Frondes (cum stipite) 6–15 poll. longæ, subtus paleis parvis adpressis fuscis conspersæ. Pinnæ plus minus acuminatæ. Indusium demum margine arcte revolutum.*—[This Fern, Ceyl. Pl. n. 1371, is identical with my n. 1072 of Gardner, which in my Sp. Fil. iii. p. 158, I have considered a small form of the very variable *A. macrophyllum*, Sw. Mr. Thwaites (see next page, n. 13) refers Gardner's plant to it.—Hook.]

7. **A. erectum**, Bory. (Hook. l. c. p. 126, cum syn.)—c.p. 3141 (1074).

HAB. Central Province, not uncommon.

8. **A. resectum**, Sm. (Hook. l. c. p. 130, cum syn.)—c.p. 427, 1336, 1337, 1338, 3269 (1075, 1077).

HAB. Very abundant in the forests of the Central Province.

An extremely variable species.

9. **A. heterocarpum**, Wall. (Hook. l. c. p. 132, t. 175.)—c.p. 1006 (1076).

HAB. Not uncommon in moist woods.

10. **A. normale**, Don, Prod. Fl. Nep. p. 7; Moore, Ind. Fil. p. 149, cum syn.—*A. multijugum*, Wall.; Hook. l. c. p. 139, t. 188, cum syn.—C.P. 1005 (1073).

HAB. Damp forests, common.

11. **A. formosum**, Willd. (Hook. l. c. p. 143, cum syn.; Moore, l. c. p. 132, cum syn.)—C.P. 3487.

HAB. Forests above Galagama, towards Horton Plains.

12. **A. caudatum**, Forst. (Hook. l. c. p. 152, cum syn.)—C.P. 1339 (1081).

HAB. Newera Ellia, *Gardner*.

13. **A. macrophyllum**, Sw. (Hook. l. c. p. 158, tt. 196, 197, cum syn.)—C.P. 1340 (1072).

HAB. Newera Ellia and other more elevated parts of the Central Province.

Is this really distinct from *A. caudatum*, or merely that species in a less advanced stage of growth?

14. **A. falcatum**, Lam. (Hook. l. c. p. 160, cum syn.)—C.P. 3140 (1080).

HAB. Common.

15. **A. planicaule**, Wall. (Hook. l. c. p. 163, cum syn.; Moore, l. c. p. 154, cum syn.)—C.P. 3289.

HAB. Not uncommon in forests of the Central Province.

16. **A. furcatum**, Thunb. (Hook. l. c. p. 165, cum syn.)—C.P. 3497.

HAB. More elevated parts of the Central Province.

17. **A. spathulinum**, J. Sm. (Hook. l. c. p. 170, cum syn.)—C.P. 2905, 1341 (1082).

HAB. Central Province, in the higher forests.

18. **A. nitidum**, Sw. (Hook. l. c. p. 172.)—C.P. 1008 (1083, 1084).

Forma magis composita. An *A. laserpitiifolium*, Lam.; Hook. Sp. Fil. iii. p. 172, t. 203?—C.P. 3801.

HAB. Common in the forests of the Central Province.

19. **A. tenuifolium**, Don. (Hook. l. c. p. 193, cum syn.)—C.P. 3268 (1079).

HAB. Newera Ellia, by the sides of streams.

20. **A. prolongatum**, Hook. l. c. p. 209; 2nd Century of Ferns, t. 42.—*A. rutafolium*, Pr., var.  $\beta$ . *furcata*, Moore, l. c. p. 163.—C.P. 1348.

HAB. Pas-doon Corle, *Gardner*. Singhe-rajah forest.

21. **A. varians**, Hook. et Grev. (Hook. l. c. p. 192, cum syn.; Moore, l. c. p. 176.)—C.P. 3139.

HAB. Oova District, not common.

#### c. *Athyrium*.

22. **A. nigripes**, Mett. (Hook. l. c. p. 222, cum syn.)—*Athyrium tenuifrons*, var.  $\gamma$ . *straminea*, Moore, l. c. p. 188.—C.P. 3067.

HAB. Forests of the Central Province, at an elevation of 4000 to 6000 feet.

23. **A. macrocarpum**, Blume (haud Fée). (Hook. l. c. p. 222, cum syn.)—C.P. 1372 (1103, 1112).

HAB. More elevated parts of the Central Province.

24. **A. aspidioides**, Schlecht. (Hook. l. c. p. 223, cum syn.)—C.P. 1346 (1065).

HAB. Newera Ellia, Maturatte, etc.

25. **A. gymnogrammoides**, Kl. (Hook. l. c. p. 227, cum syn.)—C.P. 1344, 1345 (1066–9).

HAB. Higher forests of the Central Province.

Some forms of this plant approach very nearly *A. aspidioides*.

26. **A. australe**, Brack. (Hook. l. c. p. 232, cum syn.)—C.P. 1347.

HAB. Damp shady forests in the Central Province.

The more deltoid and flaccid fronds and their smaller ultimate divisions distinguish this species from the preceding one.

#### d. *Diplazium*.

27. **A. lanceum**, Thunb. (Hook. l. c. p. 235, cum syn.)—C.P. 1335.

HAB. Forests of the Central Province.

28. **A. Zeylanicum**, Hook. l. c. p. 237; 2nd Century of Ferns, t. 16.—C.P. 3101 (1249).

HAB. Forests of the Ambagamowa and Kotmalie Districts.

A good species, and by no means the young state of a more compound form.

29. **A. sylvaticum**, Pr. (Hook. l. c. p. 248, cum syn.)—*A. elatum*, Mett.; Hook. l. c. p. 249. *Diplazium sylvaticum*, Sw.; Moore, l. c. p. 339, cum syn.—C.P. 1349.

HAB. Common in the forests of the Central and Southern Provinces.

30. **A. Thwaitesii**, A. Br. (Hook. l. c. p. 250, cum syn.; 2nd Century of Ferns, t. 45.)—C.P. 1343.

HAB. Forests of the Central Province.

31. **A. Schkuhrii**, Mett. (Hook. l. c. p. 251, cum syn.)—*Diplazium Schkuhrii*, L. Sm.; Moore, l. c. p. 337.—C.P. 3100 (1063, 1247).

HAB. Ambagamowa and Saffragam Districts.

In large specimens the lower pinnæ are pinnated with their pinnules pinnatifid.

32. **A. polypodioides**, Mett. (Hook. l. c. 257, cum syn.)—*Diplazium polypodioides*, Blume; Moore, l. c. p. 334, cum syn.—C.P. 1352, 1353, 3098 (1061, 1062).

Var.  $\beta$ ; *pinnulis* magis acuminatis, ramo inferiore venarum primariarum a costa parum distante.—*Diplazium dilatatum*, var.  $\beta$ . *minor*, Moore, l. c. p. 327.—C.P. 3332 (1245). An species distincta?

HAB. Very abundant in damp forests; var.  $\beta$ , Ambagamowa.

33. **A. dilatatum**, Hook. l. c. p. 258, cum syn.—C.P. 1350 (1060).

HAB. Very common in damp forests.

#### e. *Anisogonium*.

34. **A. esculentum**, Pr. (Hook. l. c. p. 268, cum syn.)—*Callipteris ambigua*, Moore, Ind. Fil. p. 216, cum syn.—C.P. 3270 (1058).

HAB. Common on the shady banks of rivers and streams.

## 22. ALLANTODIA, Wall.

1. **A. Brunoniana**, Wall. (Plant. As. Rar. p. 44, t. 52; Hook. Sp. Fil. iii. p. 275, cum syn.)—C.P. 2543 (1057).



HAB. Damp forests of the Central Province, at an elevation of 3000 to 6000 feet.

### 23. ACTINIOPTERIS, Link.

1. **A. radiata**, Link. (Hook. Sp. Fil. iii p. 276, cum syn.; Hook. Ic. Plant. tt. 975, 976.)—C.P. 1342.

HAB. Trincomalie, *Gardner*. Gonagama on the lower Badulla road from Kandy.

### 24. PTERIS, Linn.

#### a. *Pellæa*.

1. **P. geraniifolia**, Raddi. (Hook. Ic. Plant. x. t. 915.)—*Pellæa geraniifolia*, Fée; Hook. Sp. Fil. ii. p. 132, cum syn.—C.P. 3103 (1125).

HAB. Common in the warmer parts of the island.

2. **P. Boivini**, Hook. l. c. p. 147, t. 118, A.—C.P. 3363.

HAB. Hantani, in the Central Province, at an elevation of 4000 feet.

#### b. *Ornithopteris*.

3. **P. aquilina**, Linn.—Var.  $\beta$ . *lanuginosa*, Hook. l. c. p. 196, cum syn.—C.P. 1332 (1134).

HAB. Very common in the Central Province.

#### c. *Eupteris*.

4. **P. longifolia**, Linn. (Hook. Sp. Fil. ii. p. 157, cum syn.)—C.P. 3144.

HAB. Oova and Doombera Districts.

5. **P. Cretica**, Linn. (Hook. l. c. p. 159, cum syn.)—C.P. 3502.

Var.  $\beta$ . *P. Hookeriana*, Ag. (Hook. l. c. p. 165.)—C.P. 1329.

HAB. Oova District; var.  $\beta$  in the Saffragam District.

Var.  $\beta$  is distinguished by the more coriaceous texture of its barren fronds, their wider pinnæ, and the veins of the latter being directed more forward.

6. **P. crenata**, Sw. (Hook. l. c. p. 163, cum syn.)—C.P. 1328.

HAB. A very abundant species in the warmer parts of the island.

7. **P. patens**, Hook. l. c. p. 177, t. 137, cum syn.—An *P. (Campteria) Gardneri*, Fée, Gen. Fil. p. 139; Hook. l. c. p. 207?—C.P. 1047 (1126).

HAB. Forests of the Central Province, not very uncommon.

A most beautiful species, with large feathery fronds four or five feet in length.

8. **P. quadriaurita**, Retz. (Hook. l. c. p. 179, t. 134, B, cum syn.)—C.P. 1330 (1129).—Et forma *ludens*; *pinnulis* plus minus irregularibus, aliis elongatis, aliis abbreviatis vel abortivis.—C.P. 1351, 3060 (1133).

HAB. Very abundant.

9. **P. semipinnata**, Linn. (Hook. l. c. p. 169, cum syn.)

HAB. Ceylon; Walker, in *Herb. Hook.*

I have never seen the typical *P. semipinnata* growing in the island; some of the aberrant forms of *P. quadriaurita* would seem to make an approach to it.

d. *Campteria*.

10. *P. biaurita*, Linn. (Hook. l. c. p. 203, cum syn.)—*P. nemoralis*, Willd.; Hook. l. c. p. 202, cum syn.—c.p. 1048 (1128, 1130, 1240, 1241).

HAB. Very common in the Central Province.

Although specimens occur which appear intermediate between this plant and *P. quadriaurita*, I believe the two are specifically distinct. Very young plants of each, growing side by side under exactly similar conditions, exhibit respectively, in a marked degree, the characteristic venation of the mature plants.

e. *Litobrochia*.

11. *P. tripartita*, Sw. (Hook. l. c. p. 225, t. 138, B, cum syn.)—c.p. 1327 (1127).

HAB. Forests of the Central Province, not very common.

12. *P. aurita*, Blume. (Hook. l. c. p. 231, cum syn. An *P. incisæ*, Thunb. var. ?)—c.p. 1326.

HAB. Not uncommon in the Central Province.

## 25. CHEILANTHES, Sw.

1. *C. farinosa*, Kaulf. (Hook. Sp. Fil. ii. p. 77, cum syn.)—c.p. 2987 (1168).

HAB. More elevated parts of the Central Province.

2. *C. tenuifolia*, Sw. (Hook. l. c. p. 82, cum syn.)—c.p. 1322 (1167, 1237).

HAB. Common in the warmer parts of the island.

3. *C. Mysorensis*, Wall. (Hook. l. c. p. 94, cum syn.)—c.p. 1320 (1166).

HAB. Badulla and Doombera Districts.

4. *C. varians*, Hook. l. c. p. 89, t. 103, A, cum syn.—*C. laxa*, Moore, Ind. Fil. p. 245.—c.p. 1321.

HAB. Kornegalle District and Caloopahane.

## 26. ADIANTUM, Linn.

1. *A. lunulatum*, Burm. (Hook. Sp. Fil. ii. p. 11, cum syn.; Rheede, Hort. Mal. xii. p. 72, t. 40.)—c.p. 1323 (1238).

HAB. Hotter parts of the island.

2. *A. rhizophorum*, Sw. (Hook. l. c. p. 12, cum syn.)—c.p. 3102.

HAB. Oova and Doombera Districts.

3. *A. caudatum*, Linn. (Hook. l. c. cum syn.)—c.p. 978 (1124).

HAB. Very abundant in the warmer parts of the island.

4. *A. flabellulatum*, Linn. (Hook. l. c. p. 30, cum syn.)—c.p. 3390 (1239).

HAB. Oova District.

5. *A. hispidulum*, Sw. (Hook. l. c. p. 31, cum syn.)—c.p. 1325 (1023).

HAB. Not uncommon.

6. *A. Capillus-Veneris*, Linn. (Hook. l. c. p. 36, cum syn.)—c.p. 1324.

HAB. Oova and Doombera Districts.

27. **LINDSÆA**, Dryand.a. *Eulindsæa*.

1. **L. cultrata**, Sw. (Hook. Sp. Fil. i. p. 203, cum syn.)—c.p. 3070.

HAB. Abundant in the more elevated parts of the Central Province.

2. **L. repens**.—*L. oblongifolia*, Reinw.; Hook. l. c. p. 206. *Davallia Boryana*, Presl; Hook. l. c. p. 175, cum syn. *Acrophorus repens*, Moore, Ind. Fil. p. 3, cum syn. *Dicksonia repens*, Bory; Willd. Sp. Pl. v. p. 482.—c.p. 3389. Forma *minor*, lobis pinnarum longioribus.—c.p. 1389.

HAB. Singhe-rajah Forest. Forma *minor*, Matale, *Gardner*.

3. **L. Walkeræ**, Hook. l. c. p. 209, t. 99, A.—c.p. 1379.

HAB. Swampy grassy places in the Saffragam District.

4. **L. flabellulata**, Dryand. (Hook. l. c. p. 211, cum syn.)—c.p. 3311.

HAB. Forests of the Central Province.

5. **L. tenera**, Dryand. (Hook. l. c.)—c.p. 1381 (1259).

HAB. Not uncommon in the Central Province.

Closely allied to *L. flabellulata*, if not a form of it.

6. **L. caudata**, Hook. l. c. p. 215; Hook. Ic. Plant. x. t. 958, and Century of Ferns, t. 58.—c.p. 1380 (1262).

HAB. Forests of the Central Province, not uncommon.

b. *Schizoloma*.

7. **L. ensifolia**, Sw. (Hook. l. c. p. 220, cum syn.; Hook. et Grev. Ic. Fil. t. 3.)—c.p. 1382 (1131).

HAB. Common in the warmer parts of the island.

8. **L. nitens**, Blume, En. Fil. Jav. p. 217.—*L. recurvata*, Wall.; Hook. l. c. p. 222, cum syn. Et forma *bipinnata*, *L. obtusa*, J. Sm.; Hook. l. c. p. 224.—c.p. 982 (1260, 1261).

HAB. Very common in the forests of the Central Province.

The pinnae or pinnules, as the case may be, of this very variable plant are as frequently directed forwards as recurved, so that Blume's name is much more generally appropriated than Wallich's.

28. **DAVALLIA**, Linn.a. *Humata*.

1. **D. pedata**, Sw. (Hook. Sp. Fil. i. p. 154, t. 45, A, cum syn.)—c.p. 1383 (1113).

HAB. Common in the Central Province.

2. **D. vestita**, Blume, En. Fil. Jav. p. 233; Hook. l. c. p. 156, t. 41, C.—c.p. 3068.

HAB. Bopatalawa Plains, in the Central Province, at an elevation of 5000 to 6000 feet.

b. *Leucostegia*.

3. **D. pulchra**, Don. (Prod. Fl. Nep. p. 11; Hook. l. c. p. 160.)—*D. chærophylla*, Wall. Cat. n. 259; Hook. l. c. p. 157, t. 51, A, cum syn. *Acrophorus pulcher*, Moore, Ind. Fil. p. 3, cum syn.—c.p. 3288.

HAB. Forests above Galagama, upon rocks in damp gullies.

4. **D. affinis**, Hook. l. c. p. 158, t. 52, B.—*Acrophorus affinis*, Moore, l. c. p. 1.—c.P. 1384 (1117).

HAB. Very common in damp forests of the Central Province.

c. *Prosaptia*.

5. **D. Emersoni**, Hook. et Grev. (Ic. Fil. t. 105; Hook. l. c. p. 161, cum syn.)—c.P. 1385.

HAB. Not uncommon in forests of the Central Province.

6. **D. contigua**, Sw. (Hook. l. c. cum syn.)—c.P. 3069 (1114).

HAB. Common in the more elevated parts of the Central Province.

d. *Eudavallia*.

7. **D. elegans**, Sw. (Hook. l. c. p. 164, t. 43, A and B, cum syn.; Moore, l. c. p. 293.)—c.P. 3078.

HAB. Not very uncommon in the Central and Southern Provinces of the island, at no great elevation.

8. **D. tenuifolia**, Sw. (Hook. l. c. p. 186, cum syn.; Moore, l. c. p. 301.)—c.P. 983 (1115).

HAB. Very abundant.

e. *Microlepia*.

9. **D. strigosa**.—*D. Khasyana*, Hook. l. c. p. 173, tt. 47 A, 57 A. *Dicksonia strigosa*, Sw.; Hook. l. c. p. 81, cum syn. (Moore).—c.P. 1386 (1120).

HAB. Common in the forests of the Central Province.

10. **D. platyphylla**, Don. (Prod. Fl. Nep. p. 10.)—*D. lonchitidea*, Wall.; Hook. l. c. t. 46, B, cum syn.—c.P. 3277.

HAB. Rambodde, in the Central Province, Gardner.

11. **D. inæqualis**, Kze. (Hook. l. c. p. 180, cum syn.)

HAB. Ceylon; Walker, in Herb. Hook.

12. **D. hirta**, Kaulf. (Hook. l. c. p. 181, cum syn.)—c.P. 3272.

HAB. Forests of the Central Province, common.

13. **D. polypodioides**, Don. (Hook. l. c. cum syn.)—c.P. 1387, 1388 (1119).

HAB. Very abundant in the Southern and Central Provinces.

## 29. DICKSONIA, L'Hér.

### Subgenus PATANIA.

1. **D. deltoidea**, Hook. (Sp. Fil. i. p. 80, t. 28, A.)—*Dennstædia deltoidea*, Moore, Ind. Fil. p. 305.—c.P. 1397 (1118).

HAB. Most elevated parts of the Central Province.

## 30. ASPIDIUM, Sw.

### a. *Polystichum*.

1. **A. auriculatum**, Sw. (Hook. Sp. Fil. iv. p. 11, t. 218, cum syn.)—c.P. 1366 (1095).

HAB. Common in the forests of the Central Province.

2. **A. aculeatum**, Sw. (Hook. l. c. p. 18, cum syn.)—c.P. 1367, 3503 (1100, 1101).

HAB. More elevated parts of the Central Province.

3. **A. biaristatum**, Blume. (Hook. l. c. p. 29.)—c.P. 3275 (1102).

HAB. Maturatte, with *A. aculeatum*, of which I suspect it will prove to be merely a variety.

4. **A. amabile**, Blume. (Hook. l. c. p. 25, t. 225, cum syn.)—c.P. 3287.

HAB. More elevated parts of the Central Province.

5. **A. anomalum**, Hook. et Arn. (Hook. l. c. p. 27; excl. "c.P. 3286," formam *Polypod. Walkeræ*, Hook.)—c.P. 3504.

HAB. Horton Plains and Happootelle, at an elevation of 5000 to 6000 feet.

6. **A. aristatum**, Sw. (Hook. l. c. p. 27.)—c.P. 1373, 3384 (1098).

HAB. Very common in the Central Province.

#### b. *Euaspidium*.

7. **A. pteropus**, Kze. (Hook. l. c. p. 47, cum syn.)—c.P. 1355 (1088).

HAB. Common in damp forests.

8. **A. cicutarium**, Sw. (Hook. l. c. p. 48, cum syn.)—c.P. 1356.

Var.  $\beta$ ; *frondibus* firmioribus, *siccitate* nitidis, *venulis* margini frondis non vel rarius attingentibus.—c.P. 3331. An species distincta?

HAB. Central Province, not uncommon. Var.  $\beta$  near Badulla, *Mr. Ondaatchi*; near Galle.

9. **A. giganteum**, Blume (Hook. l. c. p. 50, cum syn.)—c.P. 1357 (1111).

Var.  $\beta$ ? *minor*, Hook. l. c.—c.P. 1358. An species distincta?

HAB. Very abundant in the Central Province. Var.  $\beta$ ? in the Doomberra district, at no great elevation.

The fronds of var.  $\beta$ ? are much smaller, more deeply lobed, and of much more delicate texture than those of the ordinary form.

10. **A. membranifolium**, Mett. *Aspid.* p. 113.—*Nephrodium* (*Lastrea*) *membranifolium*, Pr.; Hook. l. c. p. 131, t. 261, cum syn. An *A. Gardnerianum*, Mett.?—c.P. 3061, 3096 (1258).

HAB. Common in forests of the Central Province.

There would seem to be little, except the difference in the venation, to distinguish the present species from *A. giganteum*. When growing together, the two plants look precisely alike, when seen from some little distance.

11. **A. subtriphyllum**, Hook. l. c. p. 52, cum syn.—c.P. 1300.

HAB. Forests of Allagalla and Maturatte, in the Central Province.

12. **A. polymorphum**, Wall. (Hook. l. c. p. 54, cum syn.)—c.P. 1354 (1096).

HAB. Common in the Central Province.

#### c. *Pleocnemia*.

13. **A. Otaria**, Kze. (Mett. *Asp.* p. 34.)—*Nephrodium* (*Pleocnemia*) *aristatum*, Hook. l. c. p. 62, t. 238, cum syn. *Anisocampium Cumingianum*, Pr. *Epim. Bot.* p. 58.—c.P. 1299.

HAB. Bintenne, *Gardner*; Gonagama, north of Kandy.

d. *Nephrodium*.

14. **A. molle**, Sw.—*Nephrodium molle*, Desv.; Hook. l. c. p. 67, cum syn.—C.P. 714, 974.

HAB. Very abundant throughout the island.

15. **A. extensum**, Blume, En. Fil. Jav. p. 156.—*Nephrodium extensum*, Bl.; Hook. l. c. p. 72, cum syn.—C.P. 3391, 3498.

HAB. Oova District.

16. **A. Arbuscula**, Willd.—*Nephrodium Arbuscula*, Desv.; Hook. l. c. p. 74.—C.P. 1359 (1109).

HAB. Common in the Central Province.

17. **A. terminans**, Wall. Cat. n. 386.—*Nephrodium terminans*, J. Sm.; Hook. l. c. p. 73, cum syn.—C.P. 990 (1106).

HAB. Central Province, not uncommon.

18. **A. propinquum**.—*Nephrodium propinquum*, R. Br.; Hook. l. c. p. 79, cum syn.—C.P. 705.

HAB. South of the island, at no great elevation.

19. **A. unitum**, Mett. Asp. p. 107.—*Nephrodium unitum*, Sieb.; Hook. l. c. p. 81, cum syn.—C.P. 973.

HAB. A common species in the island.

20. **A. eusorum**, Thw.—*A. pennigerum*, Blume, l. c. p. 153 (haud Sw.).—C.P. 3064 (1104, 1252).

HAB. Forests of the Central Province, not uncommon.

*Caudex* brevis. *Fronde*s glabræ vel subglabræ, lanceolatae, acuminatae, pinnatae. *Pinnæ* numerosæ, lineares, acuminatae, 5–10 poll. longæ,  $\frac{1}{2}$ –1 poll. latae, sessiles, versus basin abruptam plus minus angustatae (paucæ inferiores subito parvulae, rotundatae, distantes), lobis mediocribus subintegris, abrupte truncatis, venulis pluribus, omnibus medio soriferis, 2 inferioribus infra sinum unitis. *Indusium* parvum. [This, I fear, is identical with what I consider *Nephrodium abruptum*, Sp. Fil. iv. p. 79; from which, however, the syn. of *A. pennigerum* should be excluded.—Hook.]

21. **A. abruptum**, Blume, l. c. p. 152.—*Nephrodium abruptum*, Pr.; Hook. l. c. p. 77, t. 241, B.—C.P. 3271 (1224).

HAB. Damp forests of the Central Province.

e. *Lastrea*.

22. **A. calcaratum**, Blume, l. c. p. 159.—*Nephrodium (Lastrea) calcaratum*, Hook. l. c. p. 93, cum syn. *N. (Lastrea) falcilobum*, Hook. l. c. p. 108.—C.P. 1363, 3273 (1107, 1250).

Var.  $\beta$ . Hook. l. c. p. 108.—C.P. 992, 3050.

HAB. On the sides of streams and in damp forests, common.

A most variable plant, but the extreme forms pass insensibly into one another.

23. **A. flaccidum**, Blume, En. Fil. Jav. p. 161.—*Nephrodium (Lastrea) flaccidum*, Hook. Sp. Fil. iv. p. 133, t. 263.—C.P. 3802.

HAB. More elevated parts of the Central Province.

24. **A. gracilescens**, Blume, l. c. p. 155.—*Nephrodium (Lastrea) gracilescens*, Hook. l. c. p. 93, cum syn. *A. Thelypteris*, Benth. Flor. Hongk. p. 455.—C.P. 1287 (1141).

HAB. More elevated parts of the Central Province.

25. **A. tylodes**, Kze. in Linn. xxiv. p. 281.—*Lastrea tylodes*, Moore

(fide specim. ex Herb. Ind. Orient. Hook. f. et Thoms.).—c.p. 1361 (1108).

HAB. Newera Ellia, and other more elevated parts of the Central Province.

*Sori* prope costulam siti.

26. **A. ochthodes**, Kze. l. c. p. 282.—*Nephrodium* (*Lastrea*) *ochthodes*, Hook. l. c. p. 109 partim. *A. glanduliferum*, Wall. Cat. n. 347.—c.p. 975 (1362).

HAB. Not uncommon in forests of the Central Province.

Very like *A. tylodes*, from which it is at once distinguished by the sori being situated nearer to the margin than to the costule of the pinna-lobes.

27. **A. spectabile**, Blume, l. c. p. 158.—*Nephrodium* (*Lastrea*) *spectabile*, Hook. l. c. p. 115, cum syn.—c.p. 3274.

HAB. Higher forests of the Central Province.

28. **A. hirtipes**, Blume, l. c. p. 148.—*Nephrodium* (*Lastrea*) *hirtipes*, Hook. l. c. p. 115, cum syn. *Aspidium atratum*, Wall. Cat. n. 380.—c.p. 1291 (1091).

HAB. Common in the upper forests of the Central Province.

29. **A. Filix-mas**, Sw. (Willd. Sp. Pl. v. p. 259.) — *Nephrodium* (*Lastrea*) *Filix-mas*, Rich. Var.  $\beta$ , Hook. l. c. p. 116, cum syn.—c.p. 1364 (1092).

HAB. Newera Ellia and other of the most elevated parts of the Central Province.

30. **A. concinnum**, Thw.; *caudice* brevi; *frondibus* lanceolatis vel triangulari-lanceolatis, teneris, glabris, nitidis, cum stipite stramineo basi paleaceo æquilongè bipedalibus; *pinnis* lanceolatis, acuminatis; *pinnulis* trapezoideo-lanceolatis, crenato-lobatis, *lobis* parce denticulatis, *venis* paucis paulo intra marginem terminantibus et ad apices soriferis; *soris* parvis; *indusio* glabro, rotundato.—c.p. 3798.

HAB. Singhe-rajah Forest.

Very similar in general appearance to some forms of *Aspidium sparsum*, Don, but distinguished by its smaller sori, and these being much nearer to the margin of the pinnule than in that species. [*A. sparsum*, Sp.; *A. purpurascens*, Bl., is very variable, and some of my specimens have the sori very near the margins.—Hook.]

31. **A. sparsum**, Spr. Syst. iv. p. 106.—*A. purpurascens*, Blume, l. c. p. 169. *Nephrodium sparsum*, Don, Prod. Fl. Nep. p. 6. *N. (Lastrea) purpurascens*, Hook. l. c. p. 132, t. 262, cum syn.—c.p. 1375.

Var.  $\beta$ ; *frondium* pagina superiore nitida, *pinnulis* brevioribus, magis deltoideis, *soris* margine propioribus; *indusio* siccitate plicato.—*Lastrea sparsa*, var. *lata*, Moore, MSS.—c.p. 1368, 1369? 1370, 3383. An species distincta?

HAB. Very abundant in the forests of the Central Province.

An extremely variable species.

32. **A. divisum**, Wall. Cat. n. 393.—*Nephrodium* (*Lastrea*) *divisum*, Hook. l. c. p. 133, cum syn.—c.p. 3097.

HAB. Forests at Raxawa and Happootelle, in the Central Province.

33. **A. Blumei**.—*Nephrodium* (*Lastrea*) *Blumei*, Hook. l. c. p. 135, cum syn.—c.p. 3059 (1280).

Var. ? *frondibus* firmioribus, siccitate nigrescentibus, subtus minute glanduloso-pilosis, nec paleaceis.—c.p. 3142. An species distincta?

HAB. Common in the Central Province, in damp forests. Var. ? at Newera Ellia and Maturatte.

The two forms indicated above are much alike in general appearance, but I think will prove to be specifically distinct.

34. **A. recedens**, Mett.—*Nephrodium (Lastrea) recedens*, Hook. l. c. p. 135, t. 265, cum syn.—C.P. 1374 (1110).

HAB. Forests of the Central Province.

35. **A. tenericaule**, *Nephrodium (Lastrea) tenericaule*, Hook. l. c. p. 142, t. 269, cum syn.—C.P. 1286, 1365 (1150, 1222).

HAB. Very abundant in the Central Province.

f. *Diclisodon*.

36. **A. deparioides**, Hook. Fil. Exot. i. t. 3.—*Nephrodium (Lastrea) deparioides*, Hook. Sp. Fil. iv. p. 139. *Diclisodon deparioides*, Moore, Ind. Fil. p. 316.—C.P. 3062.

HAB. Forests of the Central and Southern Provinces, up to an elevation of 3000 feet.

31. **NEPHROLEPIS**, Schott.

1. **N. tuberosa**, Pr. (Hook. Sp. Fil. iv. p. 151, cum syn.)—C.P. 1377 (1090).

HAB. Common throughout the island.

2. **N. exaltata**, Schott, (Hook. l. c. p. 152, cum syn.)—C.P. 3077 (1089).

HAB. Very abundant on the margins of woods, sides of streams, etc.

3. **N. acuta**, Pr. (Hook. l. c. p. 153, cum syn.)—C.P. 991.

HAB. Equally abundant with the preceding, of which I believe it to be only a luxuriant form.

4. **N. obliterated**, Hook. l. c. p. 154, cum syn.—C.P. 1376 (1094).

HAB. A very common plant.

32. **OLEANDRA**, Cav.

1. **O. neriiformis**, Cav. (Hook. Sp. Fil. iv. p. 156; Hook. Fil. Exot. t. 58).—C.P. 1378 (1087).

HAB. Upper forests of the Central Province.

33. **POLYPODIUM**, Linn.

a. *Eupolypodium*.

1. **P. parasiticum**, Mett. (Hook. Sp. Fil. iv. p. 167, cum syn.)—C.P. 1283.

HAB. Central Province, and in Pas-doon-Corle.

2. **P. Zeylanicum**, Mett. (Hook. l. c. p. 169, t. 272 B.)—C.P. 3074 (1138, 1283).

HAB. Common in the more elevated parts of the Central Province.



3. **P. cucullatum**, Nees. (Hook. l. c. p. 176, cum syn.)—*Calymmodon cucullatus*, Pr.—c.p. 1282.

HAB. Higher forests of the Central Province, not common.

4. **P. parvulum**, Bory, (Hook. l. c. p. 184, t. 274 B.)—c.p. 1290, 3073.

HAB. Common on the trunks of trees in the higher parts of the Central Province.

Distinguished from *P. obliquatum*, Bl., small forms of which it greatly resembles, by the sori not being immersed as in that species. The fronds are frequently hispidulous beneath.

5. **P. glandulosum**, Hook. l. c. p. 193, t. 276 A.—c.p. 1289.

HAB. Rambodde, in the Central Province, *Gardner*.

6. **P. obliquatum**, Blume, (Hook. l. c. p. 190, cum syn.)—c.p. 1284 (1146).

HAB. Very abundant on the trunks of trees in the more elevated parts of the Central Province.

7. **P. elongatum**, Wall. (Hook. l. c. p. 234).

Forma *a*. *frondibus* pinnatis, *pinnis* crenatis vel brevilobatis, lobis rotundatis subdenticulatis.—c.p. 3276 *a* (1256).

Forma *b*. *frondibus* pinnato-pinnatifidis vel subbipinnatis.—*P. Walkeræ*, Hook. l. c. p. 233.—c.p. 3276 *b* (1256).

Forma *c*. *frondibus* bi-tri-pinnatis.—c.p. 3286; an species distincta?

HAB. Var. *a* and *b*. Carawitte Kanda and Ambagamowa; var. *c*, Ambagamowa.

Var. *c*, although very closely allied to the other two varieties, and, in some instances, hardly to be distinguished from var. *b*, except by its more composite structure, is probably specifically distinct, the fronds assuming a different tint of brown when dry.

8. **P. paludosum**, Blume, (Hook. l. c. p. 245, cum syn.)—c.p. 1288 (1151).

HAB. More elevated parts of the Central Province.

9. **P. rufescens**, Blume, (Fil. Jav. p. 194, t. 91; Hook. l. c. p. 257.)—c.p. 3143 (1148).

HAB. Forests of the Central Province, at an elevation of 3000 to 5000 feet.

[A species hitherto unknown to me; Mr. Thwaites's n. 3143 sufficiently corresponds in habit and form with Blume's figure, but the specimens are smaller, less compound, and more membranaceous.—Hook.]

10. **P. rugulosum**, Labill. (Hook. l. c. p. 272, cum syn.)—c.p. 1285 (1149).

HAB. Common in the Central Province.

#### b. *Goniopteris*.

11. **P. granulosum**, Pr. (Benth. Flor. Hongk. p. 459, cum syn.)—*P. wrophyllum*, Wall. Cat. n. 299; Hook. l. c. v. p. 9.—c.p. 3063 (1137, 1223).

HAB. Forests of the Central Province, not uncommon.

#### c. *Pleopeltis*.

12. **P. Wightianum**, Wall. Cat. n. 2222. (Benth. l. c. p. 458, cum

syn.)—*P. loriforme*, Wall. ; Hook. l. c. v. p. 57. *Pleopeltis nuda*, Hook. Ex. Fl. t. 63.—C.P. 1295 (1139).

HAB. Common in forests of the Central Province.

13. *P. tridactylon*, Wall. (Hook. and Grev. Ic. Fil. t. 209 ; Hook. Sp. Fil. l. c. v. p. 75 ; Benth. l. c.)—C.P. 1301.

HAB. Muruta, *Gardner* ; Ambagamowa District, at no great elevation.

14. *P. membranaceum*, Don, Prod. Fl. Nep. p. 2 ; Hook. l. c. v. p. 79.—*Pleopeltis*, Moore.—C.P. 1298 (1145).

HAB. Common on the trunks of trees, etc., in damp forests of the Central Province.

15. *P. nigrescens*, Blume, En. Fil. Jav. p. 126 ; Hook. Fil. Exot. t. 22 ; Hook. l. c. v. p. 81.—*Phymatodes saccata*, J. Sm. Cat. Cult. Ferns, p. 9.—C.P. 1296 (1144).

HAB. Common in forests of the Central Province.

16. *P. Phymatodes*, Linn. (Willd. ; Hook. l. c. v. p. 82 ; Blume, l. c. p. 125).—C.P. 1297.

HAB. Common in the warmer parts of the island.

17. *P. oxylobum*, Wall. Cat. ; Hook. l. c. v. p. 77.—*Phymatodes hastata*, Moore ; *P. longipes*, J. Sm. l. c.—An *Polypod. hastatum*, Thunb. (*P. Cyrtoloma*, Wall. Cat.) forma?—C.P. 3291 (1143).

HAB. More elevated parts of the Central Province.

18. *P. irioides*, Poir. (Spreng. Syst. iv. p. 48 ; Hook. l. c. v. p. 67.—C.P. 3799.

HAB. Ettangwelle and Halgodde Pass, Matelle East.

#### d. *Drynaria*.

19. *P. quercifolium*, Linn. (Hook. l. c. v. p. 96).—*Drynaria quercifolia*, Bory. *Phymatodes quercifolia*, Pr.—C.P. 3265 (1142).

HAB. Very abundant upon old trees in the warmer parts of the island.

#### e. *Niphobolus*.

20. *P. Gardneri*, Mett. Polyp. p. 129 ; (Hook. l. c. v. p. 51).—C.P. 988 (1154).

HAB. Not uncommon.

*Rhizoma*, cum *stipitum* basi, paleis firmis erectis lanceolatis fusco-nigris margine fulvis tectum.

21. *P. porosum*, Wall. Cat. n. 260 ; Hook. l. c. v. p. 48.—*Niphobolus porosus*, Pr.—C.P. 3104 (1226).

HAB. Forests of the Central Province, at no great elevation, not common.

22. *P. Lingua*, Sw. (Willd. Sp. Pl. v. p. 162 ; Hook. l. c. v. p. 49).—*Acrostichum Lingua*, Thunb. Fl. Jap. p. 330, t. 33.—C.P. 1294.

HAB. Not uncommon in the warmer parts of the island, and up to an elevation of 3000 feet.

23. *P. adnascens*, Sw. ; Willd. l. c. p. 145 ; Hook. l. c. v. p. 47 ; Benth. Fl. Hongk. p. 458, cum syn.—*P. pertusum*, Roxb. ; Hook. Ex.

Fl. t. 162. *Niphobolus adnescens*, Kaulf. *N. carnosus*, Bl. *N. pertusus*, Spr.—c.p. 993, 3293 (1153).

HAB. Very common upon the trunks of trees.

[N.B. *P. acrostichoides*, Sw. Hook. l. c. v. p. 44, is considered a native of Ceylon, on the authority of a specimen from *Gardner*.]

### 34. DIACALPE, Blume.

1. **D. aspidioides**, Blume. (Hook. Sp. Fil. i. p. 59, cum syn.)—c.p. 3282.

HAB. "Black Pool" in Newera Ellia, *Dr. E. Bradford*.

### 35. CYATHEA, Sw.

1. **C. sinuata**, Hook. et Grev. Ic. Fil. t. 2; Hook. Sp. Fil. i. p. 15; Gard. Ferns, t. 21.—c.p. 3052.

HAB. Forests in the Kokoole and Hinidoon Corles, at no great elevation; most abundant in the Singhe-rajah Forest.

2. **C. Hookeri**, Thw.; *caudice* elongato; *stipitibus* basi parce muricatis et paleis rigidiusculis setoso-acuminatis munitis; *frondibus* anguste lanceolatis, acuminatis, pinnatis, apice pinnatifidis; *pinnis* lanceolatis subintegris vel crenato-lobatis, basi latioribus truncatisque, brevipe-tiolatis, superioribus sessilibus, inferioribus gradatim minoribus, subtus ad costam parce paleaceis; *venis* simplice pinnatis, ramis inferioribus supra sinum terminantibus, prope basin soriferis; indusio globoso, irregulariter fisso.—c.p. 3722.

HAB. Singhe-rajah Forest.

*Caudex* circiter  $1\frac{1}{2}$  poll. in diam., basibus nigris frondium delapsarum tectus. *Fronde*s 3-pedales vel ultra, circiter 6 poll. latæ. [A small, for an arborescent Fern, but a very distinct species, now happily in cultivation at Kew.—Hook.]

3. **C. Walkeræ**, Hook. l. c. p. 24; Hook. Ic. Plant. vii. t. 647.—c.p. 1398, 1399 (1054, 1400).—Forma *indusiis* majoribus, subregularibus.—c.p. 3053.

HAB. Common in the Central Province, up to an elevation of 6000 feet.

Varies greatly in the more or less coriaceous texture of its fronds, and in the amount of development of the indusium, which is sometimes hardly to be detected. This species seems to be very closely allied indeed to *C. spinulosa*, Wall.

### 36. ALSOPHILA, R. Br.

1. **A. gigantea**, Wall. (Hook. Sp. Fil. i. p. 53, cum syn.)—c.p. 1403.

HAB. Not uncommon in the Central and Southern Provinces, up to an elevation of 4000 feet.

2. **A. crinita**, Hook. l. c. p. 54; Hook. Ic. Plant. t. 671.—c.p. 1401.

HAB. Newera Ellia and other of the more elevated parts of the Central Province.

## 37. HYMENOPHYLLUM, Sm.

1. **H. corticola**, Hook.—*H. Lyallii*, Moore, in Litt. (but certainly not *H. Lyallii*, Hook. fil. Nov. Zeal. ii. p. 16).—c.p. 3278 (1265).

HAB. Ambagamowa District.

*Rhizoma* capillare, repens, et, cum *stipitibus* longiusculis, hispidulum. *Fronde*  $\frac{1}{2}$ –1-pollicares, cuneato-orbiculares, palmato-incisæ, glabræ, *segmentis* linearibus, retusis, costa centrali, margine non incrassata. *Sori* terminales, *involucro* poculiformi, subbilabiato.

2. **H. Tunbridgense**, Sm. (Hook. l. c. p. 95, cum syn.)—c.p. 2984.

HAB. Common in the more elevated parts of the Central Province.

3. **H. crispatum**, Wall. (Hook. l. c. p. 105, cum syn.)—c.p. 1394, 1395.

HAB. Very abundant in the forests of the Central and Southern Provinces.

4. **H. polyanthos**, Sw. (Hook. l. c. p. 106, cum syn.)—c.p. 3279, 3360 (1264).—Van den Bosch, in his second supplement to the 'Hymenophyllaceæ,' refers to Mr. Thwaites's n. 3360, for his *H. macroglossum* (Hook).

HAB. Upper forests of the Central Province.

5. **H. Blumeanum**, Spr. Syst. iv. p. 131; Blume, En. Fil. Jav. p. 220; Hook. Sp. Fil. i. p. 147; Van den Bosch, Hymenoph. Jav. p. 46, t. 36.—c.p. 1391.

HAB. Forests of the Southern and Central Provinces.

6. **H. exsertum**, Wall. (Hook. l. c. p. 109, t. 38 A, cum syn.)—c.p. 1390.

HAB. Very common in the upper forests of the Central Province.

## 38. TRICHOMANES, Linn.

1. **T. muscoides**, Sw. (Hook. Sp. Fil. i. p. 117, cum syn.)—c.p. 2986.

HAB. Raxawa, in the Central Province, *Prof. W. H. Harvey*; Palagalla.—Thwaites's n. 2986, Van den Bosch refers to *T. bimarginatum*, V. d. Bosch.

2. **T. proliferum**, Blume, (Hook. l. c. p. 118, t. 39 B, cum syn.)—c.p. 3329.

HAB. Ambagamowa District.

3. **T. intramarginale**, Hook. et Grev. Ic. Fil. t. 211; Hook. Sp. Fil. l. c. p. 120.—c.p. 3361.

HAB. Ambagamowa District.

4. **T. Filicula**, Bory, (Hook. l. c. p. 124, cum syn.)—c.p. 2985 (1266).

HAB. Common in the forests of the Central and Southern Provinces.

5. **T. glauco-fuscum**, Hook. l. c. p. 128, t. 40 A, cum syn.—c.p. 3330.

HAB. Ambagamowa District.

6. **T. rigidum**, Sw. (Hook. l. c. p. 133, cum syn.)—*T. obscurum*, Blume, En. Fil. Jav. p. 227.—c.p. 1396 (1174).

HAB. Upper forests of the Central Province.

## ADDENDA ET CORRIGENDA.

Page

2.

## "ACROTREMA, Jack."

In descriptione, post "*Carpella*" insere: regulariter vel.

3. "A. Walkeri, Wight." Adde:

Var.  $\beta$ . *argyroneurum*; foliis majoribus, costa venisque superne depressis, albis.—C.P. 3831. An species distincta?

HAB. Pasdoon Corle.

5. "Dillenia integra, Thunb." Adde: certe *D. retusæ*, Thunb., varietas, fide speciminum ex Herb. Hort. Bot. Calcuttensis.7. "Goniothalamus reticulatus, Thw." Adde: Flores, quoad formam coloremque, iis *G. salicini* simillimi, sed majores quam in illa specie.

8. "Goniothalamus Walkeri, Hook. f. et Thoms." Adde:—C.P. 3825.

HAB. Singhe-rajah Forest. Nom. vulg. "Kappooroo-gass."

Arbor parva, erecta, vix ramosa. Cortex rubro-nigrescens. Folia juniora rubentia. Flores rubri, sparsissime pilosuli. Sepala ovata, acuta, 2 lin. longa. Petala exteriora ovato-lanceolata, 8 lin. longa, interiora 5 lin. longa. Stylus clavatus, stigmate bilobo.

The fragrant roots contain camphor, and are chewed by the Cinghalese.

9. "Xylophia nigricans, Hook. f. et Thoms." Adde:

HAB. Trincomalie, Rev. S. O. Glenie. Doombera District.

9.

## "UNONA, Linn." Adde:

3. *U. elegans*, Thw.; arborea vel fruticosa, ramosissima; foliis parvis, anguste lanceolatis, longe acuminatis, subtus glaucis; floribus ut in *U. Zeylanica* sed minoribus; carpellis inter semina multum constrictis.—C.P. 3824.

HAB. Hinidoon Pattoo.

9.

## "POLYALTHIA, Blume." Adde:

Sect. 2.—*Carpella monosperma*.

(*Guatteria*, Hook. f. et Thoms., et hujus Enum., sed non Ruiz et Pavon.)

Vide Benth. et Hook. f. Gen. Plant. p. 25.

2. *P. longifolia*, Benth. et Hook. f. l. c.—*Guatteria longifolia*, Wall., hujus Enum. p. 10.—C.P. 1041.

3. *P. suberosa*, Benth. et Hook. f. l. c.—*Guatteria suberosa*, Dun., hujus Enum. p. 10.—C.P. 1035.

4. *P. Corinti*, Benth. et Hook. f. l. c.—*Guatteria Corinti*, Dun., hujus Enum. p. 10.—C.P. 1034.

Page

5. **P. persicæfolia**, Benth. et Hook. f. l. c.—*Guatteria persicæfolia*, Hook. f. et Thoms., hujus Enum. p. 10.

6. **P. coffeoides**, Benth. et Hook. f. l. c.—*Guatteria coffeoides*, Hook. f. et Thoms., hujus Enum. p. 10.—c.p. 2503.

7. **P. acuminata**, Thw.; arborea, ramulis petiolisque pilosis; foliis majusculis, oblongis vel obovato-oblongis, rostrato-acuminatis, subglabris, ad costam canaliculatis, basi angustatis, brevipetiolatis; floribus caulinis, majusculis, subsolitariis vel paucis, tomentellis, longiuscule pedicellatis; petalis ovato-lanceolatis, acuminatis; carpellis ovoideis, singulis stipite subæquilongo.—c.p. 3653.

HAB. Marai-calai, near Ratnapoora.

Cortex fusco-niger, rugosus. Folia 6–12 poll. longa, 2½–3 poll. lata, siccitate fuscescentia; petiolo 3–4 lin. longo. Pedicelli 1–2½ poll. longi, fulvo-tomentosi, infra medium bractea parva muniti. Flores ochracei. Sepala 4 lin. longa, extus fulvo-tomentosa. Petala 13 lin. longa, exteriora 5 lin., interiora 6 lin. lata. Torus convexus, tomentosus. Stylus ovario piloso æquilongus. Stigma pilosum. Carpella nucis Avellanæ magnitudine, tomentosa; semine conformi, nitido, fulvo.

11. “**ALPHONSEA**, Hook. f. et Thoms.” Adde:

3. **A. lutea**, Hook. f. et Thoms. Flor. Ind. p. 153, cum syn.; Roxb. Pl. Cor. t. 36.—c.p. 3826.

HAB. Reigam Corle.

Petala externe pilosa, interiora interne omnino, exteriora dimidio inferiore glabra.

11. “**Myristica Zeylanica**, Alph. DC.”—Certe *M. laurifolia*, Hook. f. et Thoms., varietas vel forma.

13. “**Cissampelos subpeltata**, Thw. ;” vix species distincta, sed inter varietates numerosas *C. Pareiræ* rectius includenda.

14. “**CRUCIFERÆ**.” Adde:

## 2. NASTURTIUM, R. Br.

1. **N. Indicum**, Linn. (Hook. f. et Thoms. Præcurs. ad Fl. Indic. in Linn. Soc. Journ. of Proc. v. p. 138, cum syn.)—c.p. 3747.

HAB. Central Province, not common.

## 3. BRASSICA, Linn.

1. **B. juncea**, Linn. (Hook. f. et Thoms. l. c. p. 170, cum syn.)—c.p. 3772.

HAB. Common as a weed, and cultivated. Nom. vulg. “Abba.”

14. “**CARDAMINE**, Linn.”

“**C. hirsuta**, Linn.”—Dele “var. major” et seq. ad “c.p. 1017,” una cum “The trifoliolate variety occurs in shady woods,” et adde:

2. **C. Africana**, Linn. (Hook. f. et Thoms. l. c. p. 144, cum syn.)—*C. hirsuta*, var. major, hujus Enum. p. 14.—c.p. 1017.

HAB. Shady woods in the more elevated parts of the Central Province.

15. “**CAPPARIS**, Linn.” Adde: •

11. **C. floribunda**, Wight, Illust. i. p. 35, t. 14; Walp. Rep. i. p. 199.—c.p. 3766.

HAB. Doombera District, not common.

Page

16. "**C. retusella**, Thw.;" verisimiliter *C. sepiariae*, Linn., varietas, *carpopphoro* subduplo longiore quam in forma communi illius speciei.
16. Pro "2. **PHOBEROS**, Lour." substitue:  
 2. **SCOLOPIA**, Spr.  
*Phoberos*, Lour. Et adde:  
 1. **S. Chinensis**, Clos, in Ann. des Sc. Nat. ser. 4, viii. p. 249, cum syn. — *Phoberos Chinensis*, Lour.; forma arborea, trunco spinis compositis armato. *P. Arnoltianus*, Thw. in hac Enum. p. 16.—c.p. 3526.  
 2. **S. Gærtneri**, Thw.—*S. pusilla*, Clos, l. c. *Phob. Gærtneri*, Thw., in hac Enum. p. 17. Var. *a.*—c.p. 2497. Var. *β.*—c.p. 1076. Var. *γ.*—64 (181,211).  
 3. **S. acuminata**, Clos, l. c. p. 251.—*Phob. acuminatus*, Gard. MSS., hujus Enum. p. 17.—c.p. 1077.  
 4. **S. crassipes**, Clos, l. c.—*Phob. Hookerianus*, Wight, MSS., hujus Enum. p. 17.—c.p. 629.
18. Pro "4. **ROUMEA**, Poit.," substitue:  
 4. **ABERIA**, Hochst.  
*Roumea*, Poit. Et adde:  
 1. **A. Gardnerii**, Clos, in Ann. des Sc. Nat. ser. 4, viii. p. 236.—*Roumea hebecarpa*, Gardner, hujus Enum. p. 18.—c.p. 1075.
22. "**POLYGALA**, Linn." Adde:  
 7. **P. arvensis**, Willd. (W. et A. Prod. i. p. 36, cum syn.)—*P. rosmarinifolia*, var. *a.*, hujus Enum. (non W. et A.) p. 22. Forma *a*; foliis obovato-oblongis, subglabris.—c.p. 715. Forma *b*; foliis lineari-oblongis vel linearibus, subglabris.—c.p. 1083. Forma *c*; tota planta sparsim rigide ciliata, foliis obovatis vel obovato-oblongis.—c.p. 3674.  
 HAB. Not very uncommon in the hotter parts of the island; forma *c*, Trincomalie, *Rev. S. O. Glenie*.
22. "**P. rosmarinifolia**, W. et A.;" dele descriptionem a "Var. *a.*" ad finem, et pro ea insere:  
 HAB. Common in the Oova District.  
 Formæ *b*, *P. arvensis* valde similis, sed probabiliter distincta.
22. Dele "3. **P. elongata**, Klein," et seq. ad "et 592.," et insere:  
 3. **P. glomerata**, Lour. Fl. Coch. ii. p. 426; DC. Prod. i. p. 326; Miq. Fl. Ind. Bat. i. pars altera, p. 125.—*P. densiflora*, Blume, Bijdr. p. 59.  
 Var. *a. pedunculosa*, racemis plus minus elongatis.—c.p. 592 (7, 260).
23. "**XANTHOPHYLLUM**, Roxb."  
 Dele "1. **X. flavescens**, Roxb. (W. et A. Prod. i. 39, cum syn.)" et insere:  
 1. **X. virens**, Roxb.; Fl. Ind. ii. p. 221.
23. Dele "1. **TRICHAURUS** (Arn. MSS.), W. et A.;" una cum "**T. ericoides**, Arn. MSS." et seq. ad "c.p. 1254." Et insere:

Page

1. **TAMARIX**, Linn.

1. **T. Gallica**, Linn. (W. et A. Prod. i. p. 40, cum syn.)—*T. Indica*, Willd.—c.p. 1254.

23. “**PORTULACA**, Tourn.” Adde :

4. **P. tuberosa**, Roxb. (W. et A. Prod. i. p. 356, cum syn.)—c.p. 3638.

HAB. South of the island.

26. “2. **Pavonia Zeylanica**, Cav.” Adde : c.p. 3461.

HAB. Trincomalie, *Rev. S. O. Glenie*; Colombo, *Mr. W. Ferguson*. Et dele :

“I have not met,” et seq. ad annotationis finem.

26. “**HIBISCUS**, Linn.” Adde :

8. **H. lunarifolius**, Willd. (W. et A. Prod. i. p. 49, cum syn.)—c.p. 3634.

HAB. Doombera District.

9. **H. cannabinus**, Linn. (W. et A. Prod. i. p. 50, cum syn.)—c.p. 3833.

HAB. Near Trincomalie, *Rev. S. O. Glenie*.

27. “**ABUTILON**, Dill.” Adde :

6. **A. crispum**, G. Don. (W. et A. Prod. i. p. 56, cum syn.)—c.p. 3646.

HAB. Colombo, *Mr. W. Ferguson*.

*Abutilon* (*Sida*, DC.) *Leschenaultianum*, c.p. 3521, occurs as a weed in the Botanic Garden, but I have never found it truly wild.

30. Dele “6. **KYDIA**, Roxb.” Et insere :

6. **DICELLOSTYLES**, Benth.

Benth. et Hook. f. Gen. Plant. p. 207. Et adde :

1. **D. axillaris**, Benth. l. c.—*Kydia axillaris*, Thw. in hac Enum. p. 30.—c.p. 3126.

*Petala* pollicaria, oblonga, subspathulata, parum obliqua, ochraceo-rufescentia, versus basin rubra.

31. “**CORCHORUS**, Linn.” Adde :

5. **C. fascicularis**, Lam. (W. et A. Prod. i. p. 72, cum syn.)—c.p. 3773.

HAB. Colombo, *Mr. W. Ferguson*.

6. **C. tridens**, Linn. (W. et A. Prod. i. p. 73, cum syn.)—c.p. 3791.

HAB. Trincomalie, *Rev. S. O. Glenie*.

31. “**TRIUMFETTA**, Plum.” Adde :

1. **T. pilosula**, Thw.; erecta, ramis scabriusculis; foliis caulinis rotundatis, parum trilobis, grosse crenatis, sparse pilosulis, subtus pallidioribus, basi acutis, 3- vel sub-5-nerviis, floralibus lanceolatis, utrinque acuminatis; petiolo elongatulo; cymulis densis, oppositifoliis; sepalis linearibus, apice cucullatis mucronatisque, externe pilosis; petalis calyce æquilongis; staminibus circiter 6,



Page

*filamentis* glabris; ovario 4-loculari; capsula pilosula, echinulis suberectis, uncinatis, fere ad apicem arcte ciliatis.—C.P. 2901.

HAB. Trincomalie, Rev. S. O. Glenie.

31. "GREWIA, Juss." Adde:

8. **G. rotundifolia**, Juss. (W. et A. Prod. i. p. 80, cum syn.)—C.P. 3785.

HAB. Trincomalie, Rev. S. O. Glenie.

33. "Dipterocarpus Zeylanicus, Thw." Adde: (C.P. 2398.)

34, lin. 10 ab imo. Dele: "*D. turbinatus*, Roxb., cultivated in Ceylon, is C.P. 2398."

34. "DOONA, Thw." Adde:

8. **D. macrophylla**, Thw.; glabra; foliis majusculis, oblongis, obtuse acuminatis, ad costam canaliculatis, basi rotundatis, petiolatis; paniculis axillaribus terminalibusque, apertis, folio subæquilongis; pedicellis calycem subæquantibus; calycis fructiferi (immaturi) laciniis obovato-oblongis.—C.P. 3713.

HAB. Hinidoon Pattoo.

Arbor magna. Folia firma, 5–8 poll. longa, 3 poll. lata, minutissime parallelim venulosa, venis primariis utrinque circiter 13, petiolo  $\frac{1}{2}$ –1 poll. longo. Petala oblonga, 4 lin. longa, externe et interna margine superiore pilosa.

9. **D. ovalifolia**, Thw.; glabra; foliis ovalibus, obtuse rostrato-acuminatis, superne subeveniis, subtus costa venis primariisque prominentibus, basi rotundatis, petiolatis; paniculis axillaribus terminalibusque, racemiformibus, paucifloris, folio brevioribus vel subæquilongis; pedicellis calycem subæquantibus.—C.P. 3711.

HAB. Saffragam District.

Arbor magna. Folia (acumina 4 lin. longa)  $1\frac{1}{2}$ –3 poll. longa,  $\frac{3}{4}$ – $1\frac{1}{2}$  poll. lata, venis primariis utrinque circiter 8, petiolo 3 lin. longo.

10. **D. venulosa**, Thw.; glabra; foliis oblongis vel ovato-oblongis, obtuse rostrato-acuminatis, superne creberrime et valde prominentim venulosis, ad costam canaliculatis, basi sub-5-plinerviis, rotundatis, petiolatis.—C.P. 3675.

HAB. South of the island.

Arbor magna, ramulis nigrescentibus. Folia 2–3 poll. longa, 1– $1\frac{1}{2}$  poll. lata, venis primariis utrinque circiter 7, petiolo 5–7 lin. longo. Inflorescentiam nondum vidi.

35. "Doona affinis, Thw.," adde: (C.P. 3712).

36. "SHOREA, Roxb." Adde:

3. **S. lissophylla**, Thw.; glabra; foliis ovatis vel ovato-rotundatis, abrupte breviter rostratis, obtusis, basi latis, rotundatis vel subcordatis, petiolo brevi; paniculæ ramis glabris; pedicellis brevissimis et cum alabastris minute incano-pubescentibus.—C.P. 3407.

HAB. Hinidoon Pattoo and Singhe-rajah Forest.

Arbor ingens, ramulis nigrescentibus. Folia 3–5 poll. longa,  $\frac{1}{2}$ –3 poll. lata, venis primariis utrinque circiter 7, his cum costa superne parum depressis, subtus prominentibus, venulis inconspicuis, petiolo semipollicari, ruguloso. Panicula multiflora. Antheræ barbatae.

Page  
36.

“HOPEA, Roxb.” Adde:

2. **H. jucunda**, Thw.; glabra; *foliis* ovatis vel ellipticis, longe rostrato-acuminatis, obtusis, basi rotundatis vel parce angustatis, petiolatis, *venis primariis* utrinque 4 et subtus ad axillas sæpius glanduliferis; *paniculis* axillaribus terminalibusque, folio æquilongis vel longioribus; *corolla* externe albido-pilosa.—c.p. 3709.

*Forma foliis minoribus*.—c.p. 3710.

HAB. Hinidoon Pattoo, Pasdoon Corle, and Saffragam.

*Arbor* magna, *ramulis* teretibus, nigrescentibus. *Folia* 1½–5 poll. longa, 1–2½ poll. lata, *costa* *venis primariisque* subtus prominentibus, *venulis* parum conspicuis, *petiolo* 2½–4 lin. longo. *Flores* iis *H. discoloris* simillimi.

36. “VATERIA, Linn.” Adde post descriptionem *V. Indicæ*:

Sect. 2.—*Stamina* 15, *biserialia* (*Stemonoporus*, Thw.).

2. **V. (Stemonoporus) Wightii**, Benth. in Benth. et Hook. f. Gen. Plant. p. 194.—*Stemonoporus Wightii*, Thw. in hac Enumeratione, p. 37.—c.p. 3415.

3. **V. (Stemonoporus) Gardneri**, Benth. l. c.—*Stemonoporus Gardneri*, Thw. in hac Enum. p. 38.—c.p. 1920.

4. **V. (Stemonoporus) acuminata**, Thw.; *foliis* lanceolatis vel oblongo-lanceolatis, reticulato-venulosis, gradatim (rarius subito) valde acuminatis, basi rotundatis, longiuscule petiolatis; *paniculis* folio brevioribus, racemiformibus vel parce ramosis; *bracteis* minutis, deciduis; *calycis* laciniis ovato-lanceolatis.—c.p. 3595, 3687 (3474).

HAB. Madamahanevewa, Ambagamowa, and Badulla districts.

*Arbor* magna, *ramulis* parce rugosis, rubro-nigrescentibus. *Folia* 4–7 poll. longa, ¾–2 poll. lata, *venis primariis* utrinque 10–16, *petiolo* ¾–1½ poll. longo. *Stipulæ* falcato-lanceolatæ, 3 lin. longæ, ½ lin. latæ, cito deciduæ.

5. **V. (Stemonoporus) affinis**, Benth. l. c.—*Stemonoporus affinis*, Thw. in hac Enum. p. 38.—c.p. 2430.

6. **V. (Stemonoporus) lanceolatus**, Benth. l. c.—*Stemonoporus lanceolatus*, Thw. in hac Enum. p. 38.—c.p. 2658.

7. **V. (Stemonoporus) canaliculata**, Benth. l. c.—*Stemonoporus canaliculatus*, Thw. in hac Enum. p. 38.—c.p. 3413.

8. **V. (Stemonoporus) petiolaris**, Benth. l. c.—*Stemonoporus petiolaris*, Thw. in hac Enum. p. 38.—c.p. 3151.

9. **V. (Stemonoporus) rigida**, Benth. l. c.—*Stemonoporus rigida*, Thw. in hac Enum. p. 38.—c.p. 2645.

10. **V. (Stemonoporus) oblongifolia**, Benth. l. c.—*Stemonoporus oblongifolius*, Thw. in hac Enum. p. 38.—c.p. 2646.

11. **V. (Stemonoporus) reticulata**, Benth. l. c.—*Stemonoporus reticulatus*, Thw. in hac Enum. p. 38.—c.p. 3414.

12. **V. (Stemonoporus) nitida**, Benth. l. c.—*Stemonoporus nitidus*, Thw. in hac Enum. p. 39.—c.p. 3483.

*Species dubiæ hujus Sectionis.*

13. **V. (Stemonoporus) Moonii**, Benth. l. c.—*Stemonoporus (?) Moonii*, Thw. in hac Enum. p. 39.—c.p. 1792.

Page

14. **V. (Stemonoporus) disticha**, Thw.; *ramulis* teretibus, fulvo-tomentellis; *foliis* coriaceis, oblongis, rostrato-acuminatis, ad costam canaliculatis, basi rotundatis, petiolatis; *paniculis* axillaribus terminalibusque, glabris, *junioribus* imbricato-bracteatis; *bracteis* majusculis, distichis, late ovatis, adpresse fulvo-pilosis.—c.p. 3707.

HAB. Hinidoon Pattoo and Pasdoon Corle.

*Arbor* ingens. *Folia* circiter 5 poll. longa,  $1\frac{1}{2}$ –2 poll. lata, delicatule parallelim venulosa, venis primariis utrinque circiter 12, *petiolo* 4 lin. longo.

15. **V. (Stemonoporus) scabriuscula**, Thw.; *ramulis* tomentoso-vel scabriusculo-lepidotis; *foliis* oblongo-lanceolatis, acuminatis, superne glabris, ad costam canaliculatis, subtus (cum petiolo) scabriusculo-lepidotis vel subglabris, venis primariis numerosis, versus marginem arcuato-anastomosantibus; *paniculis* axillaribus terminalibusque, tomentoso-lepidotis, *junioribus* bracteatis; *bracteis* alternis, ovato-acuminatis.—c.p. 3708 (3452).

HAB. Hinidoon Pattoo and Pasdoon Corle.

*Arbor* ingens. *Folia* 3–10 poll. longa,  $1$ – $2\frac{1}{2}$  poll. lata, *petiolo* 5–7 lin. longo. *Stipulae* 5 lin. longae, 1 lin. latae, falcatae, cito deciduae. *Bractae* 3 lin. longae, tomentoso-lepidotae.

Sect. 3.—*Stamina* 5, uniserialia (Monoporandra, Thw.).

16. **V. (Monoporandra) cordifolia**, Thw.—*Monoporandra*, hujus Enum. p. 39.—c.p. 2647.

17. (**V. Monoporandra**) **elegans**, Thw.—*Monoporandra*, hujus Enum. p. 39.—c.p. 371.

18. **V. (Monoporandra) lancifolia**, Thw.—*Monoporandra*, hujus Enum. p. 39.—c.p. 3412.

37. Pro “6. **ISAUXIS**, Arn.,” substitue:

## 6. **VATICA**, Linn.

Sect. *calycis fructiferi laciniae subaequalis* (Isauxis, Arn.).

1. **V. (Isauxis) Roxburghiana**, Benth. in Benth. et Hook. f. Gen. Plant. p. 192.—*Isauxis Roxburghiana*, Wight, Illust. p. 88; Icon. Pl. Ind. Or. t. 26.—c.p. 604.

HAB. South of the island. Nom. vulg. “Mandora-gass.”

2. **V. (Isauxis) affinis**, Thw.; *foliis* oblongo-lanceolatis, acuminatis, utrinque glabris et prominentim crebre reticulato-venulosis, venis primariis utrinque 5–7, basi parum rotundatis, petiolatis; *paniculis* axillaribus terminalibusque, *folio* brevioribus; *pedunculi ramis* nigrescentibus parce, *pedicellis alabastrisque* dense pallide fulvo-lepidotis.—*Isauxis Roxburghiana*, hujus Enum. p. 37 (haud Wight).—c.p. 3416.

HAB. Hinidoon Pattoo and Pasdoon Corle.

*Arbor* ingens, *ramulis* nigrescentibus. *Folia* semipedalia, *petiolo* pollicari, nigrescente. *Calycis laciniae* acutissimae.

Ab hac specie *V. (Is.) Roxburghiana* differt *ramulis*, *petiolis* paniculisque pallidis lepidotisque, *foliis* plures venas primarias (utrinque circiter 10) habentibus et subtus lepidotis.

3. ? **V. (Isauxis) cordifolia**, Thw.; glabra; *ramulis* teretibus; *foliis* nitidis, oblongo-vel ovato-lanceolatis, acuminatis, obscure et

Page

minute reticulato-venulosis, basi rotundato-cordatis, petiolatis; *venis primariis* utrinque circiter 7, ad axillas sæpe glanduliferis.—  
C.P. 3726.

HAB. Atakalan Corle.

*Arbor* ingens. *Folia* 5–7 poll. longa, 2–3½ poll. lata, *petiolo* pollicari. *In-*  
*florescentiam* nondum vidi.

40. Pro “1. **CLEYERA**, Thunb.,” substitue:

1. **TERNSTRÆMIA**, Linn. f.

*Cleyera*, Thunb.

Et adde:

1. **T. gymnanthera**, Benth. in Benth. et Hook. f. Gen. Plant. p. 182.—*Cleyera gymnanthera*, W. and A., hujus Enum. p. 40.—  
C.P. 778 (779).

2. **T. emarginata**, Benth. l. c.—*Cleyera emarginata*, Gardn.; hujus Enum. p. 40.—C.P. 782.

41. Pro “3. **SAROSANTHERA**, Korthals,” substitue:

3. **ADINANDRA**, Jack.

*Sarosanthera*, Korthals.

Et adde:

1. **A. lasiopetala**, Benth. l. c.—*Sarosanthera lasiopetala*, Thw. in hac Enum. p. 41.—C.P. 775 (776, 783).

43. Pro “7. **STEMONURUS**, Bl.,” substitue:

7. **LASIANTHERA**, P. de Beauv.

*Stemonurus*, Bl. (*ex parte*). *Urandra*, Thw.

Et adde:

1. **L. apicalis**, Benth. in Benth. et Hook. f. Gen. Plant. p. 350.—*Stemonurus apicalis*, Thw. in hac Enum. p. 43.—C.P. 2569.

44. Pro “8. **PLATEA**, Blume,” cum syn., substitue:

8. **GOMPHANDRA**, Wall.

*Platea*, Miers, Thw. (haud Blume). *Stemonurus*, Miers (partim).

Et adde:

1. **G. axillaris**, Wall.—*Platea axillaris*, Thw. in hac Enum. cum syn.—C.P. 251 (762).

2. **G. coriacea**, Wight.—*Platea coriacea*, Thw. in hac Enum. cum syn.—C.P. 375.

44. “**ATALANTIA**, Corr.” Adde:

2. **A. racemosa**, W. et A. Prod. i. p. 91; Oliver, in Linn. Soc. Journ. of Proc. v. Supp. ii. p. 24.—C.P. 3673.

HAB. Galagama, Gardner.

3. **A. Ceylanica**, Oliv. l. c. p. 25, cum syn.—*Sclerostylis Ceylanica*, Wight, hujus Enum. p. 46.—C.P. 1196.

Var.  $\beta$ . *rotundifolia*.—*Scl. rotundifolia*, Thw. in hac Enum. p. 46.—C.P. 3295.

4. **A. Missionis**, Oliv. l. c.—*Limonia Missionis*, Wight, hujus Enum. p. 45.—C.P. 1194.

Page  
45.

## “GLYCOSMIS, Corr.”

Pro “*G. arborea*, DC.,” et “*G. pentaphylla*, DC.,” substitue:  
*G. pentaphylla*, Corr. (Oliv. l. c. p. 37).—*G. arborea*, DC.—c.p.  
1192, 180. Et adde:

Subvar. *longifolia*, Oliv. l. c.; *foliis* sæpius unifoliolatis.—c.p.  
3822.

HAB. Singhe-rajah Forest.

45. “MURRAYA, Kœnig.” Adde:

3. *M. Glenieii*, Oliv. l. c. p. 29.—c.p. 3627.

HAB. Trincomalie, *Rev. S. O. Glenie*.

*Ovarium* 5- vel 4-loculare, *ovulis* solitariis vel geminatis, superimpositis.  
Cæteroquin *M. exotica*, var. *a*, valde similis et forsitan ejusdem varietas, *Oliv*.

3. *M. Kœnigii*, Spr. (Oliv. l. c. p. 29).—*Bergera Kœnigii*, Linn.,  
hujus Enum. p. 46.—c.p. 2547.

47. “CLAUSENA, Burm.”

“*C. Willdenovii*, W. et A.” Adde:

*Forma pubescens*.—*C. pubescens*, W. et A., hujus Enum. p. 47.  
—c.p. 2546.

Adde:

3. *C. Indica*, Oliv. l. c. p. 36, cum syn.—*Piptostylis Indica*,  
Kze. Misc. iii. p. 33, t. 2. *Bergera nitida*, Thw. in hac Enum.  
p. 46.—c.p. 2421, 1203.

47. “PARAMIGNYA, Wight.” Adde:

2. *P. armata*, Oliv. l. c. p. 43.—*Arthromischus armatus*, Thw. in  
hac Enum. p. 47.—c.p. 1197, 3115.

47. Dele “*Luvunga scandens*, Ham.,” cum descriptione.

48. “GARCINIA, Linn.” Adde:

4. *G. Zeylanica*, Roxb. Fl. Ind. ii. p. 621.

HAB. Ceylon, *Roxburgh*.

Species mihi ignota, nisi, ut suspicor, forma est *G. Cambogiæ*.

5. *G. (Discostigma) terpnophylla*, Thw.—*Terpnophyllum Zey-*  
*lanicum*, Thw. in hac Enum. *Discostigma Zeylanicum*, Planchon  
et Triana, in Ann. des Sc. Nat. ser. 4, xiv. p. 364.—c.p. 2695.

Var.  $\beta$ . *acuminata*.—*Discostigma acuminatum*, Planch. et Triana,  
l. c.—c.p. 3400.

50. “MESUA, Linn.”

“*M. ferrea*, Linn., var. *a*,” adde ut synonym.: *M. Walkeriana*,  
Planch. et Trian. l. c. xv. p. 301. “Var.  $\beta$ ,” adde ut synonym.: *M.*  
*salicina*, Planch. et Trian. l. c. p. 300.

50. Pro “2. *M. speciosa*, Chois.” cum tota descriptione, substitue  
duas species sequentes:

2. *M. Thwaitesii*, Planch. et Trian. l. c. p. 305.—*M. speciosa*,  
hujus Enum. var. *a* (haud Choisy).—c.p. 2675.

HAB. Damp forests, especially on the banks of streams, in the  
Ambagamowa, Saffragam, and Hinidoon districts.

Page

4. **M. sclerophylla**, Thw.; *foliis* valde coriaceis, subeveniis, lanceolatis vel oblongo-lanceolatis, acuminatis, subtus dense albo-pruinosis, basi rotundatis; *floribus* mediocribus, axillaribus terminalibusque, ad apicem pedunculi brevis (in meis specim.) solitariis; *calyce* glabro.—*M. speciosa*, var.  $\beta$ , hujus Enum.—c.p. 603.

HAB. Reigam Corle.

*Arbor* magna. *Folia* 3-4 poll. longa,  $\frac{3}{4}$ -2 poll. lata, *petiolo* 2-3 lin. longo.

50. Pro "3. **M. Coromandelina**, Wight, Icones, t. 117; Illust. i. 127.—c.p. 3404," substitue:

3. **M. pulchella**, Planch. et Trian. l. c. p. 307.—*M. Coromandelina* hujus Enum. (haud Wight).—c.p. 3404.

50. " **Kayea stylosa**, Thw." Adde ut synonyma: *K. cuspidata*, Planch. et Trian. l. c. p. 296.

51. " **CALOPHYLLUM**, Linn." Adde:

10. **C. cordato-oblongum**, Thw.; *ramulis* tetragonis, glabris (*junioribus* cum petiolis pedunculisque fugaci-ferrugineo-pilosis); *foliis* oblongis, obtusis, basi parum angustatis, cordatis, brevipetiolatis; *paniculis* terminalibus, folio subæquilongis; *pedicellis* alabastrisque subæquilongis pallide fulvo-tomentosis; *sepalis* 4, 2 interioribus majoribus et petaloideis; *petalis* 4.—c.p. 3823.

HAB. Hinidoon Pattoo. Floret Septembre.

*Arbor* ingens. *Folia* 4-6 poll. longa, 2-3 poll. lata, venis parallelis utrinque parum prominentibus, *petiolo* 2-3 lin. longo. *Petala* orbicularia, semipollicaria.

51. Pro "2. **C. decipiens**, Wight, Illust. i. 128; Icones, t. 106.—c.p. 3403," substitue:

2. **C. Thwaitesii**, Planch. et Trian. in Ann. des Sc. Nat. ser. 4, xv. p. 260.—*C. decipiens* hujus Enum. (haud Wight).—c.p. 3403.

*Antheræ* fere dimidio breviores quam in *C. decipiente*, Wight. *Fructus* sphæricus.

51. " **C. cuneifolium**, Thw." In floribus hujus speciei, cl. Planch. et Triana 4-8 petala invenerunt.

53. " **SALACIA**, Linn." Adde:

5. **S. terminalis**, Thw.; arborea, glabra; *ramulis* rugosulis; *foliis* ovatis vel ellipticis, acuminatis, versus apicem remote serrulatis, basi parum acutatis, petiolatis; *paniculis* ad apices ramulorum, axillaribus terminalibusque, plurifloris, folio longioribus.—c.p. 3737.

HAB. Hikkudoowe, near Galle.

*Arbor* 30-pedalis, ramosissima. *Folia* 3-4 $\frac{1}{2}$  poll. longa, 1 $\frac{1}{2}$ -3 poll. lata, superne sublævia, subtus reticulato-venosa, venis primariis utrinque circiter 6, *petiolo* semipollicari. *Pedicelli* alabastro duplo longiores. *Flores* 3-3 $\frac{1}{2}$  lin. in expan. *Calycis* segmenta brevia, rotundata vel subacuta. *Petala* orbicularia, unguiculata, reflexa. *Stamina* 3, filamentis cum ovario connatis. *Fructum* nondum vidi.

53. " **S. diandra**, Thw." Pro "*calycis segmentis* triangularibus," substitue: quam calycis segmenta triangularia.

Page  
55.**SCHMIDELIA**, Linn. Adde :

6. **S. varians**, Thw.—*S. allophylla*, var.  $\beta$ , hujus Enum. p. 55.—c.p. 562.

A *S. allophylla* differt *foliis* minoribus, *racemis* multo longioribus, *bracteis* minutissimis et brevissime pilosis.

56. Adde : 3 a. **GLENNIEA**, Hook. f.

(Benth. et Hook. Gen. Plant. p. 404.)

1. **G. Zeylanica**, Hook. f. l. c.

Var.  $\alpha$ ; *foliis* bifoliolatis.—*Sapindus unijugus*, Thw. in hac Enum. p. 56.—c.p. 2465.

Var.  $\beta$ ; *foliis* 2-4-foliolatis.—*Nephelium fuscatum*, Thw. in hac Enum. p. 58. *Sap. Glenieii*, Thw. MSS.—c.p. 3676.

HAB. Var.  $\alpha$ , Həwahetti District, at an elevation of 3000 to 4000 feet. Var.  $\beta$ . Trincomalie, Rev. S. O. Glenie.

56. “**Hemigyrosa canescens**, Thw.” Adde :

*Fructus* subsphæricus, sæpe abortu monospermus, tomentosus, ochraceus, pulposus; *epicarpio* pergamentaceo; *putamine* tenui, cartilagineo, interne adpresse piloso. *Semina* oblonga, testa tenui castanea. *Arillus* 0.

57. Adde : 6 a. **POMETIA**, Forst.

*Irina*, Blume. *Eccremanthus*, Thw. in Kew Journ.

1. **P. eximia**, Hook. f. in Benth. et Hook. Gen. Plant. p. 407.—*Nephelium eximium*, Thw. in hac Enum. p. 57, cum syn.—c.p. 1153.

6 b. **EUPHORIA**, Juss.

Hook. f. in Benth. et Hook. Gen. Plant. p. 406.

1. **E. Longana**, Bot. Mag. t. 4096.—*Nephelium Longanum*, Camb., hujus Enum. p. 58, cum syn.—c.p. 1152.

2. **E. Gardneri**, Hook. f. l. c.—*Nephelium Gardneri*, Thw. in hac Enum. p. 58.—c.p. 1154.

58. “**Nephelium fuscatum**, Thw.” est **Glenniea Zeylanica**, Hook. f., var.  $\beta$  (vide supra). *Petala* minuta, decidua.

58. Pro “8. **PTERIDOPHYLLUM**, Thw.” substitue :

8. **FILICIUM**, Thw., Benth. et Hook. f.

Et hinc (p. 59) pro “**P. decipiens**, Thw.” substitue : **F. decipiens**, Thw.

NOTA. Genus *Filicium*, fide cl. Hook. f., ad *Burseraceas* recte, ut nunc censeo, referendum est.

59. “**MELIOSMA**, Blume.”

Hoc genus, fide cl. Hook. f., ad **SABIACEAS** transferendum. Vide Benth. et Hook. f. Gen. Plant. p. 414.

59. “**MELIA**, Linn.” Adde :

2. **M. Azedarach**, Linn. (W. et A. Prod. i. p. 117, cum syn.)—HAB. Near Batticalva, Rev. S. O. Glenie.

- Page  
60. "AMOORA, Roxb." Adde:  
2. **A. Championii**, Hook. f. in Benth. et Hook. f. Gen. Plant. i. p. 335.—*Dysoxylon Championii*, Hook. f. et Thoms., hujus Enum. p. 61.—c.p. 1193.
60. "**Dysoxylon macrocarpon**, Blume, var.  $\beta$ ." Adde: c.p. 3596.
64. "OXALIS, Linn." Adde:  
2. **O. sensitiva**, Linn.—*Biophytum sensitivum*, DC., et *B. nervifolium*, hujus Enum. p. 64.—c.p. 1190, 2784.  
Var.  $\beta$ . *caule* elongato, ramoso.—*B. nudum*, Arn., et *B. proliferum*, var.  $\beta$ , hujus Enum. p. 64.—c.p. 6, 464, 2419.  
Var.  $\gamma$ . *prolifera*, *foliolis* minoribus.—*B. proliferum*, Arn., hujus Enum. p. 64.—c.p. 2418.  
HAB. Very abundant. Var.  $\beta$  and  $\gamma$  in the more elevated parts of the island.  
*Flores* dimorphi. *Stamina* utriusque formæ 10, alterna *filamentis* multo brevioribus: *forma a*, *stylosa*; *stylis* erectis elongatis, filamenta longa superantibus: *forma b*, *staminea*; *stylis* recurvis, filamenta breviora parum excedentibus.
66. "**Impatiens Walkeri**, Hook." Adde, post "6000 feet:" A variety with yellow flowers has been found at Happootelle, by Mr. W. H. Wright.
69. Adde: 1/1. **EVODIA**, Forst.  
1. **E. triphylla**, DC. Prod. i. p. 724.—*Xanthoxylon triphyllum*, Juss., hujus Enum. p. 69.—c.p. 538, 2501.
69. Pro "3. **CYMINOSMA**, Gærtn.," substitue:  
3. **ACRONYCHIA**, Forst.  
*Jambolifera*, Linn. *Cyminosma*, Gærtn. Et adde:  
1. **A. pedunculata**, Walp. Rep. i. p. 523.—*Cyminosma pedunculata*, DC., hujus Enum. p. 69.—c.p. 1249.
70. "**Ochna cordata**, Thw." Certe forma solum *O. squarrosæ*, Linn., et, ut species, delenda.
72. "**KURRIMIA**, Wall., Arnott." Adde, ut synonyma:  
*Pyrospermum*, Miq.
72. Pro "5. **CATHA**, Försk.," substitue:  
5. **GYMNOSPORIA**, W. et A., Benth. et Hook. f.  
*Catha*, Endl. (non Försk.). Et adde:  
1. **G. emarginata**, W. et A. Prod. i. p. 160.—*Catha emarginata*, G. Don, hujus Enum. p. 72.—c.p. 1235.  
2. **G. fruticosa**, Hook. f. in Benth. et Hook. f. Gen. Plant. p. 365.—*Catha fruticosa*, Thw. in hac Enum. p. 72.—c.p. 3386.



Page

74.

Pro "2. **BERCHEMIA**, Neck.," substitue :2. **SAGERETIA**, Brongn. Et adde :

1. **S. affinis**.—*S. costata*, Miq. Fl. Ind. Bat. i. p. 645. *Rhamnus affinis*, Blume, Bijd. p. 1140 (fid. Miq. l. c.) *Berchemia parviflora*, hujus Enum. p. 74 (haud Wall.).—c.p. 2477.

76. " **Semecarpus Gardneri**, Thw." Dele "2940."76. " **Semecarpus obscura**, Thw." Adde : c.p. 2940 ; et

"A *S. Gardneri*, Thw., foliis magis coriaceis et drupis erectis nec valde obliquis præcipue differt."

76. " **Semecarpus oblongifolia**, Thw." Adde :

Var.  $\beta$ . *nigro-viridis* ; floribus drupisque majoribus.—*S. nigro-viridis*, Thw. in hac Enum. p. 76.—c.p. 631, 2444 (641, 2586).

Var.  $\gamma$ . *acuminata* ; foliis drupisque acuminatis.—*S. acuminata*, Thw. in hac Enum. p. 76.—c.p. 2676.

78, 79. Genera " **PROTIUM**, Burm.," " **CANARIUM**, Linn.," et " **SCUTINANTHE**, Thw." (ut **CANARII** Sectio), fide cl. Benth. et Hook. f., ad **BURSERACEAS** transferenda.

79. Pro "1. **BLACKWELLIA**, Comm.," substitue :1. **HOMALIUM**, Jacq.Sect. *Blackwellia*. Et adde :

1. **H. Ceylanicum**, Benth. in Linn. Soc. Journ. of Proc. iv. p. 35.—*Blackwellia Ceylanica*, Gardn., hujus Enum. p. 79.—c.p. 388.

80. Adde : 3. **ELLIPANTHUS**, Hook. f.

Benth. et Hook. f. Gen. Plant. p. 434.

1. **E. unifoliolatus**, Hook. f. l. c.—*Connarus unifoliatus*, Thw. in hac Enum. p. 80.—c.p. 2443.

81. **CROTALARIA**, Linn. Adde :

23. **C. triquetra**, Dalz. in Hook. Journ. of Bot. ii. (1850), p. 34.—c.p. 3832.

HAB. Peradenia, rare.

81. " **C. evolvuloides**, W. et A." Adde : c.p. 3593.81. Dele "5. **C. bifaria**, Linn.," cum descriptione tota, et in ejus loco insere :

5. **C. globosa**, Wight. (W. et A. Prod. i. p. 190 ; Walp. Rep. i. p. 587.)—c.p. 3736.

HAB. Trincomalie, *Rev. S. O. Glenie*.82. " **C. striata**, DC." Pro "c.p. 3608," substitue : c.p. 3810, et adde :

Var.  $\beta$  ; foliis firmioribus, acutis ; racemis minus elongatis ; seminibus majoribus.—c.p. 3608.

HAB. Colombo, *Mr. W. Ferguson* ; var.  $\beta$ , common in the Kandian district.

*Legumina* pendula, brevi-stipitata, glabra, cylindrica, parum falcata clavatae, seminibus numerosis 30-40, lævia, nitida, pallide brunnea.

Page  
83.

## "INDIGOFERA, Linn." Adde:

13. *I. pentaphylla*, Linn. (W. et A. Prod. i. p. 200, cum syn.)—  
—C.P. 3524.

HAB. Near Colombo, *Mr. W. Ferguson*.

14. *I. tinctoria*, Linn. (W. et A. Prod. i. p. 202, cum syn.;  
Wight, Ic. t. 365.)—C.P. 3591.

HAB. Hotter parts of the island.

83. Pro "*I. cuneaphylla*," substitue: *I. enneaphylla*.

83. "*I. flaccida*, Kœnig." Adde:

Var.  $\beta$ . *constricta*; robustior, *foliis* majoribus, sæpius 9-foliolatis; *calycis* segmentis brevioribus, acutis, nec subulatis; *legumine* crassiore, suberecto, curvato, parum tetraquetra, marginibus parum incrassato, 4-6-spermo, inter semina constricto; *seminibus* majoribus. An *I. Kleinii*, W. et A. P.—C.P. 3811.

HAB. Matelle, East.

84. "TEPHROSIA, Pers." Adde:

7. *T. spinosa*, Pers. (W. et A. Prod. i. p. 214, cum syn.)

HAB. North of the island, *Mr. W. Ferguson*.

85. "*Uraria crinita*, DC." Adde:

In hortis culta, vix indigena.

86. "DESMODIUM, DC." Adde:

13. *D. jucundum*, Thw.; fruticosum, *ramulis junioribus*, *petiolis foliisque* subtus albido-sericeis; *foliis* pinnatim trifoliolatis; *foliolis* firmis, ellipticis vel obovatis, retusis, mucronulatis, *venis primariis* utrinque 6-14, subtus prominentibus; *racemis* terminalibus, plurifloris, sæpe valde elongatis; *pedicellis* geminis; *calycis* segmentis acuminatis, ciliatis; *legumine* lineari, complanato, parum constricto, rostrato, circiter 4-spermo, pilosulo, prominentim reticulato, articulis dimidio longioribus quam latis.—C.P. 3778.

HAB. Lakkaigalla and Ettangwella in the Matelle District, *A. Oswald Brodie, Esq.*

*Frutex* 3-4-pedalis, *ramis* rubro-nigrescentibus. *Folia* petiolo pollicari, *foliolis* 1-3½ poll. longis, ¾-1¾ poll. latis, *petiolulo* 1-2 lin. longo. *Stipulae* late lanceolatae, acuminatae, pilosulae, basi latae obliquaeque, 5 lin. longae. *Stipellae* subulatae, 1-2 lin. longae. *Peduncululi* tomentelli. *Pedicelli* 4 lin. longi, pilosi. *Petala* rubro-purpurea. *Legumen* ¾-1 poll. longum, 1½ lin. latum.

A *D. concinno*, DC., cui valde similis, hæc species differt *ramulis foliisque* magis sericeis et his firmioribus, latioribus retusisque.

14. *D. Gangeticum* DC. (W. et Arn. Prod. i. p. 225, cum syn.)  
—C.P. 3813.

HAB. Matelle, East.

87. "*D. polycarpon*, DC., var.  $\beta$ ." Adde, ut synonyma: *D. trichocaulon*, DC.

87. Adde: 21/1. HEDYSARUM, Jeaume.

1. *H. mucronatum*, Blume, Cat. Buit. p. 93; Walp. Rep. i. p. 746.—C.P. 3544.

Page

HAB. Rathoongodde, in the Maturatte District ; a common weed in the Botanic Garden.

## 21/2. ELEIOTIS, DC.

1. *E. sororia*, DC. Prod. ii. p. 348 ; W. et A. Prod. i. p. 231, cum syn.—C.P. 3765.

HAB. Near Colombo, *Mr. W. Ferguson*.

87.

## “ALYSICARPUS, Neck.” Adde :

5. *A. monilifer*, DC. (W. et A. Prod. i. p. 232, cum syn.)—C.P. 3613.

HAB. Near Colombo, *Mr. W. Ferguson*.

6. *A. longifolius*, W. et A. l. c. p. 233, cum syn.—C.P. 3814.

HAB. Near Colombo, *Mr. W. Ferguson*.

88.

## Adde : 29/1. DIOCLEA, H. B. K.

1. *D. Fergusonii*, Thw. ; alte scandens, molliter fulvo-pubes-  
cens ; *foliolis* ovatis vel ovato-lanceolatis, obtusiusculis, mucronatis,  
basi rotundatis, lateralibus parum obliquis ; *stipulis* auriculæformi-  
bus, medio affixis glandulosisque, superne lanceolatis, acutis ; *pani-  
culis* racemiformibus, multifloris, elongatis, ramis brevibus 2-3-  
floris apice incurvatis glandulosis ; *bracteis* longiusculis, anguste  
lanceolatis, sulcatis, recurvis ; *legumine* majusculo, 1-3-spermo, su-  
tura superiore anguste bialata ; *seminibus* orbicularibus, complana-  
tis, hilo lineari et valde elongato.—C.P. 3817.

HAB. Near Colombo, *Mr. W. Ferguson*.

*Folia* (cum *petiolo* 2-3 poll. longo) 9-12 poll. longa. *Foliola* 4-6 poll.  
longa, 2½-5 poll. lata, *petiolulo* circiter 2 lin. longo. *Stipulæ* 3-4 lin. longæ.  
*Flores* pollicares. *Calyx* *lacinia* superiore lata, ad apicem bifida, *lateralibus*  
acutis, falcatis, *inferiore* parum longiore et apice retusa. *Bracteolæ* 2, ad basin  
calycis, parvæ, rotundatæ. *Petala* pallida, violacea. *Ovarium* pilosum, 2-3-  
ovulatum. *Stigma* parvum, parum dilatatum. *Discus* brevis, tubulosus, mar-  
gine 10-crenato. *Legumen* oblongum, subfalcatum, 5-6 poll. longum, 2 poll.  
latum, sparse pilosum. *Semina* 11-14 lin. in diam., castanea, hilo nigro.

89.

## “PHASEOLUS, Linn.” Adde :

7. *P. calcaratus*, Roxb. Fl. Ind. iii. p. 289 ; W. et A. Prod. i. p. 245.—C.P. 1473.

HAB. Very abundant in the Central Province.

89.

“*P. radiatus*, Linn.” Pro “C.P. 1473,” substitue : C.P. 3622, et  
pro :

“HAB. Very abundant up to an elevation of 2000 feet,” sub-  
stitue :

HAB. Trincomalie, *Rev. S. O. Glenie*.

90.

“*Dunbaria Heynei*, W. et A.” Ante “*D. oblonga*,” insere : ?

91.

## “RHYNCHOSIA, Lour.” Adde :

4. *R. villosula*, Thw. ; volubilis, glanduloso-hirsutula ; *foliis*  
pinnatim 3-foliolatis ; *foliolis* late trapezoideis, acuminatis, basi  
angustis, truncatis vel subcordatis, lateralibus externe dilatatis ; *sti-*

Page

*pulis* parvis, coloratis, trapezoideis, acuminatis; *racemis* axillaribus, plurifloris, folio longioribus; *bracteis* quam pedicelli breves parum longioribus, rotundatis, acutis, basi angustatis, cito deciduis; *calycis* lobo supremo latiore, infra medium bifido; *corolla* quam calyx duplo longiore; *carina* falcata, erostellata; *stylo* versus apicem incrassato, albido, nitido.—C.P. 3790.

HAB. Trincomalie, *Rev. S. O. Glenie*.

*Foliola* 1-2 poll. longa. *Pedicelli* solitarii. *Flores* semipollicares. *Calyx* ad medium fissus, *lobis lateralibus* et *infimo* paullo longiore lanceolatis, acutis. *Corolla* ochracea, *vexillo* piloso, rubro-lineato, ad basin laminæ biauriculato, *alis* falcato-auriculatis, margine inferior dente parvo munito. *Ovarium* biovulatum, flavo-villosum. *Legumen* (immaturum) oblongum, acutum.

5. **R. acutissima**, Thw.; volubilis, breve fulvo-tomentosa, *foliis* pinnatim trifoliolatis; *foliolo terminali* trapezoideo-lanceolato, *lateralibus* parum minoribus, externe dilatatis, ovato-lanceolatis et, cum terminali, acutissime acuminatis; *stipulis* lanceolatis, deciduis; *racemis* folio subæquilongis; *pedicellis* geminis, calyce longioribus; *bracteis* lanceolatis, acuminatis, ciliatis, calyce subæquilongis, cito deciduis; *calycis* lobo supremo latiore, ad apicem bifido laciniis divaricatis, *lateralibus*, cum *infimo* dimidio longiore; lanceolatis, acutis; *corolla* quam calyx subduplo longiore; *carina* falcata, erostellata.—C.P. 3445.

HAB. Hantani, in the Central Province, at an elevation of 3000 feet.

*Foliola* 2-5 poll. longa. *Pedicelli* 4 lin. longi. *Flores* semipollicares. *Corolla* ochracea, rubro-lineata; *vexillo* biauriculato, versus apicem piloso, retuso, *alis* falcato-auriculatis, margine inferiore dente parvo acuto munito. *Ovarium* 2-ovulatum, longe albido-pilosum.

Hæc, et species præcedens, *Dunbariæ* valde proximant, sed bracteas magnas illius generis egent.

92.

## “DERRIS, Lour.”

Pro “2. **D. ovalifolia**, Benth., var.,” cum decriptione tota, substitute:

2. **D. oblonga**, Benth. in Linn. Soc. Journ. of Proc. iv. supp. p. 113, cum syn.—C.P. 1493.

HAB. Central and Southern Provinces.

*Adde species sequentes:*

4. **D. scandens**, Benth. l. c. p. 103, cum syn.—*Brachypterum scandens*, W. et A., hujus Enum. p. 93.—C.P. 1492.

5. **D. parviflora**, Benth. l. c. p. 105.—*Brachypterum elegans*, Thw. in hac Enum. p. 93 (non *Derris elegans*, Benth.).—C.P. 2508.

6. **D. Benthamii**, Thw.—*D. paniculata*, Benth. l. c. *Brachypterum Benthamii*, Thw. in hac Enum. p. 93.—C.P. 2925.

94.

“**Dalbergia monosperma**, Dalzell.” Adde:

HAB. Reigam Corle.

94.

Adde: 52/1. **PERICOPSIS**, Thw., nov. genus.

*Calyx* campanulatus, in stipitem brevem attenuatus, demum prope basin circumscisse caducus, 4-fidus, *lobis* acutis subæqualibus

## Page

æquilongis, *supremo* breviter bifido reflexo, *lateralibus* parum falcatibus, infimo erecto concavo. *Petala* unguiculata, subæquilonga, *vexillo* orbiculari, reflexo; *alis* oblongis, auriculatis, margine inferiore dente minuto munito, *carinæ* petalis dorso versus apicem parum cohærentibus, cum *alis* conformibus. *Stamina* 10, *filamentis* liberis glabris, 5 paullo brevioribus. *Discus* brevis, vaginans, truncatus. *Ovarium* stipitatum, circiter 6-ovulatum. *Stylus* gradatim attenuatus, apice incurvus. *Stigma* parvum. *Legumen* stipitatum, compressum, coriaceum, indehiscens, utraque sutura anguste marginatum, apiculatum, 1-6-spermum, inter semina parum spongiosum, sæpe abortu seminum sinuatum, demum in articulis monospermis solutum. *Semina* oblique ovalia, compressa, hilo parvo.—Arbor *ingens*; foliis *pinnatis*; foliolis *alternis*, *majusculis*; paniculis *terminalibus*.

1. **P. Mooniana**, Thw.—*Dalbergia Mooniana*, Thw. in hac Enum. p. 93, cum syn.—C.P. 439.

HAB. South of the island, at no great elevation.

*Flores* pollicares. *Calyx* 7 lin. longus. *Petala* atro-purpurea, basi flavo-viridia. *Bracteæ* cito deciduæ, et eas nondum vidi.

94. “**Sophora violacea**, Thw.” Adde ad finem descriptionis post “*rubra* :” denique *brunnea*.

95. “**CÆSALPINIA**, Linn.”

Dele “3. **C. mimosoides**, Lam.,” cum descriptione, et in ejus loco insere :

3. **C. Glenieii**, Thw.; inermis, ferrugineo-tomentella; *foliis* abrupte bipinnatis, *pinnis* circiter 10-jugis, *pinnulis* parvis circiter 15-jugis, sessilibus, oblongis, utrinque pilosulis, apice rotundatis, subretusis, obscure mucronulatis, basi obliquis, latis, parum rotundatis; *racemis* plurifloris, *pedicellis* quam alabastra brevioribus; *floribus* majusculis; *sepalis* *petalisque* externe plus minus ferrugineo-pilosis; *filamentis* ad basin ferrugineo-villosis, superne glaberrimis; *ovario*, cum ejus stipite, dense ferrugineo-tomentoso; *stylo* basi pilosulo; *stigmatibus* majusculo, subreniformi.—C.P. 3815.

HAB. Trincomalie, *Rev. S. O. Glenie*.

*Pinnæ* 2-2½ poll. longæ. *Pinnulæ* 3 lin. longæ, 1½ lin. latæ, siccitate nigrescentes. *Flores* 1½ poll. in exp., aureo-flavi.

95. Adde: 55/1. **MEZONEURUM**, Desf.

1. **M. enneaphyllum**, W. et A. Walp. Ann. iv. p. 591, cum syn.—*Cæsalpinia enneaphylla*, Roxb. Fl. Ind. ii. p. 363. *Cæsalpinia mimosoides*, hujus Enum. p. 95 (non Lam.).—C.P. 3601 (2784 partim).

HAB. Not uncommon in the hotter parts of the island.

96. “**Cassia mimosoides**, Linn., var.  $\gamma$ ,” pro “*patello*” insere: paullo.

98. Adde: 63/1. **DETARIUM**, Juss., DC.

1. **D. Zeylanicum**, Thw.; glabrum; *foliis* pinnatis, *foliolis* 2-6 (sæpissime 6), alternis vel suboppositis, ovatis vel oblongis, acuminatis, obtusis, basi parum obliquis, rotundatis, subtus pallidioribus,

Page

opacis, minutissime reticulatis; *racemis* terminalibus, plurifloris, sparse pilosulis, folio subæquilongis vel brevioribus.—C.P. 3714.

HAB. Galpaatta, near Cultura.

*Arbor* ingens. *Foliola* 2–6 poll. longa, petiolulo 2 lin. longo. *Pedicelli* 2 lin. longi, medio minute bibracteolati. *Flores* conferti. *Sepala* 2 lin. longa, rotundata, persistentia. *Stamina* 10, filamentis alternis brevioribus, antheris rotundis. *Ovarium* stipitatum, pallide fusco-tomentosum, 3–4-ovulatum. *Stylus* filiformis, glaber. *Stigma* parvum. *Legumen* (immaturum) oblongum, parum falcatum, complanatum.

99.

“ACACIA, Willd.” Adde:

8. **A. Arabica**, Willd. (W. et A. Prod. i. p. 277, cum syn.)—C.P. 3640.

HAB. Trincomalie, *Rev. S. O. Glenie*.

103.

“COMBRETUM, Löffl.” Adde:

2. **C. sarcopterum**, Thw.; scandens?; *ramulis* teretibus; *junioribus*, *petiolis pedunculisque* brevissime ferrugineo-hispidulis et, cum *foliis subtus*, creberrime glanduloso-lepidotis; *foliis* integris, oblongo-lanceolatis, obtusiusculis, basi rotundatis, superne lævibus, subtus prominentim venosis; *petiolo* brevi; *spicis* axillaribus, simplicibus, folio brevioribus; *fructu* latiuscule 4-alato, alis oblongis, utrinque angustatis, carnosus, versus marginem parum incrassatis.—C.P. 3715.

HAB. Reigam Corle.

*Folia* 3–7 poll. longa,  $1\frac{1}{4}$ – $2\frac{1}{2}$  poll. lata, *venis primariis* utrinque circiter 15, *secundariis* parallelis parum distantibus; *petiolo* 3 lin. longo. *Fructus* 2 poll. longus,  $\frac{3}{4}$  poll. latus, glanduloso-lepidotus. *Flores* nondum vidi.

110.

“MEMECYLON, Linn.” Adde:

8/1. **M. rubro-cæruleum**, Thw.; arboreum, glabrum; *foliis* mediocribus, eveniis, cuneato-ovatis, versus apicem parum angustatis, retusis, subtus pallidioribus, brevipetiolatis; *pedunculis* solitariis vel binis, umbellam compositam, glomeratam, æquilongam gerentibus; *pedicellis* calyce dimidio longioribus; *calyce* aperte campanulato, obscure 4-dentato.—C.P. 3809.

HAB. Matelle, East.

*Arbor* 30-pedalis. *Folia*  $1\frac{1}{2}$ –2 poll. longa,  $\frac{1}{2}$ –1 poll. lata, juniora rufescentia. *Pedunculi* 3 lin. longi. *Calyx* rufescens. *Petala* cærulea, margine pallida.

13/1. **M. procerum**, Thw.; arboreum, glabrum; *foliis* majusculis, ovato-lanceolatis, gradatim acuminatis, ad costam canaliculatis, venis primariis vix conspicuis, subtus pallidioribus, basi rotundatis vel parum angustatis; *petiolo* superne sulcato; *pedunculis pedicellisque* nullis vel subnullis; *baccis* sphaericis, calyce (hoc genere) majusculo coronatis.—C.P. 3347.

HAB. Pasdoon Corle.

*Arbor* 30–40-pedalis. *Folia* 6–10 poll. longa,  $1\frac{3}{4}$ –5 poll. lata, *petiolo* 3–5 lin. longo. *Calyx* 2 lin. in exp.

112.

“**M. jambosoides**, Wight.” Dele: “3347.”

Page

113. "**M. parvifolium**, Thw." Adde :  
 Forma *foliis* majoribus (1-2 poll. longis) abrupte brevi-acuminatis, obtusis.—*M. Thomsoni*, Thw. MSS.—c.p. 758.  
 HAB. Rambodde, at an elevation of 4000 to 5000 feet.
114. "**EUGENIA**, Linn." Adde :
- 2/1. **E. xanthocarpa**, Thw.; *ramulis junioribus gemmisque* ferrugineo-pilosis; *foliis* elliptico-lanceolatis, obtuse acuminatis, glabris, basi acutis, superne ad costam canaliculatis, ad venas primarias prominentibus; *pedicellis* axillaribus, fasciculatis, ferrugineo-pilosis, petiolo subæquilongis; *baccis* 1-4-spermis, subsphæricis, ferrugineo-pilosis, minute glanduloso-verrucosis.—c.p. 3834 (2623 partim).  
 HAB. Near the coast, between Galle and Colombo.  
*Arbor* humilis. *Folia* 2½-4 poll. longa, 1-1½ poll. lata, *petiolo* 3 lin. longo. *Baccæ* 3-6 lin. in diam. ochraceæ.  
 Ab *Eug. terpnophylla*, Thw., differt *venis primariis* numerosioribus, his subtus minus prominentibus et propius marginem anastomosantibus, et *baccis* ochraceis, sæpius 2-4-spermis.
- 2/2. **E. rufo-fulva**, Thw.; *ramulis junioribus gemmisque* breve rufo-vel fulvo-pilosis; *foliis* lanceolatis, longe acuminatis, acutis, superne glabris, nitidis, ad costam canaliculatis, subtus brevi rufo vel fulvo tomento adpresse tectis, denique subglabris; *pedicellis* axillaribus, fasciculatis, petiolo subæquilongis.—*E. terpnophylla*, var.  $\beta$ , Thw. in hac Enum. p. 114.—c.p. 3835.  
 HAB. Gattehatte, near Avishavelle.  
*Arbor* 30-40-pedalis. *Folia* (cum acumine) 4-5 poll. longa, 1-1½ poll. lata, *petiolo* semipollicari.  
 Differt ab *Eug. terpnophylla*, *foliis* subtus pilosis, et *venis primariis* propius marginem anastomosantibus.
- 9/1. **E. insignis**, Thw.; arborea; *partibus junioribus* fugaci-albo-villosis; *foliis* majusculis, glabris, oblongis, acuminatis, superne lævibus, ad costam canaliculatis, subtus pallidioribus, *venis primariis* utrinque circiter 12, prominentibus, procul a margine anastomosantibus, areolis lævibus subeveniis; *floribus* axillaribus rameisque, paucis, sessilibus, parvis; *calyce* externe albo-villoso, segmentis oblongis; *baccis* sphæricis albo-villosis vel floccosis.—c.p. 3677.  
 HAB. Galpara, in the Reigam Corle.  
*Arbor* humilis, ramosa. *Folia* 6-8 poll. longa, 2-3½ poll. lata, basi parum acuta, *petiolo* 6-8 lin. longo. *Calycis* segmenta 2 lin. longa. *Petala* calyce parum longiora, ciliata.
11. **E. concinna**, Thw.; arborea; *partibus junioribus* pilosis; *ramulis* teretibus; *foliis* firmis, parcissime pilosis, denique glabris, ovatis, subabrupte acuminatis, obtusis, basi parum acutatis, *venis primariis* utrinque 12-16 prope marginem anastomosantibus, areolis obscure venoso-reticulatis; *petiolo* brevi; *pedicellis* axillaribus, fasciculatis, petiolo duplo longioribus; *floribus* parvis; *baccis* sphæricis, glabris, nitidis.—c.p. 2802.  
 HAB. Ambagamowa District.  
*Folia* 3-5 poll. longa, 1½-2¼ poll. lata, *petiolo* 2-3 lin. longo. *Bacca* coccineæ, 6 lin. in diam.

- Page  
114. “**Eugenia terpnophylla**, Thw.” Adde ad descriptionem:  
*Folia* membranacea, obscure reticulatim venulosa, venis primariis utrinque 5–8, subtus prominentibus, procul a margine anastomosantibus. *Baccæ* sphaericæ, sæpius monospermæ.  
Dele “var.  $\beta$ ,” et sequent. ad “rufo-pilosis.”
116. Pro “1. **Syzygium caryophyllifolium**, DC.,” cum descriptione, substitue:  
1. **S. Jambolanum**, DC. (W. et A. Prod. i. p. 329, cum syn.) —C.P. 3644.  
Var.  $\beta$ . *microcarpum*.—*S. caryophyllifolium*, DC.—C.P. 1584.  
HAB. Not uncommon in the warmer parts of the island.
116. Pro “2. **Syzygium polyanthum**,” cum sequent. ad “p. 16?,” substitue:  
2. **S. nervosum**, DC. Prod. iii. p. 260; Wall. Cat. no. 3551 B.  
Et adde:  
Hæc species differt ab *Eugenia polyantha*, Wight, et *E. balsamea*, Wall., paniculis majoribus et pedicellis subnullis.
116. Pro “4. **Syzygium montanum**,” cum tota descriptione, substitue:  
4. **S. firmum**, Thw.; arborea, glabra; *foliis* majusculis, coriaceis, nitidis, obovatis, abrupte et brevissime acuminatis, obtusis vel retusis, utrinque prominentim subparallele plurivenosis, basi angustatis, petiolatis; *paniculis* ad apices ramulorum axillaribus terminalibusque, folio brevioribus vel subæquilongis; floribus (pro hoc genere) majusculis; *calyce* in brevem pedicellum attenuato, segmentis brevibus, rotundatis; *petalis* distinctis.—*Eugenia firma*, Wall. Cat. 3603 A.—C.P. 2694.  
HAB. Ambagamowa District.  
*Arbor* magna. *Folia* 4–8 poll. longa, 3–3½ poll. lata, *petiolo* ¾–1 poll. longo. *Calyx* (cum pedicello) ½ poll. longus, 4 lin. latus.  
Differt ab *Eugenia montana*, Wight, foliis floribusque multo majoribus.
122. Adde: 4 a. **NESÆA**, Comm.  
1. **N. triflora**, H. B. K. (W. et A. Prod. i. p. 307, cum syn.) —C.P. 1541.  
HAB. Ceylon, *Walker* (*Herb. Hook.*).
124. “**Melothria deltoidea**, Thw.” In descriptione, pro “digitata” lege: sagittata.
126. “**LUFFA**, Tourn.” Adde:  
3. **L. amara**, Roxb. (W. et A. Prod. i. p. 343, cum syn.; Naudin in Ann. des Sc. Nat. ser. 4, xii. p. 123.)—C.P. 1623.  
HAB. North of the island, *Gardner*; Trincomalie, *Rev. S. O. Glenie*.  
*Semina* multo minora quam in *L. acutangula*.
128. “**Begonia subpeltata**, Wight.” Adde:  
*Folia* viridia vel aliquando nigro-cuprea.
129. “**Kalanchoe laciniata**, DC.” Adde: C.P. 3836.  
HAB. Matelle, East.



Page

130. “**Vahlia oldenlandioides**, Roxb.” Post “c.p. 1686,” adde: 1687.

130. Dele “2. **Vahlia viscosa**, Roxb.,” cum descriptione.

132. “**Hedera Vahlia**, Thw.” In descriptione, pro “balsamiflora,” lege: balsamiflua.

133. “**LORANTHUS**, Linn.” Adde:

12 a. **L. lonchiphyllus**, Thw.; *ramulis* teretibus; *foliis* coriaceis, glabris, penniveniis, ovato-lanceolatis, gradatim acuminatis, costa venis primariisque conspicuis, basi rotundatis; *petiolo* brevi, parce alato; *pedunculis* axillaribus, 1- vel 2-nis, brevibus, flores circiter 7, subeymosos, brevipedicellatos gerentibus; *calyce* cylindrico, truncato, unibracteolato; *corolla* longiuscula, parum curvata, medio dilatata, irregulariter fissa; *baccis* ovoideis.—C.P. 3678.

HAB. Ambagamowa District.

*Folia* 4-6 poll. longa,  $1\frac{1}{2}$ - $2\frac{3}{4}$  poll. lata, *petiolo* circiter 3 lin. longo. *Inflorescentia* plus minus ferrugineo-leprosa. *Pedunculi* 2 lin. longi. *Pedicelli*  $\frac{1}{2}$  lin. longi. *Calyx* 1 lin. longus. *Bracteola* subacuta, apice parum fimbriata vel denticulata, calyce dimidio brevior. *Corolla* sesquipollicaris. *Baccæ*  $\frac{1}{2}$  poll. longæ.

134. “**Loranthus Neilgherrensis**, W. et A.,” pro “1648,” substitue: 1646.

136. “**VISCUM**, Tourn.” Adde:

6. **V. floccosum**, Thw.; ramosissimum; *partibus junioribus*, *inflorescentia foliisque* subtus pilis ramosis ochraceo-floccosis; *ramulis* teretibus, striatis; *foliis* trinerviis, late ovatis vel rotundatis, parum acuminatis, obtusis, basi rotundatis, in petiolum brevem angustatis; *floribus* paucis, terminalibus, monoicis, verticillato-spicatis. FLOR. MASC.: *antheris* subsessilibus, erectis, 8-locellatis, poris dehiscens.—*Notothixos* (?) *floccosus*, Oliv. in Linn. Soc. Journ. of Proc. vii. p. 104.—C.P. 3654.

HAB. Hinidoon Pattoo, abundant; Kokool Corle.

*Folia*  $\frac{1}{2}$ - $1\frac{1}{4}$  poll. longa, *petiolo* 2-3 lin. longo. *Spicæ* breves. *Flores* minuti,  $\frac{1}{2}$  lin. in exp. interne glabri, perianthii lobis 4 vel rarius 3, triangularibus, acutis. FLOR. FÆM.: *Stylus* conicus a basi lata subplana quadrangulari, brevis, apice parum dilatatus, stigmatosus. *Baccæ* ovoideæ, albæ, 2 lin. longæ, lobis perianthii coronatæ.

136. “**Viscum spathulifolium**, Thw.” Adde, ut synonyma: *Ginhalloa* (?) *spathulifolia*, Oliv. l. c. p. 103.

138. “**Uncaria Gambier**.” Pro “Hunt.,” substitue: Roxb.

139. “8. **AXANTHES**, Blume.” Substitue:

8. **UROPHYLLUM**, Jack., Wall.

*Axanthes*, Blume. Et adde:

1. **U. Zeylanicum**.—*Axanthes Zeylanica*, Wight, hujus Enum. p. 139.—C.P. 499 (707).

Page

2. **U. ellipticum.**—*Axanthes elliptica*, Wight, hujus Enum. p. 139.—C.P. 359 (584).

*Flores* hujus et speciei præcedentis albidæ, *baccæ* aurantiacæ.

139. “**OPHIORRHIZA**, Linn.” Adde :

8. **O. decumbens**, Gardn. MSS.; suffruticosa; *caule* scabrello, basi decumbente radicante; *foliis* lanceolatis vel ovato-lanceolatis, utrinque acuminatis; *stipulis* brevibus, setaceo-cuspidatis, interne glandulosis; *corymbis* terminalibus, pedunculatis, scabrellis; *bracteolis* parvis, angustis; *pedicellis* calyce brevioribus; *corolla* quam calyx 3-4-plo longiore; *capsula* late triangulari, angulis subacutis.—C.P. 3656.

HAB. Common in the Central Province.

Differt ab *A. nemorosa*, Thw., cui valde similis, pedicellis floribusque brevioribus, et capsula latius et acutius ad latera producta; ab *O. Mungos*, Linn., caule suffruticoso et floribus longioribus satis distincta.

140. “**HEDYOTIS**, Lam.” Adde :

18 a. **H. cinereo-viridis**, Thw.; suffruticosa, subglabra; *ramulis* parum tetragonis; *foliis* lanceolatis, utrinque acuminatis, petiolatis, venis primariis subtus parum prominentibus; *stipulis* triangularibus, bicarinatis, parum connatis, omnino vel ad carinas pilosis, ad apicem plus minus rostrato-productis glandulosisque; *floribus* glomeratis, brevipedicellatis, axillaribus vel ramulos breves terminantibus; *calyce* campanulato, 4-lobato; *lobis* acutis, brevibus vel longiusculis; *corollæ tubo* quam calyx longiore, interne piloso; *lobis* longiusculis, linearibus, subacutis, glabris.—C.P. 95 (98).

Var.  $\beta$ . *foliis* majoribus; *stipulis* longius rostratis; *floribus* omnibus axillaribus; *corolla* externe pilosa.—C.P. 2639.

HAB. Matelle, East. Var.  $\beta$ . Ambagamowa.

Hæc species olim cum *H. inamœna*, Thw., confusa.

140. “**Hedyotis evenia**, Thw.” Dele annotationem, et insere: Hæc species ab *H. Lawsoniæ*, W. et A., inflorescentia terminali satis distinguitur.

141. “**Hedyotis flavescens**, Thw.” Certe forma solum *H. Lesser-tianæ*, Arn. (var.  $\gamma$ ), in locis apricis madidis crescens.

143. “**Hedyotis inamœna**, Thw.” Dele “95,” “(98),” et a “Var.  $\beta$ ” ad “Districts” inclus. Adde ad descriptionem :

*Folia* subevenia, glaberrima. *Stipulæ* margine dilatato, membranaceo, tuberculis notato. *Calyx* pilosus, tubo corollæ parum brevior, lobis ovatis obtusis.

HAB. Near Balangodde.

144. “**Hedyotis Burmanniana**, Br.” Pro “2368,” insere: 2638. Adde: species valde variabilis.

Forma *b*; *brachypoda*; *floribus* 1-2-nis, fere sessilibus, ad axillas foliorum.—*Oldenlandia brachypoda*, DC. Prod. iv. p. 424.—C.P. 3838.

HAB. Central Province.

Forma *c*; *racemosa*; *floribus* 3-6, in racemum dispositis ad api-

Page

cem pedunculi.—*Hedyotis* (*O.*) *brachiata*, Wight ; W. et A. Prod. i. p. 415.—C.P. 3837.

HAB. Near Colombo, *Mr. W. Ferguson.*

144. "DENTELLA, Forst."

Hoc genus placentam in medio dissepimenti affixam habet, et hinc inter genera subsectionis præcedentis, post *Urophyllum*, Jack, rectius allocandum.

145. "Morinda exserta, Roxb." Adde :

*Flores* dimorphi, alii stigmati, alii antheris exsertis.

145. Pro "13. MEPHITIDIA, Reinw.

*Lasianthus*, Jack, Wight," substitue :

13. LASIANTHUS, Jack, Wight.

*Mephitidia*, Reinw. Et adde :

1. **L. Moonii**, Wight, in Calc. Journ. of Nat. Hist. vi. p. 504.—*Mephit. Moonii*, hujus Enum. p. 145.—C.P. 92.

2. **L. strigosus**, Wight, l. c. p. 512.—*Mephit. Gardneri* et *M. tomentosa* (haud *M. strigosa*), hujus Enum. pp. 145, 146.—C.P. 348, 2711 (1723).

*Calycis* lobi lanceolati, pilosi.

3. **L. rhinophyllus**, Thw.—*Mephit. rhinophylla*, Thw. in hac Enum. p. 145.—C.P. 361.

4. **L. protractus**, Thw.—*Mephit. protracta* et *M. Walkeriana*, hujus Enum. p. 146 (haud *L. Walkerianus*, Wight).—C.P. 354, 289 (340, 1728).

*Calycis* lobi triangulares.

5. **L. Walkerianus**, Wight, l. c. p. 513.—*Mephit. strigosa*, Thw. in hac Enum. p. 146 (haud *L. strigosus*, Wight).—C.P. 1725.

*Calycis* pars libera tubulosa, minute dentata.

6. **L. oliganthus**, Thw.—*Mephit. oligantha* (excl. var.  $\beta$ ), hujus Enum. p. 146.—C.P. 357 (1726, 1727).

A specie præcedenti distinguitur *petiolis* brevioribus, et *floribus* multo minoribus.

7. **L. obliquus**, Thw.—*Mephit. oligantha*, var.  $\beta$ , hujus Enum. p. 146.—C.P. 3418.

8. **L. varians**, Thw.—*Mephit. varians*, hujus Enum. p. 146.—C.P. 57 (80, 290, 422, 423).

147. "ALLÆOPHANIA, Thw." Adde :

*Bacca* 4-coeca.

149. Dele "17. PRISTIDIA, Thw. (gen. nov.)" cum descriptione, genus a GÆRTNERA, Lam., haud distinctum, et hinc "**P. divaricata**, Thw." sub nomine **Gærtnera divaricata**, Thw., posthac agnoscenda.

- Page  
 150. “**Chasalia curviflora**, Thw.” Pro “*Flores* polygamo-dioici, fertiles breves, stigmatate exserto,” substitue:  
*Flores* dimorphi, alii *stigmatate*, alii *antheris* exsertis. *Corolla* longiuscula, curvata, vel brevior recta.
151. Genera “**RUBIA**, Tourn.,” et “**GALIUM**, Linn.,” subsectionem Ordinis ovulis medio dissepimenti affixis, melius formabunt.
153. “**TIMONIUS**, Rumph., DC.” Adde, ut synonyma:  
*Bobea*, Miquel.
153. “**Timonius Jambosella**, Thw.” In descriptione, pro “*comosis*” lege: *carnosis*.
154. “**Coffea ? elliptica**, Thw.,” ad genus **STYLOCORYNE**, Cav. transferenda; vide infra.
154. “**Prismatomeris albidiflora**, Thw.” Adde:  
*Coffeae tetrandrae*, Roxb., valde affinis; an ejusdem varietas floribus pentameris?
155. “**Ixora acuminata**, Roxb.” Adde: *I. jucunda*, Bot. Mag. t. 5197 (haud Thw.).
157. “**Hyptianthera macrocarpa**, Thw.” In descriptione, pro “*lobi 5*,” insere: *lobi 4*, rarius *5*.
158. “**STYLOCORYNE**, Cav.” Adde:  
 2. **S. elliptica**, Thw.—*Coffea (?) elliptica*, Thw. in hac Enum. p. 154.—C.P. 3482.  
*Bacca* pollicaris, carnosae, loculis circiter 10-spermis. *Semina* complanata, oblonga, semipollicaria, imbricata, biserialia, pulpa immersa.
159. “**GARDENIA**, Ellis.” Adde:  
 2. **G. latifolia**, Ait. (W. et A. Prod. i. p. 395, cum syn.; Wight, Icon. t. 759.)—C.P. 3617.  
 HAB. Near Colombo, *Mr. W. Ferguson*.
159. “**Valeriana Hardwickii**, Wall.” Adde ad descriptionem:  
*Achaenia* in planta Zeylanica glabra et multo majora quam in specimine ex Herb. Ind. Orient. cl. Hook. f. et Thoms.
160. Pro “1. **Dipsacus Leschenaultii**, Coul.,” et sequent., substitue:  
 1. **D. Walkeri**, Arn. Pug. p. 26 (334); Wight, Icon. t. 1166.—C.P. 2404 (1730).  
 HAB. Newera Ellia.  
 A *D. Leschenaultii*, Coult., cui valde similis, forma palearum satis differt fide specimin. ex Herb. Ind. Or. cl. Hook. f. et Thomson.
160. “**Vernonia Neilgherryensis**, DC.” Adde:  
*Achaenia immatura* glandulis conspersa. *Pappi* series externa brevissima, parum expansa.
161. “**Vernonia Gardneri**, Thw.” Pro “*pappo exteriori* brevissimo,” substitue: *pappi* serie externa brevi, subpaleacea, distincta vel ad basin in tubum plus minus cohaerente.  
 Facie externa *Decaneuro Silhetensi*, DC., valde similis, sed forma achæni pappique satis distinguitur.

Page  
163.

## "BLUMEA, DC." Adde:

6 a. **E. balsamifera**, DC. Prod. v. p. 447, cum syn.—c.p. 3665.  
HAB. Trincomalie, *Rev. S. O. Glenie*.

165. Adde: 23 a. **XIMENESIA**, Cav.

1. **X. eucelioides**, Cav.; DC. Prod. v. p. 627.—*a. hortensis*.—  
c.p. 3839.

HAB. Trincomalie, *Rev. S. O. Glenie*.

166. "GNAPHALIUM, Linn."

Pro "**G. Indicum**, Linn. (DC. Prod. vi. p. 231; DC. l. c. p. 274)," substitue:

**G. oblongum**.—*Anaphalis oblonga* et *An. (?) elliptica*, DC. l. c. p. 274. *A. (?) elliptica*, Wight, Icon. t. 1118. Et adde:

Specimina Zeylanica veri *Gnaph. Indici*, Linn., nunquam vidi, et nomen in hac Enumeratione meo errore insertum.

166. "**G. Wightianum**, Wall. Cat." Post "c.p." dele "528, 132, 568," et dele quoque a "Suffruticosum" ad finem annotationis. Post "t. 1117" insere: An *Anaphalis araneosa*, DC. l. c. p. 275?

166. Adde: 4 a. **G. adnatum**, Wall. Cat.; DC. Prod. vi. p. 274.—An *Anaphalis leptophylla*, DC. l. c. p. 273?—c.p. 568.

Var.  $\beta$ . *spathulifolium*; fruticosum, ramosum; *foliis* (superioribus exceptis) spathulatis, parum acutis, trinerviis, superne araneosis vel subglabris, subtus dense floccoso-tomentosis; *floribus* fere ut in var. *a*, sed squamis involuc. parum majoribus.—c.p. 132, 528.—An species distincta?

166. "**G. marcescens**, Wight." Adde: An *Anaphalis Notoniana*, DC. Prod. vi. p. 273, cum syn.?

167. "**Emilia prenanthoides**, DC." Pro "sublirataque" lege: sublyrataque.

167. "**Senecio corymbosus**, Wall." Post "1130.)" dele "c.p. 565;" et post "c.p. 272" adde: 565. Post "1131.—" insere: *S. araneosa*, DC. l. c. Adde annotationem sequentem:

Nunquam specimina Zeylanica formæ (var. *a*) *S. corymbosi*, Wall., foliis subtus tomento compacto ochraceo vel ochraceo-fulvo persistente tectis vidi, sed in aliis characteribus planta Zeylanica cum hac forma omnino congruit.

168. "**Stylidium uliginosum**, Sw." Pro "c.p. 2505" lege: c.p. 3505.

169. "CAMPANULACEÆ." Adde:

## 3. CEPHALOSTIGMA, A. DC.

1. **C. spathulatum**, Thw.; herbacea, hispidula; *caule* erecto, superne paniculatim ramoso, 4-5-costato; *foliis* lanceolatis, inferioribus spathulato-lanceolatis, acutis, parum dentatis; *floribus* minutis, longe pedicellatis; *lobis calycinis* lanceolatis, acutis; *corollæ* lobis ciliatis; *stigmatibus* 3-lobo.—c.p. 3794.

HAB. Oova District, at an elevation of 4000 feet.

Herba 4-16-pollicaris. *Folia* utrinque pilosa, 5-15 lin. longa,  $1\frac{1}{2}$ - $3\frac{1}{2}$  lin. lata. *Capsula* sphaerica.

- Page  
172. “**Utricularia orbiculata**, Wall.” Pro “c.p. 2085” lege: c.p. 2084.
173. “**ARDISIA**, Linn.”  
Pro “**A. paniculata**, Roxb.,” et seq. ad “1215,” substitue:  
**A. Courtallensis**, Wight, Ic. t. 1215.—*A. Missionis*, var.  $\beta$ .  
*verticillata*, A. DC. Prod. viii. p. 130. Et adde:  
Ab *A. paniculata*, Roxb., cui valde similis et forsan ejusdem varietas, differt foliis minoribus, venis primariis minus regularibus distinctisque, calycis segmentis acutioribus et corolla glandulosa.
174. “**A. divergens**, Roxb., var.  $\beta$ .” Dele “An *A. crispa*, var.  $\beta$ , A. DC. l. c. 134, cum syn.?”
- 175, 176. Genera “**DASYAULUS**,” Thw., “**DICHOPSIS**,” Thw., et “**ISONANDRA**,” Wight,” melius, ut credo, sectiones **BASSIÆ** formabunt.
178. “**DIOSPYROS**, Dal., Linn.” Adde:  
1 a. **D. montana**, Roxb. Pl. Cor. t. 48; Roxb. Flor. Ind. ii. p. 538; Wight, Ic. t. 1225.—c.p. 3774.  
HAB. Trincomalie, *Rev. S. O. Glenie*.
178. “**D. cordifolia**, Roxb.” Dele “*D. montana*, Wight. Ic. t. 1225 (non Roxb.)” Una cum annotatione tota.
180. “**D. oocarpa**, Thw.” Post “suffultis” insere: *D. Arnottiana*, Miq., fide specim. ex Herb. Ind. Orient. cl. Hook. f. et Thoms.
180. Adde speciem sequentem *Diospyri*:  
9 a. **D. pruriens**, Dalz. in Hook. Journ. of Bot. iv. (1852) p. 110?—c.p. 2836.  
HAB. Saffragam District, at an elevation of 2000 feet.  
*Ramuli juniores* patentim longiuscule hispido-pilosi. *Folia* superne glabra, costa pilosa, subtus sparse longiuscule pilosa. Inflorescentiam nondum vidi. Forsan species nova.
183. “**MACREIGHTIA**, A. DC.”  
Adde ad char. gen. quoad species Zeylanicas:  
FLOR. MASC. *Calyx* tridentatus, uno latere profunde fissus. *Stamina* indefinita, 4–16, filamentis prope basin corollæ insertis, longitudine valde inæqualibus, antheris oblongis lateraliter dehiscentibus. *Ovarii rudimentum* parvum vel subnullum. FLOR. FÆM. *Ovarium* 6-loculare, loculis uniovulatis. *Stylus* cylindricus vel parum conicus. *Stigma* parvum, 3-partitum. Arbores mediocres, *partibus junioribus inflorescentiaque* pilosis; *gemmis squamis* distichis ovatis imbricatis; *foliis* penniveniis, superne sublaevibus, ad costam venas primariasque depressis, subtus prominentim reticulatis; *petiolo* crassiusculo; *floribus* subsessilibus, secus ramulos juniores infra folia singulatim dispositis, vel breve spicatis ab axillis foliorum annotinorum vel delapsorum.
183. “**M. oblongifolia**, Thw.” Adde ad descriptionem:  
FLOR. MASC. *Corollæ* tubus quam calyx dimidio longior. *Stamina* circiter 12. *Ovarii rudimentum* minutum, pilosum.

Page

183. Adde species sequentes *Macreightiæ* :
2. **M. ovalifolia**, Thw. ; *foliis* coriaceis, ovatis, parum acuminatis, obtusiusculis, basi rotundatis vel parce angustatis. FLOR. MASC. rameis.—C.P. 3717.  
 HAB. Hinidoon Pattoo, at no great elevation.  
*Folia* 2–3½ poll. longa, 1–1¾ poll. lata, *petiolo* 2–3 lin. longo. FLOR. MASC. *Calyx* 2½ lin. longus, tubo corollæ æquilongus. *Stamina* 13–16. *Ovarii rudimentum* subnullum.
3. **M. acuminata**, Thw. ; *foliis* membranaceis, ovatis vel ellipticis, valde acuminatis, basi rotundatis vel parum acutis, brevipetiolatis. FLOR. MASC. *Corollæ tubo* quam calyx duplo longiore ; *staminibus* 4–5 ; *ovarii rudimento* parvo, oblongo, apice attenuato, piloso.—C.P. 3718.  
 HAB. Hinidoon Pattoo.  
*Folia* (cum acumine) 3–5 poll. longa, 1–1½ poll. lata, *petiolo* 1½–2 lin. longo. *Corolla* 4 lin. longa. FLOR. FEM. *Nuculania* sphaerica, tomentosa et sparse longiuscule pilosa, 6–7 lin. in diam.
190. “**JASMINUM**, Tourn.”  
 Pro “**J. rigidum**, Zenk.,” cum sequent. ad “p. 55,” substitue :  
**J. glabriusculum**, Blume, Bijdr. p. 679 ; DC. Prod. viii. p. 303, fide specim. ex Herb. Hort. Bot. Javan.—[Conf. *J. rigidum*, Zenk. ; DC. l. c. p. 305. *J. myrtifolium*, Zenk. ; DC. l. c. p. 304. *J. ligustrifolium*, Wall. ; DC. l. c. *J. tetraphis*, Wight et Gardn. ; Walp. Ann. iii. p. 21.]
191. “**Ophioxylon serpentinum**, Willd.” Pro “1834” lege : 1836.
191. “**O. densiflorum**, Thw.” Pro “1836” lege : 1834.
193. “**Wrightia angustifolia**, Thw.” Adde ad descriptionem :  
 Folliculi cylindrici, attenuati, nigri, nitidi, 8–10 poll. longi, 2 lin. in diam.
197. “**TYLOPHORA**, R. Br.”  
 “**T. fasciculata**, Ham.” Pro “c.p. 369” substitue : c.p. 3792. Et pro “HAB. Not uncommon” et sequent. ad “7000 feet” substitue :  
 HAB. Moottettoowegama.
197. Adde species sequentes *Tylophoræ* :
- 4 n. **T. Iphisia**, A. DC. Prod. viii. p. 610, cum syn. ; Wight, Icon. t. 1276.—*T. fasciculata*, hujus Enum. p. 197 (haud Ham.).—C.P. 369.  
 HAB. Common in the more elevated parts of the Central Province.
6. **T. membranifolia**, Thw. ; volubilis, glabra ; *foliis* tenuiter membranaceis, ovatis subito breviacuminatis, acutis, mucronatis, plus minus cordatis, longiuscule petiolatis ; *petiolo* costaque superne puberulis ; *pedunculis* filiformibus, folio subæquilongis, flexuosis, ad flexuras apicemque florum fasciculos gerentibus ; *pedicellis* flore triplo longioribus ; foliolis *coronæ stamineæ* depressis, carnosis, lateraliter parum compressis ; *calycis* segmentis ovatis, parum acutatis.—C.P. 3793.  
 HAB. Colombo, Mr. W. Ferguson.

Page

Differt a *T. asthmatica*, W. et A., *foliis* plus membranaceis, pedunculis elongatis, floribus multo minoribus et segmentis calycinis latioribus.

200. “**Fagræa Zeylanica**, Thunb.” Adde: Abundant on the banks of the river at Balangodde.

201. “**STRYCHNOS**, Linn.”

Dele “3. **S. laurina**, Wall.,” cum tota descriptione, et insere:

3. **S. micrantha**, Thw.; scandens; *ramulis junioribus* teretibus, pilosulis vel subglabris; *cirrhis* gemellis ad apicem ramuli abbreviati æquilongi, medio incrassatis; *foliis* glabris, firmis, ovatis vel ellipticis vel anguste lanceolatis, plus minus acuminatis, mucronulatis, tri- vel triplinerviis, basi parum acutatis, petiolatis; *cymis* axillaribus terminalibusque, a basi ramosis, folio multum brevioribus, breve pilosis; *floribus* parvis, pentameris; *segmentis calycinis* brevibus, rotundatis, ciliatis; *corollæ* fauce villosa, tubo lobis æquilongo; *ovario* oblongo, villoso, calycem excedente, in stylum longiusculum sensim attenuato; *stigmatē* parvo.—c.p. 1866, 3540, 3720.

HAB. Warmer parts of the island, not uncommon.

*Folia* 1½–5 poll. longa, 1–2 poll. lata, *petiolo* 3–4 lin. longo. *Cymæ* pollicares. *Pedicelli* flore æquilongi vel abbreviati. *Corollæ* tubus ½ lin. longus. *Nuculania* (immat.) pollicaria.

Differt a *S. laurina*, Wall., inter alia, floribus multo minoribus.

3/1. **S. coriacea**, Thw.; scandens?; *ramulis junioribus* pilosulis; *foliis* coriaceis, ellipticis, parum acuminatis, triplinerviis, subtus pallidioribus, reticulato-venosis, brevipetiolatis; *cymis* axillaribus terminalibusque, pilosulis, folio brevioribus vel subæquilongis; *segmentis calycinis* rotundato-oblongis, obtusis vel parum acutatis, breviciliatis; *corolla* quam calyx 5–6-plo longiore, lobis interne usque ad apicem villosis, tubo æquilongis; *ovario* oblongo, villoso, calyce subæquilongo, in stylum longiusculum attenuato; *stigmatē* parum dilatato.—An *S. laurinae*, Wall., var.?—c.p. 3367.

HAB. Central Province, rare.

*Folia* 1½–3 poll. longa, 1–1½ poll. lata; *petiolo* 1 lin. longo. *Corollæ* tubus 1 lin. longus. *Nuculania* nondum vidi. Mea specimina ecirrhosa.

201. “**S. cinnamomea**, Thw.” Post “polyspermis” insere: An *S. Wallichianæ*, Steud.; DC. Prod. ix. p. 13, forma?

201. “**S. potatorum**, Linn. f.” Pro “c.p. 1866, 3367” substitue: c.p. 3719.

202. “**GÆRTNERA**, Lam.” Adde:

6. **G. divaricata**, Thw.—*Pristidia divaricata*, Thw. in hac Enum. p. 149.—c.p. 2991.

HAB. Hinidoon Pattoo.

*Nuculania* subsphærica, cyanea, semipollicaria.

204. “**Pladera pusilla**, Roxb.” Adde annotationem sequentem:

Planta Zeylanica differt a speciminibus Indicis caule robustiore, ramis plus erectis, foliis latioribus, et pedicellis brevioribus, sed vix species distincta censenda.



Page  
205.

## "OPHELIA, Don." Adde:

2. **O. Chirata**, Griseb. (DC. Prod. ix. p. 127, cum syn.)—C.P. 3723.

HAB. Central Province, rare.

207. " **Didymocarpus floccosa**, Thw." Post "consperso" insere: *D. primulaefolia*, Bot. Mag. t. 5161 (haud Gardn.).

211. "IPOMŒA, Linn."

" **I. rugosa**, Choisy." Pro "c.P. 2854" lege: c.P. 2850.

211. " **I. jucunda**, Thw." Adde ad descriptionem:

Noctiflora. *Pedunculi* 1-3-(sæpius 1-)flori. *Calyx* pollicaris, externe pilosus, basi glandulosus, segmentis ovatis mucronatis. *Corolla* hypocrateriformis, alba, 4 poll. in expans., tubo bipollicari, externe pilosula. *Stamina* alba. *Stigma* capitatum, bilobum.

212. Adde speciem sequentem *Ipomœæ*:

9/1. **I. vitifolia**, Sweet. (Choisy, in DC. Prod. ix. p. 361, cum syn.)—C.P. 3685.

HAB. Hapittigam Corle.

218. "PTEROSTIGMA, Benth." Adde:

3. **P. (Cardiosepalum) subrepens**, Thw.; villosulum; *foliis* late ovatis, crenatis, petiolatis; *floribus* axillaribus, solitariis, brevipedicellatis, bibracteolatis; *bracteolis* filiformibus, calyce dimidio brevioribus; *calycis* segmentis 3 exterioribus ovatis vel rhomboideo-ovatis, acutis, reticulato-venosis, pilosulis (postico parum majore), 2 interioribus (lateralibus) lineari-lanceolatis et parum brevioribus, glabris.—C.P. 3821.

HAB. Pasdoon Corle, on the borders of paddy-fields.

*Folia*  $\frac{3}{4}$ - $1\frac{1}{4}$  poll. longa, denique subglabra; *petiolo* 2-4 lin. longo. *Calyx* 4 lin. longus, pedicello  $\frac{1}{2}$  lin. longo. *Corolla* pallide cyaneo-purpurea, semi-pollicaris, labio superiore rotundato. *Filamenta* anteriora longiora. *Antheræ* biloculares, loculis disjunctis, inæqualibus, ambobus sæpius polliniferis.

218. " **Herpestis Monnieria**, H. B. K." Pro "c.P. 2031" lege: C.P. 2034.

219. "BONNAYA, Link et Otto." Adde:

3. **B. brachiata**, Lk. et Ott. (Benth. in DC. Prod. x. p. 420, cum syn.)—C.P. 3084.

HAB. Hotter parts of the island.

A *B. veronicaefolia*, Spr., cujus forsitan varietas, differt foliis ad basin latioribus, serraturis setaceo-apiculatis.

219. " **B. veronicaefolia**, Spr." Dele "3084."

220. Adde: 11/1. PEPLIDIUM, Delile.

1. **P. humifusum**, Del. (Benth. in DC. Prod. x. p. 422, cum syn.)—C.P. 3789.

HAB. Trincomalie, Rev. S. O. Glenie.

- Page  
222. "OROBANCHACEÆ." Adde :
3. **CAMPBELLIA**, Wight.  
Walp. Ann. iii. p. 207.
1. **C. cytinoides**, Wight. (Walp. l. c. ; Wight, Icon. t. 1425, sub nomine *C. Neilgherrica*, Gardn.)—c.p. 3770.  
HAB. Pedrotalagalla, at an elevation of 7000 feet, on roots of species of *Strobilanthes*.
230. " **Barleria Prionitis**, Linn." Adde : c.p. 2012.
233. " **Gymnostachyum hirsutum**, T. And." Adde : c.p. 2473.
233. " **Justicia Betonica**, Linn." Adde : c.p. 1990.
235. " **Asystasia Gangetica**, T. And." Pro "c.p. 1998" lege : c.p. 1988.
248. " **Euxolus polygonoides**." Adde : *Amaranthus Blitum*, Linn., fide specim. ex Herb. Bot. Calc., differt ab hac specie, cui valde similis, dehiscencia circumscissali utriculi, et semine minore gibbosioreque.
248. " **Ærva Monsonia**, Moq." Adde : c.p. 3324.
251. " **Drymispermum cauliflorum**, Thw." Pro "ab marginem" lege : ad marginem.
256. " **Actinodaphne glauca**, N. ab Es." Pro "Halagalla" lege : Pedrotalagalla.
256. " **A. stenophylla**, Thw." Pro "303" lege : 304.
257. " **A. molochina**, N. ab Es." Pro "2272 v. 2373" lege : 2273.
259. " **ELATOSTEMA**, Forst." Adde :
3. **E. sessile**, Forst. (Wedd. Mon. des Urtic. p. 294.)—*E. cuspidatum*, Wight, Icon. t. 2091, fig. 1, t. 1983.—c.p. 3767.  
HAB. Rambodde, at an elevation of 4000 feet.
262. Post " **Artocarpus integrifolia**, Linn.," insere : c.p. 2233.
263. " **Antiaris innoxia**, Bl." Post "p. 193" insere : *A. Zeylanica*, Seemann, in 'Bonplandia,' x. p. 4. Et adde :  
Planta Zeylanica variat foliis subglabris, et inter eam et *A. saccidorum*, Dalz., nullam differentiam invenire possum.
264. " **Taxotrophis Zeylanica**, Thw.," pro "refluxis" lege : reflexis.
269. **EUPHORBIA**, Linn.  
Pro "9. **E. bifida**, Hook. et Arn.," et sequent. ad "c.p. 2131," substitue :
9. **E. Atoto**, Forst. (Boissier in DC. Prod. xv. p. 12.)—*E. halophila*, Miq. ; Boiss. l. c. p. 12, cum syn.—c.p. 2131. Et adde annot. seq. :  
Hæc species differt ab *E. bifida*, Hook. et Arn., seminibus glaberrimis.
270. Dele : " \* *Monoica*."
271. Dele : " \*\* *Dioica*."

Page

274. "**Podadenia sapida**, Thw." Inter "*Flores*" et "*rubri*" insere: dioici.

274. "**MACARANGA**, Dup. Th." Adde:

2. **M. Indica**, Wight, Icon. t. 1883 et t. 1949, f. 2.—c.p. 3795.  
HAB. Very abundant up to an elevation of 4000 feet.

274. "**Gelonium lanceolatum**, Willd." Pro "695" lege: 696.

275. "**Mischodon Zeylanicus**, Thw." Adde: Nom. vulg. "*Tammanna*."

277. "**Trigonostemon diplopetalus**, Thw." Pro "2 poll. lata" ad finem descriptionis, lege:  $\frac{1}{2}$  poll. lata.

280. "**AMANOA**, Aubl." Adde, ut synonym. hujus generis: *Leiopyxis*, Miq. Fl. Ind. Bat. Supp. i. p. 445.

Dele "1. **A. patula**, Thw.," cum tota descriptione, et in ejus loco insere duas species sequentes:

1. **A. Indica**, Wight, Icon. t. 1911.—*A. robusta*, Thw. MSS. (specim. distrib.) An *Cluytia patula*, Roxb. Pl. Cor. ii. p. 38, t. 170?—c.p. 3721 (2151 partim). Forma minor, *foliis* minus coriaceis, *petalis* parce pilosis vel glabris.—c.p. 2151.

HAB. Warmer parts of the island, not uncommon.

FLOR. FÆM. *Discus* cupuliformis, profunde 4-lobus, lobis apice dentatis, quam petala longioribus.

1/1. **A. acuminata**, Thw.; glabra; *foliis* ellipticis vel rotundatis, rostrato-acuminatis, subtus venis primariis prominentibus, basi rotundatis vel parum angustatis; *floribus* confertis, axillaribus; *petalis* oblongis, glabris, apice dentatis; *flor. fæm. disco* petalis multo brevioribus, dentato vel crenulato; *capsulis* sessilibus, parum depressis, latioribus quam longis, parce pilosis.—c.p. 2164.

HAB. Colombo, *Mr. W. Ferguson*; Galle District.

*A. Indica*, Wight, valde similis, sed *foliis* abrupte et longius acuminatis, *floribus* axillaribus (rarius rameis vel glomerato-spicatis), *petalis* forma differentibus et *disco flor. fæm.* multo brevioribus.

283. **Epistylum montanum**, Thw., est forma solum *E. polyphylli* hujus Enum. p. 283, et, ut species, delendum.

292. "**CHAVICA**, Miq." Adde:

2/1. **C. Chuvya**, Miq. (Illust. Piper. p. 42, t. 39.)—c.p. 3691.

HAB. Near Colombo, *Mr. W. Ferguson*; Central Province, but probably introduced.—Nom. vulg. "*Seewiya-wel*."

293. Adde: 5. **MULDERA**, Miq.

1. **M. diandra**, Thw.; glabra; *foliis* oblongo-lanceolatis, parum acuminatis, mucronatis, basi obtusiusculis, 3- vel sub-5-nerviis; *venis primariis* supra basin 2, a costa infra medium assurgentibus et apici folii attingentibus; *spicis masc.* folio brevioribus; *floribus* parum distantibus; *staminibus* 2.—c.p. 3738.

HAB. Mapalagam, in the Hinidoon Pattoo.

Scandens, radicans. *Folia* late virentia, subcarnosa, 4-7 poll. longa,  $1\frac{1}{2}$ - $2\frac{1}{2}$  poll. lata; *petiolo* 3-7 lin. longo.

- Page  
295. Dele: "4. **Liparis odorata**, Lindl.," cum tota descriptione.
296. Adde: 2/1. **EMPUSA**, Lindl.  
1. **E. paradoxa**, Lindl. Gen. et Sp. Orchid. p. 17, cum syn.—*Liparis odorata*, hujus Enum. p. 295 (haud Lindl.). An Rheede, Hort. Mal. v. 12, t. 28?—c.p. 3180, 3375.  
HAB. Central Province, at an elevation of 2000 to 5000 feet.  
*Flores* inodori.
297. " **Microstylis purpurea**, Lindl." Adde: c.p. 3768.  
HAB. Hinidoon Patta.
297. " **DENDROBIUM**, Sw." Adde:  
6/1. **D. hæmoglossum**, Thw.—*Caulibus* elongatis, pendulis, teretibus, parum compressis, estriatis; *foliis* lineari-lanceolatis, acuminatis, firmis; *pedunculis* oppositifoliis, brevissimis, bifloris; *sepalis* *petiolisque* lanceolatis, conformibus; *labelli* limbo linguæformi, acuto, glabro, undulato; *columna* apice 5-dentata.—c.p. 3842.  
HAB. Matelle, *Gardner*; Dolosbagey District.  
*Caules* 1-1½-pedalis. *Folia* 2½-3½ poll. longa, 3-4 lin. lata. *Flores* semipollicares flavi. Labellum, cum columna, rubro-tinctum.  
Flower described from a coloured drawing in this collection.
299. " **ERIA**, Lindl." Adde:  
1/1. **E. tricolor**, Thw.—*Caule* elongato, squamis marcescentibus vaginato; *foliis* plurimis, glabris, lanceolatis; *racemis* plurifloris, quam folia brevioribus; *rachi* villosula; *bracteis* ovatis, 1-nerviis; *sepalis* *petalisque* ovatis, subacutis, æquilongis, his angustioribus, illis externe cum ovario villosulis; *labelli* trilobi lobo medio rotundato, obtuso, apice reflexo, lateralibus erectis, subfalcatis, obtusis; *columna* antice medio parum gibba.—c.p. 3840.  
HAB. More elevated parts of the Central Province.  
*Folia* 5-9 poll. longa, ½-¾ poll. lata, striata. *Rachis* cum bracteis viridis. *Flores* 6 lin. in exp. *Petala* *sepalaque* alba. *Labellum* roseum, lobo medio flavo.  
Ab *E. bicolore*, Lindl., cui valde affinis, racheos florumque colore, et his minoribus numerosioribusque, satis differt.
302. " **CYRTOPERA**, Lindl."  
Pro: " **C. Gardneri**, Thw.," substitue: **C. fusca**, Wight, Icon. t. 1690; nomen prioritatem habens.
302. **Luisia bicaudata**, Thw., est species *Vandæ*. Vide infra.
303. " **VANDA**, R. Br." Adde:  
3. **V. bicaudata**, Thw.—*Luisia bicaudata*, Thw., in hac Enum. p. 302.—c.p. 3494.
303. " **SACCOLABIUM**, Blume." Adde:  
9. **S. paniculatum**, Wight, Icon. t. 1676.—c.p. 3769.  
HAB. Doombera District.
305. " **AERIDES**, Lour." Adde:  
"4. **A. latifolium**, Thw.; *caule* brevi, *foliis* membranaccis, ob-

Page

ovato-lanceolatis, subacutis, apice parum obliquis, mucronatis, ad basin gradatim angustatis; *paniculis* patentim ramosis, plurifloris, quam folia longioribus; *sepalis* oblongis, subacutis, *petalis* conformibus; *labelli calcare* conico; *laciniis lateralibus* erectis, versus apicem parum dilatatis, truncatis, basi interne callosis; *intermedia* latiore, obovata, retusa, basi glandula bicorni appendiculata.—C.P. 3495.

HAB. On trees, Saffragam District and Hinidoon Pattoo, at no great elevation.

5. **A. viridiflorum**, Thw.; *caule* brevi; *foliis* oblongis vel ovatis, parum acutatis, basi latis; *racemis* foliis subæquilongis, 4-8-floris; *sepalis* ovatis, subacutis; *petalis* angustioribus, oblongis; *labello* saccato, lobis erectis, lateralibus oblongis, intermedio minore rotundato.—C.P. 3385.

HAB. On trees in the Central Province, Gardner.

*Folia* circiter 3 poll. longa, 1 poll. lata. *Flores* conniventes, semipollicares, virides vel flavo-virides, labello albo, interne rubro. *Bracteæ* parvæ.

Described from coloured drawings in this collection.

306. "DENDROCOLLA, Blume." Adde:

2. **D. pulchella**, Thw.—*Cylindrochilus pulchellus*, Thw. in hac Enum. p. 307.—C.P. 2354.

307. Dele: "CYLINDROCHILUS, Thw.," genus a *Dendrocolla* haud distinctum. *D. pulchellus*, Thw., est *Dendrocolla pulchella*, Thw. Vide supra.

311. "Sect. 5. ARETHUSEÆ, Lindl." Adde:

Subsect. POGONIDÆ, Lindl.

48/1. POGONIA, Juss.

1. **P. Juliana**, Wall. Cat. no. 7399.—Lindl. Gen. et Sp. Orchid. p. 414.—C.P. 3841.

HAB. Doombera District.

315. "Zingiber Wightianum, Thw." Adde ad descriptionem: *Spica* radicalis vel rarius terminalis.

316. "Curcuma albiflora, Thw." Post "spica radicali" insere: vel rarius terminali.

317. "AMOMUM, Linn." Adde:

5/1. **A. graminifolium**, Thw.; *foliis* firmis, linearibus, acuminatissimis, in petiolum brevem angustatis; *ligula* brevi, indivisa, rotundata vel truncata, marcescente.—C.P. 3820.

HAB. Abundant in the Singherajah Forest.

3-4-pedalis vel ultra. *Rhizoma* rufescens, 3-4 lin. in diam. *Folia* 12-14 poll. longa,  $\frac{1}{2}$ - $\frac{3}{4}$  poll. lata. Inflorescentiam nondum vidi.

317. "Amomum masticatorium, Thw." Pro "semilunaria," lege: semilunari.

319. "ELETTARIA, Rheede.

Sect. *Geanthus*." Adde:

5. **E. rufescens**, Thw.; *foliis* glabris, anguste lanceolatis, longiuscule acuminatis, basi acutis, sessilibus; *ligula* brevi, indivisa, rotundata; capitulis (in meis speciminibus, an semper?)

Page

terminalibus; *bracteis* floribus æquilongis, rotundatis vel ovatis, obtusis.—C.P. 3732.

HAB. Dic Oya, in the Ambagamowa District.

1½-2-pedalis, rufescens. *Folia* 5-10 poll. longa, 1-2 poll. lata, firma. *Capitulum* 1½ poll. in diam., bracteis magnis involucreto. *Flores* flavi, parce rufescentes. *Corollæ* laciniæ exteriores oblongæ, apice rotundatæ. *Labellum* æquilongum, trilobum, lobo intermedio retuso.

319. “**E. involucrata**, Thw.” Post “glabris” adde: vel subtus plus minus tomentosus.

319. “**E. floribunda**, Thw.” Post “glabris” insere: vel subtus ad costam tomentellis.

326. “**Dioscorea spicata**, Roth.” Dele “?” post “571.”

330. “**CALAMUS**, Linn.”

Dele “**C. gracilis**, Roxb.,” cum tota descriptione, et in ejus loco insere duas species sequentes:

4. **C. pachystemonus**, Thw.; *frondibus* pinnatis, *vagina* aculeis rectis sparsis armata, ad apicem rugulosa, *laminis* 2-7 (sæpius 7), secus rachin aculeatam irregulariter dispositis, obovato-lanceolatis, cuspidato-acuminatis, plurinerviis (nerv. 3-5 robustioribus); *spadicibus* elongatis, decompositis, ramis 1-4, ramulis florigeris numerosis (18-25), alternis, brevibus, confertis, recurvis; *floribus* (masc.) confertis; *corolla* quam calycem 3-4-plo longiore; filamentis (3 interioribus valde) incrassatis.—C.P. 2334.

HAB. South of the island. Nom. vulg. “Kookool-wel.”

Species minor, caulibus tenuibus scandentibus. *Fronidium* rachis 3-7 poll. longa, laminæ transverse prominentim venulosæ, 5-12 poll. longæ, 1½-2¼ poll. latæ, 2 superiores basi cohærentes. *Spadicis* rami prim. 3-5 poll. longi. *Spathæ* infundibuliformes, truncatæ, brevispidatæ. *Bracteæ* bracteolæque patelliformes. FLOR. MASC. *Calyx* ad medium fissus, lobis rotundatis. *Corollæ* tubo brevi, lobis lanceolatis subacutis. *Ovarii rudimentum* parvum, trifidum. FLORES FÆM. nondum vidi.

A *C. gracili*, Roxb., differt laminis frondis paucioribus, ramulis spadicis masc. pluribus, et forma calycis.

4/1. **C. radiatus**, Thw.; *fronidium* laminis 6-7, ad apicem racheos aculeatæ radiatim aggregatis, linearibus, acuminatis; *spadicibus* elongatis, arcte aculeatis, decompositis vel supracompositis, ramis primariis 2-6, ramulis florigeris reflexis, parum distantibus, 3-15-floris; *alabastris* (masc.) cylindricis, falcatis; *drupis* sphericis, brevirostratis.—C.P. 3805.

HAB. South of the island. Nom. vulg. “Kookool-wel.”

Polygamo-dioicus, alte scandens, caulibus tenuibus. *Fronidium* vagina aculeis simplicibus vel seriatim cohærentibus dense armata, rarius rugosa subinermis; rachis 1½-3½ poll. longa; laminæ 7-10 poll. longæ, 6-8 lin. latæ. *Spathæ* clavato-infundibuliformes, brevispidatæ. *Bracteæ* bracteolæque poculiformes, minute cuspidatæ vel obtusæ. FLOR. MASC. *Calyx* cylindricus, duplo longior quam latus, breve 3-lobus, lobis triangularibus acutis. *Corolla* calyce fere triplo longior, tubo brevi, segmentis linearibus acutis. *Filamenta* dimidio inferiore incrassata. *Ovarii rudimentum* minutum, oblongum, apice trifidum. FLOR. FÆM. *Drupæ* 4 lin. in diam., squamis latiusculis flavis nigro-rubro-marginatis.

330. “**Calamus delicatulus**, Thw.” Adde ad descriptionem:

HAB. Hinidoon Pattoo. Nom. vulg. “Narrawell.”

Page

Polygamo-dioicus, alte scandens, caulibus tenuibus. *Fronde*s 2-3-pedales (juniores fugacim nigro-ferrugineo furfuraceæ); vagina aculeis longis (versus apicem 4-5-pollicaribus) applanatis albicantibus fragilibus dense armata. *Bracteæ* poculiformes, cuspidatæ. *Bracteolæ* patelliformes. FLOR. FÆM. *Corolla* calycem vix excedens. *Ovarium* oblongum, staminibus sterilibus in forma poculi sexdentati cinctum. *Stigmata* exserta, reflexa. *Drupæ* subsphæricæ, parum oblongæ, rostratæ, semipollicares, squamis ochraceis margine nigro-rubris ornatae.

334. "LAGENANDRA, Dalz." An species hujus generis rectius sectionem **CRYPTOCORYNIS**, Fisch., formarent?

334. "ARUM, Linn." Adde:

3. **A. (Typhonium) Roxburghii**, Schott, *Aroid.* i. p. 12; Synops. *Aroid.* p. 18.—*Arum trilobatum*, Roxb. *Fl. Ind.* iii. p. 505 (haud Linn.); Wight, *Icon.* t. 803.—C.P. 3764.

HAB. Colombo, *Mr. W. Ferguson*; Peradenia.

336. "COLOCASIA, Ray." Adde:

3/1. **C. fornicata**, Kunth, *Enum.* iii. p. 41.—*Alocasia fornicata*, Schott, *Syn. Aroid.* p. 47. *Arum fornicatum*, Wight, *Icon.* t. 792.—C.P. 3830.

HAB. Colombo, *Mr. W. Ferguson*.

339. Adde: 6/1. **ROXBURGHIA**, Dryand.

1. **R. gloriosoides**, Jones in Roxb. *Pl. Corom.* i. p. 26, t. 32; Kunth, *Enum.* v. p. 287 cum syn., var.  $\beta$ . *minor*.—C.P. 3775.

HAB. Trincomalie, *Rev. S. O. Glenie*.

*Folia* 2-3 poll. longa. *Perigonii* foliola 8 lin. longa.

Differt a forma communi hujus speciei nonnisi partibus omnibus multo minoribus.

339. Adde: 7/1. **LEDEBOURIA**, Roth.

1. **L. hyacinthina**, Roth. (Kunth, *Enum.* iv. p. 336, cum syn.; Wight, *Icon.* t. 2040.)—C.P. 3829.

HAB. Trincomalie, *Rev. S. O. Glenie*.

342. "CYPERUS, Linn." Adde species sequentes:

2/1. **C. pumilus**, Linn. (N. ab Es. in Wight's *Contrib.* p. 74, cum syn.; Kunth, *Enum.* ii. p. 4, cum syn.)—C.P. 3787.

HAB. Kornegalle District.

2/2. **C. stramineus**, N. ab Es. l. c.; Kunth, l. c. p. 9.—C.P. 3776.

HAB. Kornegalle District.

25/2. **C. venustus**, R. Br. (N. ab Es. l. c. p. 86, cum syn.; Kunth, l. c. p. 68, cum syn.)—C.P. 3788.

HAB. Trincomalie, *Rev. S. O. Glenie*.

344. "Cyperus distans, Linn." Adde:

Var.  $\beta$ . *major*; culmo elatiore, *spiculis* suberectis, *squamis caryopsique* parum longioribus.—C.P. 3814.

HAB. Trincomalie, *Rev. S. O. Glenie*; Dambool, *A. O. Brodie, Esq.*

Page

345.

## "PANDANOPHYLLUM, Hassk."

"Char. emend." Pro "*Squamæ* unifloræ" insere: *Spiculæ* androgynæ. Dele "FLOR. MASC." et sequent. ad: basi incrassatus, et insere: *Antheræ* lineares, longitudinaliter dehiscentes, denique exsertæ. *Filamenta* dilatata, linearia. *Ovarium* oblongum, in stylum filiformem attenuatum. Et adde: Plantæ immersæ vel ad margines rivulorum crescentes; *culmo* brevi, basi radices copiosas cylindricas tenaces emittente; *foliis* longissimis; *pedunculis* numerosis, inter folia assurgentibus, basi vaginatis; *capitulis* subsphæricis; *squamis* ovatis, integris, denique plus minus lacerato-fissis, singulis (inferioribus vacuis exceptis) spiculam tegentibus.

345. "Pandano-phyllum Zeylanicum, Thw." Post "scabris" insere: *pedunculis* plus minus elongatis, basi squamis vaginatis, superne nudis. Post "Kokoole Corle" adde: Abundant in the Singhe-rajah Forest, and between Kornegalle and Matelle. Pro "Culmi plures" lege: *Pedunculi* plures. In adnotatione, pro "male flower" substitue: young spicula; et pro "of the flower" lege: of the spicula.

346. Adde speciem sequentem **Pandanophylli**:

2. **P. immersum**, Thw.; *foliis* anguste linearibus, subulato-acuminatis, margine lævi vel sublævi; *pedunculis* brevibus, vaginis superioribus foliaceis et capitulum excedentibus.—c.p. 3819.

HAB. Singhe-rajah Forest, in streams.

A *P. Zeylanico* differt foliis angustioribus, margine sublævibus, et pedunculis brevioribus foliatisque.

346. "**Lepironia mucronata**, Rich." Dele "In *Pandanophyllum* the flower" et sequent. ad finem adnotationis.

347. "**ABILDGAARDIA**, Vahl." Species hujus generis sensim in *Fimbristylem* transeunt, et rectius subsectionem ejusdem formabunt. Vide infra.

348. "**Fimbristylis polytrichoides**, R. Br.?" Adde: c.p. 3786.

HAB. Trincomalie, Rev. S. O. Glenie.

348. "**Fimbristylis diphylla**, Vahl." Adde:

Var.  $\beta$ . *major*.—*F. rigidula*, N. ab Es. et hujus Enum. p. 348.—c.p. 3232.

348. "**Fimbristylis rigidula**, N. ab Es." Certe varietas *F. diphyllæ*, Vahl. Vide supra.

348. Sub "\*\* Stigmatibus 3. *Trichelostylis*" insere

† *Spicæ squamis 3-pluri-stichis*.

349. "**Fimbristylis insignis**, Thw." Post "Kornegalle" adde Pasdoon Corle.

349. "**Fimbristylis complanata**, Link." Post "(in *Herb. Hook.*)" adde: Very abundant in the south of the island.

349. Post descriptionem speciei 19mæ *Fimbristylis* insere:

†† *Spica complanata vel plus minus spiraliter torta.*

*Squamis distichis. Abildgaardia.*

349. "**Fimbristylis connectens**, Thw." Post "District" adde: Abundant at Newera Ellia.



Page

350.

Adde species sequentes *Fimbristylis* :

22. **F. tenuifolia**, Thw.; *culmo* filiformi, striato, glabro, basi foliato; *foliis* angustis, *culmo* brevioribus, marginibus involutis, scabrellis; *umbellæ* radiis 2-4, glabris; *involucro* brevi, triphylo; *spicis* oblongis, acutis, parum tortis, solitariis in radiis, intermedia sessili; *squamis* distichis, paucis, laxis, glabris, late ovatis, subacutis, infra apicem mucronulatis; *floribus* triandris; *stylo* glabro; *caryopsi* late obovoidea, trigona, aspero-tuberculata.—C.P. 3780.

HAB. Ambagamowa District.

Cæspitosa, 1-2-pedalis. *Spicæ* 2-3 lin. longæ. *Squamæ* nigro-ferrugineæ, margine membranaceæ pallidiores. *Caryopsis* pallide fusca.

23. **F. nigro-brunnea**, Thw.; *culmo* rigido, filiformi, sulcato, versus apicem scabro, basi foliato; *foliis* firmis linearibus, abrupte subacutis, glabris, margine scabridis, *culmo* sæpius multo brevioribus; *umbella* decomposita, aperta; *radiis* glabris; *involucro* *involucellis*que diphyllis, brevibus; *spicis* ovato-oblongis, subacutis, parum tortis; *squamis* distichis, paucis, laxis, glabris, nitidis, ovatis, mucronatis; *floribus* triandris; *stylo* glabro; *stigmatibus* 3; *caryopsi* obovoidea, trigona, aspero-tuberculata.—C.P. 3779.

HAB. Matelle East, A. O. Brodie, Esq. South of the island, very abundant.

Cæspitosa, 1-3-pedalis. *Spicæ* 3-4 lin. longæ. *Squamæ* nigro-brunneæ. *Caryopsis* albida.

24. **F. Cinnamometorum**, Kunth.—*Abildgaardia*, hujus Enum. p. 347, cum syn.—C.P. 2752.

25. **F. fulvescens**, Thw.—*Abildgaardia*, hujus Enum. p. 347, cum syn.—C.P. 679.

HAB. Common in the Southern and Central Provinces up to 4000 feet.

26. **F. tristachya**.—*Abildgaardia*, Vahl, et hujus Enum. p. 347, cum syn.—C.P. 852.

27. **F. monostachya**.—*Abildgaardia*, Vahl, et hujus Enum. p. 347, cum syn.—C.P. 3231.

351.

“SCIRPUS, Linn.” Adde:

6. **S. tetraqueter**.—*Eleocharis tetraqueter*, N. ab Es. et hujus Enum. p. 351, cum syn.—C.P. 2397.

7. **S. ovatus**, Roth. (Vahl, Enum. ii. p. 249.)—*Eleocharis ovata*, R. Br. et hujus Enum. p. 351, cum syn.—C.P. 2635.

8. **S. capitatus**, Willd. (Vahl, l. c. p. 250.)—*Eleocharis capitatus*, R. Br. et hujus Enum. p. 351, cum syn.—C.P. 3090.

9. **S. fistulosus**, Poir. Encyc. vi. p. 749.—*Eleocharis fistulosa*, Schult, et hujus Enum. p. 351, cum syn.—C.P. 3162 (3047).

10. **S. plantagineus**, Rottb.—*Eleocharis plantaginea*, R. Br. et hujus Enum. p. 352, cum syn.—C.P. 3046.

Var.  $\beta$ ; *culmis* tenuioribus; *squamis* paucioribus, angustioribus et magis distantibus.—C.P. 3777.—HAB. Kornegalle.

11. **S. spiralis**, Rottb. (Roxb. Fl. Ind. i. p. 212.)—*Eleocharis spiralis*, R. Br. et hujus Enum. p. 352, cum syn.—C.P. 853.

Page

12. **S. laxiflorus**, Thw.; *culmis* teretibus, aphyllis, striatis, interne regulariter cellulosus nec septis nodosis, basi vaginatis; *vaginis* tubulosis, membranaceis; *spica* solitaria, angusta, cylindrica; *squamis* laxis, ovatis, crebre striatis, late hyalino-marginatis; *caryopsi* late obovata, lenticulari-biconvexa, pluri-costulata, delicatule transversim striata; *setis* 7, caryopsi duplo longioribus, retrorsum spinulosis. An *Eleocharis Sieberi*, Kunth, Enum. ii. p. 153, forma?—c.p. 3762.

HAB. South of the island, and in the Ambagamowa District.

Cæspitosus, 1-2½-pedalis. *Culmi* (sicci) 1 lin. in diam. *Vaginæ* sæpius pulchre purpurascens. *Spica* pollicaris. *Squamæ* 1½ lin. longæ, pallide virides vel ochraceæ, margine hyalino delicatissime ferrugineo-punctulato. *Caryopsis*, cum *setis*, albida vel pallide ferruginea.

13. **S. Chætaria**.—*Chætocyperus setaceus*, N. ab Es. et hujus Enum. p. 351, cum syn.—c.p. 247.

351. “**Chætocyperus setaceus**, N. ab Es.” = *Scirpus Chætaria*. Vide supra.

351. “**ELEOCHARIS**, R. Br.” Cl. Benthamio duce, species hujus generis ad *Scirpum transtuli*. Vide supra.

352. “**RHYNCHOSPORA**, Vahl.” Adde:

3/1. **R. gracillima**, Thw.; glabra, *culmis* filiformibus, trigonis, glabris, foliatis; *foliis* culmo conformibus et eodem parum brevioribus; *corymbis* axillaribus terminalibusque, his spicas plures, illis paucas gerentibus, ramis capillaribus, longiusculis, singulis basi foliolis 2 inæqualibus suffultis; *spicis* ovatis, utrinque attenuatis, flores hermaphrod. circiter 4 includentibus; *squamis* late ovatis, parum acuminatis, mucronulatis, inferioribus minoribus vacuis, mucronatis; *floribus* diandris; *stylo* bifido; *caryopsi* rotundata, lenticulari-biconvexa, transverse rugosa, longitudinaliter striolata, basi styli sursum brevi conica deorsum divaricata bifida coronata.—c.p. 3818.

HAB. South of the island.

Dense cæspitosa, 1-2-pedalis. *Spicæ* 3-4 lin. longæ, ¾-1 lin. latæ, fusco-ferrugineæ. *Caryopsis* albida vel pallide fusca.

353. “**SCLERIA**, Linn.”

Pro “2. **S. stipularis**, N. ab Es. (Steud. l. c. p. 171, cum syn.?)” insere:

2. **S. hebecarpa**, N. ab Es. in Wight's Contrib. p. 117; Kunth, Enum. ii. p. 343; Steud. Synops. p. 169. Et dele “The present plant” et sequent. ad finem adnotationis.

354. Pro “8. **S. hebecarpa**, N. ab Es. l. c.; Kunth, l. c. p. 357; Steud. l. c.—c.p. 3318,” insere:

8. **S. Ceylanica**, Poir. Encyc. vii. p. 3; Steud. Synops. p. 179; N. ab Es. in Wight's Contrib. p. 118; Kunth, Enum. ii. p. 358.—c.p. 3318.

Forma *b*, rigidior, foliis angustioribus.—c.p. 3797.

Forma *c*, paniculis pubescentibus.—c.p. 3796. Adde, post

Page

“island:” Abundant on the borders of paddy fields in the Southern Province. Ad descriptionem adde:

Dense cæspitosa,  $\frac{1}{2}$ - $1\frac{1}{2}$ -pedalis. Culmi triquetri. Folia vagina plus minus alata vel acutangula, ligula brevi rotundata aliquando fissa. Paniculae axillares terminalesque, breves, paucifloræ. Caryopsis rotundata, apiculata, lævis vel obsolete rugulosa, nitida, alba.”

358.

“**PANICUM**, Linn.” Adde:

1/1. **P. corymbosum**, Roxb. Fl. Ind. i. p. 292.—Steud. Synops. p. 38, cum syn.—C.P. 3800.

HAB. Happootelle, at an elevation of 3000 to 4000 feet.

30/1. **P. stenostachyum**, Thw.; glabrum, culmis debilibus, elongatis, parce ramosis; foliis longiusculis, linearibus, acuminatis, flaccidis, margine scabriusculis, basi subauriculatis, ligula breviliata; panicula stricta, elongatula, rachi subglabra, ramis brevibus distantibus confertifloris; spiculis lanceolatis, pedicello longioribus; gluma inferiore ovata, subacuta, trinervia; superiore parum longiore, 5-nervia, mucronulata; floribus æquilongis, gluma superiore duplo longioribus; neutro bipaleaceo, palea inferiore late ovata, 7-nervia, apice breve bifida, mucronulata; hermaphrodito palea inferiore 5-nervia, mucronulata, minute rugulosa.—C.P. 3845.

HAB.—Trincomalie, Rev. S. O. Glenie.

Dense cæspitosum. Culmi 2-4-pedales, articulis inferioribus radicanibus. Panicula 4-8 poll. longa. Spiculae  $1\frac{1}{2}$  lin. longæ.

35/1. **P. asperum**, Wight.—Steud. Synops. p. 49.—C.P. 3846.

HAB. Trincomalie, Rev. S. O. Glenie.

Gluma inferior brevis, truncatula, subnervia, pallida; superior multoties longior, valde acuminata, plurinervia, aspera,  $2\frac{1}{2}$  lin. longa, basi pallida. Flos inferior masculus, bipaleaceus; palea inferiore subglabra, lanceolato-acuminata, quam gluma superior parum brevior. Flos superior foemineus, inferiore multo brevior, paleis tenere membranaceis.

358.

“**Panicum Helopus**, Trin.” Post “cum syn.” insere: C.P. 3847. Et post “HAB.” insere: Trincomalie, Rev. S. O. Glenie. Var.  $\beta$ .

364.

“**ISCHÆMUM**, Linn.” Adde:

5/1. **I. falcatum**, N. ab Es.—*Andropogon falcatus*, Steud. Synops. p. 369, cum syn.—C.P. 3322.

HAB. South of the island.

Differt ab *I. pectinato*, Trin., apice glumæ inferioris spiculæ sessilis brevialato, nec late appendiculato, ciliis angustioribus magis distantibus et fere duplo longioribus.

365.

“**Ischæmum pectinatum**, Trin.” Dele “*Andropogon falcatus*, Steud. l. c.—C.P. 3322.” Et insere: C.P. 3848. Pro “Hotter parts of the island, in wet ground,” substitue: Trincomalie, Rev. S. O. Glenie.

366.

“**ANTHISTIRIA**, Linn.” Adde:

2/1. **A. Cymbaria**, Roxb. Flor. Ind. i. p. 251.—Steud. Synops. p. 399.—C.P. 3803.

HAB. Matelle East.

Page

367. Pro "9. *Andropogon pertusus*, Willd.," cum tota descriptione, substitue duas species sequentes, illic errore inclusas :

9. *A. pertusus*, Willd. Sp. Pl. iv. p. 922; Roxb. Fl. Ind. i. p. 258.—C.P. 951.

HAB. Abundant in the warmer parts of the island.

*Spicæ* versus apicem pedunculi aggregatæ. *Gluma exterior* spicularum omnium vel tantum sessilium foveolata. *Spicula pedicellata* mascula unipaleacea vel neutra epaleacea.

9/1. *A. fascicularis*, Roxb. l. c. p. 265.—An *A. punctatus*, Roxb. l. c. p. 264?—C.P. 411.

HAB. Hot drier parts of the island.

*Spicæ* fasciculatæ, fasciculis distantibus. *Gluma exterior* spicularum omnium efoveolata vel spicul. nonnullarum sessilium foveolata. *Spicula pedicellata* mascula unipaleacea vel neutra epaleacea.

368. "HETEROPOGON, Pers." Adde:

3. *H. insignis*, Thw.; *culmis* robustis, erectis, rigidis, tere-tibus, foliatis, ima basi ramosis; *foliis* longiusculis, linearibus, acuminatis, sparsim longe pilosis, margine scabris, vagina glabra, apice longe pilosa, ligula brevi, rotundata, glabra; *spica* longiuscula; *spiculis* inferioribus omnibus masculis, superioribus masc. et fœm. intermixtis; *spicul. masc. gluma inferiore* lanceolata, nitida, inconspicue striata, apice obliqua, *superiore* lanceolata, 1-nervia; *spicul. fœm. glumis* æquilongis coloratis, coriaceis, ferrugineo-pilosis, marginibus involutis; *inferiore* ovato-oblonga, obtusa; *superiore* angustiore, apiculata; *arista* robusta, longissima, fulvo-pilosa.—C.P. 3804.

HAB. Mahning-galla, Matelle East.

Dense cæspitosus. *Culmi* 4-5-pedales. *Folia* 1-2-pedalia, 3-4 lin. lata. *Spica* (aristis inclus.) 8-12 poll. longa. *Rachis* inferiore parte glabra, superne ferrugineo-tomentosa. *Spiculæ masc.*: 6-10 lin. longæ, flavo-virides, superiores gradatim longiores; *paleis* lanceolatis, margine longe pilosis. *Squamulæ* oblongæ, apice dentatæ et longe pilosæ. *Spiculæ fœm.*: *Glumæ* nigro-ferrugineæ, 4 lin. longæ. *Arista* 4½ poll. longæ.

375. "Arundinaria debilis, Thw." Pro "semiverticillatis" lege: semiverticillatos.

375. "Bambusa attenuata, Thw." Pro "ramoso-striata" lege: nervoso-striata.

379. "ANGIOPTERIS, Hoffm."

Pro "*A. erecta*" lege: *A. evecata*.

379. "LYGODIUM, Sw."

Pro "3. *L. Japonicum*, Sw. (Presl; Benth. l. c. p. 442.)," substitue:

3. *L. flexuosum*, Sw.; Spreng. Syst. iv. p. 29; Beddome, Ferns of Southern India, t. lxiii.

381. "Acrostichum (Gymnopteris) decurrens, Hook." Clariss. Hooker, in litt. hanc plantam ab ea in Fil. Exot. et in Flor. Hongk. descripta distinctam indicat.

Page  
381.

“**VITTARIA**, Sm.” Adde:

3. **V. elongata**, Sw. (Willd. Sp. Pl. v. p. 406; Hook. Sp. Fil. v. ined.)—c.p. 3806.

HAB. Central Province.

In Herb. Perad. species, forsan indescripta (c.p. 3807), adest parva, *caudice* brevi, subrepente, radicante; *radicibus* pilosis; *frondibus* confertis, coriaceis, abrupte acuminatis, obtusiusculis, 2–6 poll. longis, 1½–2 lin. latis; *soris* immersis, a margine frondis parum distantibus.

382. “**Meniscium Thwaitesii**, Hook.” = *Polypodium (Goniopteris) proliferum*, Pr., vide infra.

383. “**Asplenium tenerum**, Forst.; Hook. l. c.?” Certe, ut clariss. Hooker ait, forma minor *A. macrophylli*, Sw.

384. “**A. macrophyllum**, Sw.” Post “c.p. 1340” insere: 1371. Et pro adnotatione substitue:

Ab *A. caudato*, Forst., differt frondibus magis carnosis, indusio latiore et demum margine arcte revoluto.

386. “**Pteris quadriaurita**, Retz.” Post “abortivis” insere: *P. Otaria*, Beddome, Ferns of South. Ind. t. xli. An hybrida inter *P. quadriauritam*, Retz., et *P. crenatam*, Sw.?

388. “**Lindsæa nitens**, Blume.” Pro “appropriated” lege: appropriate.

389. “**DAVALLIA**, Linn.

e. *Microlepia*.” Adde:

14. **D. proxima**, Blume, En. Fil. Jav. Fasc. ii. p. 238; Hook. Sp. Fil. i. p. 183?—c.p. 3827.

HAB. Oova District, at an elevation of 3000 feet.

*Caudex* repens, tomentosus, digiti crassitudine. *Stipites* epaleacei, subglabri, rubro-fusci, ½–1 ped. longi. *Fronde*s tripinnatæ, lanceolatæ vel deltoideo-lanceolatæ, subtus ad venas sparse pilosæ, 2–3-pedales, rachi scabriuscula. *Pinnæ* (*pinnulæ*que) approximatae, lanceolatæ, caudato-acuminatæ, alternæ, pinnula (vel lobo) basali sæpius parum majore. *Pinnulæ* pinuatifidæ vel ad basin pinnatæ, lobis falcato-trapezoideis, lanceolatis, acutiusculis vel brevibus, obtusis, crenato-dentatis. *Sori* paullo intramarginales.

390. “**Aspidium aristatum**, Sw.” Planta Zeylanica ob indusia reniformia melius, ut cl. Hance in litt. notat, in sectione *Lastrea* includendum.

390. “**A. pteropus**, Kze.” Adde:

Var.  $\beta$ ?; minor, *frondibus* deltoideis, paucilobatis; *stipite* plus minus alato vel exalato; *frond. sorifer.* segmentis multo angustioribus; *sporangiis* sæpius abortivis vel vacuis.—c.p. 3808.

HAB. Var.  $\beta$ ?, Peradenia, amongst plants of *Aspid. pteropus*, Kze., var. *a*, and *Acrostichum quercifolium*, Retz. Can it possibly be a hybrid between these two species, bearing, as it does, a certain resemblance to the latter?

392. “**A. concinnum**, Thw.” Adde: *Fronde*s steriles vix ab iis *A. deparioidis*, Hook., distinguendæ, cui hæc planta valde affinis et forsan ejusdem solum varietas.

392. “**A. sparsum**, Spr.” Post “1369” pro “?” insere,

Page  
394.

“POLYPODIUM, Linn.” Adde :

3/1. **P. decorum**, Brack. (Hook. Sp. Fil. iv. p. 179, cum syn.)  
—C.P. 3816.

HAB. Singhe-rajah Forest and in the Hinidoon Pattoo.

11/1. **P. (Goniopteris) lineatum**, Colebr. (Hook. Sp. Fil. v. p. 12, cum syn.)

HAB. Ceylon, *Walker (in Herb. Hook.)*.

11/2. **P. (Goniopteris) proliferum**, Pr. (Hook. l. c. p. 13, cum syn. *Meniscium Thwaitesii*, Hook. hujus Enum. p. 382.)—C.P. 3145.



## ADDENDA ET CORRIGENDA POSTERIORA.

Page  
28.

“MALVACEÆ.” Adde:

### 11. MALVASTRUM, A. Gray.

1. *M. carpinifolium*, A. Gray, Plant. Fendler. p. 22, adnot.;  
Walp. Ann. ii. p. 152, cum syn. —C.P. 3614.

HAB. Colombo, *Mr. W. Ferguson.*

61. Lin. 5. Dele “3596.”

78. Pro “4. GLYCYCARPUS, Dalzell,” substitue:

4. NOTHOPEGIA, Blume, Benth. et Hook. f. Gen. Pl.

*Glycyarpus*, Dalzell.

1. *N. Colebrookiana*, Blume, Mus. Bot. Lugd. Batav. p. 203;  
Walp. Ann. ii. p. 281.—*Pegia* (?) *Colebrookiana*, Wight, Ic. t. 236.  
*Glycyarpus racemosus*, Dalz. (Walp. l. c. p. 286); hujus Enum.  
p. 78.—C.P. 1260.

81. “CROTALARIA, Linn.” Adde:

24. *C. Willdenowiana*, DC. (W. et A. Prod. i. p. 191, cum  
syn.)—C.P. 3853.

HAB. Colombo, *Mr. W. Ferguson.*

84. “*Sesbania aculeata*, Pers.” Adde: Var. *sericea*, Benth. MSS.  
—C.P. 3850.

HAB. Colombo, *Mr. W. Ferguson.*

87. “*Desmodium ormocarpoides*, DC.” Adde: Prod. ii. p. 327.—  
C.P. 3812.

Post “(*Hb. Hook.*)” adde: Matelle East.

91. “RHYNCHOSIA, Lour.” Adde:

1/1. *R. suaveolens*, DC. (W. et A. Prod. i. p. 240, cum syn.)—  
C.P. 3851.

HAB. Near Trincomalie, *Rev. S. O. Glenie.*

95. “CASSIA, Linn.” Adde:

15. *C. obtusa*, Roxb. (W. et A. Prod. i. p. 288, cum syn.)—  
C.P. 3843.

HAB. Trincomalie, *Rev. S. O. Glenie.*

*Mr. Glenie* has also sent me, from Trincomalie, the foliage of what I take  
to be *Cassia suffruticosa*, Koenig.



Page

119. "**Careya arborea**, Roxb." Post "100.)" adde: An *C. sphaerica*, Roxb. Fl. Ind. ii. p. 636; Wight, Ic. t. 556? Et pro "biserialia" in descript. substitue: 2-6-serialia.
124. "**Zanonia Indica**, Linn." Adde ad descriptionem:  
 FLOR. FÆM. *Calycis* lobi 3-4, valvati, late triangulares, acuti, 2 lin. longi. *Petala* 5, imbricata, carnosae, ovato-oblonga, obtusa, pallide ochracea, 4 lin. longa, apicibus æstivatione valde incurvatis. *Stamina sterilia* 5, petalis alterna, brevia, alba, truncata. *Styli* 3, lobis calycinis parum breviores, late conici, divergentes, facie interna per longitudinem sulco angusto notati, ad apicem stigmatosi, bifidi, lobis reflexis. *Ovarium* 3-loculare, obconico-cylindricum, 6 lin. longum; loculis longis, 2-ovulatis; ovulis ab apice loculi pendulis, anatropis.
128. "**Begonia Thwaitesii**, Hook." Post p. 894, adde: An *B. tenera*, Dryand. in Linn. Trans. i. 169, t. 16.
151. "**BORRERIA**, Mey., DC." Adde:  
 2. **B. tetracocca**, Thw.; herbacea, procumbens, ramis elongatis, acute tetragonis, parum compressis; foliis lanceolatis, subsessilibus, scabris; stipulis lanceolatis, marginibus nervoque dorsali dense hispido-ciliatis; floribus paucis, 2-3 ad apices ramulorum brevium axillarium; calyce hispido, dentibus 4 lanceolatis obtusiusculis; corollæ tubo longiusculo; ovario quadriloculari, loculis monospermis; stigmatate bifido, piloso.—c. p. 3859.  
 HAB. Colombo, Mr. W. Ferguson.
176. "**ISONANDRA**, Wight." Adde:  
 5/1. **I. lanceolata**, Thw.; glabra, partibus junioribus fasciculisque florum adpresse ferrugineo-pilosis; foliis lanceolatis vel oblongo-lanceolatis, obtuse acuminatis, subcoriaceis, superne planis, subtus opacis inconspicue venosis, basi acutis, longiuscule petiolatis; pedicellis petiolo brevioribus vel subæquilongis.—c.p. 3679.  
 HAB. Near Caltura.  
*Arbor* mediocris. *Folia* 3-5 poll. longa, 7-14 lin. lata, petiolo pollicari. *Fasciculi* 3-6-flori. *Calyx* 4 lin. longus.
200. "**Fagraea Zeylanica**, Thunb." Adde: c.p. 1824.
204. "**CANSCORA**, Lam." Adde:  
 Pro "4. **C. perfoliata**, Lam." et seq. substitue:  
 4. **C. Roxburghii**, Arn. (Miquel, Anal. Bot. Ind. iii. p. 11.)—*C. perfoliata*, Griseb. in DC. Prod. ix. p. 65 (haud Lam.); Wight, Ic. t. 1327, f. 2.—c.p. 1874.
209. "**SESAMUM**, Linn." Adde:  
 2. **S. occidentale**, Heer et Regel. (DC. Prod. ix. p. 250.)—c.p. 3852.  
 HAB. Trincomalie, Rev. S. O. Glenie; Colombo, Mr. W. Ferguson. A common weed in the Botanic Garden, Peradenia.

Page

A *S. Indico* tota planta pilosiore, floribus fusco-purpureis, capsula obtusa, et seminibus majoribus differt.

221. "CHRISTISONIA, Gardner." Dele:  
"et forsan *Campbellia*, Wight, et *Oligopholis*, Wight, Ic."
222. "Christisonia unicolor, Gardn." Dele: "An *Campbellia cytinoides*, Wight, Ic. t. 1425, cum syn.?"
230. "BARLERIA, Linn." Adde:  
8. *B. nitida*, N. ab Es. in DC. Prod. xi. p. 224; Wight, Ic. t. 454.—C.P. 3849.  
HAB. Trincomalie, *Rev. S. O. Glenie*.
230. "*Barleria cristata*, Linn." Adde: c.p. 2899.
246. "SALSOLACEÆ." Adde:  
2/1. SALICORNIA, Tourn., Moquin.  
1. *S. brachiata*, Roxb. (Moq. in DC. Prodr. 13-2, p. 145, cum syn.; Wight, Icon. t. 738.)—C.P. 3855.  
HAB. Jaffna, *Mr. W. Ferguson*, common.
267. "*Sponia orientalis*, Planch." Adde: c.p. 2203.
277. "EUPHORBIACEÆ." Adde:  
22/1. CROZOPHORA, A. Juss., Baillon.  
1. *C. Rottleri*, A. Juss. Euphorb. Tent. p. 28.—C.P. 3854.  
HAB. Nillevelly, near Trincomalie, *Rev. S. O. Glenie*.
291. "TRICHOPODIUM, Lindl."  
Hoc genus certe, ut clar. Klotzsch indicavit, melius DIOSCOBIACEIS adsociandum est.
335. "*Amorphophallus Zeylanicus*, Blume." Adde ad descript.:  
*A. (?) sylvaticus*, Kunth, Enum. iii. p. 34. *Brachyspatha sylvatica*, Schott, Syn. Aroid. p. 35. *Arum sylvaticum*, Roxb. (Wight, Ic. t. 812.)—C.P. 3733.  
HAB. Nillevelly, near Trincomalie, *Rev. S. O. Glenie*.
- 339 "LILIACEÆ." Adde:  
7/2. UROPETALUM, Gawl.  
1. *U. montanum*, Dalzell in Hook. Kew Journ. of Bot. v. 2 (1850), p. 142.—C.P. 3856.  
HAB. Lakkaigalla, Matelle East, *Mr. Thos. W. N. Beckett*.
355. "*Carex Walkeri*, Arn." Dele "not" ante "very common."
358. "PANICUM, Linn." Adde:  
35/2. *P. sordidum*, Thw.; fluitans, culmis elongatis, ramosis, ad nodos pilosis; foliis lanceolatis, reflexis, quam vagina glabra brevioribus, superne pilosulis, subtus glabris, margine scabris; panicula racemiformi, ramis parum confertis, suberectis, singulis spiculam unicam gerentibus et in setam scabram spicula longio-

Page

rem productis; *spicula* lanceolata, acuminata, brevipedicellata; *gluma inferiore* brevi, truncatula vel emarginata, subnervia, pallida; *superiore* multoties longiore, lanceolata, hispida, circiter 9-nervia; *flore masculo* glabro quam *gluma inferior* parum brevior; *palea inferiore* sub-7-nervia, quam *flos fœmineus* membranaceus ovato-lanceolatus brevistipitatus duplo longiore.—*Chamæruphis depauperata*, N. ab Es., Wight's List, n. 1654 (haud *P. depauperatum*, Muhl.).—C.P. 3857.

HAB. Colombo, *Mr. W. Ferguson*.

*Folia* pollicaria, circiter 1 lin. lata. *Panicula* 1-1½ poll. longa. *Spiculæ* 2 lin. longæ, pallide virides.

*P. aspero*, Wight, affinis, sed species satis distincta.

359. Pro "13. **P. cruciforme**," lege: "**P. eruciforme**."

364. "**Ischæmum barbatum**, Retz." Adde ut synonyma: *Meoschium monostachys*, W. et A., Steud. Synops. Gram. p. 369.

369. "**Dimeria ornithopoda**, Trin., var.  $\delta$ ." Adde ut synonyma: *Pterygostachium Lehmanni*, N. ab Es., Steud. Syn. Gram. p. 413.

370. "**Polypogon Monspeliensis**, Desf." Adde ut synonyma: *P. Zeylanicus*, N. ab Es. Syn. Gram. p. 183.

372. "**Gymnopogon rigidus**, Thw.," est *Dichætaria Wightii*, N. ab Es., Steud. Syn. Gram. p. 145.

372. "**Megalachne Zeylanica**, Thw.," est *Eriachne triseta*, N. ab Es., Steud. Syn. Gram. p. 237.

375. "**ARUNDINARIA**, Rich." Adde:

3. **A. Wightiana**, N. ab Es. (Steud. Syn. Gram. p. 335.)—C.P. 3860.

HAB. Pedrotalagalla, in the Central Province.

392. "**ASPIDIUM**, Sw."

"e. *Lastrea*." Adde:

31/1. **A. undulatum**, Thw.; *frondibus* tripinnatis, deltoideis, cum stipite stramineo paleaceo æquilongo bipedalibus; *rachi* geniculato-flexuosa; *pinnis primariis secundariisque* deltoideo-lanceolatis, alternis, ad racheos geniculas elegantule deflexis; *pinnulis* rhomboideo-oblongis, plus minus decurrentibus, parce crenulato-dentatis; *soris* solitariis vel paucis, a margine pinnulæ et costula æquidistantibus; *indusio* orbiculari, parum reniformi, demum reflexo nec plicato-contorto.—C.P. 3858.

HAB. Wattakelle, Kalliboka, *Mr. Thos. W. N. Beckett*.

*Aspid. sparso*, Spr. var.  $\beta$  affine, sed differt rachi flexuosa, ramis deflexis et fronde magis composita.

394. "**POLYPODIUM**, Linn."

Pro "**P. elongatum**, Wall. (Hook. l. c. p. 234)," substitue: **P. Walkeræ**, Hook. l. c. p. 233.

401. "**TRIUMFETTA**, Plum."

Pro "1. **T. pilosula**, Thw.," substitue:

3. **T. neglecta**, W. et A. (Prod. i. p. 75.)

## INDEX OF NATIVE NAMES.

	Page		Page
Aachariyapaloo . . . . .	89	Baireya-gass . . . . .	103
Ac-malla . . . . .	165	Bairoo-deyanella . . . . .	123
Ahalla-gass . . . . .	95	Bairoo-gass . . . . .	279
Ahoo-gass . . . . .	145	Bakmee-gass . . . . .	137
Aikawaireya . . . . .	191	Bala-gass . . . . .	78
Akka-paana . . . . .	129	Baloo-dang . . . . .	174
Alanga . . . . .	211	Balloonakoota . . . . .	241
Alareya-gass . . . . .	72	Bäloleyā . . . . .	86
Albadda-gass . . . . .	59	Bandoorā-wel . . . . .	291
Allandoo-gass . . . . .	263	Batta-gass . . . . .	376
Aloobo-gass . . . . .	116	Batta-domba-gass . . . . .	116
Aloo-gass . . . . .	320	Batta-kerilla-gass . . . . .	54
Amba-kaha . . . . .	316	Battoo-karawilla . . . . .	126
Ambala . . . . .	205	Bellipatta . . . . .	26
Ambaralla-gass . . . . .	78	Bim-pōl . . . . .	339
Ammoo . . . . .	358	Bin-dadakeeriya . . . . .	269
Amookkara . . . . .	217	Bindara . . . . .	203
Andanahiriya . . . . .	81	Bin-karral-habba . . . . .	249
Andara-gass . . . . .	99	Bin-kohōmba-gass . . . . .	59
Anoda . . . . .	27	Bin-nooga . . . . .	197
Appela . . . . .	31	Bin-seyembala . . . . .	96, 97
Arramana . . . . .	96	Bin-tal . . . . .	291
Aswenua . . . . .	87	Bin-tambooroo . . . . .	211
At-addeya . . . . .	161	Bo-gass . . . . .	264
Atambooroo-gass . . . . .	200	Bokaara-gass . . . . .	71
Atbamikeeriya . . . . .	320	Bōmboo-gass . . . . .	184
At-demmata . . . . .	244	Bo-mec-gass . . . . .	255
At-nairenchee . . . . .	209	Boo-ambilla-gass . . . . .	289
Attairiya . . . . .	45	Eoo-dadakeeriya . . . . .	269
Attakirilla-gass . . . . .	71	Boogatteya . . . . .	25
Atta-kirindi-wel . . . . .	201	Boo-hora-gass . . . . .	33
Attamba-gass . . . . .	75	Boo-kanda-gass . . . . .	272
Attameeriya . . . . .	213	Boo-kattoo-handa . . . . .	215
Attatora . . . . .	91	Bookinda . . . . .	12
Attawarella-gass . . . . .	59	Boolat-wel . . . . .	292
Atteekka-gass . . . . .	267	Boolo-gass . . . . .	103
Attoona . . . . .	29	Boo-nooga-gass . . . . .	265
Attora . . . . .	360	Boo-pilla . . . . .	84
Awaree . . . . .	83	Booroola-gass . . . . .	64
		Booroota-gass . . . . .	61
Babilla . . . . .	27	Boo-sairoo-gass . . . . .	243
Badoolla-gass . . . . .	76	Boo-tora . . . . .	96
Bairaleya . . . . .	35	Bootsarana . . . . .	320

	Page		Page
Bowitteya . . . . .	104, 106	Eepaatta . . . . .	133
Caahepittang . . . . .	13	Eereerajah . . . . .	313
Caaleya . . . . .	31	Eeereya-gass . . . . .	11
Cabal-mārā-gass . . . . .	100	Ella-dadakeeriya . . . . .	268
Cadjoo-gass . . . . .	75	Ella-gokatoo-gass . . . . .	50
Cadol-gass . . . . .	120	Ella-midella-gass . . . . .	119
Cahaata-gass . . . . .	119	Ella-nettol . . . . .	245
Calateya-gass . . . . .	100	Ellawel . . . . .	319
Calla-wel . . . . .	92	Ensal . . . . .	318
Caloocaara . . . . .	7	Ereeweriya . . . . .	238
Canna-gona-gass . . . . .	262	Eremoosoo . . . . .	195
Carawilla-cabilla-gass . . . . .	289	Errabadoo-gass . . . . .	89
Carrow-gass . . . . .	281	Erramineya-wel . . . . .	74
Cobbai-gass . . . . .	55	Gadoomba-gass . . . . .	267
Colong-gass . . . . .	137	Gahala . . . . .	335
Cōng-gass . . . . .	58	Gal-karanda . . . . .	97
Cora-caha . . . . .	111	Gal-kopprawalleya . . . . .	238
Daawoo-gass . . . . .	103	Gallis-gass . . . . .	159
Dada-kaha . . . . .	316	Gal-mendora-gass . . . . .	97
Dada-kaihel . . . . .	336	Gal-mora-gass . . . . .	102
Dalook-gass . . . . .	268	Gal-seyembala-gass . . . . .	97
Dang-gass . . . . .	117	Gal-weraloo-gass . . . . .	32
Dat-kateya . . . . .	9, 139	Gam-mirris-wel . . . . .	293
Dawal-kooroondoo-gass . . . . .	257	Gammāloo-gass . . . . .	92
Dawata-gass . . . . .	121	Gandapaana-gass . . . . .	43
Del-gass . . . . .	262	Gang-kolang-kola . . . . .	239
Deya-danga-gass . . . . .	206	Gang-mee . . . . .	175
Deya-eringoo . . . . .	72	Gass-dool . . . . .	261
Deya-habarella . . . . .	321	Gass-kaala . . . . .	89
Deya-kirilla . . . . .	209	Gass-kahambillya . . . . .	259
Deya-kirindi-wel . . . . .	52	Gass-kappiteya . . . . .	275
Deya-mānil . . . . .	324	Gass-karral-habba . . . . .	249
Deya-midella-gass . . . . .	119	Gass-kyla . . . . .	285
Deya-mitta . . . . .	13	Gass-nettool . . . . .	266
Deya-nā-gass . . . . .	50	Gass-nidikoomba . . . . .	99
Deyanella . . . . .	208	Gass-pinna . . . . .	243
Deyapara . . . . .	4	Gass-pitcha . . . . .	154
Deya-parandella . . . . .	331	Gatta-cola . . . . .	142
Deya-ratmal . . . . .	97	Gatta-demmata . . . . .	244
Deya-seyembala . . . . .	85	Gatta-nittool . . . . .	264
Deya-wawoolatteya . . . . .	95	Gattatoomba . . . . .	241
Diwi-addeeya . . . . .	212	Geddekillala-gass . . . . .	123
Diwi-kadooroo-gass . . . . .	192	Gendakola . . . . .	23
Diwool-gass . . . . .	48	Gherriatta-gass . . . . .	189
Dodang-paana-gass . . . . .	45	Gim-pōl . . . . .	327
Domba-keena-gass . . . . .	52	Girripatā . . . . .	321
Domba-gass . . . . .	51	Girritilla . . . . .	209
Dool . . . . .	194	Goda-kadooroo-gass . . . . .	201
Doon-gass . . . . .	35	Goda-mānil . . . . .	324
Doonoo-kaiyeya . . . . .	327	Godapara . . . . .	5
Dootoo-satootoo . . . . .	220	Goda-wawool-atteya . . . . .	95
Dorana-gass . . . . .	34	Gojabbā . . . . .	357
Dotaloo-gass . . . . .	328	Gokatoo-gass . . . . .	49
Eela-gass . . . . .	243	Goloo-bailoo . . . . .	323
		Goloo-mora-gass . . . . .	254
		Gona-gass . . . . .	180

	Page		Page
Gona-kola . . . . .	218	Imbool-gass . . . . .	28
Gonapittan-wel . . . . .	188	Indee ( <i>Phoenix sylvestris</i> ) . . . . .	329
Gona-wel . . . . .	188	Indooroo . . . . .	340
Gōng-kakiree . . . . .	127	Ingini-gass . . . . .	201
Gōng-kadooroo-gass . . . . .	201	Itta . . . . .	132
Gooranda-gass . . . . .	267		
Goraka-gass . . . . .	48	Jaawaal . . . . .	326
Goyang . . . . .	357		
Goya-wel . . . . .	340	Kaakoona-gass . . . . .	79
		Kabella-gass . . . . .	288
Haamanilla . . . . .	239	Kadoombaireya-gass . . . . .	181
Habarella . . . . .	336	Kadoopāra . . . . .	166, 167
Haddoka-gass . . . . .	274	Kadoorookettya-gass . . . . .	103
Hakambala . . . . .	128	Kaha . . . . .	316
Hakkarilla . . . . .	216	Kaha-andanaheriya . . . . .	81
Halamba-gass . . . . .	137	Kaha-kaala-gass . . . . .	179
Hal-gass . . . . .	37	Kaha-pettang . . . . .	98
Hal-mililla . . . . .	32	Kahata-kondol . . . . .	326
Hampalanda-gass . . . . .	103	Kakala-gass . . . . .	9
Hamparandella-gass . . . . .	273	Kallandooroo . . . . .	343
Hampilla . . . . .	92	Kaloo-habaraleya-gass . . . . .	183
Hampinna . . . . .	86	Kaloo-kadoombaireya-gass . . . . .	180
Hanna . . . . .	81	Kaloo-kangwaireya . . . . .	216
Haran-kaha . . . . .	316	Kaloo-midereya-gass . . . . .	180, 181
Hātāwaareya . . . . .	337	Kaloowara-gass . . . . .	180
Hawaree-maddoo . . . . .	211	Kaloowaraneya . . . . .	233
Heen-ambilla-gass . . . . .	289	Kana-goraka-gass . . . . .	49
Heen-appela . . . . .	25	Kanda-gass . . . . .	274
Heen-bintal . . . . .	324	Kandalla . . . . .	335
Heen-bowitteya . . . . .	104	Kangkoombala . . . . .	195
Heen-gendakola . . . . .	23	Kangkoombala-gass . . . . .	71
Heen-gotoocola . . . . .	130	Kangkoombala-katteya-gass . . . . .	102
Heen-himbootoo-wel . . . . .	53	Kangkoong . . . . .	211
Heen-kabarosa . . . . .	338	Kanna-kidāran . . . . .	335
Heen-moodoomahana . . . . .	164	Kāooloo . . . . .	361
Heen-maapiritta . . . . .	26	Kappooroo . . . . .	170
Heen-oondoopeyelli . . . . .	86	Kappooroo-gass . . . . .	398
Heen-talla . . . . .	236	Kappootoo-bo-gass . . . . .	265
Heen-tambala . . . . .	214	Karanda-gass . . . . .	191
Herimanadatta . . . . .	241	Karapinchee-gass . . . . .	46
Hig-gass . . . . .	78	Katakaloowa . . . . .	106
Hik-gass . . . . .	78	Katambilla . . . . .	18
Himbootoo-wel . . . . .	53	Kattakaala-gass . . . . .	279
Hingool-gass . . . . .	60	Katta-keeriya . . . . .	320
Hingooroo-wel . . . . .	99	Katteya-gass . . . . .	68
Hirritalla . . . . .	326	Kattoo-andara-gass . . . . .	90
Homedereya-gass . . . . .	181	Kattooobodde . . . . .	28
Hondala . . . . .	128	Kattoo-herileya-gass . . . . .	28
Hondapara . . . . .	5	Kattoo-keena-gass . . . . .	69
Hoolanghik-gass . . . . .	61	Kattoo-kenda-gass . . . . .	17
Hoolankeeriya . . . . .	320	Kattoo-kittool-gass . . . . .	328
Hoonoo-kirilla-gass . . . . .	285	Kattoo-kooroondoo-gass . . . . .	16, 17
Hoorakandoo-gass . . . . .	72	Kattoo-imbool-gass . . . . .	28
Hooree-mārā-gass . . . . .	100	Kattoo-oona-gass . . . . .	375
Hora-gass . . . . .	33	Kattoo-pilla . . . . .	281
		Kattoo-timbool . . . . .	263
Illook . . . . .	369	Kattoo-wel-battoo . . . . .	217

	Page		Page
Kattoo-wella . . . . .	326	Mā-banda . . . . .	209
Keekirrindee . . . . .	164, 357	Mā-bintal . . . . .	324
Keena-gass . . . . .	51	Mā-dang-gass . . . . .	116
Ken-handa . . . . .	243	Madara-gass . . . . .	280
Kerindi-wel . . . . .	80	Madateya-gass . . . . .	98
Kettula . . . . .	334	Maddeya-gass . . . . .	191
Kidāran . . . . .	335	Maddo-gass . . . . .	294
Kirri-angoona . . . . .	199	Mādeya-jāwāla . . . . .	325
Kirri-baddoo . . . . .	210	Madol-gass . . . . .	49
Kirri-handa . . . . .	247	Madooroo-taloo . . . . .	236
Kirri-kōng-gass . . . . .	61	Magoolkaranda-gass . . . . .	92
Kirri-kondōl . . . . .	326	Maha-badoolla-gass . . . . .	75
Kirri-makooloo-gass . . . . .	269	Maha-bairoo-gass . . . . .	279
Kirriwalla . . . . .	145	Maha-booloo-mora-gass . . . . .	79
Kirriwalla-gass . . . . .	194	Maha-bowitteya . . . . .	106
Kirriwarala-gass . . . . .	178	Maha-deya-dool . . . . .	260
Kirriwel . . . . .	194	Maha-doomooda . . . . .	210
Kittool-gass . . . . .	329	Maha-erramineya . . . . .	74
Kohōmba-gass . . . . .	60	Maha-gotoocola . . . . .	130
Kok-mota . . . . .	341	Maha-kabarosa . . . . .	338
Kokoon-gass . . . . .	52	Maha-kohilla . . . . .	336
Koloowala . . . . .	319	Maha-nooga-gass . . . . .	265
Koodāloo-mal . . . . .	65	Maha-oodoopeyelli . . . . .	86
Koodoo-kooroondoo-gass . . . . .	253	Maha-yakwanassa . . . . .	241
Koodoo-mirris-wel . . . . .	69	Mainairee . . . . .	360
Koombaloo-gass . . . . .	69	Makooloo-gass . . . . .	19
Koombok-gass . . . . .	104	Malabodde . . . . .	11
Koombooroo-wel . . . . .	95	Malalaboo . . . . .	292
Kooppamaineya . . . . .	271	Malkaara-gass . . . . .	70
Koorakkan . . . . .	371	Malla . . . . .	42
Koora-tampala . . . . .	248	Mallabattoo . . . . .	216
Koroondoo-gass ( <i>Cinnamomum</i> ) . . . . .	252	Mandora-gass . . . . .	37
Kopprawalleya . . . . .	238	Maoossa-gass . . . . .	259
Korasawel . . . . .	2	Marang-gass . . . . .	118
Kosbadda-gass . . . . .	254	Mā-ratmal-gass . . . . .	71
Kōs-gass ( <i>Artocarpus integrifolia</i> ) . . . . .	262	Masmoroo-gass . . . . .	98
Kota-dimboola-gass . . . . .	267	Matabimbeya-gass . . . . .	172
Kotakan-babilla . . . . .	28	Mā-waiwel . . . . .	330
Kowakka . . . . .	128	Mee-gass . . . . .	175
		Meegōng-karapinchee . . . . .	46
Lainaterri-gass . . . . .	328	Meemini-mārā . . . . .	100
Lawooloo-gass . . . . .	174	Meriya-gass . . . . .	176
Layoo . . . . .	356	Meeyan-mililla-gass . . . . .	244
Leeneya-gass . . . . .	28	Middee-gass . . . . .	242
Leeyang-gass . . . . .	79	Mililla-gass . . . . .	244
Lingoorella . . . . .	326	Mokoonoo-wanna . . . . .	250
Loloo . . . . .	214	Monera-koodimbeya . . . . .	160
Lonkenda . . . . .	316	Monera-pattang . . . . .	338
Loonoo-ankenda-gass . . . . .	69	Moodoo-bin-tambooroo . . . . .	211
Loonoo-dang-gass . . . . .	173	Moodoo-dadakeeriya . . . . .	269
Loonoo-madala-gass . . . . .	206	Moodoo-kadooroo-gass . . . . .	192
Loonoo-midella-gass . . . . .	59	Moodoo-kaiyeya . . . . .	327
Loonoo-warana . . . . .	15	Moodoo-moroonga . . . . .	94
Loonoo-weela . . . . .	218	Moonemal-gass . . . . .	175
		Moohoodoo-gattacola . . . . .	151
Maana . . . . .	367	Mooroota-gass . . . . .	122
Maassenda . . . . .	138	Moowa-keeriya . . . . .	196

	Page		Page
Mora-gass . . . . .	58	Pitcha . . . . .	189
Myla-gass . . . . .	98	Pitta-sooddoo-pala . . . . .	245
Naapiritta . . . . .	26	Pittawaaka . . . . .	282
Nadoong-gass . . . . .	93	Pöl-gass . . . . .	230
Nā-gass . . . . .	50	Pöl-hoonoopala . . . . .	248
Naha-gass . . . . .	251	Poopoola . . . . .	160
Narawella . . . . .	1	Poos-wel . . . . .	98
Natton-gass . . . . .	9	Poowak-gass . . . . .	327
Nawa-handee . . . . .	268	Pota-wel . . . . .	337
Nelli-gass . . . . .	282	Radaleya-gass . . . . .	80
Nelloo . . . . .	223	Ranawarā . . . . .	96
Neyadandoo . . . . .	316	Rankeeriya . . . . .	329
Neyadasse-gass . . . . .	41	Ranmanissa . . . . .	340
Neyanda . . . . .	338	Ranwau-keekerindi . . . . .	165
Neyangulla . . . . .	339	Rasnee . . . . .	170
Neyang-rata-coloo . . . . .	126	Rassakinda . . . . .	12
Nikka-gass . . . . .	244	Rata-boolat-wel . . . . .	292
Nil-andanahiriya . . . . .	81	Rata-ensal . . . . .	318
Nil-katarodoo . . . . .	88	Rata-gowa . . . . .	129
Nil-pitcha . . . . .	153	Ratambala . . . . .	155
Nooga-gass . . . . .	265	Rata-tora . . . . .	90
Odootalau . . . . .	239	Rat-kihiri-gass . . . . .	99
Okaiyeya . . . . .	327	Rat-nettol . . . . .	245
Okooroo-gass . . . . .	272	Ratoo-waa . . . . .	95
Olinda-wel . . . . .	91	Rat-pittawaakka . . . . .	270, 282
Oloo . . . . .	205	Rawaniddala-gass . . . . .	59
Oogoorasse-gass . . . . .	17	Rawaniddala ( <i>Wendlandia</i> ) . . . . .	159
Oolooloo-gass . . . . .	254	Ritti-gass . . . . .	264
Oona-gass . . . . .	375	Rookattana-gass . . . . .	194
Oonoonoo-gass . . . . .	103	Rook-gass . . . . .	11
Oorookannoo-gass . . . . .	43	Saarana . . . . .	23
Ooroo-tora . . . . .	95	Saayam-mool . . . . .	144
Ooyala . . . . .	326	Saipaala-gass . . . . .	190
Paawetta . . . . .	233	Samadera-gass . . . . .	70
Painaira-wel . . . . .	54	Seenook-gass . . . . .	268
Painee-bāroo . . . . .	366	Seewana-maddeya-gass . . . . .	266
Palang-gass . . . . .	72	Seyembala-gass . . . . .	95
Palenga-gass . . . . .	287	Siribo . . . . .	292
Paloo-gass . . . . .	175	Soodoo-kadoombaireya-gass . . . . .	179
Pambooroo-gass . . . . .	45	Soodoo-idda . . . . .	193
Pangaree-koorondoo-gass . . . . .	253	Soodoo-leyang-gass . . . . .	281
Panoo-alla . . . . .	335	Soodoo-nikka-gass . . . . .	244
Panoo-condōl . . . . .	326	Soodoo-pooroo-gass . . . . .	233
Panoo-kaara-gass . . . . .	117	Soodoo-toomba . . . . .	240
Pat-kaala-gass . . . . .	280	Sooreya-gass . . . . .	27
Patook-gass . . . . .	268	Soovanda-gass . . . . .	50
Pat-paadagan . . . . .	24	Taccāda . . . . .	169
Patta-appelle . . . . .	25	Taccada-gass . . . . .	189
Pawetta . . . . .	156	Taiboo-gass . . . . .	320
Pehimbia-gass . . . . .	59	Tal-gass . . . . .	329
Pehilla . . . . .	133	Talla-gass . . . . .	329
Penella-gass . . . . .	55	Tambaleya-gass . . . . .	114
Pennit-tora . . . . .	95	Tammanna-gass . . . . .	428
Pilla . . . . .	84	Tampala . . . . .	247



	Page		Page
Tanna-patā . . . . .	322	Wal-kaihel-gass . . . . .	321
Tarrana-gass . . . . .	158	Wal-kakoona-gass . . . . .	278
Telemboo . . . . .	29	Wal-karapinchee-gass . . . . .	47
Tellakeeriya-gass . . . . .	269	Wal-kidāran . . . . .	335
Tel-talla . . . . .	209	Wal-kolondoo . . . . .	165
Tetti-gass . . . . .	19	Walla-gass . . . . .	251
Tetti-wel . . . . .	12	Wallangereya . . . . .	47
Tibbootoo . . . . .	217	Walloo-keena-gass . . . . .	51
Timberee-gass . . . . .	178	Wal-loonoo . . . . .	324
Tinneya-gass . . . . .	35	Wal-maa . . . . .	89
Tippili . . . . .	292	Wal-mooroonga . . . . .	285
Titta-hondala . . . . .	127	Wal-patpaadagam . . . . .	144
Tolabo . . . . .	324	Wal-rasakinda . . . . .	124
Toomba-karawilla . . . . .	126	Wal-sappoo-gass . . . . .	5
Toompat-kooroondoo . . . . .	45	Wall-teekola . . . . .	165
Totilla-gass . . . . .	206	Wal-tibbootoo . . . . .	217
Trasta-waaloo . . . . .	212	Wal-wareka . . . . .	19
		Wanna-rajah . . . . .	314
Vatta-coloo . . . . .	126	Waranneya . . . . .	142
Velenga-gass . . . . .	30	Warra . . . . .	196
Visnoo-kraanta . . . . .	213	Wayoopaloo . . . . .	1
		Weela . . . . .	219
Waa-gass . . . . .	96	Weera-gass . . . . .	287
Waaweya . . . . .	12	Wel-ambilla . . . . .	172
Wadakaha . . . . .	337	Wel-kahambilleya . . . . .	270
Waila . . . . .	14	Wel-kappiteya . . . . .	275
Wai-wel . . . . .	330	Wel-kyla . . . . .	282
Wall-ahalla-gass . . . . .	96	Wellapeyenna-gass . . . . .	119
Wall-ahattoo . . . . .	266	Welle-wenne-gass . . . . .	278
Wal-asamodagan . . . . .	130	Wellicaha . . . . .	110
Wal-awara . . . . .	88	Weniwel . . . . .	12
Wal-biling-gass . . . . .	69	Weraloo-gass . . . . .	32
Wal-eōlloo . . . . .	91	Wissak-mal . . . . .	297
Wal-copee . . . . .	157		
Wal-gammirris-wel . . . . .	293	Yakbaireya . . . . .	81, 82
Wal-gona-gass . . . . .	266	Yak-errabadoo-gass . . . . .	89
Wal-gooranda . . . . .	243	Yak-komadoo . . . . .	126
Wall-ingoоро . . . . .	315	Yakwanassa . . . . .	240
Wal-jambo-gass . . . . .	115	Yuccahaaloo-gass . . . . .	35
Wal-kaha . . . . .	316	Yucca-naara-gass . . . . .	46
Wal-kahambilleya . . . . .	259		

## INDEX OF C.P. NUMBERS.

C. P.	Page	C. P.	Page	C. P.	Page	C. P.	Page
1	375	44	161	87	143	130	373
2	155	45	177	88	143	131	341
3	346	46	177	89	235	132	166, 422
4	131	47	177	90	237	133	150
5	282	48	48	91	141	134	101
6	64, 409	49	65	92	145, 420	135	131
7	22, 400	50	267	93	141	136	100
8	33	51	152	94	141	137	204
9	4	52	105	95	143, 419	138	106
10	255	53	251	96	141	139	142
11	114	54	282	97	142	140	65
12	81	55	91	98	143, 419	141	5
13	140	56	118	99	141	142	234
14	265	57	146, 420	100	359	143	215
15	174	58	85	101	141	144	370
16	237	59	364	102	140	145	131
17	238	60	373	103	141	146	123
18	147	61	341	104	142	147	163
19	163	62=44	161	105	143	148	72
20	160	63	107	106	143	149	170
21	162	64	17, 400	107	142	150	160
22=579	73	65	370	108	142	151	167
23	186	66	229	109	141	152	162
24	369	67	27	110	140	153	20
25	54	68	286	111	141	154	239
26	239	69	368	112	142	155	186
27	161	70	373	113	142	156	65
28	174	71	283	114=88	143	157	153
29	264	72	257	115	141	158	148
30	288	73	136	116	142	159	12
31	65	74	256	117	143	160	118
32	367	75	338	118	152	161	12
33	239	76	20	119	141	162	228
34	165	77	114	120	141	163	77
35	293	78	186	121	143	164	160
36	174	79	255	122	143	165	184
37	252	80	146, 420	123	141	166	33
38	203	81	364	124=88	143	167	256
39	136	82	150	125	144	168	13
40	255	83	252	126	141	169	13, 160
41	351	84	237	127	187	170	112
42	170	85	231	128	244	171	73
43	71	86	141	129	104	172	33

C. P.	Page	C. P.	Page	C. P.	Page	C. P.	Page
173	108	230	255	287	111	313	239
174	185	231 = 163	77	288	202	344	150
175	12	232	4	289	146, 420	345 = 265	2
176	370	233	266	290	146, 420	346	202
177	59	234	164	291	117	347	138
178	282	235	175	292 = 42	170	348	145, 40
179	177	236	360	293	59	349	109
180	45, 406	237	332	294	161	350	116
181	17, 400	238		295	136	351	255
182	130	239	2	296	282	352	207
183	195	240	224	297 = 148	72	353	109
184	22	241	376	298	184	354	146
185	158	242	52	299	5	355	149
186	232	243	94	300	59	356	257
187	201	244	6	301	235, 229	357	146, 420
188	46	245	159	302	240	358	208
189	251	246	78	303 = 3789.		359	139, 419
190	154	247	351, 435	304	256	360	255
191	177	248	4	305		361	145, 420
192	177	249	236	306	351	362	138
193	243	250	203			363	202
194	10	251	44, 405	307 = 23	186	364	224
195 = 185	158	252	274	308	338	365	114
196	273	253	3	309	227	366	327
197	126	254	228	310	230	367	82
198	134	255	231	311 = 266	229	368	185
199	352	256	92	312	371	369	197, 424
200	293	257	234	313	77	370 = 358	208
201	74	258	286	314	362	371	39, 404
202	71	259	28	315	159	372	67
203	290	260	22, 400	316	32	373	7
204	42	261	150	317	252	374	71
205	139	262	290	318 = 34	165	375	44, 405
206	251	263	252	319 = 42	170	376	174
207	310	264	237	320	283	377	341
208	105	265	2	321	256	378	341
209	91	266	229	322	49	379	266
210	152	267	156	323	80	380	118
211	17, 400	268	139	324	98	381	133
212	241	269	82	325	243	382	181
213 = 152	162	270	170	326		383	312
214 = 203	290	271	187	327	282	384	314
215	139	272	167	328	97	385	107
216	161	273	185	329	71	386	69
217	239	274	118	330 = 2516.		387	111
218	71	275	188	331	267	388	79, 410
219	346	276	108	332	67	389	161
220	341	277	172	333	149	390	149
221	11	278	171	334	138	391	150
222	54	279	92	335	49	392	253
223	255	280	81	336	136	393	253
224	91	281	374	337	100	294	177
225	131	282	361	338	5	395	207
226	310	283	239	339	4	396	270
227	308	284	104	340	146, 420	397 = 9	4
228	234	285	173	341	201	398	225
229	221	286	138	342	285	399	235

C.P.	Page	C.P.	Page	C.P.	Page	C.P.	Page
400	24	456	259	513	162	570	94
401	365	457	202	514	309	571	
402	71	458	156	515	131	572	68
403	163	459	167	516=203	290	573	183
404	234	460=164	160	517	22	574	297
405	60	461	237	518	300	575	133
406	155	462=156	65	519	264	576	5
407	186	463	115	520	1	577	276
408=203	290	464	64, 409	521	170	578	277
409	72	465	20	522	308	579	73
410=15	174	466	142	523	310	580	59
411	367, 437	467	291	524	84, 85	581	167
412	136	468	18	525	220	582	184
413	258	469	250	526	82	583	237
414	254	470	118	527	239	584	139, 419
415	19	471	78	528	166, 422	585	148
416	11	472	189	529	170	586	148
417	255	473	184	530	131	587	255
418	115	474	135	531	170	588	118
419	366	475	188	532=19	163	589	73
420	166	476	68	533	167	590	333
421	33	477	183	534=203	290	591	225
422	146, 152, 420	478	73	535	218	592	22, 400
423	146, 420	479	136	536	59	593	296
424	5	480	136	537=335	49	594	162
425	237	481	136	538	69, 409	595	138
426	107	482=216	161	539	32	596	279
427	383	483	265	540	129	597	313
428	156	484	139	541	129	598	314
429	376	485	301	542	207	599=598	314
430	167	486	107	543	296	600	105
431	140	487	160	544	202	601	115
432	283	488	304	545	28	602	50
433	150	489	304	546	335	603	50, 407
434	22	490	68	547	65	604	404
435	167	491	290	548	170	605	56
436	67	492	43	549	132	606	232
437	9	493	54	550	312	607	56
438	187	494=475	188	551	163	608	
439	93, 414	495	42	552	236	609	115
440	202	496	60	553	323	610	173
441	104	497	44	554	32	611	213
442	149	498	242	555	184	612	320
443	57	499	210	556=335	49	613	300
444	167	500=189	251	557	275	614	15
445	114	501	282	558	60	615	9
446	3	502	251	559	132	616=491	290
447	123	503	65	560	163	617	5
448	155	504	184	561	158	618	103
449 (499 in err.)		505	152	562	55, 408	619	176
	139, 418	506	105	563	71	620	328
450	373	507	106	564		621	32
451	123	508	131	565	167, 142	622	184
452	118	509	234	566	160	623	173
453	67	510	283	567	161	624	32
454	363	511=150	160	568	166, 422	625=42	170
455	65	512	203	569	71	626	185

C.P.	Page	C.P.	Page	C.P.	Page	C.P.	Page
627	188	684	165	740	174	799	343
628	252	685	68	741	173	800	342
629	17, 400	686	235	742	156	801	342
630	32	687	237	743	50	802	342
631	76, 410	688	167	744	99	803	343
632	299	689=775	41	745=658	53	804	343
633	304	690	250	746	159	805	343
634	117	691	164	747	243	806	342
635	77	692=27	161	748	155	807	342
636=201	74	693=694	3	749	80	808	343
637	42	694	3	750	140	809	344
638	102	695	138	752	305	810	344
639=335	49	696 (695 in err.)	274	753	158	811	344
640=529	170			754	308	812	342
641	76, 410	697	92	755	50	813	344
642	32	698	244	756	194	814	345
643	170	699	59	757	200	815	344
644	100	700	364	758	416	816	345
645	370	701	233	759	156	817	345
646	103	702	155	660	133	818	345
647	184	703	59	761	94	819	347
648=491	290	704	33	762	44, 405	820	355
649=557	275	705	391	763	117	821	355
650	283	706=690	250	764	111	822	355
651	272	707	139, 418	765	144	823	349
652	219	708	235	766	289	824	355
653	97	709	359	767	158	825	353
654	270	710	359	768=388	79	826	354
655	132	711	140	769=748	155	827	354
656	272	712	140	771	156	828	353
657	159	713	233, 234	772	94	829	350
658	53	714	391	773	289	830	348
659	53	715	22, 400	774	199	831	351
660	289	716	268	775	41, 405	832	348
661	88	717	196	776	41, 405	833	348
662=349	109	718=913	359	777	41	834	350
663	88	719=334	138	778	40, 405	835	350
664	153	720	90	779	40, 405	836	349
665	184	721=368	185	780	137	837	349
666	300	722	243	781=1658	137	838	349
667	300	723	218	782	40, 405	839	348
668	350	724	190	783	41, 405	840	348
669	148	725	67	784	41	841	348
670	167	726	27	785	41	842	349
671=167	256	727	26	786	41	843	349
672=638	102	728	154	787	41	844	340
673	185	729	255	788	40	845	353
674	257	730	31	789	341	846	350
675	33	731	50	790	341	847	351
676	185	732	64	791	341	848	348
677	352	733	114	792	341	849=41	351
678	343	734	28	793	341	850	351
679	347, 434	735	117	794	341	851	350
680	1	736	56	795	341	852	347, 434
681	166	737	337	796	341	853	352, 434
682	232	738	199	797	344	854	350
683	163	739	155	798	342	855	344

C. P.	Page	C. P.	Page	C. P.	Page	C. P.	Page
856	358	913	359	973	391	1030	6
857	358	914	372	974	391	1031	6
858	358	915	370	975	392	1032	6
859	358	916	372	976	382	1033	8
860	358	917	366	977	381	1034	10, 398
861	358	918	360	978	387	1035	10, 398
862	358	919	362	979=1314	380	1036	10
863	358	920	374	980	379	1037	9
864	359	921	374	981 (281)	381	1038	9
865	358	922	374	982	388	1039	11
866	368	923	374	983	389	1040	7
867	364	924	376	984	377	1041	10, 398
868	364	925	373	985	377	1042	9
869	364	926	373	986	377	1043	9
870	364	927	373	987	377	1044	9
871	368	928	373	988	395	1045	6
872	363	929	373	989	380	1046	278
873	364	930	373	990	391	1047	386
874	365	931	373	991	393	1048	387
875	365	932=925	373	992	391	1049	13
876	374	933	373, 371	993	396	1050	13
877	356	934	372	994	364	1051	12
878	357	935	374	995	368	1052	13
879	362	936	366	996	366	1053	12
880	361	937	371	997	340	1054	13
881	362	939	371	998=997	340	1055	12
882	360	940	366	999	340	1056	12
883	360	941	371	1000=796	341	1057=1050	13
884	373	942	357	1001	341	1058	15
885	361	943	363	1002=61	341	1059	15
886	359	944	363	1003	340	1060	15
887	359	945	363	1004		1061	53
888	359	947	362	1005	384	1062	15
889	359	948	370	1006	383	1063	16
890	360	949	368	1007	383	1064	15
891	359	950	369	1008	384	1065	15
892	360	951	367, 437	1009	1	1066	15
893	360	952	369	1010	383	1067	14
894	360	953	367	1011	1	1068	14
895	359	954	371	1012	1	1069	16
896	357	955	369	1013	4	1070	15
897	359	956	369	1014	2	1071	16
898	359	957	369	1015	2	1072	15
899		958	371	1016	2	1073	14
900	359	959	368	1017	14, 399	1074	264
901	359	960	362	1018=1019	14	1075	18, 400
902	359	961	366	1019	14	1076	17, 400
903	359	962	366	1020	14	1077	17, 400
904	359	963	366	1021	14	1078	41
905	359	964	360	1022	14	1079	22
906	361	965	343	1023		1080	22
907	361	966	343	1024	290	1081	22
908	361	967=241	376	1025	275	1082	22
909	361	968	369	1026	12	1083	22, 400
910	361	970	349	1027	12	1084	21
911	358	971	360	1028	5	1085	21
912	361	972	357	1029	7	1086	22

C. P.	Page	C. P.	Page	C. P.	Page	C. P.	Page
1087	20	1144	29	1202	47	1260	78, 411
1088	21	1146	30	1203	46, 406	1261	78
1089	21	1147	41	1204	47	1262	95
1090	25	1148	60	1205	48	1263	78
1091	25	1149	79	1207	67	1264	81
1092	24	1150	57, 60	1208	67	1265	81
1093	250	1151	55	1209	67	1266	81
1094	24	1152	58, 408	1210	68	1267 = 1265	81
1095	24	1153	58, 408	1211	66	1268	81
1096	24	1154	58, 408	1212	66	1269	82
1097	24	1155	58	1213	65	1270	81
1098	23	1156	55	1214	42	1271	81
1099	245	1157	27	1215	42	1272	82
1100	23	1158	59	1216	42	1273	81
1101	24	1159	59	1217	20	1274	81
1102	23	1160	61	1218	42	1275	81
1103	23	1161	78	1219	42	1276	82
1104	23	1162	61	1220	103	1277	82
1105	31	1163	61	1221	71	1278	82
1106	31	1164	52	1222	70	1279	86
1107	81	1165	53	1223	70	1280	91
1108	32	1166	53	1224	70	1281	381
1109	32	1167	53	1225	72	1282	394
1110	31	1168	53	1226 = 43	71	1283	393
1111	31	1169 = 25	54	1227	73	1284	394
1112	31	1170	51	1228	72	1285	394
1113	31	1171	51	1229 = 579	73	1286	393
1114	29	1172	48	1230	67	1287	391
1115	32	1173	50	1231	79	1288	394
1116	121	1174	62	1232	72	1289	394
1117	26	1175	63	1233	75	1290	394
1118	26	1176	63	1234	158	1291	392
1119	26	1177	62	1235	72, 409	1292	382
1120	26	1178	63	1236	74	1293	382
1121	27	1179	63	1237	42	1294	396
1122	26	1180	62	1238	75	1295	395
1123	28	1181	62	1239	75	1296	395
1124	28	1182	62	1240	74	1297	395
1125	26	1183	62	1241	74	1298	395
1126	26	1184	63	1242	74	1299	390
1127	30	1185	25	1243	74	1300	390
1128	30	1186	27	1244	23	1301	395
1129	25	1187	25	1245	79	1302	382
1130	28	1188	64	1246	20	1303	381
1131	28	1189	64	1247	20	1304	381
1132	27	1190	64, 409	1248	19	1305	382
1133	27	1191	45	1249	69, 409	1306	} = 1305 382
1134	27	1192	45, 406	1250	69	1307	
1135	27	1193	61, 409	1251	69	1308	
1136	30	1194	45, 405	1252	69	1309	382
1137	28	1195	47, 48	1253 = 1252	69	1310	380
1138	28	1196	46, 405	1254	23, 401	1311	380
1139	278	1197	47, 406	1255	78	1312	380
1140	29	1198	44	1256	75	1313	380
1141	28	1199	45	1257	76	1314	380
1142	29	1200	45	1258	76	1315	380
1143	28	1201	47	1259 = 1257	76	1316	381

C. P.	Page	C. P.	Page	C. P.	Page	C. P.	Page
1317	381	1374	393	1430	87	1487	94
1318	381	1375	392	1431	86	1488=1487	94
1319	381	1376	393	1432	87	1489	92
1320	387	1377	393	1433	87	1490	92
1321	387	1378	393	1434	87	1491	93
1322	387	1379	388	1435	86	1492	.93, 413
1323	387	1380	388	1436	86	1493	.92, 413
1324	387	1381	388	1437	85	1494	92
1325	387	1382	388	1438	85	1495	92
1326	387	1383	388	1439	91	1496	94
1327	387	1384	389	1440	91	1497	98
1328	386	1385	389	1441	91	1498	98
1329	386	1386	389	1442	91	1500	97
1330	386	1387	389	1443	84	1501	95
1331=1048	387	1388	389	1444	84	1502	95
1332	386	1389	388	1445	84	1503	96
1333	381	1390	397	1446	84	1504	96
1334	383	1391	397	1447	84	1505	95
1335	385	1392 } -1390	397	1448=1447	84	1506	96
1336	383	1393 }	397	1449	84	1507	95
1337	383	1394	397	1450	86	1508	96
1338	383	1395	397	1451	84	1509	96
1339	384	1396	397	1452	82	1510	96
1340	384	1397	389	1453	83	1512	84
1341	384	1398	396	1454	83	1513	84
1342	386	1399	396	1455	83	1514	85
1343	385	1400	396	1456	83	1515	85
1344	385	1401	396	1457	83	1516	99
1345	385	1402=1398	396	1458	83	1517	100
1346	385	1403	396	1459	83	1518	100
1347	385	1404	379	1460	83	1519	99
1348	384	1405	379	1461	83	1520	99
1349	385	1406	379	1462	83	1521	99
1350	385	1407	379	1463	83	1522	
1351	386	1408	378	1464	89	1523	99
1352	385	1409	378	1465	89	1524	94
1353	385	1410	378	1466	91	1525	95
1354	390	1411	378	1467	91	1526	96
1355	390	1412	377	1468	88	1527	95
1356	390	1413=1412	377	1469	90	1529	100
1357	390	1414	377	1470	90	1530	100
1358	390	1415 (1115 in error)	377	1471	88	1531	100
1359	391	1416	377	1472	90	1532	102
1360=1359	391	1417	377	1473	.89, 412	1533	101
1361	392	1418	377	1474	90	1534	101
1362	392	1419	377	1475	90	1535	101
1363	391	1420	378	1476	90	1536	101
1364	392	1421	378	1477	88	1537	101
1365	393	1422	378	1478	90	1538=134	101
1366	389	1423	92	1479	89	1539	123
1367	390	1424	91	1480	89	1540	123
1368	392	1425	86	1481	88	1541	417
1369	392	1426	87	1482	90	1542	121
1370	392	1427	87	1483	89	1543	23
1371=1340	384	1428	87	1484	88	1544	121
1372	384	1429	86	1485	88	1545	123
1373	390			1486	84	1546	122



C. P.	Page	C. P.	Page	C. P.	Page	C. P.	Page
1547	122	1604	103	1661	138	1718	147
1548	122	1605	103	1662	152	1719	147
1549	123	1606	126	1663	155	1720	148
1550=1102	23	1607	127	1664	138	1721	57
1551	122	1608	127	1665	157	1722=1155	58
1552	122	1609	128	1666	153	1723	145, 420
1553	122	1610	124	1667	152	1724=354	146
1554	122	1611	125	1668	145	1725	146, 420
1555	112	1612=197	126	1669	145	1726	146, 420
1556	111	1613	125	1670	208	1727	146, 420
1557=1561	111	1614=197	126	1671	139	1728	146, 420
1558	110	1615	126	1672	151	1729	160
1559	110	1616	125	1673	144	1730	163
1560	111	1617	126	1674	144	1731	163
1561	111	1618	127	1675=1674		1732	163
1562	111	1619	125	1676	144	1733	163
1563	110	1620	126	1677	200	1734	163
1564	110	1621	128	1678	144	1735	163
1565	110	1622	125	1679	144	1736	160
1566	110	1623	417	1680=1679		1737	160
1567	110	1624	126	1681	144	1738	160
1568	105	1625	127	1682	151	1739	163
1569	105	1626	127	1683	144	1740	162
1570	105	1627	128	1684	151	1741	160
1571	105	1628	124	1685	152	1742	160
1572	105	1629	127	1686	130	1743	168
1573	105	1630	18	1687	130, 418	1744	163
1574	106	1631=218	71	1688	152	1745	161
1575	104	1632	132	1689	152	1746	168
1576	104	1633	132	1690	152	1747	168
1577	109	1634	130	1691	151, 152	1748	168
1578	106	1635	130	1692	142	1749	167
1579	118	1636	136	1693=113	142	1750	166
1580	117	1637	136	1694	142	1751	167
1581	114	1638	136	1695	151	1752	164
1582	117	1639	136	1696	151	1753	164
1583	117	1640	135	1697	140	1754	164
1584	116	1641	135	1698	143	1755	165
1585=274	118	1642	135	1699	143	1756	165
1586	114	1643	133	1700=91	141	1757	164
1587	118	1644	133	1701	141	1758	164
1588	114	1645	134	1702	141	1759	165
1589	113	1646	134	1703	235	1760	165
1590	114	1647		1704	139	1761	166
1591	114	1648=198	134	1705	150	1762	166
1592	119	1649	158	1706	140	1763	166
1593	119	1650	17	1707	140	1764	166
1594	103	1651	158	1708	140	1765	166
1595	123	1652	159	1709=728	154	1766	162
1596	103	1653	152	1710	150	1767	163
1597	133	1654	154	1711	150	1768	162
1598	133	1655	159	1712	148	1769	162
1599	133	1656	137	1713	149	1770	162
1600	188	1657	137	1714	150	1771	164
1601	103	1658	137	1715	155	1772	161
1602	103	1659	137	1716	149	1773	170
1603	104	1660	137	1717	156	1774	169

C. P.	Page	C. P.	Page	C. P.	Page	C. P.	Page
1775	169	1832	192	1888	216	1947	241
1776	169	1833	192	1889	215	1948	243
1777	169	1834 (1836 in er.)	191	1890	215	1949	243
1778	221			1891	215	1950	243
1779	209	1835	191	1892	215	1951	243
1780	222	1836 (1834 in er.)	191	1893	215	1952	244
1781	222			1894	215	1953	242
1782	206	1837	193	1895	172	1954	214
1783	207	1839	193	1896	172	1955	244
1784	207	1840	193	1897	217	1956	244
1785	207	1841	196	1898	217	1957	244
1786	208	1842	199	1900	216	1958	244
1787	208	1843	194	1901	216	1959	206
1788	208	1844	197	1902	216	1960	206
1789	207	1845 = 1844.		1903	216	1961	244
1790	170	1846	198	1904	217	1962	244
1791 = 1260	78	1847	198	1905	217	1963 (1763 in er.)	121
1792	403	1848	198	1906	217		
1793	175	1849	197	1907	217	1964 (1764 in er.)	121
1794	175	1850	194	1908	181		
1795	175	1851	195	1909	178	1965	120
1796	174	1852	196	1910	178	1966	120
1797	175	1853	196	1911	179	1967	120
1798 = 1797.		1854	197	1912	180	1968	120
1799	172	1855	199	1913	180	1969	120
1800	172	1856	198	1914	180	1970	212
1801 = 1799.		1857	197	1915	178	1971	231
1802	173	1858	197	1916	183	1972	235
1803	174	1859	198	1917	183	1973	235
1804	173	1860	197	1918	37	1974	235
1805	174	1861	194	1919	35	1975	234
1806	173	1862	193	1920	38, 403	1976	234
1807	190	1863	194	1921	33	1977	232
1808	190	1864 = 1865.		1922	213	1978	235
1809	190	1865	195	1923	258	1979	235
1810	190	1866	201, 425	1924	210	1980	229
1811	190	1867	201	1925	213	1981	232
1812	189	1868	113	1926	213	1982	234
1813 = 1807.		1869	206	1927 = 1926.		1983	231
1814	190	1870	205	1928	212	1984	234
1815	181	1871	204	1929	211	1985	231
1816 = 1815.		1872 = 1871.		1930	211	1986	231
1817 = 275	188	1873	204	1931	212	1987	225
1818	188	1874	204, 442	1932	210	1988 (1998 in er.)	236
1819	185	1875	205	1933	210		
1820	185	1876	203	1934	211	1989	236
1821	188	1877	203	1935	212	1990	427
1822	191	1878	203	1936 = 1939.		1991	233
1823	191	1879	214	1937	212	1992	234
1824	442	1880	214	1938	210	1993	233
1825	193	1881	213	1939	211	1994	225
1826	200	1882	214	1940	212	1995	234
1827	192	1883	209	1942	212	1996	224
1828	251	1884	209	1943	209	1997	225
1829	191	1885	214	1944	209	1998	227
1830	196	1886	215	1945	210	1999	227
1831	196	1887	216	1946	209	2000	229

C. P.	Page	C. P.	Page	C. P.	Page	C. P.	Page
2001	226	2058	239	2116	275	2173	292
2002	226	2059	237	2117	275	2175	293
2003	230	2060	237	2118	272	2176	292
2004	231	2061	237	2119=491	290	2177	293
2005	228	2062	236	2120	287	2178	293
2006	226	2063	237	2121	287	2179	259
2007	226	2064	237	2122	287	2180	260
2008	228	2065	238	2123=2122.		2181	260
2009	227	2066	238	2124	269	2182	260
2010	228	2067	238	2125	268	2183	260
2011	230	2069	238	2126	269	2184	259
2012	227	2070	241	2127=2125.		2185	259
2013	230	2071	240	2128	269	2186	259
2014	231	2072	240	2129	269	2187	261
2015	231	2073	239	2130	284	2188	260
2016	231	2074	240	2131	269, 427	2189	260
2017	232	2075	240	2132	272	2190	261
2018	232	2076	241	2133	282	2191	260
2019	232	2077	239	2134	282	2192	261
2020	224	2078	237	2135	284	2193	259
2021	217	2079	237	2136	282	2194	261
2022	220	2080	236	2137	282	2195	261
2023	209	2081	237	2138	282	2196	260
2024	219	2082	239	2139	282	2197	260
2025	220	2083	238	2140	282	2198	260
2026	220	2084	423	2141	285	2199	261
2027	220	2085	172	2142	282	2200	259
2028	220	2086	171	2143	285	2201	262
2029	220	2087	171	2144	282	2202	258
2030	220	2088	171	2145	283	2203	443
2031	220	2089	171	2146	284	2204	185
2032	219	2090	171	2147	283	2205	119
2033	219	2091	171	2148	283	2206	250
2034	426	2092	271	2149	285	2207	293
2035	218	2093	259	2150	286	2208	289
2036	218	2094	271	2151	280, 428	2209=2208.	
2037	218	2095	271	2152	288	2210	289
2038	218	2096	271	2153	288	2211	287
2039	218	2097	271	2154	281	2212	263
2040	218	2098	271	2155	281	2213	264
2041	218	2099	270	2156	279	2214	264
2042	219	2100	270	2157=557	275	2215	263
2043	219	2101	274	2158	278	2216	191
2044	219	2102	271	2159=2158.		2217	266
2045=2044		2103	273	2160	278	2218	265
2046	218	2104	272	2161	279	2219	266
2047	219	2105	273	2162=2161.		2220	265
2048	219	2106	273	2163	280	2221	265
2049	218	2107=2105.		2164	280, 428	2222	265
2050	220	2108	273	2165=2151.		2223	265
2051	220	2109	273	2166	280	2224	266
2052	219	2110	276	2167	278	2225	265
2053	240	2111	270	2168	277	2226	265
2054	218	2112	273	2169	269	2227	266
2055	238	2113	276	2170	270	2228	266
2056	237	2114	276	2171	274	2229	266
2057	240	2115	272	2172	292	2230	267

C. P.	Page	C. P.	Page	C. P.	Page	C. P.	Page
2231	263	2289	324	2346	303	2405	13
2232	262	2290	323	2347	302	2406	8
2233	427	2291	339	2348	306	2407	280
2234	249	2292	339	2349	300	2408 = 528	166
2235	248	2293	338	2350	298	2409	82
2236	249	2294	340	2351	296	2410	95
2237	249	2295	339	2352	298	2411	177
2238	247	2296	332	2353	298	2412	71
2239	248	2297	338	2354	307, 430	2413	69
2240	247	2298	337	2355	299	2214	134
2241	249	2299	337	2356	299	2415	16
2242	249	2300	337	2357	298	2416	4
2243	249	2301	340	2358	307	2417	40
2244	247	2302	338	2359	296	2418	64, 409
2245	247	2303	326	2360	308	2419	64, 409
2246	245	2304	326	2361	303	2420	48
2248	246	2305	331	2362	299	2421	46, 406
2249	246	2306	333	2363	311	2422	233
2250	246	2307	333	2364	301	2423	34
2251	246	2308	333	2365	307	2424	287
2252	246	2309	331	2366	308	2425	66
2253	248	2310	331	2367	308	2426	66
2254	248	2311	290	2368	300	2427	66
2255	248	2312	325	2369	302	2428 = 2426.	
2256	291	2313	325	2370	302	2429 = 1900	216
2257	291	2314	321	2371	308	2430	38, 403
2258	291	2315	321	2372	309	2431	318
2259	222	2316	332	2373	310	2432	337
2260	246	2317	332	2374	310	2433	323
2261	246	2318	332	2375	296	2434	358
2262	246	2319	338	2376	295	2435	186
2263	252	2320	325	2377	313	2436	184
2264	252	2321	336	2378	331	2437	180
2265	244	2322	336	2379	331	2438	179
2266	245	2323	336	2380	333	2439	180
2267	190	2324	322	2381	267	2440	42
2268 = 1923	258	2325	322	2383	25	2441	42
2269	254	2326	322	2384	143	2442	135
2270	254	2327	323	2386	364	2443	80, 410
2271	254	2328	322	2387	113	2444	76, 410
2272 = 2269.		2329	322	2388 = 1161	78	2445	49
2273	427	2330	323	2389	218	2446	51
2274	255	2331	323	2390 = 1586	114	2447	52
2275	254	2332	323	2391	136	2448	60
2276	255	2333	327	2392	254	2449	117
2277	257	2334	330, 431	2393	372	2450	116
2278	257	2335	330	2394	370	2451	114
2279 = 413	258	2336	329	2395	355	2452	117
2280	257	2337	328	2396	352	2453	117
2281	257	2338	324	2397	351, 434	2454	61
2282	252	2339	324	2398 = 1921.		2455	155
2283	252	2340	304	2399	80	2456	155
2284	252	2341	304	2400	204	2457	142
2285 = 2284.		2342	303	2401	239	2458	154
2286	315	2343	305	2402	176	2459	271
2287	315	2344	303	2403	40	2460 (2640 in	
2288	323	2345	303	2404	160, 421	err.)	14

C. P.	Page	C. P.	Page	C. P.	Page	C. P.	Page
2461	293	2519	192	2576	235	2633	356
2462	292	2520	73	2577	58	2634	350
2463	262	2521	117	2578	195	2635	351, 434
2464	265	2522	158	2579	195	2636	5
2465	56, 408	2523	269	2580	198	2637	60
2466	195	2524	32	2581	124	2638 (2368 in err.)	
2467	194	2525	74	2582	124		144
2468	112	2526	263	2583	17	2639	143, 419
2470	152	2527	306	2584	52	2640	19
2471	92	2528	304	2585 = 2520.		2641	275
2472	153	2529	285	2586	76, 410	2642 = 2641.	
2473	427	2530	288	2587	32	2643	313
2474 = 1795	175	2531	283	2588	121	2644	109
2475	189	2532	280	2589	134	2645	38, 403
2476	33	2533	181	2590	34	2646	38, 403
2477	74, 410	2534 = 2208	289	2591	245	2647	39, 404
2478 = 1954	214	2535 = 2208.		2592	170	2648	176
2479	60	2536	256	2593	229	2649	79
2480	15	2537	265	2594	229	2650	61
2481	188	2538	112	2595	68	2651	32
2482	82	2539	118	2596	101	2652	149
2483	134	2540	116	2597	128	2653	152
2484	366	2541	242	2598	173	2654	296
2485	253	2542	42	2599	173	2655	300
2486 = 2485.		2543	385	2600	41	2656	183
2487	255	2544	6	2601	281	2657	19
2488	256	2545	55	2602	289	2658	38, 403
2489	256	2546	47, 406	2603	19	2659	2
2490	256	2547	46, 406	2604	19	2660	3
2491	256	2548	195	2605	256	2661	98
2492	135	2549	197	2606	255	2662	89
2493	118	2550	16	2607	152	2663	92
2494	118	2551	133	2608	19	2664	294
2495	117	2552	135	2609	147	2665	294
2496	117	2553	22	2610	152	2666	294
2497	17, 400	2554	135	2611	156	2667	174
2498	133	2555	251	2612	8	2668	122
2499	271	2556	76	2613	54	2669	21
2500 = 2407	280	2557	163	2614	75	2670	198
2501	69, 409	2558	230	2615	108	2671	198
2502	282	2559	230	2616	109	2672	198
2503	10, 399	2560	285	2617	109	2673	201
2504 = 1193	61	2561	285	2618	105	2674	52
2505	19	2562	266	2619	106	2675	50, 406
2506	296	2563	267	2620	11	2676	76, 410
2507	176	2564	282	2621	112	2677	77
2508	93, 413	2565	268	2622	116	2678	11
2509	15	2566	29	2623	114, 415	2679	55
2510	158	2567	26	2624	375	2680	177
2511	296	2568	31	2625	365	2681	186
2512	156	2569	43, 405	2626	373	2682	119
2513	152	2570	113	2627	354	2683 = 2685.	
2514 = 1911	179	2571		2628	355	2684	112
2515	192	2572	296	2629	355	2685	111
2516	201	2573	155	2630	355	2686	113
2517	197	2574	1	2631	355	2687	108
2518	192	2575	155	2632	355	2688	187

INDEX OF C. P. NUMBERS.

C. P.	Page	C. P.	Page	C. P.	Page	C. P.	Page
2689	174	2746	352	2801	116	2857	63
2690	1	2747	348	2802	416	2858	29
2691	193	2748	347	2803	114	2859	212
2692	34	2749	355	2804	12	2860	197
2693	114	2750	356	2805	126	2861	200
2694	116, 417	2751	355	2806	126	2862	116
2695	49, 406	2752	347, 434	2807	128	2863	119
2696	214	2753 (2573 in err.)		2808	128	2864	216
2697	214		358	2809	24	2865	216
2698	242	2754=881	362	2810	130	2866	217
2699	258	2755	361	2811	123	2867	216
2700	125	2756	363	2812	130	2868	216
2701	254	2757	13	2813	130	2869	325
2702	6	2758	31	2814	133	2870	326
2703	8	2759	31	2815	156	2871	326
2704=196	273	2760	27	2816	153	2872	326
2705	151	2761	299	2817	142	2873	330
2706	148	2762	199	2818	262	2874	330
2707	149	2763=188	46	2819	141	2875	367
2708	50	2764	51	2820	137	2876	357
2709	157	2765	101	2821	16	2877	348
2710	157	2766	70	2822	163	2878	345
2711	145, 420	2767	101	2823	335	2879	344
2712=2192	261	2768	102	2824=1761	166	2880	218
2713=2199	261	2769	102	2825	161	2881	218
2714	292	2770	102	2826	166	2882	260
2715	165	2771	102	2827	168	2883	220
2716	239	2772	81	2828	171	2884	219
2717	224	2773	254	2829	173	2885	219
2718	233	2774=2772.		2830	176	2886	220
2719	73	2775	83	2831	262	2887	220
2720	53	2776=1447.		2832	177	2888	221
2721	134	2777	85	2833	182	2889	222
2722	12	2778	86	2834	192	2890	237
2723	29	2779	86	2835	185	2891	
2724	252	2780	86	2836	423	2892=217	239
2725	63	2781	90	2837	196	2893	242
2726	247	2782	89	2838	199	2894	243
2727	11	2783	91	2839	201	2896	334
2728	220	2784	95, 414	2840	204	2897	226
2729	178	2785	96	2841	204	2898	226
2730	180	2786	96	2842	205	2899	443
2731	178	2787 (2784 in error)	64, 409	2843	207	2900	225
2732	328	2788	64	2844	208	2901	402
2733	367	2789	67	2845	210	2902	31
2734	327	2790	66	2846=1943	209	2903	33
2735	325	2791	65	2847	211	2904	234
2736	318	2792	14	2848	209	2905	384
2737	316	2793	123	2849	210	2906	22
2738	340	2794	123	2850 (2854 in err.)	211	2907	245
2739	313	2795	123		211	2908	250
2740	299	2796	121	2851	212	2909	249
2741	304	2797	121	2852=1937	212	2910	247
2742	297	2798	110	2853	213	2911	248
2743	296	2799	108	2854	215	2912	249
2744	315	2800	105	2855	284	2913	251
2745	354			2856	264	2914	258

C. P.	Page	C. P.	Page	C. P.	Page	C. P.	Page
2916	41	2975	111	3032	363	3091	109
2917	51	2976	129	3033	354	3092	177
2918	19	2977	326	3034	354	3093	147
2920	187	2978	336	3035	354	3094	147
2921	255	2979	321	3036	352	3095	143
2922	289	2980	345	3037	354	3096	390
2923	11	2981	169	3038	347	3097	392
2924	179	2982	299	3039 (3090 in		3098	385
2925	93, 413	2983	306	err.) 351, 434		3099	281
2926 = 2929.		2984	397	3040	343	3100	385
2927	112	2985	397	3041	343	3101	385
2928	110	2986	397	3042	344	3102	387
2929	110	2987	387	3043	344	3103	386
2930	272	2988	379	3044	344	3104	395
2931	265	2989	223	3045	343	3105	379
2932	124	2990	153	3046	352, 434	3106	379
2933	187	2991	149, 425	3047	352, 434	3107	212
2934 = 1879	214	2992	4	3048	364	3108	222
2935	235	2993	225	3049	360	3109	225
2936	159	2994	233	3050	391	3110	233
2937	24	2995	238	3051	378	3111	219
2938	63	2996	218	3052	396	3112	249
2939	62	2997	31	3053	396	3113	252
2940	76, 410	2998	203	3055	333	3114	3
2941 = 913	359	2999	245	3056	333	3115	47, 406
2942	344	3000	245	3057		3116	266
2943	268	3001	55	3058	310	3117	28
2944	268	3002	125	3059	392	3118	335
2945	268	3003	68	3060	386	3119	326
2946	269	3004	75	3061	390	3120	313
2947	282	3005	36	3062	393	3121	300
2948	371	3006	212	3063	394	3122	317
2949		3007	108	3064	391	3123	320
2951	270	3008	115	3065	223	3124	296
2952	260	3009	173	3066 = 1340	384	3125	36
2953	292	3010	180	3067	384	3126	30, 401
2954	366	3011	181	3068	388	3127	25
2955	113	3012	176	3069	389	3128	124
2956	292	3013	275	3070	388	3129	87
2957	24	3014	277	3071	313	3130	99
2958	54	3015	286	3072	305	3131	172
2959	112	3016	285	3073	394	3132	144
2960	5	3017	312	3074	393	3133	286
2961	5	3018	319	3075	380	3134	284
2962	14	3019	319	3076	381	3135	367
2963	29	3020	316	3077	393	3136	368
2964	311	3021	317	3078	389	3137	357
2965	356	3022	326	3079	368	3138	365
2966	355	3023	325	3080	354	3139	384
2967	350	3024 = 2329	322	3081	310	3140	384
2968	363	3025	323	3082	320	3141	383
2969	324	3026	322	3083	265	3142	392
2970 = 3215.		3027	322	3084	220, 426	3143	394
2971	222	3028	340	3085	221	3144	386
2972	221	3029	345	3086	197	3145	382, 439
2973	87	3030	353	3088	222	3146	382
2974	107	3031	353	3089	223	3147	379

INDEX OF C. P. NUMBERS.

C. P.	Page	C. P.	Page	C. P.	Page	C. P.	Page
3148	53	3205	311	3263	365	3320	375
3149	97	3206	314	3264	383	3321	369
3150	89	3207	314	3265	395	3322	365, 436
3151	38, 403	3208	306	3266	378	3323	358
3152	205	3209	306	3267	383	3324	427
3153	241	3210	324	3268	384	3325	81
3154	247	3211	324	3269	383	3326	87
3155 <sup>p</sup> = 2278.		3212	338	3270	385	3327 (3329 in er.)	
3156	286	3213	321	3271	391		87
3157	320	3214	333	3272	389	3328	25
3158	52	3215	321	3273	391	3329	397
3159	331	3216	323	3274	392	3330	397
3160	299	3217	323	3275	390	3331	390
3161	355	3218	331	3276	394	3332	385
3162	352, 434	3219	356	3277	389	3333	327
3163	373	3220	349	3278	397	3334	331
3164	369	3221	342	3279	397	3335	9
3165	364	3222	346	3280	377	3336	259
3166	364	3223	323	3281	377	3337	269
3167	365	3224	322	3282	396	3338	77
3168	365	3225	354	3283	377	3339	77
3169	119	3226	353	3284	377	3340	35
3170	206	3227 = 3327	245	3285 = 3400	49	3341	35
3171	206	3228	346	3286	394	3342	202
3172	329	3229	347	3287	390	3343	167
3173	334	3230	347	3288	388	3344	140
3174	334	3231	347, 434	3289	384	3345	109
3175	316	3232	348, 433	3290	382	3346	111
3176		3233	350	3291	395	3347	112, 415
3177	295	3234	345	3292	380	3348	246
3178	295	3235	365	3293	396	3349	287
3179	295	3236	359	3294	55	3350	172
3180	295, 429	3237	359	3295	46, 405	3351	229
3181	295	3238	361	3296	121	3352	227
3182	297	3239	358	3297	123	3353	226
3183	297	3240	360	3298	123	3354	234
3184	298	3241	359	3299	123	3355	235
3185 = 3072	305	3242	361	3300	99	3356	232
3186	299	3243	360	3301	82	3357	234
3187	300	3244	358	3302	110	3358	321
3188	301	3245	371	3303	162	3359	376
3189	301	3546	361	3304	200	3360	397
3190	301	3247	372	3305	267	3361	397
3191	305	3248	366	3306	151	3362	45
3192	306	3249	358	3307	171	3363	386
3193	304	3251	373	3308	171	3364	3
3194	307	3252	375	3309	219	3365	19
3195	305	3253	374	3310	271	3366	222
3196	308	3254	364	3311	388	3367	201, 425
3197	309	3255	375	3312	320	3368	207
3198	356	3256	369	3313	322	3369	208
3199	309	3257	366	3314	322	3370	240
3200	309	3258	368	3315	334	3371	257
3201	310	3259	373	3316	342, 343	3372	261
3202	310	3260	362	3317	349	3373	316
3203	311	3261	369	3318	354, 435	3374	319
3204	312	3262	362	3319	353	3375	295, 429



C. P.	Page	C. P.	Page	C. P.	Page	C. P.	Page
3376	304	3433	288	3490	69	3546	94
3377	313	3434	176	3491	214	3547	99
3378	305	3435	238	3492 (3392 in err.)		3548	67
3379	308	3436	177		303	3549	55
3380	340	3437	208	3493	311	3550	22
3381	333	3438	116	3494	303, 429	3551	23
3382	341	3439	114	3495	430	3552	23
3383	392	3440	115	3496	334	3553	28
3384	390	3441	112	3497	384	3554	31
3385	430	3442	110	3498	391	3555	26
3386	72, 409	3443	106	3499	89	3556	368
3387	212	3444	77	3500	125	3557	347
3388	330	3445	413	3501	212	3558	343
3389	388	3446	176	3502	386	3559	343
3390	387	3447	177	3503	390	3560	342
3391	391	3448	212	3504	390	3561	343
3392	3	3449	257	3505 (2505 in err.)		3562	341
3393	3	3450	62		168	3563	315
3394	181	3451	62	3506	125	3564	309
3395	183	3452	404	3507	166	3565	314
3396	183	3453	134	3508	56	3566	302
3397	7	3454	186	3510	123	3568	269
3398	7	3455	70	3511	152	3569	270
3399	6	3456	69	3512	87	3570	277
3400	49, 406	3457	291	3513	83	3571	281
3401		3458	121	3514	83	3572	240
3402	52	3459	251	3515	301	3573	238
3403	51, 407	3460	129	3516	227	3574	241
3404	50, 407	3461	383	3517	227	3575	224
3405	34	3462	297	3518	256	3576	231
3406	34	3463	311	3519	270	3577	226
3407	36, 402	3464	334	3520	259	3578	232
3408	36	3465	320	3521	401	3579	211
3409	35	3466	317	3522	166	3580	211
3410	35	3467 = 3, var. $\beta$ .		3523	163	3581	204
3411	35	3468	346	3524	411	3582	
3412	39, 404	3469	352	3525	216	3583	165
3413	38, 403	3470	371	3526	16, 400	3584	129
3414	38, 403	3471	362	3527	6	3585	92
3415	37, 403	3472	189	3528	26	3586	90
3416	37, 404	3473	177	3529	27	3587	89
3417	139	3474	403	3530	302	3588	87
3418	146, 420	3475	279	3531	333	3589	88
3419	152	3476	182	3532	318	3590	85
3420	153	3477	180	3533	291	3591	411
3421	142	3478	182	3534	127	3592	83
3422	136	3479	112	3535	211	3593	81, 410
3423	157	3480	139	3536	211	3594	82
3424	155	3481	158	3537	225	3595	403
3425	313	3482	154, 421	3538	235	3596	409
3426	260	3483	39, 403	3539	166	3597	63
3427	260	3484	183	3540	201, 425	3598	84
3428	274	3485	96	3541		3599	85
3429	279	3486 = 3484	2	3542	143	3600	85
3430	280	3487	384	3543	143	3601	95, 414
3431	280	3488	54	3544	411	3602	95
3432	285	3489	377	3545	115	3603	96

INDEX OF C.P. NUMBERS.

C. P.	Page	C. P.	Page	C. P.	Page	C. P.	Page
3604	97	3661	226	3718	424	3774	423
3605	98	3662	233	3719	425	3775	432
3606	98	3663	234	3720	425	3776	432
3607	99	3664	232	3721	428	3777	434
3608	82, 410	3665	422	3722	396	3778	411
3609	90	3666	334	3723	426	3779	434
3610	119	3667	336	3724	335	3780	434
3611	120	3668	314	3725	336	3781	356
3612	120	3669	319	3726	405	3782=919	362
3613	412	3670	338	3727	315	3783	353
3614	441	3671	362	3728	319	3784	367
3615=1131	28	3672	264	3729	319	3785	402
3616	156	3673	405	3730	320	3786	433
3617	421	3674	400	3731	321	3787	432
3618	159	3675	402	3732	431	3788	432
3619	151	3676	408	2733	443	3789	426
3620	152	3677	416	3734	324	3790	413
3621	143	3678	418	3735	324	3791	401
3622	412	3679	442	3736	410	3792	424
3623	236	3680	339	3737	407	3793	424
3624	236	3681	369	3738	428	3794	422
3625	293	3682	362	3739	327	3795	428
3626	229	3683	358	3740	327	3796	435
3627	406	3684	370	3741(3471 in err.)	327	3797	435
3628	225	3685	426			3798	392
3629	247	3686	321	3742	329	3799	395
3630	165	3687	403	3743	329	3800	436
3631	168	3688	293	3744	330	3801	384
3632	169	3689	294	3745	337	3802	391
3633	161	3690	292	3746	336	3803	436
3634	401	3691	292, 428	3747	399	3804	437
3635	221	3692	292	3748	371	3805	431
3636	290	3693	306	3749	371	3806	438
3637	276	3694	307	3750	343	3807	438
3638	401	3695	297	3751	342	3808	438
3639	184	3696	297	3752	343	3809	415
3640	415	3697	299	3753	345	3810	410
3641=3461	(in error)	3698	297	3754	345	3811	411
		3699	315	3755	345	3812	441
3642	247	3700	316	3756	347	3813	411
3643	248	3701	317	3757=3737	348	3814	412
3644	417	3702	316	3758	348	3815	414
3645	242	3703	319	3759	349	3816	439
3646	401	3704	317	3760	349	3817	412
3647	210	3705	316	3761	350	3818	435
3648	211	3706	320	3762	435	3819	433
3649	212	3707	404	3763	353	3820	430
3650	213	3708	404	3764	432	3821	426
3651	217	3709	403	3765	412	3822	406
3653	399	3710	403	3766	399	3823	407
3654	418	3711	402	3767	427	3824	398
3655	269	3712	402	3768	429	3825	398
3656	419	3713	402	3769	429	3826	399
3657	299	3714	415	3770	427	3827	438
3658	246	3715	415	3771	370	3828=3848.	
3659	339	3716=2814	133	3772	399	3829	432
3660	228	3717	424	3773	401	3830	432

C.P.	Page	C.P.	Page	C.P.	Page	C.P.	Page
3831 . . .	398	3839 . . .	422	3846 . . .	436	3853 . . .	441
3832 . . .	410	3840 . . .	429	3847 . . .	436	3854 . . .	443
3833 . . .	401	3841 . . .	430	3848 . . .	436	3855 . . .	443
3834 . . .	416	3842 . . .	429	3849 . . .	443	3856 . . .	443
3835 . . .	416	3843 . . .	441	3850 . . .	441	3857 . . .	444
3836 . . .	417	3844 . . .	432	3851 . . .	441	3858 . . .	444
3837 . . .	420	3845 . . .	436	3852 . . .	442	3859 . . .	442
3838 . . .	419						

# INDEX ORDINUM ET GENERUM.

	Page		Page
Abelmoschus, <i>Med.</i>	26	Aganosma, <i>G. Don</i>	194
Aberia, <i>Hochst.</i>	400	Ageratum, <i>Linn.</i>	161
Abildgaardia, <i>Vahl</i>	347, 433	Agrimonia, <i>Tourn.</i>	102
Abrus, <i>Linn.</i>	91	Agrostis, <i>Linn.</i>	370
Abutilon, <i>Dill.</i>	27, 401	Agrostistachys, <i>Dalz.</i>	279
Acacia, <i>Willd.</i>	99, 415	Agyneia, <i>Linn.</i>	283, 288
Acalypha, <i>Linn.</i>	271	Ailanthus, <i>Desv.</i>	69
Acampe, <i>Lindl.</i>	303	ALANGIACEÆ	133
ACANTHACEÆ	223	Alangium, <i>Linn.</i>	133
Acanthonotus, <i>Benth.</i>	83	Albizzia, <i>Durazz.</i>	99
Acanthophippium, <i>Bl.</i>	307	Alchemilla, <i>Tourn.</i>	102
Acanthus, <i>Linn.</i>	232	Alisma, <i>Juss.</i>	332
Achyranthes, <i>Linn.</i>	249	ALISMACEÆ	332
Aclisia, <i>E. Mey.</i>	323	Allantodia, <i>Wall.</i>	385
Acmena, <i>DC.</i>	118	Allium, <i>Linn.</i>	339
Acorus, <i>Linn.</i>	337	Allæanthus, <i>Thw.</i>	263
Acranthera, <i>Arn.</i>	138	Allæophania, <i>Thw.</i>	147, 420
Acronychia, <i>Forst.</i>	409	Alocasia	336, 432
Acrostichum, <i>Linn.</i>	380, 395, 437	Alphonsea, <i>Hook. f. and T.</i>	11, 399
Acrotrema, <i>Jack.</i>	2, 398	Alpinia, <i>Linn.</i>	319
Actephila, <i>Bl.</i>	280	Alseodaphne, <i>N. ab Es.</i>	254
Actiniopteris, <i>Link</i>	386	Alsodeia, <i>Thouars</i>	21
Actinodaphne, <i>N. ab Es.</i>	256, 427	Alsophila, <i>R. Br.</i>	396
Actinostachys	379	Alstonia, <i>R. Br.</i>	193
Adansonia, <i>Linn.</i>	28	Alternanthera, <i>Mart.</i>	250
Adenanthera, <i>Linn.</i>	98	Alwisia, <i>Lindl.</i>	300
Adelia, <i>Schreb.</i>	272, 275	Alysicarpus, <i>Neck.</i>	87, 412
Adenochlæna, <i>Boiv.</i>	270	Alyxia, <i>R. Br.</i>	191
Adenosma, <i>N. ab Es.</i>	224	Amanoa, <i>Aubl.</i>	280, 428
Adenostemma, <i>Forst.</i>	162	AMARANTACEÆ	247
Adhatoda	233	Amarantus, <i>Tourn.</i>	247, 248
Adiantum, <i>Linn.</i>	387	AMARYLLIDACEÆ	324
Adinandra, <i>Jack.</i>	405	Amblogyne	248
Æchmandra, <i>Arn.</i>	125	Ambrosinia	334
ÆGICERACEÆ	174	Ameletia, <i>DC.</i>	122
Ægicera, <i>Gærtn.</i>	174	Ammannia, <i>Houst.</i>	121
Æginetia, <i>Linn.</i>	221	Amomum, <i>Linn.</i>	316, 430
Æluropus, <i>Trin.</i>	374	Amoora, <i>Roxb.</i>	60, 409
Aerides, <i>Lour.</i>	305, 429	Amorphophallus, <i>Bl.</i>	335, 443
Aerva, <i>Forsk.</i>	238, 427	AMPELIDÆ	62
Æschynanthus, <i>Jack.</i>	206	Amphidonax, <i>N. ab Es.</i>	370
Æschynomene, <i>Linn.</i>	85	Amphirhapis, <i>DC.</i>	162
Ætheilema, <i>R. Br.</i>	225	Anacardium, <i>Rottb.</i>	75

	Page		Page
Anagallis, <i>Tourn.</i> . . . . .	172	Artabotrys, <i>R. Br.</i> . . . . .	9
Anamirta, <i>Colebr.</i> . . . . .	12	Artanema, <i>Don</i> . . . . .	219
<i>Anaphalis</i> . . . . .	166	Artemisia, <i>Linn.</i> . . . . .	165
Anatherum . . . . .	368	Arthrocnemum, <i>Moq.</i> . . . . .	246
Anaxagorea, <i>St. Hil.</i> . . . . .	10	<i>Arthromiscus</i> , <i>Thw.</i> . . . . .	47, 406
ANCISTROCLADEÆ . . . . .	188	Arthrostylis, <i>R. Br.</i> . . . . .	352
Ancistrocladus, <i>Vahl</i> . . . . .	188	Artocarpus, <i>Linn.</i> . . . . .	262, 427
Andrographis, <i>Wall.</i> . . . . .	232	Arum, <i>Linn.</i> 334, 335, 336, 432, 443	
Andropogon, <i>Linn.</i> , 364, 365, 366,	368, 437	Arundina, <i>Bl.</i> . . . . .	301
Aneilema, <i>R. Br.</i> . . . . .	322	Arundinaria, <i>Rich.</i> . . . . .	375, 437
Anemone, <i>Linn.</i> . . . . .	1	Arundinella, <i>Raddi</i> . . . . .	362
Angiopteris, <i>Hoffm.</i> . . . . .	379, 437	<i>Arundo</i> . . . . .	370
Angræcum, <i>Thouars</i> . . . . .	306	<i>Ascarina</i> . . . . .	293
<i>Anguillaria</i> . . . . .	339	ASCLEPIADACEÆ . . . . .	195
Ania, <i>Lindl.</i> . . . . .	301	Asparagus, <i>Linn.</i> . . . . .	339
Aniseia, <i>Choisy</i> . . . . .	212	Aspidium, <i>Sw.</i> . . . . .	389, 438, 444
<i>Anisocampium</i> . . . . .	390	Asplenium, <i>Linn.</i> . . . . .	383, 438
Anisochilus, <i>Wall.</i> . . . . .	238	<i>Asteracantha</i> . . . . .	225
<i>Anisogonium</i> . . . . .	385	<i>Astylis</i> , <i>Wight.</i> . . . . .	287
Anisomeles, <i>R. Br.</i> . . . . .	240	Asystasia, <i>Bl.</i> . . . . .	235, 427
<i>Anisonema</i> , <i>A. Juss.</i> . . . . .	282	Atalantia, <i>Corr.</i> . . . . .	44, 405
Anisophyllea, <i>R. Br.</i> . . . . .	119	Ate, <i>Lindl.</i> . . . . .	309
<i>Anisophyllum</i> , <i>Don</i> . . . . .	119	<i>Athyrium</i> . . . . .	384
Anodendron, <i>A. DC.</i> . . . . .	194	Atriplex, <i>Gærtn.</i> . . . . .	246
Anæctochilus, <i>Bl.</i> . . . . .	314	Atylosia, <i>W. and Arn.</i> . . . . .	91
<i>Anogeissus</i> , <i>Wall.</i> . . . . .	103	AURANTIACEÆ . . . . .	44
<i>Anomospermum</i> , <i>Dalz.</i> . . . . .	280	Avena, <i>Linn.</i> . . . . .	372
ANONACEÆ . . . . .	6	Avicennia, <i>Linn.</i> . . . . .	244
Anstrutheria, <i>Gardn.</i> . . . . .	121	<i>Axanthes</i> , <i>Bl.</i> . . . . .	139, 418
Anthistiria, <i>Linn.</i> . . . . .	366, 436	Axinandra, <i>Thw.</i> . . . . .	122
<i>Anthocometes</i> . . . . .	234	Azadarachta, <i>A. Juss.</i> . . . . .	59
Antiaris, <i>Lesch.</i> . . . . .	263, 427	Azima, <i>Lam.</i> . . . . .	191
Antidesma, <i>Linn.</i> . . . . .	289	Azolla, <i>Lam.</i> . . . . .	378
<i>Antrophyum</i> . . . . .	382	Balanophora, <i>Forst.</i> . . . . .	293
Apaturia, <i>Lindl.</i> . . . . .	301	BALANOPHOREÆ . . . . .	293
<i>Aphylleia</i> , <i>Champ.</i> . . . . .	294	BALSAMINEÆ . . . . .	65
Apluda, <i>Linn.</i> . . . . .	364	Bambusa, <i>Schreb.</i> . . . . .	375, 376, 437
Apocopis, <i>N. ab Es.</i> . . . . .	365	Barleria, <i>Linn.</i> . . . . .	230, 427, 443
APOCYNACEÆ . . . . .	191	Barringtonia, <i>Forst.</i> . . . . .	119
Apodytes, <i>E. Mey.</i> . . . . .	42	BARRINGTONIACEÆ . . . . .	119
Apollonias, <i>N. ab Es.</i> . . . . .	253	Bassia, <i>Kön.</i> . . . . .	175, 423
Aponogeton, <i>Thunb.</i> . . . . .	333	Batatas, <i>Rumph.</i> . . . . .	210
Aporosa, <i>Bl.</i> . . . . .	288	<i>Batis</i> . . . . .	263
Apostasia, <i>Bl.</i> . . . . .	315	<i>Batratherum</i> . . . . .	368
APOSTASIACEÆ . . . . .	315	Bauhinia, <i>Linn.</i> . . . . .	98
Appendicula, <i>Bl.</i> . . . . .	306	Baumea, <i>Gaud.</i> . . . . .	353
AQUIFOLIACEÆ . . . . .	183	Begonia, <i>Linn.</i> . . . . .	128, 417, 442
ARALIACEÆ . . . . .	131	BEGONIACEÆ . . . . .	128
Ardisia, <i>Linn.</i> . . . . .	173, 423	BERBERIDEÆ . . . . .	13
Areca, <i>Linn.</i> . . . . .	327	Berberis, <i>Linn.</i> . . . . .	13
Argyreia, <i>Lour.</i> . . . . .	209, 210	<i>Berchemia</i> , <i>Neck.</i> . . . . .	74, 410
Arisæma, <i>Mart.</i> . . . . .	335	<i>Bergera</i> , <i>Kön.</i> . . . . .	46, 406
Aristida, <i>Linn.</i> . . . . .	370	<i>Berghausia</i> . . . . .	363
Aristolochia, <i>Tourn.</i> . . . . .	291	Bergia, <i>Linn.</i> . . . . .	23
ARISTOLOCHIACEÆ . . . . .	291	Berrya, <i>Rorb.</i> . . . . .	32
AROIDEÆ . . . . .	334	<i>Bhesa</i> , <i>Arn.</i> . . . . .	72

	Page		Page
Bidaria, <i>Endl.</i> . . . . .	198	Callitriche, <i>Linn.</i> . . . . .	290
Bidens, <i>Linn.</i> . . . . .	165	Calonyction, <i>Choisy</i> . . . . .	211
BIGNONIACEÆ . . . . .	206	Calophanes, <i>Don</i> . . . . .	225
<i>Biophytum</i> , <i>DC.</i> . . . . .	64, 409	Calophyllum, <i>Linn.</i> . . . . .	51, 407
Bixa, <i>Linn.</i> . . . . .	16	Calosantes, <i>Bl.</i> . . . . .	206
Blachia, <i>Baill.</i> . . . . .	277	Calotropis, <i>R. Br.</i> . . . . .	196
<i>Blackwellia</i> , <i>Comm.</i> . . . . .	79, 410	<i>Calymmodon</i> . . . . .	390
Blainvillea, <i>Cass.</i> . . . . .	164	<i>Calyptranthes</i> . . . . .	116, 117
Blechnum, <i>Linn.</i> . . . . .	381	<i>Cambogia</i> . . . . .	48, 49
Blepharis, <i>Juss.</i> . . . . .	231	<i>Cameraria</i> . . . . .	191
Blumea, <i>DC.</i> . . . . .	163, 422	Campanula, <i>Linn.</i> . . . . .	169
Blyxa, <i>Thouars</i> . . . . .	332	CAMPANULACEÆ . . . . .	169, 422
<i>Bobea</i> , <i>Miq.</i> . . . . .	421	Campbellia, <i>Wight</i> . . . . .	427, 443
Böhmeria, <i>Jacq.</i> . . . . .	259, 260	<i>Campium</i> . . . . .	380
Bœrhaavia, <i>Linn.</i> . . . . .	245	Camptosperma, <i>Thw.</i> . . . . .	78
Bolbophyllum, <i>Thouars</i> . . . . .	298	<i>Campteria</i> . . . . .	386, 387
Bonnaya, <i>Lk. and Otto</i> . . . . .	219, 426	Canarium, <i>Linn.</i> . . . . .	79, 410
BORAGINACEÆ . . . . .	213	Canavalia, <i>DC.</i> . . . . .	88
Borassus, <i>Linn.</i> . . . . .	329	Canna, <i>Linn.</i> . . . . .	320
Borreria, <i>Mey.</i> . . . . .	151, 442	Canscora, <i>Lam.</i> . . . . .	204, 442
Botrychium, <i>Sw.</i> . . . . .	378	Cansjera, <i>Juss.</i> . . . . .	251
Boucerosia, <i>W. and A.</i> . . . . .	200	<i>Cantharospermum</i> . . . . .	91
Bouchea, <i>Cham.</i> . . . . .	241	Canthium, <i>Linn.</i> . . . . .	152
Brachypodium, <i>Beaun.</i> . . . . .	374	CAPPARIDACEÆ . . . . .	14
<i>Brachypterum</i> , <i>Benth.</i> . . . . .	93, 513	Capparis, <i>Linn.</i> . . . . .	15, 399
Brachyramphus, <i>DC.</i> . . . . .	168	CAPRIFOLIACEÆ . . . . .	136
<i>Brachyspatha</i> . . . . .	335, 443	Carallia, <i>Roxb.</i> . . . . .	120
<i>Bradleia</i> , <i>Gærtn.</i> . . . . .	285	Caralluma, <i>R. Br.</i> . . . . .	200
Bragantia, <i>Lour.</i> . . . . .	291	Cardamine, <i>Linn.</i> . . . . .	14, 399, 400
Brassica, <i>Linn.</i> . . . . .	399	Cardiospermum, <i>Linn.</i> . . . . .	54
Breweria, <i>R. Br.</i> . . . . .	213	Carex, <i>Linn.</i> . . . . .	354, 442
Briedelia, <i>Willd.</i> . . . . .	279	Careya, <i>Roxb.</i> . . . . .	119, 442
Bruguiera, <i>Lam.</i> . . . . .	120	Carissa, <i>Linn.</i> . . . . .	191
Bryonia, <i>Linn.</i> . . . . .	124, 125, 126, 127	<i>Carria</i> , <i>Gardn.</i> . . . . .	40
Bryophyllum, <i>Salisb.</i> . . . . .	129	CARYOPHYLLACEÆ . . . . .	24
Buchanania, <i>Roxb.</i> . . . . .	78	Caryota, <i>Linn.</i> . . . . .	329
Buchnera, <i>Linn.</i> . . . . .	220	Casearia, <i>Jacq.</i> . . . . .	19, 275
Bupleurum, <i>Tourn.</i> . . . . .	131	Cassia, <i>Linn.</i> . . . . .	95, 414, 441
Burmannia, <i>Linn.</i> . . . . .	325	Cassytha, <i>Linn.</i> . . . . .	258
BURMANNIACEÆ . . . . .	325	Catenaria, <i>Benth.</i> . . . . .	86
BURSERACEÆ . . . . .	408, 410	<i>Catha</i> , <i>Försk.</i> . . . . .	72, 409
Bursinopetalum, <i>Wight</i> . . . . .	42	CEDRELACEÆ . . . . .	61
Butea, <i>Roxb.</i> . . . . .	89	CELASTRACEÆ . . . . .	71
<i>Buterea</i> . . . . .	226	Celastrus, <i>Linn.</i> . . . . .	68, 72
BYTTNERIACEÆ . . . . .	29	Celosia, <i>Linn.</i> . . . . .	247
		Celsia, <i>Linn.</i> . . . . .	217
CACTACEÆ . . . . .	129	Celtis, <i>Tourn.</i> . . . . .	267
Cadaba, <i>Försk.</i> . . . . .	15	Centotheca, <i>Desv.</i> . . . . .	374
Cæsalpinia, <i>R. Br.</i> . . . . .	94, 414	Centranthera, <i>R. Br.</i> . . . . .	220
Cajanus, <i>DC.</i> . . . . .	90	Centrosthachys, <i>Wall.</i> . . . . .	249
<i>Calamagrostis</i> . . . . .	370	<i>Centrostylis</i> , <i>Baill.</i> . . . . .	270
Calamintha, <i>Benth.</i> . . . . .	239	<i>Cephaloschænus</i> . . . . .	352
Calamus, <i>Linn.</i> . . . . .	330, 431	Cephalostigma, <i>A. DC.</i> . . . . .	422
Calanthe, <i>R. Br.</i> . . . . .	308	Cerastium, <i>Linn.</i> . . . . .	24
Callicarpa, <i>Linn.</i> . . . . .	243	<i>Ceratogynum</i> , <i>Wight</i> . . . . .	284
<i>Callipteris</i> . . . . .	385	CERATOPHYLLACEÆ . . . . .	290
CALLITRICHACEÆ . . . . .	290	<i>Ceratophyllum</i> , <i>Linn.</i> . . . . .	290

	Page		Page
<i>Ceratopsis</i> , Lindl. . . . .	311	<i>Cœlachne</i> , R. Br. . . . .	373
<i>Ceratopteris</i> , Brongn. . . . .	379	<i>Cœlogyne</i> , Lindl. . . . .	300
<i>Cerbera</i> , Linn. . . . .	192	<i>Coffea</i> , Linn. . . . .	154, 421
<i>Ceriops</i> , Arn. . . . .	120	<i>Coix</i> , Linn. . . . .	357
<i>Ceropegia</i> , Linn. . . . .	199	<i>Coldenia</i> , Linn. . . . .	215
<i>Chætocarpus</i> , Thw. . . . .	274	<i>Coleus</i> , Lour. . . . .	238
<i>Chætocyperus</i> , N. ab Es. . . . .	351, 435	<i>Colocasia</i> , Ray . . . . .	335, 432
CHAILLETIACEÆ . . . . .	79	<i>Colubrina</i> , Rich. . . . .	75
<i>Chamabainia</i> , Wight . . . . .	260	COMBRETACEÆ . . . . .	103
<i>Chamissoa</i> , Kunth . . . . .	247	<i>Combretum</i> , Löffl. . . . .	103, 415
<i>Championia</i> , Gardn. . . . .	208	<i>Commelyna</i> , Dill. . . . .	321
<i>Chasalia</i> , Comm. . . . .	150, 421	COMMELYNACEÆ . . . . .	321
<i>Chavica</i> , Miq. . . . .	292, 428	COMPOSITÆ . . . . .	160
<i>Cheilanthes</i> , Sw. . . . .	387	CONNARACEÆ . . . . .	80
<i>Cheirostylis</i> , Bl. . . . .	313	<i>Connarus</i> , Linn. . . . .	80, 410
<i>Chenopodium</i> , Tourn. . . . .	246	<i>Conocarpus</i> , Gærtn. . . . .	104
<i>Chickrassia</i> , A. Juss. . . . .	61	<i>Conocephalus</i> . . . . .	261
<i>Chilocarpus</i> . . . . .	191	CONVOLVULACEÆ . . . . .	209
<i>Chionachne</i> , R. Br. . . . .	357	<i>Convolvulus</i> . . . . .	210
<i>Chionanthus</i> , Linn. . . . .	188	<i>Conyza</i> , Less. . . . .	163
<i>Chirita</i> , Ham. . . . .	207	<i>Corchorus</i> , Linn. . . . .	31, 401
<i>Chironia</i> . . . . .	203	<i>Cordia</i> , Plum. . . . .	213
CHLORANTHACEÆ . . . . .	293	<i>Cordyline</i> . . . . .	338
<i>Chloranthus</i> , Sw. . . . .	293	<i>Corymbis</i> , Thouars . . . . .	314
<i>Chloris</i> , Linn. . . . .	371	<i>Corypha</i> , Linn. . . . .	329
<i>Chlorophytum</i> , Ker . . . . .	339	<i>Cosciniun</i> , Colebr. . . . .	12
<i>Chloroxylon</i> , DC. . . . .	61	<i>Cosmostigma</i> , Wight . . . . .	192
<i>Chondrachne</i> . . . . .	346	<i>Costus</i> , Linn. . . . .	320
<i>Chonemorpha</i> , G. Don . . . . .	194	<i>Cottonia</i> , Wight . . . . .	303
<i>Choripetalum</i> , A. DC. . . . .	173	<i>Covellia</i> , Gasparr. . . . .	266
<i>Chorisandra</i> , Wight . . . . .	281	CRASSULACEÆ . . . . .	129
<i>Christisonia</i> , Gardn. . . . .	221, 443	<i>Cratæva</i> , Linn. . . . .	14
<i>Chrysophyllum</i> , Linn. . . . .	174	<i>Crawfurdia</i> , Wall. . . . .	204
<i>Chrysopogon</i> , Trin. . . . .	366	<i>Cressa</i> , Linn. . . . .	213
<i>Cicca</i> . . . . .	281	<i>Crinum</i> , Linn. . . . .	324
<i>Cicendia</i> . . . . .	204	<i>Crossandra</i> , Salisb. . . . .	231
<i>Cinnamomum</i> , Burm. . . . .	252	<i>Crotalaria</i> , Linn. . . . .	81, 410, 441
<i>Cirrhopetalum</i> , Lindl. . . . .	299	<i>Croton</i> , Linn. . . . .	272, 275, 277, 278
<i>Cissampelos</i> , Linn. . . . .	13, 399	<i>Crozophora</i> , A. Juss. . . . .	443
<i>Cissus</i> , Linn. . . . .	62	CRUCIFERÆ . . . . .	14, 399
<i>Citrullus</i> , Neck. . . . .	126	<i>Cryptocarya</i> , R. Br. . . . .	254
<i>Cladium</i> , P. Browne . . . . .	353	<i>Cryptocoryne</i> , Fisch. . . . .	334, 432
<i>Claoxylon</i> , A. Juss. . . . .	271	<i>Cryptolepis</i> , R. Br. . . . .	195
<i>Clausena</i> , Burm. . . . .	47, 406	<i>Cryptophragmium</i> . . . . .	232
<i>Cleghornia</i> , Wight . . . . .	194	<i>Cucumis</i> , Linn. . . . .	126, 127
<i>Cleidion</i> , Bl. . . . .	272	CUCURBITACEÆ . . . . .	124
<i>Cleisostoma</i> , Bl. . . . .	304	<i>Cudrania</i> , Trécul . . . . .	262
<i>Clematis</i> , Linn. . . . .	1	<i>Cudranus</i> , Rumph. . . . .	262
<i>Cleome</i> , DC. . . . .	14	<i>Cullenia</i> , Wight . . . . .	28
<i>Clerodendron</i> , Linn. . . . .	243	<i>Cupania</i> . . . . .	56
<i>Cleyera</i> , Thunb. . . . .	40, 41, 405	<i>Curculigo</i> , Gærtn. . . . .	324
<i>Clitorea</i> , Linn. . . . .	88	<i>Curcuma</i> , Linn. . . . .	316, 430
<i>Cluytia</i> . . . . .	279, 280	<i>Cuscuta</i> , Tourn. . . . .	213
<i>Cnemidia</i> , Lindl. . . . .	314	<i>Cyanospermum</i> , W. and A. . . . .	92
<i>Coccinea</i> , W. and A. . . . .	128	<i>Cyanotis</i> , Don . . . . .	323
<i>Cocculus</i> , DC. . . . .	13	<i>Cyathea</i> , Sw. . . . .	396
<i>Cocos</i> , Linn. . . . .	330	<i>Cyathocalyx</i> , Champ. . . . .	9

	Page		Page
Cyathula, <i>Lour.</i>	249	Dichilanthe, <i>Thw.</i>	136
CYCADEÆ	294	Dichopsis, <i>Thw.</i>	176, 423
Cycas, <i>Linn.</i>	294	Dichrocephala, <i>DC.</i>	162
Cyclea, <i>Arn.</i>	13	Dichrostachys, <i>DC.</i>	99
Cyclostemon, <i>Bl.</i>	286	Dicksonia, <i>L'Her.</i>	389
<i>Cylindrochilus</i> , <i>Thw.</i>	307, 430	Dicliptera, <i>Juss.</i>	235
<i>Cylindropus</i>	354	<i>Diclisodon</i>	393
Cymbidium, <i>Sw.</i>	302, 307	Dicræa, <i>Thouars</i>	222
<i>Cyminosma</i> , <i>Gærtn.</i>	69, 409	Dictyospermum	322
Cymodocea, <i>Kön.</i>	333	Didymocarpus, <i>Wall.</i>	207, 426
<i>Cynanchum</i>	195, 196, 198	Dienia, <i>Lindl.</i>	297
Cynoctonum, <i>E. Mey.</i>	195	Digera, <i>Forsk.</i>	249
Cynodon, <i>Pers.</i>	371	<i>Dilivaria</i>	232
Cynoglossum, <i>Tourn.</i>	215	Dillenia, <i>Linn.</i>	5, 398
Cynometra, <i>Linn.</i>	97	DILLENACEÆ	2
CYPERACEÆ	342	Dimeria, <i>R. Br.</i>	369, 444
Cyperus, <i>Linn.</i>	342, 351, 432	<i>Dimocarpus</i>	58
CYTANDRACEÆ	206	Dimorphocalyx, <i>Thw.</i>	278
Cyrtopera, <i>Lindl.</i>	302, 429	Dioclea, <i>H. B. K.</i>	412
Cyrtosia, <i>Bl.</i>	311	Dioscorea, <i>Linn.</i>	325, 431
		DIOSCOREACEÆ	325, 442
<i>Dactylis</i>	374	Diospyros, <i>Del.</i>	178, 423
Dactyloctenium, <i>Willd.</i>	371	Diplacrum, <i>R. Br.</i>	354
Dædalacanthus, <i>T. And.</i>	229	<i>Diplazium</i>	385
Dæmia, <i>R. Br.</i>	196	Diplocentrum, <i>Lindl.</i>	306
Dalbergia, <i>Linn.</i>	93, 413, 414	<i>Diploclinium</i>	129
Dalechampia, <i>Plum.</i>	270	<i>Diplosiphon</i>	332
Dalzellia, <i>Wight</i>	223	DIPSACEÆ	160
<i>Damaonium</i>	332	Dipsacus, <i>Tourn.</i>	160, 421
<i>Daphne</i>	250	<i>Dipteracanthus</i>	225
Daphnidium, <i>N. ab Es.</i>	257	DIPTEROCARPEÆ	33
Daphniphyllum, <i>Bl.</i>	290	Dipterocarpus, <i>Gærtn.</i>	33, 402
Dasyaulus, <i>Thw.</i>	175, 423	Dischidia, <i>R. Br.</i>	198
DATISCACEÆ	252	Discospermum, <i>Dalz.</i>	156
Datura, <i>Linn.</i>	217	<i>Discostigma</i> , <i>Hassk.</i>	49, 406
Davallia, <i>Linn.</i>	388, 438	Disperis, <i>Sw.</i>	311
<i>Debregeasia</i>	261	Disporum, <i>Salisb.</i>	338
<i>Decaneurum</i>	161	<i>Dithyrocarpus</i> , <i>Kunth</i>	323
Delima, <i>Linn.</i>	2	Dodonæa, <i>Linn.</i>	59
Dendrobium, <i>Sw.</i>	297, 299, 429	Dolichos, <i>Linn.</i>	90
Dendrocalamus, <i>N. ab Es.</i>	376	Doodia, <i>R. Br.</i>	382
Dendrocolla, <i>Bl.</i>	306, 430	Doona, <i>Thw.</i>	34, 402
Dendrolobium, <i>Benth.</i>	86	Dopatrium, <i>Ham.</i>	218
<i>Dennstædtia</i>	389	Doronicum, <i>Linn.</i>	167
Dentella, <i>Forst.</i>	144, 420	Dorstenia, <i>Plum.</i>	264
Derris, <i>Lour.</i>	92, 413	Dracæna, <i>Vand.</i>	338
Desmanthus, <i>Benth.</i>	98	<i>Dracontium</i>	336
<i>Desmochæte</i>	249	Drosera, <i>Linn.</i>	21
Desmodium, <i>DC.</i>	85, 86, 411, 441	DROSERACEÆ	21
Desmostemon, <i>Thw.</i>	278	Drymaria, <i>Willd.</i>	25
Detarium, <i>Juss.</i>	414	Drymispermum, <i>Reinw.</i>	251, 427
Diacalpe, <i>Bl.</i>	396	Drymoglossum, <i>Prest</i>	381
Dialium, <i>Linn.</i>	97	<i>Drynaria</i>	395
Dianella, <i>Lam.</i>	338	Dumasia, <i>DC.</i>	88
Dicellostyles, <i>Benth.</i>	401	Dunbaria, <i>W. and A.</i>	90, 412
Dicerma, <i>DC.</i>	86	<i>Dyschoriste</i> , <i>N. ab Es.</i>	225
<i>Dichætaria</i> , <i>N. ab Es.</i>	444	<i>Dysodidendron</i> , <i>Gardn.</i>	150



	Page		Page
Dysophylla, <i>Bl.</i>	239	ERYTHROXYLÆ	53
Dysoxylon, <i>Bl.</i>	60, 409	<i>Erythroxyton</i>	54
EBENACEÆ	178	<i>Eugenia, Linn.</i>	113, 114, 416, 417
<i>Ebermaiera, N. ab Es.</i>	224	<i>Eulalia, Kunth</i>	368
<i>Eccremanthus, Thw.</i>	57, 408	<i>Eulophia, R. Br.</i>	310
<i>Echinolæna?</i>	359	<i>Euphorbia, Linn.</i>	268, 427
<i>Echinolytrum</i>	350	EUPHORBIACEÆ	268, 442
<i>Eclipta, Linn.</i>	164	<i>Euphoria, Juss.</i>	408
<i>Ehretia, Linn.</i>	214	<i>Eupyrena, W. and A.</i>	153
ELÆAGNACEÆ	252	<i>Eurya, Thunb.</i>	41
<i>Elæagnus, Linn.</i>	252	<i>Euxolus, Rafin.</i>	248, 427
ELÆOCARPEÆ	32	<i>Evia, Comm.</i>	78
<i>Elæocarpus, Linn.</i>	32	<i>Evodia, Forst.</i>	69, 409
<i>Elæodendron, Jacq.</i>	73	<i>Evolvulus, Linn.</i>	213
<i>Elaphoglossum</i>	380	<i>Evonymus, Linn.</i>	73
<i>Elatostema, Forst.</i>	259, 427	<i>Exacum, Linn.</i>	203
<i>Eleiotis, DC.</i>	412	<i>Excæcaria, Linn.</i>	269
<i>Eleocharis, R. Br.</i>	351, 435	<i>Fagraea, Thunb.</i>	200, 425, 442
<i>Eleogiton</i>	350	<i>Falconeria, Royle</i>	270
<i>Eleogynus</i>	351	<i>Feronia, Corr.</i>	48
<i>Elephantopus, Cass.</i>	161	<i>Ficus, Linn.</i>	264, 265, 266
<i>Elettaria, Rheede</i>	318, 430	FILICES	378
<i>Eleusine, Gærtn.</i>	371	<i>Filicium, Thw.</i>	408
<i>Ellipanthus, Hook. f.</i>	410	<i>Fimbristylis, Vahl.</i>	347, 348, 433, 434
<i>Elytraria, Vahl</i>	224	<i>Firmiana, Marsili.</i>	29
<i>Elytrophorus, Beauv.</i>	374	<i>Flacourtia, Comm.</i>	17
<i>Embelia, Burm.</i>	172	FLACOURTIACEÆ	16
<i>Emblica, Gærtn.</i>	282	<i>Flagellaria, Linn.</i>	340
<i>Emilia, Cass.</i>	167, 422	<i>Flemingia, Roxb.</i>	92
<i>Empusa, Lindl.</i>	429	<i>Fleurya, Gaud.</i>	258
<i>Endespermum</i>	94	<i>Floscopa, Lour.</i>	323
<i>Endopogon</i>	226	<i>Flüggia, Kunth</i>	339
<i>Enhalus, L. C. Rich.</i>	332	<i>Flüggia, Willd.</i>	281
<i>Entada, Adans.</i>	98	<i>Freycinetia, Gaud.</i>	327
<i>Epaltes, Cass.</i>	164	<i>Fuirena, Linn.</i>	347
<i>Epicarpurus, Bl.</i>	264	<i>Gærtnera, Lam.</i>	201, 420, 425
<i>Epiphanes</i>	311	<i>Galactia, P. Browne</i>	88
<i>Epipogium, Gmel.</i>	311	<i>Galera, Bl.</i>	311
<i>Epistylum, Sw.</i>	283, 428	<i>Galium, Linn.</i>	151, 421
<i>Epithema, Bl.</i>	208	<i>Garcinia, Linn.</i>	48, 406
<i>Epithinia, Jack.</i>	157	<i>Gardenia, Ellis.</i>	159, 421
<i>Eragrostis, Beauv.</i>	373	<i>Garnotia, Brongn.</i>	363
<i>Eranthemum, Linn.</i>	229, 239	<i>Gastrodia, R. Br.</i>	311
<i>Eria, Lindl.</i>	299, 300, 429	<i>Gaultheria, Kalm.</i>	170
<i>Eriachne, N. ab Es.</i>	444	<i>Geanthus, Reinw.</i>	319
ERICACEÆ	170	<i>Gelonium, Roxb.</i>	274, 428
<i>Erigeron</i>	163	<i>Gendarussa</i>	233
<i>Eriocaulon, Linn.</i>	341	<i>Geniosporum, Wall.</i>	236
<i>Eriochloa, Humb. and Kunth</i>	358	<i>Gentiana, Tourn.</i>	204
<i>Eriococcus, Hassk.</i>	283	GENTIANACEÆ	203
<i>Eriodendron, DC.</i>	28	<i>Geodorum, Jacks.</i>	308
<i>Erycibe, Roxb.</i>	213	<i>Geophila, Don</i>	150
<i>Erythrina, Linn.</i>	89	<i>Georchis, Lindl.</i>	314
<i>Erythrocarpus</i>	274	GERANIACEÆ	64
<i>Erythrochilus, Reinw.</i>	271	<i>Geranium, Linn.</i>	64
<i>Erythrospermum, Lam.</i>	18		

	Page		Page
Gerardinia, <i>Gaud.</i>	259	<i>Haagea</i>	128
Gieseckia, <i>Linn.</i>	250	Habenaria, <i>Willd.</i>	309, 310
<i>Ginalloa?</i>	418	<i>Hæmatospermum</i> , <i>Wall.</i>	272
Gironniera, <i>Gaud.</i>	267	HALORAGEÆ	123
Givotia, <i>Griff.</i>	278	<i>Haloragis</i>	123
Gleichenia, <i>Sw.</i>	379	<i>Halostemma</i> , <i>Wall. mss.</i>	345
Glenniea, <i>Hook. f.</i>	408	Hapalosia, <i>Wall.</i>	25
Glinus, <i>Linn.</i>	24	<i>Haplachne</i>	369
Globba, <i>Linn.</i>	315	<i>Haplostylis</i>	352
Glochidion, <i>Forst.</i>	285	Harpullia, <i>Roxb.</i>	56
Gloriosa, <i>Linn.</i>	339	<i>Hebradendron</i>	49
Glycine, <i>Linn.</i>	88	Hedera, <i>Linn.</i>	131, 418
Glycosmis, <i>Corr.</i>	45, 406	Hedyotis, <i>Lam.</i>	140, 419
<i>Glycycaurus</i> , <i>Dalz.</i>	78, 441	Hedychium, <i>Kön.</i>	319
Glyptopetalum, <i>Thw.</i>	73	Hedysarum, <i>Jeauime</i>	411
Gmelina, <i>Linn.</i>	244	Helicia, <i>Lour.</i>	250
Gnaphalium, <i>Linn.</i>	166, 422	<i>Helichrysum</i>	166
<i>Gnidia</i>	250	Helicteres, <i>Linn.</i>	28
<i>Goldfussia</i>	228	Heligme, <i>Bl.</i>	193
Gomphandra, <i>Wall.</i>	44, 405	Heliophytum, <i>DC.</i>	215
Gomphia, <i>Schreb.</i>	71	Heliotropium, <i>Tourn.</i>	215
<i>Goniopteris</i>	394, 439	<i>Helminthospermum</i> , <i>Thw.</i>	267
Goniothalamus, <i>Bl.</i>	6, 398	Helminthostachys, <i>Kaulf.</i>	378
<i>Gonyanthes</i>	325	Helosciadium, <i>Koch</i>	130
GOODENOVIÆ	169	Hemarthria, <i>R. Br.</i>	363
Goodyera, <i>R. Br.</i>	313	Hemicyclia, <i>W. and A.</i>	287
Gordonia, <i>Ellis</i>	40	Hemidesmus, <i>R. Br.</i>	195
Gouania, <i>Jacq.</i>	75	Hemigyrosa, <i>Bl.</i>	56, 408
<i>Goughia</i> , <i>Wight</i>	290	Hemionitis, <i>Linn.</i>	382
<i>Govindovia</i> , <i>Wight</i>	314	Heracleum, <i>Linn.</i>	131
GRAMINEÆ	356	Heritiera, <i>Ait.</i>	28
Grammitis, <i>Sw.</i>	382	Hernandia, <i>Plum.</i>	258
Grangea, <i>Adans.</i>	163	Herpestes, <i>Gærtn. f.</i>	218, 426
<i>Grayia</i>	362	Heteropogon, <i>Pers.</i>	368, 437
Grewia, <i>Juss.</i>	31, 402	Heterostemma, <i>W. and A.</i>	198
Griffithia, <i>W. and A.</i>	158	<i>Heracentris</i>	223
Grislea, <i>Löfl.</i>	122	Heylandia, <i>DC.</i>	81
Grumilea, <i>Gærtn.</i>	147, 149	Hibiscus, <i>Linn.</i>	26, 401
<i>Guarea</i>	60	Hippocratea, <i>Linn.</i>	52
<i>Guatteria</i> , <i>Ruiz and Pav.</i>	10, 398	HIPPOCRATEACEÆ	52
Guazuma, <i>Plum.</i>	29	Hiptage, <i>Gærtn.</i>	53
Guettarda, <i>Linn.</i>	153	Holarrhena, <i>R. Br.</i>	194
Guilandina, <i>Linn.</i>	94	<i>Holcus</i>	362, 366
GUTTIFERÆ	48	<i>Hologamium</i>	365
Gymnema, <i>R. Br.</i>	197	<i>Holoptelæa</i>	267
<i>Gymnogramme</i>	382	HOMALINEÆ	79
Gymnopetalum, <i>Arn.</i>	127	Homalium, <i>Jacq.</i>	410
Gymnopogon, <i>Beauv.</i>	371, 372, 444	Hopea, <i>Roxb.</i>	36, 403
<i>Gymnopteris</i>	381, 437	Hortonia, <i>Wight</i>	11
Gymnosporia, <i>W. and A.</i>	409	Hoya, <i>R. Br.</i>	198
Gymnostachyum, <i>N. ab Es.</i>	232, 427	Hugonia, <i>Linn.</i>	25
Gynandropsis, <i>DC.</i>	14	HUGONIACEÆ	25
<i>Gynoon</i> , <i>A. Juss.</i>	285	<i>Humata</i>	388
Gynura, <i>Cass.</i>	166, 167	Humboldtia, <i>Vahl.</i>	97
<i>Gyrandra</i> , <i>Wall.</i>	290	Hunteria, <i>Roxb.</i>	191
Gyrinops, <i>Gærtn.</i>	251	<i>Hyalisma</i> , <i>Champ.</i>	294
Gyrocarpus, <i>Jacq.</i>	258	Hydnocarpus, <i>Gærtn.</i>	18

	Page		Page
Hydrilla, <i>Rich.</i>	331	Kadsura, <i>Juss.</i>	5
Hydrobryum, <i>Endl.</i>	223	Kæmpferia, <i>Linn.</i>	316
Hydrocera, <i>Bl.</i>	68	Kalanchoe, <i>Adans.</i>	129, 417
HYDROCHARIDÆ	331	Kanilia, <i>Bl.</i>	120
Hydrocotyle, <i>Tourn.</i>	130	<i>Karivia</i>	125
<i>Hydroglossum</i>	379	Kayea, <i>Wall.</i>	50, 407
Hydrolea, <i>Linn.</i>	209	Kirganelia, <i>Juss.</i>	282
HYDROLEACEÆ	209	Klugia, <i>Schlecht.</i>	208
Hydrophylax, <i>Linn.</i>	151	Knoxia, <i>Linn.</i>	151
Hygrophila, <i>R. Br.</i>	225	Kokoona, <i>Thw.</i>	52
Hygroryza, <i>N. ab Es.</i>	356	Kurrimia, <i>Wall.</i>	72, 409
<i>Hymenochæte</i>	351	<i>Kydia</i> , <i>Roxb.</i>	30, 401
<i>Hymenolepis</i>	381	Kyllingia, <i>Linn.</i>	345
Hymenophyllum, <i>Sm.</i>	397		
HYPERICACEÆ	48	LABIATÆ	236
Hypericum, <i>Linn.</i>	48	Lablab, <i>Adans.</i>	90
Hypolytrum, <i>Rich.</i>	346	<i>Lactuca</i>	168
<i>Hypoporum</i>	354	Lagenandra, <i>Dalz.</i>	334, 432
HYPOXIDACEÆ	323	Lagenophora, <i>Cass.</i>	162
Hypoxis, <i>Linn.</i>	323	Lagerströmia, <i>Linn.</i>	122
Hyptianthera, <i>W. and A.</i>	157, 421	Lagunea, <i>Cav.</i>	27
Hyrtaandra, <i>Miq.</i>	261	<i>Langsdorffia</i>	293
		Lantana, <i>Linn.</i>	242
Ichnanthus, <i>Beauv.</i>	361	Laportea, <i>Gaud.</i>	259
Ichnocarpus, <i>R. Br.</i>	194	Lappago, <i>Schreb.</i>	362
Ilex, <i>Linn.</i>	183	Lasia, <i>Lour.</i>	336
Ilysanthes, <i>Rafin.</i>	219	Lasianthera, <i>Beauv.</i>	44, 405
Impatiens, <i>Linn.</i>	65, 409	Lasianthus, <i>Jack.</i>	145, 420
Imperata, <i>Cyr.</i>	369	Lasiosiphon, <i>Fresen.</i>	250
Indigofera, <i>Linn.</i>	83, 411	<i>Lastrea</i>	391
<i>Inga</i>	100	LAURACEÆ	252
Ionidium, <i>Ventn.</i>	20	<i>Lawia</i> , <i>Tulasne</i>	223
Iphigenia, <i>Kunth</i>	339	Lawsonia, <i>Linn.</i>	122
Ipomea, <i>Linn.</i>	211, 426	Ledebouria, <i>Roth.</i>	432
Ipsca, <i>Lindl.</i>	301	Leca, <i>Linn.</i>	64
<i>Irina</i> , <i>Bl.</i>	408	Leersia, <i>Soland.</i>	356
Isachne, <i>R. Br.</i>	361, 373	<i>Legnotideæ</i>	120
Isanthera, <i>N. ab Es.</i>	208	LEGUMINOSÆ	81
<i>Isauxis</i> , <i>Arn.</i>	37, 404	<i>Leiopyxis</i> , <i>Miq.</i>	428
Ischæmum, <i>Linn.</i>	364, 365, 436, 444	Lemna, <i>Linn.</i>	331
Isolepis, <i>R. Br.</i>	350	LENTIBULARIÆ	171
Isonandra, <i>Wight</i>	176, 423, 442	Leonotis, <i>R. Br.</i>	241
Ixora, <i>Linn.</i>	154, 421	<i>Lepidenia</i> , <i>N. ab Es.</i>	254
		Lepidagathis, <i>Willd.</i>	231
<i>Jambolifera</i> , <i>Linn.</i>	409	<i>Lepidostachys</i> , <i>Wall.</i>	288
Jambosa, <i>Rumph.</i>	115	Lepironia, <i>Rich.</i>	346, 433
JASMINACEÆ	189	<i>Leptacanthus</i>	229
Jasminum, <i>Tourn.</i>	189, 424	Leptadenia, <i>R. Br.</i>	198
Jatropha, <i>Kunth</i>	277	Leptaspis, <i>R. Br.</i>	357
Jonesia, <i>Roxb.</i>	97	Leptochloa, <i>Beauv.</i>	371
Josephia, <i>Wight</i>	307	<i>Leptogramme</i>	382
Julostylis, <i>Thw.</i>	30	<i>Leptostachya</i>	233
JUNCACEÆ	340	<i>Lepurandra</i>	263
Juncus, <i>DC.</i>	340	<i>Lettsomia</i>	210
Jussiaea, <i>Linn.</i>	123	Leucas, <i>Benth.</i>	240
Justicia, <i>Linn.</i>	233, 235, 427	Leucocodon, <i>Gardn.</i>	138
		<i>Leucostegia</i>	388

	Page		Page
Ligustrum, <i>Tourn.</i>	188	MARSILEACEÆ	378
LILIACEÆ	337	<i>Martynia</i>	207, 228
Limacia, <i>Lour.</i>	12	<i>Marumia</i>	110
Limnanthemum, <i>Gmel.</i>	205	Medinella, <i>Gaud.</i>	106
<i>Limnocharis</i>	352	Megalachne, <i>Steud.</i>	372, 444
Limnophila, <i>R. Br.</i>	218	Melanthesa, <i>Bl.</i>	285
Limonia, <i>Linn.</i>	45, 405	Melanthium	339
Lindenbergia, <i>Lehm.</i>	218	Melastoma, <i>Linn.</i>	106
Lindsæa, <i>Dryand.</i>	388, 438	MELASTOMACEÆ	104
<i>Linociera</i>	188	<i>Melia, Linn.</i>	59, 408
LINEÆ	25	MELIACEÆ	59
Linum, <i>Linn.</i>	25	<i>Melica</i>	371
Liparis, <i>Rich.</i>	295, 306, 429	Meliosma, <i>Bl.</i>	59, 408
Lipocarpha, <i>R. Br.</i>	347	Melochia, <i>Linn.</i>	30
Lippia, <i>Linn.</i>	241	Melothria, <i>Linn.</i>	124, 417
<i>Litobrochia</i>	387	Memecylon, <i>Linn.</i>	110, 415, 416
Litsæa, <i>Juss.</i>	257	<i>Memorialis, Ham.</i>	261
Lobelia, <i>Linn.</i>	169	Meniscium, <i>Schreb.</i>	382, 438
LOGANIACEÆ	200	MENISPERMACEÆ	12
Lomaria, <i>Willd.</i>	381	Mentha, <i>Linn.</i>	239
Lophatherum, <i>Brongn.</i>	374	<i>Menyanthes</i>	205, 206
LORANTHACEÆ	133	<i>Meoschium</i>	364, 444
Loranthus, <i>Linn.</i>	133, 418	<i>Mephitidia, Reinw.</i>	145, 420
<i>Loxogramme</i>	382	Mesua, <i>Linn.</i>	50, 406, 407
Ludwigia, <i>Roxb.</i>	123	<i>Methonica</i>	339
Luffa, <i>Tourn.</i>	126, 417	<i>Meyenia</i>	224
Luisia, <i>Gaud.</i>	302, 429	Mezoneurum, <i>Desf.</i>	414
Lumnitzera, <i>Willd.</i>	103	Michelia, <i>Linn.</i>	5
Luvunga, <i>Ham.</i>	47, 406	Microcarpæa, <i>R. Br.</i>	220
LYCOPODIACEÆ	377	<i>Micrococca, Benth.</i>	271
Lycopodium, <i>Linn.</i>	377	<i>Microlepia</i>	389, 438
Lygodium, <i>Sw.</i>	378, 437	Micromelum, <i>Bl.</i>	46
Lysimachia, <i>Linn.</i>	172	Microrhynchus, <i>Less.</i>	168
LYTHRARIACEÆ	121	Microstachys, <i>A. Juss.</i>	270
		Microstylis, <i>Nutt.</i>	296, 429
		Microtropis, <i>Wall.</i>	71
Maba, <i>Forst.</i>	183	<i>Milium</i>	358
Macaranga, <i>Thouars</i>	274, 428	Miliusa, <i>Lesch.</i>	10
Machilus, <i>Rumph</i>	254	<i>Millingtonia, Roxb.</i>	59
<i>Maclura, Bl.</i>	262	Milnea, <i>Roxb.</i>	60
<i>Macræa, Wight</i>	282	Mimusops, <i>Linn.</i>	175
Macreightia, <i>A. DC.</i>	183, 423	<i>Miquelia</i>	363
Mæsa, <i>Forsk.</i>	172	Mischodon, <i>Thw.</i>	275, 428
MAGNOLIACEÆ	5	Mitrasacme, <i>Labill.</i>	200
Malacochæte	351	Mitrephora, <i>Bl.</i>	8
Mallea, <i>A. Juss.</i>	60	Mnesithea, <i>Kunth</i>	365
MALPIGHIACEÆ	53	<i>Mnianthus, Walp.</i>	223
MALVACEÆ	25	Moacurra, <i>Roxb.</i>	79
Malvastrum, <i>A. Gray</i>	441	Modecca, <i>Linn.</i>	128
Mangifera, <i>Linn.</i>	75	Mærua, <i>Forsk.</i>	15
Manisuris, <i>Linn.</i>	364	<i>Molinæa</i>	56
Mappia, <i>Jacq.</i>	43	Mollugo, <i>Linn.</i>	24
Maranta, <i>Plum.</i>	320	Momordica, <i>Linn.</i>	126
MARANTACEÆ	320	MONIMIACEÆ	11
Marattia, <i>Sw.</i>	379	Monocera, <i>Jack.</i>	33
<i>Mariscus</i>	344, 345	Monochilus, <i>Wall.</i>	313
Marsdenia, <i>R. Br.</i>	197	Monochoria, <i>Presl</i>	321
Marsilea, <i>Linn.</i>	378		

	Page		Page
<i>Monogramme</i> . . . . .	381	<i>Nymphæa, Linn.</i> . . . . .	14
<i>Monoporandra, Th.</i> . . . . .	39, 404	NYPHÆACEÆ . . . . .	14
<i>Monothecium, Hochst.</i> . . . . .	234	<i>Oberonia, Lindl.</i> . . . . .	296
<i>Moonia, Arn.</i> . . . . .	164	<i>Ochlandra, Thw.</i> . . . . .	376
<i>Morinda, Vaill.</i> . . . . .	144, 420	<i>Ochna, Linn.</i> . . . . .	70, 409
<i>Morisia</i> . . . . .	352	OCHNACEÆ . . . . .	70
<i>Morocarpus, Sieb. and Zucc.</i> . . . . .	261	<i>Ochrosia, Juss.</i> . . . . .	192
<i>Moschosma, Reich.</i> . . . . .	237	<i>Ocimum, Linn.</i> . . . . .	236
<i>Mucuna, Adans.</i> . . . . .	89	<i>Octarrhena, Thw.</i> . . . . .	305
<i>Mukia, Arn.</i> . . . . .	125	<i>Odina, Roxb.</i> . . . . .	78
<i>Muldera, Miq.</i> . . . . .	428	ŒCEOCLADES, <i>Lindl.</i> . . . . .	306
<i>Munronia, Wight</i> . . . . .	59	OLACACEÆ . . . . .	41
<i>Murraya, Kön.</i> . . . . .	45, 406	<i>Olax, Linn.</i> . . . . .	42
<i>Musa, Schreb.</i> . . . . .	321	<i>Olea, Tourn.</i> . . . . .	188
MUSACEÆ . . . . .	321	OLEACEÆ . . . . .	188
<i>Mussænda, Linn.</i> . . . . .	138	<i>Oleandra, Cav.</i> . . . . .	393
<i>Myriactis, Less.</i> . . . . .	162	ONAGRACEÆ . . . . .	123
<i>Myriogyne, Less.</i> . . . . .	165	<i>Oncosperma, Bl.</i> . . . . .	328
<i>Myriophyllum, Willd.</i> . . . . .	123	<i>Ophelia, Don</i> . . . . .	205, 426
<i>Myristica, Linn.</i> . . . . .	11, 399	<i>Ophioglossum, Linn.</i> . . . . .	378
MYRISTICACEÆ . . . . .	11	<i>Ophiopogon, Ker.</i> . . . . .	339
MYRSINACEÆ . . . . .	172	<i>Ophiorrhiza, Linn.</i> . . . . .	139, 419
<i>Myrsine, Linn.</i> . . . . .	173	<i>Ophioxylon, Linn.</i> . . . . .	191, 424
MYRTACEÆ . . . . .	114	<i>Opilia, Roxb.</i> . . . . .	41
<i>Myrtus, Tourn.</i> . . . . .	114	<i>Oplismenus</i> . . . . .	359
<i>Nageia, Roxb.</i> . . . . .	287	ORCHIDACEÆ . . . . .	295
NAIADACEÆ . . . . .	333	<i>Orchis</i> . . . . .	310
<i>Naraveli, DC.</i> . . . . .	1	<i>Oreocnide, Miq.</i> . . . . .	261
<i>Nasturtium, R. Br.</i> . . . . .	399	<i>Ormocarpum, Beauv.</i> . . . . .	85
<i>Nauclea, Linn.</i> . . . . .	137	<i>Ornithopteris</i> . . . . .	386
<i>Necamandra, Planch.</i> . . . . .	332	OROBANCHACEÆ . . . . .	221, 427
<i>Nelitris, Gærtn.</i> . . . . .	153	<i>Oropetium, Trin.</i> . . . . .	363
<i>Nelsonia, R. Br.</i> . . . . .	224	<i>Orophea, Bl.</i> . . . . .	8
NELUMBIACEÆ . . . . .	14	<i>Orthosiphon, Benth.</i> . . . . .	237
<i>Nelumbium, Juss.</i> . . . . .	14	<i>Oryza, Linn.</i> . . . . .	357
<i>Nematostigma, Planch.</i> . . . . .	267	<i>Osbeckia, Linn.</i> . . . . .	104
NEPENTHACEÆ . . . . .	290	<i>Osmelia, Thw.</i> . . . . .	20
<i>Nepenthes</i> . . . . .	290	<i>Osmunda, Linn.</i> . . . . .	379
<i>Nephelium, Linn.</i> . . . . .	57, 408, 408	<i>Osyris, Linn.</i> . . . . .	251
<i>Nephrodium</i> . . . . .	390-393	<i>Otonychium, Bl.</i> . . . . .	56
<i>Nephrolepis, Schott</i> . . . . .	393	<i>Ottelia, L. C. Rich.</i> . . . . .	332
<i>Neptunia, Lour.</i> . . . . .	99	OXALIDACEÆ . . . . .	64
<i>Nesæa, Comm.</i> . . . . .	417	<i>Oxalis, Linn.</i> . . . . .	64, 409
<i>Neurocalyx, Hook.</i> . . . . .	138	<i>Oxystelma, R. Br.</i> . . . . .	196
<i>Niehbuhria</i> . . . . .	15	<i>Pachycentria, Bl.</i> . . . . .	107
<i>Nipa, Rumph.</i> . . . . .	327	<i>Pachygone, Miers</i> . . . . .	13
<i>Niphobolus</i> . . . . .	395	<i>Palenga, Thw.</i> . . . . .	287
<i>Nomismia, W. and A.</i> . . . . .	91	<i>Palimbia, Bess.</i> . . . . .	131
<i>Norysca</i> . . . . .	48	PALMACEÆ . . . . .	327
<i>Notelæa</i> . . . . .	188	<i>Pancratium, Linn.</i> . . . . .	324
<i>Nothopegia, Bl.</i> . . . . .	441	PANDANEÆ . . . . .	327
<i>Nothosæerva, Wight</i> . . . . .	248	<i>Pandanophyllum, Hassk.</i> . . . . .	345, 433
<i>Notonia, DC.</i> . . . . .	168	<i>Pandanus, Linn.</i> . . . . .	327
<i>Notothiros?</i> . . . . .	418	PANGIACEÆ . . . . .	18
NYCTAGINACEÆ . . . . .	245	<i>Panicum, Linn.</i> 358, 361, 362, 436, 443	
<i>Nyctanthes, Juss.</i> . . . . .	190		

	Page		Page
PAPILIONACEÆ . . . . .	18	Pityranthe, <i>Thw.</i> . . . . .	29
Paramignya, <i>Wight</i> . . . . .	47, 406	Pladera, <i>Roxb.</i> . . . . .	204, 425
<i>Paratropia</i> . . . . .	132	PLANTAGINACEÆ . . . . .	245
Paritium, <i>A. Juss.</i> . . . . .	26	Plantago, <i>Linn.</i> . . . . .	245
Parochætus, <i>Ham.</i> . . . . .	82	Platanthera, <i>Rich.</i> . . . . .	310
PARONYCHIACEÆ . . . . .	25	<i>Platea</i> , <i>Bl.</i> . . . . .	44, 405
<i>Parsonia</i> . . . . .	193	Plecospermum, <i>Trécul</i> . . . . .	263
Paspalum, <i>Linn.</i> . . . . .	357	Plectranthus, <i>L'Her.</i> . . . . .	237
PASSIFLORACEÆ . . . . .	128	<i>Pleocnemia</i> . . . . .	390
<i>Patania</i> . . . . .	389	<i>Pleopeltis</i> . . . . .	394
Pavetta, <i>Linn.</i> . . . . .	155	Pleurostylia, <i>W. and A.</i> . . . . .	71
Pavonia, <i>Cav.</i> . . . . .	26, 401	PLUMBAGINACEÆ . . . . .	244
Pedaliium, <i>D. Roy</i> . . . . .	209	Plumbago, <i>Tourn.</i> . . . . .	244
Pedicularis, <i>Linn.</i> . . . . .	221	Poa, <i>Linn.</i> . . . . .	372, 373
<i>Pegia?</i> . . . . .	441	Podadeuia, <i>Thw.</i> . . . . .	273, 428
<i>Pellæa</i> . . . . .	386	<i>Podanthera</i> , <i>Wight</i> . . . . .	311
Pellionia, <i>Gaud.</i> . . . . .	259	Podochilus, <i>Bl.</i> . . . . .	306
Peltandra, <i>Wight</i> . . . . .	281	PODOSTEMACEÆ . . . . .	222
Pemphis, <i>Forst.</i> . . . . .	122	Podostemon, <i>Mich.</i> . . . . .	222
<i>Pentaloba</i> . . . . .	21	<i>Pæcilopteris</i> . . . . .	380
Pentatropis, <i>R. Br.</i> . . . . .	196	Pogonatherum, <i>Beauv.</i> . . . . .	365
Peperomia, <i>Ruiz and Pav.</i> . . . . .	292	Pogonia, <i>Juss.</i> . . . . .	430
Peplidium, <i>Del.</i> . . . . .	426	Pogonotrophe, <i>Miq.</i> . . . . .	266
Pericopsis, <i>Thw.</i> . . . . .	413	Pogostemon, <i>Desf.</i> . . . . .	239
Periostrophe, <i>N. ab Es.</i> . . . . .	234	Polanisia, <i>Rafin.</i> . . . . .	14
Peristylus, <i>Bl.</i> . . . . .	310	Pollia, <i>Thunb.</i> . . . . .	323
Perotis, <i>Ait.</i> . . . . .	369	Pollinia, <i>Trin.</i> . . . . .	368
Pestalozzia, <i>Moritzi</i> . . . . .	124	Polyalthia, <i>Bl.</i> . . . . .	9, 398
<i>Peucedanum</i> . . . . .	131	<i>Polybotrya</i> . . . . .	380
Phaius, <i>Lour.</i> . . . . .	300	Polycarpæa, <i>Lam.</i> . . . . .	25
Phalangium? . . . . .	339	Polygala, <i>Linn.</i> . . . . .	22, 400
Pharbitis, <i>Choisy</i> . . . . .	210	POLYGALACEÆ . . . . .	22
Phaseolus, <i>Linn.</i> . . . . .	89, 412	POLYGONACEÆ . . . . .	245
<i>Phoberos</i> , <i>Lour.</i> . . . . .	16, 400	Polygonum, <i>Linn.</i> . . . . .	245
Phœnix, <i>Linn.</i> . . . . .	329	<i>Polyodontia?</i> . . . . .	102
Pholidota, <i>Lindl.</i> . . . . .	300	Polypodium, <i>Linn.</i> . . . . .	393, 439, 444
Photinia, <i>Lindl.</i> . . . . .	100	Polypogon, <i>Desf.</i> . . . . .	370, 444
Phragmites, <i>Trin.</i> . . . . .	370	Polystachya, <i>Hook.</i> . . . . .	307, 308
Phreatia, <i>Lindl.</i> . . . . .	299	<i>Polystichum</i> . . . . .	389
Phyllanthus, <i>Linn.</i> . . . . .	282, 284, 285	Pometia, <i>Forst.</i> . . . . .	408
Phyllodium, <i>Desv.</i> . . . . .	86	Pongamia, <i>Linn.</i> . . . . .	92
<i>Phymatodes</i> . . . . .	395	<i>Pontedera</i> . . . . .	321
Physalis, <i>Linn.</i> . . . . .	217	PONTERACEÆ . . . . .	321
Physurus, <i>L. C. Rich.</i> . . . . .	314	Portulaca, <i>Tourn.</i> . . . . .	23, 401
PHYTOLACCACEÆ . . . . .	250	PORTULACACEÆ . . . . .	23
Pilea, <i>Lindl.</i> . . . . .	259	Posidonia, <i>Kön.</i> . . . . .	333
Piliostigma, <i>Hochst.</i> . . . . .	98	<i>Potamochoa</i> . . . . .	356
Pimpinella, <i>Linn.</i> . . . . .	131	Potamogeton, <i>Linn.</i> . . . . .	333
Piper, <i>Linn.</i> . . . . .	292	Potentilla, <i>Linn.</i> . . . . .	101
PIPERACEÆ . . . . .	292	Poterium, <i>Linn.</i> . . . . .	102
<i>Piptostylis</i> . . . . .	406	Pothomorphe, <i>Miq.</i> . . . . .	292
Pisonia, <i>Plum.</i> . . . . .	245	Pothos, <i>Linn.</i> . . . . .	336
Pistia, <i>Linn.</i> . . . . .	331	Pouzolzia, <i>Gaud.</i> . . . . .	260, 261
PISTIACEÆ . . . . .	331	Premna, <i>Linn.</i> . . . . .	242
Pithecolobium, <i>Mart.</i> . . . . .	100	PRIMULACEÆ . . . . .	172
PISTOSPOREÆ . . . . .	68	Prismatomeris, <i>Thw.</i> . . . . .	154, 421
Pittosporum, <i>Sol.</i> . . . . .	68	<i>Pristidia</i> , <i>Thw.</i> . . . . .	149, 420, 425

	Page		Page
<i>Procris, Comm.</i> . . . . .	260	<i>Rostellularia, Reichen.</i> . . . . .	234, 235
<i>Prosaptia</i> . . . . .	389	<i>Rotala, Linn.</i> . . . . .	122
<i>ProSORUS, Dalz.</i> . . . . .	281	<i>Rothia, Pers.</i> . . . . .	82
PROTEACEÆ . . . . .	250	<i>Rottböllia, Linn.</i> . . . . .	361, 363, 364
<i>Protium, Burm.</i> . . . . .	78, 410	<i>Rottlera, Roxb.</i> . . . . .	272, 274
<i>Pseudanthus, Wight</i> . . . . .	248	<i>Roumea, Poit.</i> . . . . .	18, 400
<i>Pseudarthria, W. and A.</i> . . . . .	87	<i>Rourea, Aubl.</i> . . . . .	80
<i>Psilotrichum, Bl.</i> . . . . .	248	<i>Roxburghia, Dryand.</i> . . . . .	432
<i>Psilotum, Sw.</i> . . . . .	378	<i>Rubia, Tourn.</i> . . . . .	151, 421
<i>Psoralea, Linn.</i> . . . . .	84	RUBIACEÆ . . . . .	137
<i>Psychotria, Linn.</i> . . . . .	147, 148	<i>Rubus, Linn.</i> . . . . .	101
<i>Pteridophyllum, Thw.</i> . . . . .	58, 408	<i>Ruellia, Linn.</i> . . . . .	225, 227
<i>Pteris, Linn.</i> . . . . .	381, 386, 438	<i>Rungia, N. ab Es.</i> . . . . .	234
<i>Pterocarpus, Linn.</i> . . . . .	92	<i>Ruppia, Linn.</i> . . . . .	333
<i>Pteroloma, Desv.</i> . . . . .	86	SABIACEÆ . . . . .	408
<i>Pterospermum, Schreb.</i> . . . . .	30	<i>Saccharum, Linn.</i> . . . . .	368, 369
<i>Pterostigma, Benth.</i> . . . . .	218, 426	<i>Saxolabium, Bl.</i> . . . . .	303, 429
<i>Pterygostachyum, N. ab Es.</i> . . . . .	444	<i>Sageræa, Dalz.</i> . . . . .	6
<i>Pterygota, Endl.</i> . . . . .	29	<i>Sageretia, Brongn.</i> . . . . .	410
<i>Ptilotus, R. Br.</i> . . . . .	248	<i>Salacia, Linn.</i> . . . . .	53, 407
<i>Ptychosperma, Labill.</i> . . . . .	328	<i>Salicornia, Linn.</i> . . . . .	443
<i>Ptyssiglottis, T. And.</i> . . . . .	235	<i>Salmalia, Sch. and Endl.</i> . . . . .	28
<i>Pupalia, Juss.</i> . . . . .	249	<i>Salomonina, Lour.</i> . . . . .	22
<i>Putranjiva, Wall.</i> . . . . .	287	SALSOLACEÆ . . . . .	246
<i>Pycnospora, R. Br.</i> . . . . .	92	<i>Salvadora, Linn.</i> . . . . .	190
<i>Pygeum, Gærtn.</i> . . . . .	102	SALVADORACEÆ . . . . .	190
<i>Pyrenacantha, Hook.</i> . . . . .	290	<i>Samadera, Gærtn.</i> . . . . .	70
<i>Pyrospermum, Miq.</i> . . . . .	409	<i>Samara, Linn.</i> . . . . .	173
<i>Pyrularia, Mich.</i> . . . . .	25	SAMYDACEÆ . . . . .	19
<i>Randia, Houst.</i> . . . . .	159	<i>Sanicula, Tourn.</i> . . . . .	130
RANUNCULACEÆ . . . . .	1	<i>Sansevieria, Thunb.</i> . . . . .	338
<i>Ranunculus, Linn.</i> . . . . .	1	SANTALACEÆ . . . . .	251
<i>Reichenheimia</i> . . . . .	128	SAPINDACEÆ . . . . .	54
<i>Reidia, Wight</i> . . . . .	283	<i>Sapindus, Linn.</i> . . . . .	55, 408
<i>Remirea, Aubl.</i> . . . . .	345	<i>Sapium, Jacq.</i> . . . . .	259
<i>Remusatia</i> . . . . .	336	<i>Sapota, Plum.</i> . . . . .	175
RESTIACEÆ . . . . .	341	SAPOTACEÆ . . . . .	174
<i>Rhabdia, Mart.</i> . . . . .	214	<i>Sarcandra</i> . . . . .	293
RHAMNACEÆ . . . . .	73	<i>Sarcanthus</i> . . . . .	303
<i>Rhamnus, Linn.</i> . . . . .	74, 282, 410	<i>Sarcoelinium, Wight</i> . . . . .	279
<i>Rhamphidia, Lindl.</i> . . . . .	313	<i>Sarcococca, Lindl.</i> . . . . .	290
<i>Rhaphidophora</i> . . . . .	336	<i>Sarcostemma, R. Br.</i> . . . . .	196
<i>Rhaphidospora</i> . . . . .	234	<i>Sarosantha, Korth.</i> . . . . .	41, 405
<i>Rhesa, Walp.</i> . . . . .	72	<i>Satyrium, Sw.</i> . . . . .	308
<i>Rhinacanthus, N. ab Es.</i> . . . . .	234	<i>Sauropus, Bl.</i> . . . . .	284
<i>Rhipsalis, Gærtn.</i> . . . . .	129	<i>Savia, Willd.</i> . . . . .	280
<i>Rhizophora, Linn.</i> . . . . .	120	SAXIFRAGACEÆ . . . . .	130
RHIZOPHORACEÆ . . . . .	120	<i>Scævola, Linn.</i> . . . . .	169
<i>Rhododendron, Linn.</i> . . . . .	170	<i>Scopa, Lindl.</i> . . . . .	288
<i>Rhus</i> . . . . .	59	<i>Schizachyrium</i> . . . . .	368
<i>Rhynchosia, Lour.</i> . . . . .	91, 412, 441	<i>Schizæa, Sm.</i> . . . . .	379
<i>Rhynchospora, Vahl</i> . . . . .	352, 435	SCHIZANDRACEÆ . . . . .	5
<i>Rhytiglossa</i> . . . . .	235	<i>Schizoloma</i> . . . . .	388
<i>Rissoa</i> . . . . .	46	<i>Schizostigma, Arn.</i> . . . . .	139
<i>Rivea, Choisy</i> . . . . .	209	<i>Schleichera, Willd.</i> . . . . .	58
ROSACEÆ . . . . .	100	<i>Schmidelia, Linn.</i> . . . . .	55, 408

	Page		Page
Schumacheria, <i>Vahl</i> . . . . .	4	Spiranthes, <i>L. C. Rich</i> . . . . .	312
Sciaphila, <i>Bl.</i> . . . . .	294	<i>Splizgerbera</i> . . . . .	260
Scindapsus, <i>Schott</i> . . . . .	336	Spodiopogon, <i>Trin.</i> . . . . .	365
Scirpus, <i>Linn.</i> 347, 350, 351, 352, 434		Sponia, <i>Comm.</i> . . . . .	267, 443
Scleria, <i>Linn.</i> . . . . .	353, 435	Sporobolus, <i>R. Br.</i> . . . . .	370
<i>Sclerostylis</i> , <i>Bl.</i> . . . . .	46, 405	Stachytarpha, <i>Vahl</i> . . . . .	241
Scolopia . . . . .	400	STAPHYLEACEÆ . . . . .	71
SCROPHULARIACEÆ . . . . .	217	<i>Stegnogramme</i> . . . . .	383
Scutellaria, <i>Linn.</i> . . . . .	239	Stellaria, <i>Linn.</i> . . . . .	24
Scutia, <i>Comm.</i> . . . . .	75	<i>Stemonoporus</i> , <i>Thw.</i> . . . . .	37, 403
Scutinanthe, <i>Thw.</i> . . . . .	78, 410	<i>Stemonurus</i> , <i>Bl.</i> . . . . .	43, 405
Scyphellandra, <i>Thw.</i> . . . . .	21	<i>Stenochlæna</i> . . . . .	380
Scyphostachys, <i>Thw.</i> . . . . .	157	Stenosiphonium, <i>N. ab Es.</i> . . . . .	225, 228
Secamone, <i>R. Br.</i> . . . . .	195	Stenotaphrum, <i>Trin.</i> . . . . .	361
Selaginella, <i>Spring.</i> . . . . .	377	Stephania, <i>Lour.</i> . . . . .	13
Semecarpus, <i>Linn.</i> . . . . .	75, 410	Sterculia, <i>Linn.</i> . . . . .	29
<i>Senacia</i> . . . . .	68	STERCULIACEÆ . . . . .	28
Senecio, <i>Less.</i> . . . . .	167, 422	Stereospermum, <i>Cham.</i> . . . . .	206
Serissa, <i>Comm.</i> . . . . .	150	<i>Stratiotes</i> . . . . .	332
Serpicula, <i>Linn.</i> . . . . .	123, 331	<i>Stravadium</i> , <i>Juss.</i> . . . . .	119
SESAMEÆ . . . . .	209	Streblus, <i>Lour.</i> . . . . .	264
Sesamum, <i>Linn.</i> . . . . .	209, 442	Streptogyne, <i>Beauv.</i> . . . . .	374
Sesbania, <i>Pers.</i> . . . . .	84, 441	<i>Streptostigma</i> , <i>Thw.</i> . . . . .	56
SESUVIACEÆ . . . . .	23	Striga, <i>Linn.</i> . . . . .	220
Sesuvium, <i>Linn.</i> . . . . .	23	Strobilanthes, <i>Bl.</i> . . . . .	226
<i>Setaria</i> . . . . .	361	Strombosia, <i>Bl.</i> . . . . .	42
Sethia, <i>Kunth</i> . . . . .	53	Strongylocalyx, <i>Bl.</i> . . . . .	116
Shorea, <i>Roxb.</i> . . . . .	36, 402	Strongilodon, <i>Vogel</i> . . . . .	89
Shutereia, <i>Choisy</i> . . . . .	212	Strychnos, <i>Linn.</i> . . . . .	201, 425
Shuteria, <i>W. and A.</i> . . . . .	88	STYLIDIÆ . . . . .	168
Sida, <i>Linn.</i> . . . . .	27	Stylidium, <i>Sw.</i> . . . . .	168, 422
Siegesbeckia, <i>Linn.</i> . . . . .	164	Stylocoryne, <i>Cav.</i> . . . . .	158, 421
SIMARUBEÆ . . . . .	70	Stylosanthes, <i>Sw.</i> . . . . .	84
<i>Sium</i> . . . . .	130	Suæda, <i>Forsk.</i> . . . . .	246
Slegotia, <i>Reich.</i> . . . . .	204	Susum, <i>Bl.</i> . . . . .	340
Smilax, <i>Tourn.</i> . . . . .	338	<i>Sykesia</i> . . . . .	202
Smithia, <i>Ait.</i> . . . . .	85	Symphorema, <i>Roxb.</i> . . . . .	242
Soja, <i>Mæench</i> . . . . .	88	<i>Symphyllia</i> , <i>Baill.</i> . . . . .	270
SOLANACEÆ . . . . .	216	SYMPLOCEÆ . . . . .	184
<i>Solandra</i> . . . . .	200	Symplocos, <i>Jacq.</i> . . . . .	184
Solanum, <i>Tourn.</i> . . . . .	216	Syzygium, <i>Gærtn.</i> . . . . .	116, 417
<i>Solidago</i> . . . . .	162		
Sonchus, <i>Linn.</i> . . . . .	168	Tabernæmontana, <i>Plum.</i> . . . . .	191, 192
Sonerila, <i>Roxb.</i> . . . . .	107	Tacca, <i>Forst.</i> . . . . .	325
Sonneratia, <i>Linn.</i> . . . . .	123	<i>Tæniopsis</i> . . . . .	381
Sophora, <i>Linn.</i> . . . . .	94, 414	Tæniophyllum, <i>Bl.</i> . . . . .	305
Sopubia, <i>Ham.</i> . . . . .	220	Tamarindus, <i>Linn.</i> . . . . .	95
Spathiostemon, <i>Bl.</i> . . . . .	272	TAMARISCINEÆ . . . . .	23
Spathodea, <i>Beauv.</i> . . . . .	206	Tamarix, <i>Linn.</i> . . . . .	401
Spermacoce, <i>Linn.</i> . . . . .	151	Taxotrophis, <i>Bl.</i> . . . . .	264, 427
Sphæranthus, <i>Vaill.</i> . . . . .	162	Tephrosia, <i>Pers.</i> . . . . .	84, 411
<i>Sphærocarya</i> . . . . .	251	TEREBINTHACEÆ . . . . .	75
Sphenoclea, <i>Gærtn.</i> . . . . .	170	Terminalia, <i>Linn.</i> . . . . .	103
SPHENOCLEACEÆ . . . . .	170	Ternströmia, <i>Linn. f.</i> . . . . .	405
<i>Sphragidia</i> , <i>Thw.</i> . . . . .	286	TERNSTRÖMIACEÆ . . . . .	40
Spilantes, <i>Jacq.</i> . . . . .	165	<i>Terpnophyllum</i> , <i>Th.</i> . . . . .	49, 406
Spinifex, <i>Linn.</i> . . . . .	362	Tetracera, <i>Linn.</i> . . . . .	2



	Page		Page
<i>Tetracrypta</i> , Gardn. and Champ.	119	<i>Uralespis</i> , Nutt.	372
<i>Tetramelus</i> , R. Br.	252	<i>Urandra</i> , Thw.	43
<i>Tetranthera</i> , N. ab Es.	254	<i>Uraria</i> , Desv.	85, 411
<i>Teucrium</i> , Linn.	241	<i>Urena</i> , Linn.	25
<i>Thalassia</i> , Soland.	333	<i>Urochloa</i>	358
<i>Thalictrum</i> , Linn.	1	<i>Uropetalum</i> , Gawl.	443
<i>Thamnopteris</i>	383	<i>Urophyllum</i> , Jack.	58
<i>Thespesia</i> , Corr.	27	<i>Urostigma</i> , Gasparr.	264
<i>Thismia</i> , Griff.	325	<i>Urtica</i>	259, 261, 262
<i>Thouarea</i> , Pers.	362	URTICACEÆ	258
<i>Thunbergia</i> , Linn. f.	223	<i>Utricularia</i> , Linn.	171, 423
THYMELACEÆ	250	<i>Uvaria</i> , Linn.	6
<i>Tiaridium</i>	215	VACCINIEÆ	170
TILIACEÆ	31	<i>Vaccinium</i> , Linn.	170
<i>Tiliacora</i> , Colebr.	12	<i>Vaginularia</i> , Fée	381
<i>Timonius</i> , Rumph.	153, 421	<i>Vahlia</i> , Thunb.	130, 418
<i>Tinospora</i> , Miers	12	<i>Valeriana</i> , Neck.	159, 421
<i>Toddalia</i> , Juss.	69	VALERIANACEÆ	159
<i>Torenia</i> , Linn.	219	<i>Vallisneria</i>	331, 332
<i>Tournefortia</i> , Linn.	214	<i>Vanda</i> , R. Br.	303, 305, 429
<i>Toxicarpus</i> , W. and A.	195	<i>Vandellia</i> , Linn.	219
<i>Trachys</i> , Pers.	362	<i>Vanilla</i> , Plum.	311
<i>Tragia</i> , Plum.	270, 271	<i>Vateria</i> , Linn.	36, 403
<i>Trapa</i> , Linn.	124	<i>Vatica</i> , Linn.	404
<i>Trewia</i> , Linn.	272	<i>Ventilago</i> , Gærtn.	74
<i>Trianthema</i> , Sawv.	23	VERBENACEÆ	241
<i>Tribrachys</i> , Champ.	325	<i>Vernonia</i> , Schreb.	160, 421
<i>Tribulus</i> , Tourn.	68	<i>Viburnum</i> , Linn.	136
<i>Trichadenia</i> , Thw.	19	<i>Vicoa</i> , Cass.	164
<i>Trichaurus</i> , Arn.	23, 400	<i>Vigna</i> , Savi	90
<i>Trichelostylis</i>	348	<i>Villebrunia</i> , Gaud.	261
<i>Trichodesma</i> , R. Br.	216	<i>Viola</i> , Linn.	20
<i>Trichomanes</i> , Linn.	397	VIOLACEÆ	20
<i>Trichopodium</i> , Lindl.	291, 443	<i>Visania</i>	188
<i>Trichopus</i> , Gærtn.	291	<i>Viscum</i> , Tourn.	136, 418
<i>Trichosanthes</i> , Linn.	126	<i>Vitex</i> , Linn.	244
<i>Tridens</i>	372	<i>Vitis</i> , Linn.	62, 63
<i>Trigonostomon</i> , Bl.	277, 428	<i>Vittaria</i> , Sm.	381, 438
<i>Trimeriza</i>	291	<i>Volkameria</i>	243
<i>Tripogon</i> , Roth	374	Wahlenbergia, Schr.	169
<i>Tripterospermum</i> , Bl.	204	<i>Walkera</i>	71
<i>Tristicha</i>	223	<i>Walsura</i> , Roxb.	61
<i>Triticum</i> , Linn.	376	<i>Waltheria</i> , Linn.	30
<i>Triumfetta</i> , Plum.	31, 401, 444	<i>Webera</i>	158
TRIURIDACEÆ	294	<i>Wedelia</i> , Jacq.	165
<i>Trophis</i> , Willd.	262, 264	<i>Wendlandia</i> , Bartl.	159
<i>Tropidia</i> , Lindl.	314	<i>Wikströmia</i> , Endl.	250
<i>Turpinia</i> , Vent.	71	<i>Willughbeia</i> , Roxb.	191
<i>Tylophora</i> , R. Br.	196, 424	<i>Winchia</i>	191
<i>Typha</i> , Linn.	331	<i>Wissadula</i> , Med.	27
TYPHACEÆ	331	<i>Withania</i> , Duval	217
<i>Typhonium</i> , Schott	334	<i>Wollastonia</i> , DC.	165
<i>Ulmus</i> , Linn.	267	<i>Wormia</i> , Rottb.	40
UMBELLIFERÆ	130	<i>Wrightia</i> , R. Br.	193, 424
<i>Uncaria</i> , Schreb.	138, 418		
<i>Unona</i> , Linn.	9, 398		

	Page		Page
Xanthium, <i>Tourn.</i> . . . . .	164	Zanonia, <i>Linn.</i> . . . . .	124, 442
Xanthochymus, <i>Roxb.</i> . . . . .	46	Zehneria, <i>Endl.</i> . . . . .	125
Xanthophyllum, <i>Roxb.</i> . . . . .	23, 400	Zeuxine, <i>Lindl.</i> . . . . .	312
XANTHOXYLACEÆ . . . . .	69	Zingiber, <i>Gærtn.</i> . . . . .	315, 430
Xanthoxylon, <i>Linn.</i> . . . . .	69, 409	ZINGIBERACEÆ . . . . .	315
Ximenesia, <i>Cav.</i> . . . . .	422	Zizyphus, <i>Tourn.</i> . . . . .	73
Xylocarpus, <i>Kön.</i> . . . . .	61	Zornia, <i>Gmel.</i> . . . . .	84
Xylophia, <i>Linn.</i> . . . . .	9, 398	Zosterostylis, <i>Bl.</i> . . . . .	312
XYRIDACEÆ . . . . .	340	Zoysia, <i>Willd.</i> . . . . .	370
Xyris, <i>Linn.</i> . . . . .	340	ZYGOPHYLLÆ . . . . .	68
Youngia, <i>Cass.</i> . . . . .	168	<i>Zygospermum</i> , <i>Thw.</i> . . . . .	281

THE END.