

No. 3.—*On the Mammals and Winter Birds of East Florida, with an Examination of certain assumed Specific Characters in Birds, and a Sketch of the Bird-Faunæ of Eastern North America.* By J. A. ALLEN.

INTRODUCTION.

THE present paper embraces five more or less distinct parts. The first consists of introductory remarks respecting the topographical, climatic, and faunal features of that part of the peninsula of Florida usually known as East Florida. The second is an annotated list of the mammals of this region. The third is devoted to a consideration of individual, seasonal, age and geographical variation among birds, with reference to certain characters commonly assumed to be specific. The fourth contains a list of the winter birds of East Florida, with field and revisionary notes. The fifth is given to an examination of the geographical distribution of the birds and mammals (more particularly of the birds) of Eastern North America, in which is considered the number of the natural faunæ of this region, their distinctive features and their boundaries.

The enumeration of the mammals and birds, forming Parts II and IV, is based partly on my own observations and partly on notes kindly furnished me by Messrs. C. J. Maynard and G. A. Boardman. These observations may be considered as equivalent collectively to the labors of a single observer constantly in the field for at least four or five winters.

My own observations were made during a three months' exploration of the country bordering the St. John's River, between Jacksonville and Enterprise, in the winter of 1868 and 1869, under the auspices of the Museum of Comparative Zoölogy. The greater part of January was passed at Jacksonville, where I also spent the last week of March. Five weeks were also passed in the vicinity of Enterprise, and the balance of the time at various intermediate points.

Mr. Maynard's explorations were made during the same winter, mainly in portions of the country unvisited by myself, a large part of his collection coming from the Upper St. John's and Indian Rivers. He also spent several weeks at Dummitt's, twenty miles south of New Smyrna. During most of December and January he collected

in the vicinity of Jacksonville, at which point one of his assistants, Mr. Charles Thurston, remained during April and a portion of May, collecting, among other things, the later arriving birds. Nearly all the birds and mammals collected by these gentlemen, and by Mr. J. F. Le-Baron, a third member of Mr. Maynard's party, have been added to the collection of the Museum of Comparative Zoölogy, and are accompanied by measurements carefully taken before skinning.

Mr. Boardman's observations were continued through three successive winters, during which he spent considerable time at the following points: St. Augustine and Fernandina on the coast, Jacksonville, Green-Cove-Springs and Enterprise on the St. John's River. Although the numerous specimens he collected at these and intermediate points were presented by him to the Smithsonian Institution, I am indebted to him for an annotated manuscript list of the species he met with. I am also indebted to the Rev. Thomas Marcy, who accompanied me on my Florida trip, for valuable assistance in collecting, and to Mr. J. E. Brundage for similar aid.

Having made use of the reports of previous visitors on the faunæ of this region, the following lists are believed to embrace all the species of mammals thus far known from East Florida, and all the birds regularly present in winter, of nearly all of which I have examined specimens from Florida. A few other birds not included in my list doubtless occasionally visit this region from the North, and others may linger here which usually pass the winter further south. In order to increase the value of the bird list as a faunal record, those species known to be resident throughout the year have been indicated by an asterisk (*), and those known only as winter visitors by an obelisk (†). The date of the first appearance of the strictly spring visitors is also noted, so far as such arrivals were observed.

The specimens on which the investigations detailed in Part III are based, as well as the revisionary notes of Parts II and IV, are mainly those of the Museum of Comparative Zoölogy, which embrace, among others, nearly a thousand specimens of birds from Florida.*

The topics discussed in Part III, namely, individual and climatic variation, necessarily involve the question of the nature of species, as well as the validity of various diagnostic characters. Many details

* I have also made use of measurements, taken by Mr. Wm. Brewster and Mr. C. J. Maynard, of hundreds of specimens not in the collection of the Museum.

in reference to these variations are given in this part, but a large proportion are recorded in the general and revisionary notes of Part IV. The conclusions arrived at, it may be here premised, are mainly the following: (1.) That the majority of nominal species originate in two principal sources of error; namely, (*a*) an imperfect knowledge of the extent and character of individual variation, and (*b*) of geographical variation. (2.) That this imperfect knowledge is mainly due to the neglect of zoölogists to study with sufficient care the common species of their respective countries, whence has arisen a faulty method of investigation and erroneous ideas respecting species and specific characters. (3.) Instead of the method at present pursued by a large school of descriptive naturalists — the analytic, or the search for differences — being the proper one, that synthesis should be duly combined with analysis, and that general principles should be sought as well as new forms, or so-called “new species” and “new genera.” (4.) It is claimed that nothing is to be gained by giving binomial names to climatic or other forms, in cases where, however considerable the differences between them may be, a complete transition from the one to the other can be traced in specimens from intermediate localities, notwithstanding the plea sometimes urged that their use affords “convenient handles to facts.”

In accordance with such views a partial revision of the species of certain groups is incidentally attempted in Part IV, more especially of the *Icteridæ*, the raptorial birds, and the genera *Parus*, *Turdus*, *Passerculus*, etc.

PART I.

The Topographical, Climatic, and Faunal Characteristics of East Florida.

No part of the Florida Peninsula, as is well known, is much elevated above the level of the sea, the greater portion being extremely low and large areas swampy. The surface is slightly undulating, but the higher ridges rarely attain a height of more than fifty or seventy-five feet, and the highest eminence is less than two hundred. A large part of Northern Florida, including what is usually termed East and West Florida, is covered with open pine forests, constituting the so-called “pine barrens.” These barrens frequently rise into dry knolls, but they likewise embrace considerable tracts that are so low as to be more or less submerged during a portion of the year, especially in wet

seasons; they are also interspersed with cypress swamps of varying extent. Such swamps usually border the St. John's on its upper course, sometimes extending back from the river for several miles. Other portions of the low grounds support a mixed forest of live-oak, water-oak, elm, bitter-nut hickory, maple, laurel, sweet gum, etc., with a more or less dense undergrowth, such forests forming the so-called "hummocks." Some portions of these forests are swampy; others are dry, and slightly elevated. The saw and dwarf palmettos (*Sabal serrulata* R. & S. and *S. Adansonii* Guerns.) frequently render the former difficult to penetrate, and extensive groves of the cabbage palm (*Chamærops palmetto* Michx.; *Sabal palmetto* R. & S.) here and there occupy the banks of the streams. At intervals in the pine barrens extensive thickets of low trees and thickly growing shrubs are met with, which are exceedingly difficult to enter, and are appropriately termed "scrubs." Each of these kinds of country, as would be naturally expected, forms the favorite haunt of certain species of birds and mammals, the grassy or open pineries being frequented by some that rarely visit the swamps and hummocks, and the latter by others that rarely visit the open pineries. The extensive savannas which occur along the upper portion of the St. John's River and elsewhere form the favorite haunts of numerous wading birds; and the numerous lakes are congenial to the swimming birds.

East Florida hence differs but little in its general character from the lower portions of Georgia and the low lands of the coast of South Carolina. The trees, especially of the hummocks and swampy forests, are usually covered with the pendant *Tillandsia usnoides*, or "Spanish moss," and the abundance of epiphytic orchids and other plants, as well as the palms, clearly indicates the subtropical and peculiar character of the climate.

From the great extent in latitude of the Florida peninsula — from 25° to 31° , or about four hundred miles — considerable differences necessarily exist between the fauna and flora of the northern and southern portions. Although the change in these features from the north southward is more or less gradual, it seems to be appreciably greater near Lake George than elsewhere. At this point so well-marked a change occurs in the vegetation as to attract the attention of unscientific observers, and a corresponding change in the fauna is readily traced. Above Lake George the general aspect of both the flora and fauna is decidedly more southern than it is below the lake. The

boundary between the Floridian and Louisianian faunæ and floræ, it would hence seem, may be properly regarded as passing near this point, the portion of the State to the southward being alone properly Floridian, the northern resembling more the Louisianian type.*

As already observed, Florida, from its excessively marshy character, is pre-eminently suited to the wants of the grallatorial birds. Immense numbers of the heron tribe hence make it their permanent home, while it is the favorite winter resort of numerous species of *Grallæ* that pass the breeding season far to the northward. Ibises and egrets abound in its swamps and savannas, forming at all times, by their numbers and showy plumage, a characteristic feature of the fauna. In winter the abundance of snipe and other species of *Grallæ* and ducks render it at that season a sportsman's paradise. Florida hence attracts great numbers of sportsmen in winter, through whose reckless and often wanton waste of life the water-fowl, especially of late years, are annually decimated.

The summer bird fauna of Florida is probably not better represented in species than that of the temperate parts of the continent generally; but this State being the winter resort of numerous species of sparrows and warblers, and of those smaller land birds generally that pass the summer in much higher latitudes, its winter bird fauna, as compared with that of the Northern States, is extremely rich. In New England the number of species of birds that can be regarded as "common" in winter does not exceed fifteen,† but in Florida at that season at least five times that number can be so regarded. This, however, accords with a general law of distribution in respect to the relative number of species found at different points in latitude from the arctic zone southward, the number increasing in proportion to the decrease of the latitude, or with the increase of the mean temperature. In winter, through the southward migration of many species, the minimum number of species which in summer is characteristic of the arctic zone is carried down nearly to the Northern States, there being a marked decrease from summer to winter as far south as the warm temperate or subtropic belt; within the tropics, on the contrary, the number of species is far greater in winter than in summer, through the temporary influx of species from colder regions.

* For a further definition of the Floridian bird fauna, as distinguished from the Louisianian, see beyond, Part V.

† See American Naturalist, Vol. I, p. 47, March, 1867.

In consequence of the subtropical character of the climate of Florida certain peculiarities occur in respect to the development of vegetation at the vernal period, and in the time of breeding of the resident birds, that seem in this connection worthy of record. The mildness of the winter climate is such that the verdure of the forests is to a greater or less degree perennial, severe frosts being of rare occurrence. Some of the early flowering trees, such as the maples, ashes, and elms, begin to bloom and to gradually unfold their leaves early in January. Although the forest trees in general put forth their leaves in February, and a few have acquired their full summer dress by the 1st of March, their development is slow and irregular. I observed peach-trees in flower at the same locality (Jacksonville) in January and in April; and the flowering period of some of the forest trees is nearly as protracted. The development of vegetation is hence as great during a single week in May, in New England, as during any four weeks in February and March, in Florida.

A similar irregularity is observed in respect to the pairing and breeding of the resident birds. Some of the rapacious species, as the fish-hawk and the white-headed eagle, commence incubation in January, and, as I have been informed, occasionally in December; other members of the same species delay breeding till February or March. The great blue heron and the egrets nest in February, as do also the courlans, several of the hawks, the sandhill crane, the wood-duck and the blue-bird; the mocking-bird and other resident song-birds, in March and April.

In the Northern States the vivacity of the birds during the pairing season is as much greater than it is in Florida as is the rapidity of the development in vegetation. In spring at the North the woods, the fields, and the hedgerows are ever vocal with bird music; but in Florida no such outburst of song marks the arrival of the vernal season. The brown thrush, the blue-bird, the cat-bird, the towhee, and the various kinds of sparrows that are common in the breeding season to both New England and Florida, seem to lose at the latter locality the vivacity which characterizes them at the North, their attempts at song being listless and feeble. The songs of some are also much abbreviated, and so different from what they are at the North as to be sometimes scarcely recognizable as proceeding from the same species. Even the mocking-bird sings far less than in the Middle States, and

with much less power. Such at least is the general fact as indicated by my own limited experience in Florida, which accords, I find, with that of various other observers.

In recounting the faunal peculiarities of Florida it is necessary to allude further to a few facts that will be more fully presented in the following chapters, namely, the differences which distinguish the Florida representatives of species that have a wide distribution to the northward from the northern ones. It has for some time been well known that a difference in size in birds and mammals usually accompanies differences of locality in respect to latitude and elevation. Other differences, however, are found to accompany these with considerable uniformity; namely, a relative increase in the length or general size of the bill, and an increase in the intensity of the general color of the plumage.* Florida birds, in short, usually differ considerably in these respects from their New England conspecific representatives; so much so, indeed, that in many cases the majority of ornithologists would probably regard the two forms as distinct species, though few of them have as yet been specifically separated.

Hence not only do birds of the same species living at distant points differ considerably in size, color, and other features, but also in their habits, notes, and songs. With the decrease in size to the southward there seems to be a corresponding decrease in vivacity, — a fact which accords with the general law of the distribution of the higher forms of life in the temperate latitudes. Although a few structurally high types are, from certain peculiarities of their conformation, necessarily tropical, the highest races of men, whether considered physically, intellectually, or morally, are inhabitants of a medium climate, and gradually decline in rank both to the northward and southward from this favored region, animal and vegetable life reaching, as a whole, its highest manifestation in the temperate latitudes. The excessive variety of forms within the tropics mainly results from the addition of those of comparatively low or medium grades, only a few of the exclusively tropical forms being of absolutely high rank. Generally, too, the forms to be properly regarded as temperate are represented in the tropics by only their lower members, while, conversely, many of the higher types of the tropics are really cosmopolitan.

* See Annual Report of the Mus. Comp. Zool., 1869, p. 16.

PART II.

List of the Mammals of East Florida, with Annotations.

FELIDÆ.

1. *Felis concolor* Linné. PANTHER.

Not very unfrequent in the more unsettled parts of the State. I saw several hunter's skins of it at Jacksonville, said to have been taken up the river.

2. *Lynx rufus* Rafinesque. BAY LYNX.

Abundant. Especially numerous on the Upper St. John's and Indian Rivers, according to Mr. Maynard and others.

CANIDÆ.

3. *Canis lupus* Linné. GRAY WOLF.

Canis lupus LINNÉ, Syst. Nat., I, 58, 1767. — ALLEN, Bull. Mus. Comp. Zool., I, 154, October, 1869

Canis lupus, occidentalis RICH., Fauna Bor. Amer., I, 60, 1829.

Canis occidentalis et var. BAIRD, Mam. N. Amer., 104, 111, 113, 1857.

Not numerous. They were described to me as being very dark colored, or black.* This account tends to confirm the statement of Audubon and Bachman in respect to this point.† After citing the comparative frequency of this form of the common wolf in Kentucky, and in several of the Southern Atlantic and Gulf States, as compared with its occurrence in regions more to the northward and westward, they observe: "The varieties with more or less of black continue to increase, as we proceed farther to the south; and in Florida the prevailing color of the wolves is black."‡

4. *Vulpes virginianus* Richardson. GRAY FOX.

Canis virginianus ERXL., Syst. Reg. Anim., 567, 1777. — "SCHREBER, Säugeth., III, 361, pl. xcii, 1778."

Canis cinereo-argentatus ERXL., Syst. Reg. Anim., 567, 1778. — "SCHREBER, Säugeth., 360, pl. xcii." — GODMAN, Am. Nat. Hist., I, 280, 1826.

Canis griseus BODD., Elenchus Anim., I, 77, 1784.

* Since writing the above, I have received a letter from Mr. G. A. Boardman, of Milltown, Me., in which he also refers to the dark color of the Florida wolves.

† Quad. N. Amer., Vol. II, p. 130.

‡ Respecting the distribution of the different color races of the common wolf in North America, see my paper on the Mammals of Massachusetts, Bulletin Mus. Comp. Zool., Vol. I, p. 156, 1869.

Canis (Vulpes) virginianus RICH., Faun. Bor. Am., I, 96, 1829.

Vulpes virginianus DEKAY, New York Fauna, I, 45, pl. vii, fig. 2, 1842. —

AUD. & BACH., Quad. N. Am., I, 162, pl. xxi, 1849.

Vulpes (Urocyon) virginianus BAIRD, Mam. N. Am., 138, 1857.

Common.

MUSTELIDÆ.

5. *Putorius lutreolus* Cuvier. MINK.

Mustela lutreola LINN., Syst. Nat., 66, 1766.

Putorius lutreolus CUV., Règ. Anim., I, 148, 1817. — ALLEN, Bull. Mus. Comp. Zoöl., I, 175, October, 1869.

Putorius vison GAPPER, Zoöl. Journ., V, 202, 1830.

Putorius nigrescens AUD. & BACH., Quad. N. Am., III, 104, pl. cxxiv, 1853.

"Not common." — Boardman. I did not meet with it. It is well known to be common, however, in the adjoining States. Audubon and Bachman speak of it as being very numerous in the rice-fields of South Carolina

6. *Lutra canadensis* Sabine. OTTER.

Abundant. Its fur, however, is of little value, compared with that of northern specimens, and the animal is hence not much hunted.

7. *Mephitis mephitis* Baird. COMMON SKUNK.

Viverra mephitis SHAW, Mus. Lever., 172, 1792. — IBID., Gen. Zoöl., I, 390, 1809.

Mephitis chinga TIEDEM., Zoöl., 362, 1808.

Mustela (Mephitis) americana DESM., Mamm., I, 186, 1820.

Mustela varians GRAY, Charlesw. Mag. Nat. Hist., I, 581, 1837.

"*Mustela mesomelas* LICHT., Darst. Säugeth., I, fig. 2." — GEOFF. ST. HIL., Voy. de la Venus, Zool., I, 133, 1855. — MAX. ZU WIED, Archiv für Naturgesch., XXVII, 218, 1861. — BAIRD, Mam. N. Am., 199, 1857.

Mephitis macroura AUD. & BACH., Quad. N. Am., III, 11, 1853.

Mephitis mephitis BAIRD, Mam. N. Amer., 195, 1857. — ALLEN, Bull. Mus. Com. Zoöl., I, 178, October, 1869.

Mephitis occidentalis BAIRD, Mam. N. Amer., 194, 1857.

Common on the Lower St. John's, but, according to Mr. Maynard, quite unknown on the Indian River.

8. *Mephitis bicolor* Gray. LITTLE STRIPED SKUNK.

Mephitis bicolor GRAY, Charlesw. Mag. Nat. Hist., I, 581, 1837. — BAIRD, Mam. N. Amer., 196, 1857.

Mephitis zorrilla LICHT., Abhand. Akad. Wiss. Berlin, for 1836, 281, 1838. — AUD. & BACH., Quad. N. Amer., III, 276, 1854.

Mephitis interrupta LICHT., Abhand. Akad. Wiss. Berlin, for 1836, 283, 1838.

This beautiful little animal was obtained by Mr. C. J. Maynard at Captain Dummitt's, where it was said to be common in the scrub. Mr. Maynard says they are domesticated and used there as cats, the odor glands being removed when the animals are young; they become very tame and are quite efficient in destroying the mice (*Hesperomys* sp.) that infest the houses. I am not aware that this animal has been reported before from any point east of the Mississippi River. It has been recently ascertained to extend northward in the interior as far as Central Iowa.*

URSIDÆ.

9. *Procyon lotor* Storr. RACCOON.

Ursus lotor LINNÉ, Syst. Nat., 48, 1758.

Procyon lotor BAIRD, Mam. N. Amer., 209, 1857. — ALLEN, Bull. Mus. Comp. Zoöl., I, 181, October, 1869.

Procyon Hernandezii WAGLER, Isis, XXIV, 514, 1831. — BAIRD, Mam. N. Amer., 212, 1857. — IBID., U. S. and Mex. Bound. Surv., II, Mam., 22, 1859.

Exceedingly numerous.

10. *Ursus arctos* Linné. COMMON BEAR.

Ursus arctos LINNÉ, Syst. Nat., 69, 1766. — CUVIER, Règ. Anim., I, 142, 1817.

— BLAINVILLE. — MIDDENDORFF, Sibirische Reise, II, ii, 1854. — GRAY, Proc. London Zoöl. Soc., 1864, 682. — ALLEN, Bull. Mus. Comp. Zoöl., I, 184, October, 1869.

Ursus americanus PALLAS, Spicilegium Zoöl., XIV, 6, 1780. — GMELIN, Syst. Nat., I, 101, 1788. — RICHARDSON, Faun. Bor. Amer., I, 14, 1829. — AUD. & BACH., Quad. N. Amer., III, 187, 1853. — MAX. ZU WIED & MAYER, Verhandl. Akad. der Naturf., XXVI, i, 33, 1857. — BAIRD, Mam. N. Amer., 225, 1857.

Ursus (Euarctos) americanus GRAY, Proc. Lond. Zoöl. Soc. 1864, 692.

Ursus horribilis ORD, "Guthrie's Geog., 2d Amer. ed., II, 291, 299, 1815." — SAY, Long's Exped., II, 53, 1823. — BAIRD, Mam. N. Amer. 219.

Ursus horribilis, var. *horriaceus*, BAIRD, U. S. & Mex. Bound. Survey, Rep., II, Mam., 24, 1859.

Ursus cinereus DESM., Mam., I, 164, 1820.

Ursus (Danis) cinereus GRAY, Proc. Lond. Zoöl. Soc., 1864, 690.

Ursus ferox RICHARDSON, Faun. Bor. Amer., I, 24, 1829. — MAX. ZU WIED, Reise in das innere Nord Amer., I, 488, 1839. — MAX. ZU WIED & MAYER, Verhandl. Akad. der Naturforsch., XXVI, 39.

Ursus cinnamomeus BAIRD, U. S. & Mex. Bound. Survey Rep., II, Mam., 29.

* See H. W. Parker, in Amer. Nat., Vol. IV, 376, August, 1870.

Numerous and often troublesome, occasionally destroying swine, of which they are exceedingly fond. Judging from their tracks in the swamps, they must not only be exceedingly numerous, but some of them of enormous size. The several skins seen by me were all intensely black.*

CERVIDÆ.

11. *Cariacus virginianus* Gray. VIRGINIA DEER.

Cervus virginianus BODDERT, Elench. Animal., I, 136, 1784. — GMELIN, SCHREBER, DESMEREST, AUD. & BACH., BAIRD, &c.

Cariacus virginianus GRAY, Cat. of Bones in Brit. Mus., 266, 1862.

Abundant almost everywhere. Not so numerous along the Lower St. John's as in the more unsettled districts further south. As remarked by Professor Baird, the Florida deer are considerably smaller than those of the Northern States; so much so that it is a fact of common observation.

MANATIDÆ.

12. *Trichechus manatus* Linné. MANATEE.

Trichechus manatus LINNÉ, Syst. Nat., I, 34, 1758.

"*Manatus australis* TILESIIUS, Jahrb. der Naturg., I, 23." — GRAY, Cat. Seals and Whales, 358, 1866. — MURRAY, Geog. Distr. Mam., 202, 1866.

Manatus amer' anus DESM., Mam., 507, 1822.

Manatus latirostris HARLAN, Journ. Phil. Acad. Nat. Sci., III, 390, pl. xii, fig. 1-3, 1824. — IBID., Faun. Amer., 277, 1825.

I learn from Mr. Maynard that the manatee is still quite common in Indian River, where they are often caught in nets. They come into the river at night to feed on the mangrove bushes. Mr. Maynard did not meet with them in Mesquito Lagoon, which he traversed nearly its whole length, and he thinks they do not occur there.

The manatees of America and Africa seem to be very closely allied, and to number at most but two species. Those of the same species also appear to be exceedingly variable in their osteological characters. Dr. J. E. Gray,

* In my recent paper in this Bulletin, cited above, I have discussed the mutual relationship of the numerous supposed species of land bears of the northern hemisphere. The close affinity between the bears of Northwestern America and Northeastern Asia is especially noticed; but at that time I was not aware that Temminck, in the Fauna Japonica, had referred the large land bear of Japan to the *U. ferox* of authors, or to the so-called "grizzly bear" of Western America. This indicates the very close affinity, in this author's opinion, of the Japan and American bears.

in a valuable paper entitled "On the Species of Manatees (*Manatus*), and on the Difficulty of distinguishing such Species by Osteological Characters,"* states that he finds the African and American species are distinguished by only a single character, — the absence of the nasal bones in the African species. Concerning the individual variation in the skulls of the two species, he observes as follows: "When Cuvier had a skull of the American and one of the African Manatee, he gave eight characters by which the African skull could be known from the American. Now we have a series of skulls of each kind, we find that not one of these characters will separate the skulls of the two countries from one another. Indeed, the skulls of each kind are so variable that, after having them laid out before me for two or three days, studying them every now and then, and inducing two proficient in the study of bones, and in observing minute characters, to give me their assistance, we came to the conclusion that we believed there was no character, common to all the skulls of each kind, which could be used to separate them. As a proof of the difficulty of so doing, I may state that there was one skull in the series which had been long in the collection, and had been received without any habitat, and neither of the three could decide to which of the series this skull should be referred; and it was not until I accidentally observed the character, derived from the absence of the nasal bones in the African kind, that this question could be settled."

Having myself been struck with the variability of osteological as well as external characters in individuals of the same species, in both birds and mammals, — a matter to which I have already often called attention, and the consideration of which occupies a considerable portion of Part III of the present paper, — I can hardly refrain, in this connection, from citing further the judicious remarks of Dr. Gray on this point. "The examination," he says, "of a large series of skulls of the bears (*Ursus*) and *Paradoxuri*, shows how difficult it is to distinguish species by the study of the skulls alone. Thus, when we have a series of skulls of bears from different localities, which, from their external form and habits, are known to be distinct species, it is easy to say which is the skull of *U. thibetanus*, *U. syriacus*, *U. arctos*, *U. cinereus*, and *U. americanus*, when we have the habitat marked on each; but the true test of the power of distinguishing the one from the other is to determine to what species a skull belongs, of which we have no information as to its origin; and we have several skulls in the British Museum under these circumstances, and I cannot, with the best assistance at my command, determine to which species they ought to be referred. And it is the same with the *Paradoxuri*." "If this is the case with the skulls," he continues, "how must the difficulty of distinguishing species with certainty be increased when we have only fossil bones, which are generally more or less imperfect,

*Ann. and Mag. Nat. Hist., 3d Ser., Vol. XV, pp. 130 - 139, 1865.

to examine and compare, or of which only a limited number of examples are to be obtained and compared? They [the skulls] vary in most genera much more than was expected, before series of the skulls of each species were collected and compared."

These observations by Dr. Gray are fully confirmed by my own studies; and I hence believe that, as the number of specimens of different species increases in our museums, many species now believed to be valid will be found to rest merely on individual characters.

VESPERTILIONIDÆ.

13. *Lasiurus noveboracensis* Gray. RED BAT.

Vespertilio noveboracensis ERXL., Syst. Règ. Anim., 135, 1717.

Vespertilio lasiurus GMEL., Syst. Nat., 1788.

Vespertilio rubellus PAL. DE BEAUV., Cat. Peale's Mus., 1796.

? *Vespertilio cinereus* PAL. DE BEAUV., Ibid.

? *Vespertilio pruinosus* SAY, Long's Exped., 67, 1823. — RICH., Faun. Bor. Am., I, 1, 1829.

Taphozous rufus HARLAN, Faun. Amer., 23, 1825.

Lasiurus rufus GRAY, List Mam. Brit. Mus., 32, 1842.

Lasiurus noveboracensis TOMES, Proc. Lond. Zool. Soc., 1857, 34.

? *Lasiurus pruinosus* TOMES, Ibid., 37.

Lasiurus noveboracensis H. ALLEN, Mon. N. Am. Bats, 15, 1864. — J. A. ALLEN, Bull. Mus. Comp. Zool., I, 207, 1869.

? *Lasiurus cinereus* H. ALLEN, Mon. N. Am. Bats, 21.

Common. All of the several specimens obtained, both by myself and Mr. Maynard, are of a deep cherry red, with but a slight skirting of ash, and are uniformly much darker or deeper colored than any I have seen from the Northern States. All examined (nine specimens) were males.

14. *Scotophilus fuscus* H. Allen. CAROLINA BAT.

Vespertilio fuscus PAL. DE BEAUV., Cat. Peale's Mus., 14, 1796. — LECONTE, Proc. Phil. Acad. Nat. Sci., VII, 437, 1855.

Vespertilio carolinensis GEOFF. ST. HIL., Ann. du Mus., VIII, 193, 1806, pl. xlvii, fig. 7. — HARLAN, North Am. Jour. Geol. & Nat. Sci., I, 218, 1831 — LECONTE, Proc. Phil. Acad. Nat. Sci., VII, 437.

Vespertilio arcuatus SAY, Long's Exped., 167, 1823.

Vespertilio phaiops RAF., Amer. Month. Mag., 445, 1818.

Vespertilio ursinus TEMM., Mam., II, 234, 1835.

Scotophilus fuscus H. ALLEN, Mon. N. Am. Bats, 31, 1864.

Scotophilus carolinensis H. ALLEN, Ibid., 28.

Common. Several specimens taken.

15. *Scotophilus georgianus* H. Allen. GEORGIA BAT.

Scotophilus georgianus H. ALLEN, Mon. N. Am. Bats, 35, 1864, *nec. syn.* —
J. A. ALLEN, Bull. Mus. Comp. Zool., No. 8, 1809.

This species doubtless occurs in Florida, at least in the northern part, since the capture of specimens at different localities in Georgia and at New Orleans is on record.*

16. *Nycticejus crepuscularis* H. Allen.

Vespertilio crepuscularis LECONTE, McMurtrie's Cuv. An. King., I, 432, 1831.
IBID., Proc. Phil. Acad. Nat. Sci., VII, 438, 1855.
Nycticejus crepuscularis H. ALLEN, Mon. N. Am. Bats, 12, 1864.

A specimen collected by Mr. Maynard at Jacksonville, in January, but afterwards lost, I refer from his measurements and description of it to this species. There is also a specimen (No. 731) in the Museum of Comparative Zoölogy, collected in Florida by Mr. Chas. Belknap.

17. *Corynorhinus macrotis* H. Allen. BIG-EARED BAT.

Plecotus macrotis LECONTE, McMurtrie's Cuv. An. King., I, 431, 1831.
Plecotis LeContei COOPER, Ann. Lyc. Nat. Hist., IV, 12, 1837.
Synotis macrotis H. ALLEN, Mon. N. Am. Bats, 63, 1864.
Corynorhinus macrotis H. ALLEN, Proc. Phil. Acad. Nat. Sci., XVII, 173, Aug. 1865.

A specimen of this species from Micanopy, Florida, collected by Dr. Bean, is cited by Dr. Allen.† This Southern species ranges northward along the coast nearly or quite to the Middle States, it being comparatively common, according to authors, in South Carolina.

NOCTILIONIDÆ.

18. *Nyctinomus nasutus* Tomes.

Molossus nasutus SPIX, Sim. et Vesp. Bras., 60, pl. xxxv, fig. 7, 1823.
Nyctinomus nasutus TOMES, Proc. Lond. Zool. Soc., 1861, 68. — H. ALLEN, Mon. N. Am. Bats, 7, 1867.

This widely distributed southern species should unquestionably be included among the mammals of Florida. It has been reported from Texas, Louisiana, South Carolina, and the West Indies,‡ as well as from South America, as far south even as Buenos Ayres.§ Specimens in

* Dr. H. Allen, Monograph of North American Bats, p. 38.

† Ibid., p. 55.

‡ Ibid., p. 10.

§ Tomes, Proc. Lond. Zool. Soc., 1861, p. 68.

the Museum of Comparative Zoölogy from Hayti, collected by Mr. P. R. Uhler, have been identified by Dr. Harrison Allen as of this species.

19. *Megadermatidæ* *Sp.?*

A large species of bat was noticed by both Mr. Maynard and myself, but as it always flew very high, neither of us obtained it. It was very much larger than any other species yet described from the United States, and is doubtless a West Indian form; probably a species of *Megadermatidæ*.

SORECIDÆ.

20. *Blarina brevicauda* *Baird*. MOLE SHREW.

Sorex brevicaudus SAY, Long's Exped., I, 164, 1862-63.

Sorex parvus SAY, *Ibid.*, 164.

Sorex talpoides GAPPER, *Zoöl. Journ.*, V, 208, pl. viii, 1830.

Sorex carolinensis BACH., *Journ. Phil. Acad. Nat. Sci.*, VII, 366, pl. xxvi, fig. 3, 1837.

Sorex cinereus BACH., *Ibid.*, 373, fig. 3.

Sorex Dekayi BACH., *Ibid.*, 377, fig. 4.

Corsira (*Blarina*) *talpoides* GRAY, *Proc. Lond. Zoöl. Soc.*, V, 124, 1837.

Blarina brevicauda BAIRD, *Mam. N. Am.*, 42, pl. xxx, fig. 6, 1857.—ALLEN, *Bull. Mus. Comp. Zoöl.*, I, 212, October, 1869.

Blarina talpoides BAIRD, *Mam. N. Am.*, 37, pl. xxx, fig. 5.

Blarina carolinensis BAIRD, *Ibid.*, 45, pl. xxx, fig. 8.

Blarina cinerea BAIRD, *Ibid.*, 48, pl. xxx, figs. 9 and 10, young.

Blarina exilipes BAIRD, *Ibid.*, 51, pl. xxviii, young.

Blarina Berlandieri BAIRD, *Ibid.*, 53, pl. xxviii, young.

A single specimen of *Blarina* from Indian River, Florida, collected by Mr. G. Wurdemann, is mentioned under "*Blarina cinerea*," by Professor Baird, as having been received at the Smithsonian Institution.* While it may be of a species distinct from *B. brevicauda*, it seems more probable that it is the young of that species, as I have elsewhere stated.† *Sorex cinereus* of Bachman,‡ which Professor Baird cites as a synonyme of his *Blarina cinerea*, Dr. Bachman subsequently regarded as the young of his *S. carolinensis*,§ which is the same as *B. talpoides et brevicauda* of recent writers.

* North American Mammals, p. 50.

† *Bull. Mus. Com. Zoöl.*, Vol. I, No. 8, p. 212.

‡ *Journ. Phil. Acad. Nat. Sci.*, Vol. VII, 1837, p. 373, pl. xxiii, fig. 3.

§ Quadrupeds of North America, Vol. III, p. 344.

TALPIDÆ.

21. *Scalops aquaticus* Fischer. SHREW MOLE.

Several specimens of this species from Indian River and Jacksonville, Florida, are mentioned by Professor Baird in his list of the specimens of this species in the Museum of the Smithsonian Institution, in his Report on North American Mammals. Mr. Boardman has also informed me that it is not uncommon there.

SCIURIDÆ.

22. *Sciurus niger* Linné. SOUTHERN FOX SQUIRREL.

Sciurus niger LINNÉ, Syst. Nat., I, 64, 1758.

Sciurus vulpinus GMEL., Syst. Nat., I, 147, 1788.

Sciurus vulpinus et syn. BAIRD, Mam. N. Am., 246, 1857.

Common. Confined chiefly to the pine woods. Extremely variable in general color, the variations in this respect ranging from pale yellowish gray to black. The specific name *niger* of Linné is the one which has unquestionably the priority, as observed by Professor Baird, and its applying only to a single stage of coloration, inasmuch as it is a common one, does not seem to be sufficient reason for rejecting it, since it is as applicable as any name referring to its color can be, and is not likely to seriously mislead.*

23. *Sciurus carolinensis* Gmelin. GRAY SQUIRREL.

Sciurus carolinensis GMEL., Syst. Nat., I, 143, 1788. — BAIRD, Mam. N. Am., 256, 1857.

"*Sciurus cinereus* SCHREBER, Säugeth., IV, 766, pl. cccxiii, 1792."

Sciurus niger GODMAN, Am. Nat. Hist., II, 133, 1826.

Sciurus leucotis GAPPER, Zoöl. Journ., V, 206, pl. xxi, 1830.

Exceedingly abundant, and generally very tame. Two of my party shot a dozen one evening in less than half an hour at Hawkinsville. They are considerably smaller than at the North, and also differ somewhat in color from northern specimens, the gray being more suffused with brownish than in the gray northern type.

The fifty or sixty specimens carefully examined were quite uniform in color and generally so in size. The yellowish-brown patch on the back usually present in the gray type of this species was of greater extent and less distinctly defined than in northern examples. No

* See Baird, North American Mammals, p. 248.

dusky or black varieties were noticed, nor could I learn that they existed here. Their voice is not so heavy as that of the northern animal, but in no other respects than in those above mentioned do they differ from it. Professor Baird has quite fully described the gradual transition from the common gray to the glossy black type of coloration seen at the North, where the dark varieties are most common.*

Measurements of Florida Specimens.

M. C. Z No.	Original No.	Sex.	Locality.	Date.	Collected by	Total Length.	Nose to Occiput	Nose to Tail.	Tail to end of Vertebrae	Tail to end of Hairs.	Length of Fore Foot	Length of Hind Foot
2455	203	♂	Jacksonv'l	Jan. 12	C.J. Maynard	20.00	2.50	9.00	8.00	11.00	2.00	2.45
2454	206	♀	"	" 10	"	18.20	2.40	9.00	7.75	10.45	1.35	2.35
2453	207	♀	"	" 12	"	17.75	2.45	8.45	7.50	10.25	1.45	2.45
2040	351	♀	"	" 25	J. A. Allen	19.50	—	9.50	7.50	10.00	—	—
2041	352	♂	"	" 25	"	20.50	—	10.50	7.50	10.00	—	—
	221	♀	Dummitt's	Mar. 16	C.J. Maynard	20.00	2.45	9.50	8.15	10.50	1.60	2.56
	222	♂	"	" 18	"	18.56	2.55	10.20	8.00	10.56	1.35	2.40
2054	365	♂	Welaka	Feb. 6	J. A. Allen	21.00	—	10.00	8.25	11.00	—	—
2055	366	♀	"	"	"	21.50	—	10.00	8.75	11.50	—	—
2056	367	♂	"	"	"	19.00	—	10.00	8.40	9.00	—	—
2057	368	♂	"	"	"	19.00	—	9.00	7.50	10.00	—	—
2058	369	♀	"	"	"	19.00	—	9.15	7.60	9.85	—	—
2059	370	♂	"	"	"	21.50	—	10.15	9.60	11.35	—	—
2066	377	♂	Hawkinsv'l	Mar. 12	"	19.50	2.40	9.50	8.00	10.50	1.50	2.15
2067	378	♀	"	"	"	19.50	2.42	9.25	8.00	9.25	1.60	2.23
	379	♀	"	"	"	19.25	2.33	8.25	8.75	11.00	1.35	2.22
	380	♀	"	"	"	20.00	2.50	9.25	8.00	9.75	1.40	2.20
	381	♀	"	"	"	18.75	2.45	8.75	7.60	10.00	1.50	2.25
2068	382	♂	"	"	"	19.50	2.35	8.90	7.80	10.60	1.50	2.05
	384	♂	"	"	"	19.00	2.45	9.00	7.60	10.00	1.50	2.25
	385	♂	"	"	"	18.50	2.30	9.50	6.75	9.00	1.40	2.20
	386	♀	"	"	"	18.50	2.35	8.50	8.00	10.00	1.40	2.15
	387	♂	"	"	"	19.75	2.45	9.00	8.50	10.75	1.45	2.20
	388	♂	"	"	"	20.50	2.50	10.00	8.00	10.50	1.45	2.37
	389	♂	"	"	"	19.75	2.35	8.75	8.00	11.00	1.42	2.25
	390	♂	"	"	"	19.25	2.40	9.00	8.00	10.25	1.45	2.22
	391	♂	"	"	"	19.00	2.35	9.25	7.65	9.75	1.40	2.30
	392	♂	"	"	"	18.50	2.40	9.00	7.50	9.50	1.45	2.32
	393	♂	"	"	"	19.25	2.40	9.15	7.85	10.10	1.45	2.22
	394	♀	"	"	"	20.00	2.38	9.25	8.15	10.75	1.45	2.20
	395	♂	"	"	"	19.50	2.40	9.00	7.90	10.50	1.50	2.40
	396	♂	"	"	"	20.00	2.50	9.75	8.15	10.25	1.55	2.35
	397	♂	"	"	"	19.25	2.40	9.00	7.75	10.25	1.40	2.30
	398	♂	"	"	"	18.35	2.40	8.75	7.95	9.60	1.55	2.20
	399	♀	"	"	"	20.50	2.50	10.15	7.85	10.35	1.65	2.45
	400	♂	"	"	"	19.00	2.40	8.90	7.35	10.10	1.60	2.15

* N. Am. Mam., p. 259. See further on this point my remarks on this species in No. 8 of the first volume of this Bulletin, already cited.

24. *Geomys pineti* Rafinesque. "SALAMANDER."*Geomys pinetis* RAF., Amer. Month. Mag., II, 45, 1817.*Pseudostoma floridana* AUD. & BACH., Quad. N. Am., III, 242, pl. cl, fig. 1, 1853.

Common, but mainly confined to the drier portions of the pine woods. The five specimens collected by me differ very much in size, and considerably in color, some of them being plumbeous and others brownish-plumbeous; in other words, some are much darker than others. The difference in size appears to be mainly due to age. This species extends southwards at least as far as Lake Harney, and at some localities is particularly numerous, the little hillocks of earth it throws up sometimes nearly covering the ground.

MURIDÆ.

25. *Mus decumanus* Pallas. BROWN RAT.

Abundant at Jacksonville, but not observed by any of my party on the Upper St. John's, nor by Mr. Maynard on Indian River.

Although no other species of *Mus* was observed, it is not improbable that the common mouse (*M. musculus*) occurs in the vicinity of the towns. It was not found on the Upper St. John's (to which locality it probably has not yet extended), where the common house mice are a species of *Hesperomys*, as are also the house mice on Indian River, according to Mr. Maynard. Neither was any species of *Reithrodon* obtained. The *R. humilis*, which occurs in Georgia and South Carolina, is certainly to be expected in Northern Florida; but it has not yet to my knowledge been reported from there.

26. *Hesperomys leucopus* Wagner. WHITE-FOOTED MOUSE.*Mus sylvaticus*, var. ERXL., Syst. Reg. An., I, 390, 1775.*Mus leucopus* DESM., Mam., II, 307, 1822. — AUD. & BACH., Quad. N. Am., I, 300, pl. xlv, 1849.*Mus agrarius* GODMAN, Am. Nat. Hist., II, 1826.*Mus noveboracensis* SELYS-LONGCH., Étude Micromam., 67, 1839.*Mus Emmonsii* DEKAY, Emmon's Rep. Quad. Mass., 61, 1840.*Cricetus myoides* GAPPER, Zoöl. Journ., 1830, 204.*Hesperomys polionotus* WAGNER, Wieg. Arch., 1843, ii, 52.? *Hesperomys cognatus* LECONTE, Proc. Phil. Acad. Nat. Sci., VI, 442, 1852.*Hesperomys leucopus* LECONTE, Ibid., 413. — BAIRD, Mam. N. Am., 459, 1857.

— ALLEN, Bull. Mus. Comp. Zoöl., I, 227, October, 1869.

Hesperomys myoides BAIRD, Mam. N. Am., 472.*Hesperomys indianus* MAX. ZU WIED., Arch. für Naturg., XXVIII, i, 111, 1862.

A mouse provisionally referred to this species was abundant, especially at certain localities. At my first camp, about twenty-five miles above Jacksonville (near Hibernia), an *Hesperomys* and the wood rat (*Neotoma floridana*) were excessively numerous. At evening they began scampering over the leaves, their little footsteps being heard in every direction; at times they approached so near the camp-fire as to be distinctly seen. They ascended the bushes, and could be heard on the lower branches of the trees. Some of my party being unaccustomed to such manifestations of nocturnal life, were at first filled with apprehension as to the character of their visitors, and could scarcely be convinced that the place was not infested with poisonous snakes or other dangerous animals. Depending upon my traps for specimens, which unfortunately for me the mice avoided, I secured but two or three examples of the *Hesperomys* so abundant here. These, with several others obtained by me elsewhere, as also others obtained on Indian River by Mr. Maynard, including both young and adult, are undistinguishable from the common *H. leucopus* of the North, the young being deep plumbeous.

I observed at this place a fact in respect to the habits of the *Hesperomys* I had not previously noticed nor seen pointed out, though it was noticed in all the parts of Florida I visited. I refer to its habit of cutting off the branches and main stems of the young saplings. I at first supposed this work to be that of the wood-boring larvæ of some coleopterous insect, so nearly did the "pruning" resemble that of the so-called "oak-pruners" (*Cerambycidæ* sp.). A closer examination, however, showed that, instead of the twigs being smoothly cut, as by a boring insect working from within outwards and severing the bark last, the cutting was begun from without, and that a considerable portion of wood had been gnawed away, both the cut surfaces being highest at the middle. Marks of the teeth of these little gnawers were also generally clearly distinguishable. No traces of boring by insect larvæ could be detected near the severed point. The branches thus cut are generally of about the size of one's finger, and are usually the main stem of a young sapling. Various species of trees are thus mutilated; but as they are usually destitute of fruit, the purpose of these animals in this work is not apparent. It is a habit that may be common to the *Hesperomys* of the North, but I have never seen it referred to. These little animals being a hundred-fold more numerous

in East Florida than they generally are in the Northern States, their work would here be of course much more noticeable.

27. *Hesperomys aureolus* Wagner. GOLDEN MOUSE.

Arvicola Nuttalli HARLAN, Month. Amer. Journ. Geol. & Nat. Sci., I, 446, 1842. — IBID., Med. & Phys. Researches, 55, pl. —, 1835.

Mus (Calomys) aureolus AUD. & BACH., Journ. Phil. Acad. Nat. Sci., VIII, 302, 1842. — AUD. & BACH., Quad. N. Am., II, 305, pl. xcv, 1851.

Hesperomys aureolus WAGNER, Wieg. Archiv, 1843, ii, 51.

Hesperomys Nuttalli BAIRD, Mam. N. Am., 467, 1857.

A single specimen which I refer to this so-called species was obtained by Mr. Maynard at Dummitt's. While this example is of the size and general proportions of *H. leucopus*, it is markedly different in color, being of a bright golden yellow above, which color reaches on the outside of the legs to the feet; the under surface has also a yellowish wash. It also differs in the texture of its fur, which is remarkably soft and fine. It is a little lighter colored than Audubon and Bachman's description and figure of *H. aureolus* represent that animal to be, but the distribution of the colors is the same, the specimen in question being not orange, but bright yellowish-cinnamon. It is, however, much nearer this than to Dr. Harlan's *Arvicola Nuttalli*. The latter does not differ very appreciably, judging from Dr. Harlan's very unsatisfactory description and his wretched figure of it, which was evidently made from a badly stuffed skin. Mr. Maynard believes the specimen referred to above to be a young animal, and states that it was so regarded by the people in whose house it was caught. He further informs me that he captured another of the same color, but very much larger, which was lost. This he regards as merely the adult of the same species. His measurements show the latter to have agreed in size and proportions with the so-called *H. gossypinus*. The texture of the fur of the small specimen above referred to agrees with that of the plumbeous, immature stage of *H. leucopus*. This form, whether a valid species or not,* is now known to occur in Southeastern Pennsylvania, Southern Illinois, Georgia, Florida, Mississippi, Missouri, and at several intermediate points.

28. *Hesperomys gossypinus* LeConte. COTTON MOUSE.

Hesperomys gossypinus LECONTE, Proc. Phil. Acad. Nat. Sci., VI, 411, 1853.

— BAIRD, Mam. N. Am., 469, 1857.

* This and the following species are only provisionally adopted. See a previous number of this Bulletin (Vol. I., No. 8, p. 227) for a fuller expression of my views as to the number of North American species of this group, and their mutual affinities.

Several specimens were obtained, corresponding in size and color with what LeConte and Baird have described under this name. It is apparently common. As I have previously stated elsewhere,* these Florida specimens have well-developed cheek-pouches.

The specimens in question are rather larger than any examples of *H. leucopus* I have seen from the Northern States, they agreeing very well in measurements with the two specimens cited by Professor Baird.† The large size of these specimens, conjoined with their southern habitat, would seem at first to clearly indicate their being distinct from *H. leucopus*, as they are at least one third larger than the average size of the latter at the North. Professor Baird in speaking of this species observes: "There is every reason to consider this mouse as specifically distinct from *H. leucopus* of the North; although skins, when much stretched (as Nos. 1105, 1112, from Middleboro', Massachusetts), of the latter, may measure as much as those recorded here, yet they are certainly actually smaller, as shown by the feet, which never attain anything of the length of .45 for the anterior and .90 for the posterior." But he is "hardly satisfied," he adds, "that this animal is different from the smaller *H. leucopus*, as the difference in size is no greater than is to be seen in a series of *Hesperomys* from more northern localities. The tail is duskier beneath than in *H. cognatus*, and the sides more rusty; otherwise I can realize only the larger size. Should both [*H. cognatus* and *H. gossypinus*] prove to be the same, the name *H. gossypinus* must of course take precedence."

As already observed, the prevailing form of the *Hesperomys* of East Florida is not essentially different from a large proportion of the *H. leucopus* of the North, either in measurements, proportions, or color, although it is unmistakably referable to the so-called *H. cognatus*, which has been supposed to replace in the Southern Atlantic and Gulf States the *H. leucopus* of the more northern ones. If, as I have elsewhere suggested (*loc. cit.*), as Professor Baird admits may be, and as the facts seem to indicate, *H. gossypinus* is inseparable from *H. cognatus*, and the latter being most unquestionably referable to *H. leucopus*, it would seem that *H. gossypinus* must also be referred to the *H. leucopus*.

Respecting the variations in this species and the affinities of the *H. gossypinus*, Audubon and Bachman observe as follows: "That a species so widely distributed and subject to so many variations in size, length of tail, and color, should have been often described under different names is not surprising. We have ourselves often been in a state of doubt on obtaining some striking variety. The name *Hypudæus gossypinus* of our

* Bulletin Mus. Comp. Zoöl., Vol. I, No. 8, p. 229, 1869.

† Mam. N. Am., p. 469.

friend Major LeConte (see Appendix to McMurtrie's translation of Cuv. An. Kingd., Vol. I, p. 434) was intended for this species, as it is found in the Southern States. We were for several years disposed to regard it as distinct, and have, not without much hesitation, and after an examination of many hundred specimens, been induced to set it down as a variety only." These authors also remark that they are considerably larger in the Carolinas than in the Eastern States.*

29. *Hesperomys palustris* Wagner. RICE-FIELD MOUSE.

Mus palustris HARLAN, Am. Journ. Sci., XXXI, 386, 1837.

"*Hesperomys palustris* WAGNER, Supplem. Schreb. Säugeth., III, 543, 1843."

Hesperomys (Oryzomys) palustris BAIRD, Mam. N. Am., 482, 1857.

Arvicola oryzivora AUD. & BACH., Quad. N. Am., III, 214, pl. cxliv, fig. 3, 1857.

No specimens of this species were obtained by either Mr. Maynard or myself. Its habitat is usually given as South Carolina and Georgia, but Audubon and Bachman state: "The late Dr. Leitner brought us a specimen obtained in the Everglades of Florida."† It in all probability occurs also in East Florida. The above-mentioned authors give it as somewhat common in the salt-marshes near Savannah and Charleston. Professor Baird has received it from Columbus and St. Simon's Island, Georgia, and Society Hill, South Carolina.

30. *Neotoma floridana* Say & Ord. WOOD RAT.

Mus floridanus ORD, Bull. Soc. Philom., 1818, 181. — SAY, Long's Exped., I, 54, 1823.

Arvicola floridana HARLAN, Faun. Amer., 141, 1825.

Neotoma floridana SAY & ORD, Journ. Phil. Acad. Nat. Sci., IV, ii, 352, 1825. — BAIRD, Mam. N. Am., 487, 1857.

I found this species very abundant on the Lower St. John's, especially around Jacksonville and Hibernia, but I did not meet with it above Lake George. The old residents about Hawkinsville seemed wholly unacquainted with it. Mr. Maynard also failed to meet with it on Indian River. It hence appears probable that it may not occur very frequently in the southern part of the peninsula. Professor Baird, however, has recorded a specimen from "Indian River, Fla.," collected by Dr. Wurdemann.

The present usual northward range of this species does not appear to extend beyond North Carolina; but Professor Baird, writing in 1857,‡

* Quad. N. Amer., Vol. I, pp. 301, 305.

† Mam. N. Am., p. 489.

‡ Ibid., Vol. III, p. 216.

remarks: "A few specimens of unusually large size were captured some years ago by J. G. Bell, near Piermont, on the Hudson River, but I have not heard of any in intermediate localities [New York and Society Hill, South Carolina]." Mr. George Gibbes states that he "caught a specimen, many years ago, in Massachusetts."* Audubon and Bachman remark that specimens of it have been obtained in North Carolina, and that they had "observed a few nests in the valleys of the Virginia mountains," and that they had "somewhere heard it stated that one or two had been captured as far to the north as Maryland."†

31. *Sigmodon hispidus* Say & Ord. COTTON RAT.

Arvicola hispidus GODMAN, Am. Nat. Hist., II, 68, 1826.

Arvicola hortensis HARLAN, Faun. Am., 138, 1825.

Arvicola ferrugineus HARLAN, Am. Journ. Sci., X, 285, 1826.

Sigmodon hispidum SAY & ORD, Journ. Phil. Acad. Nat. Sci., IV, ii, 354, pl. x, figs. 5-8, 1825. — BAIRD, N. Am. Mam., 503, 1857.

Sigmodon Berlandieri BAIRD, Proc. Phil. Acad. Nat. Sci., VII, 333, 1855.

IBID., N. Am. Mam., 504.

Abundant throughout the country along the St. John's River, and also on Indian River, whence Mr. Maynard brought fifteen specimens. They are quite a pest to the farmers, who often successfully resort to poison to reduce their numbers. By scattering grain poisoned with strychnine about their fields they are able to destroy hundreds with slight trouble. Different specimens vary considerably in color, from gray through yellowish-brown to rufous. The so-called *Sigmodon Berlandieri*, from Texas and New Mexico, seems undistinguishable from *S. hispidus*.

In its general economy, the cotton rat represents the *Arvicolæ* of the North, especially *A. riparius*.

Concerning *S. Berlandieri*, Professor Baird remarks: "This species is readily distinguishable from *S. hispidus* by the much lighter color above, where it is grayish-yellow brown instead of distinct reddish-brown; the tail is considerably longer and covered by finer annuli. The toes are shorter, and the metatarsus shorter, while the feet are nearly the same length. The claws, however, are much weaker." The tail in this species is said to be "equal to or longer than the trunk"; the "color above grayish-yellow brown, lined with black"; while *S. hispidus* is said to have the

* Nat. Hist. Wash. Terr., Zoöl., p. 128, 1860.

† Quad. N. Am., Vol. I, p. 36.

tail "less than the trunk," and "the color above reddish brown, lined with very dark brown." The specimens from Florida examined by me are mainly of the gray type, and hence like *S. Berlandieri*, but some were decidedly rufous, or like *S. hispidus*. In "Mammals of North America" measurements of specimens of the so-called *S. Berlandieri* are given, and of twelve of *S. hispidus*. In the latter the length of the tail to the length of the trunk is as 69 to 100; in the former (*S. Berlandieri*) as 63 to 100! It hence appears from Professor Baird's own measurements that the *S. Berlandieri* is far from having the tail relatively the longer. The other distinctions are based on too few specimens to have much value, since individual variations of the same character are common.

32. *Arvicola pinetorum* LeConte. PINE MOUSE.

Psammomys pinetorum LeConte, Ann. N. Y. Lyceum Nat. Hist., III, 132, pl. ii, 1829.

Arvicola scalopsoides Aud. & Bach., Journ. Phil. Acad. Nat. Sci., VIII, 299, 1842.

Arvicola pinetorum Aud. & Bach., Quad. N. Am., II, 216, pl. lxxx, 1851.

Arvicola (Pitymys) pinetorum Baird, N. Am. Mam., 544, 1857.

Included on the authority of Audubon and Bachman, who state that they had received it from Florida, Alabama, Mississippi, and Georgia. Professor Baird also cites specimens from Georgia and Louisiana. This is the most southern of the *Arvicolæ*, and the only one, except *A. austerus*, whose habitat includes the Gulf States.

LEPORIDÆ.

33. *Lepus sylvaticus* Bach. GRAY RABBIT.

Lepus americanus Desm., Mam., II, 351, 1822. — Harlan, Faun. Amer., 193, 1825.

Lepus sylvaticus Bach., Journ. Phil. Acad. Nat. Sci., VII, 1837. — Waterh., Nat. Hist. Mam., II, 116, 1848. — Aud. & Bach., Quad. N. Am., I, 173, pl. xxii, 1849. — Baird, Mam. N. Am., 597, 1857.

Abundant. Mr. Maynard obtained a specimen but a few weeks old, at Dummitt's, as early as the 16th of February

34. *Lepus palustris* Bachman. MARSH RABBIT.

Lepus palustris Bach., Journ. Phil. Acad. Nat. Sci., VII, 194, 336, pl. xv, xvi, 1837; Ibid., VIII, 79, 1839. — Aud. & Bach., Quad. N. Am., I, 151, pl. xviii, 1849. — Baird, Mam. N. Am., 615, 1857. — Coues, Proc. Bost. Soc. Nat. Hist., XIII, 86, 1869.

Common, especially on the Lower St. John's.*

* See on the habits and anatomy of this species a paper by Dr. Elliott Coues, Proceed. Bost. Soc. Nat. Hist., Vol. XIII, pp. 86 - 101, June, 1869.

DIDELPHIDÆ.

35. *Didelphys virginiana* Shaw. OPOSSUM.

Didelphys virginiana SHAW, Gen. Zoöl., I, 473, pl. cvii, 1800. — DESMAREST, HARLAN, TEMMINCK, WATERHOUSE, BAIRD, and most other authors.

? "*Didelphys marsupialis* SCHREB., Säugeth., III, pl. cxlv, 1778."

Didelphys californica BENNETT, Proc. Lond. Zoöl. Soc., I, 40, 1833. — Also WAGNER, WATERHOUSE, AUD. & BACH. (from Bennett). — BAIRD, Mam. N. Am., 233, 1857. — BAIRD, U. S. & Mex. Bound. Surv. Rep., II, Zoöl., 32, 1859.

Didelphys breviceps BENNETT, Proc. Lond. Zoöl. Soc., I, 40, 1833. — WATERHOUSE, Nat. Hist. Mam., I, 477, 1846 (from Bennett?). — AUD. & BACH., Quad. N. Am., III, 330, 1851 (from Bennett).

Didelphys pruinosus WAGNER, Wiegmann's Archiv, 1842, 358. — WATERHOUSE, Nat. Hist. Mam., I, 477, 1846, (from Wagner).

Abundant.

This species is quite variable in its color-markings, and remarkably so in many other features, especially in the length and size of the nose, and in the size and proportions of the skull, even in specimens from the same locality.* Slight and quite inconstant differences also occur between examples from the Southern States, Texas, Mexico, and California. It would, in fact, be quite unusual if specimens of any species ranging so widely should not be found to differ somewhat at localities so widely separated. Two supposed species of North American *Didelphys* described by Mr. Bennett, as cited above, have been quoted by numerous other authors, and by them currently adopted, without apparently an examination of their merits. Professor Baird, rejecting one of them, has endeavored to separate the opossums occurring west of the Mississippi valley from those living farther eastward, designating the western one as *D. californica*. The distinctions claimed are somewhat similar to those urged as distinguishing the so-called *Procyon Hernandezii* of the western half of the continent from the *P. lotor* of the Atlantic States. They are equally slight and unsatisfactory, and at most mark but a geographical race, so intimately allied to and intergrading with the better-known eastern form that the point at which the one supplants the other is thus far undetermined. The *Didelphys breviceps* of Bennett was founded on a single specimen from California, which differed from the so-called *D. californica* only in having a relatively shorter head.

* Since writing the above I have been incidentally informed by Dr. Coues that, in preparing his memoir on the anatomy of *Didelphys virginiana* (now publishing in the Mem. of the Bost. Soc. Nat. Hist., Vol. II, Pt. I), he had occasion to examine a large number of specimens, and that he found the variation in size and proportions to amount to nearly twenty per cent.

PART III.

On Individual and Geographical Variation among Birds, considered in respect to its bearing upon the Value of certain assumed Specific Characters.

A systematic investigation of the extent and character of individual variation in birds seems not to have hitherto been attempted; in fact, few collections exist that furnish the material necessary to such a work. In occasional instances considerable differences between individuals of the same species, other than those that result from age and sex, have, however, already been pointed out, but these instances seem to have been generally, but improperly, regarded as exceptional cases.

The collection of birds in the Museum of Comparative Zoölogy now offers unusual facilities for a general investigation of this subject, most of the common species of Eastern North America being each represented by fifty to one hundred and fifty or more specimens. The greater part of them having been collected in Southern New England, and a large proportion in Eastern Massachusetts, they are the more valuable for this purpose, from their having been collected essentially from the same locality. The examination of this material has disclosed a hitherto unsuspected range of purely individual differentiation in every species thus far studied. At the same time regard has been had to the more obscure seasonal variations in color, and to the general differences that depend upon age, including such as result from senility as well as from immaturity. Local or geographical variations have likewise been carefully considered, with results that a short time since were unsuspected. These several lines of investigation have shown that in many instances what have been regarded as reliable characteristics of species have in not a few cases really little or no value; that the importance of many diagnostic features has been too highly estimated, and that consequently a careful revision of our published faunæ will be necessary for the elimination of the merely nominal species. In the following pages many of the data which have led to these conclusions will be presented.

Individual variation not only affects color and size, but the proportions of different parts, as the relative size and form of the wings, tail, bill, toes, and tarsi, including the skeleton as well as the external organs;

of the soft parts no account can as yet be given. Geographical variation has an equally universal range, but is most strikingly exhibited in the color, in size, and in the form of the bill. Individual variation will be first considered, and subsequently geographical variation. In each case each prominent phase of variation will be more or less fully described.

1. INDIVIDUAL VARIATION.

Individual Variation in Color. — In birds of whatever age, two lines of variation from the average or medium type of coloration are readily distinguishable, the variation depending essentially on differences in the depth or intensity of the general tint. On the one hand, individual variation in color results from a greater than the average amount of coloring matter in the integuments; on the other hand, from an amount less than the average amount. The difference in this respect between the extremes of a series of fifty or one hundred specimens of any species, collected in course at a single locality, and nearly at the same season of the year, is often as great as occurs between truly distinct species. But the difference is here solely one of *intensity of color*, while in allied species there is almost always an appreciable variation in the *style of coloration*. In individual variation the differences usually extend alike to all parts of the integuments; that is, if the plumage of the upper surface of the body is brighter or paler than usual, the same difference extends to the plumage of the lower surface of the body, and also to the bill and the feet. This is noticeable not only in species that have the color in uniform masses, differing in tint on different regions of the body, as in the robin (*Turdus migratorius*), the blue-bird (*Sialia sialis*), the Maryland yellow-throat (*Geothlypis trichas*), the mocking-bird (*Mimus polyglottus*), and species generally of that type of coloration, but also in spotted birds, as in the various spotted species of *Fringillidæ*, *Turdus*, *Dendræca*, etc., where the plumage on certain regions of the body is marked with numerous streaks and spots differing from the ground color, in which case the intensity of the color of the markings correlates in its variations with that of the ground color. Closely allied species, on the contrary, usually vary more or less, not only in respect to the ground color, but also to a greater or less degree in the style of the markings. In illustration of this point the familiar group of the small, spotted-breasted wood-thrushes of Eastern

North America — the group *Hylocichla* of Professor Baird — may be taken. Three of these species (*Turdus fuscescens*, *T. Swainsoni*, and *T. Pallasi*) are so closely related that for many years they were variously confounded with each other by almost all who wrote of them, one of them not being clearly recognized as distinct from the others till thus established by Dr. T. M. Brewer,* in 1844, and also at about the same time by Mr. J. P. Giraud,† each apparently independently of the other. Yet they are so distinct that there seems to be not the slightest excuse for again confounding them. While they all agree so closely in general size, in form, and in proportions, that a series of detailed measurements of many specimens of each species gives in the average no constant differences in any of these particulars, each differs from the other radically and constantly in style of coloration, and somewhat in general tints, in habitat, nidification, habits and song. Two of these species (*T. fuscescens* and *T. Swainsoni*) agree in the *style* of the coloration of the dorsal surface, but differ so much in the *color* of this part, that this character alone is always sufficient to separate them, while a still wider difference is seen in the color and markings of the ventral surface, a glance at this part of *T. fuscescens* being sufficient to invariably distinguish it from either of its above-named allies. The third species (*T. Pallasi*) differs markedly from both the other two in the style of coloration of the *dorsal surface*, the rump and tail being conspicuously different in color from the anterior part of the body, whilst the others exhibit no contrast of color between these regions. But the under surface of *T. Swainsoni* is so like that of *T. Pallasi* that frequently specimens cannot readily be referred to the one species rather than to the other from a view of this surface alone. This group serves as a fair general illustration of the kind of variation in color usually seen in closely allied species, but there occur occasional exceptions, where a difference in the relative proportions of different parts, or a wide difference in size, is the prominent specific distinction, the smaller species, so far as color is concerned, being a diminutive representative of the larger.

Taking the present group of *Hylocichla* (for reasons that will appear hereafter ‡) as a group illustrative also of individual variation, it is found that the differences in color in different individuals of either species

* Proc. Boston Soc. Nat. Hist., Vol. I, p. 191, July, 1844.

† Birds of Long Island, p. 91, 1843-44.

‡ See the remarks on these species in Part IV.

results from the amount of rufous pervading the plumage. Individuals of *Turdus Swainsoni* of the rufous or bright-colored type have the dorsal surface of a uniform brownish-olivaceous tint, and the sides of the head and breast strongly suffused with yellowish-brown, which tint is also traceable throughout the lower plumage, in the brighter color of the basal brownish band on the inside of the wings, and in the color of the mouth and base of the bill. In other individuals the upper plumage is of a dark olivaceous tint, without any trace of brownish, the sides of the head, neck, and breast being ashen, with often no appreciable tinge of ferruginous; specimens of this type thus differing widely in general aspect from those of the other. Between these extremes, of which examples are not unfrequent, nor confined to any particular locality or season of the year, there is every degree of intergradation, specimens intermediate between the two being by far the most frequent, and constituting the average or common form.

Turdus Pallasi and *Turdus fuscescens* present precisely similar variations. They are also seen in *Turdus mustelinus*, in *Turdus migratorius*, in *Sialia sialis*, in *Seiurus noveboracensis*, in many species of *Dendroica*, sparrows, and other species which I have especially investigated in reference to this point, embracing examples of all the leading families of birds. The ruffed grouse (*Bonasa umbellus*), as is well known, varies in the color of the upper parts from reddish-brown to gray; the great horned owl (*Bubo virginianus*) from dusky through numerous shades of rufous and fulvous to nearly white, the fulvous suffusion so commonly present in this species varying from ferruginous on the one extreme to its complete obsolescence on the other. In such common and thoroughly known species as the robin, blue-bird, etc., the true character of these variations is recognized, but in groups where the species are not well known, and especially in specimens from partially explored regions, they are frequently regarded as of specific value, and the addition of numerous nominal species is the result.

Besides the variation in the *depth* of color already noticed, birds having the plumage varied with streaks and spots differ exceedingly in different individuals of the same species in respect to the size, shape, and number of these marks, and in the general aspect of the plumage resulting from such variations. Generally, as already stated, such differences correlative with the variations in the intensity of the ground color, the darker or more deeply colored birds being usually those with the mark-

ings largest and brightest. A wide range of variation in this respect is seen in all birds which have the breast and lower plumage marked with dark streaks and spots on a lighter ground, or that have the whole plumage streaked. In the common song sparrow (*Melospiza melodia*), the fox-colored sparrow (*Passerella iliaca*), the swamp sparrow (*Melospiza palustris*), the black and white creeper (*Mniotilta varia*), the water wag-tail (*Seiurus noveboracensis*), in *Turdus fuscescens* and its allies, etc., the difference in the size of the streaks is often very considerable. In the song sparrow they vary to such an extent that in some cases* they are reduced to narrow lines; in others so enlarged as to cover the greater part of the breast and sides of the body, sometimes uniting on the middle of the breast into a nearly continuous patch. Variation in this respect is equally great in the fox-colored sparrow and in the grass finch (*Poocætes gramineus*). Massachusetts specimens of the savanna sparrow (*Passerculus savanna* auct.) also present variations exactly parallel with those of the song sparrow. Yet these differences, with other variations to be hereafter mentioned,† have been regarded, as in the case of *Passerculus savanna*, as of specific value. Similar variations in the *Hylocichla* group are very marked, as in *Turdus (Hylocichla) fuscescens* especially. In some specimens of this species the colors are on all parts not only very pale, but the markings on the breast are reduced to indistinct narrow lines; in others, in which the general color of the plumage is darker, the markings on the breast are dark, broad, and triangular. Two specimens taken in Cambridge the same day (early in May), both of which are males, exhibit these extremes. Average male specimens of the black and white creeper (*Mniotilta varia*), in which the plumage is varied with longitudinal black and white streaks, have the black streaks about a third broader than the white ones; but other specimens occur in which the white ones are equal to and even broader than the black ones; others have the black streaks so much broader than they usually are, — the white ones of course being proportionally reduced, — that the general aspect of the plumage at a short distance is nearly black. The difference between these two extremes is strikingly great. Yet similar variations, scarcely less in degree, occur in nearly all of the striped-breasted warblers.

In birds which have the ground color of certain areas of the body

* Perfectly mature specimens only are here referred to.

† See the remarks on the genus *Passerculus* in Part IV.

black spotted with white, as in some of the woodpeckers (*Picus villosus* and *P. pubescens*, for example), the white markings vary in size most notably, and sometimes in number. The white markings so common on the wings and tails of birds, as the bars formed by the white tips of the greater wing coverts, the white patch occasionally present at the base of the primary quills, or the white band crossing them, and the white patch near the end of the outer tail feathers, are also extremely liable to variation in respect to their extent and the number of feathers to which, in the same species, these markings extend. Variation in the tail markings is particularly common, as may be seen by comparing numerous specimens of almost any species of *Dendræca*, *Junco*, *Pipilo*, of *Mimus polyglottus*, *Chordeiles popetue*, etc. In the latter species the white patch on the wing does not ordinarily encroach upon the outer vane of the first primary, and rarely upon its shaft, but in several specimens before me it covers not only the shaft of the first primary, but extends completely across its outer vane! The black subterminal bar on the upper surface of the tail of the ruffed grouse (*Bonasa umbellus*) ordinarily crosses all but the middle pair of feathers, on which there is usually no trace of this bar; in many specimens, however, it is barely traceable on them, and in others it is as distinct and perfect on the middle pair as on the others.*

The *Parula americana* presents also remarkable examples of individual color variation. The colors of the males are usually much brighter than those of the females, but cases are frequent where the sex cannot be determined by the color of the plumage. Adult males also vary greatly in the *style* of coloration. They are generally bright yellow anteriorly below, with a broad band of dusky reddish-brown across the breast, varying in tint from nearly pure chestnut to dusky reddish-brown, and even black, and also greatly in extent. In some, however, this band is partially obsolete, in which case the whole plumage is generally paler than in average specimens. More rarely large, brightly colored males are taken, even in New England, with the whole breast bright yellow, the brownish pectoral band being entirely absent. This condition, however, seems to be more frequent in specimens of *Parula* collected in Mexico, and Central and Northern South America, which on this account have been regarded as distinct from the *Parula* of the North; yet all the conditions of color seen in specimens from

* See remarks on color variations in other species in Part IV.

the North are also common to those from the South, and *vice versa*.

In species in which the female usually differs from the male in being paler colored, the pattern of coloration being the same in both sexes, females occur more or less frequently which are as brightly colored as the brightest males, and males that are paler than the generality of the females.

Variation in Color depending on Season. — A word in this connection seems necessary concerning some of the more obscure variations depending upon season and age, since it is sometimes difficult to avoid confounding these differences with those resulting from individual variation. In many species there is a marked change in the color of the plumage without a change of the plumage itself. No experienced collector can have failed to notice the much brighter and livelier tints the plumage of most song birds presents immediately after the autumnal moult, in species in which there is no marked seasonal change of color, in comparison with the faded appearance they exhibit towards the close of the breeding season. This brighter autumnal tint is particularly marked in the *Vireos*, the different species of *Empidonax*, *Sayornis*, *Contopus*, and in some of the *Sylvicolidæ*, and is clearly traceable in hundreds of other species. But almost as great a difference is seen when specimens of any species taken in spring, on its first arrival at its breeding station, are compared with those collected several weeks later, or just before the autumnal renewal of the plumage. In this case the variation results in part from an actual fading of the color, and in part from the wearing of the edges of the feathers. Seasonal differences of this character are often only readily appreciable to the experienced eye, and the failure to recognize the cause of these differences has led in many instances to their being regarded as of specific value. Especially noteworthy instances of such mistakes will be noticed later. Collectors, and even naturalists, generally place little value on faded or dull-colored specimens, so that ordinarily in collections of our native birds only fine-looking specimens are preserved. But travellers and explorers of new localities are often compelled to content themselves with any representative they may be able to get, so that the "closet" or exclusively "museum naturalist" has not usually the material necessary to furnish him with a clue to the cause of these variations.

Generally, aside from the paler tints of late-collected birds, as compared with those taken early in the season, there is a total absence of the grayish, yellowish, brownish, or rufous suffusions (the particular tint varying of course in different species) that tinges the feathers early in the season. The general aspect of the plumage at the two periods in question is thus essentially different. The common chickadee (*Parus atricapillus*) will illustrate this point, in which the brownish tint so conspicuous on the lateral portions of the ventral plumage in autumn and winter is gradually lost as spring approaches, and in summer is almost entirely wanting, especially in nesting females, which at this season have the plumage generally much more worn than the males. The savanna sparrow will also illustrate the differences resulting simply from the fading of the color during the breeding season. In spring both sexes have a greenish-yellow, superciliary stripe, varying more or less in intensity in different specimens, but rarely or never of the pale soiled-whitish so frequently met with late in the breeding season. In the large series of specimens before me collected at that season in Massachusetts, few if any have this stripe so bright as average spring specimens have it, in many it having faded to soiled white. Scores of similar cases might be cited, but the above are sufficient for illustration.

Variations in Color depending upon Age. — So well known are many of the variations depending upon age, that it seems necessary to advert to only a few of the lesser known phases. In many species there is no marked difference between old and young birds, after the moulting of the first or nestling plumage, which usually occurs in the oscine groups in a few weeks after they leave the nest. But even in these, in many cases, sufficient marks of immaturity remain for a time to enable any one acquainted with such features to recognize birds of one or two years of age from those that are older. Yearling birds of this group are often recognizable by their having more or less well-defined bars across the wings, formed of light-colored, hastate, or drop-shaped spots on the ends of the greater wing-coverts and inner secondaries, which in many genera are peculiar to yearling birds, though in other respects, so far as the plumage is concerned, they are not distinguishable from adults, — a difference which in some instances has been considered specific. Similar marks are also seen in older birds, in species that do not obtain their adult colors till later in life.

Yearling and two-year-old birds are also often distinguishable from older ones by the presence, after the spring moult, of a greater than the ordinary amount of ferruginous, ashy, or yellowish edging to the feathers, such as is often seen in the winter plumage of adult birds. In some cases such a bordering to the clothing feathers, especially those of the back, is often strictly distinctive of young birds, and is, moreover, a feature of common occurrence.

Generally speaking, several years elapse before the purity of the colors and the definiteness of outline of the markings characteristic of maturity is fully obtained, especially in highly colored species. In birds of variegated colors the contrasts of color become for a time more and more decided with each moult, and the markings better and better defined, especially in respect to the white bars of the wings and the spots on the tail common to a large number of species. The latter markings usually gradually increase in extent for a considerable period. A good illustration of this is seen in many of the gulls, particularly in the genus *Larus*. In *L. argentatus* the following gradual change with age occurs in the white markings on the tips of the primaries. At first, as ornithologists are aware, the plumage of this species is uniformly dusky, the adult colors not being acquired before the second year, and apparently frequently not before the third, there being in the breeding season usually a large proportion of individuals in the brown plumage.* But there are wide differences in the intensity of the color in different individuals in this stage of plumage, some being but slightly dusky and others extremely dark, — differences that probably result mainly from differences in age, the darker birds being probably yearling birds and the lighter ones two years old, though part of the difference is doubtless due to individual differentiation. In this stage the wings and tail are of nearly the same uniform dusky tint as the general plumage. In what may be considered as the second stage, the general color is somewhat lighter, the tail much lighter, and the primaries much darker, with a distinct paler apical margin. At a third stage the tail becomes white, the dorsal plumage begins to assume the blue tint characteristic of maturity, the primaries change from dull blackish brown to black, and a small white spot appears near the end of the inner vane of the first

* Generally the large parties that spend the summer on the coast of Massachusetts, where none of these birds now breed, consist almost wholly of birds in the brown stage of plumage. See American Naturalist, Vol. III, p. 640, 1870.

primary, separated from the white at the extreme tip by a broad space of black. A subsequent gradual increase occurs in the purity of the colors and in the extent and form of the wing markings. The complete series of the changes in the latter is as follows: At first, as previously remarked, the primaries are dull brownish black, a little darker than the general plumage, with their extreme apical margins lighter. At the next stage the three inner primaries have become much lighter, and the light border to all broader and whiter. Later the three inner primaries and the distal portions of most of the others become wholly ashy white, and the outer portion of the other primaries much blacker. The subapical dark portion of the wing now embraces only the seven outer primaries, and is of a triangular form, the first primary forming the base of the triangle. The black on the outer vane of the first primary reaches nearly to the base of the outer vane of the second, and is more and more restricted on the others, till on the sixth (or, more rarely, on the seventh) it forms only a narrow bar near the tip. In other words, the black, if present on the seventh primary, exists as a narrow transverse subapical bar, which bar increases in distal extension on the sixth, fifth, fourth, third, and second, to the first, and embraces the whole outer vane of the first primary. The basal outline of the black area being an oblique one, a much larger portion of the outer than of the inner vane of each feather is embraced in the black space. All the primaries are now terminated with a narrow white border, the first primary having also an oval white spot on the inner vane, near the end of the feather. Subsequently this spot enlarges so as to embrace a part of both vanes, the white at the tip of the feather also meantime increasing somewhat in extent, and the two being separated by a broad bar of black. Coincident with this increase in the amount of white on the first primary, a small white spot appears on the inner vane of the second primary. Subsequent increase in the extent of these white markings goes on until the white area on the second primary extends to both vanes, and the two white spots on the first primary are separated by only a narrow bar of black. Later still this bar becomes broken, through the partial union of the two white spots, and finally becomes entirely obsolete, leaving the first primary with a single continuous white apical area, an inch and a half to two inches in length. It is probable that not all individuals reach this final stage, though most doubtless do in old age. A large series of specimens of mature birds usually exhibit the gradual

change above described, and indicate the inconstancy of these markings and their unreliability as specific characters. Often, as is well known, these markings in the gulls differ considerably in the two wings of the same bird.

Although the *L. argentatus* has been taken as a general illustration, the same variations with age, or in different individuals, are exhibited by most species of the genus *Larus*. Generally they are admitted to have no value as specific characters, even by those who in the case of *L. argentatus* have accorded to them this importance.

In some of the species of *Junco* and *Pipilo*, in *Mimus*, in numerous species of *Dendræca*, in *Parula*, *Mniotilta*, etc., there is a similar increase with age in the extent of the white markings on the tail, sometimes three and sometimes four pairs of feathers being spotted or terminated with white in different specimens of the same species. In short, these variations occur in so many species that they may be looked upon as indicating a general law of variation in color depending upon age, namely, *an increase in the purity or intensity of the general color, and an increase in the size of the wing and tail markings, for a time, with age.*

After complete maturity is attained there is, however, unquestionable evidence of a decline in color, which in many cases, and especially in bright-colored species, is quite marked. So general is such a decline in other groups of the animal kingdom that a citation of evidence on this point seems wholly needless. Yet in birds, in numerous instances, it is scarcely appreciable, and doubtless is in most species too slight to be readily traced. This obscurity may result, however, more from an absence of favorable conditions for such a decline to be recognized than from its real absence. It can hardly be doubted, in fact, that a share of the color variation seen in mature birds is attributable to this cause. It is well known that young mammals in their first pelage are, as a general rule, much darker colored than the adults of the same species. At a later period the color fades more slowly, but in old age the hair often becomes more or less gray, the blanching being in some cases very marked. Nearly all birds are also darker in their nestling and immature stages of plumage than after they arrive at maturity, especially if in the adult stage the plumage is light colored; and it is more than probable, and in some cases certain, that the decline in color

continues in a slight degree through life. The change of *Falco candicans* from dusky when young to nearly white when fully mature may be hardly referable wholly to the blanching of age; but the gradual obsolescence of the dusky mottlings of the snowy owl (*Nyctea nivea*), as it advances in age, seems strictly parallel to the blanching of the gray colt to a white horse. Hence a second law of variation in color in old age, namely, that of *senile decline*.

INDIVIDUAL VARIATION IN GENERAL SIZE AND IN THE RELATIVE SIZE OF DIFFERENT PARTS.

Individual Variation in General Size and Form. — Measurements of scores of specimens of birds of the same species and sex, collected at the same locality and season, show the existence of a large range of individual variation, both in size and in general proportions; the variation extending to every external part of the body, and implying a corresponding variation in the internal anatomy. In birds size has usually been regarded, from its comparative constancy in the same species, as an important specific character. But from the fact that specimens of closely allied species often differ but little from each other in this respect, it has been justly looked upon as being in some cases more or less unreliable; but from the great importance commonly attached to it, it is evident that such instances are usually regarded as exceptional. Individual variation in this respect having been formerly regarded as too slight to have any significance, the size of a single specimen has usually been given as that of the species to which it belonged; hence subsequent variations from it discovered in other specimens of the same species has sometimes led to the recognition of the latter as specifically distinct. Especially has this been the case when a difference in size has been associated with a wide difference of locality. The facts in the case, however, show that a variation of fifteen to twenty per cent in general size, and an equal degree of variation in the relative size of different parts, may be ordinarily expected among specimens of the same species and sex, taken at the same locality, while in some cases the variation is even greater than this. Table A (p. 198) shows to some extent the general variation in size, but it does not always give, nor even generally, the extreme differences in the size of similar parts, as the wing, tail, etc., since those averaging the largest or

smallest for the four measurements given are often not those having the longest or the shortest wing, tail, or tarsus, or which measure the most or the least in length or alar extent. The extremes of variation in the size of the wing and tail is given in Tables B, C, and D.*

Table A.—Variation in General Size.

Mus. Comp. Zoology No.	Collector's Number.	Sex.	Name.	Locality.	Date.	Collected by	Length.	Alar Extent.	Wing.	Tail.
8844	693	♂	Turdus Swainsoni	Belmont, Mass.	May 27, '68	C. J. Maynard	7.76	12.65	4.20	3.00
1520	—	—	Turdus Swainsoni	Springfield, "	May 14, '62	J. A. Allen	6.62	11.40	3.80	2.83
—	250	♂	Turdus Pallasi	Newton, "	Apr. 25, '68	C. J. Maynard	7.30	12.83	3.80	3.83
—	608	♂	Turdus Pallasi	Newton, "	May 25, '68	C. J. Maynard	7.00	10.64	3.00	2.45
8830	337	♂	Turdus fuscescens	Newton, "	May 5, '68	C. J. Maynard	7.81	13.70	4.16	4.00
8834	553	♂	Turdus fuscescens	Watertown, "	May 20, '68	C. J. Maynard	7.00	11.95	3.58	3.55
—	621	♂	Dendroica striata	Newton, "	May 27, '68	C. J. Maynard	5.45	8.75	2.80	1.70
—	641	♂	Dendroica striata	Newton, "	May 27, '68	C. J. Maynard	5.51	9.30	3.10	2.00
—	543	♂	Dend. pennsylvanica	Newton, "	May 20, '68	C. J. Maynard	5.40	8.25	2.61	2.07
—	451	♂	Dend. pennsylvanica	Newton, "	May 20, '68	C. J. Maynard	5.00	7.56	2.35	1.75
4374	1011	♂	Spizella pusilla	Newton, "	July 22, '68	C. J. Maynard	6.00	8.32	2.62	2.60
4373	—	♂	Spizella pusilla	Waltham, "	Sept. 19, '67	C. J. Maynard	5.06	7.62	2.00	2.62
4990	1411	♂	Sayornis fuscus	Newton, "	Sept. 30, '68	C. J. Maynard	7.50	11.40	3.60	3.00
—	104	♂	Sayornis fuscus	Waltham, "	Mar. 28, '68	C. J. Maynard	7.25	12.60	3.82	2.20
4819	5	♂	Sayornis fuscus	Newton, "	Oct. 9, '69	C. J. Maynard	6.51	10.32	3.20	2.85
4701	317	♂	Passerculus savanna	Newton, "	Apr. 5, '68	C. J. Maynard	5.85	9.73	3.00	2.26
5093	873	♂	Passerculus savanna	Ipswich, "	June 17, '68	J. A. Allen	5.83	7.75	2.80	2.10
—	890	♀	Passerculus savanna	Ipswich, "	June 17, '68	J. A. Allen	5.35	7.75	2.51	1.95
5088	851	♀	Passerculus savanna	Ipswich, "	June 17, '68	J. A. Allen	5.75	9.75	2.78	2.05
—	714	♂	Vireo olivaceus	Waltham, "	May 30, '68	C. J. Maynard	6.55	10.65	3.48	2.46
—	713	♂	Vireo olivaceus	Waltham, "	May 30, '68	C. J. Maynard	6.25	9.88	3.22	2.10
4624	1407	♂	Chrysomitris tristis	Newton, "	Sept. 28, '68	C. J. Maynard	5.35	9.40	3.00	2.00
4926	150	♂	Chrysomitris tristis	Newton, "	Mar. 23, '68	C. J. Maynard	5.00	8.60	2.65	1.85
—	106	♂	Melospiza melodia	Newton, "	Mar. 12, '68	C. J. Maynard	6.75	9.15	2.65	2.80
—	55	♂	Melospiza melodia	Newton, "	Nov. 24, '67	C. J. Maynard	6.00	8.25	2.35	2.68
1456	—	♂	Sialia sialis	Worthington, "	July —, '63	C. H. Hamlin	6.50	11.10	3.75	2.45
2378	—	♂	Sialia sialis	Waterville, Me.	Apr. 15, '64	W. H. Niles	7.00	12.25	4.10	2.60
4371	334	♂	Mniotilta varia	Newton, Mass.	Apr. 20, '68	C. J. Maynard	5.40	9.00	3.00	2.15
4376	372	♂	Mniotilta varia	Waltham, "	May 6, '68	C. J. Maynard	5.35	8.25	2.87	2.05
—	887	♂	Cotyle riparia	Ipswich, "	June 17, '68	J. A. Allen	5.45	11.00	4.65	2.10
5111	868	♂	Cotyle riparia	Ipswich, "	June 17, '67	J. A. Allen	5.25	10.55	3.75	1.86
—	1082	♂	Passerella iliaca	Springfield, "	Mar. 22, '68	J. A. Allen	7.50	11.65	3.65	2.98
—	—	♂	Passerella iliaca	Newton, "	Mar. 27, '68	C. J. Maynard	6.80	10.80	3.40	2.50
—	783	♂	Icterus Baltimore	Newton, "	June 6, '68	C. J. Maynard	8.00	12.25	4.00	3.10
4852	653	♂	Icterus Baltimore	Newton, "	May 27, '68	C. J. Maynard	7.30	11.15	3.56	2.55
—	946	♂	Sterna hirundo	Muskeget Isl.	June 23, '68	J. A. Allen	15.50	31.85	11.30	7.00
10485	916	♂	Sterna hirundo	Muskeget Isl.	June 23, '68	J. A. Allen	14.90	29.30	10.40	5.50
10464	932	♂	Sterna arctica	Muskeget Isl.	July 2, '68	J. A. Allen	16.1	32.15	11.60	7.50
—	992	♂	Sterna arctica	Muskeget Isl.	July 2, '68	J. A. Allen	14.40	29.00	10.70	6.30
4009	—	♂	Pipilo erythrophthal.	Milton, Mass.	—	H. C. Daring	8.55	12.25	3.57	3.85
10151	—	♂	Pipilo erythrophthal.	Cambridge, "	—	L. Agassiz	7.50	10.20	3.34	3.60

As a large proportion of the specimens mentioned in some of the following tables (most of Tables A to G) were taken during the season of their migration, they may have originated at widely different localities, and thus the differences indicated may be in some measure due to geographical causes. In other cases, however, all the specimens

* The measurements given in this paper were all taken either from fresh specimens by the collector, or by myself from specimens preserved in spirits.

were taken in the breeding season; while in still other instances (Tables H to P) the species were purposely chosen from among such as find their northern limit of distribution near the locality where all were taken. Of ten species of the latter class, twenty perfect male specimens have been carefully measured,* the measurements embracing a series of eighteen to twenty distinct parts; under such circumstances the variation in general size, in length, in alar extent, in the length of the folded wing, the tail, the tarsus, the head, the bill, etc., etc., commonly ranged from twelve to eighteen per cent.

In respect to the differences in the general form of the body, two leading styles of variation from the average form may be recognized in nearly all species, namely, a relatively robust form, in which the stoutness extends to all parts, and a relatively slender form, in which the slenderness is equally general. Variations of this general and symmetrical character are remarkable only for their extent, since in such cases there are no marked discrepancies between the relative size of different parts. Contrary, however, to our usual notions of exact symmetry in animals, the unsymmetrical variations are by far the most frequent and important.

Variation in the Relative Size of Different Parts. — In specimens of average size of any given species, considerable differences exist in the relative size of different parts. In individuals of the average alar extent of their species, for example, the length of the folded wing may vary very considerably, in consequence of a difference in the length of the primary quills as compared to the length of the bones of the wing. The length of the folded wing or the alar extent may vary with reference to the whole length of the specimen, in consequence of differences in the relative length of the tail, the neck, or the body. The tarsus also varies independently of variation in the general size, as do also the toes to the tarsi, relatively short toes being found to accompany tarsi of ordinary length, and, conversely, long toes short tarsi.† The wing varies in its form in consequence of the different relative development of the primary and secondary quills.‡ The tail varies in respect to its form, especially in regard to the degree of its emargination or graduation, and, in some groups, in respect to the number of its feathers. The bill also varies greatly in size and form. The variations in these various parts will be considered separately and in detail.

* See below, Tables H to P, pp. 210-219.

† See Table E, p. 204.

‡ See Table F, p. 205.

Variation in the Length of the Folded Wing and the Tail.—The measurements given in the following table (Table B) sufficiently illustrate the variation in the length of the folded wing in fully mature specimens of the same sex and species, while Table C indicates the variation in the length of the tail, in specimens of a similar character. All the specimens, with a few exceptions, were taken within a few miles of Cambridge; the others are mainly also from Eastern Massachusetts, a few * being from a single locality in Florida. The series from which these extremes are taken embrace ordinarily not more than twenty-five or thirty specimens; with larger suites the differences would in many cases doubtless be much increased. The largest and smallest only are taken, between which, however, there is every gradation. The difference between these extremes is indicated, and also the percentage of the variation, based on the average of the two extremes. The amount of the variation in the length of the folded wing ranges, as will be seen from the table, from twelve to twenty-one per cent of the average length. In the tail the amount of variation in respect to length ranges from fourteen to twenty-three per cent. The different species vary considerably in respect to the amount of variation each presents, some being much more variable than others. It should be stated, however, that as a general rule the widest extremes, or the highest percentages of variation, occur in those species of which the greatest number of specimens has been examined. It will also be noticed that the tail usually varies more than the wings. In species with a relatively long tail the percentage of variation in the length of this member is found to be greater than in those species in which it is of medium length or short, as would have been naturally expected. In several cases the greater differences occur between females, but this may be a mere coincidence.

In this connection it may be added that the variation proves to be much less between specimens of the same species and sex when taken at a single locality in the breeding season than when taken during the period of migration. In many instances specimens of the same species may be obtained at one locality which shall represent the whole range of its geographical variation, as well as its individual variation, as in the case of those species which breed far to the North, but migrate in winter to the tropics, being thus but transient visitors to the temperate portions of the United States.

* Those of *Mimus polyglottus*, *Cardinalis virginianus*, *Picus borealis*.

Table B.—*Individual Variation in the Length of the Folded Wing.*

M. C. Z. No.	Orig. No.	Sex		Folded Wing	Difference.	Percent of Variation.
10596	2510	♀	Mimus polyglottus	4.75	.75	17.0
—	2485	♀	Mimus polyglottus	4.00		
10716	1987	♂	Cardinalis virginianus	3.85	.55	14.6
—	1993	♂	Cardinalis virginianus	3.30		
—	316	♂	Passerculus savanna	2.95	.40	14.5
—	820	♂	Passerculus savanna	2.55		
8830	367	♂	Turdus fuscescens	4.16	.61	15.8
8834	556	♂	Turdus fuscescens	3.55		
4821	148	♂	Sayornis fuscus	3.87	.67	19.0
4819	5	♂	Sayornis fuscus	3.20		
9057	618	♂	Geothlypis trichas	2.56	.50	21.0
5020	703	♂	Geothlypis trichas	2.06		
4648	1389	♂	Carpodacus purpureus	3.70	.60	17.6
4655	751	♂	Carpodacus purpureus	3.10		
9696	—	♂	Pipilo erythrophthalmus	3.68	.51	14.6
1421	—	♂	Pipilo erythrophthalmus	3.17		
—	170	♂	Junco hyemalis	3.20	.45	18.0
4910	140	♂	Junco hyemalis	2.75		
1568	—	?	Tyrannus carolinensis	4.85	.68	15.0
10025	—	?	Tyrannus carolinensis	4.17		
10014	—	?	Galeoscoptes carolinensis	3.85	.60	17.0
2734	—	?	Galeoscoptes carolinensis	3.25		
786	—	♂	Icterus Baltimore	4.00	.58	16.0
1334	—	♂	Icterus Baltimore	3.42		

Table C.—*Individual Variation in the Length of the Tail.*

M. C. Z. No.	Orig. No.	Sex		Tail.	Difference.	Percent of Variation.
—	2474	♂	Mimus polyglottus	5.15	.95	20.5
10592	2372	♂	Mimus polyglottus	4.20		
—	1955	♂	Cardinalis virginianus	4.30	.90	23.4
—	2460	♂	Cardinalis virginianus	3.40		
—	317	♂	Passerculus savanna	2.26	.41	19.5
5086	846	♂	Passerculus savanna	1.85		
8830	528	♂	Turdus fuscescens	3.00	.45	14.4
8835	556	♂	Turdus fuscescens	2.55		
—	—	♀	Parus atricapillus	2.63	.48	20.0
—	—	♀	Parus atricapillus	2.15		
9056	454	♂	Geothlypis trichas	2.15	.45	23.4
5020	703	♂	Geothlypis trichas	1.70		
4651	1071	♀	Carpodacus purpureus	2.57	.52	22.5
4653	1371	♀	Carpodacus purpureus	2.05		
4614	1330	♂	Pipilo erythrophthalmus	4.00	.71	19.5
4727	415	♂	Pipilo erythrophthalmus	3.29		
—	160	♂	Junco hyemalis	2.78	.38	15.0
4917	201	♂	Junco hyemalis	2.40		
10646	1972	♀	Picus borealis	3.75	.50	14.0
10633	41	♀	Picus borealis	3.25		
1317	—	?	Tyrannus carolinensis	2.93	.61	19.0
1568	—	?	Tyrannus carolinensis	3.54		
2734	—	?	Galeoscoptes carolinensis	3.35	.75	20.0
10014	—	?	Galeoscoptes carolinensis	4.10		
1334	—	♂	Icterus Baltimore	2.70	.40	13.8
2289	—	♂	Icterus Baltimore	3.10		

Variation in the Relative Length of the Wings and Tail.—Table D illustrates the irregularity of the variation in the wings and the tail. The first column of measurements gives the length of the folded wing, and

Table D.—Individual Variation in the relative Length of the Folded Wing and Tail.

M. C. Z. No.	Original No.	Sex.		Wing.	Tail.	Diff. betw'n Wing and Tail.	Amount of Variation.
—	2429	♂	Mimus polyglottus	4.35	4.35	.00	1.20
10590	2342	♀	Mimus polyglottus	3.25	4.35	+1.00	
—	2560	♀	Mimus polyglottus	4.15	4.35	+ .25	
—	2614	♂	Mimus polyglottus	4.40	4.90	+ .50	
—	2340	?	Mimus polyglottus	4.40	4.50	+ .10	
—	2478	♀	Mimus polyglottus	4.40	4.20	— .20	.45
—	2374	♂	Mimus polyglottus	4.30	4.16	— .14	
8881	441	♂	Galeoscoptes carolinensis	3.60	3.60	.00	
—	1376	♀	Galeoscoptes carolinensis	3.70	3.60	— .10	
8879	412	♂	Galeoscoptes carolinensis	3.75	4.10	+ .35	
8841	495	♂	Turdus fuscescens	4.00	3.00	1.00	1.15
8832	332	♂	Turdus fuscescens	4.10	4.00	.10	
8835	581	♂	Turdus fuscescens	4.15	2.90	1.25	
8821	374	♂	Seiurus aurocapillus	3.00	2.00	1.00	
—	423	♂	Seiurus aurocapillus	3.00	2.66	1.66	
8851	322	♀	Turdus Pallasi	3.50	2.60	.90	.67
—	301	♂	Turdus Pallasi	3.43	3.17	.23	
4301	514	♂	Dendroeca aestiva	2.85	1.80	1.05	
—	362	♂	Dendroeca aestiva	2.45	1.98	.47	
5058	707	♂	Dendroeca striata	2.85	2.00	.85	
—	1341	♂	Dendroeca striata	3.00	1.75	1.25	.79
5062	734	♀	Dendroeca striata	2.45	1.98	.46	
—	741	♀	Dendroeca striata	2.80	1.80	1.00	
5041	665	♂	Setophaga ruticilla	2.60	2.10	— .50	
—	698	♂	Setophaga ruticilla	2.43	2.50	+ .07	
—	693	♀	Regulus satrapa	2.20	1.52	.68	.49
—	50	♀	Regulus satrapa	1.94	1.75	.19	
4808	711	♂	Contopus virens	3.35	2.36	.99	
4994	1116	♂	Contopus virens	3.15	2.70	.45	
10645	1924	♂	Picus borealis	4.80	3.32	1.48	.48
10646	1972	♀	Picus borealis	4.75	3.75	1.00	
4587	323	♂	Agelaius phoeniceus	4.85	3.40	1.45	
4589	214	♂	Agelaius phoeniceus	4.60	3.82	.78	
4654	1069	♂	Carpodacus purpureus	3.85	2.00	1.35	
4655	286	♂	Carpodacus purpureus	3.03	2.32	.71	.64
—	288	♂	Poocætes gramineus	3.55	2.41	1.14	
—	846	♀	Poocætes gramineus	3.10	2.50	.60	
—	881	♂	Passerculus savanna	2.75	1.85	.90	
—	127	♂	Passerculus savanna	2.74	2.25	.49	
—	115	♂	Passerella iliaca	3.75	2.65	1.10	.78
—	55	♀	Passerella iliaca	3.32	3.00	.32	
—	177	♂	Melospiza melodia	2.35	2.68	+ .33	
—	2363	♂	Melospiza melodia	2.60	2.40	— .20	
—	2369	♂	Cardinalis virginianus	3.60	3.40	— .20	
—	—	♂	Cardinalis virginianus	3.60	4.10	+ .50	.70
2293	—	♂	Dolichonyx oryzivora	3.75	2.78	.96	
5741	—	♂	Dolichonyx oryzivora	4.00	2.72	1.28	
10107	—	♂	Hedymeles ludoviciana	4.20	2.93	1.27	
9787	—	♂	Hedymeles ludoviciana	3.83	2.95	.88	

the second the length of the tail, of the same specimens; the third column shows the difference in length between the tail and the wing, and the fourth column the amount of the difference between the two extremes. In *Mimus polyglottus* the tail is usually one fourth to one half an inch longer than the wing; but in many specimens the wings and tail are equal, and in a small proportion the tail is *shorter* than the wing. In the seven specimens of this species cited in the table, the variation ranges from the tail being one fifth of an inch shorter than the folded wing to one inch longer. In the three specimens which agree in the length of the tail (4.35 in.), the variation in the length of the folded wing ranges from 3.25 in. to 4.25 in., or is nearly twenty-seven (26.85) per cent. The larger specimen, however, is a male, while the others are females; but between the two females the difference is over twenty-four (24.3) per cent. Similar differences have been met with in various other species, but it has not been deemed necessary to cite a larger list of examples.

Variation in the Form of the Wing. — By the form of the wing is meant its general outline when expanded, which is mainly determined by the relative length of the remiges. The form of the wing, and especially the relative length of the different primary remiges, has direct relation to the power of flight. In strong, swift-flying birds, the outer primaries are the longest, giving a narrow pointed form to the expanded wing, as in the swifts, the swallows, in *Chordeiles*, in the *Sterninæ* and in most of the *Procellariidæ*. In birds of medium powers of flight, as in most of the true finches (*Coccothraustinæ*) and *Turdinæ*, the *Tyrannidæ*, the *Sylvicolidæ*, etc., etc., the third, fourth, and fifth primaries are the longest, the wing being less pointed and broader. In species with low power of flight, as the *Troglodytidæ*, several genera of sparrows, the grouse, etc., the outer primaries are still more reduced, the wing is much more rounded and shorter, and the power of flight is in each case correspondingly less. In birds of the first class, which live almost wholly on the wing, little variation is seen in the relative length of the primaries. In those of the second and third classes, slight variations affect in less degree the particular habits of life, so that among the latter would be naturally expected the greatest range of individual variation.

Correlating with the variation in the form of the wing, as determined by the relative length of the outer primaries to the length of the inner

primaries are similar variations in the relative length of the inner secondaries as compared with the outer secondaries. Relatively short inner secondaries (generally improperly called "tertiaries") hence accompany long primaries, and, conversely, long inner secondaries, short outer primaries. The particular form of the wing in any group depending upon the relative development of these several elements, they hence afford excellent generic characters; but while thus important, they are subject to a considerable range of individual variation. The form of the wing being readily determined by measurements, and easily expressed mathematically, the amount of the variation is easily measured and tabulated. In the following table (Table E) the extent and character of this variation is to some degree illustrated. In the first column of measurements is given the length of the folded wing; in the second the extent of the longest primary beyond the outer (or shortest) secondary, and in the third the extent of the longest primary beyond the inner (or longest) secondary. The fourth column gives the amount of variation in each specimen cited.

Table E.—Variation in the Form of the Wing.

M. C. Z. No.	Sex.	Species.	Length of the Wing.	Ext. of Pr. beyond Outer Sec.	Ext. of Pr. beyond Inner Sec.	Amount of Variation.
2319	♂	Icterus Baltimore	3.75	.77	.90	.13
2290	♂	Icterus Baltimore	3.83	.67	.81	.14
1333	♂	Icterus Baltimore	3.64	.57	1.06	.49
1567	♂	Icterus Baltimore	3.80	.56	.92	.36
2964	♂	Icterus Baltimore	3.85	.77	1.07	.30
2299	♂	Icterus Baltimore	3.85	.87	1.12	.25
2296	♂	Dolichonyx oryzivora	3.80	.98	1.42	.44
5741	♂	Dolichonyx oryzivora	4.00	1.20	1.40	.20
119	♂	Dolichonyx oryzivora	3.82	.78	1.23	.45
9854	♂	Dolichonyx oryzivora	3.53	.98	1.14	.16
284	?	Tyrannus carolinensis	4.30	.85	1.15	.30
113	?	Tyrannus carolinensis	4.60	.90	1.45	.55
1317	?	Tyrannus carolinensis	4.25	.76	1.10	.34
4008	?	Tyrannus carolinensis	4.60	1.35	1.62	.27
10107	♂	Hedymeles ludoviciana	4.20	.90	1.05	.15
590	♂	Hedymeles ludoviciana	4.00	.90	1.25	.35
9935	♂	Hedymeles ludoviciana	4.00	.60	1.06	.40
1456	♂	Sialia sialis	3.75	1.00	1.10	.10
1945	♂	Sialia sialis	3.90	1.03	1.10	.07
333	♂	Sialia sialis	4.07	1.30	1.30	.00
5606	♂	Sialia sialis	4.05	1.25	1.40	.15
10292	♂	Sialia sialis	3.90	.95	1.15	.20
256	?	Galeoscoptes carolinensis	3.37	.55	.50	— .05
1790	?	Galeoscoptes carolinensis	3.75	.55	.70	+ .15
5858	?	Galeoscoptes carolinensis	3.55	.35	.57	+ .22
10014	?	Galeoscoptes carolinensis	3.85	.70	.75	+ .05
2274	?	Galeoscoptes carolinensis	3.75	.67	.70	+ .03

Variation in the relative Length of the Primary Quills. — From the great stress laid upon the relative length of the outer primaries by descriptive ornithologists in determining genera and species, one would be led to expect but a slight amount of variation in this respect in specimens of the same species. On the contrary, however, it is soon found, on giving special attention to this character, that a considerable amount of individual variation in this regard really exists. That the wing formula, so generally introduced of late years into specific diagnoses, is in a great degree unreliable as a specific character, is sufficiently shown by the subjoined table (Table F, p. 206) of the relative proportions of the primaries. The comparison, extended in the table to only a few species has been carried to scores of others with similar results.

In general, in species of the *Oscines* which have the second primary usually the longest, it is sometimes the first and sometimes the third that is the longest. In those which have the third ordinarily the longest, the second and third, the third and the fourth, or the second, third, and fourth are frequently equal. In those in which the first (or the second when the first is very short) is intermediate to the second and fourth or to the third and fifth, it may be equal to or longer than the second or third, or only equal to the fourth or fifth.

Variation in the Form of the Tail, and in the Number of the Rectrices. — Individual variation in the form of the tail is often quite marked. In species with the tail deeply forked, different specimens vary considerably in respect to the depth of the fork. Those with the tail rounded and much graduated differ greatly in respect to the amount the middle feathers exceed the outer ones in length. In species with a normally nearly even tail, the tail is sometimes distinctly emarginate, and sometimes as distinctly rounded in different specimens of the same species.

In regard to the number of rectrices, in those groups in which the number exceeds twelve, as in the *Rasores*, the *Lamellirostres*, etc., the number is frequently variable. The rectrices of the common ruffed grouse (*Bonasa umbellus*) are usually eighteen in number, but an examination of numerous specimens shows that the number varies from sixteen to twenty. The usual number in *Tetrao canadensis* is sixteen, but the number varies from fourteen to eighteen. In *Cupidonia cupido*, and in other species of grouse, similar variations also occur. They are also frequent in the *Anserinæ*. In *Bernicla canadensis*, for example, the usual number of rectrices is eighteen, but the number

Table F. — Variation in the relative Length of the Primaries.

M. C. Z. No	Species.	Longest.	2d in Length.	3d in Length.	4th in Length.	5th in Length.	6th in Length.	7th in Length.	8th in Length.	9th in Length.
3289	<i>Turdus fuscescens</i>	4	3	2	5	6	7			
6764	<i>Turdus fuscescens</i>	3	4	2	5	6	7			
8837	<i>Turdus fuscescens</i>	4 } 3 }	2	5	6	7	8			
8848	<i>Turdus Pallasi</i>	4	3	5	6	2	7			
5197	<i>Turdus Pallasi</i>	4	5	3	6	2	7			
8205	<i>Turdus Pallasi</i>	4	3	5	6 } 1 }	7	8			
8206	<i>Turdus Pallasi</i>	4	3 } 5 }	6	2	7	8			
10698	<i>Myiarchus crinitus</i>	3	4	2	5	6	1	7		
10699	<i>Myiarchus crinitus</i>	3	2	4	5	6	1	7		
8166	<i>Myiarchus crinitus</i>	4	3	2	5	6	1 } 7 }	8		
10700	<i>Myiarchus crinitus</i>	3 } 4 }	2	5	6	1	7	8		
10701	<i>Myiarchus crinitus</i>	3	4 } 2 }	5	6	1	7	8		
12420	<i>Tyrannus carolinensis</i>	2	3	1	4	5	6			
4612	<i>Tyrannus carolinensis</i>	2	3	4	1	5	6			
6457	<i>Tyrannus carolinensis</i>	2	3	1 } 4 }	5	6	7			
4816	<i>Contopus borealis</i>	2	1	3	4	5	6			
6665	<i>Contopus borealis</i>	2	3	1	4	5	6			
6938	<i>Sayornis fuscus</i>	3	4	2	5	6	1	7		
6932	<i>Sayornis fuscus</i>	3 } 4 }	2 } 5 }	6	1	7	8	9		
5364	<i>Lophophanes bicolor</i>	4	5	6	3	7	8	9	2	1
5248	<i>Lophophanes bicolor</i>	5	4	6	7	3	8	9	2	1
—	<i>Lophophanes bicolor</i>	5 } 4 } 6 }	7	3	8	9	2	1		
5080	<i>Dendroeca coronata</i>	2	3	1	4	5	6			
5176	<i>Dendroeca coronata</i>	3	2	4	1	5	6			
6678	<i>Dendroeca coronata</i>	3	2 } 4 }	1	5	6	7			
3412	<i>Dendroeca coronata</i>	2 } 3 }	4	1	5	6	7			
10533	<i>Dendroeca coronata</i>	2 } 1 } 3 } 4 }	5	6	7	8	9			
5056	<i>Dendroeca striata</i>	2	1	3	4	5	6			
5057	<i>Dendroeca striata</i>	1	2	3	4	5	6			
3390	<i>Dendroeca striata</i>	2	1 } 3 }	4	5	6	7			
6675	<i>Dendroeca striata</i>	2	3	1	4	5	6			
10958	<i>Pinicola enucleator</i>	3	4	2	5	1	6	7		
10963	<i>Pinicola enucleator</i>	3	2	4	1	5	6	7		
10960	<i>Pinicola enucleator</i>	3	2	1	4	5	6	7		
10962	<i>Pinicola enucleator</i>	3 } 2 }	4	1	5	6	7	8		
8114	<i>Pinicola enucleator</i>	3	4 } 2 }	1	5	6	7	8		
4843	<i>Ampelis cedrorum</i>	2	3	1	4	5				
4633	<i>Ampelis cedrorum</i>	1	2	3	4	5				
4844	<i>Ampelis cedrorum</i>	2	1	3	4	5				

varies from fourteen to twenty. Specimens with sixteen are tolerably frequent. Yet one of the principal characters urged as separating the *B. Hutchinsii* from the *B. canadensis* is the possession of two more feathers in the tail by the latter than the so-called *B. Hutchinsii* is assumed to have. In *Bernicla brenta* the usual number is sixteen, but in different specimens they vary from fourteen to eighteen. A greater, or less amount of variation in the number of the feathers of the tail is more or less common to numerous other species of the duck tribe. An odd number is even quite frequent, one half of the tail having normally one more feather than the other.

Variation in the Relative Length of the Tarsus and Toes.—A common feature in modern generic and specific diagnoses is a statement of the ratio the length of the tarsus bears to the length of the middle toe or to the hallux, and the relative length of the hallux to the outer or inner toe, as though we had here constant structural proportions. The following table (Table G) shows that such is not the case, the varia-

Table G.—Relative Length of Tarsi and Toes.

M. C. Z. No.	Sex.	Species.	Tarsus.	Middle Toe.	Outer Toe.	Hallux.
5858	?	Galeoscoptes carolinensis	1.08	1.04	.77	.75
2273	?	Galeoscoptes carolinensis	1.15	.98	.70	.70
10356	?	Galeoscoptes carolinensis	1.00	1.00	.70	.73
5857	?	Galeoscoptes carolinensis	1.18	1.08	.70	.75
2229	?	Galeoscoptes carolinensis	1.07	.93	.68	.67
5605	♂	Sialia sialis	.77	.77	.57	.58
1456	♂	Sialia sialis	.74	.84	.62	.65
5766	♂	Sialia sialis	.83	.80	.56	.60
1883	♂	Sialia sialis	.80	.91	.77	.65
1946	♂	Sialia sialis	.80	.84	.77	.61
1881	♂	Sialia sialis	.77	.85	.56	.72
1771	♂	Pipilo erythrophthalmus	.98	.95	.73	.80
1399	♂	Pipilo erythrophthalmus	1.05	1.05	.80	.78
350	♂	Pipilo erythrophthalmus	1.05	1.12	.76	.84
1476	♂	Pipilo erythrophthalmus	1.10	1.03	.75	.78
2985	♂	Pipilo erythrophthalmus	1.13	1.00	.80	.80
9854	♂	Dolichonyx oryzivora	.98	1.17	.83	.82
5585	♂	Dolichonyx oryzivora	1.15	1.27	.98	.93
9894	♂	Dolichonyx oryzivora	1.00	1.00	.83	.81
10219	♂	Dolichonyx oryzivora	1.03	1.25	.98	.76
2320	♂	Icterus Baltimore	.83	.68	.88	.72
9793	♂	Icterus Baltimore	1.02	.85	.70	.70
1567	♂	Icterus Baltimore	.97	1.00	.75	.80
10025	?	Tyrannus carolinensis	.67	.73	.53	.54
10027	?	Tyrannus carolinensis	.80	.85	.55	.61
10028	?	Tyrannus carolinensis	.70	.87	.53	.57
5546	?	Tyrannus carolinensis	.70	.80	.60	.60

tion being as great between different specimens of the same species as between different species of the same genus, and even of different genera. The variation in the length of the toes is often due to an increase or a decrease in the length of the nail, but by no means rarely to variations in the length of the phalanges themselves. As already stated, and as appears from the table, toes of less than the average length accompany tarsi of the average or of more than the average length, and toes of more than the average length accompany tarsi of medium or less than the medium length. In compiling the above table the specimens mentioned have been selected in each case from a series of only twenty specimens of the species to which they respectively belong, and represent the longest and shortest tarsus, middle toe, outer toe, and hind toe met with in each series, and also the greatest and least amount of difference in these several elements. They are all taken from Tables H to Q (see pp. 210-219), which serve to show the usual range of variation, in respect to size and proportions, in ten species.*

Individual Variation in other Parts.—In addition to the instances already mentioned, individual variation of a similar character and equal extent occurs in the relative size of other parts. The length of the bill, for instance, is often compared to the length of the head, or to that of the tarsus in specific diagnoses. Table G¹ (see next page) serves to show the individual variation in respect to the proportion of length to alar extent ordinarily met with in specimens of the same species.

To show more fully, however, the exact nature and extent of what may be considered as purely individual variation, tables of detailed measurements of about twenty specimens of each of a number of species are herewith appended (Tables H to Q). Care has been taken to not only select specimens of the same sex, collected at the same locality, and as nearly as possible at the same season, but also such species as find their northern limit so near the locality at which they were taken as to obviate the complication of individual with geographical variation, which would result if the range of the species extended far to the northward of the locality in question. In general, the specimens are all from Eastern Massachusetts, and

* *Icterus* Baltimore, *Dolichonyx oryzivorus*, *Pipilo erythrophthalmus*, *Sialia sialis*, *Galeoscoptes carolinensis*, *Pyrrhuloxia rubra*, *Geothlypis trichas*, *Harporhynchus rufus*, *Tyrannus carolinensis*, *Hedymeles ludoviciana*.

Table G¹ — Individual Variation in the Proportion of Length to Alar Extent.

M. C. Z. No.	Orig. No.	Sex.	Species.		
5056	668	♀	Dendroeca striata	5.45	9.70
—	777	♀	Dendroeca striata	5.50	8.68
5087	848	♂	Passerculus savanna	5.50	9.13
—	981	♂	Passerculus savanna	5.83	7.75
—	—	♂	Passerculus savanna	6.00	8.27
—	1987	♂	Cardinalis virginianus	9.00	11.50
—	2394	♀	Cardinalis virginianus	8.00	11.75
9901	—	♂	Dolichonyx oryzivorus	6.65	11.50
2295	—	♂	Dolichonyx oryzivorus	7.50	11.50
—	2340	?	Mimus polyglottus	9.60	14.25
—	2374	♀	Mimus polyglottus	9.75	14.00
—	2371	♀	Mimus polyglottus	9.80	13.00
5757	—	?	Turdus Swainsoni	7.25	12.15
2930	—	?	Turdus Swainsoni	7.75	11.20
1829	—	?	Turdus Swainsoni	6.90	11.20
307	—	?	Turdus Swainsoni	7.24	11.00
9691	—	?	Turdus Pallasi	7.00	10.50
145	—	?	Turdus Pallasi	7.00	11.40
5756	—	?	Turdus Pallasi	7.38	11.05
—	314	♂	Turdus Pallasi	7.38	12.33
—	363	♂	Turdus Pallasi	7.23	11.94
—	26	♂	Turdus Pallasi	6.80	11.28
—	367	♂	Turdus fuscescens	7.81	13.70
—	495	♂	Turdus fuscescens	7.87	11.91
—	551	♂	Turdus fuscescens	7.00	11.95
—	112	♂	Parus atricapillus	5.50	8.12
4946	268	♀	Parus atricapillus	5.00	8.60
11714	114	♀	Parus atricapillus	5.75	7.88
95	—	♂	Agelæus phæniceus	9.00	15.10
93	—	♂	Agelæus phæniceus	9.20	14.40
5723	—	♂	Agelæus phæniceus	8.45	14.45

from within a short distance of Cambridge. A very few are from Southern Maine and from the Connecticut valley at Springfield; but the general faunal character of all these localities is essentially the same.*

In addition to the measurements given in these tables, several others are sometimes taken by collectors, as the relative posterior extent of the outstretched feet and the wing, as compared with the tail. As they are, however, among the most variable of proportions, and are likewise among the most difficult measurements to take with accuracy, they have been here neglected.

* In consequence of the small size of these pages, it has been found impracticable to include the names of the localities, the date of collecting, and the name of the collector in the tables, as would have been desirable.

Table H. — *ICTERUS BALTIMORE Daud.*

M. C. Z. Number.	Sex.	Length.	Alar Extent.	Wing.	Tail.	Extent of Primaries beyond Inner Secondaries.	Extent of Primaries beyond Outer Secondaries.	Extent of Tail beyond Upper Coverts.	Extent of Tail beyond Lower Coverts.	Tarsus.	Hind Toe.	Middle Toe.	Outer Toe.	Inner Toe.	Head.	Bill.				
																Culmen.	Commissure.	Gonys.	Height.	Width.
1332	♂	7.45	11.75	3.70	2.90	.70	.97	1.50	1.40	.96	.72	.89	.67	.65	1.62	.76	.77	.47	.32	.33
9793	♂	7.50	11.60	3.70	2.95	.76	1.07	1.67	1.33	1.02	.70	.85	.70	.66	1.58	.76	.77	.47	.38	.35
1333	♂	7.20	11.20	3.64	2.88	.57	1.06	1.52	1.40	.93	.67	.85	.70	.66	1.62	.78	.81	.50	.35	.32
2289	♂	8.00	11.90	3.85	3.10	.87	1.12	1.77	1.65	.91	.69	.95	.75	.73	1.63	.75	.77	.52	.37	.34
9884	♂	7.00	10.40	3.54	2.82	.86	.93	1.37	1.15	.87	.65	.93	.70	.69	1.60	.73	.46	.46	.33	.35
2733	♂	7.55	11.30	3.75	3.00	.82	1.03	1.82	1.18	.87	.70	.92	.70	.68	1.60	.80	.83	.48	.34	.35
2962	♂	7.50	11.50	3.72	3.00	.68	.95	1.63	1.38	.90	.70	.87	.70	.65	1.60	.77	.78	.48	.37	.35
1567	♂	7.60	11.75	3.80	3.00	.56	.92	1.53	1.40	.97	.75	1.00	.80	.78	1.63	.77	.80	.50	.32	.32
1334	♂	7.20	11.00	3.45	2.70	.60	.84	1.45	1.20	.90	.70	.90	.73	.73	1.55	.75	.77	.48	.35	.35
1566	♂	7.40	11.50	3.60	2.90	—	—	1.55	1.10	.90	.73	.86	.73	.70	1.60	.82	.82	.53	.35	.34
2993	♂	7.60	11.90	3.75	2.97	.68	.96	1.43	1.30	.91	.70	.90	.70	.67	1.58	.78	.78	.52	.34	.33
388	♂	7.40	11.00	3.68	2.78	.85	1.00	1.58	1.27	.95	.70	.88	.75	.72	1.62	.78	.77	.51	.37	.35
2320	♂	7.30	11.50	3.60	2.75	.77	.95	1.30	1.25	.83	.68	.88	.72	.68	1.50	—	—	—	—	—
2964	♂	7.75	11.90	3.85	3.00	.77	1.07	1.57	1.50	.90	.62	.82	.73	.68	1.62	.75	.76	.50	.32	.33
2290	♂	7.80	11.75	3.75	2.95	.62	.85	1.67	1.35	.86	.68	.87	.64	.61	1.60	.77	.78	.45	.34	.34
2499	♂	7.75	11.75	3.78	2.96	.90	1.03	—	—	.94	.70	.82	.68	.65	1.55	.72	.74	.44	.38	.38
2290	♂	7.55	11.75	3.83	2.95	.67	.81	1.58	1.37	.95	.72	.80	.67	.67	1.62	.82	.84	.50	.40	.38
2319	♂	7.65	11.75	3.75	2.85	.77	.90	1.54	1.40	.95	.70	.98	.70	.73	1.58	—	—	.50	.35	.33
9885	♂	7.50	11.75	3.55	3.00	.78	.95	1.87	1.65	.90	.70	.90	.75	.75	1.60	.80	.82	.47	.35	.35
2500	♂	7.65	12.00	3.83	3.00	—	—	—	—	.98	.72	.90	.75	.72	1.53	.75	.78	.50	.37	.37

Table I. — *DOLICHONYX ORYZIVORUS Swain.*

M. C. Z. Number.	Sex.	Length.	Alar Extent.	Wing.	Tail.	Extent of Primaries beyond Inner Secondaries.	Extent of Primaries beyond Outer Secondaries.	Extent of Tail beyond Upper Coverts.	Extent of Tail beyond Lower Coverts.	Tarsus	Hind. Toe.	Middle Toe.	Outer Toe.	Inner Toe.	Head	Bill.				
																Culmen.	Commissure.	Gonyx.	Height.	Width.
2293	♂	7.50	11.50	3.75	2.68	.93	1.27	1.10	1.00	1.05	.87	1.12	.77	.80	1.32	.60	.63	.34	.36	.36
10342	♂	7.00	11.00	3.64	2.45	.88	1.20	.80	—	1.07	.92	1.17	.73	.75	1.37	.62	.65	.35	.37	.37
10343	♂	7.30	11.80	3.87	2.80	.90	1.30	.95	.90	1.12	.95	1.28	.87	.87	1.37	—	—	.33	.34	.37
5585	♂	7.35	11.55	3.80	2.66	.97	1.40	—	.83	1.15	.98	1.27	.93	.87	1.35	.62	.62	.35	.36	.36
2295	♂	7.50	11.50	3.71	2.73	1.05	1.33	.93	.90	1.00	.92	1.18	.85	.85	1.37	.62	.62	.35	.36	.36
119	♂	7.25	11.65	3.82	2.57	.78	1.23	—	.80	1.00	.83	1.12	.76	.78	1.34	.60	.60	.35	.38	.37
5743	♂	7.35	11.75	3.65	2.68	1.03	1.33	—	.80	1.10	.92	1.13	.78	.80	1.38	.60	.63	.36	.37	.40
9901	♂	6.65	11.50	3.70	2.53	.82	1.25	.95	.90	1.00	.85	1.07	.70	.76	1.35	.57	.60	.35	.35	.36
389	♂	7.20	11.20	3.55	2.50	.85	1.20	.95	.87	1.08	.86	1.10	.75	.78	1.40	.58	.62	.35	.37	.37
9854	♂	6.80	11.15	3.53	2.57	.98	1.14	1.03	.82	.98	.83	1.17	.82	.82	1.37	.60	.62	.35	.38	.38
2298	♂	7.70	11.80	3.75	2.78	1.03	1.35	1.07	.95	1.05	.90	1.12	.82	.85	1.37	.62	.64	.40	.38	.38
5741	♂	7.15	11.65	4.00	2.72	1.20	1.40	.97	.82	—	.87	1.13	.83	.82	1.36	.58	.60	.37	.38	.37
2296	♂	7.15	11.50	3.80	2.75	.98	1.42	.98	1.00	1.05	.82	1.20	.85	.88	1.33	.57	.63	.37	.38	.37
9893	♂	7.35	11.75	3.90	2.73	.98	1.37	1.15	.92	1.10	.95	1.27	.85	.85	1.36	.58	.65	.35	.38	.37
9892	♂	7.25	11.90	3.86	2.82	.97	1.35	1.10	.90	1.04	.87	1.23	.82	.83	1.33	.57	.62	.35	.37	.38
1482	♂	7.10	11.90	3.85	2.47	1.00	1.35	1.10	.95	1.05	.90	1.20	.88	.88	1.35	.60	.64	.35	.38	.37
9896	♂	7.25	12.15	3.90	2.67	1.05	1.40	1.08	.91	1.07	.85	1.17	.78	.78	1.36	.60	.60	.35	.36	.35
9894	♂	6.90	11.75	3.75	2.70	.95	1.32	.90	1.00	1.00	.83	1.00	.81	.82	1.35	.62	.64	.38	.39	.39
10219	♂	7.55	12.00	3.83	2.75	.98	1.33	1.23	1.05	1.03	.98	1.25	.76	.76	1.33	.60	.60	.37	.35	.39
5744	♂	7.45	11.35	3.85	2.82	1.08	1.43	1.00	1.05	1.10	.95	1.13	.82	.83	1.35	.60	.62	.37	.37	.39

Table J. — PIPILLO ERYTHROPHthalmus Vieillot.

M. C. Z. Number.	Sex.	Length.	Alar Extent.	Wing.	Tail.	Extent of Primaries beyond Inner Secondaries.	Extent of Primaries beyond Outer Secondaries.	Extent of Tail beyond Upper Coverts.	Extent of Tail beyond Lower Coverts.	Tarsus	Hind. Toe.	Middle Toe.	Outer Toe.	Inner Toe.	Head.	Bill.				
																Culmen.	Commissure.	Gonys.	Height.	Width.
350	♂	7.80	10.50	3.20	3.35	.33	.43	2.05	2.25	1.05	.76	1.12	.84	.82	1.37	.60	.67	—	.37	.37
1366	♂	8.25	10.65	3.43	3.70	.40	.50	2.25	2.15	1.03	.78	1.02	.85	.80	1.47	.56	.73	.40	.40	.37
1543	♂	8.00	11.00	3.60	3.90	.43	.48	2.35	2.27	1.05	.77	1.08	.88	.88	1.50	.60	.68	.40	.42	.34
1476	♂	7.80	10.30	3.20	3.40	.45	.45	2.38	2.18	1.10	.75	1.03	.78	.77	1.50	.60	.72	.37	.39	.35
1397	♂	8.25	10.25	3.45	3.70	.53	.58	2.60	2.13	1.12	.80	1.10	.82	.77	1.48	.53	.67	.34	.34	.38
1421	♂	8.10	10.60	3.17	3.48	.30	.20	—	1.85	1.07	.80	1.00	.78	.73	1.45	.53	.66	.40	.40	.33
349	♂	8.15	10.70	3.50	3.75	.43	.50	—	1.82	1.08	.80	1.00	.78	.77	1.46	.55	.70	.36	.38	.30
1399	♂	7.75	10.00	3.35	3.57	—	—	—	2.15	1.05	.80	1.05	.78	.77	1.46	.57	.70	.37	.35	.35
2985	♂	8.35	11.05	3.43	3.65	.35	.52	2.23	2.00	1.13	.75	1.00	.80	.80	1.50	.53	.72	.37	.35	.35
2965	♂	8.25	10.50	3.50	3.70	.45	.55	2.40	2.40	1.06	.81	1.08	.83	.78	1.49	.60	.70	.35	.37	.35
4009	♂	8.55	12.25	3.57	3.85	.47	.47	2.58	2.20	1.07	.77	1.04	.83	.78	1.40	.56	.70	.35	.35	.35
1783	♂	8.25	11.20	3.54	3.68	.45	.53	2.20	2.15	1.05	.80	1.04	.76	.68	1.50	.55	.73	.42	.35	.35
1612	♂	8.00	10.45	3.22	3.45	.30	.41	2.20	2.50	1.10	.80	1.10	.75	.72	1.50	.62	.73	.37	.32	.37
1771	♂	7.75	10.85	3.38	3.70	.35	.50	2.35	2.85	.98	.73	.95	.80	.73	1.45	.57	.68	.35	.33	.32
1784	♂	7.90	10.80	3.43	3.62	.40	.57	2.30	2.17	1.10	.78	1.08	.78	.73	1.50	.59	.72	.38	.38	.35
81	♂	8.25	11.15	3.54	3.80	.40	.45	2.53	2.30	1.08	.80	1.04	.80	.75	1.45	.55	.66	.38	.34	.35
83	♂	8.50	11.15	3.52	3.75	.42	.53	2.22	2.18	1.06	.80	1.05	.82	.75	1.50	.60	.73	.37	.41	.42
10151	♂	7.50	10.20	3.34	3.60	.46	.55	2.58	1.93	1.08	.75	1.12	.77	.75	1.45	.53	.62	.35	.37	.35
9696	♂	8.20	11.00	3.68	3.93	.50	.50	2.25	2.05	1.10	.78	1.07	.82	.82	1.50	.60	.73	.37	.42	.39
84	♂	8.15	10.25	3.30	3.70	.38	.47	—	2.06	1.02	.75	1.00	.83	.80	1.40	.60	.67	.37	.37	.35

Table K. — *SIALIA SIALIS* *Hild.*

M. C. Z. Number.	Sex.	Length.	Alar Extent.	Wing.	Tail.	Extent of Primaries beyond Inner Secondaries.	Extent of Primaries beyond Outer Secondaries.	Extent of Tail beyond Upper Coverts.	Extent of Tail beyond Lower Coverts.	Tarsus.	Hind Toe.	Middle Toe.	Outer Toe.	Inner Toe.	Head.	Bill.				
																Culmen.	Commissure.	Gonys.	Height.	Width.
2331	♂	7.00	12.30	4.00	2.50	1.00	1.20	1.30	1.20	.75	.63	.87	.68	.60	1.48	.48	.77	.30	.18	.35
1945	♂	6.75	12.20	3.90	2.60	1.03	1.11	1.56	1.04	.77	.56	.85	.72	.65	1.48	.50	.70	.33	.20	.34
2378	♂	7.00	12.25	4.10	2.60	1.20	1.26	1.27	1.05	.77	.60	.85	.64	.65	1.45	.50	.76	.32	.21	.33
2479	♂	6.10	11.35	3.87	2.50	1.10	1.20	1.25	1.05	.78	.60	.87	.65	.58	1.50	.56	.77	.30	.18	.35
1946	♂	6.85	12.10	4.03	2.58	1.03	1.20	1.33	1.13	.80	.77	.84	.61	.50	1.50	.53	.80	.32	.18	.37
1883	♂	6.85	12.55	4.07	2.56	1.17	1.26	1.20	1.15	.80	.77	.91	.65	.60	1.48	.50	.80	.30	.20	.38
10292	♂	6.45	11.75	3.90	2.60	.95	1.15	—	—	.76	.85	.82	.62	.55	1.35	.49	.72	.30	.20	.38
1881	♂	6.70	11.65	3.82	2.43	.98	1.13	1.20	1.08	.77	.67	.85	.56	.50	1.50	.56	.80	.33	.20	.33
72	♂	6.80	12.00	3.87	2.33	1.03	1.12	1.17	1.03	.75	.60	.85	.68	.62	1.50	.55	.78	.33	.20	.33
10226	♂	6.75	12.40	4.00	2.77	1.15	1.25	1.50	1.08	.77	.62	.83	.63	.58	1.50	.52	.80	.34	.20	.33
10231	♂	6.75	12.10	3.90	2.50	1.10	1.30	1.37	1.23	.75	.55	.80	.60	.54	1.42	.48	.76	.27	.18	.32
1456	♂	6.50	11.10	3.75	2.45	1.00	1.10	1.15	.95	.74	.62	.84	.65	.60	1.53	.55	.80	.35	.20	.35
5605	♂	6.70	12.00	3.85	2.35	1.10	1.25	1.08	.82	.77	.57	.77	.58	.54	1.50	.47	.77	.30	.18	.35
5763	♂	6.68	12.25	4.03	2.63	1.10	1.25	1.15	.90	.78	.55	.83	.58	.53	1.53	—	—	.32	—	—
5765	♂	6.65	11.80	3.90	2.63	1.05	1.17	1.58	1.23	.76	.57	.80	.61	.56	1.50	.55	.80	.31	.17	.30
5606	♂	6.50	11.65	4.05	2.68	1.25	1.40	1.27	1.18	.78	.60	.85	.63	.60	1.47	.53	.83	.32	.17	.34
338	♂	6.40	11.10	4.07	2.63	1.30	1.30	1.25	1.07	.80	.60	.88	.68	.62	1.50	.45	.77	.30	.21	.35
5760	♂	7.00	12.25	3.85	2.50	1.15	1.30	1.20	1.05	.80	.60	.83	.60	.55	1.45	.45	.70	.29	.18	.32
5766	♂	6.80	12.00	3.85	2.48	1.00	1.20	1.20	1.12	.83	.56	.80	.60	.58	1.45	.52	.70	.32	.16	.35
77	♂	6.75	11.75	3.95	2.55	1.15	1.30	—	—	.82	.60	.87	.65	.62	1.45	.45	.77	.30	.18	.34

Table L. — GALEOSOPTES CAROLINENSIS Cab.

M. C. Z. Number.	Sex.	Length.	Alar Extent.	Wing.	Tail.	Extent of Primaries beyond Inner Secondaries.	Extent of Primaries beyond Outer Secondaries.	Extent of Tail beyond Upper Coverts.	Extent of Tail beyond Lower Coverts.	Tarsus	Hind Toe.	Middle Toe.	Outer Toe.	Inner Toe.	Head.	Bill.				
																Culmen.	Commissure.	Gonys.	Height.	Width.
5858	?	8.75	11.35	3.55	3.75	.35	.57	2.20	2.05	1.08	.75	1.04	.77	.73	1.75	.80	1.00	.45	.23	.37
1790	?	8.75	11.75	3.75	3.85	.55	.70	2.50	1.85	1.10	.70	1.06	.76	.72	1.90	.73	.98	.43	.24	.37
10132	?	7.80	10.50	3.35	3.70	.42	.55	—	—	1.10	.73	1.03	.76	.73	1.67	.65	.92	.40	.24	.38
10014	?	8.75	11.25	3.85	4.10	.70	.75	2.20	1.95	1.12	.71	1.10	.77	.74	1.80	.78	1.00	.45	.27	.38
2273	?	8.50	11.25	3.40	3.65	.48	.65	—	1.85	1.15	.70	.98	.74	.70	1.75	.74	1.00	.45	.25	.37
234	?	8.40	10.80	3.60	3.68	.45	.62	2.10	1.87	1.10	.70	.96	.75	—	1.75	.75	1.00	.43	.24	.37
257	?	9.00	11.75	3.75	4.10	.58	.67	2.40	2.05	1.10	.72	1.03	.74	.70	1.73	.78	.96	.45	.24	.37
2229	?	8.65	10.75	3.65	3.90	.50	.62	2.35	2.00	1.07	.68	.93	.67	.64	1.70	.80	.95	.47	.24	.39
2734	?	8.00	10.50	3.25	3.35	—	—	—	—	1.15	.70	1.05	.70	.65	1.65	.70	.92	.43	—	—
10356	?	8.35	11.00	3.35	3.57	.53	.65	—	—	1.00	.70	1.00	.73	.70	1.68	.72	.95	.42	.23	.36
1754	?	8.80	11.45	3.62	3.87	.65	.70	2.40	1.75	1.08	.69	1.02	.73	.71	1.70	.73	.93	.45	.23	.36
2274	?	9.00	11.85	3.75	4.00	.67	.70	2.40	1.20	1.12	.70	1.00	.73	.70	1.76	.78	.97	.50	.25	.37
256	?	8.25	11.25	3.37	3.55	.55	.50	—	1.60	1.10	.68	.97	.72	.68	1.75	.80	.97	.45	.26	.35
1638	?	8.80	11.40	3.55	3.70	.47	.60	—	—	1.06	.70	1.00	.77	.73	1.75	.75	1.00	.46	.24	.36
2281	?	8.85	11.50	3.75	3.80	—	—	—	—	1.10	.71	1.04	.74	.71	1.70	.75	.98	.47	.24	.32
5857	?	8.75	10.75	3.42	3.78	.46	.52	—	1.82	1.18	.70	1.08	.75	.72	1.75	.76	1.00	.46	.23	.37
235	?	8.62	11.20	3.50	3.63	—	—	—	—	1.05	.72	1.07	.75	.70	1.75	.73	.96	.44	.23	.35
1481	?	8.80	11.25	3.63	3.95	.53	.45	2.50	2.00	1.13	.71	1.05	.75	.72	1.75	.80	.94	.46	.21	.33
254	?	8.70	10.90	3.40	3.67	.40	.53	—	—	1.10	.66	1.00	.70	.67	1.72	.67	.97	.48	.23	.37
2986	?	8.50	10.75	3.40	3.45	.55	.60	—	—	1.05	.66	1.02	.75	.70	1.62	.63	.93	.37	.22	.37

Table M. — TYRANNUS CAROLINENSIS Temm.

M. C. Z. Number.	Sex.	Length.	Alar Extent.	Wing.	Tail.	Extent of Primaries beyond Inner Secondaries.	Extent of Primaries beyond Outer Secondaries.	Extent of Tail beyond Upper Coverts.	Extent of Tail beyond Lower Coverts.	Tarsus.	Hind Toe.	Middle Toe.	Outer Toe.	Inner Toe.	Head.	Bill.				
																Culmen.	Commissure.	Gony.	Height.	Width.
1071	?	8.00	13.50	4.37	3.25	1.00	1.40	1.50	1.30	.75	.50	.80	.62	.56	1.67	.80	1.00	.44	.32	.53
10027	?	8.20	14.25	4.63	3.20	.87	1.42	1.70	1.30	.80	.55	.85	.61	.58	1.67	.80	1.02	.47	.30	.54
10026	?	7.85	14.20	4.67	3.50	1.05	1.53	1.80	1.46	.70	.55	.77	.60	.56	1.66	.77	.98	.47	.27	.45
10023	?	8.00	14.25	4.55	3.50	1.06	1.48	1.78	1.50	.70	.53	.87	.57	.53	1.58	.77	.92	.46	.28	.37
5546	?	8.15	14.15	4.63	3.43	1.09	1.56	1.62	1.36	.70	.60	.80	.60	.57	1.71	.83	1.02	.51	.27	.43
114	?	8.00	14.10	4.45	3.22	1.03	1.40	1.77	1.38	.80	.54	.77	.55	.50	1.73	.80	1.05	.50	.30	.50
112	?	8.20	13.50	4.47	3.33	.95	1.44	1.71	1.40	.70	.53	.77	.60	.56	1.60	.74	.98	.42	.30	.47
10025	?	7.00	13.00	4.17	3.15	.96	1.22	1.80	1.43	.67	.53	.73	.54	.50	1.58	.70	.97	.42	.28	.44
10376	?	7.75	13.75	4.42	3.25	.90	1.38	1.77	1.57	.70	.55	.77	.60	.53	1.68	.80	1.02	.50	.27	.47
10029	?	8.00	13.80	4.45	3.25	.90	1.30	1.73	1.73	.70	.72	.85	.62	.60	1.72	.78	1.05	.50	.27	.47
284	?	8.00	13.50	4.30	3.15	.85	1.15	1.70	1.47	.75	.67	.75	.60	.58	1.63	.67	1.00	.40	.28	.46
113	?	8.25	14.15	4.60	3.20	.90	1.45	1.55	1.20	.78	.58	.85	.70	.65	1.68	.80	1.05	.46	.27	.45
1317	?	7.40	12.50	4.25	2.93	.76	1.10	1.55	1.20	.73	.57	.78	.62	.55	1.57	.62	.90	.37	.20	.46
4003	?	8.20	14.15	4.60	3.30	1.35	1.62	1.78	1.42	.71	.57	.87	.65	.58	1.68	.82	1.04	.48	.26	.49
285	?	7.80	13.00	4.38	3.25	.83	1.15	1.77	1.45	.73	.54	.77	.64	.60	1.73	.82	1.05	.45	.29	.50
1318	?	7.85	13.25	4.20	3.25	.87	1.22	1.68	1.10	.75	.54	.78	.60	.53	1.63	.73	.95	.37	.27	.43
1758	?	8.10	13.80	4.50	3.30	.95	1.43	1.75	1.68	.70	.55	.77	.58	.55	1.68	.83	1.00	.50	.28	.46
1563	?	8.40	14.80	4.85	3.54	1.05	1.47	1.83	1.30	.76	.62	.83	.66	.60	1.70	.80	1.00	.45	.30	.45
283	?	8.65	14.25	4.58	3.47	.98	1.45	1.75	1.37	.70	.58	.82	.57	.63	1.75	.84	1.07	.48	.25	.48
4003	?	8.30	13.50	4.60	3.45	1.00	1.40	1.97	1.44	.73	.56	.77	.62	.60	1.63	.75	.90	.45	.28	.45

Table N. — PYRANGA RUBRA Vieillot.

M. C. Z. Number.	Sex.	Length.	Alar Extent.	Wing.	Tail.	Extent of Primaries beyond Inner Secondaries.	Extent of Primaries beyond Outer Secondaries.	Extent of Tail beyond Upper Coverts.	Extent of Tail beyond Lower Coverts.	Tarsus.	Hind Toe.	Middle Toe.	Outer Toe.	Inner Toe.	Head.	Bill.				
																Culmen.	Commissure.	Gonys.	Height.	Width.
—	♂	6.75	11.75	3.57	2.62	.80	1.13	—	—	.77	.52	.75	.53	.45	—	.58	.73	.44	.33	.38
10300	♂	6.90	11.30	3.75	2.75	.85	1.25	1.50	1.05	.82	.60	.70	.57	.50	1.53	.58	.75	.45	.35	.37
2929	♂	7.25	11.30	3.65	2.62	.75	1.12	1.23	.85	.73	.56	.67	.53	.47	1.48	.62	.75	.45	.33	.36
9795	♂	7.25	11.30	3.70	2.55	.92	1.12	1.41	1.13	.86	.60	.80	.57	.57	1.50	.64	.76	.40	.35	.40
10119	♂	7.25	11.60	3.80	2.73	.95	1.25	1.25	1.00	.68	.57	.80	.57	.53	1.48	.65	.73	.42	.35	.40
589	♂	7.30	11.75	4.00	2.73	1.07	1.30	1.35	1.00	.72	.57	.75	.58	.52	1.52	.63	.76	.47	.37	.40
10366	♂	7.30	11.75	3.85	2.85	1.00	—	—	—	.76	.58	.76	.53	.51	1.52	.63	.75	.45	.36	.39
10388	♂	7.20	11.60	3.87	2.75	.95	1.20	1.25	1.33	.77	.53	.79	.55	.50	—	—	—	—	—	—
10133	♂	7.00	11.00	3.77	2.77	.85	1.00	1.45	.88	.78	.58	.75	.55	.53	1.48	.57	.73	.40	.35	.36
1834	♂	6.75	11.30	3.60	2.50	.85	1.17	1.12	1.12	.70	.60	.73	.56	.54	1.46	.62	.73	.45	.37	.42
190	♂	6.75	10.75	3.80	2.64	.90	1.20	1.50	1.60	.67	.55	.70	.56	.56	1.38	.64	.70	.42	.33	.38
1833	♂	6.90	11.25	3.70	2.55	.85	1.05	1.33	1.05	.73	.55	.70	.57	.52	1.50	.65	.77	.43	.35	.40
1642	♂	7.00	10.65	3.85	2.75	1.07	1.23	1.50	1.10	.75	.53	.76	.55	.48	1.50	.65	.77	.45	.33	.37

Table O. — HEDYMELES LUDOVICIANUS Cab.

M. C. Z. Number.	Sex.	Length.	Alar Extent.	Wing	Tail.	Extent of Primaries beyond Inner Secondaries.	Extent of Primaries beyond Outer Secondaries.	Extent of Tail beyond Upper Coverts.	Extent of Tail beyond Lower Coverts.	Tarsus.	Hind Toe.	Middle Toe.	Outer Toe.	Inner Toe.	Head.	Bill.				
																Culmen.	Commissure.	Gonys.	Height.	Width.
1438	♂	7.50	12.25	3.87	3.05	.80	1.17	1.25	1.00	.92	.70	.75	.75	.70	1.50	.72	.78	.43	.54	.45
590	♂	7.75	12.30	4.00	3.00	.90	1.25	1.55	1.24	.93	.70	.74	.74	.72	1.55	.77	.80	.48	.55	.47
9787	♂	7.75	11.75	3.83	2.95	.85	1.13	1.50	1.40	.90	.70	.67	.67	.62	1.53	.73	.75	.45	.50	.43
2882	♂	8.30	12.50	4.15	3.05	.93	1.15	1.45	1.15	.88	.70	.75	.75	.74	1.55	.73	.77	.45	.51	.46
9935	♂	7.55	11.90	4.00	3.02	.60	1.06	1.28	.95	.87	.67	.75	.75	.70	1.55	.70	.78	.48	.56	.45
9580	♂	8.12	12.50	4.25	3.05	1.06	1.24	1.30	1.00	.88	.72	.65	.65	.55	1.58	.75	.88	.50	.60	.50
1518	♂	7.75	11.80	3.85	2.75	.70	1.00	1.20	1.00	.80	.70	.65	.65	.60	1.54	.75	.82	.40	.55	.45
1065	♂	7.60	12.00	3.83	2.88	.82	.97	1.24	.98	.90	.60	.65	.65	.58	1.55	.70	.81	.45	.55	.43
1547	♂	7.50	12.10	3.86	2.70	.95	1.10	1.36	—	.82	.67	.63	.63	.56	1.50	.65	.75	.44	.51	.45
187	♂	7.75	11.50	3.85	2.77	.80	1.07	1.40	1.42	.82	.65	.67	.67	.63	1.52	.70	.77	.45	.53	.45
9581	♂	7.85	12.50	3.95	2.82	.77	1.00	—	1.12	.84	.67	.68	.68	.61	1.42	.66	.70	.43	.53	.42
5591	♂	7.75	11.90	4.00	3.08	.82	1.00	—	1.18	.87	.67	.68	.68	.58	1.50	.73	.82	.45	.54	.44
5592	♂	8.10	12.90	4.15	3.00	.75	1.15	1.70	1.35	.92	.65	.72	.72	.65	1.55	.70	.77	.45	.54	.43
2942	♂	7.85	12.15	3.97	2.93	.80	1.08	1.92	1.10	.90	.68	.65	.65	.60	1.50	.68	.72	.45	.55	.50
9934	♂	8.10	12.75	4.00	3.00	.83	1.05	1.40	1.22	.83	.70	.67	.67	.57	1.58	.75	.77	.46	.52	.47
1823	♂	7.15	12.00	4.05	2.85	.90	1.07	1.25	1.00	.70	.62	.78	.78	.74	—	—	—	—	—	—
10107	♂	7.65	11.75	4.20	2.93	.90	1.05	1.05	1.10	.86	.70	.73	.73	.60	1.57	.75	.73	.45	.50	.48

Table P. — GEOTILYPIS TRICHAS Cab.

M. C. Z. Number.	Sex.	Length.	Alar Extent.	Wing.	Tail.	Extent of Primaries beyond Inner Secondaries.	Extent of Primaries beyond Outer Secondaries.	Extent of Tail beyond Upper Coverts.	Extent of Tail beyond Lower Coverts.	Tarsus	Hind Toe.	Middle Toe.	Outer Toe.	Inner Toe.	Head.	Bill.				
																Culmen.	Commissure.	Gonys.	Height.	Width.
9976	♂	5.00	7.00	2.10	2.00	.35	.42	.92	—	.82	.55	.68	.52	.51	1.20	.45	.57	.31	.17	.22
358	♂	5.17	7.25	2.20	2.07	—	—	—	—	.80	.53	.75	.54	.53	1.20	.47	.53	.30	.17	.27
1808	♂	5.25	7.25	2.25	2.05	.36	.32	1.04	.75	.77	.53	.68	.50	.48	1.17	.48	.59	.30	.17	.27
2534	♂	5.15	7.20	2.12	1.88	.35	.35	—	.64	.74	.55	.75	.55	.52	1.25	.55	.67	.30	.18	.26
9775	♂	4.65	6.75	2.37	2.05	.25	.33	—	—	.78	.52	.73	.54	.53	1.13	.43	.55	.27	.15	.24
1809	♂	5.25	6.75	2.20	2.00	.26	.33	.87	1.00	.77	.52	.72	.51	.51	1.28	.48	.58	.31	.16	.27
2997	♂	5.25	7.10	2.18	1.95	.22	.35	—	.70	.77	.53	.73	.55	.53	1.22	.45	.57	.31	.17	.26
9858	♂	4.88	6.45	2.20	2.10	—	—	1.06	.82	.77	.52	.72	.55	.53	1.22	.47	.57	.34	.18	.27
7515	♂	5.37	7.25	2.27	2.10	—	—	—	—	.74	.50	.72	.53	.51	1.18	.50	.60	.32	.18	.22
330	♂	5.10	6.75	2.13	2.10	.33	.37	—	—	.72	.47	.68	.48	.48	1.22	.50	.60	.32	.18	.27
1444	♂	5.63	7.50	2.32	2.17	.35	.40	—	—	.77	.52	.75	.55	.53	1.26	.52	.65	.32	.17	.28
1971	♂	4.60	6.30	1.95	1.78	—	—	—	—	.75	.51	.62	.46	.44	1.16	.45	.54	.28	.15	—
2996	♂	5.37	7.30	2.27	2.08	.32	.38	.97	.70	.78	.54	.73	.53	.52	1.22	.48	.60	.32	.18	.25
2361	♂	4.87	6.75	2.12	1.82	—	—	—	—	.82	.44	.70	.52	.52	1.18	.43	.60	.31	.13	.26
2427	♂	4.88	6.75	2.03	1.88	—	—	—	—	.79	.50	.66	.47	.45	1.15	.43	.58	.30	.15	.28
9669	♂	5.05	6.75	2.13	2.05	—	—	—	—	.80	.48	.73	.53	.51	1.22	.48	.57	.32	.15	.22
9836	♂	5.00	6.85	2.05	1.85	—	—	—	—	.76	.51	.66	.52	.50	1.15	.48	.57	.30	.14	.21
2398	♂	5.20	7.00	2.25	2.00	—	—	—	—	.76	.50	.70	.53	.51	1.20	.49	.59	.32	.15	.23
1445	♂	5.33	6.90	2.15	2.05	—	—	—	—	.78	.48	.73	.55	.54	1.23	.48	.57	.32	.16	.24
2436	♂	5.10	6.80	2.10	1.95	—	—	—	—	.74	.52	.67	.52	.50	1.15	.44	.56	.27	.15	.24

Table Q. — HARPORHYNCHUS RUFUS *Cab.*

M. C. Z. Number.	Sex.	Length.	Alar Extent.	Wing.	Tail.	Extent of Primaries beyond Inner Secondaries.	Extent of Primaries beyond Outer Secondaries.	Extent of Tail beyond Upper Coverts.	Extent of Tail beyond Lower Coverts.	Tarsus.	Hind Toe.	Middle Toe.	Outer Toe.	Inner Toe.	Head.	Bill.				
																Culmen.	Commissure.	Gonys.	Height.	Width.
1752	?	11.85	14.00	4.25	5.05	.50	.65	—	3.20	1.42	.93	1.25	.85	.80	2.30	1.13	1.25	.70	.35	.45
1625	?	10.55	12.75	3.90	4.60	.43	.60	3.15	2.95	1.27	.90	1.25	.84	.77	2.15	1.00	1.24	.60	.28	.41
237	?	11.00	12.55	3.80	4.78	.38	.56	3.13	2.75	1.33	.84	1.17	.86	.80	2.30	1.06	1.33	.76	.27	.38
10320	?	11.25	13.30	3.95	4.95	.43	.68	3.35	2.95	1.42	.87	1.22	.78	.72	2.25	1.05	1.34	.72	.29	.43
1670	♂	11.50	13.25	4.15	5.18	.47	.60	3.60	3.20	1.36	.93	1.25	.85	.80	2.18	—	1.28	.65	.27	.44
2585	?	10.85	12.90	3.95	4.86	.50	.65	—	2.65	1.25	.87	1.17	.80	.70	2.26	1.03	1.35	.68	.28	.40
1439	♂	11.10	12.80	4.00	4.93	.36	.56	3.17	2.95	1.35	.92	1.28	.90	.85	2.25	1.12	1.33	.67	.27	.43
1067	?	10.85	13.35	4.15	5.00	.32	.55	3.38	3.05	1.32	.86	1.25	.90	.78	2.30	1.06	1.35	.73	.30	.45
8887	♂	11.21	13.50	4.20	4.50	—	—	—	—	1.25	.79	1.25	.75	.73	2.25	1.05	1.35	.74	.30	—
8886	♂	11.41	13.00	4.07	4.59	—	—	—	—	1.30	.85	1.23	.82	.75	2.15	1.00	1.23	.67	.27	.40
8888	♀	11.31	13.30	4.16	5.00	—	—	—	—	1.30	.82	1.23	.85	.82	2.23	1.00	1.25	.65	—	—
8889	♀	11.30	13.00	5.00	5.10	—	—	—	—	1.20	.78	1.25	.90	.84	—	1.00	1.28	.72	.27	.40
8885	♂	11.35	13.35	4.11	5.00	—	—	—	—	1.22	.85	1.30	.87	.78	2.22	1.04	1.33	.70	.27	.37
8891	♂	11.60	13.55	4.25	5.00	—	—	—	—	1.27	.85	1.24	.87	.77	—	1.00	1.26	.67	.25	.38
8996	♂	11.50	13.25	4.10	5.00	—	—	—	—	1.35	.87	1.25	.86	.78	2.23	1.05	1.32	.72	.27	.38
8897	♂	11.75	13.35	4.32	5.30	—	—	—	—	1.24	.85	1.30	.83	.79	2.22	1.05	1.28	.73	.28	.36
8896	♂	11.60	13.00	4.15	5.20	—	—	—	—	1.35	.85	1.27	.83	.75	—	.94	1.30	.68	.25	.40

Individual Variation in the Size and Form of the Bill. — That considerable variation occurs in the size and shape of the bill, in specimens of the same sex and species living together at the same locality, is evident from a glance at some of the preceding tables of measurements. The variation in this organ is further illustrated in the accompanying plates (Plates IV – VIII), in which are given figures of the bills of several specimens of each of a number of species. Much greater differences are here shown to exist in conspecific specimens of the same sex and from the same locality than occur between those supposed to be distinct, of which comparative figures of the bills have been published with a view of demonstrating their specific diversity. In only a few groups in fact, and mainly in the long-billed *Grallæ*, is the bill generally admitted to be too variable to afford an important basis for the discrimination of species.

The principal points of variation in the form of the bill consist in variations in its general size, without corresponding variations in the general size of the individual, and in the details of its form in regard to thickness and length. There are also other variations in respect to the emargination or dentation of the terminal portion, especially in the vast group of the insectivorous species, and in the “festooning” of the bill in many of the hawks.* In respect to the size of the bill, it is a noteworthy fact that birds specifically and sexually identical vary in such a way that specimens much below the average size possess bills above the average size for their respective species, and, conversely, that specimens above the average size have bills much smaller than the average for their respective species, the general proportions of the bill in each case being essentially the same. In such cases, with the increase or decrease in length, there are corresponding differences in the thickness of the bill, both in the vertical and transverse directions. In other cases with the increase in length there is no corresponding increase in thickness, such a differentiation thus resulting in a relatively attenuated form of the bill. In other cases the bill is shortened without a corresponding decrease in its thickness, from which results a short, thick, or robust bill. The variation in thickness is again sometimes relatively greater in the vertical

* In respect to this point, see Dr. Henry Bryant's paper on “Variations in the Plumage in *Buteo borealis* auct. and *B. Harlani* Aud.?” (Proc. Boston Soc. Nat. Hist., Vol. VIII, p. 107 *et seq.*, 1861, where the variation in this feature is especially noticed.

than in the transverse direction, and sometimes the reverse, thus giving in some cases a deep, narrow bill, and in others a broad, depressed bill. In the latter case the differences are especially important, as will be more fully shown later. In regard to the tooth-like indentation near the tip of the bill in so many of the insectivorous birds, it is found that in some species which usually have it strongly developed, specimens occasionally occur with the indentation nearly or quite obsolete. Again in other cases where this feature is usually but slightly developed, some specimens have the notch at the tip of the bill exceedingly prominent. Similar variations occur in regard to the development of the so-called "festoon" of the upper mandible in the hawks, as Dr. Bryant has already sufficiently shown.

The greatest range of individual differentiation in any given organ occurs, as would be naturally expected, in those species which have that organ more than ordinarily developed, and also in species of a low grade of structure. In the long-billed *Grallæ* both these conditions exist, and it is in such genera as *Numenius*, *Gambetta*, *Limosa*, *Scolopax*, *Philohela*, and *Gallinago*, that the maximum of bill variation is seen. It is less marked in the song-birds, though in many members of this group the variation is by no means small. In the typical woodpeckers, on the other hand, which have the bill especially adapted to a peculiar function, that of digging into wood, the variation is scarcely appreciable, since any considerable variation from its usual form would seriously impair its efficiency. In the semi-frugivorous and terrestrial *Picidæ*, however, we again meet with the usual range of variation.

In the accompanying plates illustrative of variation in the bill, representatives from the higher types of the *Oscines* have mainly been chosen, several representatives from widely different families having been selected. Plate IV, figures 1 and 1*a*, 2 and 2*a*, give a view of the bills of two specimens of the common king-bird (*Tyrannus carolinensis*), from Eastern Massachusetts, which differ from each other as much as the bills of different genera sometimes do. One of them, as will be seen, is so much narrower and deeper than the other as to give very different proportions and outlines. The skulls of these two specimens vary in the same manner as do the bills, the one having a broad, flat skull, and the other a narrow, high one. Two specimens of *Myiarchus crinitus*, one of which is from South Carolina and the other from Western New York, differ as much from each other, and in nearly the

same way, as do those of the king-bird. Similar and nearly as great variations occur also between different specimens of *Contopus borealis*, *C. virens*, *Empidonax minimus*, *E. flaviventris*, *Sayornis fuscus*, and in several species of the South American *Tyrannidæ*. But between these two extremes are found in other specimens nearly every possible degree of gradation.

Figures 3 and 3a to figures 7 and 7a (same plate) represent different forms of the bill in *Troglodytes aëdon*. Between these specimens there are great differences both in respect to absolute size and to general form, greater than would be deemed necessary by most ornithologists for the differentiation of species. These examples are all from Florida, and essentially from the same locality. Other specimens in the Museum come between these extremes in such a way as to show the inconstancy of all these forms. The variation in color, which is considerable in this species, does not accord with the variation in the bill, specimens exhibiting the extremes of color as often having the bills alike as otherwise, and, conversely, those with bills alike differ widely in color.

Figures 8 and 8a to 11 and 11a (same plate) indicate the variability of the bill, especially in respect to length, in Massachusetts specimens of *Seiurus noveboracensis*. The first corresponds essentially with, and unquestionably is, an example of the so-called *Seiurus ludovicianus*, which, in all probability, is but the darker colored, longer-billed southern form of *S. noveboracensis*. This species varies also remarkably in color, but the variation in color, as in the case of *Troglodytes aëdon*, and as is commonly the case in other species, does not accord with the variation in the bill, some of the long-billed specimens being in color almost undistinguishable from some of the short-billed ones, while some of those with medium bills present the extreme degrees of variation in respect to color.

Figures 12 and 12a to 14 and 14a (same plate) represent the bills of three male specimens of *Mniotilta varia* from the vicinity of Cambridge, which present as great differences as modern ornithologists would ordinarily deem sufficient, if the specimens had come from Mexico instead of from Massachusetts, to warrant their recognition as types of three distinct species. The correspondingly great variations in color in this species have already been adverted to (p. 190). The bill, however, in specimens presenting extreme forms of color variation, unfortunately for ultra-divisionists, may be either of the ordinary form

or of either of the forms figured, or of any intermediate form, as exemplified by the specimens of this species in the collection of the Museum. Figure 15 and 15*a* and 16 and 16*a* (same plate) are accurate representations of the bills of two Massachusetts males of *Dendræca striata*. The differences between these specimens, though so great, are not greater than occur in different conspecific examples of several other species of this genus contained in the Museum.

Massachusetts specimens of *Certhia familiaris* differ even more in the form of the bill than do the specimens above figured of either *Troglodytes ædon* or *Mniotilta varia*. They also present a similar range of color variation in the plumage, and one equally at variance with the variation in the bill.

Figures 19 and 19*a*, 20 and 20*a* (same plate), show how widely two Florida specimens (both males) of *Pyranga æstiva* vary in respect to the size of the bill, the specimens in question differing but little in general size. If these figures are compared with the figures recently published of the bills of certain supposed species of *Pyranga*,* they will be found to vary more than some of the latter do, and indicate how unsatisfactory the nature of species must be when based mainly upon differences in the bill. Other conspecific specimens of *Pyranga* in the Museum exhibit great difference in the size, form, and position of the tooth-like processes of the upper mandible, and in the color of the bill, — differences that have been regarded as specific characters. The color of the bill in many species of birds, in fact, varies greatly in specimens of the same species taken at the same season, and generally in those taken at different seasons; yet it is a character that has been relied upon for the distinction of species.

Figures 1 and 1*a*, 2 and 2*a*, 4 and 4*a*, and 5 and 5*a*, Plate V, illustrate variations in the bill in Massachusetts representatives of *Ægiolais linarius*. Figures 3 and 3*a*, and 5 and 5*a*, are drawn from specimens from Arctic America, the first being an original specimen of the *Æ. fuscescens* Coues ex auct., and the other a similar specimen of the *Æ. exilipes* Coues. Figures 7 and 7*a* to 10 and 10*a*, inclusive (same plate), represent variations of the bill in male specimens of *Chrysomitris tristis*, a species allied to *Æ. linarius*. It will be seen that the two series are nearly parallel in respect to the amount and character of the variations in the bill. Figures 11 and 11*a* and 12 and 12*a* indicate similar variations in an-

* Proceed. Phil. Acad. Nat. Sci., June, 1869, pp. 130-133.

other allied species, the *Chrysomitris pinus*, and figures 13 and 13a to 15 and 15a, inclusive (same plate) similar variations in another species (*Curvirostra americana*), of the same sub-family. In the latter case the specimens are also all males, and all from the vicinity of Cambridge, they having been killed in fact from the same flock. In the *Ægiothus* group numerous so-called "species" have been described by different writers, six or seven of which were recognized by Dr. Coues a few years since in his monograph of that genus.* A considerable number of these species have been generally looked upon as equivocal, and the exact number in the group and their distinctive characteristics have been a matter of much uncertainty. Recently the writer above referred to has again revised the group,† and arrives at the conclusion that if more than one species exists, all the forms previously recognized by him as species are valid species. I can readily grant this alternative, being fully convinced that the genus consists of but a single known species, which has a circumpolar distribution. The alleged specific distinctions have consisted in differences in general size, in the relative size of the bill, the length of the tarsus, wing, and tail, and in color. Some of these differences are doubtless climatic and local, while others may be due to age, but the greater part I believe to be to a great degree purely individual, inasmuch as they are paralleled in allied species, whose standing has not been and cannot reasonably be questioned. But the special consideration of the variations presented by the *Ægiothi* and similar groups will be reserved till after the facts relating to geographical variation have been presented, since they can then be more appropriately discussed.

Figures 16 and 16a to 18 and 18a, inclusive (Plate V), represent the bills of three male specimens of *Passerculus savanna*, from different localities on the Atlantic coast. The specimen represented in figures 18 and 18a, has the bill of minimum size, being in bulk less than half that of the one represented in figures 17 and 17a.‡ Figure 17, it will be observed, corresponds nearly with the so-called *P. sandwichensis* § of

* A Monograph of the genus *Ægiothus*, etc., Proc. Phil. Acad. Nat. Sci., Vol. XII, p. 373, 1861, Vol. XV, p. 40, 1863.

† On variations in the plumage of the *Ægiothi*, Ibid., Vol. XXI, p. —, 1869.

‡ Other specimens received from Grinnell, Iowa, from Professor H. W. Parker, since the above was written, have bills still smaller than any of those here figured.

§ Baird's Birds of N. Amer., p. 444, 1858.

the Pacific coast, and figure 17 with the so-called *Passerculus alaudinus*,* also of the Pacific coast.

Plate VI, although designed more especially to illustrate local variation, indicates to some extent the individual variation existing in *Agelaius phoeniceus*. Figures 1 and 1*a* represent the average type of the bill in this species in Massachusetts, and figures 3 and 3*a*, and 4 and 4*a*, unusually long and unusually short forms of the bill found at the same locality. Figures 2 and 2*a*, 5 and 5*a*, and 6 and 6*a*, represent a similar series from the St. John's River, Florida. All the specimens of the two series are adult males.

Plate VII represents similar variations of the bill in *Quiscalus purpureus*. Figures 1 and 1*a*, 3 and 3*a*, 4 and 4*a*, and 6 and 6*a*, represent the average and the extreme types of the bill met with in Massachusetts males. The latter also represents an inflexed type of bill, a modification seen in many species, it being especially common in the *Quiscali* and other genera having the bill of a similar form. It is unmistakably an individual peculiarity, evidently depending mainly upon age, and resulting from the upper mandible outgrowing and overhanging the lower. In *Quiscalus purpureus* such specimens are more or less frequent at probably all localities, they having been received at the Museum from Maine, Massachusetts, New Jersey, Florida, and Illinois, and I have seen them from the West Indies. It often gives rise to the name *inflexirostris*, which is found so frequently a synonyme.† The figures of the bills of four females of *Sturnella ludoviciana* (Plate VIII), from Florida, indicate the character of the bill variation exhibited by different individuals of this species at the same locality, independently of any variation attributable to sex. Figures 5 and 5*a*, and 6 and 6*a* (same plate) show that like variations occur in *Colaptes auratus*, the figures being drawn from two Massachusetts females.

Similar comparisons, with similar results, might be made with scores of other species, but the above illustrations will doubtless suffice to show that individual variation in the form of the bill is not only great, but that it exists in groups having a high grade of structure. Other groups might have been chosen in which the individual variation in the form of the bill, as already stated, is far greater than in the instances above

* Bonaparte, Comptes Rendus, Vol. XXXVII, p. 918, 1853.

† Concerning *Quiscalus inflexirostris* Swainson, see below (Part IV), under *Q. purpureus*.

cited. The *Grallæ* have already been referred to as presenting remarkable examples of bill variation. In some of the *Anatidæ*, however, it is scarcely less, whilst it is especially great among many of the *Longipennes*. Hence some authors evidently attach too high importance to the exact form of the bill in these groups.

All the illustrations referred to above have been drawn, with one or two exceptions, from fully adult specimens. One of these is a specimen of *Passerculus savanna* (Plate V, fig. 18), which is a bird of the year, killed in Labrador in August, before it had quite completed its first moult. Another is the smallest billed specimen of *Chrysomitris tristis* (Plate V, fig. 10), which is also evidently a bird of the year. The other is an autumnal specimen of *Dendræca striata* (Plate IV, fig. 15). They all, however, would be ordinarily considered as adult in size.

VARIATIONS IN THE SIZE AND FORM OF THE BILL, WING, ETC., RESULTING FROM AGE.

In the foregoing remarks on the variations in general size, in proportions, and in the form of different parts, exclusive reference has been had to adult specimens. It is easy, however, to confound difference depending upon age with those strictly resulting from individual differentiation. The form of the bill is especially subject to variation by age in specimens that upon casual inspection would seem to be full-grown. In long-billed birds the bill increases in length for several months after the bird is full-fledged, and even after it has once moulted. In short- and thick-billed birds, the bill increases considerably in thickness as well as in length after the individual seems to have acquired its adult size and proportions. As a general rule, then, "birds of the year" possess a relatively shorter and thicker bill than those fully adult, or three or four years of age. In old age an abnormal elongation of the upper mandible occasionally occurs, especially in species in which the tip of the upper mandible is decurved and projects slightly beyond the lower, as in *Corvus*, *Quiscalus*, *Vireo*, *Turdus*, *Larus*, etc. Since, however, great differences occur in the form of the bill in specimens of the same age, in birds of the year as well as in those unquestionably adult, it is sometimes difficult to determine how much of the difference in certain cases is to be considered as due to age and how much to individual variation.

The wing also varies considerably in form with age. In many of the song-birds, at least, and also in the raptorial birds, the wing becomes more pointed with the second and third moultings of the remiges. Birds of the first year hence have, even after the flight feathers are fully grown, a shorter and more rounded fore-wing, as a general rule, than birds of two or three years of age. These differences of course result from variations in the relative length of the primaries, the outer primaries being the last to acquire their ultimate proportions, as they are also the last primaries to be renewed in the annual moult. A similar change with age occurs in the form of the inner point of the wing, or that formed by the inner secondaries. These, like the primaries, are subject to a gradual increase in length for a time with each moult, they likewise being the latest of the secondaries to acquire their mature size, as they are also the last of the secondaries changed in each normal moult. Thus, through the gradual elongation of the outer primaries and the inner secondaries, a slight change is produced in the general form of the wing. It is, however, only slight, and since some young birds have as pointed wings as any of the same species which are fully adult, and some adult birds have wings as much rounded as the full-grown young, the rule is subject to many exceptions. The sexes of the same species also often differ similarly with the young and old in respect to the form of the wing. This is more especially the case in those species in which the female is much smaller and much duller colored than the male, the structural inferiority of the female to the male being thus evident in various features.

While the wing may be regarded, as already stated, as generally smaller and more rounded in the younger individuals, it not unfrequently happens that the specimens having the greatest alar extent are immature birds. This has been particularly noticed in the eagles and hawks, as well as in some of the gulls, in which it is so frequent as to have attracted the attention of numerous observers.* The feathers of the wings and tail are not only longer, but they are also broader, and hence in the expanded wing present a greater resisting surface to the air. Two explanations of this fact present themselves. First, in the cases referred to, the birds may have been born at a very northern locality, whence only the younger birds ever descend so far south. Second, the greater lack of power in the muscles of flight in the young birds, as

* See American Naturalist, Vol. III, 1869, p. 517.

compared with those fully mature, may be counterbalanced by a relatively larger supporting surface in the wings and tail. Whatever the explanation may be, the facts seem to be unquestionably as above stated.

Other variations in the plumage and in other characters depending upon age, but which are liable to be confounded with individual differentiation, might be cited, but none seem to be of sufficient importance to require a special description.

GENERAL REMARKS ON INDIVIDUAL VARIATION.

After the preceding remarks on this subject, I should perhaps state expressly what I regard to be the bearing of the facts above discussed, otherwise I might be understood as in a great measure discarding the majority of the characters used in the diagnoses of species and genera. Nothing, however, is further from my purpose. What I urge is simply this: that the extent of purely individual variation is far greater than has usually been recognized, and that as a result numerous strictly nominal species have found their place in our systems, from naturalists having mistaken these differences for true specific characters. Individual variation, however, is so complicated with geographical variation, that the general bearings of the whole subject will be deferred till the end of the discussion of the latter topic.

As regards the general cause of individual differences in animals, it is too evidently constitutional to allow of any other hypothesis, and akin to that seen in domestic animals, and which in man gives to each individual his unlikeness in temperament and physical structure to all other men. While individuality is so patent and so universal in the human species, and scarcely less so in domesticated animals, it is one of the most surprising facts in zoölogy that so many naturalists should have entertained the idea that there is an almost total absence of it in feral animals, and that the description of a single specimen will suffice for that of its species. Practically, however, this has been the fact, and eminently so with that large class of "species hunters," who have not inaptly been characterized as "closet naturalists"; for to this class and not to the field naturalists are we mainly indebted for the long lists of synonymes that form so vexatious a burden to zoölogical science.

Certain secondary causes that share in producing individual variation

are doubtless more or less obscurely traceable. Among these are certain circumstances attending the time of hatching, as well as, of course, the vigor of the parent. Not unfrequently the first attempts of birds to rear their brood are unsuccessful, from their eggs or young being destroyed by their enemies. Persisting, however, in their efforts, it is late in the season before their brood is fledged, several sets of eggs or young having been previously destroyed. The birds of such broods are found to be smaller and paler colored than those hatched earlier in the season. In cases where several broods are reared each year, as a general rule the birds of the earlier brood seem in all respects the most perfect and vigorous. Various other causes operating during their infancy doubtless more or less affect their general size, their proportions, and colors when mature. Food has doubtless much to do with variation in color, though but few facts bearing upon this point have been yet recorded. Professor Agassiz informs me, however, that many years since, in Switzerland, he raised many *Pyrrhula vulgaris*, and found that by feeding them on the seeds of hemp the red on the breast changed to black. The well-known fact that certain brightly colored birds, as the purple finch (*Carpodacus purpureus*) and the crossbills (*Curvirostra*), change, when kept in cages, from bright red to dull olive with their first moult, and never again, or at least so long as kept in confinement, regain their original color, shows how susceptible the color of birds is to the influences of food and artificial conditions of life.

CLIMATIC VARIATION.

Climatic variation involves as completely all parts of the animal as does individual variation. It is more marked, however, in some features than in others. The three most prominent phases of climatic variation in birds are the following: variation in general size, variation in the size and form of the bill, variation in color.

Climatic Variation in Size. — Variation in the size of individuals of the same species with differences in the latitude and altitude of their respective places of birth is a fact already so well known as to be quite generally recognized; hence any demonstration of such a variation is in the present connection unnecessary. A few tables of comparative measurements of New England and Florida specimens given in Part IV serve to illustrate its general character and extent. Similiar illustrations are abundantly afforded by the tables of measurements published in Pro-

fessor Baird's Birds of North America,* in the text of which work frequent reference is made to the differences in size between northern and southern specimens of the same species. The same author also subsequently called attention to the subject, and explicitly announced a general law of geographical variation in size; namely, a gradual decrease in size in individuals of the same species with the decrease in the latitude and altitude of their birth-places.†

In some species, and throughout some entire families, climatic variation is more marked than in others; generally, however, it is very appreciable, and amounts, in respect to size, not unfrequently to from twelve to twenty per cent ‡ of the average dimensions of the species.

Climatic Variation in the Bill. — The climatic variation in the size of the bill is, in general, inverse to that of the general size of the individual. In some species, as in the *Sittæ* and the typical members of the *Picidæ*, I have as yet been unable to trace an independent variation in the size of the bill to that of the body; but in many species there is not only a marked *relative* increase in the size of the bill to the southward, but, in some, an *absolute* increase, especially in its length.

* Pacific Railroad Explorations and Surveys, Vol. IX, Birds. By Professor S. F. Baird, with the co-operation of Mr. John Cassin and Mr. George N. Lawrence. 1858. Subsequently republished under the title of "The Birds of North America," with an Atlas of one hundred plates.

† Proc. Phila. Acad. Nat. Sci., Vol. XI, p. 300, November, 1859. Also in Am. Journ. Sci. and Arts, 2d Ser., Vol. XLI, p. 190, March, 1866.

‡ Variation in size with differences in habitat is by no means confined to birds. In mammals it is well known to be as great, if not greater, than among birds. In some wide-ranging species of mammals there appears to be a double decadence in size, — a diminution to the northward, in those non-migratory species whose habitats extend into the arctic regions, as well as a diminution to the southwards of the point where in general the maximum of size is attained, — as I have elsewhere had occasion to remark. (Bull. Mus. Comp. Zool., Vol. I, p. 199.) But in these exceptional cases of a decline in size to the northward, the cause of such a decline must result from climatic conditions the reverse of those producing the decline at the southward, — from the excessive rigor of the arctic climate instead of from the enervating influence of warm temperate and sub-tropical latitudes.

In the case of reptiles, the larger representatives of a given species are generally found at the North, as has also been observed to be the case with the edible marine and fluviatile fishes. (I am credibly informed that this is markedly the case with the codfish and the halibut.) In some groups of crustacea and mollusca the same fact has been repeatedly observed; but in insects, as in plants, the increase in size is generally to the southward, as is especially noticeable in the diurnal Lepidoptera. In plants, however, the increase is a purely vegetative one, the northern representatives of a given species being generally far the most prolific, in proportion to the size of the plant, near the northern limit of their respective habitats.

An increase in the length of the bill is most frequent in long-billed species, while in short-billed ones the increase is in general size, without material change in its proportions. With the increased length and slenderness of the bill there is in many cases also a tendency to greater curvature.

An increase in the length of the bill is quite marked in the genera *Quiscalus*, *Agelæus*, *Geothlypis*, *Troglodytes*, *Seiurus*, *Harporhynchus*, *Galeoscoptes*, etc. *Quiscalus purpureus* and *Agelæus phæniceus* afford good illustrations of geographical variation in the size and shape of the bill. Notwithstanding that the northern specimens are the larger, the southern ones have, in the average, bills as long, though slenderer, than the northern, and occasionally even longer. These differences are shown to some extent in Plates VI and VII, where the figures of the bills of Massachusetts and Florida specimens of these species are given side by side. In Plate VI, figures 1 and 1*a* represent the bill of an average Massachusetts male *A. phæniceus*, and figures 2 and 2*a* the bill of an average Florida male of the same species. The latter, while much less thick, is fully as long as the former. Figures 4 and 4*a* represent the shortest bill of a considerable series of Massachusetts specimens, and figures 6 and 6*a* the shortest or thickest bill of a similar series of Florida specimens. Figures 3 and 3*a* give the longest bill of the Massachusetts series, and figures 5 and 5*a* the longest of the Florida series, the specimens being in each case adult males. Plate VII, figures 3 and 3*a* represent the bill in average Massachusetts males of *Quiscalus purpureus*, and figures 2 and 2*a* that of average Florida specimens, while figures 1 and 1*a*, and 4 and 4*a*, show respectively the longest and the shortest bills of a considerable series of Massachusetts specimens. Figures 5 and 5*a* are from a New Jersey specimen, and figures 6 and 6*a* from a Florida specimen, the latter showing an inflection of the upper mandible more or less frequent in the various species of *Quiscalus*. The figures, as in the previous plate, were all drawn from adult males. In each of these species the average difference in the bills of Florida and Massachusetts birds is as great as is frequently considered to be sufficient to constitute specific differentiation, and between the extremes, especially of *A. phæniceus*, even subgeneric. Yet specimens from intermediate localities present such a gradual and complete transition between the two forms as to render their specific identity unquestionable.

A similar difference between Massachusetts and Florida examples,

with a gradual transition from the one to the other, through specimens from intermediate localities, is seen in *Troglodytes ædon*, *Geothlypis trichas*, and *Seiurus noveboracensis*. In *Pipilo erythrophthalmus*, *Ortyx virginianus*, *Corvus americanus*, and *Cyanura cristata* the bill is appreciably larger in the Florida than in the northern form. In *Corvus americanus* this difference was long since noticed by Professor Baird, the larger bill of South Florida specimens having led him to recognize a variety *floridanus* of this species, based chiefly on this difference.* The same author has also referred to the larger size of the bill in Florida specimens of *Ortyx virginianus*.†

In some species individual variation is so great that it is unsafe to draw conclusions respecting geographical variation from the examination of a small number of specimens. This is notably the case in *Sturnella ludoviciana*, in which the bill varies greatly in size and form, as does the bird in general bulk, at all localities. In the average, however, Florida specimens of this species seem to have a relatively longer and slenderer bill than those from the Northern States.

As already noticed, variation in the bill is not equally marked in all species, but it occurs in too many to admit of the supposition that the numerous cases wherein it is clearly marked are exceptional, or that it does not follow a general law of geographical variation. The observations above detailed are based on specimens collected on the Atlantic coast, from New England southward to Florida, and refer exclusively to species breeding within that range. But specimens of species which breed entirely to the northward of this range, collected during their semi-annual migrations, corroborate the law already stated, namely, an increase in the size of the bill to the southward in specimens of the same species from different breeding stations. In the *Anatidæ* and *Tringæ*, which breed far to the northward and pass the winter in lower latitudes, it is noticeable that, while those which arrive first in the fall, and those which return north latest in the spring, are smaller than those that arrive later and depart earlier, they have, nevertheless, relatively larger bills. This has been especially noticed in species of *Fulix*, *Bernicla*, *Actodromas*, and *Macrorhamphus*. Professor Baird has also referred to the larger size of the bill of the southern representatives of *Lagopus albus* as compared with those from further north,

* Birds of North America, p. 568, 1858.

† Am. Journ. Sci. and Arts, 2d Ser., Vol. XLI, p. 191, 1866.

"those from Eastern Labrador and Newfoundland," he says, appearing "to have decidedly broader, stouter, and more convex bills than those from the Hudson's Bay and more northern countries."* In the writings of various authors on the birds of Southern Mexico, Central America, Southern Asia, and Northern Africa, frequent mention is incidentally made of the larger size of the bills of southern representatives of northward ranging species. Although such statements record what have been apparently regarded as only isolated facts, their frequency indicates that the increase in the size of the bill to the southward is not confined to the birds of Eastern North America, nor exclusively to those of temperate and sub-tropical countries, but that it is a general geographical law, similar to that of the variation with locality in the general bulk of the individual.

Geographical Variation in Color. — Geographical variation in color in birds may be regarded as of two kinds, which may be termed, from their different geographical relations, latitudinal variation and longitudinal variation. The first is coincident with differences in latitude, and the second with differences in longitude. Both are due, however, to climatic peculiarities, and are hence, strictly speaking, climatic. The latitudinal is perhaps at present the best known, and will be first considered.

(a) *Latitudinal Variation.* — In those species of North American birds whose breeding range extends over a wide range of latitude, the southern-born specimens are, as a general rule, appreciably darker or brighter, or more intensely colored, than northern-born ones of the same species; in many instances the difference being so great as to impress even the casual observer. Dark colored birds, like the *Quiscalis*, *Agelaius phœniceus*, etc., become blacker towards the southern limit of their respective habitats, where those with metallic reflections have the iridescence more intense and of a darker hue, greenish and bronzy reflections changing to purple. The slaty, ferruginous, and olive tints, and the various shades of red and yellow of others, become also far more intense. In species barred transversely with dark and light colors, the dark bands, as a general rule, become broader, and the light ones narrower. Those with white spots on a black ground have the spots reduced in size and number, the smaller ones becoming obsolete. White bars on the wings and terminal white spots on the tail feathers

* Birds of N. Amer., p. 634.

are also of less extent in southern specimens. There hence results, as already observed, a generally darker aspect in the plumage of the southern representatives of wide-ranging species; the bill and the feet also usually sharing in the general accession of coloring matter in the integuments. The difference in color between the extremely northern and the extremely southern representatives of a given species is often so great that, taken in connection with other differences, as in general size and in the size and form of the bill, the two extremes might be excusably taken for distinct species, especially if viewed aside from the connecting series between the two types formed by specimens from successively intermediate points, which beyond question show their specific identity.

As in the case of climatic variation in the bill and in general size, the variation in color differs greatly in degree in different species. Climatic difference in color is particularly striking in *Agelæus phæniceus*. In the males the black is greatly intensified and more lustrous at the South, and the red on the shoulders becomes equally heightened. Instead of the *light* red shoulder-patch, bordered externally with *whitish* or *pale yellowish-whitish*, seen in Massachusetts specimens, the shoulder-patch in the Florida males is of a brilliant *dark* red, with a *rich cream-colored* or *orange-yellow* border. While the differences in the bills of the two types might in extreme cases be taken as indicative of different sub-genera, the difference in color is as great as occurs between the northeastern type of *A. phæniceus*, and either the so-called *A. tricolor* or *A. gubernator* of the Pacific slope, or between any of these *inter se*. *Quiscalus purpureus* also affords a similar example of climatic variation, as well in color as in the bill and general size. In the males the change in general tint is in the black becoming more intense at the South, and the iridescence being dark purple or bluish instead of bronzy or greenish. The change in the females is as great as that in the males. At the North their plumage is nearly lustreless brownish-black, but at the South it becomes nearly as black as that of the northern males, and has considerable iridescence, so that the northern collector, judging from color alone, would at first be likely to mistake the southern females for males.

In *Ortyx virginianus*, through the increased breadth of the transverse bars of black at the South, on the dorsal as well as on the ventral surface, the general aspect of the plumage is very much darker in Florida

specimens than in New England ones. In *Sturnella ludoviciana* the yellow of the ventral surface in Florida specimens is far more intense than it is in northern ones; the slate color of *Galeoscoptes carolinensis* is correspondingly darker, and the ferruginous of *Harporhynchus rufus* is much redder. In *Centurus carolinus* not only are the black transverse bars on the back broader and darker, but the red on the head and abdomen becomes more extended and lustrous. In *Picus pubescens* the white spots on the wings become smaller and fewer, with a greater tendency to black streaks on the sides of the breast, a variation in the direction of *P. Gairdneri* and *P. Harrisii*, as will be noticed at length in the remarks on *P. pubescens* and *P. villosus* in Part IV. Similar differences occur between northern and southern specimens of *Picus borealis*, which are so great as to have led Mr. Cassin to regard the southern type as specifically distinct from the northern. Similar differences to those above described occur between northern and southern specimens of *Thryothorus ludovicianus*, *Troglodytes ædon*, *Geothlypis trichas*, *Colaptes auratus*, *Buteo lineatus*, and various other species, as will be described more in detail in Part IV.

The climatic variation in respect to the relative size of the white spaces on the rectrices and primary remiges may be illustrated by a single example. In northern specimens of *Pipilo erythrophthalmus* the terminal white spots of the tail feathers are found on the *four* outer feathers of each side; but in Florida-born ones they occur on only the *three* outer feathers on each side; and are correspondingly reduced in length. The white area on the tail of Florida specimens hence has only about the extent that would be presented in northern specimens if the outer pair of feathers were removed. The extent of the white space at the base of the primaries is correspondingly reduced in size in the southern type.

Extending the examination to northern species, it is found that similar color differences with the latitude of the birthplace are of frequent occurrence. In *Bernicla brenta* and *Bernicla canadensis* the smaller southern-born birds are, as a general rule, considerably darker than the larger northern-born ones. The same is true of *Fulix marila* and *Bucephala americana*, the so-called *Bucephala islandica* being the larger northern type of *B. americana*, in which the white markings on the wings and head occupy a somewhat larger area. It is altogether probable also that the so-called *Anser frontalis* holds a similar relation

to *A. Gambeli* (= *A. albifrons* ?), and the *Anser cærulescens* to the *A. hyperboreus*, though by some the former has been regarded as the young of the later. In *Larus argentatus* the southern specimens are not only smaller, with the "mantle" somewhat darker, but as a general rule the white spots at the tips of the first and second primary quills are more restricted.

The changing of the pelage to white in winter in certain northern mammals, and of the plumage in certain birds, as the ptarmigans, correlates perfectly with these geographical differences in color; and since in some species of mammals only the northern representatives change to white in winter, while the southern ones are of the same color throughout the year, this seasonal change seems evidently to come under the above-stated general law of geographical or climatic color variation, namely, a *gradual increase in color to the southward in individuals of the same species*.

A comparison of Florida birds with West India specimens of the same species shows that the difference between them in color (and, it may be added, in size and other general features) are generally not greater, and in some cases far less, especially between Cape Florida and Cuba specimens, than obtains between Florida and Massachusetts examples, and that it is of precisely the same character. West Indian specimens of course differ more from Massachusetts examples of the same species than the latter do from others from East Florida, yet by means of the South Florida specimens, which differ but slightly from the Cuba type, a gradual transition is evident from the extreme northern to the extreme southern forms. Of late many Jamaican, Porto Rican, and Cuban forms have been regarded, by many writers, as specifically distinct from their representatives in the Northern States, and in many cases they might well be so regarded, were there not a succession of intermediate forms connecting them,—a fact which seems to have been hitherto overlooked. The earlier writers considered the *Ortyx*, the *Sturnella*, the *Strix*, the *Circus*, several of the *Buteos*, etc., of the West Indies as specifically identical with the *Ortyx virginianus*, *Sturnella ludoviciana*, *Strix flammea*, *Circus hudsonius*, *Buteo borealis*, etc., of the United States, and doubtless justly, notwithstanding that the comparison of specimens reveals certain relatively slight but constant differences in color and size, and to some extent in other features.

(b) *Longitudinal Variation*. — In comparing the birds of the Atlantic States with specimens specifically identical from the interior of the continent, one is soon struck with the brighter colors of the latter, and especially with a tendency, in many species, to more ferruginous tints, and to melanism in others. In comparing again the birds of the Mississippi valley with those of the Pacific slope, especially that portion north of the fortieth parallel, a similar difference is also noticeable, the extremes of color variation in truly continental species being met with (especially to the northward of this parallel) at the Atlantic seaboard on the one hand, and the Pacific on the other, between which there is a gradual and, with an exception soon to be noticed, a uniform increase in intensity of color to the westward. This tendency to more ferruginous and melanic colors to the westward is especially marked in *Falco peregrinus*,* *Accipiter fuscus*, *Circus hudsonius*, *Buteo lineatus*, *Buteo borealis*, *Archibuteo lagopus*, *Hypotriorchis columbarius*, *Otus vulgaris*, and other species of *Strigidae*, *Tetrao canadensis*, *Bonasa umbellus*, *Bernicla canadensis*, *Bernicla brenta*, *Larus argentatus*, *Parus atricapillus*, *Carpodacus purpureus*, etc., etc. The western representatives of *Melospiza melodia*, *Passerella iliaca*, *Junco hyemalis*, *Pipilo erythrophthalmus*, *Parus hudsonicus*, etc., differ mainly from their Eastern congeners in their more ferruginous or darker colors, according to the species.

While the general tendency from the East westward is thus to darker or deeper colors in specimens of the same species, and in representative species of the same genus, the rule is not without exceptions, nor is the transition quite uninterrupted. On the arid sterile plains the representatives of not a few, and probably of most, species are much lighter colored than their relatives either to the eastward or to the westward. Also at the southward on the Pacific slope there is not the tendency to deeper colors seen farther to the northward, specimens from Northwestern Texas, New Mexico, much of the Colorado basin and Lower California, being lighter than others of the same species from Northern California, Oregon, and Washington, an explanation of which will be suggested later.†

In comparing again the European representatives of circumpolar species with their representatives in Eastern North America, a difference

* For the synonymy and other remarks on these species, see Part IV.

† See below, p. 239 *et seq.*

similar to, but hardly so great as, that between the Atlantic and Pacific coast examples of indentical species is likewise seen, the American being in general several shades darker than the European. In certain cases there is also a difference in the markings, as in some of the hawks, in which in the European the transverse bars are broader and better defined, and the longitudinal ones less so than in the American. This is illustrated in *Astur palumbarius* and *A. atricapillus*, in *Accipiter nisus* and *Ac. fuscus*, etc. In many instances the only tangible differences between so-called representative American and European species consists in the darker, brighter, or intenser color of the American, the differences being oftentimes less than that between specimens of the same species from the Atlantic States and the Mississippi valley, or between those from the Mississippi valley and the Pacific coast. Not unfrequently, however, are American and European specimens so nearly alike, even of species that have rarely been considered as identical, that without a knowledge of the locality whence they came it would be impossible to confidently refer them to the one species rather than to the other.

There are also indications of various local differences in color in specimens specifically identical within the larger areas above considered, and which are in a measure exceptional to the general law of a westward increase in color. The data at hand are at present too few either to limit these exceptional areas or to indicate to what extent they are exceptional. They appear, however, to be coincident with peculiar climatic conditions, the exact nature and extent of which are likewise imperfectly known.*

Variation in the Length of the Tail and in other Characters.—At certain localities, and more especially to the southward, there are well-known instances of an increase in the length of the tail, without an appreciable modification of other parts. Marked examples of this are seen in *Icteria virens*, *Harporhynchus rufus*, and *Mimus polyglottus*, as has been pointed out by Professor Baird and other writers,† each of which species has a western long-tailed variety. The *Quiscalus macrura* is also little else than a long-tailed variety of *Q. major*. A tendency is seen to this variation in *Geothlypis trichas* at the southward,

* See on this point below, p. 239 *et seq.*

† See especially Prof. Baird in Amer. Journ. of Science and Arts, 2d Series, Vol. XLI, p. 191.

while it seems to be a marked characteristic of many of the birds of Lower California. The tendency in southern forms to an elongation of the tail seems, however, less general than the southward decrease in size and the increase in color, or the tendency to an elongation of the bill.

Among other local variations may be mentioned the white instead of a red iris in the South Florida representatives of *Pipilo erythrophthalmus*; the yellow instead of a black bill in the magpies of the coast of California; the white basal half of the feathers of the neck of the raven of Southwestern Texas and Mexico, by which it is chiefly distinguished from the common species; the greater continuation anteriorly of the superciliary stripe in the western forms of *Zonotrichia leucophrys*, by which alone it is distinguishable from the eastern form; the white frontlet of one of the western forms of the *Parus atricapillus* group, etc. There appears frequently to be also a locally greater development of the foot in western and southern forms of wide-ranging species, and occasionally an exceptional increase in general size under identical isothermes.

Causes of Climatic Variation. — The facts respecting climatic variation are at present too imperfectly known to be fully explained. There are, however, certain peculiarities of climatic variation, especially in color, coincident with certain meteorological peculiarities of the regions where they occur, that demand attention. The increase in color to the southward, especially the tendency to darker tints above shown to be so general, coincides with the increase in the intensity of the solar rays to the southward, and in the humidity of the climate. The southward increase in depth of color and in iridescence in birds specifically identical coincides also with the general increase in brilliancy of color in birds, taken as a whole, in the lower latitudes (as well as in insects and animals generally), the maximum being reached in the tropics.

The longitudinal variation, or the westward increase in color, seems to be also coincident with the increased humidity to the westward, the darker representatives of any species occurring where the annual rain-fall is greatest, and the palest where it is least. This coincidence is clearly illustrated in the birds of the United States, where the darkest representatives of a species, as a general rule, (indeed without exception so far as known to me,) come from regions of maximum annual rain-fall, and the palest from those of minimum annual rain-fall. In the Northeastern States the amount of rain is only one half to two thirds

what it is in the Northwestern States, while on the Great Plains it is less than one half what it is in the Northeastern States. In the lower part of the Mississippi basin and in the Southeastern States it is much greater than to the northward under the same meridians. Within the tropics, in America and Asia at least, the humidity, as well as the intensity of the solar rays, reaches the maximum, as does the intensity of color in both birds and other animals. In Europe, as is well known, the birds from near the Scandinavian coast, where the annual rain-fall reaches forty inches, are darker than in Central Europe, where the yearly rain-fall is only half this amount. So much darker, in fact, are the Scandinavian forms, that by some writers they have been regarded as specifically distinct from their representatives in Southern Germany, the Scandinavian forms of circumpolar species being as dark as their Eastern North American allies. There is again a striking parallelism between the relative humidity of Western Europe and Eastern North America, and the relative depth of color in the representatives of circumpolar species living in these two countries, the rain-fall of the latter region being double that of the former, and the birds of darker and livelier colors. As already intimated, this coincidence is not confined to the birds of these different regions, the same correlation of livelier, brighter, deeper tints with increased humidity being also exhibited by the mammals of these various districts, the Europeo-North American species being higher colored, as a general rule, in Eastern North America than in Europe, as the western forms of the continentally distributed American species are often higher colored than the eastern.

It is a most striking fact that the birds, and even the mammals and reptiles, of the almost rainless districts of Lower California, the Gila and Colorado deserts, are almost all so much paler in color than their relatives of the better-watered neighboring districts, that many of them have been described as distinct species, and the others referred to as strongly marked varieties, they all being characterized to a greater or less degree by a faded or bleached aspect. The birds and mammals of the arid plains of the middle region of the continent exhibit also the same bleached appearance, but in a somewhat less degree.

I had long suspected that hygrometric conditions had much to do with local variations in color in individuals of the same species, but I was not a little surprised when I came to compare the known areas

most prolific of dark and light local forms with rain-fall charts, — which may be assumed as indicating relatively the hygrometric conditions of different regions, — to find the distribution of the light-colored races so strictly coincident with the regions of minimum mean annual rain-fall, and the dark forms with those of maximum mean annual rain-fall, as seems to be the case.

Humidity has hence apparently far more to do with climatic variation in color than solar intensity, though the latter has undoubtedly an influence upon color. The occurrence of a light-colored race of *Arvicola riparius* on Muskeget Island and the sandy sea-beaches of the coast of Massachusetts shows clearly that the intense light caused by reflection from a sandy surface tends to the diminution rather than to an increase of color in animals, and even plants, since the foliage of the latter in arid districts so commonly assumes a dull grayish tint. The capture on Muskeget Island last season (July, 1870), by Messrs. Maynard and Brewster, of two pairs of the short-eared owl (*Otus brachyotus*) with the color of the plumage so pale as at first to suggest their being albinos, is additional evidence of the bleaching effect of strong light upon the colors of animals. Such facts render it doubtful whether the increased intensity of the light in the tropics has really much to do with the brighter colors of tropical birds and insects, and suggest that humidity alone may be the principal agent in producing this accession of color.

In regard to the cause of other climatic variations, certain other facts are naturally recalled. In the remarks on the climatic and faunal peculiarities of East Florida,* attention was called to the less degree of vivacity and energy exhibited by the southern as compared with the northern members of the same species, and the general higher physiological development of essentially extra-tropical species in the temperate portions of their habitats. Is it hence improbable that the southward deterioration in size seen in such species is directly related to the enervating influence of increased heat? And why is it that so large a proportion of the birds pre-eminently singing-birds are found in temperate latitudes?

In the increased size of the bill and tail to the southward, especially of the former, we have a fact somewhat parallel to what is not unfrequently seen in mammals. The ears, for example, of the arctic repre-

* See above, p. 166.

sentatives of species ranging to warm-temperate latitudes are smaller at the northward than at the southward, as is seen in the native dogs, the foxes, and the wolves, and in the arctic races of man. The explanation generally given of this seems possibly applicable to the beaks of birds, namely, a greater activity in the circulation of the blood in the peripheral parts of the body in the temperate latitudes.

SPECIES, VARIETIES, AND GEOGRAPHICAL RACES.

The foregoing remarks on individual and geographical or climatic variation necessitates a brief consideration of the character of species, varieties, and races, and the propriety of applying binomials to such forms as can be clearly shown to be connected by intergrading links with others previously known. As preparatory to what follows, it seems proper to refer briefly to the origin of the excessive synonymy with which our descriptive ornithological works are burdened.

Ornithological synonymes may be arranged, as regards their origin, under four primary heads, namely: (1) Those arising from the description of immature and adult birds of the same species for different species, (2) from authors mistaking sexual for specific differences, (3) individual variation for specific differentiation, and (4) climatic differentiation for specific. A fifth source of error, and one which has given rise to a large class of synonymes, results from a combination of the causes indicated under (3) and (4).

Synonymes arising from the first two causes mainly preceded the others in regard to the relative frequency of their occurrence, especially so far as regards the birds of this continent. During the previous century, and the first two decades of the present, our birds were mainly described by European naturalists, who had no acquaintance with them in life, and whose resources often consisted of single and imperfect specimens received from chance travellers, without any indication of their sex or age. Later they were studied by resident naturalists, by whom the mistakes of their predecessors in this respect were to a great extent corrected. The laws of sexual and age variation becoming gradually known, errors from this source were soon far less frequent than in earlier times. When at a comparatively recent date critical comparisons were made of specimens from distant localities before regarded as specifically identical, it was found that occasionally distinct species had been confounded. Such results led in the end to undue importance

being attached to trivial differences, so that assumed species were frequently based solely on either individual or climatic variation, but oftener on both combined.

As the rage for describing new species increased, differences seemed alone to be sought; and so long as a given species was usually deemed sufficiently represented, even by the best ornithologists of the day, by a single pair,* the subject of individual and climatic variation was necessarily almost wholly neglected, the custom of many naturalists being to describe species from single specimens, as though all the representatives of a species were cast after an unvarying pattern. As the number of specimens of well-known species increased in our large museums, it was soon seen that some of the supposed most reliable diagnostic features were subject to considerable variation. The collections brought together from various parts of the continent by the Pacific Railroad surveying parties and from other sources, and the reports published thereon, formed the beginning of a new era in the history of the ornithology of North America, and in ornithological science. The facts thus disclosed in respect to geographical range, and individual and climatic variation, opened new fields of inquiry. Old theories and blind adherence to authorities, however, still impeded progress and led to frequent inconsistencies, which only time and further investigations could correct. Hence has gradually dawned the fact of the existence of a range of individual variation previously unsuspected, and of general laws of climatic variation, the full scope of which, as bearing upon the character of species, is yet to be determined.

Nearly half a century since it was discovered that the North American representatives of what were then commonly regarded as circumpolar species could not in all cases longer be regarded as identical with the European. Further comparisons showed that in most cases of the supposed circumpolar distribution of species, specimens from the Old World and the New could be more or less readily distinguished, yet the differences were in most cases slight, more or less inconstant, and not unfrequently due more to differences in the latitude whence the specimens came than to other causes. Yet a precedent for specific

* Not many years since amateur ornithologists were kindly informed, by one of the leaders in the science of ornithology, that his collection of the birds of a certain country, numbering over two thousand species, required for their convenient storage a space equal to only about one hundred cubic feet, the specimens averaging less than two to a species!

separation in such cases having been established by recognized authorities, it was followed till all the land-birds and a large proportion of the water-birds of the two continents were separated, in many cases, it would appear, on purely theoretical or geographical grounds.* When the comparison was carried to specimens of continentally distributed species from distant localities, differences between these were also detected, and the theory of specific diversity assumed, till the Pacific representatives of such species were separated from the Atlantic ones, and in like manner the southern from the northern, and those of particular areas, as insular, peninsular, and interior basins, from the others. In some cases such separations were of course properly made, but a high percentage of such forms are now found to intergrade through specimens from the intermediate localities.

Not a few of the species of our faunal lists have been based on, and are still only known from, single specimens, and often on differences manifestly within the range of individual variation; others represent local races, which only appear distinct when extremes alone are considered, the intermediate stages being unknown or ignored. The increase of synonymes from this fruitful source appears to have not yet culminated, a large proportion of the "new species" now annually described being but slight local differentiations of previously known specific forms, from which they often differ only in being a little smaller, a little darker or brighter colored, and in the individual peculiarities of the single specimens on which some of them are based. In many cases this process of ultra subdivision has furnished stepping-stones to later generalizations; in too many other cases it has been in its results only unmitigatedly injurious.

So large a proportion of the commonly recognized species are virtually nominal, or rest on a false basis, it is not surprising that in the reaction consequent upon a fuller knowledge of the birds of this continent, which has already commenced, the reality of species should be to some extent ignored. Whether, however, species are considered as entities or only as arbitrary inventions, convenience demands some established definition of them.

* Audubon, writing in 1838 (Orn. Biog., Vol. IV, p. 608), refers to the Prince of Musignano (by whom a large part of the circumpolar and cosmopolitan species were separated into numerous assumed species) as "having altered his notions so far as to seem desirous of proving that the same species of birds cannot exist on both the continents"; and there seems to have been good reason for the remark, only instead of *proving* them distinct, he in most cases merely *assumed* them to be so.

Not a few naturalists have hence adopted the test of intergradation, which seems a reasonable and an unobjectionable one. The question of species and of specific synonymy is thus simplified to this: that whenever two forms which have both received names are found to intergrade, the more recent name shall become a synonyme of the older. Some, however, still urge that every recognizable form, however closely allied to others, and even intergrading, should be recognized by a binomial epithet, and that whether we call them species, or varieties, or races, or simply *forms*, that such names are none the less convenient expressions for certain facts. It seems to me, however, that there are insuperable objections to this course; for however distinct the extreme geographical forms of a species may be, a vast proportion of its representatives are intermediate to them, and could never be but doubtfully referred to the one rather than to the other. Ordinarily, for instance, in the birds of the Atlantic slope, the representatives of a given species at the extreme north of its breeding range almost always differ very tangibly from its representatives at the extreme southern limit, sometimes more, sometimes less, according to the species. Those living only a little to the northward of the middle region differ less from the extreme southern type than the extreme northern type does, and those a little to the southward of the middle region differ still less from the southern type, and are quite distinguishable from the extreme northern form. In other words, in species ranging from Southern Labrador or Northern New England to Florida, of which there are numerous unquestioned instances, specimens from Southern New England differ somewhat from the more northern ones; those from Southern New England from those of Southern New Jersey and Eastern Maryland, and these latter from those of Georgia and Florida. It hence depends entirely upon individual predilection whether two, three, or four "species" or "binomial forms" shall be recognized; and in either case there is the same difficulty in disposing of the intermediate types. Again, specimens from the Mississippi valley differ more or less from their relatives from the Atlantic coast, the central plains, and the Pacific slope. Here again similar difficulties are encountered. Hence it is necessary to decide between recognizing a single binomial form, with a considerable but definite range of climatic variation, or three, or six, or nine, or even more, which cannot be rigidly defined, and between each of which will always be found a greater or less proportion of intermediate types,

doubtfully referable to one of the binomial forms rather than to another. Another important objection may be urged against giving binomial names to intergrading forms. In faunal and nominal lists of the species of a large or continental area, scarcely distinguishable forms take equal rank with the most distinct congeneric species. For instance, in a list of the birds of North America, *Turdus Aliciæ* and *Turdus Swainsoni*, *Turdus Auduboni* and *Turdus Pallasi*, stand side by side with *Turdus mustelinus* and *Turdus fuscescens*, though in the former cases *Turdus Aliciæ* and *T. Auduboni* are founded at best on slight, and in the one case on inconstant individual or local differences, while in the latter no two congeneric species need be more distinct. In the one case only experts can distinguish the forms, and frequently they only by an actual comparison of specimens, and then too frequently but doubtfully, while in the other case a casual observer need not mistake them. The names alone give no clew to their real character, and are hence in a great measure meaningless when separated from the most explicit diagnoses, and whose affinities can frequently only be settled by the arbitrary criterion of locality. But it is urged that cognizance should in some way be taken of these differences; and "How can they be better recognized," it may be asked, "than in the way proposed?"

As already shown, and as I trust a large proportion of ornithologists are willing to admit, these local forms occur in accordance with recognizable laws of climatic variation, similar variations with locality occurring, to a greater or less extent, in all species having nearly the same geographical range. Eventually, then, will not the recognition of these laws be sufficient, and should not a statement of the tendencies to variation with locality, and the degree to which it is developed, be embraced in the specific diagnosis of each species as a part of its specific description? Is not this, in fact, actually essential to the proper characterization of a species? The average characters being given, a line or two would suffice for a statement of its variations, both geographical and individual. Then only in one case where now there are hundreds would there be instances of doubtful identification. Till within a very recent period, perhaps, no other course could have been pursued than that of giving binomial names to each apparently distinct form, however slightly it may have differed from others previously known. In many cases, indeed, the differences between strictly intergrading geographical forms are very great,—greater, indeed, if they were not thus serially con-

nected, than would be deemed necessary for specific separation ; and so long as the extremes only were known, no one could have regarded them otherwise than as well-defined species. But the time has already come, it seems to me, for a different and a more philosophic method, and that to further increase synonymy by giving new names to slightly different local forms of the same species is worse than useless.

It is important, in this connection, to observe that the species occurring at any point on the Atlantic coast, or on the Pacific coast, or in the Mississippi valley, or on the Great Plains, in short, at any restricted locality, have, as compared with each other, with scarcely an exception, an unequivocal character ; they are based on differences that place them beyond controversy. It is not so, however, when we compare the species of distant localities with each other, whether the localities differ in latitude or longitude. In such cases we constantly meet with controverted species. At the South are species admitted as doubtfully distinct from others found farther north ; at the West, those holding the same relation to others of the East ; while at intermediate points either both the disputed forms occur with greater or less frequency, or there is a gradual transition of the one into the other, neither form being typically represented. This is evidently what should be expected to occur, if what has been said above in respect to climatic variation be correct, and is evidently a suggestive and important fact. Is the theory of hybridization, so often appealed to in such cases, necessary to explain these facts ? and is it, in fact, true ? By uniting the intergrading forms, the number of species occurring at any single locality is not essentially reduced, but such a union would considerably reduce the total number recognized, as well as the number usually assigned to the several continents, as at present not a few are repeatedly counted.

The many facts bearing upon individual and geographical variation, presented in the foregoing pages form but an imperfect exposition of the subject. They are, nevertheless, eminently suggestive of interesting results, and the conclusions above deduced I can but believe will be only the more fully confirmed by further research. Additional details are given in the general remarks embraced in Part IV, where various facts merely hinted at above are more fully presented, and an application is made in many cases of the principles deducible from them.

As previously stated, individual and geographical variations are in

some cases difficult to distinguish. They can be satisfactorily investigated only from extensive suites of specimens taken from the same locality in the *breeding season*, and sufficiently extensive suites of this character are, with rare exceptions, still wanting. In specimens taken during migration it is difficult to determine what share of the variation is due to birthplace and what to individuality. Whilst, however, the variations noticed cannot be always traced with certainty to their origin, their bearing upon the general subject of variation within specific limits is in no way vitiated. In considering hypothetical species, it is frequently clearly evident that they are based in part upon slight and tolerably constant climatic differences, and in part and sometimes wholly upon the individual peculiarities of the single specimen upon which the original description of the species was based; in part, too, upon seasonal differences, and upon characters of immaturity. It seems to me that in the numerous closely allied species of the *Ægiothus* group, to cite a case in point, some are based in part upon one and in part upon other of these differences of a single circumpolar species. As already shown, the bill in different specimens of *Æ. linarius* varies greatly in size, yet an examination of a considerable series of specimens of several of its allies shows an amount of variation in the bill closely approximate to that seen in the specimens of the various assumed species of *Ægiothus*. Much of the variation in color seen in the flocks of *Ægiothi* that visit the Northern States in winter is due to age, yet it has been taken as characteristic of different species. These birds only visiting us in winter, those inhabiting widely distant localities in the breeding season are probably then more or less associated. The light-colored specimens are doubtless in part old or fully mature birds, or inhabitants in summer of more northern districts than the browner or more fulvous ones, a large portion of which, however, are unquestionably young birds. The short-billed ones have also relatively longer setæ at the base of the bill, which, by concealing a large portion of it, give it the appearance of being shorter than it really is. Analogy would lead us to infer that those with the shorter and more heavily clothed bills have a more northern habitat than the others.

The persistency with which nominal species when characterized by "authorities" are retained in our literature is not a little remarkable. If specimens from the original localities cannot be found to exactly fit the descriptions, the diagnosis is slightly amended to suit examples that

somewhat approach them, and the name retained. In other cases the species is retained without its character being questioned, the name and the original description being copied by succeeding writers, till the species becomes traditionally accepted without its claims to recognition having been critically examined.

Another noteworthy coincidence in regard to nominal species is the fact of their most frequent occurrence in obscurely known groups, which obscurity usually results from the difficulty of obtaining specimens of the forms in question, — either from the remoteness of their habitat, their scarcity, or the peculiarities of their habits, — or from preconceived notions of the intimate relationship of the species of such groups.

Since the above was put in type, I have for the first time met with some important and timely remarks by an eminent English botanist concerning variation within specific limits in plants, which are so apropos to what has been said above in regard to individual and climatic variation in birds, and contains, moreover, such judicious strictures on various practices indulged in by botanists, and of which zoölogists are equally guilty, that a short abstract of them forms a fitting conclusion to the present paper. Says Dr. J. D. Hooker, in the introductory essay to his "*Flora Novæ-Zelandiæ*" (Part I, pp. xii, xiii, xv, 1853): —

"Some naturalists consider every minute character, if only tolerably constant or even prevalent, as of specific value; they consider two or more doubtful species to be distinct till they have been proved to be one; they limit the ranges of distribution, and regard plants from widely severed localities as almost necessarily distinct; they do not allow for the effects of local peculiarities in temperature, humidity, soil, or exposure, except they can absolutely trace the cause to the effect; and they hence attach great importance to habit, stature, color, hairiness, period of flowering, etc. These views, whether acknowledged or not, are practically carried out in many of the local floras of Europe, and by some of the most acute and observant botanists of the day; and it is difficult to overestimate the amount of synonymy and confusion which they have introduced into some of the commonest and most variable of plants. . . . In working up incomplete floras especially, I believe it to be of the utmost importance to regard dubious species as varieties, to take enlarged views of the range of variation in species, and to weigh characters not only *per se*, but with

reference to those which prevail in the order to which the species under consideration belong; and to resist steadily the temptation to multiply names; for it is practically very difficult to expunge a species founded on an error of judgment or observation. The state of the British flora proves not only this, but further, that one such error leads to many more of the like kind; students are led to overestimate inconstant characters, to take a narrow view of the importance and end of botany, and to throw away time upon profitless discussions about the differences between infinitely variable forms of plants, of whose identity really learned botanists have no doubt whatever. There is, further, an inherent tendency in every one occupied with specialties to exaggerate the value of his materials and labors. . . .

"To the amateur these questions are perhaps of very trifling importance, but they are of great moment to the naturalist who regards accurately defined floras as the means of investigating the great phenomena of vegetation; he has to seek the truth amid errors of observation and judgment, and the resulting chaos of synonymy which has been accumulated by thoughtless aspirants to the questionable honor of being the first to name a species. The time, however, has happily passed when it was considered to be an honor to be the namer of a plant; the botanist who has the true interests of science at heart not only feels that the thrusting of an uncalled-for synonyme into the nomenclature of science is an exposure of his own ignorance and deserves censure, but that a wider range of knowledge and a greater depth of study are required to prove those dissimilar forms to be identical, which any superficial observer can separate by words and a name."

The above remarks are as strictly applicable to zoölogy and zoölogists as they have ever been to botany and to botanists. The present state of ornithology, and the tendency the majority of ornithologists have to multiply species on improper grounds, find here a fitting rebuke.

PART IV.

*List of the Winter Birds of East Florida, with Annotations.**

TURDIDÆ.

1.† *Turdus migratorius* Linné. ROBIN.

Seen daily, sometimes in considerable flocks, till about the first of March, after which time few were observed. It was shot by me at

* An asterisk (*) prefixed to the name of a species indicates that it is a constant resident; an obelisk (†), that it is a winter visitor.

Jacksonville, April 1st, but according to general report it does not breed in the State.

In this species the females are commonly supposed to be paler colored than the males, which is undoubtedly usually the case, but specimens as brightly colored as any I ever saw proved on dissection to be *females*, and other specimens as palely colored as any I ever met with have likewise proved on dissection to be *males*. This shows the importance of determining the sex in all cases by dissection, and not from external appearances. It also indicates a wide range of variation in color in the present species, as great as is seen between typical representatives of the so-called *Turdus Swainsoni* and *T. Aliciae*, and which is, moreover, of the same character, namely, simply a variation in intensity.

2† *Turdus Swainsoni* *Cabanis*. OLIVE BACKED THRUSH.

Turdus minor GMELIN, Syst. Nat., I, 817, 1788; in part only. — VIEILLOT, Ois. Am. Sept., II, 7, pl. lxiii, 1807; in part only. — BONAPARTE, Geog. and Comp. List, 1838.

Turdus solitarius WILSON, Am. Orn., V, pl. xiii, fig. 2; not the text.

Turdus nanus AUDUBON, Birds of Amer., III, pl. cxlvii; * not the text. — SAMUELS, Am. Nat., II, 218, 1868

Turdus olivaceus GIRAUD, Birds of Long Island, 92, 1843-44. Not the *T. olivaceus* of Linné.

Turdus Swainsonii CABANIS, "in Tschudi's Fauna Peruana, 188, 1844-46." — BAIRD, Birds N. Am., 216, 1858. — SCLATER, Cat. Am. Birds, 2, 1862. — ALLEN, Proc. Essex Inst., IV, 56, 1864. — BAIRD, Rev. Am. Birds, I, 19, 1864. — ALLEN, Mem. Bost. Soc. Nat. Hist., I, 514, 1868. — RIDGWAY, Proc. Phil. Acad. Nat. Sci., XXI, 128, 1869.

Turdus Aliciae BAIRD, Birds N. Am., 217, 1858. — COUES and PRENTISS, Smithsonian Rep., 1861, 405. — COUES, Proc. Phil. Acad. Nat. Sci., XIV, 217, 1861. — BAIRD, Rev. Am. Birds, I, 21, 1864. — RIDGWAY, Proc. Phil. Acad. Nat. Sci., XXI, 128, 1869.

Merula Wilsoni SWAINSON, Faun. Bor. Am., I, 182, 1831.

Merula olivacea BREWER, Proc. Bost. Soc. Nat. Hist., I, 191, 1844.

Rare. Given on the authority of Mr. Boardman, who writes me he obtained one specimen at Enterprise, February 18th, and another at St. Augustine, in the same month. The greater part pass the winter farther south.

* The plates in "Birds of America" are too poorly colored, as is well known, to be recognizable representations of the species whose names they bear, including all those representing wood-thrushes, they having but little resemblance to those of the folio edition. The figures of "*Turdus nanus*," *Turdus solitarius*, and *Turdus mustelinus*, might all pass for the *Turdus Swainsoni*, so far as the color of the dorsal surface is concerned.

In my "Catalogue of the Birds of Massachusetts,"* published in 1864, I first advanced the opinion that the so-called *Turdus Aliciæ* Baird was the paler form of *T. Swainsoni*. To this view other writers have taken exception. Professor Baird, in his "Review of American Birds" (p. 21), summarily disposes of the matter by presuming that I had not seen what he called *T. Aliciæ*. In 1868, in my "Notes on the Birds of Iowa, Illinois," etc.,† I again reviewed the subject, having in the mean time examined some twenty specimens sent out by the Smithsonian Institution to different scientific institutions, labelled respectively, "*Turdus Aliciæ*," "*Turdus Aliciæ?*" "*Turdus Aliciæ? hybrid?*" "*Turdus Swainsoni*," "*Turdus Swainsoni?*" "*Turdus Swainsoni? hybrid?*" After having examined these authentic specimens of the bird in question, and also large numbers of Massachusetts examples of what I called *Turdus Swainsoni*,—among which are a considerable number that correspond in every particular respectively with the typical, authentic specimens of "*Turdus Swainsoni*" and "*Turdus Aliciæ*" of Baird, the larger number, however, being intermediate in character between them, and agreeing with specimens sent out from the Smithsonian Institution as "*T. Swainsoni?*" "*Turdus Aliciæ?*" "*Turdus Aliciæ? hybrid?*" etc.,—I state in this paper that the opinion I had previously expressed in respect to *Turdus Swainsoni* and *Turdus Aliciæ* was fully confirmed. In this paper I discussed at some length the variations presented, not only by this species, but by *Turdus Pallasi* and *Turdus fuscescens*, and the character of their supposed allies, *T. Auduboni*, *T. nanus*, and *T. ustulatus*, and their supposed respective habitats. I gave also some details in respect to the variations in general size, form of the bill, proportions of the primary quills of the wing, etc., as well as in color, and concluded that *Turdus Aliciæ* was based on simply individual variation in color, the other differences, as of size, form of bill, etc., supposed at first to characterize it, being rarely coincident with the variations in color, they occurring as frequently in the one type of coloration as in the other. *Turdus nanus* and *Turdus ustulatus* I also deemed to hold the same relationship to *T. Pallasi* and *T. fuscescens* that *T. Aliciæ* does to *T. Swainsoni*. Though described as exclusively western, I stated I had found specimens in Massachusetts that accorded with them in every particular. After having given the subject still further attention, I am but the more fully confirmed in these opinions.

Dr. Coues, thus far one of the most strenuous advocates of the validity of these nominal species, in a somewhat recent paper of his,‡ after stating

* Proceedings of the Essex Institute, Vol. IV, p. 56.

† Memoirs of the Bost. Soc. Nat. Hist., Vol. I, p. 507.

‡ "A List of the Birds of New England," Proceedings Essex Institute, Vol. V, p. 267, 1868.

that he had shown the *T. Aliciæ* to be "a very common eastern bird, having a range of habitat as extensive as, and nearly identical with, that of *T. Swainsoni*," says, in referring to my earlier remarks on this subject, that they "illustrate very fully the well-known seasonal and other variations to which *T. Swainsoni* and *T. fuscescens* are subject," and adds that I appear to have been "autoptically unacquainted" with *T. Aliciæ* at the time of writing them. In respect to this supposition of Dr. Coues, I will merely add that one of the numerous specimens considered by me to typically represent the supposed *T. Aliciæ* has been sent to the Smithsonian Institution, and pronounced by Professor Baird himself to "typically represent the *T. Aliciæ*."

The measurements given below of this species and the two following indicate the average size and the usual range of variation in this respect in these species as represented in the Atlantic States. These measurements embrace twenty-four specimens of *Turdus Swainsoni*, nearly fifty of *T. Pallasi*, and about forty of *T. fuscescens*, nearly all of which are from New England, and by far the greater part from Eastern Massachusetts.

The following is the range of variation in the series of twenty-four specimens of *T. Swainsoni*: Length, 6.62 to 7.75; alar extent, 10.75 to 12.65; wing, 3.47 to 4.30; tail, 2.40 to 3.40 (4.00?); tarsus, 1.02 to 1.27. The average dimensions are as follows: Length, 7.17; alar extent, 11.65; wing, 3.86; tail, 2.88; tarsus, 1.15.

Measurements of New England Specimens of TURDUS SWAINSONI.

M. C. Z. No.	Collector's Number.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.	Tarsus.
2877	—	—	Springfield, Mass.	May 14, '63	J. A. Allen	6.75	11.90	3.19	2.78	1.12
—	—	—	" "	May 14, '63	"	6.62	11.40	3.80	2.83	1.11
—	18	—	" "	May 27, '61	"	6.75	—	3.92	2.78	1.10
—	19	♂	" "	May 27, '61	"	6.98	—	4.30	3.40	1.22
2930	—	—	" "	May 25, '63	"	7.60	12.50	4.12	3.10	1.20
—	37	—	" "	May 25, '63	"	7.06	11.40	3.65	2.86	1.13
2940	—	—	" "	May 29, '63	"	7.10	12.00	4.00	2.83	1.13
—	44	♂	" "	May 29, '63	"	7.25	12.00	4.00	2.95	1.10
2827	—	—	" "	May 14, '63	"	7.15	11.10	3.90	2.70	1.05
1829	29	—	" "	May 30, '62	"	6.90	11.20	3.55	2.58	1.02
1830	23	—	" "	May 30, '62	"	7.75	12.20	3.95	3.07	1.17
1831	1	—	" "	May 30, '62	"	7.35	11.50	3.80	2.73	1.10
2930	—	—	" "	May 25, '63	"	7.75	12.20	3.95	3.07	1.17
2940	—	—	" "	May 29, '63	"	7.38	11.37	3.93	2.87	1.14
5757	—	—	Concord, "	—	H. Mann	7.25	12.15	4.10	2.95	1.16
8844	646	♂	Belmont, "	May 27, '68	C. J. Maynard	7.76	12.65	4.20	4.00	—
4999	1326	—	" "	Sept. 21, '68	"	7.75	—	4.05	3.07	1.20
—	17	♀	Watertown, "	Oct. 2, '69	Wm. Brewster	7.12	12.00	3.94	3.00	1.19
—	213	♀	" "	May 24, '69	"	7.12	—	4.00	2.87	1.27
282	—	—	Malden, "	—	D. Higgins	7.10	11.00	3.70	2.48	1.08
397	—	—	Norway, Maine	—	A. E. Verrill	7.24	11.00	3.47	2.73	1.14
308	—	—	" "	—	"	7.00	10.75	3.48	2.40	1.12
5963	—	—	Upton, "	—	J. G. Rich	7.30	11.10	3.71	2.68	1.03
1520	—	—	Glen House, W. Mts.	—	S. H. Scudder	7.00	11.50	3.84	2.74	1.10

3.† *Turdus Pallasi* Cabanis. HERMIT THRUSH.

Turdus solitarius WILSON, Am. Orn., V, 95, 1812. Not the figure (pl. xliii, 2), which is of *T. Swainsoni*. Not *T. solitarius* LINNÉ. — BONAPARTE, Geog. and Comp. List, 17, 1838. — AUDUBON, Synop., 91, 1839. — IBID., Birds of Amer., III, 29, pl. cxlvi, 1841.

Turdus minor BONAPARTE, Obs. on Wilson's Nomenclature, Journ. Phil. Acad., IV, 33, 1824. — NUTTALL, Man. Am. Orn., I, 346, 1830. — AUDUBON, Orn. Biog., I, 303, pl. lviii, 1831. — IBID., V, 445, 1839. — GAMBEL, Proc. Phil. Acad. Nat. Sci., III, 113, 1846. — GIRAUD, Birds of Long Island, 90, 1843-44.

Turdus Pallasi CABANIS, Wieg. Archiv, I, 205, 1847. — BAIRD, Birds N. Am., 212, 1858. — SCLATER, Cat. Am. Birds, 2, 1862. — BAIRD, Review Am. Birds, Part I, 14, 1864. — ALLEN, Mem. Bost. Soc. Nat. Hist., I, 514, 1868. — RIDGWAY, Proc. Phil. Acad. Nat. Sci., XXI, 128, 1869.

Turdus nanus AUDUBON, Orn. Biog., V, 201, pl. ccccxix, 1839 (*T. minor* on the plate). — IBID., Birds of Am., III, 32, 1841. — BAIRD, Birds N. Am., 213, 1858. — SCLATER, Cat. Am. Birds, 2, 1862. — BAIRD, Rev. Am. Birds, I, 15, 1864. — RIDGWAY, Proc. Phil. Acad. Nat. Sci., XXI, 129, 1869. — COOPER and BAIRD, Orn. Cal., I, 4, 1870.

Turdus Audubonii BAIRD, Rev. Am. Birds, I, 16, 1864. — RIDGWAY, Proc. Phil. Acad. Nat. Sci., XXI, 129, 1869.

Merula solitaria SWAINSON, Faun. Bor. Amer., II, 184, pl. xxxvii, 1831. — BREWER, Proc. Bost. Soc. Nat. Hist., I, 191, 1844.

Merula silens SWAINSON, Faun. Bor. Amer., II, 186, 1831. — SCLATER, Cat. Am. Birds, 2, 1862.

Common. Last seen about March 25th.

As already observed in the remarks under *Turdus Swainsoni*, I regard the *Turdus nanus* of authors as identical with *T. Pallasi*. The assumed differences are slight and inconstant, and seem to be principally individual variation in color. Although of late supposed to be exclusively western, representing on the Pacific slope the *T. Pallasi* of the Atlantic and Central States, Audubon's original specimen came from Pennsylvania, though he subsequently received it from the valley of the Columbia River. In his "Synopsis" he gives its habitat as "Columbia River. Accidental in the United States." His description of its color is identical with that he gives of *T. Pallasi* (*T. solitarius* Aud.), even the words used being almost entirely the same throughout each description. In size, however, he gives *T. nanus* as being one inch less in length and one inch less in extent than *T. Pallasi*. Since Professor Baird, in 1858, recognized the *T. nanus* as a valid species and its habitat as "Pacific coast of North America to the Rocky Mountains," and restricted the *T. Pallasi* to "Eastern North America to the Mississippi River," the validity of *T. nanus* has been generally accepted. Professor Baird himself, however, speaks of it in this work

as though it was in his opinion doubtfully distinct, and observes that, "if really distinct, is so closely allied to *T. Pallasi* as to render a separation of the two exceedingly difficult." The *T. Pallasi* was formerly recognized as inhabiting California by good authorities. Dr. Gambel, in his "Remarks on the Birds of Upper California," etc.,* after stating that "the dwarf thrush of Audubon was founded upon specimens from the Atlantic States, and no doubt upon the true hermit thrush," remarks: "An examination of specimens of the *T. minor* [= *T. Pallasi*] from the Atlantic and Pacific coasts of North America shows no difference in any way, except that perhaps the western one is somewhat smaller, yet the difference is scarcely appreciable. From the measurement of many western specimens I found its length to be $6\frac{1}{4}$ inches, and the extent of wings $10\frac{1}{2}$ inches; the tail, wings, and relative length of quills the same as in our eastern one, and, in fact, I think it can in no possible way be distinguished as specifically different." California specimens, however, seem to average a little smaller than New England ones, so that the *T. nanus* seems best entitled to recognition of any of the several disputed forms of this group.

The habits of *T. nanus*, as described by Dr. Cooper, are exactly like those of the *T. Pallasi* of the East, except in regard to the situation of its nest, *his account of its nest and eggs according exactly with those of T. Swainsoni*, and not at all with those of *T. Pallasi*, its nearest ally.†

The *Turdus Auduboni* of Baird, of the Rocky Mountains, I have already also referred to *T. Pallasi*, from average specimens of which it differs only in being slightly larger. My reasons for this opinion have been given with sufficient detail elsewhere.‡

It is difficult to reconcile the account given by Wilson,§ and corroborated by Audubon,|| of the breeding habits of this species with what is now known of the distribution in the breeding season of this group (subgenus *Hylocichla*) of thrushes. The account given by these authors of the situation and structure of the nest is applicable to only *T. Swainsoni*, which, as well as the *T. Pallasi*, is not known to breed so far south by several hundred miles as the localities they give. The only species which may probably breed there is the *T. fuscescens*; but this species does not nest on trees. To determine to which species of thrush these authors refer

* Proc. Phil. Acad. Nat. Sci., Vol. III, p. 14, October, 1844. Also Journal Phil. Acad. Nat. Sci., 2d Series, Vol. I, p. 41, 1847.

† According to Professor A. E. Verrill, the *T. Pallasi* nests on the ground, and lays "bright-blue" eggs. Proc. Essex Inst., Vol. III, p. 145.

‡ Mem. Bost. Soc. Nat. Hist., Vol. I, p. 512.

§ Am. Orn., Vol. V, p. 91.

|| Orn. Biog., Vol. I, p. 303: Birds of America, Vol. III, p. 30

as breeding in this manner on the Lower Mississippi would solve an interesting problem.

The following table will indicate the average size of *Turdus Pallasi* in the Atlantic States. The extremes in size of forty-six specimens are as follows: Length, 6.50 and 7.65; alar extent, 10.00 and 12.25; wing, 3.30 and 3.90; tail, 2.47 and 3.17; tarsus, 1.12 and 1.33. The average dimensions of these specimens are as follows: Length, 7.04; alar extent, 11.17; wing, 3.79; tail, 2.72; tarsus, 1.15.

Measurements of Specimens of TURDUS PALLASI.

M. C. Z. No.	Collector's Number.	Sex	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.	Tarsus.
9835	—	—	Milltown, Maine	June —, '64	G. A. Boardman	6.98	—	3.50	2.68	1.17
4071	—	—	Upton, "	June —, '65	J. G. Rich	6.88	10.00	3.48	2.67	1.16
1990	—	—	Norway, "	—	A. E. Verrill	6.90	10.50	3.30	2.51	1.15
1991	—	—	"	—	"	7.00	10.60	3.55	2.65	1.17
5610	—	—	"	—	S. I. Smith	7.25	11.55	3.73	2.93	1.20
5611	—	—	"	—	"	6.80	10.95	3.45	2.78	1.12
312	—	—	"	—	"	7.15	11.50	3.70	2.92	1.15
321	—	—	"	—	"	7.32	11.00	3.59	2.98	1.17
1897	—	—	Waterville, "	Apr. 20, '62	C. E. Hamlin	7.20	10.77	3.66	2.93	1.07
1943	—	—	"	Apr. 14, '62	"	6.80	10.75	3.49	2.65	1.18
4235	—	—	"	Oct. 21, '63	"	6.90	10.25	3.37	2.60	1.18
4251	—	—	"	Oct. 24, '63	"	6.80	10.75	3.57	2.93	—
5754	—	—	Concord, Mass.	—	H. Mann	7.50	11.82	3.83	2.95	1.23
5755	—	—	"	—	"	7.15	10.15	3.45	2.57	1.14
4960	—	—	"	—	"	7.25	11.62	3.77	2.94	1.18
2584	—	—	Woburn, "	—	J. G. Shute	6.80	10.25	3.36	2.57	1.17
2832	—	—	Springfield, "	May 8, '63	J. A. Allen	7.25	10.70	3.52	2.83	1.20
9690	1002	—	"	Oct. 17, '63	"	7.10	11.20	3.57	2.70	1.13
9691	1021	—	"	Oct. 29, '63	"	7.00	10.50	3.47	2.74	1.18
—	49	♂	Watertown, "	Dec. 10, '69	Wm. Brewster	7.00	11.63	3.63	2.63	1.33
—	58	♂	"	Nov. 6, '69	"	6.87	11.19	3.61	2.52	1.20
—	60	♂	"	Nov. 10, '69	"	7.12	11.12	3.74	2.58	1.27
—	62	♂	Belmont, "	Nov. 22, '69	"	7.12	11.50	3.80	2.71	1.20
—	333	♂	Cambridge, "	Apr. 16, '70	"	7.23	11.94	3.71	2.85	1.29
—	59	♂	Waltham, "	Oct. 26, '69	"	7.20	11.12	3.74	2.58	1.27
—	61	♂	"	Nov. 1, '69	"	7.06	11.37	3.72	2.59	1.25
—	62	♂	Watertown, "	Oct. 26, '69	"	6.50	10.50	3.59	2.37	1.15
—	283	♂	"	Nov. 22, '69	"	6.56	11.00	3.50	2.58	1.16
—	353	♂	"	Apr. 26, '70	"	6.75	11.30	3.45	2.50	1.17
8845	8	♂	Newton, "	Oct. 12, '67	C. J. Maynard	6.83	11.00	3.32	2.65	—
—	13	♂	"	Oct. 12, '67	"	7.00	10.77	3.37	2.47	—
8848	250	♂	"	Apr. 18, '68	"	7.21	11.32	3.60	2.80	—
8852	322	♂	"	Apr. 25, '68	"	7.27	11.00	3.50	2.60	—
8847	338	♂	"	May 5, '68	"	7.00	11.30	3.59	2.73	—
—	24	♂	"	Oct. 19, '68	"	7.00	11.47	3.45	2.75	—
—	14	♂	"	Oct. 16, '68	"	7.00	11.43	3.50	2.74	—
8846	26	♂	"	Oct. 16, '68	"	6.80	11.28	3.57	2.75	—
8849	311	♂	"	Apr. 25, '70	"	7.60	11.75	3.43	3.17	—
—	314	♂	"	Apr. 25, '70	"	7.38	12.83	3.80	2.83	—
8853	333	♂	"	Apr. 28, '70	"	7.45	11.85	3.90	2.90	—
5129	—	?	Jacksonville, Fla.	Jan. 21, '68	J. A. Allen	7.40	12.25	3.85	—	—
5145	—	?	"	Jan. 25, '68	"	7.00	11.60	3.60	—	—
5146	—	?	"	Jan. 25, '68	"	6.75	11.10	3.45	—	—
5147	—	?	"	Jan. 25, '68	"	7.00	11.50	3.60	—	—
5197	—	?	Hibernia, "	Feb. 3, '68	"	7.65	11.87	3.65	—	—
5320	—	?	Enterprise, "	Mar. 1, '68	"	6.75	10.90	3.40	—	—

4.† *Turdus fuscescens* Stephens. WILSON'S THRUSH.

Turdus mustelinus WILSON, Am. Orn., V, 98, pl. xliii, 1812. (Not *T. mustelinus* GMELIN.)

Turdus fuscescens STEPHENS, Shaw's Gen. Zoöl., X, i, 182, 1817. — G. R. GRAY, Gen. Birds, 1849. — BAIRD, Birds N. Am., 214, 1858. — SCLATER, Cat. Am. Birds, 2, 1862. — BAIRD, Rev. Am. Birds, I, 17, 1864. — ALLEN, Mem. Bost. Soc. Nat. Hist., I, 514, 1868. — RIDGWAY, Proc. Phil. Acad. Nat. Sci., XXI, 127, 1869.

Turdus Wilsonii BONAPARTE, Obs. on Wilson's Nomenclature. — NUTTALL, Man. Am. Orn., I, 349, 1832. — AUDUBON, Orn. Biog., II, 362, pl. clxvi, 1834. Ibid., V, 446. — GIRAUD, Birds L. Island, 89, 1843-44.

Turdus ustulatus NUTTALL, Man. Am. Orn., I, (2d ed.) 400, 1840. — BAIRD, Birds N. Am., 215, 1858. — IBID., Rev. Am. Birds, I, 18, 1864. — RIDGWAY, Proc. Phil. Acad. Nat. Sci., XXI, 127, 1869. — COOPER & BAIRD, Orn. Cal., I, 5, 1870.

Merula minor SWAINSON, Faun. Bor. Am., II, 179, pl. xxxvi, 1831.

Merula Wilsonii BREWER, Proc. Bost. Soc. Nat. Hist., I, 191, 1844.

Not common, the greater part passing the winter in the tropics. A few specimens were taken by Mr. Boardman at Green Cove Springs, February 20th and 22d. I did not meet with it.

The considerable variation in color exhibited by different specimens of this species have perhaps been already sufficiently adverted to. It may be added that some of the brightest colored specimens of this species proved on dissection to be females, as well, also, as some of the palest. As in *T. migratorius*, *T. Swainsoni*, etc., these variations in color do not depend entirely upon sex, age, nor season. The latter, however, doubtless has much to do with it, as has also age, as already explained; * but the variation is in the main strictly the result of individual differentiation.

Dr. Cooper says † that in habits this species is the "exact counterpart of *T. nanus*," the resemblance extending to the situation and structure of the nest, and also to the color of the eggs. In this connection it may be remarked that it is not a little remarkable that the eggs and nests of both the so-called *T. ustulatus* and *T. nanus* should so exactly coincide with those of *T. Swainsoni* (which breeds where the other species are said to), when the birds themselves are scarcely distinguishable respectively from *T. fuscescens* and *T. Pallasi*, both of which nest on the ground and lay unspotted eggs, while *T. Swainsoni* nests in trees and lays spotted eggs. The nests and eggs I have seen purporting to be those of *T. ustulatus* and *T. nanus* (and also of *T. Aliciæ*) were so closely like those of *T. Swainsoni*, — not differing more from those of this species than those of the same species usually differ, — as to at once raise the suspicion in my mind that they might all be really those of *T. Swainsoni*, and that they may have been in some accidental way wrongly identified by the collector.

* In Part III, pp. 193 *et seq.*

† Ornithology of California, Vol. I, p. 5.

In the following table are given the measurements of forty specimens, some twenty-five of which were taken in Massachusetts during the breeding season. The extremes of the series are as follows: Length, 6.95 and 7.87; alar extent, 11.05 and 12.70; wing, 3.55 and 4.16; tail, 2.63 and 3.02; tarsus, 1.06 and 1.18. The average dimensions are as follows: Length, 7.38; alar extent, 11.83; wing, 3.82; tail, 2.88; tarsus, 1.13.

Measurements of Specimens of TURDUS FUSCESCENS.

M. C. Z. No.	Collector's Number.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent	Wing.	Tail.	Tarsus.
2272	—	—	Waterville, Maine	June 2, '62	C. E. Hamlin	7.12	11.25	3.90	2.87	1.10
2275	—	—	" "	June 5, '62	"	7.39	11.90	3.79	2.84	1.18
2276	—	—	" "	June 2, '62	"	7.53	11.73	3.86	2.88	1.06
2277	—	—	" "	June 2, '62	"	7.40	12.00	3.83	2.87	1.13
9607	—	—	Canton, St. Lawrence Co., N. Y.	June —, '60	J. S. Foley	7.36	11.40	3.72	2.92	1.12
9608	—	—	" "	June —, '60	"	7.50	11.75	3.93	3.00	1.10
9609	—	—	" "	June —, '60	"	7.50	—	3.92	2.93	1.18
10382	—	—	" "	June —, '60	"	7.30	11.43	3.77	2.97	1.13
10383	—	—	" "	June —, '60	"	6.95	11.05	3.55	2.63	1.09
10384	—	—	" "	June —, '60	"	7.45	11.75	3.69	2.80	1.12
10385	—	—	" "	June —, '60	"	7.12	11.75	3.68	2.67	1.11
8830	367	♂	Newton, Mass.	May 5, '68	C. J. Maynard	7.81	12.70	4.16	3.00	—
8831	382	♂	" "	May 6, '68	"	7.75	12.55	4.10	3.00	—
8841	495	♂	" "	May 15, '68	"	7.87	11.91	4.00	3.00	—
8832	528	♂	" "	May 16, '68	"	7.70	12.45	4.00	3.00	—
8833	538	♂	Wayland, "	May 18, '68	"	7.35	11.91	3.91	2.73	—
8834	556	♂	Weston, "	May 20, '68	"	7.00	11.95	3.55	2.55	—
8835	581	♂	Newton, "	May 22, '68	"	7.41	12.50	4.15	2.90	—
8337	611	♂	" "	May 25, '68	"	7.50	12.45	3.80	2.90	—
8838	683	♂	" "	May 28, '68	"	7.50	12.30	4.15	3.17	—
8839	692	♂	" "	May 28, '68	"	7.45	12.16	3.76	2.85	—
8336	610	♂	" "	May 25, '68	"	7.30	11.33	3.55	2.60	—
8840	768	♂	" "	June 5, '68	"	7.50	12.15	3.90	2.89	—
2876	2876	—	Springfield, "	June 14, '62	J. A. Allen	7.00	11.70	3.75	2.80	1.17
1749	1749	—	" "	June 14, '62	"	7.00	11.35	3.63	2.65	1.10
1828	1828	2	" "	May 29, '62	"	7.50	11.50	3.99	2.86	1.08
1832	1832	3	" "	May 29, '62	"	7.50	11.55	3.79	2.80	1.12
—	408	—	Newton, "	May 18, '70	Wm. Brewster	7.50	12.00	3.74	2.74	1.14
2873	2873	—	Springfield "	May 12, '63	J. A. Allen	7.75	12.40	3.98	3.02	1.08
2876	2876	—	" "	May 14, '63	"	7.12	11.30	3.73	2.91	1.12
2937	2937	—	" "	May 29, '63	"	7.45	11.80	3.65	2.78	1.12
2938	2938	—	" "	May 29, '63	"	7.65	11.50	3.86	2.99	1.15
1431	—	—	Malden, "	May 22, '62	"	7.15	11.87	3.75	2.93	1.14
1432	—	—	" "	May 22, '62	"	7.52	11.80	3.93	3.00	1.15
280	—	—	" "	—	D. Higgins	7.35	11.75	3.76	2.89	1.10
281	—	—	" "	—	"	7.25	11.90	3.68	2.73	1.15
143	—	—	" "	June —, '61	"	7.30	11.72	3.72	2.83	1.12
144	—	—	" "	June —, '61	"	7.25	11.75	3.78	2.70	1.15
145	—	—	" "	June —, '61	"	7.00	11.40	3.63	2.68	1.11
146	—	—	" "	June —, '61	"	7.45	11.75	3.69	2.80	1.12

5.* *Harporhynchus rufus* Cabanis. BROWN THRUSH.

Very abundant. The specimens examined were smaller and much brighter colored than any I have seen from the Northern States. Commences nesting the last week in March.

6.* *Galeoscoptes carolinensis* Cabanis. CAT-BIRD.

Abundant. Smaller and darker colored than at the North. Some

evidently remain and breed. Audubon states that none breed so far south as South Carolina, and that few remain so far north as Florida in winter; but Dr. Coues, in his "Synopsis of the Birds of South Carolina,"* gives it as abundant and resident in that State.

7.* *Mimus polyglottus* Boie. MOCKING-BIRD.

Common. Contrary to my anticipations, I failed to hear this bird sing during my three months' stay in Florida, except in a few instances near Jacksonville early in April, at which time they were nesting, although everywhere more or less common. It was more frequent along the borders of the forest and about clumps of bushes in the pine barrens than in the hummocks. It differed from its relatives, the brown thrush and cat-bird, in avoiding the denser thickets, which are the favorite resorts of the latter. The resemblance of the mocking-bird to the loggerhead shrike, in mode of flight and general appearance, which must strike every observer, has been properly referred to by Dr. Coues.†

Different specimens of the mocking-bird from Florida differ considerably from each other in intensity of color, some being much darker than others, and in the extent of the white on the outer tail feathers, and also in the length, thickness, and curvature of the bill. Some have the commissure but slightly curved and the tip of the bill moderately depressed; others have the commissure much arched and the tip much decurved. Several specimens before me from Cape Florida are smaller than those from the St. John's River, with longer, slenderer, and more curved bills. There seems to be as much difference between specimens from South Florida and the Middle States, as between the numerous so-called species of the West Indies, which, many of them at least, are scarcely more than local forms of the original or first-described *M. polyglottus*.

The following measurements of forty-four Florida specimens of this species indicates its usual range of variation in size and proportions. The extremes of this series are as follows: Length, 9.25 and 11.00; alar extent, 13.00 and 14.75; wing, 4.00 and 4.75; tail, 4.10 and 5.15. The average dimensions are as follows: Length, 9.91; alar extent, 13.69; wing, 4.28; tail, 4.87.

* Proc. Bost. Soc. Nat. Hist., Vol. XII, p. 113.

† "Synopsis of the Birds of South Carolina," Proc. Bost. Soc. Nat. Hist., Vol. XII, p. 113, October, 1868.

Measurements of Florida Specimens of MIMUS POLYGLOTTUS.

M. C. Z. No.	Original No.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent	Wing.	Tail.
5118	—	—	Jacksonville, Fla.	Jan. 19, '68	J. A. Allen	9.75	14.00	4.60	
5124	—	—	"	Jan. 21, '68	"	9.75	14.35	4.50	
5350	—	—	Enterprise, "	Nov. 4, '69	"	10.00	14.15	4.25	
—	—	—	Hawkinsville, "	Mar. 15, '69	"	10.60	14.75	4.50	
5415	—	—	"	Mar. 15, '69	"	9.85	14.00	4.40	
5395	—	—	"	Mar. 14, '69	"	9.85	14.00	4.30	
5355	—	—	"	Mar. 10, '69	"	10.12	13.00	4.26	
5185	—	—	Hibernia, "	Jan. 30, '69	"	10.30	14.15	4.30	
10586	1955	—	Jacksonville, "	Dec. 31, '68	C. J. Maynard	10.00	13.00	4.05	4.65
—	2005	—	"	Jan. 10, '69	"	10.20	14.00	4.50	4.70
—	2407	—	Dummitt's, "	Mar. 2, '69	"	11.00	14.65	4.50	4.60
—	2614	—	"	Mar. 13, '69	"	10.15	14.00	4.40	4.90
10589	2341	—	"	Feb. 16, '69	"	9.75	13.50	4.25	4.75
—	2449	—	"	Feb. 24, '69	"	9.25	14.00	4.35	4.35
—	2613	—	"	Mar. 13, '69	"	10.20	14.00	4.40	4.75
—	2374	—	"	Mar. 13, '69	"	9.80	14.00	4.30	4.10
—	2370	—	"	Feb. 17, '69	"	9.50	13.25	4.25	4.40
10592	2372	—	"	Feb. 17, '69	"	9.75	13.75	4.20	4.50
—	2528	—	"	Mar. 5, '69	"	10.30	14.00	4.45	4.95
—	2474	—	"	Feb. 16, '69	"	10.15	14.30	4.60	5.15
10595	2486	—	"	Feb. 16, '69	"	10.00	13.00	4.00	4.38
10596	2518	—	"	Mar. 2, '69	"	9.75	13.50	4.75	4.50
—	2428	—	"	Mar. 2, '69	"	10.00	13.50	4.50	4.45
—	2429	—	"	Mar. 2, '69	"	10.00	13.50	4.35	4.35
—	2478	—	"	Mar. 11, '69	"	10.50	13.50	4.18	4.75
—	2419	—	"	Mar. 11, '69	"	10.00	13.35	4.25	4.75
10590	2310	—	"	Mar. 13, '69	"	9.60	14.35	4.40	4.50
10594	2345	—	"	Mar. 16, '69	"	9.75	13.60	4.25	4.35
10597	2569	—	"	Mar. 11, '69	"	9.30	13.00	4.00	4.17
—	2507	—	"	Mar. 11, '69	"	9.60	13.35	4.10	4.60
10587	2339	—	"	Feb. 10, '69	"	9.40	13.50	4.05	4.20
10588	2333	—	"	Feb. 17, '69	"	9.75	13.50	3.40	4.50
10591	2509	—	"	Mar. 13, '69	"	9.50	13.25	4.20	4.30
—	2560	—	"	Mar. 8, '69	"	9.50	13.20	4.15	4.35
10593	2375	—	"	Feb. 17, '69	"	9.50	13.00	4.10	4.60
—	2478	—	"	Feb. 16, '69	"	9.50	13.50	4.40	4.20
—	2485	—	"	Feb. 16, '69	"	10.00	13.00	4.00	4.38

SAXICOLIDÆ.**8.* *Sialia sialis* Haldemann.* BLUE-BIRD.**

Common. In this species the smaller size of the Florida specimens, as compared with those from Massachusetts, is very marked, as is also the greater intensity of color.

SYLVIADÆ.**9.† *Regulus calendula* Lichtenstein. RUBY-CROWNED KINGLET.**

Abundant. One of the most numerous of the winter birds. Chiefly confined to the swamps and hummocks.

10.† *Regulus satrapa* Lichtenstein. GOLDEN-CRESTED KINGLET.

Not common. A single pair was collected by Mr. Maynard at Jacksonville in January.

* *Sialia siolis* HALDEMANN, Trego's Geography of Pennsylvania, p. 77, 1843.
—BAIRD, Birds of N. Am., 222, 1858. See American Naturalist, Vol. III, p. 159, 1869.

11.* *Polioptila cærulea* *Sclater*. BLUE-GRAY GNATCATCHER.

Common. Generally seen in the same situations as *R. calendula*.

PARIDÆ.

12.* *Lophophanes bicolor* *Bonaparte*. CRESTED TITMOUSE.

Common.

13.* *Parus atricapillus* *Linne*. BLACK-CAPPED TITMOUSE. CHICKADEE.

Parus atricapillus LINNE, Syst. Nat., I, 341, 1766. — WILSON, Am. Orn., I, 137, 1808. — BONAPARTE, Obs. Nom. Wils. Orn., Journ. Phil. Acad. Nat. Sci., IV, 254, 1825. — RICH. & SWAIN., Faun. Bor. Am., II, 226, 1831. — AUDUBON, Birds Am., II, 146, pl. cxxvi, 1841. — CASSIN, Ill. Birds Cal. I, 17, 1853. — BAIRD, Birds N. Am., 390, 1858. — SCLATER, Cat. Am. Birds, 13, 1862. — BAIRD, Rev. Am. Birds, I, 80, 1864.

Parus palustris NUTTALL, Man. Orn., 241, 1832.

Parus carolinensis AUDUBON, Orn. Biog., II, 341, 1837; V, 474, pl. clx, 1839. — AUDUBON, Birds Am., II, 152, pl. cxxvii, 1841. — CASSIN, Ill. Birds Cal., I, 17, 1853. — BAIRD, Birds N. Am., 392, 1858. — SCLATER, Cat. Am. Birds, 14, 1862. — BAIRD, Rev. Am. Birds, I, 81, 1864.

Parus septentrionalis HARRIS, Proc. Phil. Acad. Nat. Sci., II, 300, 1845. — CASSIN, Ill. Birds Cal., I, 17, 80, pl. xiv, 1853. — BAIRD, Birds N. Am., 389. — SCLATER, Cat. Am. Birds, 14. — BAIRD, Rev. Am. Birds, I, 82.

Parus meridionalis SCLATER, Proc. Lond. Zool. Soc., 1856, 293. — BAIRD, Birds N. Am., 392. — SCLATER, Cat. Am. Birds, 14. — BAIRD, Rev. Am. Birds, I, 81.

Parus occidentalis BAIRD, Birds N. Am., 391, 1858. — SCLATER, Cat. Am. Birds, 14, 1862. — BAIRD, Rev. Am. Birds, I, 81, 1864.

Pacila atricapilla BONAP., Consp. Av., 230, 1850.

Pacila carolinensis BONAP., Ibid.

Seen by Mr. Marcy at Jacksonville, where also specimens of it were collected by Mr. Maynard. Not observed by any of us up the river. Audubon speaks of having found it abundant in the Floridas in the winter of 1831 and 1832, and "breeding in the swamps as early as the middle of February."*

The common titmouse (*P. atricapillus*), although not more subject to geographical variation than many other birds, is one of the species in which such differences were first detected, though not recognized at the time as such. Audubon, in 1833, upon returning to Charleston, South Carolina, from a visit to the Eastern States, the British Provinces, and Labrador, noticed a considerable difference in size between the examples of this bird he met with at the North, and those of the lowlands of the

* Birds of America, Vol. II, p. 153.

Carolinas. Though no other difference was appreciable, he and his friend Bachman thought this was sufficient to warrant the description of the southern form as specifically distinct from the northern. He accordingly thus separated them in the second volume of his "Ornithological Biography." But if the black-capped titmice of the Carolinas, the lower parts of Virginia, Maryland, and Southern New Jersey are distinct from those of Massachusetts, on precisely the same grounds are those of Massachusetts distinct from those of Northern Maine. Even the titmice of Massachusetts are not just the same in winter that they are in summer, those which breed here doubtless mainly going south in winter, while their place is filled by others that spend the summer more to the northward. This at least is what the slight average difference in size between summer and winter specimens seems to indicate. But the Carolina titmouse (*P. carolinensis*) has been recognized as valid by most subsequent writers, and in accordance with the principle upon which this supposed species was admitted, several others have been added by other authors.

The titmice from the middle, elevated regions of the continent, in accordance with a general law of geographical variation among both birds and mammals, are a little larger than those of either the Mississippi valley or the Pacific coast, and have also, apparently, a relatively slightly longer tail and paler colors, — variations which occur in a number of other birds that have a similar distribution. The titmice of this region form the *Parus septentrionalis* of authors. Specimens labelled "*Parus septentrionalis*," collected near Chicago, have been received at the Museum of Comparative Zoölogy from the Chicago Academy. They do not differ, however, from numerous others collected in Massachusetts, though the true *P. septentrionalis*, or the black-capped titmice of the Rocky Mountains, does have a slightly longer tail than those from the other parts of the continent.

Those which occur on the Pacific slope of the continent, though forming the *P. occidentalis* of authors, are admittedly the same in size and general appearance as the *P. atricapillus* of the Atlantic States, this species having been introduced to the world with the following suggestive remarks: "It is rather a hazardous undertaking to add another to the list of North American black-capped and throated titmice; but if we have three good species now, instead of one, then the present is equally entitled to specific distinction with *carolinensis* and *septentrionalis*."

The *P. meridionalis* was first made known from a single specimen from Mexico, and of which very few specimens seem to have been recognized as belonging to it. The original type certainly recalls only a worn summer specimen of the common titmouse, though its darker color^o may be due to

its southern habitat. Towards the end of the breeding season specimens of *P. atricapillus*, more especially females, have the plumage, particularly that of the lower surface of the body, much darker than in fall and winter, simply from the wearing off of the rufous and ashy extremities of the feathers, July specimens generally differing much in color from winter ones.

In respect to *P. carolinensis*, as already observed, the only difference urged as distinguishing it from *P. atricapillus* is that of its smaller size. Yet this difference is so slight that it is admitted that if *P. carolinensis* and *P. atricapillus* were "separated by a wide interval of locality, it might be a question whether it [*P. carolinensis*] might not be a variety. As, however," it is urged, "both are found together in the Middle States, and preserving together their characteristics, there will be little risk in considering them distinct." Since the larger birds are, in the main, either northern or occupy the elevated regions of the Alleghanies, the two forms must necessarily be found associated together, especially in winter, through their migrations. Unfortunately, in the work where this group has been most elaborately considered,* but two examples of each are cited, with a statement of their measurements; the two of *P. atricapillus* being from Carlisle, Pennsylvania, and the two of *P. carolinensis* from Washington, D. C. From the annexed table of measurements of *P. atricapillus* from Massachusetts and Maine, it will be seen that a few are small enough to be regarded as belonging to the *P. carolinensis*. There is, also, a larger amount of seasonal difference in the color and general character of the plumage than has been either admitted or suspected, as well as in size. No one who has previously written on this group appears yet to have compared many specimens of these supposed two species, or to have examined a sufficiently large number of either to become aware of the wide differences that exist between specimens from the same locality.

Variations similar to those assumed to specifically distinguish *P. carolinensis* from *P. atricapillus* occur in *P. hudsonicus* between specimens from localities quite distant in latitude. Dr. Bryant has already called attention to such differences in the *P. hudsonicus*, and at the same time proposed for the southern "variety" the name of "*P. hudsonicus* var. *littoralis*." Concerning this variety and the general subject in question, he remarks as follows: "The specimens of *Parus hudsonicus* from Yarmouth [Nova Scotia] and those from the Hudson Bay territory present as great, if not greater, differences in size than exist between *P. carolinensis* and *P. atricapillus*, and in color, between *P. septentrionalis* and *P. atricapillus*. I am inclined myself to consider *P. atricapillus*, *septentrionalis*, *meridionalis*, and *occidentalis* as varieties of one species; but, if they are considered as specifically distinct, there can be little question of the propriety of

* Baird's Birds of North America.

separating the Yarmouth bird from those found in the Hudson Bay territory."*

In the following table of measurements of twenty-seven specimens, all taken within ten miles of Cambridge, and all but two in December and January, the extremes of size are as follows: Length, 4.70 and 5.75, both specimens being females; alar extent, 7.50 and 8.60, both specimens being also females; wing, 2.33 and 2.63, also both females; tail, 2.15 (female) and 2.67 (male); tarsus, .62 (male) and .77 (female). The average size of these specimens is as follows: Length, 5.38; alar extent, 8.37; wing, 2.47; tail, 2.50; tarsus, .70. The females average a little smaller than the males, but the difference is only slight.

The largest specimen of the group of black-capped and black-throated titmice cited by Professor Baird† measures as follows: Length, 5.75; alar extent, 8.37; wing, 2.75; tail, 2.86 (*Parus septentrionalis*, from the Black Hills, Neb., Sm. Inst. No. 8827). A specimen of the *P. carolinensis*, cited by the same author, measures as follows: Length, 4.62; alar extent, 7.00; wing, 2.50; tarsus, .60 (Sm. Inst. No. 706, from Washington, D. C.). So far as the length of the wing and tail are concerned, specimens are fre-

Measurements of Massachusetts Specimens of PARUS ATRICAPILLUS.

M. C. Z. No.	Collector's Number.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.	Tarsus.
11703	52	♂	Cambridge, Mass.	Dec. 10, '69	Wm. Brewster	5.38	8.12	2.62	2.67	.75
11704	86	♂	"	Dec. 14, '69	"	5.25	8.12	2.55	2.37	.75
11705	87	♂	"	Dec. 14, '69	"	5.06	7.88	2.62	2.50	.62
11708	96	♂	"	Dec. 14, '69	"	5.00	7.50	2.44	2.30	.75
11707	94	♂	"	Dec. 17, '69	"	5.06	7.87	2.50	2.50	—
—	99	♂	"	Dec. 17, '69	"	5.12	8.12	2.55	2.43	.72
—	104	♂	"	Dec. 20, '69	"	5.25	8.00	2.55	2.50	.70
11711	103	♂	"	Dec. 20, '69	"	5.06	8.00	2.62	2.43	.75
11710	101	♂	"	Dec. 20, '69	"	5.12	8.06	2.55	2.43	.75
—	153	♂	"	Jan. 7, '70	"	4.94	7.80	2.43	2.25	.69
—	112	♂	"	Dec. 24, '69	"	5.50	8.12	2.62	2.56	.69
—	206	♂	"	Jan. 20, '70	"	5.40	8.21	2.58	2.55	.70
11706	88	♂	"	Dec. 14, '69	"	4.94	7.50	2.43	2.31	—
—	95	♂	"	Dec. 17, '69	"	5.00	8.00	2.50	2.37	.75
11709	97	♂	"	Dec. 17, '69	"	5.06	7.83	2.43	2.42	—
11712	105	♂	"	Dec. 20, '69	"	5.19	7.75	2.50	2.25	.67
—	100	♂	"	Dec. 20, '69	"	5.19	8.12	2.56	2.50	.75
11713	114	♂	"	Dec. 24, '69	"	5.75	7.88	2.50	2.33	.63
—	161	♂	Watertown, "	Jan. 7, '70	"	5.45	8.00	2.63	2.64	.69
—	163	♂	"	Jan. 7, '70	"	4.94	7.50	2.33	2.25	.69
—	179	♂	Belmont, "	Jan. 13, '70	"	4.84	7.55	2.35	2.15	.77
—	205	♂	"	Jan. 20, '70	"	5.41	8.17	2.54	2.42	.70
—	239	♂	"	Jan. 26, '70	"	5.25	7.52	2.35	2.30	.67
—	203	♂	Arlington, "	Jan. 20, '70	"	4.70	7.75	2.45	2.40	.68
4946	268	♂	Newton, "	Apr. 21, '68	C. J. Maynard	5.00	8.60	2.58	2.30	—
5011	1216	♂	"	Sept. 8, '68	"	5.35	7.85	2.50	2.40	.65
4945	269	♂	"	Apr. 21, '69	"	5.00	8.00	2.41	2.35	—

* Proc. Bost. Soc. Nat. Hist., Vol. IX, p. 368, April, 1865.

† Birds of North America, p. 390.

quently taken in Massachusetts (and of which I have measurements before me) that are considerably smaller than this one from Washington, or than any given in the above table.

SITTIDÆ.

- 14.* *Sitta carolinensis* Gmelin. WHITE-BREASTED NUTHATCH.

Common; especially in the pineries.

- 15.* *Sitta pusilla* Latham. BROWN-HEADED NUTHATCH.

Common in the pineries; rarely seen elsewhere.

TROGLODYTIDÆ.

- 16.* *Troglodytes aëdon* Vieillot. COMMON WREN.

Troglodytes aëdon VIEILLOT, Ois. Am. Sept., II, 52, pl. cvii, 1807. — BONA-PARTE, RICHARDSON & SWAINSON, AUDUBON. — BAIRD, Birds N. Am., 367, 1858. — IBID., Rev. Am. Birds, I, 138, 1864. — MAYNARD, Naturalist's Guide, Part II, p. 95, 1870.

Troglodytes fulvus, NUTTALL, Man. Am. Orn., I, 422, 1832.

Troglodytes americanus AUDUBON, Orn. Biog., II, 452, pl. clxxix, 1834. — BAIRD, Birds N. Am., 368. — IBID., Rev. Am. Birds, I, 141.

Troglodytes Parkmani AUDUBON, Orn. Biog., V, 310, 1839. — BAIRD, Birds N. Am., 367. — IBID., Rev. Am. Birds, I, 140.

Troglodytes sylvestris GAMBEL, Proc. Phil. Acad. Nat. Sci., III, 113, 1864.

Sylvia domestica WILSON, Am. Orn., I, 129, pl. viii, fig. 3, 1808.

Abundant, occurring everywhere. It keeps so closely concealed that it is difficult to shoot, except when on the wing. Both this and the Carolina wren are exceedingly quick in their movements, and if they are watching the collector when he is about to shoot at them, they are pretty sure to dodge the charge; although he finds the bushes and foliage where the bird sat riddled by the shot, he usually searches in vain for the specimen he is sure he ought to have killed. When approached in old grassy fields or pine openings, they will allow one to almost tread on them before attempting to get away, and then, instead of taking to wing, often seek to escape by running off like a mouse beneath the grass. The term "house" wren, usually applied to this bird, is decidedly a misnomer, since it frequents the fields the thickets, and even the forest, as much as the vicinity of houses. In the wilds of Florida, where human habitations are few, there is nothing whatever in its habits to suggest this name.

The "wood wren," *Troglodytes americanus* of Audubon, I am sure is only the brighter colored form of *T. ædon*; in size or proportions there is nothing, though the contrary has been claimed, to distinguish them. Specimens equally large and equally small occur in each state of plumage, in which the same general range of variation in proportions is presented. There is also an intergradation in color, and no observable difference in habits. Both forms were common in Florida; both also occur in New England, whence Audubon obtained the first specimen of his supposed new species. Audubon admits that it "can hardly be distinguished in description" from the house wren. The large size assumed by him as characterizing it may be readily accounted for by the fact of his obtaining his first specimens at Eastport in Maine, which is the extreme northern limit of the habitat of this species.

The following measurements of fifteen Florida specimens indicates the usual range of variation in respect to size and proportions found in specimens from the same locality. The extremes of this series are as follows: Length, 4.30 and 5.10, both specimens being females; alar extent, 6.10 and 6.95, both specimens being males; wing, 1.90 and 2.44; tail, 1.30 and 2.40; tarsus, .50 and .68; bill, .47 and .60 (.80?). The differences between these extremes, it will be noticed, are very great, considering the small size of the bird. The average dimensions are as follows: Length, 4.89; alar extent, 6.61; wing, 2.05; tail, 1.80; tarsus, .52.

Measurements of Florida Specimens of TROGLODYTES AËDON.

M. C. Z. No.	Collector's Number.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.	Tarsus.	Bill.
10681	1900	♂	Jacksonville	Jan. 1, '69	C. J. Maynard	4.70	6.60	2.44	2.40	.50	.47
—	1942	♂	"	Jan. 1, '69	"	5.00	6.50	2.00	1.70	.57	.50
—	1966	♂	"	Jan. 3, '69	"	5.00	6.75	2.05	1.75	.65	.50
10682	1967	♂	"	Jan. 3, '69	"	4.75	6.75	2.05	1.95	.55	.52
—	1968	♂	"	Jan. 3, '69	"	4.50	6.50	2.05	1.65	.61	.50
—	2790	♂	"	Mar. 20, '69	"	5.65	6.95	2.00	1.64	.54	.51
—	2576	♂	Dummitt's	Mar. 10, '69	"	5.00	6.50	2.10	2.00	.62	.50
—	4	♂	Jacksonville	Mar. 29, '69	"	4.60	6.10	2.00	1.80	.61	.60
—	2033	♂	"	Jan. 5, '69	"	5.70	6.75	2.10	1.75	.60	.80
—	1979	♂	"	"	"	4.30	6.50	2.00	1.30	.60	.56
—	2588	♂	Dummitt's	Mar. 11, '69	"	5.00	6.70	1.90	1.70	.65	.50
5178	—	—	Hibernia	Jan. 20, '69	J. A. Allen	5.20	6.75	2.03	1.70	.65	.60
5179	—	—	"	Jan. 20, '69	"	4.75	6.50	2.00	1.65	.67	.55
—	—	—	Hawkinsville	Mar. 10, '69	"	5.00	6.50	2.00	—	—	—
5361	—	—	"	Mar. 10, '69	"	4.87	6.75	2.00	—	.68	—

17.* *Thryothorus ludovicianus* Bonaparte. CAROLINA WREN.

Common. Rarely seen outside of thickets.

In few species is the difference in color between northern and southern specimens greater than in this. Florida specimens have the reddish-brown

of the dorsal surface many shades deeper than Maryland ones, and the under surface strongly rufous. The tail and wings, besides being much darker, have the dark bars black, they being deep black on the tail, and consequently far more conspicuous. The crissum, however, is lighter than in the Maryland specimens, with the black bars broader. The Florida specimens have also a much longer bill, they closely agreeing in every particular with the so-called *Thryothorus Berlandieri* of Northeastern Mexico, the Florida specimens even possessing the interrupted black bars on the sides of the body said to occasionally characterize that species as distinguished from the *T. ludovicianus*. The differences between Florida and Maryland specimens of *T. ludovicianus* in the length of the bill, as well as in color, are very striking. They are paralleled, however, in *Harporhynchus rufus* and in other species. The *T. Berlandieri* hence appears to be only the smaller, darker form of *T. ludovicianus*, — the Mexican homologue of the Florida representatives of this species.

The *Thryothorus Bewickii*, from what is known of its range, doubtless occurs as a resident bird in Florida, but is probably rare there, as it generally is elsewhere.

18.† *Anorthoura hyemalis* Rennie. WINTER WREN.

Rare. — Boardman.

19.† *Cistothorus stellaris* Cabanis. SHORT-BILLED MARSH WREN.

Rare. Enterprise, February. — Boardman.

The *Telmatodytes palustris* doubtless also occurs as a winter resident.

MOTACILLIDÆ.

20.† *Anthus ludovicianus* Lichtenstein. TITLARK.

Common. Several were usually seen in company, but along the river I saw no large flocks. According to Mr. Maynard, however, they occurred in large flocks in the "old fields" away from the river.

SYLVICOLIDÆ.

21.† *Mniotilta varia* Vieillot. BLACK AND WHITE CREEPER.

Not uncommon throughout the winter, but much more numerous in March.

22.† *Parula americana* Bonaparte. BLUE YELLOW-BACKED WARBLER.

Occasional during the winter months, but very numerous after the 1st of March, soon after which time they were in full song.

23.† *Helminthophaga celata* Baird. ORANGE-CROWNED WARBLER.

"Enterprise, 15th of February. Rare." — Boardman.

24.* *Dendræca pinus* Baird. PINE WARBLER.

Abundant. Is much on the ground at this season, as it sometimes is at the north in spring; on the whole, however, it is much less terrestrial in its habits than is *D. palmarum*. In full song in February.

25.† *Dendræca palmarum* Baird. YELLOW REDPOLLS WARBLER.

Extremely abundant. Probably the most numerous of the winter birds in East Florida, where it is more or less common in all situations. Exceedingly terrestrial in its habits, being generally seen hopping along the ground or fallen timber. At the 1st of April they had considerably decreased in numbers, but many were at that time observed at Jacksonville.

There is some indication that the males and females, and possibly the adult and young, frequent separate districts at this season. When at Jacksonville in January I saw only males; on the Upper St. John's, in February and March, only females or immature males; but these were in excessive abundance, as were also the males at the earlier date around Jacksonville. Is it not probable that the old males either do not go quite so far south as the females and immature males, or that the species was already on its way north? As is well known, the males in the species of this family, as probably in most other birds, precede the females in their journey northward.

26.† *Dendræca coronata* Gray. YELLOW-CROWNED WARBLER.

More or less common till the 1st of April, and probably some remained still later. During the last half of March they began to moult, but at the end of the month a large part were still in winter dress. The same remarks in respect to moulting apply also to *D. palmarum*.

27.* *Dendræca dominica* Baird. YELLOW-THROATED WARBLER.

Seen at Jacksonville in January, but much more abundantly up the river in February and March. March 5th I found them in great numbers in the cypress and maple swamps near Lake Munroe, at which time the spring migration had commenced.

28.* *Dendræca discolor* Baird. PRAIRIE WARBLER.

Abundant at Jacksonville, April 1st, and occasionally seen at earlier dates. This species is undoubtedly resident in Florida the whole year.

29.† *Seiurus aurocapillus* Swainson. GOLDEN-CROWNED WAGTAIL.

Not common. A few were seen in February, as well as later.

30.† *Seiurus noveboracensis* Nuttall. WATER WAGTAIL.

Rare. Found at Dummitt's by Mr. Maynard in February.

31.* *Geothlypis trichas* Cabanis. MARYLAND YELLOW-THROAT.

Abundant. Though somewhat brighter colored throughout, they differ mainly from the northern type in the greater breadth of the black facial band. There is but little difference in general size, that is, so far as I have had an opportunity of observing; occasionally a Florida example has a bill considerably longer than the average in northern examples, but this does not appear to be a very constant difference between the southern and northern specimens. It would probably be more marked in specimens from South Florida.

Other species of this family were seen in March that are not to be reckoned as winter residents. Among them are the following: *Dendræca maculosa*, *D. virens*, and *D. pennsylvanica*, *Euthlypis canadensis*, *Setophaga ruticilla*, and *Helminthophaga ruficapilla*, all of which began to appear on the Upper St. John's, near Enterprise, about the middle of March, and most of them were also seen later at lower points on the river. *Helmitherus vermivorus* and *H. Swainsoni* were taken at St. Augustine, by Mr. L. L. Thaxter, in April.

HIRUNDINIDÆ.

32.† *Tachycineta bicolor* Cabanis. WHITE-BELLIED SWALLOW.

More or less numerous, but observed at irregular intervals. Large flocks were seen near the St. John's River in January. It probably does not breed in Florida.

33.† *Cotyle riparia* Boie. BANK SWALLOW.

Not observed by either Boardman, Maynard, or myself prior to the last of March, but Mr. Audubon saw it in immense flocks "in winter," first at St. Augustine, and afterwards in other parts of the State.*

The *Stelgidopteryx serripennis* was seen about Jacksonville the first week in April, and specimens of it were obtained. Several pairs were seen flying about some bluffs a few miles below the town, apparently with the intention of selecting breeding-places.

* Birds of America, Vol. I, p. 187.

VIREONIDÆ.

34.† *Lanivireo solitarius* Baird. SOLITARY VIREO.

Rather common. In full song early in March.

35.* *Vireo noveboracensis* Bonaparte. WHITE-EYED VIREO.

Common. In full song in March.

36.† *Vireosylvia olivacea* Bonaparte. RED-EYED VIREO.

"A few all winter." — Boardman. Common after the 1st of March, on the Middle St. John's.

The Yellow-throated Vireo, *Lanivireo flavifrons*, was quite common early in March, and is undoubtedly a winter resident in South Florida.

AMPELIDÆ.

37.† *Ampelis cedrorum* Baird. CEDAR BIRD.

Common. Perhaps resident.

LANIIDÆ.

38.* *Collurio ludovicianus* Baird. LOGGERHEAD SHRIKE.

Lanius ludovicianus LINNÉ, Syst. Nat., I, 184, 1766. — BONAPARTE, NUTTALL, AUDUBON. — GAMBEL, Proc. Phil. Acad. Nat. Sci., III, 200, 1847.

Lanius garrulus BARTRAM, Travels, 289, 1791 (no description).

? *Lanius ardosiaceus* VIEILLLOT, Ois. Am. Sept., I, 81, pl. li, 1807. — BONAPARTE, Obs. on Wils. Nomencl., Journ. Phil. Acad. Nat. Sci., III, 358, 1824.

Lanius carolinensis WILSON, Am. Orn., III, 57, pl. xxii, fig. 5, 1811.

Lanius excubitoroides SWAINSON, Faun. Bor. Am., II, pl. xxxiv, 1831.

Lanius elegans SWAINSON, Ibid., 122. — NUTTALL, Man. Am. Orn., I, 2d ed., 287, 1840. — GAMBEL, Proc. Phil. Acad. Nat. Sci., I, 261, 1843.

Lanius mexicanus BREHM, Cab. Journ. für Orn., II, 145, 1854. — SCLATER, Catal. Am. Birds, 46, 1861.

Collurio ludovicianus BAIRD, Birds of N. Am., 325, 1858. — ALLEN, Amer. Nat., III, 579, 1869. — BAIRD, Rev. Am. Birds, I, 443, 1866.

Collurio excubitoroides BAIRD, Birds N. Am., 337. — BAIRD, Rev. Am. Birds, I, 445. — COOPER & BAIRD, Orn. Cal., I, 138, 1870.

Collurio elegans BAIRD, Birds N. Am., 328. — BAIRD, Rev. Am. Birds, I, 444. COOPER & BAIRD, Orn. Cal., I, 140, 1870.

Not very numerous.

I have already referred to the questionable distinctness of the so-called *C. excubitoroides* from the present species.* Further examination of the

* See a series of articles in the "American Naturalist," entitled "Notes on some of the Rarer Birds of Massachusetts," Vol. III, 1869.

subject has only confirmed me in the opinion that they are not distinct, and that in all probability the *C. elegans* of California should also be referred to the *C. ludovicianus*.*

TANAGRIDÆ.

The *Pyrrhula æstiva* became common on the Lower St. John's April 1st to 5th, but was not observed previously. *P. rubra* was not seen at all.

A considerable number of specimens of this species (*P. æstiva*) in the Museum, from the Atlantic States, present great differences in the size of the bill in respect to vertical and lateral thickness, as well as in the position and distinctness of the "tooth" of the bill, and in the curvature of the commissure, as indicated by the accompanying figures (Plate IV, figs. 19, 20). They also vary greatly in intensity of color, both of the bill and plumage, as do different specimens of *P. rubra* from Massachusetts. Hence species based solely on such distinctions should be accepted, if at all, with great hesitancy.†

FRINGILLIDÆ.

39.† *Chrysomitris tristis* Bonaparte. YELLOW BIRD.

Common throughout the winter, and as numerous the first week in April as earlier.

I am sure I heard the notes of the Pine Finch (*Chrysomitris pinus*), but as I obtained no specimens of it and do not find it reported by others, I do not include it in the present list. It is not improbable that this species and the Purple Finch (*Carpodacus purpureus*) are occasional winter visitors.

* Since writing the above I have met with the following observations on this group, made by Dr. Gambel, in his "Remarks on the Birds observed in Upper California" (Proc. Phil. Acad. Nat. Sci., Vol. III, p. 200, 1847): "In the shrikes we are presented with a group of birds closely allied to each other, and undergoing such changes in plumage as renders them difficult to discriminate. Although examined with great care by Swainson in the Fauna Boreali-Americana, yet he appears to have laid too much stress upon characters subject to great variation, as size, relative length of quills, and color. . . . The relative length of quills in the shrikes is an uncertain character, and differs very much according to age. In the young of this species the second quill is generally much shorter than the sixth, but in the adult equals and may even exceed the sixth in length; the proportion of the third, fourth and fifth to each other is also exceedingly various, and indeed in each wing of the same bird it is very common to find the proportions of the quills differing very materially. This I have found to be the case in the European and both American species [*Collurio ludovicianus* and *C. borealis*]."

† See some remarks on the "Uniformly red species of *Pyrrhula*," in Proceed. Phil. Acad. Nat. Sciences, p. 127. June, 1869.

40.† *Passerculus savanna* Bonaparte. SAVANNA SPARROW.

Emberiza sandwichensis GMELIN, Syst. Nat., I, 875, 1788.

Emberiza arctica LATHAM, Ind. Orn., I, 414, 1790.

Emberiza chrysops PALLAS, Zool. Rosso-Asiat., II, 45, pl. xlvi, fig. 2, 1811.

Fringilla savanna WILSON, Am. Orn., III, 55, pl. xxii, fig. 2, 1811.

Passerculus savanna BONAPARTE, Geog. and Comp. List., 33, 1838. — BAIRD, Birds N. Am., 442, 1858. — SCLATER, Cat. Am. Birds, 112, 1862.

Passerculus alaudinus BONAPARTE, Compte Rendu, XXXVII, 918, 1853. — BAIRD, Birds N. Am., 446, 1858. — SCLATER, Cat. Am. Birds, 112, 1862. — COUES, Proc. Phil. Acad. Nat. Sci., XVIII, 84, 1866. — COUES, Proc. Essex Inst., V, 281, 1868. — COOPER & BAIRD, Orn. Cal., I, 181, 1870.

Passerculus anthinus BONAPARTE, Compte Rendu, XXXVII, 919, 1853. — BAIRD, Birds N. Am., 445, 1858. — SCLATER, Cat. Am. Birds, 112, 1862. — COOPER & BAIRD, I, 183.

Passerculus sandwichensis BAIRD, Birds N. Am., 444, 1858. — SCLATER, Cat. Am. Birds, 112, 1862. — COOPER & BAIRD, Orn. Cal., I, 180.

Abundant, especially on the savannas, where it was the principal sparrow seen.

This species, like all the sparrows, varies considerably in color with the season of the year. Fall specimens, and especially the young of the year, have the yellow superciliary stripe very indistinctly defined, it being in numerous cases entirely obsolete. The general plumage is also much browner, with the streaks on the dorsal surface suffused and obscured with ferruginous, and those below, as in fall specimens of *Melospiza melodia*, bordered with the same tint. Different individuals also vary considerably in the breeding season, some being much grayer above than others; the superciliary line varies from bright yellow to grayish white, with the yellow either entirely wanting or limited to a slight wash on the part anterior to the eye. This grayer plumage and faded condition of the superciliary stripe is more especially seen towards the end of the breeding season. The spots below also vary so much in size as to give very different aspects to the plumage of the lower surface of the body in different specimens. In some they form little more than a narrow line along the middle of the feathers of the breast and sides of the body; in others they are quite broad, occupying relatively a much larger surface; occasionally, also, they are aggregated on the lower part of the breast, forming a large conspicuous patch, as distinct as is ever seen in *Melospiza melodia*. The general size of the bird also varies considerably, as is indicated in the accompanying table of measurements, and the bill is subject to very marked variations in respect to length, size, thickness, and slenderness, as substantiated by a series of nearly one hundred specimens now before me, including some thirty specimens taken at Ipswich, Massachusetts, in the breeding season.

These specimens are separable to some extent into several series, which may be based either upon difference in general size, the character of the bill, or upon coloration; but these several kinds of variation fail to corroborate each other. If separated upon differences in size, the two or more series thus separated embrace every combination of the other differences; and similar incongruities result when the separation is made upon differences in coloration or other characters. Yet the Massachusetts specimens present among themselves differences as well marked and of the same character as is assumed to distinguish several of the so-called species from the Pacific coast, that have been proposed and adopted by different authors.

Alexander Wilson was the first naturalist who gave any adequate description of the species in question, though the *Emberiza sandwichensis* of Gmelin unmistakably refers to this bird, and this name having been given long before that of Wilson, should, in accordance with the rule of priority, supplant Wilson's more euphonious and familiar one of *savanna*. The first supposed species recognized by modern writers after the well-known one of Wilson was the *P. alaudinus*, described by Bonaparte in 1853, in his notes on the Delattre collection,* from a specimen from California. He says it is not easily distinguished from *P. savanna*, but differs from it in being smaller, with the bill shorter and slenderer, and in wanting the yellow superciliary line.† Professor Baird redescribed it in his Birds of North America in similar language, and cites under it five specimens, which came respectively from Brownsville, Texas; Tamaulipas, Mexico; Petaluma, Cal., and Shoalwater Bay, W. T. He remarks respecting it as follows: "This species, if really distinct from *P. savanna*, differs in the rather smaller size, although the difference is not great, and in the considerably paler colors. The superciliary stripe shows a very faint trace of yellow, especially anteriorly near the bill. In some specimens, as 4342, there is none at all." Bonaparte, in his paper just cited, added another "new species" from Kodiak, Alaska, which he called *Passerculus anthinus*, and described as follows: "*Passerculus anthinus*, Bp., ex Kodiak, Am. Ross. *Simillimus præcedenti, sed rostro etiam graciliore et capite flavo induto; subtus albo-rufescens magis maculatus.*" He says it is still smaller and has the bill slenderer even than the other, and that it appears to live farther north. Professor Baird also redescribes this species, and is much more explicit in his account of it. He says: "Similar to *P. savanna*, but smaller. . . . Breast and upper part of belly thickly spotted with sharply defined sagittate brown spots, exhibiting a tendency to aggregation on the

* Comptes Rendus, Tome XXXVII, p. 918.

† "*Passerculus alaudinus*, Bp., ex Wils., mais plus petite sans jaune aux sourcils et à bec plus court et plus effilé."

middle of the belly," etc. He adds: "This species is the smallest of its group, and differs from all in the much greater amount of spotting on the under parts. The streaks, indeed, extend over the whole breast and upper part of the abdomen, instead of being mainly confined to the jugulum." It differs, he says, from *P. alaudinus* "in the strong shade of yellow on the head, the much darker tints above, and the thick crowding of larger and better defined spots beneath, with a faint tinge of reddish." He refers to it three specimens from San Francisco, Benicia, and Petaluma, California.

In 1858 Professor Baird added still another species of *Passerculus* to those previously recognized, through the redescription of the original type of this group, the *Emberiza sandwichensis* of Gmelin, based upon Latham's Sandwich Bunting* and Pennant's Unalaska Bunting.† The name Sandwich, as Professor Baird has remarked, refers not to the Sandwich Islands, but to Sandwich Sound, on the northern coast. To this species Baird judiciously refers the *Emberiza arctica* of Latham‡ and Vigors,§ and the *E. chrysops* of Pallas.|| Professor Baird's description of it is as follows: "Almost exactly like *P. savanna*, but half an inch larger, with much larger bill. Length, 6.12; wing, 3.00; tail, 2.55. Habitat, north-western coast, from the Columbia River to Russian America." He also further observes: "This species is extremely similar to the *P. savanna*, and is only distinguishable by its greater size and more western locality. The tail feathers also are rather more acutely pointed. There is also a greenish-yellow shade on the top and sides of the head, brighter than is seen in *P. savanna*. The bill is considerably larger and longer, measuring .51 of an inch above instead of .44." To this is referred one specimen from "Russian America," one from Fort Steilacoom, W. T., and three from Shoalwater Bay, W. T., three of which measure as is indicated in the above-quoted description, and the other nearly three fourths of an inch less.

In respect to size, then, it appears that the so-called *P. sandwichensis* is the larger, the *P. savanna* the next in size, *P. alaudinus* the third, and *P. anthinus* the smallest. So, at least, it is claimed; but from the measurements published in Birds of North America, a female of *P. savanna* from Carlisle, Pa. (No. 780), is, with one exception (No. 4340, from Brownsville, Texas), the smallest of the specimens of this genus of which measurements are there given; two others from Pennsylvania are below the average of *P. alaudinus*. No. 10,203, from Russian America, referred to *P. sandwichensis*, is scarcely larger than an average *P. savanna*. The

* Latham's Synopsis, Vol. II, p. 202, 1783.

† Pennant's Arctic Zoology, Vol. II, Species No. 229, pp. 320, 368.

‡ Indian Ornithology, Vol. I, p. 414, 1790.

§ Zoology of the Blossom, p. 20, 1839.

|| Zoographia Rosso-Asiatica, Vol. II, p. 45, pl. xlviii, fig. 1, 1811.

accompanying series of measurements shows that specimens occur in Massachusetts as large and as small as any specimens of the genus of which measurements are given by Professor Baird.

In respect to the geographical distribution of these different supposed species, it will be observed that of the three West Coast species, the larger, *P. sandwichensis*, is northern, and the others, *P. alaudinus* and *P. anthinus*, southern, which perfectly explains the difference in size that occurs between them.* In respect to *P. alaudinus* and *P. anthinus*, one is only the paler colored and the other the brighter colored form of the common savanna sparrow as represented in the Pacific States; the three supposed species together forming a series similar to what is seen when a large number of specimens of this bird from the Atlantic States are compared. In other words, the characters whereon these species are based are evidently only individual differences. The *P. alaudinus* is the form with narrow streaks and generally paler tints, or that having a minimum intensity of color; the *P. anthinus* is that with the brighter tints, or with the maximum intensity of color, the greater breadth of the streaks, and the rufous suffusion below correlating with the generally brighter tints. Aside from this normal range of variation referred to at length in Part III as obtaining in all species, there is that of season to be taken into account, as the fading of the superciliary stripe and the grayer aspect of the plumage above towards the end of the breeding season, through the natural wearing and bleaching of the plumage,† and also the rufous suffusion and greater amount of color characteristic of the renewed plumage in fall. It will be noticed that authors report the occurrence of all the western species either actually at or near the same points,‡ while *P. savanna* was not until recently supposed to occur on the Pacific slope of the continent.§ But one of the others have been announced from the plains as far east as Nebraska,|| and from Brownsville, Texas.¶

In respect to the habits of these supposed species, there is nothing attributed to the western one that is not equally applicable to the eastern bird. Dr. Coues, it is true, says that in Southern California *P. anthinus* seemed confined to the moist salt grass and sedgy weeds of the sea-shore

* Since the above was written, Mr. Dall has given, not only *P. savanna* and *P. sandwichensis*, but also *P. alaudinus* and *P. anthinus* in his list of the birds of Alaska. (See Trans. Chicago Acad. Sciences, Vol. I, pp. 283, 284.)

† See Part III, p. 193.

‡ See Professor Baird, "Birds of North America," Dr. Coues, "Notes on the Birds of Arizona Territory," and Cooper's Ornithology of California.

§ It has recently been reported by Mr. Dall as common in Alaska.

|| *P. alaudinus*, Sclater's Catalogue of American Birds, p. 112.

¶ *P. alaudinus*, Baird, in Birds of North America, p. 446.

itself. "When with difficulty it was flushed, its flight was," he remarks, "very rapid and irregular; and it would alight again almost immediately, and run with great celerity among the roots of the thick grasses. It was then exceedingly difficult to procure."* All of which is quite true of *P. savanna* when frequenting the salt marshes, which form its most favorite resort in Massachusetts. "*P. alaudinus*," he says, "was common two or three miles away from the coast, but on the sea-shore itself I never found one mixing with *P. anthinus*; it is a bush-and-weed rather than a grass species." *P. savanna* also frequents similar localities. Mr. Dall, under *P. anthinus*, has also accurately indicated the habits of the eastern *Passerculus*. Under *P. savanna*, however, he mentions a fact in respect to the breeding habits of this species I have never before seen mentioned as characterizing any of the *Passerculi*, namely, *its nesting in bushes*. I have met with many nests of the eastern savanna sparrow, and have always found them placed on the ground, usually in a tuft of grass.

To recur again to the series in the Museum of Comparative Zoölogy, I may add that while some of the Ipswich specimens, taken late in June, have a decidedly yellow superciliary stripe, none have it so bright as it is usually in spring specimens; in a considerable proportion it is very pale, and in Nos. 4700, 10668, etc., it is grayish-white, with no perceptible trace of yellow. No. 5099, and some others, have the spots on the breast and sides very narrow, occupying but a small share of the surface; on the other hand, in No. 5088, as also in several others of the series, the spots are so broad as to occupy more space than the enclosing white portion. In other specimens, taken at a different season of the year, the "rufous tinge" surrounding the spots referred to in the above-quoted description of *P. anthinus* is very marked. There is likewise great difference in the color of the upper surface in different specimens. In some the black central spots of the interscapularies are so broad as to give to the dorsal aspect a very dark tint; in others, taken the same day at the same locality, they are so restricted that the general aspect of this surface is gray. The bills of the different specimens vary as much in length and robustness as they are represented to do in the two extremes in this respect in the western bird. Some of the long-billed ones have the bill slender; others have it thick and stout. Occasionally one has the upper mandible projecting considerably beyond the lower, but only in cases where it is abnormally developed. A specimen from Fort Bridger, Utah (No. 11115 of the Smithsonian Catalogue), in the Museum, labelled *Passerculus alaudinus* at the Smithsonian Institution, is of this character, the upper mandible being very much abnormally developed and decurved, and projecting much beyond the lower.

* Ibis, July, 1866, 268.

In short, while not denying that there may be a slight average difference between eastern and western specimens, as I know there is between those of the Central Plains and those of the Atlantic States, I cannot allow that it is at all sufficient to substantiate a specific difference. On the contrary, I am confident that the above-named supposed species of the Pacific States are based chiefly on individual variation perfectly parallel with that seen in a large series of specimens from the Atlantic States. No one, in fact, seems to have felt very confident that any of them were distinct from the eastern *P. savanna*. Dr. Coues has even repeatedly expressed his belief that *Passerculus alaudinus* is not permanently distinct from that species. "In a large series of the latter," he says, "shot about Washington, I have found fully as great differences as I have ever detected in comparing the eastern with the western forms."*

Dr. Cooper also refers as follows to the close resemblance of the *P. alaudinus* to the *P. sandwichensis*. He says, "I think it very doubtful whether these specimens (which measure larger than the dimensions given by Baird, though otherwise agreeing) are anything more than a southern form of *P. sandwichensis*, though collected near San Diego. . . . Baird considers it almost identical with *P. savanna* of the east, and says that *P. sandwichensis* differs from that species in its larger size. Spring specimens have the superciliary stripe more decidedly yellow, so that there only remains a more slender bill to distinguish this from *P. savanna*, and the larger size (characteristic of northern specimens generally), with darker hues, from *P. sandwichensis*."† Respecting *P. anthinus* Dr. Cooper remarks, "This species appears better marked, as compared with *P. savanna*, than the preceding [*P. alaudinus* and *P. sandwichensis*], although I am not entirely satisfied that it is different."‡

The following measurements of twenty-six specimens (fourteen males and twelve females), all taken at Ipswich during June and July, 1868, and measured before skinning, indicates the range of individual variation presented by this species. The extremes are as follows: Length, 5.20 and 6.00, both males; alar extent, 7.61 and 9.75, both females; wing, 2.44 and 2.95; tail, 1.64 and 2.25; tarsus, .75 and .88. The average dimensions are: Length, 5.20; alar extent, 8.79; wing, 2.70; tail, 1.96; tarsus, .84. The following are the extremes of the series of measurements of the western *Passerculi*, given in Birds of North America: Length, 5.00 (*P. alaudinus* Tamaulipas, Mex.) and 6.12 (*P. sandwichensis*, Fort Steilacoom, W. T.); alar extent, 8.50 and 9.37 (same specimens); wing, 2.50 and 2.95 (same specimens); tail, 2.00 and 2.57 (same specimens). It thus appears that

* Ibis, July, 1866, p. 289.

† Ornithology of California, Vol. I, p. 182.

‡ Ibid., p. 183.

specimens taken in the breeding season in Massachusetts, overlap in two out of the four measurements given, all the so-called western species, while specimens taken in Massachusetts at other seasons, vary still more than the specimens cited in the following table.

Measurements of Massachusetts Specimens of PASSERCULUS SAVANNA,
taken in the Breeding Season.

M. C. Z. No.	Collector's Number.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.	Tarsus.
5083	811	♂	Ipswich	June 12, '68	Allen & Maynard	5.76	8.32	2.72	2.07	—
—	820	♂	"	June 13, '68	"	5.76	9.25	2.55	2.00	.85
5086	846	♂	"	June 14, '68	"	5.65	9.15	2.75	1.85	.85
5087	848	♂	"	June 14, '68	"	5.50	9.13	2.75	2.07	.84
5089	852	♂	"	June 15, '68	"	5.40	9.10	2.65	2.00	.85
5090	853	♂	"	June 15, '68	"	5.70	9.25	2.63	2.10	.87
5094	854	♂	"	June 15, '68	"	5.70	9.25	2.64	1.95	.88
—	855	♂	"	June 15, '68	"	5.50	9.15	2.75	1.85	.84
5901	856	♂	"	June 15, '68	"	5.40	9.25	2.60	1.93	.80
5092	857	♂	"	June 15, '68	"	5.20	9.37	2.95	2.05	—
5092	858	♂	"	June 15, '68	"	5.40	9.25	2.63	2.06	.83
5096	862	♂	"	June 17, '68	"	5.75	8.00	2.90	2.00	—
5098	873	♂	"	June 17, '68	"	5.83	7.75	2.80	2.10	—
—	881	♂	"	June 17, '68	"	6.00	8.27	2.74	2.25	—
5082	810	♂	"	June 12, '68	"	5.42	8.81	2.71	1.81	—
5084	819	♂	"	June 13, '68	"	5.75	8.85	2.57	1.83	.83
5084	847	♂	"	June 14, '68	"	5.54	8.55	2.70	1.90	.80
5088	851	♂	"	June 15, '68	"	5.75	9.75	2.70	2.05	.75
5094	859	♂	"	June 17, '68	"	5.45	8.90	2.65	2.05	.85
5095	860	♂	"	June 17, '68	"	5.25	8.50	2.44	1.85	—
5096	862	♂	"	June 17, '68	"	5.70	8.90	2.70	1.90	—
5097	878	♂	"	June 17, '68	"	5.75	7.61	2.60	1.80	—
5099	877	♂	"	June 17, '68	"	5.75	8.05	2.75	2.00	—
—	890	♂	"	June 17, '68	"	5.35	9.75	2.77	1.95	—
5100	1006	♂	"	July 15, '68	C. J. Maynard	5.65	7.95	2.72	2.00	.87
—	1158	♂	"	Aug 19, '68	"	5.36	8.50	2.70	1.64	.85

41.† *Pooecetes gramineus* Baird. GRASS FINCH.

Abundant, especially in and about the old fields. The most numerous sparrow in East Florida in winter.

42.† *Junco hyemalis* Selater. SNOW BIRD.

"Common in January." — Boardman. Not seen by either Mr. Maynard or myself. Probably of somewhat irregular occurrence so far south.

43.† *Spizella socialis* Bonaparte. CHIPPING SPARROW.

Common. A large proportion of those seen were young birds.

44.* *Spizella pusilla* Bonaparte. FIELD SPARROW.

Common. More numerous than the preceding species (*S. socialis*). They appeared to be breeding at Jacksonville the first week in April.

The songs of the males were so different from those of the northern bird that the species was almost unrecognizable by me from its notes.

45.† *Zonotrichia albicollis* Swainson. WHITE-THROATED SPARROW.
Generally more or less common.

46.† *Melospiza melodia* Baird. SONG SPARROW.
Not numerous. At least comparatively few were seen.

47.† *Melospiza palustris* Baird. SWAMP SPARROW.
Common, frequenting the hummocks and swamps.

48.† *Passerella iliaca* Swainson. FOX-COLORED SPARROW.
A single specimen was seen by Mr. G. A. Boardman at Enterprise.
None were seen by Mr. Maynard or myself.

49.† *Ammodromus maritimus* Swainson. SEASIDE FINCH.
"Abundant at Fernandina." — Boardman.

50.† *Ammodromus caudacutus* Swainson. SHARP-TAILED FINCH.
"Abundant, with the preceding." — Boardman. Although I have marked as winter visitors both these species of *Ammodromus*, they may be resident.

51.† *Coturniculus Henslowi* Bonaparte. HENSLOW'S SPARROW.
Stated by Audubon to be abundant in winter on the grassy pine barrens of Florida.*

52.* *Peucæa æstivalis* Baird. PINE-WOOD SPARROW.
Fringilla æstivalis LICHTENSTEIN, Verzeich. Doubleder Zool. Mus. der königl. Univ. zu Berlin, 25, 1823.
Fringilla Bachmani AUDUBON, Orn. Biog., II, 366, pl. clxv, 1834.
Fringilla æstiva NUTTALL, Man. Orn., I, 2d ed., 568, 1846.
Peucæa Bachmani AUDUBON, Syn. Am. Birds, 112, 1839.
Peucæa æstivalis CABANIS, Mus. Hein., 132, 1850.
Zonotrichia Cassinii WOODHOUSE, Proc. Phil. Acad. Nat. Sci., 1852, 60.
Peucæa Cassinii BAIRD, Birds N. Am., 485, 1858.

Common, but confined to the pine woods.

The twenty-two specimens, collected by Mr. Maynard's party and myself, now in the Museum, present considerable differences. Several are so different in color from most of the others as to almost have the appearance of being a different species, the general color of the upper

* Birds of America, Vol. III, p. 76.

parts being rufous instead of gray. These are all females, the others being males. But the males differ greatly in color, few of our sparrows being more variable in this respect than the present species.

The following measurements of twenty-two Florida specimens indicate quite a constancy in size, much greater than in color. The extremes of this series are as follows: Length, 5.75 and 6.20; alar extent, 7.60 and 8.30; wing, 2.17 and 2.55; tail, 2.25 and 2.68. Average: Length, 5.88; alar extent, 8.99; wing, 2.40; tail, 2.49.

Measurements of Florida Specimens of PEUCÆA ÆSTIVALIS.

M. C. Z. No.	Collector's Number.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.	Tarsus.
5377	5377	♂	Hawkinsville	Mar. 13, '69	J. A. Allen	6.00	8.20	2.35	2.50	.76
5393	5393	♂	"	Mar. 15, '69	"	5.85	8.05	2.28	2.30	.75
5425	5425	♂	Jacksonville	Apr. 2, '69	"	5.90	8.00	2.40	2.40	.76
5426	5426	♂	"	Apr. 2, '69	"	5.90	8.30	2.55	2.50	.70
5427	5427	♂	"	Apr. 2, '69	"	5.60	7.85	2.50	2.25	.76
5428	5428	♂	"	Apr. 2, '69	"	5.80	8.20	2.40	2.50	.74
5429	5429	♂	"	Apr. 2, '69	"	5.90	7.85	2.45	2.47	.76
5430	5430	♂	"	Apr. 2, '69	"	5.60	7.85	2.40	2.50	.68
10616	18	♂	"	Apr. 3, '69	C. Thurston	5.80	8.00	2.45	2.50	.70
10617	21	♂	"	Apr. 3, '69	"	5.90	7.75	2.17	2.50	.67
—	24	♂	"	Apr. 5, '69	"	5.75	8.00	2.45	2.35	.66
10618	27	♂	"	Apr. 6, '69	"	5.75	8.00	2.50	2.55	.65
10619	28	♂	"	Apr. 6, '69	"	5.90	8.00	2.45	2.65	.62
10620	36	♂	"	Apr. 7, '69	"	5.90	8.00	2.25	2.55	.68
10624	72	♂	"	Apr. 24, '69	"	6.00	8.00	2.40	2.55	.70
10623	68	♂	"	Apr. 16, '69	"	6.00	8.00	2.40	2.56	.75
10622	67	♂	"	Apr. 16, '69	"	6.00	8.00	2.45	2.50	.74
10625	73	♂	"	Apr. 24, '69	"	6.20	8.15	2.40	2.50	.70
—	651	♂	"	Apr. 15, '69	"	6.00	8.00	2.40	2.53	.72
10621	45	♂	"	Apr. 18, '69	"	6.10	8.10	2.35	2.60	.73
—	37	♂	"	Apr. 7, '69	"	5.90	8.00	2.45	2.68	.75
10626	2934	♂	"	Apr. 13, '69	"	5.85	7.60	2.25	2.34	.70

53.* *Cardinalis virginianus Bonaparte.* CARDINAL BIRD.

Exceedingly numerous. Their clear, musical, loud call-note was heard everywhere, this being the most noisy bird of the forest.

None of the specimens I have seen from Florida are as large as those from the Middle States. The colors of the former are also somewhat brighter, especially in the females, in which the brownish-yellow of the lower parts is not only much deeper, but a large proportion have the breast and middle of the abdomen strongly tinged with bright red, giving a very different appearance from northern females.

The following measurements of fifty-eight specimens shows the amount of variation in size in specimens from the same locality. The females, it will be seen, average a little smaller than the males, but the sexual difference in this respect is not very great. The range of variation, which is much less in this species than in many, is as follows: In the males: Length, 7.75 to 9.10; alar extent, 11.00 to 11.78; wing, 3.50 to

3.85; tail, 3.40 to 4.20; tarsus, .62 to .80. In the females: Length, 7.50 to 8.75; alar extent, 10.70 to 11.75; wing, 3.25 to 3.85; tail, 3.40 to 4.10; tarsus, .62 to .75. Average size of the males: Length, 8.46; alar extent, 11.43; wing, 3.63; tail, 3.87. Average of the females: Length, 8.27; alar extent, 11.27; wing, 3.53; tail, 3.77.

Measurements of Florida Specimens of CARDINALIS VIRGINIANUS.

M. C. Z. No.	Collector's Number.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.
5164	5164	♂	Hibernia	Jan. 30, '69	J. A. Allen	8.60	11.45	3.55	4.10
5165	5165	♂	"	Jan. 30, '69	"	8.45	11.70	3.65	3.83
5166	5166	♂	"	Jan. 30, '69	"	8.45	11.50	3.60	4.10
5167	5167	♂	"	Jan. 30, '69	"	8.75	11.50	3.55	3.95
5189	5189	♂	"	Feb. 3, '69	"	8.60	11.25	3.50	3.95
5192	5192	♂	"	Feb. 3, '69	"	8.75	11.75	3.50	4.10
5193	5193	♂	"	Feb. 3, '69	"	8.45	11.35	3.60	3.83
5230	5230	♂	Volusia	Feb. 12, '69	"	8.15	11.00	3.30	3.78
5311	5311	♂	Enterprise	Mar. 1, '69	"	8.75	11.60	3.60	3.90
5312	5312	♂	"	Mar. 1, '69	"	8.50	11.35	3.40	3.78
5347	5347	♂	"	Mar. 4, '69	"	9.10	11.50	3.58	4.15
—	—	♂	Hawkinsville	Mar. 13, '69	"	7.75	10.70	3.25	—
—	—	♂	Jacksonville	Mar. 31, '69	"	7.75	11.15	3.55	—
5424	5424	♂	"	Apr. 2, '69	"	8.50	11.50	3.65	3.90
—	—	♂	"	Apr. 2, '69	"	8.55	11.10	3.55	—
—	1955	♂	"	Jan. 2, '69	C. J. Maynard	9.00	11.50	3.65	4.30
—	1987	♂	"	Jan. 5, '69	"	9.00	11.50	3.85	4.05
10706	1988	♂	"	Jan. 5, '69	"	8.50	11.51	3.75	4.20
10707	1989	♂	"	Jan. 5, '69	"	8.50	11.45	3.75	3.80
—	2003	♂	"	Jan. 10, '69	"	8.05	11.60	3.75	4.15
—	2041	♂	"	Jan. 7, '69	"	8.00	11.00	3.75	4.00
—	2430	♂	Dummitt's	Feb. 24, '69	"	8.00	11.25	3.50	3.40
—	2418	♂	"	Feb. 22, '69	"	8.75	11.10	3.80	3.90
—	2533	♂	"	Feb. 7, '69	"	8.60	11.00	3.65	4.10
—	2537	♂	"	Feb. 9, '69	"	8.70	11.50	3.65	4.00
—	2447	♂	"	Feb. 24, '69	"	8.00	11.56	3.60	3.60
10709	2337	♂	"	Feb. 16, '69	"	8.50	11.50	3.60	3.80
—	2328	♂	"	Feb. 25, '69	"	8.25	11.50	3.50	3.80
10710	2338	♂	"	Feb. 16, '69	"	8.50	11.50	3.60	3.80
10713	2398	♂	"	Feb. 16, '69	"	8.50	11.50	3.65	3.65
—	2324	♂	"	Feb. 25, '69	"	8.60	11.50	3.60	3.50
—	2339	♂	"	Feb. 25, '69	"	8.75	11.78	3.65	3.90
—	2368	♂	"	Feb. 25, '69	"	8.40	11.50	3.60	4.10
—	2335	♂	"	Feb. 17, '69	"	8.00	11.50	3.60	3.90
—	2364	♂	"	Feb. 17, '69	"	8.50	11.50	3.60	3.60
—	2363	♂	"	Feb. 17, '69	"	8.50	11.50	3.60	4.00
—	2538	♂	"	Mar. 10, '69	"	8.15	11.18	3.60	3.40
—	2535	♂	"	Mar. 4, '69	"	8.40	11.00	3.57	3.95
—	2459	♂	"	Feb. 25, '69	"	8.50	11.50	3.75	3.95
—	2008	♂	Jacksonville	Jan. 9, '69	"	8.25	11.25	3.85	3.85
—	2042	♂	"	Jan. 6, '69	"	8.75	11.24	3.50	3.70
—	2334	♂	"	Jan. 11, '69	"	8.00	11.00	3.40	3.40
—	2579	♂	Dummitt's	Mar. 10, '69	"	7.75	10.75	3.40	3.95
—	2337	♂	"	Feb. 24, '69	"	7.50	11.00	3.30	3.50
—	2594	♂	"	Feb. 11, '69	"	8.50	11.10	3.50	3.91
—	2595	♂	"	Feb. 11, '69	"	8.50	11.05	3.50	3.85
—	2334	♂	"	Feb. 24, '69	"	8.75	11.75	3.67	4.00
—	2415	♂	"	Feb. 20, '69	"	8.00	11.10	3.50	3.50
—	2394	♂	"	Feb. 16, '69	"	8.00	11.75	3.30	3.55
10716	2324	♂	"	Feb. 15, '69	"	8.50	11.50	3.70	4.10
—	2458	♂	"	Feb. 25, '69	"	8.10	11.15	3.75	3.65
—	2336	♂	"	Feb. 17, '69	"	8.50	11.50	3.60	3.70
—	2474	♂	"	Feb. 16, '69	"	8.00	11.25	3.45	3.55
—	2475	♂	"	Feb. 26, '69	"	8.00	11.00	3.40	3.75
10117	2489	♂	"	Mar. 1, '69	"	8.05	11.15	3.64	3.40
10715	2488	♂	"	Mar. 1, '69	"	8.50	11.20	3.60	4.00
10714	2427	♂	"	Mar. 4, '69	"	8.20	11.25	3.55	3.60
—	2043	♂	"	Feb. 11, '69	"	8.50	11.50	3.75	3.90

54.* *Pipilo erythrophthalmus* Vieillot. CHEWINK.

Exceedingly numerous.

Mr. C. J. Maynard detected an interesting local race or variety of this bird at Dummitt's. Besides having the irides *yellowish-white* instead of *red*, there is less white at the base of the primaries, less skirting the secondaries, and much less on the tail. The whole bird is also smaller. The white on the tail generally extends only to the three outer pairs of feathers; in the common northern form it extends over the four outer pairs, and on the first is much more extended than in the Florida one. The tail of the common form, with the outer pair of feathers removed, would resemble, in respect to the distribution and extent of the white, that of the Florida bird. The song of this bird, as I heard it at Jacksonville in April, is quite different from that of the northern bird, it being ordinarily only about half as long, and uttered with much less spirit. As is well known, the song of the towhee, or chewink, at the north consists of two parts, nearly equal in length but otherwise quite different. In the Florida bird the last half is almost entirely omitted. According to Mr. Maynard, this variety is almost the only one occurring on Indian River, and of which he brought home some forty or more specimens. I found also one among half a dozen I shot at Jacksonville in January. In April, among a few towhees exposed in cages for sale in the market, were several of this kind. There is probably a large proportion of northern birds among the *Pipilones* of Northern Florida in winter, while probably in summer the majority are of the southern type above described, as are those of Middle and Southern Florida, doubtless, at all seasons.

Had this form been discovered ten, or even five years since, it would probably have been regarded by most ornithologists as entitled to specific rank, and not as a local race of *P. erythrophthalmus*, as it evidently is. Indeed, there are many species still on our lists that are far less entitled to rank as species than this, but which, though at first only provisionally adopted, have become traditionally established as valid species.

The two tables of measurements of specimens of this species given below, with Table J (p. 212), show the difference in size that obtains between Massachusetts and Florida specimens. The first table embraces twenty-nine specimens (nineteen males and ten females) of the white-eyed Florida type; the second table embraces sixteen specimens (ten males and six females) of the common northern type from Eastern Massachusetts; the measurements of twenty other Massachusetts males having been also already given in Table J, on p. 212. The following are the extremes of the two series. Northern type, males: Length, 7.50 and 8.80; alar extent, 10.00 and 12.25; wing, 3.17 and 3.90; tail, 3.30 and 3.93;

*Measurements of Specimens of PIPILLO ERYTHROPHthalmus from
Indian River, Florida.*

M. C. Z. No.	Collector's Number.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.	Tarsus.
10722	2477	♂	Dummitt's	Feb. 26, '69	C. J. Maynard	8.10	10.25	3.25	3.60	1.00
—	2476	♂	"	Feb. 26, '69	"	8.00	10.50	3.20	3.65	.95
—	2531	♂	"	Mar. 5, '69	"	8.00	10.00	3.00	3.60	1.00
—	2530	♂	"	Mar. 5, '69	"	7.20	10.10	3.00	3.65	.90
—	2529	♂	"	Mar. 6, '69	"	8.30	10.20	3.25	3.70	1.00
—	2559	♂	"	Mar. 7, '69	"	7.90	9.85	3.42	3.50	1.02
10729	2669	♂	"	Mar. 12, '69	"	8.20	11.00	3.45	3.57	1.01
—	2592	♂	"	Mar. 11, '69	"	8.50	11.00	3.50	3.60	.87
—	2417	♂	"	Feb. 22, '69	"	8.00	10.00	3.05	3.70	.95
—	2416	♂	"	Feb. 22, '69	"	8.00	10.25	3.25	3.35	—
—	2394	♂	"	Feb. 25, '69	"	8.00	10.10	3.10	3.50	.95
—	2426	♂	"	Feb. 20, '69	"	7.78	10.00	3.00	3.25	.90
—	2514	♂	"	Mar. 2, '69	"	7.70	9.50	2.80	3.40	.80
—	2512	♂	"	Mar. 2, '69	"	7.25	10.25	3.10	3.70	.80
10724	2511	♂	"	Feb. 20, '69	"	7.75	10.50	3.20	3.50	1.09
10721	2395	♂	"	Feb. 18, '69	"	7.40	10.00	2.92	3.90	—
10729	2668	♂	"	Feb. 22, '69	"	7.40	10.75	3.00	3.45	.95
—	2044	♂	"	Feb. 17, '69	"	8.50	11.30	3.00	3.50	.95
10728	2516	♂	"	Mar. 2, '69	"	8.00	10.25	3.05	3.45	.90
—	2481	♂	"	Feb. 24, '69	"	7.50	9.48	3.00	3.60	.90
—	2483	♂	"	Feb. 20, '69	"	7.50	9.50	3.00	3.54	.90
—	2590	♂	"	Mar. 11, '69	"	7.65	10.05	3.00	3.75	.92
—	2593	♂	"	Mar. 11, '69	"	8.20	10.10	3.05	3.58	1.00
—	2591	♂	"	Mar. 11, '69	"	7.90	9.95	3.03	3.60	.92
—	2578	♂	"	Mar. 10, '69	"	7.40	9.75	3.00	3.80	.95
10727	2515	♂	"	Mar. 2, '69	"	7.50	9.75	3.10	3.05	.95
10725	2513	♂	"	Mar. 2, '69	"	7.35	10.50	3.35	3.10	.85
—	2445	♂	"	Feb. 18, '69	"	8.05	10.53	3.50	3.45	.95
10726	2514	♂	"	Mar. 2, '69	"	7.50	9.50	3.90	3.05	.95

*Measurements of Specimens of PIPILLO ERYTHROPHthalmus from
Eastern Massachusetts.*

M. C. Z. No.	Collector's Number.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.	Tarsus.
—	384	♂	Newton	May 6, '68	C. J. Maynard	8.30	12.35	3.30	3.60	—
—	414	♂	Weston	May 9, '68	"	8.25	11.00	3.35	3.64	—
—	415	♂	"	May 9, '68	"	8.20	11.30	3.45	3.49	—
—	430	♂	Newton	May 11, '68	"	8.25	11.00	3.65	3.76	—
—	439	♂	Waltham	May 13, '68	"	8.45	11.30	3.51	3.85	—
—	539	♂	"	May 18, '68	"	8.35	11.23	3.46	3.55	—
—	1008	♂	Weston	July 10, '68	"	8.50	11.00	3.41	3.46	—
4616	1329	♂	Newton	Sept. 21, '68	"	8.80	11.65	3.42	3.70	1.10
—	1330	♂	"	Sept. 21, '68	"	8.51	11.41	3.51	4.00	1.00
4615	1295	♂	"	Sept. 17, '68	"	8.50	11.50	3.55	3.75	1.00
4725	496	♂	"	May 15, '68	"	7.60	10.45	3.31	3.35	—
4726	527	♂	Waltham	May 16, '68	"	8.25	11.76	3.35	3.60	—
—	—	♂	"	May 16, '68	"	8.26	10.55	3.55	3.65	—
4724	555	♂	Weston.	May 20, '68	"	8.00	10.56	3.25	3.36	—
4617	1028	♂	Newton.	July 22, '68	"	8.50	11.00	3.41	3.52	1.10
4613	1328	♂	"	Sept. 21, '68	"	8.45	11.00	3.35	3.65	1.05

tarsus, .98 and 1.13. Southern type, males: Length, 7.20 and 8.50; alar extent, 9.50 and 11.30; wing, 2.80 and 3.50; tail, 3.25 and 3.90; tarsus,

.80 and 1.09. The females in both cases average a little smaller than the males. The average dimensions of thirty northern males are as follows: Length, 8.19; alar extent, 11.32; wing, 3.43; tail, 3.66; tarsus, 1.06. Of nineteen southern males: Length, 7.88; alar extent, 9.88; wing, 3.13; tail, 3.56; tarsus, .94. The measurements given in the two preceding tables were all taken by Mr. Maynard from fresh specimens.

Other species of *Fringillidæ* that from their general distribution one naturally expects to meet with in East Florida in winter, but which, so far as I can learn, have not yet been met with there, are the Yellow-winged Sparrow (*Coturniculus passerinus*), Black-throated Bunting (*Euspiza americana*), Indigo Bird (*Cyanospiza cyanea*), and the Nonpareil (*C. ciris*). Specimens of the latter, collected at Cape Florida in winter, have been received at the Museum, and it was taken in April at Jacksonville and St. Augustine by Mr. Thurston and Mr. L. L. Thaxter.

ICTERIDÆ.

55.† *Molothrus pecoris* Swainson. COW BLACKBIRD.

Not numerous. Sometimes seen in small parties by themselves, but more frequently associating with the red-wings and grackles.

56.* *Agelæus phœniceus* Vieillot. RED-WINGED BLACKBIRD.

Abundant. Apparently chiefly Florida born birds seen, especially in February and March. The sexes were usually in separate flocks.

The differences in respect to size and color between Florida and New England specimens usually seen in individuals of the same species from these localities are very marked in the present species, especially in respect to color. In no group, in fact, is it generally more so than in the *Icteridæ*.

In the Florida red-wings the general form is slenderer and more delicate, the bill relatively longer and more pointed, and the general color more intense and lustrous. The difference is particularly marked in the shoulder-patch, in which the red of its anterior portion is darker, approaching bright orange, and the posterior part, which in the northern bird is usually pale cream-color, whitish, or even nearly pure white, is orange-yellow, — very nearly as in the *A. gubernator* of the Pacific States. The difference in color, size, and especially in the form of the bill, is much greater than the differences existing between many currently received species of North American birds, and it is surprising that the two forms have not been specifically separated. I can only account for it on the supposition that specimens from Florida and the Gulf States have not fallen

into the hands of the assiduous species hunters. As remarked in Part III (p. 234), Florida and New England specimens are as different from each other as are the so-called *Agelæus phæniceus* of the Northeastern States, the *A. tricolor* and the *A. gubernator* from each other.

Specimens of *A. phæniceus* from Louisiana I find correspond very nearly in every respect with the specimens from Florida. I have also before me one specimen from Maine with the shoulder-patch as highly colored, and with nearly as long a bill as is found in the specimens from Florida.

Plate VI shows the average form of the bill in Florida and Massachusetts specimens, and the annexed table of measurements the difference in general size. They also illustrate individual variation.

The following measurements of seventy specimens of this species from Massachusetts (forty males and thirty females), eighteen specimens from South Carolina and Florida (eleven males and seven females), and thirteen specimens from California (four males and nine females), exhibit, besides the average size and the individual variation at the same locality (especially in the case of those from Massachusetts), several interesting facts in respect to geographical variation. While the northern specimens (see the summary of these measurements given below) are somewhat larger than the southern ones, the latter have the longer head (including the bill), and also the longer bill. The height and width of the bill at the base remaining essentially the same in both, the southern ones have the bill relatively more attenuated. The difference in this respect is more striking than the measurements given seem to indicate. The California specimens closely resemble those from Florida, not only in respect to size, but in regard to the size and form of the bill, and also in respect to color; these, as well as the Florida ones, belonging to the southern type. As previously remarked, they bear a much closer resemblance to the Florida form in every respect than to that found in New England.*

The individual variation in this species seems to be very great everywhere, the variation in specimens of the same sex from the same locality being fully fifteen per cent of the average size at that locality.

* The affinities of *Agelæus gubernator* and *A. tricolor* with *A. phæniceus* are acknowledged to be exceedingly close. Professor Baird cites, in his *Birds of North America*, one specimen of the *A. phæniceus* from San José, California, and five from Fort Steila-coom, W. T. He also cites specimens of *A. gubernator* from Petaluma and San Francisco, Cal.; but Dr. Cooper regards this species as "limited to the interior of the State" (California), while those found along the coast, he says, clearly resemble the eastern bird. (*Ornithology of California*, Vol. I, p. 264.) From the close resemblance, already alluded to, of both the *A. gubernator* and *A. tricolor* to *A. phæniceus*, and their occurrence mainly in the hot valleys of California and the region more to the southward, I can scarcely doubt that these forms, especially *A. gubernator*, are the southern smaller, brighter colored, more attenuated billed western homologues of the similar eastern form from Florida and the Gulf States.

Measurements of Northern Specimens of AGELÆUS PHŒNICEUS.

M. C. Z. No.	Collector's Number.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.	Head.	Bill.		
											Culmen.	Height.	Width.
1274	—	♂	Vassalboro', Me.	—	Mr. Becker	9.40	14.90	4.65	3.82	1.82	.87	.45	.37
9577	—	♂	Waterville, "	June 20, '64	C. E. Hamlin	9.45	14.75	4.82	3.88	1.77	.87	.46	.40
391	—	♂	Malden, Mass.	1859	D. Higgins	9.35	14.75	4.67	3.75	1.76	.86	.47	.40
96	—	♂	"	"	"	9.15	14.35	4.55	3.50	1.75	.80	.50	.37
392	—	♂	"	"	"	9.20	14.75	4.70	3.66	1.82	.89	.47	.38
92	—	♂	"	"	"	9.25	14.50	4.75	3.82	1.80	.92	.50	.40
93	—	♂	"	"	"	9.20	14.40	4.50	3.45	1.85	.88	.45	.40
94	—	♂	"	"	"	9.00	14.25	4.60	3.55	1.76	.90	.50	.38
95	—	♂	"	"	"	9.00	15.10	4.65	3.65	1.83	.95	.50	.40
393	—	♂	"	"	"	9.00	14.55	4.65	3.35	1.82	.84	.45	.37
394	—	♂	"	"	"	8.40	14.10	4.50	3.40	1.60	.75	.46	.40
5727	—	♂	Concord, "	"	H. Mann	8.58	13.95	4.50	3.46	1.78	.92	.48	.40
5723	—	♂	"	"	"	8.45	14.45	4.55	3.35	1.75	.90	.43	.45
5724	—	♂	"	"	"	8.40	14.25	4.43	3.12	1.80	.84	.45	.38
5729	—	♂	"	"	"	9.00	14.51	4.54	3.33	1.82	.85	.47	.39
5726	—	♂	"	"	"	9.20	14.95	4.75	3.75	1.80	.87	.44	.38
5720	—	♂	"	"	"	9.05	15.00	4.86	3.86	1.84	.88	.44	.37
5721	—	♂	"	"	"	9.70	14.88	4.68	3.80	1.77	.92	.46	.40
5722	—	♂	"	"	"	9.00	14.50	4.65	3.65	1.74	.83	.43	.38
5725	—	♂	"	"	"	9.85	14.25	4.73	3.76	1.85	.93	.43	.35
5728	—	♂	"	"	"	9.25	14.50	4.60	3.81	1.78	.88	.44	.33
5732	—	♂	"	"	"	9.58	14.62	4.74	3.85	1.83	.91	.48	.38
10096	—	♂	Ipswich, "	June 14, '68	J. A. Allen	9.15	14.50	4.70	3.78	1.79	.92	.44	.40
1674	—	♂	Springfield, "	June 26, '62	"	9.00	14.00	4.59	3.42	1.62	.87	.45	.40
1675	—	♂	"	June 26, '62	"	9.25	15.00	4.82	3.73	1.84	.93	.46	.43
1782	—	♂	"	July 12, '62	"	9.50	14.50	4.62	3.65	1.94	.97	.45	.37
1781	—	♂	"	July 12, '62	"	9.00	14.60	4.50	3.38	1.80	.86	.43	.40
626	—	♂	Auburndale, "	Mar. 23, '57	S. Tenney	9.62	15.35	4.87	3.77	1.85	.91	.43	.40
1022	—	♂	Wenham, "	May —, '61	J. Bartlett	9.25	15.00	4.67	3.55	1.84	.89	.48	.43
—	114	♂	Newton, "	Mar. 13, '68	C. J. Maynard	8.75	14.83	4.70	3.53	—	—	—	—
—	180	♂	"	Mar. 28, '68	"	9.38	15.00	4.75	3.60	—	—	—	—
—	214	♂	"	Apr. 11, '68	"	8.90	14.85	4.60	3.52	—	—	—	—
—	251	♂	Weston, "	Apr. 18, '68	"	9.52	15.00	4.82	3.80	—	—	—	—
—	302	♂	Newton, "	Apr. 23, '68	"	9.00	15.00	4.66	3.65	—	—	—	—
—	323	♂	"	Apr. 25, '68	"	9.00	15.00	4.85	3.40	—	—	—	—
—	352	♂	Weston, "	May 1, '68	"	9.16	15.10	4.80	3.70	—	—	—	—
—	371	♂	Newton, "	May 5, '68	"	9.00	15.00	5.00	3.90	—	—	—	—
—	351	♂	"	May 1, '68	"	9.50	15.25	4.92	3.85	—	—	—	—
—	417	♂	Weston, "	May 9, '68	"	9.50	15.15	4.90	3.77	—	—	—	—
—	3047	♂	Newton, "	Mar. 23, '70	"	9.50	15.00	4.60	3.56	—	—	—	—
9843	—	♂	Milltown, Me.	—	G. A. Boardman	8.00	12.50	3.95	3.05	1.57	.73	.38	.32
9844	—	♂	"	—	"	8.00	12.50	4.08	3.10	1.54	.79	.40	.30
399	—	♂	Malden, Mass.	1859	D. Higgins	7.50	11.75	3.60	2.70	1.48	.72	.42	.33
396	—	♂	"	"	"	7.75	12.30	3.90	3.06	1.60	.82	.43	.35
97	—	♂	"	"	"	7.35	11.75	3.63	2.80	1.55	.70	.40	.37
402	—	♂	"	"	"	8.55	13.55	4.26	3.15	1.66	.75	.44	.41
398	—	♂	"	"	"	8.05	13.50	4.16	3.10	1.67	.77	.43	.36
98	—	♂	"	"	"	7.75	12.10	3.74	2.97	1.58	.72	.37	.33
403	—	♂	"	"	"	7.42	11.55	3.87	2.73	1.62	.73	.37	.35
397	—	♂	"	"	"	7.50	11.25	3.70	2.98	1.58	.76	.34	.33
395	—	♂	"	"	"	7.40	12.50	4.11	2.68	1.50	.70	.38	.38
99	—	♂	"	"	"	7.45	11.50	3.75	2.90	1.55	.75	.42	.43
5730	—	♂	Concord, "	—	H. Mann	7.75	12.50	3.80	3.02	1.54	.70	.40	.33
1611	—	♂	Springfield, "	July 15, '62	J. A. Allen	7.50	12.00	3.73	2.90	1.68	.79	.40	.34
1639	—	♂	"	June 26, '62	"	7.65	11.75	3.67	2.82	1.54	.70	.38	.37
1672	—	♂	"	June 26, '62	"	7.85	11.82	3.75	2.89	1.60	.75	.40	.38
1673	—	♂	"	June 26, '62	"	7.75	12.00	3.77	2.85	1.57	.78	.38	.37
1679	—	♂	"	June 26, '62	"	8.00	12.25	3.79	3.00	1.55	.74	.39	.36
1680	—	♂	"	June 26, '62	"	8.00	12.10	3.85	2.93	1.53	.78	.38	—
—	350	♂	Weston, "	Apr. 30, '68	C. J. Maynard	7.73	12.61	3.95	2.95	—	—	—	—
—	850	♂	Ipswich, "	June 15, '68	"	7.45	13.60	4.00	3.05	—	—	—	—
—	891	♂	Essex, "	June 17, '68	"	8.09	12.35	4.00	3.00	—	—	—	—
—	893	♂	"	June 17, '68	"	7.75	12.54	3.90	2.90	—	—	—	—
—	1093	♂	Waltham, "	Aug. —, '68	"	7.45	12.37	3.80	2.75	—	—	—	—
—	1075	♂	"	Aug. —, '68	"	7.67	12.30	3.85	2.85	—	—	—	—
—	1096	♂	"	Aug. —, '68	"	7.50	12.40	3.95	2.72	—	—	—	—
—	1097	♂	Newton, "	Aug. —, '68	"	7.75	12.30	3.71	2.80	—	—	—	—
—	1098	♂	Waltham, "	Aug. —, '68	"	7.50	12.00	3.85	2.65	—	—	—	—
—	1099	♂	Weston, "	Aug. —, '68	"	7.50	12.10	3.67	2.80	—	—	—	—
—	2830	♂	Newton, "	June 8, '69	"	7.50	12.00	4.00	2.90	—	—	—	—

Measurements of Southern Specimens of AGELÆUS PHŒNICEUS.

M. C. Z. No.	Coll. No.	Sex.	Locality.	Date.	Collector.	Length.	Al. Ext.	Wing.	Tail.	Head.	Bill.		
											Cul.	Hgt.	Wid.
4126	—	—	Charleston, S. C.	—	L. Agassiz	9.55	14.75	4.75	3.65	1.90	1.00	.45	.37
4127	—	—	" "	—	"	8.80	14.30	4.50	3.55	1.74	.87	.47	.40
4128	—	—	" "	—	"	9.45	14.50	4.60	3.72	1.80	.90	.50	.40
4129	—	—	" "	—	"	9.05	13.50	4.37	3.45	1.78	.85	.43	.42
4125	—	—	" "	—	"	9.05	14.12	4.42	3.35	1.94	.95	.46	.35
—	—	—	Hawkinsville, Fla.	Mar. 15, '69	J. A. Allen	8.25	13.60	4.34	—	—	—	—	—
—	1928	—	Jacksonville, "	Dec. 31, '69	C. J. Maynard	9.10	14.90	4.75	3.58	—	—	—	—
10563	1929	—	" "	Dec. 31, '69	"	9.20	14.80	4.80	3.90	—	—	—	—
10561	2013	—	" "	Dec. 31, '69	"	8.80	14.15	4.55	3.58	—	—	—	—
10574	2552	—	Dummitt's, "	Mar. 8, '69	"	9.50	14.20	4.75	3.90	—	—	—	—
10573	2450	—	" "	Feb. 24, '69	"	8.50	14.00	4.75	3.45	—	—	—	—
5153	—	—	Hibernia, "	Jan. 30, '69	J. A. Allen	7.65	12.60	3.85	3.05	—	—	—	—
5154	—	—	" "	Jan. 30, '69	"	7.85	12.50	3.90	3.07	—	—	—	—
5155	—	—	" "	Jan. 30, '69	"	7.80	12.85	—	3.20	—	—	—	—
4141	—	—	" "	Jan. 30, '69	"	8.00	12.25	3.80	3.05	—	—	—	—
5209	—	—	Welaka, "	Feb. 8, '69	"	7.65	12.50	3.75	2.90	—	—	—	—
5208	—	—	" "	Feb. 8, '69	"	7.50	11.85	3.63	2.75	—	—	—	—
5210	—	—	" "	Feb. 8, '69	"	7.65	12.55	3.95	—	—	—	—	—

Measurements of California Specimens of AGELÆUS PHŒNICEUS.

M. C. Z. No.	Sex.	Locality.	Date.	Collector.	Length.	Al. Ext.	Wing.	Tail.	Tarsus.
5885	—	San Francisco, Cal.	Winter '59-'60	A. Agassiz	8.50	14.98	4.83	3.50	1.63
5884	—	" "	Winter '59-'60	"	8.75	15.05	4.95	3.35	1.74
566	—	" "	Winter '59-'60	"	8.60	14.55	4.47	3.09	1.90
2188	—	Gulf of Georgia, W. T.	Sept. —, '60	"	8.71	13.50	4.45	3.26	1.75
5889	—	San Francisco, Cal.	Winter '59-'60	"	7.58	12.80	4.03	2.73	1.63
5893	—	" "	Winter '59-'60	"	7.55	12.35	3.95	2.47	1.46
5887	—	" "	Winter '59-'60	"	7.81	12.80	4.25	3.86	1.54
5890	—	" "	Winter '59-'60	"	7.50	12.75	3.94	2.47	1.56
5886	—	" "	Winter '59-'60	T. G. Cary	7.82	12.77	4.04	2.62	1.56
2075	—	" "	Winter '59-'60	"	8.29	13.27	4.32	3.00	1.62
2074	—	" "	Winter '59-'60	"	8.18	13.25	3.85	2.95	1.67
2078	—	" "	Winter '59-'60	"	8.50	13.00	4.15	3.10	1.65
5888	—	" "	Winter '59-'60	A. Agassiz	7.25	12.25	3.90	3.71	1.50

Summary of the above Measurements of Specimens of AGELÆUS PHŒNICEUS.

Locality.	Sex.	No. of Specimens	Aver.	Length.	Al. Ext.	Wing.	Tail.	Head.	Culmen.	Height.	Width.
Massachusetts	♂	40	Aver.	9.16	14.71	4.69	3.63	1.79*	.88*	.46*	.39*
South Carolina and Florida	♂	28	Aver.	7.53	12.24	3.86	2.93	1.57†	.75†	.395†	.357†
	♂	11	Aver.	9.02	14.41	4.62	3.61	1.83	.91†	.46†	.39†
	♂	7	Aver.	7.73	12.44	3.83	2.99	—	—	—	—
California	♂	7	Aver.	8.64	14.52	4.67	3.30	1.75	—	—	—
	♂	9	Aver.	7.83	12.70	4.00	2.99	1.57	—	—	—
	♂	40	Max.	9.85	15.35	5.00	3.90	1.94*	.97*	.50*	.45*
Massachusetts	♂	40	Min.	8.40	13.95	4.43	3.12	1.60*	.75*	.43*	.33*
	♂	28	Max.	8.55	13.55	4.26	3.15	1.68†	.82†	.44†	.43†
	♂	23	Min.	7.35	11.25	3.63	2.65	1.48†	.70†	.37†	.30†
	♂	11	Max.	9.55	14.90	4.80	3.90	1.94†	1.00†	.50†	.40†
South Carolina and Florida	♂	11	Min.	8.25	13.60	4.34	3.35	1.74†	.85†	.43†	.35†
	♂	7	Max.	8.00	12.85	3.90	3.20	—	—	—	—
	♂	7	Min.	7.50	11.85	3.63	2.75	—	—	—	—
	♂	7	Max.	8.75	15.05	4.95	3.50	1.90	—	—	—
California	♂	7	Min.	8.50	13.50	4.45	3.09	1.63	—	—	—
	♂	9	Max.	8.50	13.27	4.32	3.86	1.67	—	—	—
	♂	9	Min.	7.25	12.25	3.85	2.47	1.46	—	—	—

* 29 specimens.

† 19 specimens.

‡ 5 specimens.

57.* *Sturnella ludoviciana* Swainson. MEADOW LARK.

- Alauda magna* LINNÉ, Syst. Nat., I, 167, 1758. — WILSON, Am. Orn., III, 20, pl. xix, 1811.
- Sturnus ludovicianus* LINNÉ, Syst. Nat., I, 290, 1766. — BONAP., Journ. Phil. Acad. Nat. Sci., IV, 180, 1824. — NUTTALL, Man. Orn., I, 147, 1832. — AUDUBON, Orn. Biog., II, 216, 1834.
- Sturnus collaris* WAGLER, Syst. Avium, I, 1827.
- Sturnella ludoviciana* SWAINSON, Faun. Bor. Am., II, 282, 1831. — BONAP., Geog. and Comp. List, 1838. — AUDUBON, Synop. Am. Birds, 148, 1839. — CABANIS, Mus. Hein., 192, 1851. — SCLATER, Cat. Am. Birds, 139, 1862. — CASSIN, Proc. Phil. Acad. Nat. Sci., 1866, 23.
- Sturnella magna* SWAINSON, Phil. Mag., I, 436, 1827. — BAIRD, Birds N. Am., 535, 1858. — ALLEN, Mem. Bost. Soc. Nat. Hist., I, 496, 1868.
- Sturnella collaris* VIEILLOT, Analyse, 1816.
- Sturnella hippocrepis* WAGLER, Isis, 1832, 281. — LAWRENCE, Ann. N. York Lyceum N. Hist., VII, 266, 1860. — SCLATER, Ibis, 1861, 79. — CASSIN, Proc. Phil. Acad. Nat. Sci., 1866, 24.
- Sturnella neglecta* AUDUBON, Birds of Am., VII, 339, pl. cccclxxxvii, 1843. — BAIRD, Birds of N. Am., 537, 1858. — CASSIN, Proc. Phil. Acad. Nat. Sci., 1866, 23.
- Sturnella mexicana* SCLATER, Ibis, 1861, 79. — CASSIN, Proc. Phil. Acad. Nat. Sci., 1866, 24.
- Sturnella meridionalis* SCLATER, Ibis, 1861, 79. — CASSIN, Proc. Phil. Acad. Nat. Sci., 1866, 24.

Abundant. Found chiefly in the moister parts of the pinneries.

Somewhat smaller than in the Northern States, but in most cases with longer and larger bills, brighter colors, and a quite different song. The latter somewhat resembles that of the western meadow lark, but is still as distinct from it in its general character as it is from that of the New England bird. The present species has a wide geographical range, throughout the greater part of which it is resident. The Alleghanian fauna forms its northern limit, from which it mostly retires during winter. To the southward it extends to Cuba and the other larger West India Islands, throughout most of Central America, and to the elevated parts of Northern South America. It ranges westward over the elevated arid plains of the middle of the continent to the Pacific. As might be expected, it is not quite uniform in its characters at all points. The main differences, however, consist merely in the lighter color of those from the plains, and the smaller size of those from the south. The former constitute the *Sturnella neglecta* of Audubon and most other writers since his time. In Cuba it is the *S. hippocrepis* of Wagler and others, and the Mexican and Guatemalan form is the *S. mexicana* of Sclater, and the South American form the *S.*

meridionalis of the same author. Yet the distinctions between them are trivial, all of these so-called species having been generally looked upon as doubtfully distinct from the *S. ludoviciana* of the United States, especially the three last named. The *S. collaris* of Vieillot has very generally been referred by subsequent writers to the *S. ludoviciana*. The main distinctive feature of the *S. neglecta* has been its song, — a very doubtful basis on which to found a species. The Florida specimens are intermediate in size and other characters between the Cuban and New England representatives of this species. As already remarked, the song of the Florida birds is as widely different from that of the New England bird as the song of the latter is from that of the western ones. Concerning the affinities of *S. neglecta* I have already remarked.* Concerning those of the other supposed species, I may well borrow the appropriate remarks of the late Mr. Cassin, who observes in respect to them, in his "Study of the *Icteridæ*,"† as follows:—

"This bird [*Sturnella ludoviciana*] is nearly related to the next four species of this genus [*S. neglecta*, *S. hippocrepis*, *S. mexicana*, *S. meridionalis*], equally in structure and in colors, and it would be difficult to describe by positive characters either species of this group, so as to insure recognition absolutely, or without comparative characters being given. . . . No other genus or sub-genus of this family presents so many species of such uniformity of structure and similarity of color, and there are, assuredly, few such in the kingdom of birds." Under *S. neglecta* he further remarks in respect to the transition that is so apparent between it and *S. ludoviciana*: "In the central regions of North America it is possible that a hybrid race between the two species may be produced, to be referred with about equal propriety to either." *S. hippocrepis*, he says, is very nearly related to *S. mexicana*, "and can scarcely be distinguished from it by any characters which seem to be reliable." He thinks it to be somewhat more distinct, however, from *S. neglecta*. Mr. Lawrence had previously remarked that the *S. hippocrepis* is somewhat smaller than *S. ludoviciana* of the United States, and that he "thinks it is specifically distinct"; although he adds, "it would be difficult to point out any reliable differences in coloration, especially of the upper plumage, as individuals even of the same species are very variable."‡ He says, further, that specimens of it from Jalapa, Mexico, differ "only in the pectoral band appearing broader in the Mexican bird, and the tertials much shorter than the primaries, but this last may not be a reliable character." In the

* See Memoirs of the Boston Soc. Nat. Hist., Vol. I, p. 494, 1868.

† Proceedings of the Phil. Acad. Nat. Sciences, 1866, p. 23.

‡ Annals of New York Lyceum of Nat. Hist., Vol. VII, 266, 1860.

following year, however, Mr. Sclater separated the Mexican bird from those of Cuba and the United States, under the name *S. mexicana*, and also the South American under the name *S. meridionalis*. Mr. Cassin says of the latter: "Very nearly related to the preceding (*S. hippocrepis*), if distinct, and I give it, at present, as a species provisionally only. . . . The colors of the upper parts seem to be less clearly defined, and of a slightly different style and pattern from the preceding, and it may bear about the same relation to that species (*S. hippocrepis*) that *S. neglecta* does to *S. ludoviciana*. Such relation I hold to be rather probable from the specimens now at hand."

Having given the views of the describers of these several "species," I may add that I have seen examples of each, and do not question that they should all be referred to one. As is evident from the above-quoted remarks, these different species gradually pass into each other,—the *S. magna* into the *S. neglecta*, the *S. neglecta* into the *S. mexicana*, and the *S. mexicana* into the *S. hippocrepis*, which is their exact geographical relation.

In regard to the Florida specimens, as compared with New England ones, the most striking differences consist in their smaller size and much brighter colors, especially of the ventral surface.

The following tables of measurements indicate the individual and sexual differences in size, and also the difference in size between specimens from the Northern States and from Florida.

Measurements of Northern Specimens of STURNELLA LUDOVICIANA.

M. C. Z. No.	Coll. No.	Sex.	Locality.	Date.	Collector.	Length.	Al. Ext.	Wing	Tail.
4862	416		Newton, Mass.	May 8, '68	C. J. Maynard	10.75	16.59	5.13	3.50
—	1100		Waltham, "	Aug. 6, '68	"	10.20	16.30	4.98	2.90
4863	1134		"	Aug. 19, '68	"	10.25	15.85	4.80	2.92
—	2696		Newton, "	May 15, '69	"	10.75	16.75	5.00	3.35
—	2698		"	May 15, '69	"	11.00	17.00	5.15	3.35
—	1700		Waltham, "	Aug. 6, '68	"	10.20	16.30	4.98	2.90
—	4045		Newton, "	Aug. 2, '69	"	11.00	17.00	5.00	3.40
—	4061		"	Aug. 2, '69	"	11.00	16.00	4.80	3.35
—	2738		"	May 15, '69	"	9.75	15.00	4.55	2.65
362	—		Malden, "	—	D. Higgins	9.25	13.50	4.17	2.50
363	—		"	—	"	9.58	14.00	4.35	2.90
364	—		"	—	"	10.50	15.33	4.82	3.11
365	—		"	—	"	10.35	15.65	4.83	3.13
366	—		"	—	"	10.00	15.05	4.75	3.14
367	—		"	—	"	10.75	15.50	4.82	3.30
568	—		"	—	"	8.90	14.00	4.35	2.60
569	—		"	—	"	9.50	15.68	5.00	3.05
9764	—		Evanston, Ill.	—	O. Marcy	9.25	13.92	4.15	2.82
9765	—		"	—	"	9.75	14.65	4.50	3.10
9766	—		"	—	"	10.00	15.50	4.74	3.15
2646	—		Lawn Ridge, "	—	K. Butler	9.60	14.75	4.55	2.84
4042	—		Concord, Mass.	—	F. C. Brown	10.25	16.00	4.77	2.83
4102	—		"	—	"	10.33	15.65	4.67	3.08

Measurements of Florida Specimens of STURNELLA LUDOVICIANA.

M. C. Z. No.	Coll. No.	Sex.	Locality.	Date.	Collector.	Length.	Al. Ext.	Wing.	Tail.
—	2037	♂	Jacksonville	Jan. 20, '69	C. J. Maynard	9.55	15.60	4.50	2.85
—	2817	♂	Dummitt's	May 15, '69	"	10.20	15.10	4.50	3.20
—	2816	♂	"	May 15, '69	"	10.00	15.15	4.60	2.95
5335	—	♂	Enterprise	Mar. 4, '69	J. A. Allen	9.75	14.75	4.50	—
5336	—	♂	"	Mar. 4, '69	"	9.85	15.20	4.40	2.89
5337	—	♂	"	Mar. 4, '69	"	9.70	14.80	4.45	2.82
5368	—	♂	Hawkinsville	Mar. 12, '69	"	9.75	15.00	4.50	3.05
5339	—	♂	"	Mar. 12, '69	"	9.50	14.75	4.25	4.83
5371	—	♂	"	Mar. 12, '69	"	10.00	15.75	4.50	3.07
—	—	♂	Volusia	Mar. 17, '69	"	8.75	13.75	4.05	—
5370	—	♂	Hawkinsville	Mar. 12, '69	"	8.90	14.15	4.10	2.65
5372	—	♂	"	Mar. 12, '69	"	9.50	14.65	4.20	2.90
5125	—	♂	Jacksonville	Jan. 19, '69	"	8.75	14.25	4.20	2.70
—	2072	♂	"	Jan. 20, '69	C. J. Maynard	8.75	14.00	4.40	2.50
—	2070	♂	"	Jan. 20, '69	"	8.50	13.55	3.90	2.55
—	2070	♂	"	Jan. 20, '69	"	8.75	13.00	4.00	2.55
—	2068	♂	"	Jan. 20, '69	"	9.25	14.75	4.50	2.80
—	2069	♂	"	Jan. 20, '69	"	8.76	14.25	4.20	2.40
—	2071	♂	"	Jan. 20, '69	"	9.00	14.00	4.30	2.80
—	2051	♂	"	Jan. 20, '69	"	9.50	14.75	4.65	2.50
—	2791	♂	"	Apr. 15, '69	"	9.05	14.00	4.10	2.88

The following is a tabulated summary of the two preceding tables:—

No. of Speci- mens.	Sex.	Locality.		Length.	Alar Extent.	Wing.	Tail.
15	♂	Northern States	Average	10.43	16.30	4.91	3.16
8	♂	"	Average	9.55	14.43	4.29	2.82
15	♂	"	Maximum	11.00	17.00	5.15	3.50
15	♂	"	Minimum	10.00	15.05	4.74	2.83
8	♂	"	Maximum	9.75	15.65	4.55	3.10
8	♂	"	Minimum	8.90	13.50	4.15	2.50
9	♂	Florida	Average	9.81	15.70	4.47	2.85
12	♂	"	Average	8.93	14.09	4.22	2.57
9	♂	"	Maximum	10.20	15.75	4.60	3.20
9	♂	"	Minimum	9.50	14.75	4.25	2.82
12	♂	"	Maximum	9.50	14.75	4.65	2.90
12	♂	"	Minimum	8.50	13.00	3.90	2.40

58.† *Scolecophagus ferrugineus* Swainson. RUSTY GRACKLE.

Abundant. Occasionally met with in large flocks.

59.* *Quiscalus purpureus* Cassin. PURPLE GRACKLE.

Gracula quiscula LINNÉ, Syst. Nat., I, 165, 1766. — WILSON, Am. Orn., III, 44, pl. xxi, fig. 4, 1811.

Gracula barita LINNÉ, Syst. Nat., 165, 1766. — ORD, Journ. Phil. Acad. Nat. Sci., I, 253, 1818.

Gracula purpurea BARTRAM, Travels, 289, 1791. (No description.)

? *Oriolus ludovicianus* GMELIN, Syst. Nat., 387, 1788.

Quiscalus baritus VIEILLOT, Nouv. Dict., XXVIII, 487, 1819. — BAIRD, Birds North Amer., 556, pl. xxvii, 1858. — CASSIN, Proc. Phil. Acad. Nat. Sci., 1866, 405.

Quiscalus versicolor VIEILLOT, Nouv. Dict., XXVIII 488, 1819. — BONAPARTE, SWAINSON, NUTTALL, AUDUBON, BAIRD.

- Quiscalus purpureus* CASSIN, Proc. Phil. Acad. Nat. Sci., 1866, 403. — RIDGWAY, Ibid., 1869, 133.
- Quiscalus purpuratus* SWAIN., Lardner's Cab. Cyclop., 299, 1838 (female).
- ? *Quiscalus lugubris* SWAIN., Lardner's Cab. Cyclop., 299, 1838. — ? CASSIN, Proc. Phil. Acad. Nat. Sci., 1866, 408.
- Quiscalus inflexirostris* SWAIN., Lardner's Cab. Cyclop., 300, 1838. — CASSIN, Proc. Phil. Acad. Nat. Sci., 1866, 407.
- Quiscalus crassirostris* SWAIN., Lardner's Cab. Cyclop., 355, 1838. — GOSSE, Birds of Jamaica, 217, 1847.
- Quiscalus agleus* BAIRD, Amer. Journ. Sci. and Arts, XLI, 87, 1866. — CASSIN, Proc. Phil. Acad. Nat. Sci., 1866, 404. — RIDGWAY, Ibid., 1869, 135.
- Quiscalus aeneus* RIDGWAY, Ibid., 134.
- Quiscalus mexicanus* CASSIN, Ibid., 1866, 408.
- Quiscalus Gundlachii* CASSIN, Ibid., 406.
- Quiscalus brachypterus* CASSIN, Ibid., 406.
- Quiscalus niger* CASSIN, Ibid., 407.
- ? *Quiscalus rectirostris* CASSIN, Ibid., 409.
- Chalcophanes quiscalus* WAGLER, Syst. Avium, 1827. — CABANIS, Mus. Hein., 197, 1851.
- Chalcophanes baritus* WAGLER, Syst. Avium, 1827. — CABANIS, Mus. Hein., 197, 1851.

Very abundant everywhere. Flocks containing many hundreds were frequently met with.

As already remarked in Part III, few species present such marked climatic variations as the present, or better illustrate the three principal laws of geographical variation already enumerated; namely, a decrease in general size from the north southward, and at the same time an increase in the length and slenderness of the bill, and an increase in the intensity and brilliancy of the color of the plumage. Far to the north, as in Labrador, the colder parts of Canada, and Northern New England, the bill is shortest and thickest, the size of the bird at its maximum, and the colors of the plumage least brilliant, with the metallic reflections of a light tint, tending to green rather than to blue. In Southern New Jersey the change from the northern type is already considerable; even between summer specimens from Calais (Maine) and Eastern Massachusetts there is an appreciable difference. In the lowlands of South Carolina and Georgia the divergence from the northern type is still greater, and it goes on rapidly increasing in Florida, especially in South Florida, the maximum of divergence from the northern type being attained in the West Indies. In East Florida, while the general size of the bird is less than in New England, the bill is considerably longer, much slenderer and much more decurved, as is shown by the accompanying figures (Plate VII). The

change in color is equally marked. Not only do the reflections become much darker at the south, but form prismatic bars across the interscapularies and the feathers of the rump, especially in the South Atlantic States. In South Florida and the West Indies these prismatic bars, in some specimens at least, seem to lose their distinctness, evidently through the continued darkening or increased intensity of the general color. The difference in size between Florida and Massachusetts specimens is considerable, especially between those from South Florida and Massachusetts. Those from the West Indies are still smaller; and in comparing specimens of these with others from Northern New England, the difference is so striking that it seems impossible at first to believe that both can belong to the same species, yet a gradual transition between the two, through the individuals inhabiting the intermediate region, fully proves it. Even between Florida and New England specimens the difference is so great that, were there no transition from one to the other, the two extremes might well be regarded as not only valid species, but as well-marked ones. Being familiar with the so-called *Quiscalus aglaeus* before visiting Florida, through specimens in the Museum of Comparative Zoölogy from Cape Florida, I had no doubt that it was a species distinct from the *Q. purpureus*. But a subsequent study of these birds in Florida, and an examination of specimens from various points between Florida and Northern Maine, and also from the West Indies, has forced me to the conclusions indicated in the above table of synonymes.

The purple grackles of the Mississippi Valley have recently been separated as specifically distinct from those of the Atlantic States, under the name *Q. æneus*, *Q. purpureus* being retained for the latter. The range of *Q. purpureus* is given as "Atlantic and Gulf? States, north to Nova Scotia, west to the Alleghanies." The New England type, however, is entirely referable to the *Q. æneus*, as defined by its describer. The same writer also follows some of his predecessors in separating those of South Florida from the *Q. purpureus*, under the name of *Q. aglaeus*. But Cape Florida specimens differ but little — being, in fact, scarcely distinguishable except in size — from those from the St. John's River.

Mr. Cassin, in one of his latest papers,* took the ground that each of the larger West India Islands has a distinct species of this group, peculiar to itself. That these forms, many of them evidently difficult of recognition, should be distinct species is quite contrary to general principles. These islands are generally separated by a distance of rarely more than a hundred miles; yet a near ally of these "species," the *Q. purpureus* (or *Q. æneus* as recently restricted), is admitted to range from the Gulf of Mexico

* "A Second Study of the Icteridæ," Proc. Phil. Acad. Nat. Sci., 1866, pp. 403 - 417.

to the arctic regions, so that those that breed farthest north make annually a journey of fully a thousand miles to reach their breeding-grounds. As I have already observed, individuals of species possessing a very northern habitat usually present a great uniformity of character, while those of species ranging farther to the southward are more variable; also that within the warm-temperate and tropical latitudes, islands but slightly separated from each other or the mainland, and peninsulas which, like Florida, are almost insular in their geographical relations, present each peculiar modifications of species ranging throughout not only all of them, but portions of the adjoining continents, which render the individuals from these different localities more or less readily distinguishable. This results partly, doubtless, from the isolation of these different districts, partly from the more sedentary habits of birds in warm countries, as compared with those of cold latitudes, and partly from the greater tendency to variation in species inhabiting tropical and sub-tropical countries.

In the subjoined tables measurements are given of thirteen males and eight females from the Northern States, and of twenty-three males and seven females from Florida, of which the following is a tabulated summary:—

No. of Specimens.	Sex.	Locality.		Length.	Alar Extent.	Wing.	Tail.
13	♂+♂+♂+♂+♂+♂+♂+♂+♂+♂+♂+♂	Northern States	Average	12.63	17.73	5.66	5.30
8		"	Average	11.45	15.76	4.94	4.49
13		"	Maximum	13.50	18.43	6.05	6.00
13		"	Minimum	12.00	17.00	5.20	4.58
8		"	Maximum	12.05	16.30	5.20	4.85
8		"	Minimum	10.90	15.38	4.60	4.10
23		Florida	Average	12.19	16.64	5.42	5.22
7		"	Average	11.12	14.86	4.75	4.55
23		"	Maximum	13.00	17.80	5.75	5.50
23		"	Minimum	11.00	15.25	5.00	4.55
7		"	Maximum	11.75	16.75	5.00	4.77
7		"	Minimum	10.25	13.75	4.50	4.45

Measurements of Northern Specimens of QUISCALUS PURPUREUS.

M. C. Z. No.	Coll. No.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.
1234	—	♂	Red River, B. A.	—	S. H. Scudder	12.50	17.50	5.65	5.68
—	187	♂	Watertown, Mass.	Apr. 3, '68	C. J. Maynard	12.50	17.80	5.77	5.20
—	186	♂	"	Apr. 3, '68	"	12.45	18.25	5.85	5.30
—	185	♂	"	Apr. 3, '68	"	12.80	17.53	5.57	5.43
—	—	♂	"	Mar. 28, '68	"	12.80	18.10	5.86	5.50
—	3097	♂	Ipswich, "	Aug. 28, '69	"	13.10	18.00	5.85	5.60
9768	—	♂	Evanston, Ill.	—	O. Marcy	12.40	17.75	5.62	5.07
9770	—	♂	"	—	"	12.25	17.05	5.42	4.87
2643	—	♂	Lawn Ridge, "	—	K. Butler	12.50	17.25	5.60	5.20
1401	—	♂	"	—	"	13.50	18.43	6.05	6.00
1871	—	♂	Springfield, Mass.	July 29, '62	J. A. Allen	12.48	17.25	5.50	5.00
1874	—	♂	"	July 29, '62	"	12.12	17.00	5.20	4.58
2574	—	♂	"	July 29, '62	"	12.77	17.50	5.68	5.50
1602	—	♂	"	July 12, '62	"	11.47	15.75	5.00	4.37
1873	—	♂	"	July 29, '62	"	10.90	15.45	4.75	4.10
9769	—	♂	Evanston, Ill.	—	O. Marcy	11.30	15.38	4.60	4.25
9767	—	♂	"	—	"	11.48	15.67	4.95	4.40
9598	—	♂	Waterville, Me.	—	C. E. Hamlin	11.50	16.00	5.00	4.65
2284	—	♂	"	May 3, '62	"	11.40	16.00	4.98	4.46
2271	—	♂	"	June 9, '62	"	12.05	16.30	5.20	4.85
2501	—	♂	Lynn, Mass.	—	S. Jillson	11.50	15.50	5.05	4.85

Measurements of Florida Specimens of QUISCALUS PURPUREUS.

M. C. Z. No.	Coll. No.	Sex	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.
5201	—	♂	Welaka	Feb. 6, '69	J. A. Allen	12.75	17.80	5.65	5.20
5202	—	♂	"	Feb. 6, '69	"	12.40	16.87	5.45	5.00
5264	—	♂	Hawkinsville	Feb. 18, '69	"	12.80	17.15	5.50	5.37
5251	—	♂	"	Feb. 18, '69	"	11.75	17.00	5.50	4.87
5266	—	♂	"	Feb. 18, '69	"	13.00	17.60	5.55	5.40
5267	—	♂	"	Feb. 18, '69	"	11.50	16.70	5.50	—
—	—	♂	Enterprise	Feb. 21, '69	"	12.85	16.87	—	—
5345	—	♂	"	Mar. 5, '69	"	12.30	17.38	5.50	5.25
5346	—	♂	"	Mar. 5, '69	"	12.37	16.60	5.20	5.25
10604	2583	♂	Dummitt's	Mar. 9, '69	C. J. Maynard	12.40	17.50	5.55	—
—	2344	♂	"	Feb. 26, '69	"	12.25	17.50	5.75	5.50
10602	2469	♂	"	Feb. 25, '69	"	12.50	17.30	5.50	5.55
—	2470	♂	"	Feb. 25, '69	"	12.50	17.00	5.50	5.00
10603	2471	♂	"	Feb. 25, '69	"	11.75	16.75	5.75	5.55
6848	—	♂	Cape Florida	Mar. 31, '58	G. Wurdemann	11.50	15.50	5.50	5.12
6851	—	♂	"	Apr. 10, '58	"	12.00	16.25	5.25	5.00
6852	—	♂	"	Apr. 22, '58	"	11.75	16.25	5.75	5.00
—	10335*	♂	"	Mar. 31, '58	"	11.50	16.00	5.50	—
—	10336*	♂	"	Apr. 15, '58	"	11.50	15.25	5.00	—
—	10337*	♂	"	Apr. 15, '58	"	12.00	15.50	5.00	—
—	10340*	♂	"	Apr. 22, '58	"	12.00	16.50	5.12	—
—	10341*	♂	"	Apr. 9, '58	"	11.00	15.25	5.25	—
—	10342*	♂	"	May 18, '58	"	11.75	16.25	5.00	—
—	2342	♂	"	Feb. 26, '69	"	11.50	15.50	5.00	4.77
—	2344	♂	"	Feb. 26, '69	"	11.00	15.50	5.00	4.40
10601	2468	♂	"	Feb. 26, '69	"	11.50	16.00	4.50	4.60
5263	—	♂	Hawkinsville	Feb. 18, '69	J. A. Allen	11.45	15.25	4.85	4.55
6853	—	♂	Cape Florida	Apr. 22, '58	G. Wurdemann	11.00	14.50	4.50	4.45
—	10338*	♂	"	Apr. 22, '58	"	11.12	14.50	4.75	—
—	10339*	♂	"	Mar. 31, '58	"	10.25	13.75	4.75	—

The specimens from Cape Florida are considerably smaller than those from the St. John's River; but the same difference occurs in other species between specimens from these two localities. The Cape Florida specimens of *Quiscalus purpureus* differ from others from North Florida also in having a relatively longer, slenderer, and more decurved bill, but not appreciably in color.

60.* *Quiscalus major* Vieillot. BOAT-TAILED GRACKLE.

Abundant. Particularly numerous along the St. John's River. According to Dr. Bryant they breed about the first of April. He says that about Lake Monroe some of the birds, as late as the 6th of April, had not commenced laying, "though the majority had hatched, and the young of others were almost fledged." † He notes also their sandpiper-like habit of running along the edge of the water. At Lake Dexter I observed great numbers of them walking on the floating aquatic plants.

The females of this species present very singular variations in color. Of four specimens collected at Lake Dexter, in March, one is pale ashy-

* Smith. Inst. No. Copied from Baird's Birds of North America, p. 557.

† Proc. Bost. Soc. Nat. Hist., Vol. VII, p. 9, January, 1859.

brown below, on the throat and breast nearly white, and dull dusky-brown above; while another is deep reddish-brown below and proportionally darker above, and the others are intermediate to these.

Between the two extremes there is more difference than usually obtains between valid congeneric species. The series of twenty-four males, on the other hand, are quite uniform in color, there being only a slight difference in its intensity and in the prevailing tint of the iridescence.*

The average dimensions of the thirty-three specimens of which measurements are given below are as follows:

Length (males): 16.51; alar extent, 22.48; wing, 7.19; tail, 7.00.

Length (females): 12.95; alar extent, 17.94; wing, 5.67; tail, 5.11.

The individual variation is as follows.

Males, length, 15.50 to 16.80; alar extent, 21.10 to 23.50; wing, 6.25 to 8.35; tail, 6.25 to 7.60.

Females, length, 12.10 to 13.40; alar extent, 17.25 to 18.25; wing, 5.25 to 5.95; tail, 4.75 to 5.60.

Measurements of Florida Specimens of QUISCALUS MAJOR.

M. C. Z. No.	Coll. No.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.
5272	—	♂	Blue Springs	Feb. 21, '69	J. A. Allen	16.00	22.15	7.25	6.80
5252	—	♂	Enterprise	Mar. 1, '69	"	16.25	22.50	7.15	7.15
5283	—	♂	"	Feb. 22, '69	"	16.25	21.75	7.15	7.10
5332	—	♂	"	Mar. 4, '69	"	15.50	22.00	6.85	6.70
5333	—	♂	"	Mar. 4, '69	"	15.75	22.30	7.20	6.85
5334	—	♂	"	Mar. 4, '69	"	15.60	21.85	7.00	—
5407	—	♂	Hawkinsville	Mar. 15, '69	"	16.00	23.00	7.00	7.00
5243	—	♂	"	Feb. 18, '69	"	15.75	22.25	7.30	—
5408	—	♂	"	Mar. 15, '69	"	16.50	22.50	7.30	7.15
5244	—	♂	"	Feb. 18, '69	"	17.30	23.50	7.80	—
5409	—	♂	"	Mar. 15, '69	"	16.00	22.75	7.15	7.00
5410	—	♂	"	Mar. 15, '69	"	16.00	22.75	7.20	6.90
5411	—	♂	"	Mar. 15, '69	"	16.35	22.50	7.50	—
—	—	♂	"	Mar. 15, '69	"	16.50	23.25	7.25	7.40
—	2408	♂	Dummitt's	Feb. 19, '69	C. J. Maynard	16.50	23.00	7.40	7.10
10607	2405	♂	"	Feb. 19, '69	"	16.75	23.50	7.50	7.25
—	2406	♂	"	Feb. 19, '69	"	17.50	23.00	8.35	7.60
—	—	♂	"	Mar. 17, '69	"	16.00	21.10	6.75	6.60
10610	2345	♂	"	Mar. 9, '69	"	16.90	23.00	7.70	7.50
—	2409	♂	"	Mar. 9, '69	"	16.25	22.00	7.00	7.20
—	2586	♂	"	Mar. 9, '69	"	16.75	22.00	7.00	7.00
—	2399	♂	"	Mar. 9, '69	"	15.75	22.00	6.90	7.00
—	2431	♂	"	Mar. 9, '69	"	16.00	22.25	7.25	6.75
—	2404	♂	"	Mar. 9, '69	"	16.17	20.75	6.50	6.50
—	2345	♂	"	Feb. 16, '69	"	16.50	22.30	6.25	6.25
10609	2563	♂	"	Mar. 9, '69	"	13.00	17.50	5.85	5.60
—	2343	♂	"	Mar. 9, '69	"	13.00	18.25	5.80	5.00
—	2464	♂	"	Mar. 9, '69	"	13.00	18.25	5.95	5.25
5290	—	♂	Enterprise	Feb. 25, '69	J. A. Allen	13.40	18.25	5.85	4.75
5334	—	♂	"	Mar. 4, '69	"	12.75	17.50	5.50	5.00
5412	—	♂	Lake Dexter	Mar. 23, '69	"	13.00	18.05	5.60	5.20
5413	—	♂	"	Mar. 23, '69	"	12.10	17.25	5.25	5.00
5414	—	♂	"	Mar. 23, '69	"	12.50	17.60	5.45	—

* For a very full biography of this species, see an article by Dr. Elliott Coues in the Ibis. Vol. VI, pp. 367 - 378, 1870.

The present species is hence not only remarkable for variation in size between specimens of the same sex, but especially so for its sexual variation in size, the sexual difference in this respect being greater than in any other species of insessorial bird with which I am acquainted, and it is rarely, if ever, exceeded in any group.

CORVIDÆ.

61.* *Corvus americanus* Audubon. COMMON CROW.

Corvus corone WILSON, Am. Orn., IV, 79, pl. xxv, fig. 3, 1811. — NUTTALL, Man. Orn., I, 209, 1832.

Corvus americanus AUDUBON, Orn. Biog., II, 317, 1834. — BAIRD, Birds N. Am., 566, 1858.

Corvus americanus var. *floridanus* BAIRD, Ibid., 568, 1858.

Corvus minimus GUNDLACH, Cabanis's Journal für Ornithologie, IV, 97, 1856.

Everywhere abundant.

In the average, while the general size of Florida specimens is smaller than New England ones, the bill is somewhat larger. As is well known, the crow is exceedingly variable in the size and shape of its bill even in specimens collected from the same flock. There is, however, an appreciable average difference in the size of the bill, as in general size, between northern and southern examples. This was some time since observed by Professor Baird in comparing a single specimen from the southern point of the Florida peninsula with others from the Northern States, and so strongly was he impressed by it that he thought if his Florida specimen did not represent a distinct species, it did at least a distinct variety, and as such he characterized it, calling it *Corvus americanus* var. *floridanus*. He at the same time referred to the little crow of Cuba, described by Dr. Gundlach as *Corvus minutus*, to which he said it was more nearly allied than either are to *C. americanus*. I have no examples of the latter, but from descriptions of it see no reason why it should be regarded as other than the extreme southern form of *C. americanus*.

62.* *Corvus ossifragus* Wilson. FISH CROW.

Abundant. Perhaps rather more numerous than the common crow.

63.* *Cyanurus cristatus* Swainson. BLUE JAY.

Very abundant and unsuspicious. It frequents the towns, where it seems half domestic.

The same difference occurs in this species between Florida and northern specimens in size and shape of bill as has been already pointed out in

respect to *Corvus americanus*, but it is far less marked than in *Agelaius phoeniceus*, *Quiscalus purpureus*, and *Sturnella ludoviciana*. The brilliancy of its colors seems not much greater than in New England specimens.

The difference in size between northern and southern specimens is as follows: Average of eighteen Massachusetts specimens (eleven males and seven females): Length, 11.71; alar extent, 16.87; wing, 5.13; tail, 4.89. Average of eleven Florida specimens (proportion of males and females nearly the same as in the previous case): Length, 10.98; alar extent, 15.11; wing, 4.75; tail, 5.00. The maxima and minima of the eleven males from Massachusetts are as follows: Length, 12.25 and 11.35; alar extent, 17.50 and 16.30; wing, 5.50 and 5.00; tail, 5.65 and 4.25.

Measurements of Specimens of CYANURA CRISTATA.

M. C. Z. No.	Coll. No.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.
—	34	♂	Newton, Mass.	Oct. 25, '67	C. J. Maynard	11.62	16.30	5.32	5.06
—	90	♂	" "	Feb. 5, '68	"	11.35	17.00	5.00	4.78
—	94	♂	" "	Feb. 8, '68	"	12.00	17.00	5.00	5.00
—	93	♂	" "	Feb. 8, '68	"	11.55	17.20	5.25	4.80
—	—	♂	" "	Feb. 21, '68	"	12.00	16.80	5.00	5.00
—	687	♂	" "	May 28, '68	"	12.16	17.00	5.45	5.40
—	1667	♂	" "	—	"	12.25	17.25	5.65	5.65
12393	—	♂	Springfield, "	Feb. 25, '70	Irving Allen	12.00	17.20	5.15	5.15
12392	—	♂	" "	Feb. 25, '70	"	11.50	17.00	5.30	5.10
12388	—	♂	" "	Feb. 25, '70	"	12.00	17.00	5.00	4.25
12385	—	♂	" "	Feb. 25, '70	"	12.00	17.50	5.50	5.40
12389	—	♂	" "	Feb. 25, '70	"	12.00	17.00	4.40	4.45
12392	—	♂	" "	Feb. 25, '70	"	11.00	16.50	4.33	4.80
12391	—	♂	" "	Feb. 25, '70	"	11.00	17.00	5.25	4.75
12386	—	♂	" "	Feb. 25, '70	"	11.50	17.00	5.50	5.15
—	33	♂	Newton, "	Oct. 25, '67	C. J. Maynard	11.40	16.32	5.30	5.30
4875	688	♂	" "	May 28, '68	"	11.62	16.53	4.75	4.77
—	1685	♂	" "	Nov. 4, '68	"	11.75	16.00	5.20	4.35
10733	1951	♂	Jacksonville, Fla.	Jan. 2, '69	"	11.15	16.00	5.00	5.00
10734	1973	♂	" "	Jan. 3, '69	"	11.00	15.50	4.80	4.80
10734	1974	♂	" "	Jan. 3, '69	"	11.00	14.75	4.00	4.80
5522	—	♂	Blue Springs, "	Feb. 21, '69	J. A. Allen	10.75	15.75	4.20	—
5128	—	♂	Jacksonville, "	Jan. 21, '69	"	10.75	15.50	4.70	5.12
5190	—	♂	Welaka, "	Feb. 3, '69	"	10.70	15.60	5.10	5.10
—	—	♂	Enterprise, "	Mar. 1, '69	"	11.00	15.75	5.00	—
—	—	♂	" "	Mar. 4, '69	"	10.70	15.15	4.50	—
5348	—	♂	" "	Mar. 4, '69	"	11.00	16.00	5.00	5.05
5162	—	♂	Hibernia, "	Jan. 30, '69	"	11.25	15.75	5.00	5.15
5163	—	♂	" "	Jan. 30, '69	"	11.50	15.50	5.00	—

64.* *Cyanocitta floridana* Bonaparte. FLORIDA JAY.

Corvus floridanus BARTRAM, Travels, 291, 1791. — AUDUBON, Orn. Biog., I, 444, pl. lxxxvii, 1831.

Garrulus floridanus BONAP., Am. Orn., II, 11, pl. ix, 1828.

Garrulus caerulescens ORD, Journ. Phil. Acad. Nat. Sci., I, 347, 1818.

Garrulus californicus VIGORS, Zool. Beechey's Voyage, 21, pl. v, 1839.

Cyanocitta floridana BONAP., Consp. Gen. Avium, 377, 1850.

Cyanocitta superciliosa STRICKLAND, Ann. & Mag. Nat. Hist., XV, 260, 1845.

Cyanocitta californica STRICKLAND, Ibid., 342.

Cyanocitta Woodhousei BAIRD, Birds N. Am., 585, 1858.

Numerous in the scrub, but does not appear to frequent the pine woods the hummocks or swamps. I saw none along the St. John's, except at Blue Springs, but they occur in numbers a few miles back from the river.

On comparing a number of specimens of the so-called *Cyanocitta californica* with numerous others from Florida, I find, as previous writers have observed, that the differences between them are very slight, and not so great as obtain between Florida and New England specimens of *Pipilo erythrophthalmus*, *Agetæus phæniceus*, and other species where there is no reason to question their specific identity. The so-called *C. Woodhousei* is described as, and is, intermediate in character between *C. floridana* and *C. californica*. The habitat of *C. Woodhousei* is also intermediate between those of the other two, but adjoins that of *C. californica*, to which it is most nearly allied. How great the interval is between the habitats of *C. floridana* and *C. Woodhousei* I have not been able to accurately determine. Bonaparte* reported the former as being found in Louisiana and northward to Kentucky, and the latter occurs in Western Texas.

In the following measurements of twelve specimens of this species (six males and six females) the extremes are as follows: Length, 11.00 and 12.50 (both specimens being females); alar extent, 13.50 (female) and 15.00 (male); wing, 4.00 and 4.75 (both specimens females); tail, 4.25 and 5.35 (both specimens females). The average dimensions of these specimens are as follows: Length, 11.74; alar extent, 14.44; wing, 4.42; tail, 4.80. The females average slightly smaller than the males.

Measurements of Florida Specimens of CYANOCITTA FLORIDANA.

M. C Z No.	Coll. No.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.
10739	2480	♂	Dummitt's	Feb. 22, '69	C. J. Maynard	11.50	14.50	4.30	4.35
—	2377	♂	"	Feb. 22, '69	"	12.00	15.00	4.45	4.75
—	2421	♂	"	Feb. 15, '69	"	12.00	15.00	4.75	5.00
10736	2326	♂	"	Feb. 15, '69	"	12.00	14.50	4.50	4.60
—	2329	♂	"	Feb. 15, '69	"	11.50	14.25	4.50	4.25
—	2379	♂	"	Feb. 22, '69	"	11.50	14.25	4.50	5.35
10737	2328	♂	"	Feb. 15, '69	"	12.50	14.50	4.75	4.90
—	2378	♂	"	Feb. 15, '69	"	11.50	14.10	4.30	5.15
—	2375	♂	"	Feb. 15, '69	"	11.60	14.40	4.60	4.25
5271	—	♂	"	Feb. 21, '69	J. A. Allen	11.00	13.50	4.00	5.35
5272	—	♂	"	Feb. 21, '69	"	12.00	14.50	4.30	4.75
5523	—	—	"	Feb. 21, '69	"	11.75	14.80	4.20	—

TYRANIDÆ.

65.† *Sayornis fuscus* Baird. PEWEE.

Abundant all winter, and a few remain till into April.

The king-bird (*Tyrannus carolinensis*), the great-crested flycatcher (*Myiarchus crinitus*), and the wood pewee (*Contopus virens*) became

* Am. Orn., Vol. II, p. 60, 1828.

common the last week in March, as also, according to Mr. Boardman, the least flycatcher (*Empidonax minimus*).

Several specimens of the gray king-bird (*Tyrannus dominicensis*) were obtained by Mr. L. L. Thaxter at St. Augustine, about the first of May.

ALCEDINIDÆ.

66.* *Ceryle alcyon* Boie. KINGFISHER.

Abundant. As shy and distrustful here as in the more thickly settled parts of the country. Begins to breed very early. Mr. Maynard saw them forming their holes in the coquina rock, in the banks of the canal connecting Indian River with Mosquito Lagoon, the first week in February.

CAPRIMULGIDÆ.

67.* *Antrostomus carolinensis* Gould. CHUCKWILL'S WIDOW.

Abundant. Not observed till about the first of March, when its notes are usually first heard. Said by Audubon to be resident; which statement is confirmed by the testimony of old residents of the State.

68.* *Antrostomus vociferus* Bonaparte. WHIPPOORWILL.

Apparently not numerous in winter. I heard it once in February, and Mr. Maynard took it at Dummitt's in the same month. The inhabitants along the St. John's agree with Audubon that this species is also a winter resident.

The night hawk (*Chordeiles popetue* * Baird) was collected at Jacksonville by Mr. Thurston as early as April 20th.

* *Caprimulgus virginianus* BRISSON, Orn., II, 477 (in part).

Caprimulgus popetue VIEILLOT, Ois. Am. Sept., I, 56, pl. liv, 1807.

Caprimulgus americanus WILSON, Am. Orn., V, 65, pl. cxl, 1812.

Caprimulgus (Chordeiles) virginianus SWAIN., Faun. Bor. Am., II, 62, 1831.

Chordeiles virginianus BON., Geog. & Comp. List, 8, 1838. — GOSSE, Birds of Jamaica, 33, 1847.

Chordeiles sapiti BONAP., Consp. Gen. Avium, I, 63, 1849. — CASSIN, Ill. N. Am. Birds, 238, 1855.

Chordeiles brasilianus LAW., Ann. N. Y. Lyceum Nat. Hist., V, 114, 1851.

Chordeiles Henryi CASSIN, Ill. N. Am. Birds, 239. — BAIRD, Birds N. Am., 153.

Chordeiles Gundlachii LAW., Ann. N. Y. Lyc. Nat. Hist., VI, 167, 1856.

Chordeiles tezensis LAW., Ibid., 165. — BAIRD, Birds N. Am., 154.

Chordeiles minor CABANIS, Journ. für Orn., 5, 1856.

Chordeiles popetue BAIRD, Birds N. Am., 151.

This widely distributed species presents only the usual variations in size and color

CYPSELIDÆ.

The chimney swift (*Chaetura pelasgia*) arrives about the last week in March. It was common at Jacksonville, April 1st.

TROCHILIDÆ.

The ruby-throated humming-bird (*Trochilus colubris*) became common about March 1st. Some probably spend the winter in South Florida.

PICIDÆ.

69.* *Campephilus principalis* Gray. IVORY-BILLED WOODPECKER.

Picus principalis LINNÉ, Syst. Nat., I, 173, 1767.

Campephilus principalis GRAY, Genera of Birds, 1840.

Campephilus Bairdii CASSIN, Proc. Phil. Acad. Nat. Sci., 1863, 322. (West Indian form.)

Rather rare; at least far less numerous than most of the other species of woodpecker.

With only Florida specimens of this species before me, I am unable to give comparisons between them and specimens from other localities. According to the late Mr. Cassin, those found in Cuba differ from those of the Southern States, in being smaller, as would be expected, with very slight deviations in color-markings. He has, however, given to the Cuba race the name of *Campephilus Bairdii*, remarking that it appears to be "one of those singular insular species which have become well known to naturalists."

Measurements of Florida Specimens of CAMPEPHILUS PRINCIPALIS.

M. C. Z. No.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.
5221	♂	Volusia.	Feb. 12, '69	J. A. Allen	20.00	32.25	10.40	6.90
5222	♂	"	Feb. 12, '69	"	19.50	32.50	10.25	6.90
5229	♀	"	Feb. 12, '69	"	19.30	31.50	10.60	6.85
5354	♂	Enterprise	Mar. 5, '69	"	19.25	30.50	9.70	6.40
5399	♀	Hawkinsville	Mar. 15, '69	"	19.50	31.50	10.25	6.75

seen in other species of our birds. Yet these variations have in the present case been mistaken as indicating numerous species. The southern representatives of it are appreciably smaller than the northern, and have the white markings on the wings more restricted, — variations that have already been pointed out in this paper as occurring in numerous others similarly distributed. Those from the central arid region of the continent are also lighter in general color than those from the eastern or western portions; also a common color variation in other species. The latter type forms the so-called *Chordeiles Henryi*; the southern ones have been variously characterized as *C. sapiti*, *C. texensis*, *C. Gundlachii*, etc., as indicated in the above-cited synonymes.

70.* *Hylotomus pileatus* Baird. PILEATED WOODPECKER.

Abundant. Much smaller than at the north, but not otherwise appreciably different.

The average dimensions of fourteen Florida specimens (seven males and seven females) are as follows:—

Males, length, 17.48; alar extent, 28.07; wing, 9.21; tail, 6.82.

Females, length, 16.44; alar extent, 26.80; wing, 8.98; tail, 6.54.

The individual variation is as follows:—

Males, length, 17.25 to 17.75; alar extent, 27.50 to 28.50; wing, 9.00 to 9.50; tail, 6.20 to 6.75.

Females, length, 15.50 to 16.80; alar extent, 26.00 to 27.75; wing, 8.50 to 9.50; tail, 5.85 to 6.80.

Measurements of Florida Specimens of HYLOTOMUS PILEATUS.

M. C. Z. No.	Coll. No.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.
5118	—	♂	Hibernia	Jan. 30, '69	J. A. Allen	17.75	28.25	9.25	6.65
5203	—	♂	Welaka	Feb. 7, '69	"	17.25	28.00	9.00	6.50
5215	—	♂	"	Feb. 10, '69	"	17.50	28.50	9.25	6.75
—	—	♂	Hawkinsville	Mar. 10, '69	"	17.25	27.50	9.50	—
—	1937	♂	Jacksonville	Dec. 31, '69	C. J. Maynard	17.75	28.50	9.50	6.40
—	2076	♂	"	Jan. —, '69	"	17.25	27.75	9.00	6.20
—	2543	♂	Dummitt's	Feb. 15, '69	"	17.60	28.00	9.00	6.45
—	2334	♂	"	Mar. 11, '69	"	15.50	26.40	8.70	5.85
—	2602	♂	"	Mar. 5, '69	"	16.60	27.75	9.00	6.75
5204	—	♀	Welaka	Feb. 7, '69	J. A. Allen	16.75	26.25	8.50	6.75
5214	—	♀	"	Feb. 10, '69	"	16.35	26.75	9.15	6.60
5216	—	♀	"	Feb. 10, '69	"	16.30	27.25	9.00	6.80
5274	—	♀	Blue Springs	Feb. 21, '69	"	16.75	27.20	9.50	6.50
—	—	♀	Hawkinsville	Mar. 10, '69	"	16.80	26.00	9.00	—

71.* *Picus villosus* Linné. HAIRY WOODPECKER.

Picus villosus LINNÉ, Syst. Nat., I, 175, 1767. — FORSTER, Philosoph. Transact., LXII, 383, 1772. — WILSON, Am. Orn., I, 150, pl. ix, fig. 3, 1808. — AUDUBON, Orn. Biog., V, 164, pl. ccccxvii, 1837. (Northern form.)

Picus leucomelanus WAGLER, Syst. Av., No. 18, 1827. (Immature male.)

Picus Auduboni SWAINSON, Faun. Bor. Am., II, 306, 1831. (Immature male.)

— TRUDEAU, Journ. Phil. Acad. Nat. Sci., 404, 1837. (Immature male). —

AUDUBON, Orn. Biog., V, 194, 1839. (Same as the last.)

Picus Martinæ AUDUBON, Ibid., 181, pl. ccccxvii. (Very immature.)

Picus Phillipsii AUDUBON, Ibid., 186, pl. ccccxvii. (Immature.)

Picus Harrisii AUDUBON, Ibid., 191, same plate. (Northwestern form.) — BAIRD, Birds N. Am., 87.

Picus septentrionalis NUTTALL, Man. Orn., I (2d Ed.), 685, 1840.

Picus rubricapillus NUTTALL, Ibid., 684. (Immature male.)

Picus Cuvieri MALHERBE, Mon. Picidæ, I, 85, pl. xxii, fig. 3. (Young female.)

Picus Jardinei MALHERBE, Ibid., I, 85, pl. xxv, fig. 4, 5. — CASSIN, Proc. Phil. Acad. Nat. Sci., 1863, 201.

Not numerous in Florida in comparison with the other species of *Picidæ*.

The difference in size between northern and southern specimens of all the species of the *Picidæ* is greater than obtains in most other families of birds. So great is it in *Picus villosus* and *Picus pubescens* that it was in these species that such variations were first noticed. This difference is well pointed out by Professor Baird in his work on the North American Birds, and fully demonstrated in his table of measurements. On this ground he distinguished three varieties of *P. villosus*, — *P. villosus major*, occupying the northern and western portions of the continent; *P. villosus medius*, occupying the Middle States; and *P. villosus minor*, occupying the Southern States. Audubon regarded the two former as distinct species. In addition to these variations in size, my Florida specimens indicate a well-marked variation in color between the northern and extreme southern races, the Florida specimens differing from New England ones in having the white markings of relatively less extent, which gives to the plumage a considerably darker aspect. Through this variation there is an approach in the Florida examples of *P. villosus* to the so-called *P. Harrisii* of the Pacific coast and Rocky Mountain regions of the continent, and in the Florida examples of *P. pubescens* to the so-called *P. Gairdneri*, also of the middle and western regions of the continent. These, as is well known, differ respectively from *P. villosus* and *P. pubescens* almost solely in a general darker aspect, resulting simply from the relatively greater predominance of the black color of the plumage over the white markings in the western type; there being no change whatever in the general style of coloration, though some of the smaller white spots seen in the eastern are entirely obsolete in the western type. Under *Picus Gairdneri* Professor Baird thus describes these variations. "There is," he says, "the same series in specimens of *Picus Gairdneri* that were indicated under *P. Harrisii*. Thus the most northern from Washington Territory and Oregon have the under parts more brown, with faint black streaks, the white spots above smaller and less numerous. In specimens from California and farther east the white is purer, the spots more conspicuous." "The almost perfect parallelism," he further observes, "with appreciable differences between the markings of the northwestern and southeastern varieties of *Picus Harrisii* and *Gairdneri*, and their relationship to *P. villosus* and *pubescens*, is a remarkable fact in American ornithology, and may possibly indicate the necessity either of dividing the dark ones into a Pacific and Rocky Mountain series, or of considering all as variations of two species, a larger [*P. villosus*] and a smaller [*P. pubescens*], changing their character with longitudinal distribution." And he aptly adds, "Many other supposed species are involved in the

same consideration." * Professor Baird in his account of these species, expressly refers to California specimens that have less white on the wings than the one form and more white than the other.† This with the color differences existing between Florida specimens and New England ones, similar in character to these, though less in degree, seems to confirm the necessity alluded to by Professor Baird of regarding the small spotted woodpeckers in question as forming only two species, — the *Picus villosus* and *Picus pubescens*, — with parallel and remarkable geographical variations. So great is the difference, however, between typical representatives of the two leading forms of each, that their discoverers, with too few specimens of each to enable them to detect the gradual passage of the one into the other, — a fact which now seems well substantiated, — were quite excusable in regarding them as distinct species. Several other supposed species, as indicated by the synonymes given above, and previously by other authors, have been based on phases of immaturity. The young of either sex often have the crown spotted with red or yellow, while the mature male alone has red on the head, and in which it is usually confined to a narrow occipital transverse band. In respect to the number, shape, position, and size of the white spots on the wings, however, there is always considerable variation in specimens from the same locality, these variations being dependent upon neither sex nor age.

Florida specimens of not only *Picus pubescens* and *P. villosus*, but of *Centurus carolinus*, *Sitta carolinensis*, and *Sitta pusilla*, often have the plumage of the lower surface of the body so much soiled and darkened by running over the blackened trees in recently burnt districts as to materially alter their appearance, so that they might almost be taken for distinct species, as previously noted by Audubon.‡

72.* *Picus pubescens* Linné. DOWNY WOODPECKER.

Picus pubescens LINNÉ, Syst. Nat., I, 175, 1766. — WILSON, AUDUBON, BONAPARTE, NUTTALL, BAIRD, CASSIN, etc.

Picus (Dendrocopus) pubescens SWAINSON, Faun. Bor. Am., II, 307, 1831.

Picus (Dendrocopus) medianus SWAINSON, Ibid., 308. (Described from New Jersey specimens).

* Birds of North America, p. 91.

† In accounting for these intermediate forms, Mr. Cassin adopts the very convenient but, as it seems to me, uncalled-for and incorrect theory of hybridity, so often resorted to in similar cases. Under *Picus villosus*, he says that *P. villosus* and *P. Harrisii* probably associate in a region intermediate between the proper ranges of the two species, "and produce hybrids, which present difficulties to naturalists." Under *Picus pubescens* he makes similar remarks in respect to *P. pubescens* and *P. Gairdneri*. *Proc. Phil. Acad. Nat. Sci.*, 1863, pp. 200, 201.

‡ Orn. Biog., Vol. II, p. 82.

Picus (Dendrocopus) meridionalis SWAINSON, Ibid. (Southern race.)

Picus Gairdneri AUDUBON, Orn. Biog., V, 317, 1839. (Northwestern form.) — BAIRD, Birds N. Am., 91, 1858.

Picus meridionalis NUTTALL, Man. Orn., I, (2d Ed.) 690, 1840. (Not of Swainson).

Picus Lecontei JONES, Ann. N. York Lyc. Nat. Hist., IV, 489, pl. xviii, 1848. (Three-toed specimen.)

Picus Turati MALHERBE, Mon. Pic., I, 125, pl. xxix, fig. 5, 6. — CASSIN, Proc. Phil. Acad. Nat. Sci., 1863, 202.

Common. Much more numerous than *Picus villosus*.

The difference in size and color between northern and southern specimens has been sufficiently detailed under the previous species.

73.* *Picus borealis* Vieillot. RED-COCKADED WOODPECKER.

Picus borealis VIEILLOT, Ois. Am. Sept., II, 66, pl. cxxii, 1807. — CASSIN, Proc. Phil. Acad. Nat. Sci., 1863, 203.

Picus querulus WILSON, Am. Orn., II, 103, pl. xv, fig. 1, 1810. — CASSIN, Proc. Phil. Acad. Nat. Sci., 1863, 203.

Common in the pineries.

Mr. Cassin regards the Carolina and Georgia representatives of this species as specifically distinct from the Pennsylvania ones. He says that they are as distinct and as easily recognized as are *Picus villosus* and *P. Harrisii*, which he of course regards as valid species. He assigns Vieillot's

Measurements of Florida Specimens of PICUS BOREALIS.

M. C. Z. No.	Coll. No.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.
10641	1919	♂	Jacksonville	Dec. 31, '68	C. J. Maynard	8.40	14.20	4.75	3.52
10642	1920	♂	"	Dec. 31, '68	"	8.30	14.20	4.76	3.62
10643	1921	♂	"	Dec. 31, '68	"	8.30	14.80	4.80	3.56
—	1922	♂	"	Dec. 31, '68	"	8.50	14.50	4.75	3.69
10644	1923	♂	"	Dec. 31, '68	"	8.20	14.45	4.75	3.39
—	1924	♂	"	Dec. 31, '68	"	8.50	15.00	4.80	3.32
10645	1925	♂	"	Dec. 31, '66	"	8.50	14.75	4.85	3.60
—	1971	♂	"	Jan. 3, '69	"	8.50	15.00	4.85	3.50
10646	1972	♂	"	Jan. 3, '69	"	8.50	14.30	4.75	3.75
10631	29	♂	"	Apr. 11, '69	"	8.00	14.75	4.90	3.45
10632	30	♂	"	Apr. 6, '69	"	8.50	15.00	4.90	3.35
10633	31	♂	"	Apr. 6, '69	"	8.30	14.90	4.85	3.35
10634	41	♂	"	Apr. 7, '69	"	8.15	14.50	4.70	3.25
10637	47	♂	"	Apr. 8, '69	"	8.60	15.15	4.87	3.40
10638	48	♂	"	Apr. 8, '69	"	8.50	15.00	4.95	3.46
10639	58	♂	"	Apr. 13, '69	"	8.50	14.10	4.75	3.59
—	49	♂	"	Apr. 8, '69	"	8.50	14.15	4.85	3.49
10640	59	♂	"	Apr. 12, '69	"	8.50	15.00	4.80	3.50
10636	44	♂	"	Apr. 7, '69	"	8.30	15.00	4.80	3.60
10635	43	♂	"	Apr. 7, '69	"	8.35	14.60	4.60	3.60
—	32	♂	"	Apr. 3, '69	"	8.30	14.90	4.85	3.50
—	42	♂	"	Apr. 7, '69	"	8.20	14.70	4.75	3.29
5116	—	♂	"	Jan. 19, '69	J. A. Allen	8.50	15.20	4.40	3.30
5137	—	♂	"	Jan. 25, '69	"	8.38	14.75	4.57	3.42
5375	—	♂	Hawkinsville	Mar. 12, '69	"	8.55	14.55	4.50	3.40
5393	—	♂	"	Mar. 15, '69	"	8.50	14.50	4.45	3.20
5394	—	♂	"	Mar. 15, '69	"	8.25	14.50	4.40	3.15
5414	—	♂	Volusia	Mar. 25, '69	"	7.90	14.60	4.45	3.25

name *borealis* to the Pennsylvania type, and Wilson's name *querulus* to the more southern form. In recognizing two species of red-cockaded woodpecker in the Atlantic States, Mr. Cassin differs from all previous writers. Having only Florida specimens, a series of twenty-two, before me, I cannot state from personal observation as to how they differ from northern ones. They appear, however, to be merely a little smaller and darker.

The average size of the twenty-eight Florida specimens of which measurements are given in the foregoing table is as follows: Length, 8.34; alar extent, 14.46; wing, 4.71; tail, 3.41.

74.† *Sphyrapicus varius* Baird. YELLOW-BELLIED WOODPECKER.
Common.

75.* *Centurus carolinus* Bonaparte. RED-BELLIED WOODPECKER.
Picus carolinus LINNÉ, Syst. Nat., I, 174, 1767.
Picus griseus VIEILLOT, Ois. Am. Sept., II, 52, pl. cxvi, 1807.
Centurus carolinus BONAP., Geog. & Comp. List, 40, 1838.

Abundant. The most numerous species of its family in Florida. Specimens in the Museum from Cape Florida, taken the 8th of May by Mr. G. Wurdemann, indicate it as resident throughout Florida, though considered by Audubon and others as only a winter visitant to this and the other Gulf States.

The Florida specimens are all very much brighter colored than others before me from Maryland, Indiana, Illinois, and Michigan, the Michigan specimens being the palest. Professor Baird has remarked, in regard to a specimen from Amelia Island, Florida,* that it was not only very much smaller than northern ones, but had the white transverse bands on the back much narrower, the black ones being three times the breadth of the white ones, instead of twice, as in the northern specimens. These differences my large series from the St. John's River indicate as constant. A similar increase in the breadth of the black bands over the white ones in southern specimens as compared with northern ones, in species banded transversely, is seen in numerous other species. It is well marked in *Colaptes auratus* (where the bands are dark and light brown), in *Sphyrapicus varius*, and, as I shall show more fully subsequently, in *Ortyx virginianus*. The extent and intensity of the red on the abdomen and head, and especially its brilliancy on the head, is much greater in the Florida specimens of *C. carolinus*. In this respect there is also a well-marked difference between Cape Florida specimens and those from the St. John's River, the Cape Florida ones being much the brighter. These seem to accord in every particular with

* Birds of North Amer., p. 109.

the so-called *Centurus subelegans* of Lower California and Mexico. It is interesting to note that variations in color occur between the northern and southern representatives of *Centurus flaviventris* similar to those exhibited by northern and southern examples of *C. carolinus*. The southern forms of *C. flaviventris* were long since characterized by Wagler, Swainson, and Bonaparte as specifically distinct from the northern, under the names of *C. elegans*, *C. santacruzi*, etc., etc., which many authors still rank as species.

76.* **Melanerpes erythrocephalus**, Swainson. RED-HEADED WOOD-PECKER.

Rare in winter; said to be common in summer. I saw two only, about March 15th. Mr. Boardman also gives it as rare, while Mr. Maynard did not meet with it at all. Audubon speaks of its being very abundant in winter in Louisiana, and Dr. Coues gives it as resident in South Carolina; but it is certainly not common in winter in East Florida.

77.* **Colaptes auratus** Swainson. GOLDEN-WINGED WOODPECKER.

Abundant.

Considerably smaller than at the north, with the colors much more intense, and the transverse black bars on the back relatively broader. The individual variations in this species, even at the same locality, are very considerable, especially in respect to the bill. Figures 5 and 6, Plate VIII, illustrate the variation in the form and size of the bill of two specimens from Massachusetts, both of which are females.

The following summary of the subjoined tables indicates the difference in size between Massachusetts and Florida specimens, and the individual differentiation in the same respect at each locality. The sexes seem not to differ essentially in size.

No. of Specimens.	Sex.	Locality.		Length.	Alar Extent.	Wing.	Tail.
18	—	Massachusetts.	Average	12.45	19.94	6.24	4.35
11	—	Florida	Average	11.66	18.82	5.84	4.40
18	—	Massachusetts.	Maximum	13.00	20.75	6.60	4.70
18	—	"	Minimum	12.00	19.00	6.00	4.00
11	—	Florida.	Maximum	12.75	19.75	6.25	4.85
11	—	"	Minimum	10.60	17.60	5.60	4.10

While the Florida specimens are considerably smaller than the northern in three of the measurements, the tail is actually longer in the Florida birds, and hence relatively much longer. In most of the species of which comparative tables of measurements are given in the present paper, there is a decided tendency to an elongation of the tail at the southward, the tail decreasing less in length than the wing or the general size.

Measurements of Massachusetts Specimens of COLAPTES AURATUS.

M. C. Z. No.	Coll. No.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.
—	29	♂	Watertown	Oct. 14, '67	C. J. Maynard	12.50	20.15	6.00	4.15
—	232	♂	"	Mar. 17, '68	"	12.60	20.00	6.31	4.56
—	252	♂	Waltham	Apr. 18, '68	"	13.00	20.75	6.60	4.60
—	281	♂	Newton	Apr. 21, '68	"	12.30	19.60	6.00	4.24
4880	280	♂	"	Apr. 21, '68	"	12.25	19.90	6.25	4.62
—	325	♂	"	Apr. 25, '68	"	12.55	20.05	6.27	4.35
—	—	♂	"	Aug. 5, '68	"	12.67	19.80	6.10	4.00
4881	356	♂	"	May 1, '68	"	12.00	19.00	6.10	4.10
—	958	♂	"	June 2, '68	"	12.00	19.45	6.10	4.25
5460	1011	♂	Waltham	Aug. 6, '68	"	12.50	19.90	6.20	4.05
—	2902	♂	Newton	June 12, '69	"	12.50	20.50	6.40	4.30
—	4028	♂	Waltham	Aug. 22, '69	"	12.50	20.00	6.25	4.70
—	4029	♂	Newton	Aug. 22, '69	"	13.00	20.60	6.25	4.70
—	4034	♂	"	Aug. 26, '69	"	12.50	20.27	6.25	4.35
—	2913	♂	"	June 30, '69	"	12.10	20.00	6.45	4.55
—	2916	♂	"	June 22, '69	"	12.00	19.50	6.15	4.25
—	2915	♂	"	June 22, '69	"	12.50	19.50	6.25	4.20
—	2939	♂	"	May 1, '69	"	12.50	20.00	6.30	4.45

Measurements of Florida Specimens of COLAPTES AURATUS.

M. C. Z. No.	Coll. No.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.
—	2075	♂	Jacksonville	Jan. 20, '69	C. J. Maynard	10.75	18.50	5.75	4.55
—	2074	♂	"	Jan. 20, '69	"	11.00	17.60	5.50	4.50
10612	2346	♂	Dummitt's	Feb. 16, '69	"	11.75	19.50	6.25	4.10
10614	2601	♂	"	Mar. 11, '69	"	12.00	19.00	5.90	4.60
10611	2584	♂	"	Mar. 9, '69	"	10.60	17.75	5.70	4.30
10613	2542	♂	"	Mar. 5, '69	"	12.75	19.10	6.00	4.85
10610	2335	♂	"	Mar. 5, '69	"	12.00	19.20	6.00	4.25
5196	—	♂	Welaka	Feb. 5, '69	J. A. Allen	12.20	19.10	5.85	4.30
5321	—	♂	Enterprise	Mar. 1, '69	"	11.50	18.75	5.60	4.15
—	—	♂	Volusia	Mar. 25, '69	"	12.25	19.75	6.00	—
—	—	♂	"	Mar. 25, '69	"	11.50	18.75	5.65	—

Of the eight species of woodpecker mentioned above as occurring in Florida in winter, all but one or two (*Melanerpes erythrocephalus* and *Campephilus principalis*) are numerously represented. Most of them are exceedingly abundant, the woodpeckers hence forming a conspicuous element in the bird-fauna of East Florida. All of them are resident, according to Dr. Coues, in South Carolina. Audubon, however, states that two of them (*Sphyrapicus varius*, *Centurus carolinus*) do not breed south of Maryland, but Dr. Coues gives them as resident the whole year in South Carolina.

PSITTACIDÆ.

78. *Conurus carolinensis* Bonaparte. CAROLINA PAROKEET.

Common. Hundreds are captured every winter on the Lower St. John's by professional bird-catchers and sent to the northern cities. Thousands of others are destroyed wantonly by sportsmen. Concerning

this needless slaughter Mr. Boardman thus writes: "The little parokeet must soon be exterminated. Some of our Enterprise party would sometimes shoot forty or fifty at a few discharges, for sport, as they hover about when any are shot until the whole flock is destroyed." From its habit of feeding upon the tender maize in autumn, it is sometimes somewhat injurious to the farmer, and for this cause many are also killed. It is also more or less hunted as a game-bird. It is well known that the parokeet formerly inhabited large portions of the United States where it is now never seen, and the cause of its disappearance has been deemed a mystery. Such facts as these, however, seem to render clear what its ultimate fate must be in the United States, — extermination.

I could learn nothing from the inhabitants in regard to the time, manner, or place of breeding of this species, even old residents professing total ignorance in regard to these points.

The following table of measurements of specimens of this species serves to indicate its average size and proportions in Florida. In mature specimens the sexual difference in color and size is very slight. Neither sex acquires its adult colors before the second or third year.

The average size of the nineteen specimens (six males and thirteen females) cited below is as follows: Length, 13.10; alar extent, 21.76; wing, 7.59.

The extremes are as follows: —

Length, 12.50 and 13.60 (both specimens females); alar extent, 21.10 (female) and 22.50 (male); wing 7.00 and 7.85. These specimens seem to indicate a tolerable constancy in general size and proportions.

Measurements of Florida Specimens of CONURUS CAROLINENSIS.

M. C. Z. No.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.
5205	♂	Welaka	Feb. 8, '69	J. A. Allen	13.25	22.00	7.70	6.05
5206	♀	"	Feb. 8, '69	"	13.55	22.30	7.85	6.75
5207	♀	"	Feb. 8, '69	"	12.90	21.50	7.45	6.10
5225	♀	Volusia	Feb. 12, '69	"	13.00	21.75	7.00	5.80
5226	♀	"	Feb. 12, '69	"	13.00	21.60	7.35	5.80
5227	♀	"	Feb. 12, '69	"	13.00	21.75	7.30	6.00
5228	♀	"	Feb. 12, '69	"	13.00	21.50	7.50	6.00
5294	♀	Enterprise	Feb. 25, '69	"	13.25	21.50	7.40	—
5295	♀	"	Feb. 25, '69	"	13.00	22.45	7.60	6.00
5296	♀	"	Feb. 25, '69	"	13.60	22.00	7.34	6.60
5297	♀	"	Feb. 25, '69	"	13.45	22.00	7.50	—
—	♀	Hawkinsville	Mar. 13, '69	"	13.25	22.50	7.75	—
—	♀	"	Mar. 13, '69	"	13.15	21.25	7.50	—
—	♀	"	Mar. 13, '69	"	12.50	21.35	7.30	—
—	♀	Orange Bluffs	Mar. 24, '69	"	12.85	21.75	7.40	—
—	♀	"	Mar. 24, '69	"	13.60	22.30	7.75	—
—	♀	"	Mar. 24, '69	"	13.05	21.10	7.50	—
—	♀	"	Mar. 24, '69	"	13.25	21.30	7.50	—
—	♀	"	Mar. 24, '69	"	13.25	21.50	7.55	—

VULTURIDÆ.

79.* *Cathartes aura* Illiger. TURKEY VULTURE.

Vultur brasiliensis BRISSON, Orn., I, 468, 1760.

Vultur aura LINNÉ, Syst. Nat., I, 122, 1767. — VIEILLOT, Ois. Am. Sept., I, 25, pl. 2 bis, 1807. — WILSON, Am. Orn., IX, pl. lxiv, fig. 1, 1814.

Cathartes aura ILLIGER, Prodrum, 283, 1811. — BONAPARTE, Ann. N. Y. Lyc. Nat. Hist., II, 23, 1828. — AUDUBON, Orn. Biog., II, 296, pl. clii, 1835. — BONAPARTE, Geog. and Comp. List, I, 1838. — D'ORBIGNY, Voy. dans l'Amer. Merid., IV, iii, 38, 1844. — CASSIN, Proc. Phil. Acad. Nat. Sci., 1849, 159. — BONAPARTE, Consp. Gen. Av., I, 9, 1850.

Vultur jota MOLINA, Saggio sul stor. nat. del Chile, 1782.

Cathartes ruficollis SPIX, Av. Spec. Novæ, 2, 1824.

Vultur jota MOLINA, Sagg. sul stor. nat. del Chile, 235, 1782. — GMELIN, Syst. Nat., I, 347, 1788.

Cathartes jota BONAPARTE, Consp. Gen. Av., I, 9, 1850. — CASSIN, U. S. Nav. Astr. Exp., II, 172, 1855.

Cathartes septentrionalis PR. MAXIMILIAN, Reise in das Nord-Amer., I, 162, 1839.

? *Cathartes Burrovianus* CASSIN, Proc. Phil. Acad. Nat. Sciences, 1843, 212. — ? CASSIN, Baird's Birds of N. Am., 6, 1858.

Abundant. Collect in large companies about the dead alligators so numerous in the St. John's River.

Both this species and the following (*Carthartes atratus*) paid us frequent visits at our camps at Enterprise and Hawkinsville, and whenever we left them they did not fail to gather up and devour the carcasses of the birds and mammals thrown away by us after skinning. We found them, in fact, rather troublesome neighbors, since on more than one occasion they proceeded, in our absence, to investigate the character of the specimens we had left in the sun to dry, and in a manner so unsatisfactory to ourselves that one of the party was frequently obliged to stay in camp to protect them while the others were away collecting.

Both this and the following species were represented as breeding late in the season, and as frequenting the palmetto swamps as well as some of the islands above Enterprise for this purpose.

The synonymy here given of the present and following species indicates clearly the confusion which several continental European authors have introduced through their descriptions of these species, to which attention has been previously called by Mr. Cassin.* While a *Vultur* (or *Cathartes*) *aura* has been described by most authors who have written of the two species in question, the name *aura* has been applied sometimes to the one and

* Proc. Phil. Acad. Nat. Sci., 1849, 159.

sometimes to the other, but when given to the true *aura* of Linné, Vieillot, and Wilson, the *atratus* of Bartram and Wilson has been cited as a synonyme, and the true *atratus* described under a new name. The name *jota* has likewise been repeatedly applied to both species by different authors, and in some cases even by the same author, as has been also the name *brasiliensis*. The description given by Linné in the twelfth edition of his *Systema Naturæ*, under *V. aura*, clearly refers to the *V. aura* of Wilson, of which the *V. jota* of Molina and Gmelin are synonyms; although some of Linné's synonyms may refer to the *C. atratus* of modern writers. Bonaparte, however, in both his *Synopsis of the Birds of the United States* and in his *Geographical and Comparative List*, strangely applied the name *jota* to the *atratus* of Wilson, in which he was for a time followed by other writers. By those who have regarded the South American representatives of *C. aura* as distinct from its North American ones, the name *jota* has latterly been applied to the supposed distinct South American representative of the supposed true or northern *C. aura*.

The distinctions between the so-called *C. jota* and *C. aura* seem, judging from the published accounts, to be by no means clear. Mr. Cassin, in his report on the birds of Lieutenant Gilliss's Expedition, says the *C. jota* "is apparently, or so far as can be ascertained from prepared specimens, a more slender bird, and longer in all its measurements." The last character is particularly applicable to its wings.* In his *Illustrations of the birds of California and Texas*, published the following year, he reverses this statement, and says: "The South American species [*C. jota*] is the smaller," and "is the more slender in all its members"; and adds: "All the specimens that we have seen have been of a more uniform clear black color." Having myself examined numerous specimens, both in Brazil and in Florida, I find the difference in the average exceedingly slight, and nearly as stated by Mr. Cassin in his later work; that is, the Brazilian are slightly smaller, and have the plumage appreciably darker.

Bonaparte, in his *Conspectus*, gives the *jota* of Molina as being simply smaller and with a shorter tail than *aura* of Linné. The differences are indeed very slight; they are, moreover, strictly in accordance with the well-known general laws of variation between specimens of the same species from northern and southern localities, and by no means indicate a diversity of species. Because formerly not known to occur in some of the West India Islands, it was at one time supposed by some that the habitats of the two supposed species did not meet, or that there was a region in Central and Northern South America where neither existed. As I have elsewhere stated,† this is a mistake, both this species and the *C. atratus* ranging from

* U. S. Naval Astronomical Expedition, Vol. II, p. 173, 1855.

† *Memoirs Bost. Soc. Nat. Hist.*, Vol. I, p. 500, 1868.

the middle and northern portions of the United States nearly to the southern extremity of South America; the *C. aura* also extending as much beyond the southern limit of the *C. atratus* in South America as it does to the north of it in North America.

The *Cathartes Burrovianus* of Cassin, described in 1843, from a single specimen from Mexico, is referred by Bonaparte, in his *Conspectus*, to *C. jota*, or to what I regard as the typical form of *C. aura*, and evidently with good reason. It differs from *C. aura* only in being smaller. I am therefore disposed to regard it as based on an unusually small specimen of that species. Though Dr. Gambel supposed he had seen it with the other species in Lower California, but two specimens seem to have been known to Mr. Cassin, one of which was from an unknown locality.

Measurements of Florida Specimens of CATHARTES AURA.

M. C. Z. No.	Coll. No.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.
5143	—	♂	Jacksonville	Jan. 25, '69	J. A. Allen	27.50	72.50	22.50	11.10
5180	—	♂	Hibernia	Feb. 1, '69	"	27.50	72.00	22.00	11.75
5187	—	♂	"	"	"	—	—	21.00	12.00
10746	2541	♂	Dummitt's.	Mar. 11, '69	C. J. Maynard	26.50	68.00	21.00	11.00
—	2603	♂	"	Mar. 11, '69	"	—	68.00	20.00	10.50
—	2433	♂	"	Mar. 10, '69	"	27.50	72.00	21.75	11.25

80.* *Cathartes atratus* Swainson. BLACK VULTURE.

? *Vultur brasiliensis* aut *mexicanus* RAY, Synop. Meth. Avium, 10, 1713.

Vultur atratus BARTRAM, Travels, 289, 1791.

Cathartes atratus SWAINSON, Faun. Bor. Am., II, 6, 1831. — AUDUBON, Synopsis, 3, 1839. — BONAPARTE, Consp. Gen. Av., I, 9, 1850. — CASSIN, Illust. Birds Cal., Texas, etc., 58, 1854. — CASSIN, Gilliss's U. S. Nav. Astr. Exp., II, 173, 1855.

Vultur jota WILSON, Am. Orn., IX, 104, pl. lxxv, fig. 2, 1814. (Not of Molina; not of Gmelin.)

Cathartes jota BONAPARTE, Ann. N. Y. Lyc. Nat. Hist., II. — AUDUBON, Orn. Biog., II, 33, 1835. — BONAPARTE, Geog. and Comp. List, I, 1838.

Vultur urubu VIEILLOT, Ois. Am. Sept., I, 53, pl. ii, 1807.

Cathartes urubu LESSON, Voy. autour du Monde, 614. — D'ORBIGNY, Voy. dans l'Amer. Merid., 1844.

Cathartes aura SPIX, Av. Spec. Novæ, 2, 1824.

Cathartes brasiliensis BONAP., Consp. Gen. Av., I, 9, 1850.

Abundant. On the whole, probably about as numerous as the preceding, but the two species occur in different proportions at different localities, and at different times at the same locality. None were seen about Jacksonville during the two weeks I spent there in January, and none were met with for some distance up the river. Above Lake

George it was generally common, and sometimes outnumbered the other species, as it did often at Hawkinsville during my stay there. The younger birds appear to be generally not so highly colored as the fully mature, nor to have the naked skin of the head and neck so rugose and corrunculated as the older. The differences in these respects are very considerable between individuals of the same flock.

A comparison of Florida specimens with Brazilian ones shows that the latter are slightly smaller than the former; in color or other general features they do not appear to differ. Most writers have regarded the South American as identical with the North American, but Mr. Cassin,* apparently on the authority of Bonaparte,† says the South American bird "is the *Vultur brasiliensis* Ray," and that "it is considerably smaller, and otherwise quite distinct." But he only refers definitely to the difference in size. The year preceding the publication of these remarks, however, he gives *C. atratus* as inhabiting Chili.‡ In speaking of the Chili specimen, he says: "A single specimen in mature plumage and excellent condition is exactly identical in size and other characters with the common species [*C. atratus*] of the southern parts of North America." He adds: "It is the only specimen presenting this similarity that we have ever seen from South America, and is larger and in other respects different from the allied *Cathartes brasiliensis*, which is an inhabitant also of that division of this continent." *C. atratus*, he says, is "not abundant in Chili, though represented to be occasionally met with in the interior"; these larger individuals referred to being doubtless the birds that inhabit the more elevated districts. Whatever Mr. Cassin's *Cathartes brasiliensis* may prove to be, it remains unquestionable that the *C. atratus* is a general inhabitant of South America, and that Bonaparte's *brasiliensis* is merely the southern type of this species. The exact parallelism of its range on the two continents as compared with that of *C. aura* has already been alluded to.

The PAINTED or SACRED VULTURE ("*Vultur sacra*"), § an apocryphal species described by Bartram || as inhabiting Florida, demands in this connection a passing notice. Though not identified by any succeeding author (by some, however, it has been referred to the king vulture, *Sarcorham-*

* Illust. Birds of Cal. and Texas, p. 58, 1856.

† Conspectus Generum Avium, Tom I, p. 9, 1850.

‡ U. S. Naval Astronomical Expedition, Vol. II, p. 173, 1855.

§ Travels in Florida, etc., p. 150, 1790.

|| *Vultur sacra* BARTRAM, Travels, pp. 150, 289, 1791. — VIEILLOT. — NUTTALL, Man. Orn. I, 42.

Sarcorhamphus sacer CASSIN, Illust. Birds of Cal. and Texas, 59, 1856.

See also BONAPARTE, Conspectus Gen. Av., I, 9.

phus papa), Bartram's account of it leads one to infer that he found it quite abundant. His description of it is given with satisfactory detail. He says it is "near the size of the turkey-buzzard, but his wings are much shorter, and consequently he falls greatly below that admirable bird in sail. I shall call this bird the painted vulture. The bill is long and straight almost to the point, where it is hooked, or bent suddenly down, and sharp; the head and neck bare of feathers nearly down to the stomach, where the feathers begin to cover the skin, and soon become long and of a soft texture, forming a ruff or tippet, in which the bird, by contracting his neck, can hide that as well as his head; the bare skin on the neck appears loose and wrinkled, which is of a deep bright yellow color, intermixed with coral red; the hinder part of the neck is nearly covered with short, stiff hair; and the skin of this part of the neck is of a dun-purple color, gradually becoming red as it approaches the yellow of the sides and fore part. The crown of the head is red; there are lobed lappets of a reddish orange color, which lay on the base of the upper mandible. But what is singular, a large portion of the stomach hangs down on the breast of the bird, in the likeness of a sack or half wallet, and seems to be a duplicature of the craw, which is naked and of a reddish flesh color; this is partly concealed by the feathers of the breast, unless when it is loaded with food (which is commonly, I believe, roasted reptiles), and then it appears prominent. The plumage of the bird is generally white or cream color, except the quill feathers of the wings, and two or three rows of the coverts, which are of a beautiful dark brown; the tail, which is large and white, is tipped with this dark brown or black; the legs and feet of a clear white; the eye is encircled with a gold-colored iris; the pupil black.

"The Creeks or Muscogulgees," he continues, "construct their royal standard of the tail feathers of this bird, which is called by a name signifying the eagle's tail; this they carry with them when they go to battle, but then it is painted with a zone of red within the brown tips, and in peaceable negotiations it is displayed new, clean, and white; this standard is held most sacred by them on all occasions, and is constructed and ornamented with great ingenuity. These birds seldom appear but when the deserts are set on fire (which happens almost every day throughout the year in some part or other, by the Indians, for the purpose of rousing up game, as also by the lightning), when they are seen at a distance soaring on the wing, gathering from every quarter, and gradually approaching the burnt plains, when they alight upon the ground yet smoking with hot embers; they gather up the roasted serpents, frogs, and lizards, filling their sacks with them. At this time a person may shoot them with pleasure, they not being willing to quit the feast, and indeed seem to brave all danger."

Mr. Cassin * refers the species described as above by Bartram to the genus *Sarcorhamphus* (*S. sacer* Cassin = *Vultur sacra* Bartram), believing it to be a valid species, and remarks that its identification "may be considered as one of the most important services to be performed in North American ornithology." It is related, Mr. Cassin continues, "to the king vulture (*S. papa*), but that species has a black tail, and in case of mistake or misprint in Bartram's description, it may be presumed, at any rate, to relate to an occurrence of that species within the United States.† There is no more interesting nor more singular problem in North American ornithology." Two years later, in Baird's *Birds of North America*, Mr. Cassin again refers to the subject, and says that "recent information renders it probable that this [*Vultur sacra* Bartram], or a species different from the vultures just described [*Cathartes aura*, *C. atratus*, *C. Burrovianus*], is found about Lake Okechobee in Southern Florida, where it is called king buzzard. The verification of this statement by actual specimens would be one of the most important discoveries yet to be made in North American ornithology."

Although the description of Bartram's "*Vultur sacra*" accords more nearly with the *Sarcorhamphus papa* than with any other known species, I cannot avoid the conclusion that it is in the main a *purely mythical species*, notwithstanding the high reputation for veracity generally accorded to Mr. Bartram. I mainly so regard it for the reason that Florida has of late been too often traversed by naturalists, and especially all the parts visited by Bartram, for a bird of so striking an appearance, and so numerous as Bartram represented his *V. sacra* to be, to remain undiscovered if such a species exists there. While it nearly accords with the *S. papa* in size and general color, it is most radically different from this species, in the color of the tail, and in having a "large portion of the stomach hanging down on the breast, in the likeness of a sack or half-wallet." In the latter feature it is structurally widely different from any known American bird. It is mentioned as though it was an abundant species on, at least, the upper portion of the St. John's River, inasmuch as he speaks of large flights of them. As to the feathers of its tail being used by the Creek Indians for a royal standard, and to which feathers they give a "name signifying an eagle's tail," it seems to me more probable that they were really feathers of the white-headed eagle (*Haliaëtus leucocephalus*), since it is well known that the tail feathers of that bird are very generally used for this and similar purposes by the Indian tribes of this continent. whereas the tail feathers of so foul a bird as the vulture must in all

* Illustr. of Birds of Cuba and Texas, p. 59.

† The *S. papa*, a Central and South American species, appears to have not yet been seen north of Mexico.

probability be too ill scented to suit even the unfastidious taste of an Indian. As to Mr. Cassin's supposition that the word *white* in the description of the tail should perhaps read *black*, the context wholly forbids its probability. If thus changed the passage referred to would read, "the tail which is rather large and *black*, is tipped with this *dark brown or black!*" which makes simply an absurdity. Besides this, the tail is again mentioned in the following paragraph as being painted by the Indians, when used in their war standards, etc., "with a zone of red within the brown tips," and afterwards as being "displayed new, clean, and *white*." As to the information referred to by Mr. Cassin as having been received by him respecting a "king buzzard" existing in Southern Florida, it may be remarked that this is the name by which the caracara eagle (*Polyborus tharus* Cassin) is commonly known in Florida, and which is undoubtedly the bird of which, under the name of "king buzzard," Mr. Cassin had heard.

On the whole, it seems evident that Bartram's account of the *Vultur sacra* is a confused mixture either of pure fiction and truth, with the former largely in preponderance, or of the characters of several different species. The description would seem to have been mainly drawn from an example of *Sarcoramphus papa* that he may have somewhere met with, but with which he combined certain features of this or other species which he had only observed at a distance, and that he thus misjudged their exact character (as in respect to the strange external food-pouch) or else added them solely on popular, fabulous rumors. The flights of these birds, which he observed assembling over recently burned districts, I think must refer to the *Polyborus tharus*, which is well known to have this habit, while the tail feathers he speaks of as used by the Indians in their councils were more probably either those of the *Haliaëtus leucocephalus* or *Polyborus tharus* than of any species of vulture, since a white-tailed American vulture, I believe, is a bird thus far unknown. If the "*V. sacra*," then, is to be regarded as anything else than a myth, it should in all probability be identified with the *S. papa*, as already stated, and as was done by Bonaparte in his *Conspectus*.

FALCONIDÆ.

81† **Falco peregrinus** Linné. DUCK HAWK.

Falco peregrinus GMELIN, Syst. Nat. I, 272, 1788. — WILSON, Am. Orn., IX, 120, 1814. — BONAPARTE, Journ. Phil. Acad. Nat. Sci., 1st Ser., I, 342, 1824. — AUDUBON, Orn. Biog., I, 85, 1832; V, 365, pl. xvi. — NUTTALL, Man. Orn., I, 53, 1832.

Falco anatum BONAPARTE, Geog. and Comp. List, I, 1838. — CASSIN, Illust. Birds Cal. and Texas, 86, 1853. — CASSIN, Baird's Birds of N. Am., 7, 1858. — ALLEN, Proc. Essex Inst., IV, 153, 1865.

Falco nigriceps CASSIN, Illust. Birds of Cal., 87, 1853. — CASSIN, Baird's Birds of N. Am., 1858.

"One instance, St. Augustine, February, 1868." *Boardman*. Mr. Maynard found it rather common near Dummitt's, where he observed its peculiar manner of capturing the ducks. Also well known to occur in winter in Cuba and other of the West India Islands.

In 1838, Bonaparte, in his "Geographical and Comparative List," gave to the American peregrine or duck hawk the name *Falco anatum*. Previous to this time all writers had considered it, and it seems to me justly, as identical with the European peregrine, or *F. peregrinus*, — an opinion still held by many eminent ornithologists. Until about this date the peregrine falcon was believed to have a nearly cosmopolitan distribution, but since then the Australian and other supposed species have been separated from it on grounds that it now seems should be reconsidered. Among these supposed species is the *Falco nigriceps* of Cassin, first described in 1858, from specimens received from California and Chili. These first specimens were smaller, with the rufous color of the under parts in the young of a stronger tint than in the so-called *F. anatum*, they more resembling the African, Australian, and especially the Indian type of *F. peregrinus*. Specimens since obtained from farther north, however, fully equal those from Eastern North America, and the slight differences found to really exist between them seem to be by no means of specific value.

Formerly a difference in breeding habits was supposed to obtain between the American and European peregrines, the American peregrine being for a long time believed to breed in trees, whilst the European was well known to nest on cliffs. Recently, however, the American bird has been repeatedly found nesting in similar situations, but never yet in trees.*

82.† *Falco columbarius* Linné. PIGEON HAWK.

Falco columbarius WILSON and subsequent American writers generally.

Falco æsalon SWAINSON, Fann. Bor. Am., II, 35, pl. xxv, 1831. — NUTTALL, Man. Orn., I, 60, 1832.

Falco temerarius AUDUBON, Orn. Biog., I, 381, pl. lxxv, 1832.

"Frequent." *Boardman*.

* For an account of the breeding habits and nesting-places of the American bird in the Atlantic States, see the author's papers in Proc. Essex Inst., Vol. IV, pp. 153 - 161, and American Naturalist, Vol. III, p. 514. The past summer (1870) its eggs have been received by Mr. C. W. Bennett from Vermont. Prof. S. S. Haldeman was not only the first naturalist who made known the fact of its breeding on cliffs, but of its breeding in the United States. See Proc. Phil. Acad. Nat. Sci., Vol. I, p. 54, July, 1841.

Many of the earlier ornithological writers regarded, as is well known, a considerable proportion of the rapacious birds of North America as identical with species inhabiting the Old World. More accurate comparisons of specimens from the two continents, however, eventually revealed appreciable differences between them, and one after another of those of the American continent were regarded as specifically distinct from their Old World relatives; and now there is not one of the diurnal species that has not been separated by one author or another. The owls of the two continents, with two exceptions, have also been similarly separated. While in many of these cases there are appreciable differences that seem more or less constant, in the majority of instances there appears to be no just cause for the separation. Especially is this the case in respect to *Falco peregrinus* (as already observed), *Falco candicans*, *Archibuteo lagopus*, *Aquila chrysaëtos*, *Pandion haliaëtus*, *Otus vulgaris*, *Brachyotus palustris*, *Nyctale Tengmalmi*, and *Strix flammea*, in all of which species the American birds have been specifically separated from the European. *Buteo borealis*, *Astur atricapillus*, and *Falco columbarius* present stages of plumage that are scarcely distinguishable from certain stages of respectively *Falco æsalon*, *Buteo vulgaris*, and *Astur palumbarius*, and it is hence not strange that each of these European species have been described by many good authorities as occurring in the northern parts of North America. Certain styles of plumage presented by *Falco columbarius*, especially at northwestern localities, so strongly resemble common phases of *F. æsalon*, that one is readily puzzled to know whether to recognize the latter as also inhabiting North America, or whether, since these types imperceptibly grade into the so-called typical *F. columbarius*, all should not be regarded as forming a single species, since they differ essentially only in coloration, and never very widely. The specimens of *F. æsalon* before me (all immature) mainly differ from average specimens of *F. columbarius* of corresponding age in being less ferruginous, the style of coloring being the same in both.

83.* **Falco sparverius** Linné. SPARROW HAWK.

Falco sparverius LINNÉ, Syst. Nat., 128, 1766; and of subsequent writers generally.

Falco dominicensis GMELIN, Syst. Nat., I, 285, 1788.

Falco gracilis SWAINSON, Lardner's Cab. Cyc., 281, 1838.

Falco cinnamominus SWAINSON, Ibid., 281.

Falco isabellinus SWAINSON, Ibid., 281.

Falco sparveroides VIGORS, Zool. Journ., III, 436, 1827.

Abundant. Breeds in March. As has been previously pointed out, though not observed by all writers, the sexes differ greatly in color, the

adult females being banded transversely above, much as the young birds are.

Florida specimens are considerably smaller than New England ones, the former being intermediate in size between the latter and the West Indian and South American representatives of this species, which have been regarded as distinct species, and to which various names have been applied by different writers. Audubon observes that he found this species in the Southern States, and more especially in Florida, so much smaller than the northern birds that he was at first inclined to consider them specifically distinct, but finally felt sure they were the same. The colors, as usual in other species, are generally brighter in the more southern examples. Wide variations in the color of the plumage in this species have been long recognized, but, as Mr. Cassin has remarked, "they do not appear to be constant, nor peculiar to any locality." *

84 * **Accipiter fuscus** Bonaparte. SHARP-SHINNED HAWK.

Falco fuscus GMELIN, Syst. Nat., I, 280, 1788.

Accipiter fuscus BONAPARTE, Geog. and Comp. List, 5, 1838.

Astur fuscus AUDUBON, Syn., 18, 1839

Falco dubius GMELIN, Syst. Nat., I, 281, 1788.

Falco velox WILSON, Am. Orn., V, 116, 1812.

Falco pennsylvanicus WILSON, Ibid., VI, 13, 1812.

Accipiter striatus VIEILLOT, Ois. Am. Sept., I, 42, 1807.

Accipiter fringilloides VIGORS, Zool. Journ., III, 434, 1827.

Accipiter pennsylvanicus RICH. & SWAIN., Faun. Bor. Am., II, 44, 1831.

Nisus Malfini LESSON, Traité d'Ornithol., I, 58, 1831.

Common. I was unable, however, to obtain specimens.

In this species, as in the hawks generally, but more especially in the group to which the present species belongs, there are wide variations in color and size, not only with age and sex, but independently of either. One of the most interesting features in the specimens before me, in respect to these variations, is the much brighter color of the several western and southwestern examples in the collection of the Museum, as compared with New England ones. In one from Cheltenham, Missouri, the color of the lower parts is nearly uniformly red; the transverse dark lines, which in adult eastern specimens usually occupy half the exposed surface of the feathers, and often more, being in this specimen almost obsolete. The tibial feathers are especially bright, while the tints are livelier throughout the plumage. Other specimens from Fort Steilacoom, received from the Smithsonian Institution, present nearly the same appearance. Although the western representatives of the present species yet await some enter-

* Illust. Birds of California and Texas, etc., p. 93

prising divisionist to give them a distinctive name, they are interesting as indicating a rufous western race, corresponding with the *Accipiter mexicanus* form of the *A. Cooperi*, the *Falco nigripes* form of the *F. peregrinus*, the *Archibuteo ferrugineus* form of the *A. lagopus*, and the western rufous forms of *Buteo borealis* and *Circus hudsonius*.*

Although the *Accipiter fuscus* has always been regarded as closely related to the *Accipiter nisus* of the Old World, they have, with one or two exceptions,† been regarded by all authors as specifically distinct. The only distinctive difference between them, however, has been properly regarded as a slight difference in color, which difference is merely one of tint, the style of coloration being precisely the same in both. In the Museum of Comparative Zoölogy are several specimens of *A. nisus* from Germany and Switzerland, which represent both the adult and the young. The brown transverse markings on the lower plumage of the mature *A. nisus* are rather darker and broader than in most New England specimens of *A. fuscus*; but they still more closely resemble average New England specimens than the latter do any specimens of *A. fuscus* I have seen from the western parts of the United States. The western form of *A. fuscus*, as already stated, is brighter colored or more rufous than the eastern, while the eastern differs similarly from the European, the latter being much duller colored than the eastern form of *A. fuscus*. So closely, however, does one of the immature examples of *A. nisus* resemble several of the immature New England specimens of *A. fuscus*, that, if their origin was unknown, few ornithologists would probably consider them as otherwise than specifically identical; especially if placed in a large series composed of both eastern and western specimens of the *A. fuscus*. As I have previously remarked, the transverse markings on the lower plumage in the adult stage are broader and more regular and distinct in *A. nisus* than in *A. fuscus*. This, it may be added, is also the only difference observable between *A. palumbarius* and *A. atricapillus*. Such a coincidence of parallel differences between *Accipiter nisus* and *Accipiter fuscus*, and between *Astur palumbarius* and *A. atricapillus*, is a point of much interest to any

* For further remarks concerning the rufous western races of several of these species see the following pages.

† Prince Max zu Wied, in his "Beiträge zur Naturgeschichte von Brasilien," referred a hawk, probably of this species, of which he obtained a single immature male in Eastern Brazil, to the *Falco nisus* Linn. Respecting this species he observes: "Der Vogel dieser Beschreibung scheint von dem europäischen Sperber nicht abzuweichen. . . . Dieser Sperber ist mir selbst in Brasilien nicht vorgekommen, allein Freireifs hat mir ein Exemplar davon mitgetheilt, welches in der Gegend von Camamu, südlich von Bahia, geschossen wurde. So viel ich von diesem einzigen Individuo urtheilen kann, so scheint es identisch mit dem europäischen *Nisus* zu seyn; denn sowohl seine Verhältnisse als sein Gefieder stimmen vollkommen überein." Vol. III, pp. 112, 114.

one interested in geographical color variations in animals; the more so, perhaps, from the two latter species being so intimately related as to have been at one time generally regarded as identical. Yet so far as can be judged from a limited number of specimens, *Astur palumbarius* differs more from *A. atricapillus* than *Accipiter nisus* does from *Accipiter fuscus*, which latter species have never been considered as identical.*

85.* **Accipiter Cooperi** Cassin. COOPER'S HAWK.

Falco Cooperi BONAP., Am. Orn., II, 1, 1828.

Falco Stanleyi AUDUBON, Orn. Biog., I, 186, 1831 (young).

Astur Cooperi BONAP., Geog. and Comp. List, 5, 1838.

Accipiter Cooperi CASSIN, Illust. Birds of Cal., etc., 96, 1854.

Accipiter mexicanus SWAIN., Faun. Bor. Am., II, 45, 1831. — CASSIN, Baird's Birds N. Am., 17, 1858.

Accipiter Gundlachi LAW., Ann. N. Y. Lyc. Nat. Hist., VII, 252, 1860.

Common.

Mr. Cassin has very properly indicated the variations in size and color commonly seen in this species in the following remarks: "Rather a difficult species to the ornithologists, on account of the great variations in its colors, and in size also. It is, in fact, unusual to find two alike in a dozen specimens." † Its relationship to *Accipiter fuscus* is of course well understood, it holding a similar relationship to that species that *Picus villosus* does to *Picus pubescens*, the essential difference between them being mainly a great difference in size. But the specific distinctness of *A. mexicanus* from it is not so clear. Being without authentic specimens of *A. mexicanus*, and having only New England specimens of *A. Cooperi*, ‡ I cannot speak confidently respecting the character and affinities of the former. According to authors, however, it seems to differ from *A. Cooperi* in being somewhat smaller and more highly colored. It is also more southern in its distribution. Hence these variations, being in accordance with the general laws of geographical variation in size and color, do not necessarily

* In this connection I wish to cite some interesting variations in color presented by Massachusetts and Maine specimens of *Astur atricapillus*. Ordinarily this species has each feather below centred with a longitudinal dark shaft-line, with several transverse broader but somewhat irregular dark ashy-brown bars on a lighter ground. Some specimens, however, as one from Maine, have the transverse bars so narrow and broken that the lower surface presents a nearly uniform, minutely mottled appearance. Another specimen (from Springfield, Mass.) represents the opposite extreme, it having the transverse bars broad, regular, and quite far apart, so that its resemblance to average specimens of *Astur palumbarius* is very close. The color in this specimen is much darker throughout than is usual in this species.

† Illustrations of Birds of California, etc., p. 93, 1854.

‡ Since the above was written, specimens have been received at the Museum from Jalapa, Mexico, from Sn. R. Montes-de-Oca.

imply a diversity of species; they only accord with what would naturally be expected to occur if *A. mexicanus* and *A. Cooperi* were known to constitute but a single species.*

Accipiter Cooperi, as is well known, is not only closely allied in general structure to *Buteo lineatus*, but also in style of coloration in both the immature and adult stages. It may be fair, then, to test the value of the distinctive characters assigned to *A. mexicanus* by what obtains as geographical variations in size and color in *Buteo lineatus*. Of this species I have fortunately a large number of specimens, including some from localities similarly separated to those whence *A. Cooperi* and *A. mexicanus* respectively come. In the case of *Buteo lineatus* there is no reason whatever to doubt that my specimens from Florida and New England are specifically identical. Yet the Florida specimens are very much brighter colored, and very much smaller; the difference in the length of the folded wing between two males, one of which is from Maine and the other from Florida, being *two and one half inches*, with corresponding differences in general measurements. This is relatively much greater than the difference in size between specimens of the so-called *A. Cooperi* and *A. mexicanus*. Similar variations in color and size to those between *A. Cooperi* and *A. mexicanus* also occur between northeastern and southwestern specimens of *A. fuscus*, the latter, as already noted under *A. fuscus*, being smaller than the former, and very much brighter colored; the difference in color between specimens from Maine and the State of Missouri being greater than is represented to occur between *A. Cooperi* and *A. mexicanus*, and of a parallel kind. In accordance with the evident inference that may be drawn from these facts, I provisionally include *A. mexicanus* among the synonymes of *A. Cooperi*. The *A. Gundlachi* of Cuba differs from the southern *A. Cooperi* in the way southern birds usually differ from the northern ones of the same species, that is, in being smaller and brighter colored, and in having the dark transverse bars on the under plumage increased in breadth at the expense of the alternating light ones.

86.* ***Buteo borealis* Bonaparte.** RED-TAILED HAWK.

Falco borealis GMELIN, Syst. Nat., I, 266, 1788. — WILSON, Am. Orn., VI, 75, pl. lii, fig. 2, 1812. — RICH. & SWAIN., Faun. Bor. Am. II, 50, 1831. — AUDUBON, Orn.-Biog., I, 265, pl. II, 1832.

Buteo borealis BONAPARTE, Geog. and Comp. List, 3, 1838. — GOSSE, Birds of Jamaica, II, 1847. — LEMBEYE, Av. de la Isla de Cuba, 18, 1850. — CASSIN, Syn. N. A. Birds (Illust. Birds Cal. and Texas, etc.), 97, 1854. — BREWER, N. Am. Oölogy, 21, 1857. — CASSIN, Baird's Birds of N. Am., 25, 1858. — BRYANT, Proc. Bost. Soc. Nat. Hist., VIII, 109, 1861. — ALLEN, Memoirs Bost. Soc. Nat. Hist., I, 499, 1868.

* Bonaparte indeed long since cited *A. mexicanus* Swainson as a synonyme of *A. Cooperi*.

- Falco leverianus* GMELIN, Syst. Nat., I, 266, 1788. — WILSON, Am. Orn., VI, 78, pl. lii, 1812.
- Falco jamaicensis* GMELIN, Syst. Nat., I, 266, 1788.
- Falco aquilinus* BARTRAM, Travels, 290, 1791.
- Falco Harlani* AUDUBON, Am. Orn., I, 441, 1831.
- Accipiter ruficaudus* VIEILLOT, Ois. Am. Sept., I, 47, 1807.
- Buteo ferrugineicaudus* VIEILLOT, Ibid., 32.
- Buteo fulvus* VIEILLOT, Nouv. Dict. Hist. Nat., IV, 472, 1816.
- Buteo americanus* VIEILLOT, Ibid., 477.
- Buteo vulgaris* RICH. & SWAINSON, Faun. Bor. Am., II, 47, pl. xxvii, 1831. — AUDUBON, Syn., 5, 1839.
- Buteo buteoides* NUTTALL, Man. Orn., I, 100, 1832.
- Falco buteo* AUDUBON, Orn. Biog., IV, 108, 1838.
- Buteo Swainsoni* BONAPARTE, Geog. and Comp. List, 3, 1838. — CASSIN, Illust. Birds Cal. Texas, etc., 98, 1854. — BREWER, N. Am. Oölogy, 24, 1857. — COOPER & BAIRD, Orn. Cal., I, 476, 1870.
- Buteo Harlani* BONAPARTE, Geog. and Comp. List, 3, 1838. — CASSIN, Illust. Birds Cal., Texas, etc., 101, 1854. — CASSIN, Baird's Birds N. Am., 14. — ? BRYANT, Proc. Bost. Soc. Nat. Hist., VIII, 115, 1861. — COOPER & BAIRD, Orn. Cal., I, 473.
- Buteo montanus* NUTTALL, Man. Orn. I (2d ed.), 112, 1840. — CASSIN, Baird's Birds N. Am., 26. — COUES, Proc. Phil. Acad. Nat. Sci., 1866, 43. — COOPER & BAIRD, Orn. Cal., I, 469.
- Buteo Bairdii* HOY., Proc. Phil. Acad. Nat. Sci., 1853, 451. — CASSIN, Baird's Birds N. Am., 21.
- Buteo insignatus* CASSIN, Birds Cal. and Texas, 102, pl. xxi, 1854. — CASSIN, Baird's Birds N. Am., 23. — COOPER & BAIRD, Orn. Cal., I, 474.
- Buteo calurus* CASSIN, Proc. Phil. Acad. Nat. Sci., 1855, 281. — CASSIN, Baird's Birds N. Am., 22. — COOPER & BAIRD, Orn. Cal., I, 471.

Not apparently uncommon, but far less numerous than the next species.

The *Buteoninæ*, or the group of hawks to which the present and the two following species belong, is well known to embrace species more variable in color than those of any other section of the *Falconidæ*, although all the members of this family are more or less remarkable for individual and other variations of plumage. The present species, however, admitting for it the wide variation in this respect herein claimed, scarcely equals the immense range of color variation well known to characterize its near ally and representative in the Old World, the *Buteo vulgaris* auct. (*Falco buteo* Linné). Six specimens of this species in the Museum from Switzerland and Germany, received under the name *Falco buteo*, vary in color as follows: One is almost entirely black; another is nearly black throughout, with obscure narrow transverse bands of ferruginous on the

crissum and abdomen; another is mainly black, but varied below bars of pale rufous and blotches of white; a fourth is also nearly black, very dark brown, but considerably more relieved with white below than the last; a fifth is mainly white below, with longitudinal stripes of dark brown, and so nearly resembles a common immature stage of the American *Buteo borealis* that if placed together the most discriminating observer could not tell which specimen was the European or which the American one. The sixth is very light colored throughout, with only a few dusky longitudinal spots on the breast. This specimen is also not readily distinguishable from certain common phases of *B. borealis*. Another specimen of *B. vulgaris*, in the La Fresnaye collection in the Museum of the Boston Society of Natural History, is still lighter than this, being nearly uniform whitish below, and very light colored, almost white above. The latter specimen and the first-mentioned dark specimen present as great differences in color as two specimens of one species can well be conceived to exhibit.

The variations presented by the American *B. borealis* have already been fully detailed by the late Dr. Henry Bryant, in his "Remarks on the Variations of the Plumage of *Buteo borealis* auct., and *B. Harlani* Aud."* He observes that the variation in plumage of the species of *Buteo*, common in the Atlantic States, "are so slight that it is not to be wondered at that the first specimens from other parts of the country, presenting as they did such extraordinary variations in color, should have been described as distinct species. At present, however," he continues, "the number of specimens known is so large that on careful examination it seems to me necessary to adopt one of two conclusions, namely, either to increase the number of species indefinitely, or to reduce them to a much smaller number than are now supposed to exist. As the European buzzard, *Buteo vulgaris*, is well known to present the greatest variety of color, it seems to me more reasonable to adopt the last conclusion."† With the above opinions and remarks I in the main agree, but do not regard the variations presented by the *Buteo borealis* as by any means slight, even in the Atlantic States. Although instances of such excessive variation as are seen in the Central and Pacific States are apparently more rare in the Atlantic States, speci-

* Proc. Bost. Soc. Nat. Hist., Vol. VIII, p. 107, 1861.

† In respect to the variety of color in the *B. vulgaris*, Dr. Bryant makes the following quotation from Naumann's Natural History of the Birds of Germany (Vol. I, p. 347): "In the coloring of the feathers of the bird there prevails a most extraordinary difference, and one which is not often seen in other birds of prey. From the darkest uniform blackish-brown to the purest white, we find all the shades, and also both colors mixed and spotted, in such various ways that the countless transitions cannot be described; this difference is independent of age and sex." Many other European writers, it may be added, have made similar remarks in respect to its astonishing range of variation in color.

mens from Massachusetts now before me vary as follows: Some are nearly unspotted beneath, others, sparsely spotted, have the spots mainly restricted to the pectoral region; others, in which the spots are equally few, have them mainly accumulated on the abdominal region, while still others have them so numerous as to occupy the greater part of the lower surface, sometimes covering the abdomen in an almost unbroken broad band. They likewise vary in the amount of rufous tint in the plumage, in some it being very slight, while others are as strongly ferruginous as any of the California specimens (*B. montanus*) I have yet seen.

The *Buteo borealis* was first described by Latham in his "General Synopsis of Birds,"* in 1781, under the names of "cream-colored buzzard" and "American buzzard," the first name being applied to the young,† and the last to the adult stage of plumage. Pennant, in his "Arctic Zoölogy,"‡ also redescribes the immature bird as the "Leverian falcon," and to these several descriptions of Latham and Pennant, Gmelin, in his "Systema Naturæ," gave respectively the names *Falco jamaicensis*, *F. borealis*, and *F. Leverianus*. Some twenty years later the *Buteo borealis* was redescribed by Vieillot, in his "Histoire des Oiseaux de l'Amerique Septentrionale," as *Accipiter ruficaudus* and *Buteo ferrugineicaudus*, both names evidently referring to the mature or nearly mature bird; and again ten years later, in the "Nouveau Dictionnaire d'Histoire Naturelle," as *Buteo fulvus* and *B. americanus*. Audubon, in 1831, figured and described a specimen from Louisiana under the name *Falco Harlani*. This specimen, which was finally sent to the British Museum, has been regarded by Mr. G. R. Gray and others as only a very dark-colored example of *B. borealis*.§ In the same year Richardson and Swainson reported the *Buteo vulgaris*, in their "Fauna Boreali-Americana," as an inhabitant of North America, and of which they figure an immature male. As already remarked, the *B. vulgaris*, in certain stages of plumage, is not readily distinguishable from *B. borealis*, so that the mistake is a perfectly excusable one. This form, however, was for some time currently received by most writers as a species distinct from the *B. borealis*, and to which the name *B. Swainsoni* was given by Bonaparte. In 1832 Nuttall described a *Buteo buteoides*, which, though referred by Bonaparte to *B. lineatus*, and by Cassin to *B. pennsylvanicus*, seems to me to much more nearly agree with *B. borealis*. In 1840 the same writer described a *B. montanus*, which was subsequently

* Vol. I, pp. 49, 50, Nos. 30 and 31.

† Latham observes: "This beautiful specimen was sent to me from Jamaica by an intelligent friend and a good naturalist, who did not hint the least of its being a variety of the common buzzard [*Buteo vulgaris* auct.], which I should have otherwise suspected."

‡ Vol. II, p. 206, No. 101.

§ Cat. of Birds in British Museum.

referred by Bonaparte to his *B. Swainsoni*, but has since been recognized as a valid species by Cassin and other recent American authors. In 1853 Mr. P. R. Hoy described a *Buteo Bairdii*, and in 1854 Mr. Cassin added *B. insignatus*, in 1855 *B. calurus* and *B. oxypterus*, and in 1856 *B. Cooperi*. In 1861 Dr. Bryant made a revision of the group, then containing eight or nine species currently recognized by American ornithologists, and reduced the number of species to two, one of which he called *B. borealis* and the other *B. Harlani*; which latter, however, is not the *Harlani* of Cassin, and probably not the *Harlani* of Audubon.

Dr. Bryant, in the above-cited paper, describes in detail the leading variations presented by our red-tailed hawks, and the character of the numerous supposed species of this group that had then been recently described. He having at his command all the specimens of this group contained in the Museums of the Philadelphia Academy and the Smithsonian Institution, including the original types of Mr. Cassin's species, as well as the specimens in his own collection, his opportunities for investigating the subject were unusually favorable. The results of his examination of this material may be briefly stated in his own words. He says that after examining this large series of specimens, he found "that of all those belonging to *Harlani*, *insignatus*, *Swainsoni*, *Bairdii*, *oxypterus*, *borealis*, *montanus*, *calurus*, and perhaps *Cooperi*," could be "easily reduced to two very distinct groups, each of which is distinguishable by definite external characters, and in which the variations of plumage, though apparently so great, if the extremes of the series only are taken into consideration, can, it seems to me, be arranged in a series, in which the connecting of the different members may be readily traced. Of these two groups, or rather species, one, which should be called *B. borealis*, as the first described, consists of that species, *montanus*, *calurus*, *Harlani*, and probably *Cooperi*, and is characterized by a very muscular body,* stronger and longer bill, longer and more powerful tarsi, and a more rounded wing, the fourth quill generally the longest, the fifth little if any shorter than the third, and the first always longer than the eighth. The other species, to which *Harlani*?, *insignatus*, *Swainsoni*, *Bairdii*, and *oxypterus* belong, is distinguishable by a more slender body, shorter and weaker tarsi, and a more pointed wing, the third quill generally the longest, the fifth considerably shorter than the third, and the first always longer than the eighth."

"On making the examinations which led to the conclusion above stated," he further observes, "I was struck by the small number of specimens in which all the feathers were equally developed, and when they were so, the variation in the proportions of the primaries, and of the wings and

* Stuffed skins evidently afford rather unsatisfactory data for the determination of the relative muscularity of the body.

tail, in specimens of the same variety, was much greater than I had expected to find"; a result which indicates how unreliable such features are as specific distinctions, as I have already repeatedly remarked, and also, of course, the fallacy of the belief so generally held, that they are really among the most trustworthy.* After detailing some of the instances of variation in this respect in the specimens in question, he makes the following remarks on variations in other characters: "The variation in the number and shape of the tarsal scales is considerable, as is usual in birds of this order. The development of the festoon of the lower edge of the upper mandible, *one of the principal generic characters*,† varies particularly in *B. montanus*, the series of which is the largest, *from a sharp, almost tooth-like process to an entire absence of it.*"‡

Dr. Bryant described each of the so-called species of the later authors, and generally several authentic specimens of each, showing the variations of color they present. *B. montanus* is the so-called "western red-tail," replacing, it is supposed, *B. borealis* in the western half of the continent, and differing from it in the main only in being more rufous or brighter colored. Some specimens, however, from California and Oregon are not appreciably different from others from the Atlantic States, and among them is one received at the Museum from the Smithsonian Institution labelled "*B. borealis.*" *B. calurus* differs from these in being much darker throughout, and especially below. It has, however, according to Dr. Bryant, two varieties, one of which is much darker than the other. The *B. Harlani* of Cassin, Dr. Bryant says, "resembles very closely the dark variety of *calurus*, with the exception of its tail, which resembles *montanus.*" Respecting the single known specimen of *B. Cooperi*, he says there is nothing in its coloration "that would make the supposition of its being a variety of *montanus* improbable." The tail presents the greatest dissimilarity and "has very much the appearance it would have in a semi-adult of this species, if the color were partially washed out." The tarsus, though long, he says is not longer than in some specimens of *montanus*; but observes that the scutellation of the tarsus presents certain peculiarities not seen in the others, there being but *two* rows of lateral scales instead of *three or four*, and two more than the usual number of transverse scales.§ In respect to these supposed species he then observes: "After

* See the remarks on this point in Part III.

† The italicizing is my own.

‡ On differences of this kind the several supposed species of the *B. borealis* group have been arranged *in different subgenera*!

§ Since writing the above I have learned from Professor Baird that he is inclined to regard this specimen as "only an *Archibuteo ferrugineus* without feathers on the tarsus; at any rate, hardly a species." It is hence omitted in Cooper and Baird's "Ornithology of California," which has just appeared.

carefully examining the birds described above, I do not see, if *Buteo borealis*, *montanus*, and *calurus* are to be considered distinct species, that we can avoid increasing the number by separating from *montanus* two species, — one the dark Steilacoom variety, and the other that from Cape St. Lucas (which, by the way, is the most distinct variety that I have seen); from *calurus*, one species, the ferruginous variety from Fort Tejon; and adding to this group one species based on the adult *Harlani* of the Academy [*Harlani* of Cassin, not of Audubon], making in all seven species of this group. I have not included in this list the young *Harlani* of the Academy, which differs as much from the adult as from any other specimen of this group; or *Cooperi*," etc. After next describing in detail *Buteo Harlani* (*B. Harlani* of Bryant, not *B. Harlani* of Cassin, nor of Audubon), and its several varieties, which form the "species" *B. insignatus*, *Swainsoni*, and *oxypterus* of Cassin and the *B. Bairdii* of Hoy and Cassin, with several varieties under each, some of which he clearly shows are connecting links to others, Dr. Bryant concludes his paper with the following summary: "Taking color, therefore as a sufficient ground for specific distinction, we find that we have in the red-tailed group seven species, and in the other nine, which, with the young *Harlani* of the Academy, *Cooperi*, *fuliginosus*, *albonotatus*, *lineatus*, *elegans*, and *pennsylvanicus*, give a total of twenty-three species of this genus which are found in the United States."

But Dr. Bryant by no means admits color in this group to be a specific characteristic, and, as I have already remarked, in reducing the number of species of the red-tailed hawks to two, he takes general size and the proportions of the primary quills of the wing as the basis of distinction. He has accordingly given a table of comparative measurements and proportions of the two species, in which he has arranged, as he says and doubtless supposed, the larger specimens under *B. borealis*, and the smaller under *B. Harlani*. Size and the proportions of the quills, however, it seems to me, are equally arbitrary grounds for their separation, as an examination of his tables and descriptions evidently proves. It happens that in the first, or *B. borealis* series, nearly all the specimens are fully adult, as indicated by the tail being uniformly red, with a subterminal black band, — a stage of plumage which characterizes only adult individuals. In the second, or *B. Harlani* series, but one specimen (which does not appear in the table of measurements), is described that is not evidently somewhat immature, while the greater part of them are quite so.* Respecting the so-called *Buteo Bairdii*, of which numerous specimens have been reported, some from quite eastern localities, Dr. Bryant

* They have at least the tail numerously banded, as all immature *B. borealis* do have, and their general diagnosis is that of immature birds.

remarks that a single specimen in the Museum of the Philadelphia Academy is the only one he had seen "presenting the least appearance of adult plumage." In regard to the size of the specimens of the two series, adopting the length of the folded wing as the basis of comparison, — the best element in the tables available for comparison, in this respect, — the smallest and the largest specimens, measuring 370 and 438 millimetres respectively, occur in the *B. borealis* series. The average length of wing in twenty specimens of *B. borealis* is 409 millimetres, and in fourteen * specimens of *B. Harlani* Bryant, 405. The difference of 4 millimetres is an amount too trivial to be of account, as the addition of a single specimen to either series might reverse the difference. Hence the impression possessed by Dr. Bryant of an average difference in size between the two series was evidently an erroneous one.

There hence remains but a single difference, that in respect to the form of the wing, or the relative length of the primaries, by which to distinguish the two series, which is at best one of doubtful value. My present opinion is that all the so-called species of these two groups may be safely referred to the original *Buteo borealis*, except the *B. oxypterus*, which should be undoubtedly referred to the *B. pennsylvanicus*.

87.* *Buteo lineatus* Jardine. RED-SHOULDERED HAWK.

Falco lineatus GMELIN, Syst. Nat., I, 268, 1788. — WILSON, Am. Orn., VI, 86, pl. liii, fig. 3, 1812. — AUDUBON, Orn. Biog., I, 296, pl. lvi, 1832.

Buteo lineatus JARDINE, Am. Orn., I, 1832. — AUDUBON, Syn., 7, 1839. — CASSIN, Baird's Birds N. Am., 28, 1858. — VERRILL, Proc. Essex Institute, III, 141, 1862.

Falco hyemalis GMELIN, Syst. Nat., I, 274, 1788. — WILSON, Am. Orn., IV, 73, 1812. — NUTTALL, Man. Orn., I, 106, 1832. — AUDUBON, Orn. Biog., V, pl. lxxi, 1832 (young).

Buteo Cooperi ALLEN, Amer. Nat., III, 518, 1869.

Circus hyemalis BONAP., Journ. Phil. Acad. Nat. Sci., 1st Ser., III, 305, 18

Buteo elegans CASSIN, Proc. Phil. Acad. Nat. Sci., 1855, 281. — CASSIN, Baird's Birds of N. Am. 28, 1858.

Very abundant. By far the most numerous species of the family.

Generally smaller and much brighter colored than New England specimens. The dark line along the shaft of the feathers below, especially on the throat and breast, is very distinct, in this respect and in the bright colors greatly resembling the so-called *Buteo elegans* of Cassin. *B. elegans*,

* The *B. oxypterus*, referred to the *B. Harlani* by Bryant, is very much smaller than any other specimen in either series, and it seems to me has decided affinities, in its small size as in other features, with the *B. pennsylvanicus*, as stated by Mr. Cassin, and it is hence excluded in my computation of the average length of the folded wing.

however, has been generally considered as the western representative of *B. lineatus*, but it differs from the latter only in being brighter colored, or in having the ferruginous of the under parts more intense. In this it resembles the western representatives of the *B. borealis*, *Archibuteo lagopus*, *Accipiter fuscus*, *Circus cyaneus*, *Falco peregrinus*, and other species of this family, the western specimens of which are ordinarily more rufous than the eastern, though in only a part of them have the eastern and western races as yet been separated as distinct species.

The considerable difference in size between specimens of this species from New England and Florida has led to the supposition that the former may be specifically distinct from the latter, or at least that they form well-marked varieties.* The following measurements, however, show that specimens occur in Florida, in winter at least, nearly as large as average-sized New England specimens. But these may have been merely winter visitors, since two of the three specimens taken in February on the St. John's River are larger than any of the others, all of which were taken later in the season. Those taken by Dr. Würdemann at Cape Florida and Indian Key are also smaller than those from the St. John's River.

Measurements of Florida Specimens of BUTEO LINEATUS.

M. C. Z. No.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.
5223	♂	Volusia	Feb. 12, '69	J. A. Allen	22.25	41.50	13.00	7.75
5224	♂	"	Feb. 12, '69	"	20.00	39.50	12.25	7.50
5276	♂	Blue Springs	Feb. 21, '69	"	20.00	42.00	13.00	8.40
5310	♂	Enterprise	Mar. 1, '69	"	17.65	39.15	12.25	8.00
5331	♂	"	Mar. 1, '69	"	17.75	40.25	12.30	7.50
5398	♂	Hawkinsville	Mar. 15, '69	"	18.00	40.50	12.85	7.75
10744	♂	Jacksonville	Dec. 31, '68	C. J. Maynard	19.20	41.50	12.60	8.50
10743	♂	"	Jan. 11, '69	"	19.20	40.05	12.60	8.50
6899	♂	Cape Florida	Apr. 5, '58	G. Würdemann	15.75	35.75	11.00	6.75
8630†	♂	Indian Key	Aug. 31, '57	"	17.50	37.00	11.20	—
6898	♂	"	Aug. 1, '58	"	15.50	34.50	10.50	7.15
8629†	♂	"	Nov. 10, '57	"	17.75	40.00	12.00	—
8631†	♂	"	Aug. 31, '57	"	17.50	37.00	11.10	—

88.† *Buteo pennsylvanicus* Bonaparte. BROAD-WINGED HAWK.

Falco pennsylvanicus WILSON, Am. Orn., VI, 22, 1812.

Buteo pennsylvanicus BONAP., Geog. and Comp. List, 3, 1838. — AUDUBON, Syn., 6, 1839. — CASSIN, Illust. Birds Cal., Texas, etc., 100, 1854. — CASSIN, Baird's Birds N. Am., 29, 1858.

Falco latissimus WILSON, Am. Orn., VII, 22, 1812. (Later published copies.)‡

* See Prof. A. E. Verrill in Proc. Essex Institute, Vol. III, p. 141, 1862.

† Smithson. Inst., No. (Copied from Cassin in Baird's Birds of North America, p. 28.)

‡ Concerning the names *F. pennsylvanicus* and *F. latissimus* given by Wilson to this species, see Mr. Cassin's remarks, Illust. Birds of Cal., Texas, etc., p. 101.

Falco Wilsoni BONAP., Journ. Phil. Acad. Nat. Sci., III, 348, 1824.

Sparvius platypterus VIEILLOT, Encyc. Meth., III, 1273, 1823.

Buteo oxypterus CASSIN, Proc. Phil. Acad. Nat. Sci., 282, 1855. — CASSIN, Baird's Birds of N. Am., 31, 1858.

"Common." — Boardman. Audubon, however, gives it as rare south of the Middle States, and it is not mentioned by Dr. Coues in his list of the birds of South Carolina. There is, however, a specimen in the Museum of Comparative Zoölogy labelled as having been taken in Florida.

As previously observed, it appears to me that the *Buteo oxypterus* of Cassin, described from a single specimen taken at Fort Filmore, New Mexico, corresponds more nearly with the young of this species than with any known stage or form of *B. borealis*.

89.* *Circus cyaneus* Boie. MARSH HAWK.

Falco cyaneus LINN., Syst. Nat. I, 126, 1766. — BONAP., Am. Orn., II, 30. — AUDUBON, Orn. Biog., IV, 396, pl. ccclvi, 1838.

Circus cyaneus BOIE, Isis, 1822, 549. — AUDUBON, Synop., 19, 1839. — G. R. GRAY, Gen. of Birds, I, p. 32. — IBID., Cat. Brit. Birds, 17, 1863.

Falco hudsonius LINN., Syst. Nat., I, 128, 1766.

Falco uliginosus GMELIN, Syst. Nat., I, 278, 1788.

Circus uliginosus VIEILLOT, Ois. Am. Sept., I, 37, 1807.

Falco uliginosus WILSON, Am. Orn., VI, 67, pl. li, fig. 2, 1812.

Buteo (Circus) cyaneus ? var. ? *americanus*, RICH. and SWAIN., Faun. Bor. Am., II, 55, pl. xxix, 1831.

Circus hudsonius VIEILLOT, Ois. Am. Sept., I, 36, 1807. — CASSIN, Ill. Birds Cal., Texas, etc., 108, 1854. — BREWER, N. Am. Oöl., 42, 1857. — CASSIN, Baird's Birds N. Am., 38, 1858.

Circus variegatus VIEILLOT, Ois. Am. Sept., I, 37, 1807.

Strigiceps uliginosus BONAP., Geog. and Comp. List, 5, 1838.

Strigiceps pygargus BONAP., Ibid.

Common about the savannas.

The present species has been considered by most writers as identical with the *C. cyaneus* of the Old World. It was first separated as a distinct species by Bonaparte in 1838, in his Geographical and Comparative List. Mr. Cassin also regarding it as distinct, this opinion has been generally adopted by recent American ornithologists. They seem to be, however, quite identical.

The same variation in color between eastern and western specimens is seen in this species that has been noted in others of this family, the young western ones especially being much brighter colored than the eastern.

The great variation in plumage attending differences of age and sex in this species have given rise to numerous synonymes, of which twenty are cited by Mr. G. R. Gray in his Catalogue of British Birds.

90.* **Pandion haliaëtus** Cuvier. FISH HAWK. OSPREY.

- Falco haliaëtus* LINNÉ, Faun. Suec., 22, 1735. — WILSON, Am. Orn., V. 13, pl. xxxvii, 1812. — BONAP., Ann. N. Y. Lyc. N. Hist., II, 26, 1828. — AUDUBON, Orn. Biog., I, 415, pl. lxxxii, 1832. — NUTTALL, Man. Am. Orn., I, 78, 1832. *Pandion haliaëtus* CUV., Règ. An., I, 316, 1817. — AUDUBON, Synopsis, 12, 1839. — G. R. GRAY, Cat. Brit. Birds, 5, 1863. — PELZELN, Ornithol. Brasiliens, 4, 1868. — HEUGLIN, Ornithol. Nordost-Afrika's, 54, 1869. *Falco arundinaceus* GMELIN, Syst. Nat. I, 263, 1788. *Falco carolinensis* GMELIN, Ibid. *Pandion carolinensis* BONAP., Geog. and Comp. List, 3, 1838. — CASSIN, Illust. Birds Cal., Texas, etc., 112, 1854. — BREWER, N. Am. Oöl., 53, 1857. — CASSIN, Baird's Birds N. Am., 44, 1858. *Falco cayanensis* GMELIN, Syst. Nat. I, 268, 1788. *Aquila piscatrix* VIEILLOT, Ois. Am. Sept., I, 29, 1807. *Pandion fluviatilis* SAVIG., Descr. de l'Égypte, Hist. Nat., I, 96, 1809. *Pandion americanus* VIEILLOT, Gal. des Ois., I, 33, 1828. *Pandion indicus* HODGSON, Journ. As. Soc. Bengal, 366, 1837.

Abundant everywhere ; especially so around the lakes of the Upper St. John's. Commences nesting in January. At Lake Monroe I counted six nests from a single point of view. Their nests were also frequent all along the river. They generally selecting a dead tree in which to build, and often those situated in cleared fields, their nests were conspicuous objects, and could usually be seen from a long distance. Even these harmless birds do not fail to attract the fire of the numerous sportsmen who visit this region in winter, some of whom are ignorant enough to believe that when shooting them they are killing "bald eagles."

Gmelin, in his "Systema Naturæ," described the present species not only as *Falco haliaëtus*, but he gave to it also the names *F. carolinensis*, *F. arundinaceus*, and *F. cayanensis*, apparently indicating under them, however, what he regarded as varieties rather than as distinct species. For many years, however, the common fish-hawk was generally regarded as having an almost cosmopolitan distribution. Bonaparte spoke of it in 1826, in his Synopsis of the Birds of the United States,* as follows: "Inhabits almost every part of the globe near waters; much more common in North America than in Europe." Ten or twelve years later, however, he seems to

* Annals of the N. Y. Lyceum of Nat. History, Vol. II, p. 26.

have changed this opinion, since in his Geographical and Comparative List of the Birds of Europe and the United States (to which paper, by the way, we are indebted for the separation of eight of the American species of raptorial birds previously considered identical with the European,* embracing all thus separated up to the present time, except two†) he calls the American fish-hawk *Pandion carolinensis*, and gives its habitat as "America generally." Other authors have since separated the West Indian and South American as a third, the Asiatic as a fourth, and the Australian as still another. The numerous specimens in the Museum show that considerable variation obtains in color, size, and proportions among those recognized by authors as belonging to the *P. carolinensis*, much greater differences in color — the main ground on which they have been separated from the European — being presented among the Florida specimens alone than obtains in the average between Brazilian and New England specimens, or American and European. Generally the feathers of the breast are each centred with a broad longitudinal spot or stripe of brown, which spots sometimes cover the greater part of the breast; but they are often simply narrow lines, and are not unfrequently entirely wanting. Sometimes these spots are uniform dark-brown, at others suffused or broadly margined with ferruginous, and are occasionally altogether of the latter color. In reuniting the American fish-hawk with the osprey of the Old World, I but adopt the view always held by a large number of ornithologists, though by all American authors they have for the last fifteen years been commonly considered as distinct.

Measurements of Florida Specimens of PANDION HALIAËTUS.

M. C. Z. No.	Sex	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.
5268	♂	Blue Springs	Feb. 21, '69	J. A. Allen	21.75	64.00	19.50	8.75
5298	♂	Enterprise	Feb. 25, '69	"	24.25	68.75	20.25	10.00
5330	♂	"	Mar. 4, '69	"	22.00	63.50	19.25	9.00
5356	♂	Hawkinsville	Mar. 10, '69	"	20.75	63.00	18.75	8.60
5355	♂	"	Mar. 10, '69	"	—	—	20.25	7.80
—	♂	"	Mar. 15, '69	"	24.25	66.25	19.00	—
—	♂	"	Mar. 15, '69	"	23.50	68.50	20.25	—

91. *Haliaëtus leucocephalus* Savigny. WHITE-HEADED EAGLE.

Falco leucocephalus GMELIN, Syst. Nat., I, 255, 1788. — WILSON, Am. Orn., IV, 89, pl. xxxvi, 1811. — AUDUBON, Orn. Biog., I, 58, pl. xxi, 1832; II, 160; V, 354, pl. cxxvi.

* *Pandion carolinensis* from *P. haliaëtus*, *Butastur* (or *Archibuteo* as now called) *Sancti-Johannis* from *B. lagopus*; *Buteo Swainsoni* from *B. vulgaris*; *Falco anatum* from *F. peregrinus*; *Astur atricapillus* from *A. palumbarius*; *Strigiceps* (*Circus* as now called) *uliginosus* from *S. pygargus* (*cyaneus* auct.); *Otus americanus* (or "*Wilsonianus*") from *O. vulgaris*; *Nyctale Richardsoni* from *N. Tengmalmi*; *Strix pratincola* from *S. flammea*.

† *Aquila chrysaëtos*, *Brachyotus palustris*.

Haliaëtus leucocephalus SAVIGNY. — BONAPARTE, Geog. and Comp. List, 3, 1838. — AUDUBON, Synop., 10, 1839. — CASSIN, Illust. Birds Cal., Texas, etc., 111, 1854. — CASSIN, Baird's Birds N. Am., 43, 1858.

Falco ossifragus WILSON, Am. Orn., VII, 16, pl. lv, 1813.

Aquila (Haliaëtus) leucocephalus RICH. & SWAIN., Faun. Bor. Am., II, 15, 1832.

Falco Washingtoni AUDUBON, Orn. Biog., I, 58, pl. xi, 1831 (plate published 1827).

Falco Washingtoniana AUDUBON, Loudon's Mag. N. Hist., I, 115, 1828.

Haliaëtus Washingtoni AUDUBON, Synop., 10, 1839. — CASSIN, Baird's Birds N. Am., 42, 1859.

Common. Breeds in January and later. Very abundant on the Upper St. John's, and especially at Lake Monroe. Saw them repeatedly dive and catch their own fish, though usually depending upon robbing the fish-hawks for them. The same fact has been reported by other observers,* although it was formerly supposed they never caught any fish themselves.

The large specimen of an eagle taken by Audubon in Kentucky, and figured and described by him as *Falco Washingtoni*, seems not to have been preserved; it is at least not known to be extant, and appears to have never been examined by any other naturalist. Audubon states that he altogether saw not "more than eight or nine" specimens,† and deemed it very rare. He does not appear, however, to have really examined but the one figured. Numerous local observers have reported it as occasional at different localities, and Mr. Cassin has doubtfully referred specimens to it taken in New Jersey. Nuttall believed the young were more or less common near Boston every winter, and considered it as "probably also indigenous to northern Europe, but confounded with the ordinary sea eagle."‡ But, as remarked by Mr. Cassin, "No specimen precisely corresponding to Mr. Audubon's bird has been obtained since its discovery, and it has latterly been looked upon by naturalists, especially in Europe, as an unusually large specimen of the white-headed eagle."§ The important point of difference between Audubon's bird and other representatives of this genus consists in the scutellation of the tarsi, which are covered in front with broad transverse scales, instead of with a great number of small irregular ones, as in other sea eagles. This, Mr. Cassin

* WILLIAM COUPER, Massachusetts Ploughman, August 26, 1870. CHARLES H. NAUMAN, on his own authority and that of Professor S. S. Haldeman, *ibid.*, September 24, 1870. HENRY REEKS, *Can. Nat.*, Vol. V, No. 1, p. 43, 1870.

† Loudon's Mag. of Nat. Hist., Vol. I, p. 116, April, 1828.

‡ Mem. Am. Acad., 1st Ser., Vol. I, p. 92, 1831.

§ Illustrations of Birds of California, Texas, etc., p. 111, 1854.

has observed, is "a character quite unusual in any rapacious bird," * though I do not see that in this respect it differs essentially from *Buteo lineatus*, *B. pennsylvanicus*, or *Circus cyaneus*, etc. Its other main point of difference from the *H. leucocephalus* is its greater size. Audubon described his bird as measuring "3 feet 7 inches in length," "10 feet 2 inches" in extent of wings, and the folded wing "32 inches." In this series of measurements there is no discrepancy between the different dimensions given — the proportions being exactly the same as in *H. leucocephalus* — that might lead to the suspicion of a typographical or other accidental error, as some writers have suggested there may be in respect to the alar extent. It is, then, either a valid species or a large individual of *H. leucocephalus*, or a large immature *H. albicilla*. Since known specimens of *H. leucocephalus* sometimes nearly approach the supposed *H. Washingtoni* in size, it seems not unreasonable, on the whole, to regard it as really a remarkably large example of *H. leucocephalus* in immature plumage. Audubon describes his bird as breeding within the United States, and hence it is hardly probable it could have been the arctic *H. albicilla*, which has never, so far as known to me, been observed so far south at any season of the year. In reference to its fishing habits, supposed by Audubon to distinctively characterize it, it is now well known that the *H. leucocephalus* will occasionally capture its own fish, instead of depending wholly upon robbing the fish-hawk for them.

Mr. Cassin further observes, † respecting the *H. Washingtoni*, that he believes it to be more nearly related to his *H. pelagica*, which he describes as "the largest of eagles," than to any other. In the same connection he judiciously remarks respecting the numerous apocryphal species of eagles on record as follows: "But there is no end to the accounts of strange eagles given by travellers and naturalists. Some of them may have reference to peculiar species which have in later times escaped attention, but the probability is they more frequently allude to accidental varieties, or that the authors describe from such reports as they had heard at second hand, or fell into error from insufficient personal observation." Many of these reports he alludes to in detail, including the reference by Captain Cook ‡ to a "black eagle" with a "white breast" seen by him at Kay's Island, on the northwest coast of America. A specimen of the *H. leucocephalus* in peculiar (probably albinic) plumage in the Museum of Comparative Zoölogy, taken in Eastern Massachusetts, seems to indicate that the eagle of Captain Cook may have been but an unusual stage of coloration of the common white-headed eagle. The Massachusetts specimen

* Baird's Birds of N. America, p. 42.

† Illust. Birds of Cal. and Texas, p. 36.

‡ Cook's Voyages, II, 352, 1784.

above referred to has the general color of the under parts white, with most of the feathers centred with spots of dusky brown of varying size, but with a nearly uniform dusky brown patch on the middle of the breast. The interscapulars are also mainly white, and the general plumage above, except the wings, more or less varied with the same color. The tail below is mottled with irregularly shaped specks and spots of dusky or black on a white ground, and above with white on a nearly black ground, and tipped with dusky. The appearance of the under side of the bird at a distance would be nearly uniform whitish.

Mr. Cassin having stated repeatedly that his *Haliaeetus pelagicus* (the *Aquila pelagica* Pallas*) is the largest and most powerful of all known eagles,† I was greatly surprised, in critically studying his description, to find it in every respect evidently far inferior in size to Audubon's bird of Washington, and scarcely equalling the *H. albicilla*, as described by himself; the folded wing, in fact, of his *H. pelagicus* is *one inch* shorter than the folded wing of his *H. albicilla*, *four inches* shorter than the wing of the *H. Washingtoni*, as measured by Audubon, and *two inches* shorter than the folded wing of several different Massachusetts specimens of *H. leucocephalus*! The length he gives of "a skin from Behring's Strait" — the only specimen, he says, at that time in America — is "about 3 feet 8 inches," which exceeds by *one inch only* the length of Audubon's *H. Washingtoni*, as given by Audubon, doubtless from the fresh bird. But the length given by Mr. Cassin for his *H. pelagicus* is evidently too great, as, taken in connection with the other measurements of the same specimen given by Cassin, if correct, it would indicate a bird of the most anomalous and improbable proportions. Mr. Cassin's erroneous conception of the gigantic size of his bird was doubtless formed from the length of his specimen, which if a flat or unfilled skin, as it probably was, must have measured several inches more than the natural length of the bird.‡ While I do not in the least question the sincerity of Mr. Cassin's belief in the large size of his bird, I have felt it proper to call the attention of future investiga-

* Zoographia Rosso-Asiatica, I, p. 343.

† "The bird which is the subject of our present article is the largest and most powerful of the eagles." — *Illust. Birds Cal. and Texas*, p. 32, first paragraph. "Even the famous condor of the Andes, the largest of vultures, scarcely exceeds him in size," etc. *Ibid.*, p. 32, third paragraph. "The largest of all known eagles, and nearly related to *H. Washingtoni* (Aud.). It differs from the latter as described by Audubon in being generally *larger*," etc. *Ibid.*, p. 38. "It is the largest of the eagles and appears to be related to the species immediately succeeding" (*H. Washingtoni*). *Ibid.*, p. 110.

‡ Pallas says of his *Aquila pelagica*, which Cassin makes identical with his *H. pelagicus*: "Caudæ 1' 1''; longitudo alæ compositæ 1'', 11'', 2''' "; which dimensions do not indicate a bird larger than average examples of *H. leucocephalus* or *H. albicilla*.

tors of this group to this evident discrepancy of proportions in Mr. Cassin's description. An error in Mr. Cassin's figure also demands attention, which is doubtless due to an inadvertency of the artist. This consists in the scales on the front of the tarsus being arranged as Mr. Cassin says he never saw in any rapacious bird, namely, continued to the toes in broad, *unbroken transverse plates*, nearly as in Audubon's figure of the *H. Washingtoni*!

92.* **Polyborus brasiliensis** Audubon. CARACARA EAGLE. "KING BUZZARD."

Milvus brasiliensis PAY, Synop. Method. Av. et Pisc., 17, No. 6, 1713.

Circus brasiliensis BRISSON, Ornithologie, I, 116, No. 31, 1760.

Falco brasiliensis GMELIN, Syst. Nat., I, 262, 1788.

Falco tharus MOLINA, Sagg. sul. Storia Nat. del Chile, 1782.

Polyborus tharus CASSIN, Illust. Birds of Cal. and Texas, 113, 1856. — CASSIN, Baird's Birds N. Am., 45, 1858.

Polyborus vulgaris VIEILLOT, Nouv. Dict., V, 257, 1816. — AUDUBON, Orn. Biog., II, 350, pl. clxi (young).

Polyborus brasiliensis AUDUBON, Synop., 4, 1839. — BONAP., Consp. Gen. Av., 13, 1850.

"Frequent at Enterprise, associating with the vultures." — Boardman.

The swallow-tailed hawk (*Nauclerus furcatus*) became more or less common early in March. I also saw a specimen of the Mississippi kite (*Ictinia mississippiensis*) at Hawkinsville, March 15th.

STRIGIDÆ.

93. **Bubo virginianus** Swainson. GREAT-HORNED OWL.

Strix virginiana GMELIN, Syst. Nat., I, 287, 1788. — WILSON, NUTTALL, AUDUBON.

Strix (Bubo) virginiana SWAINSON, Faun. Bor. Am., II, 82, 1831.

Bubo virginianus BONAPARTE, Geog. and Comp. List, 6, 1838. — AUDUBON, Synop., 29, 1839 — CASSIN, Illust. Birds Cal. and Texas, 177, 1854. — CASSIN, Baird's Birds of N. Am., 1858.

Strix bubo, var. *magellanicus* GMELIN, Syst. Nat., I, 286, 1788

Strix pythaulæ BARTRAM, Travels, 289, 1791.

Bubo ludovicianus DAUDIN, Traité d'Orn., II, p. 210, 1800.

Bubo pinicola VIEILLOT, Ois. Am. Sept., I, 51, 1807.

Strix (Bubo) arctica SWAINSON, Faun. Bor. Am., II, 86, pl. xxx, 1831.

Bubo sub-arcticus HOY, Proc. Phil. Acad. Nat. Sci., VI, 211, 1852.

Not apparently numerous. Mr. Boardman states that he saw only a single specimen, which was killed at Enterprise. I did not observe it

above Lake George, and only heard its notes a few times below. Mr. Maynard gives it as rather common about Jacksonville, and says he frequently observed it elsewhere.

Mr. Cassin has very properly remarked that different specimens of this widely distributed species vary materially in size and color, and states that after having examined a large number of specimens from many localities he believed that they were all of one species. He thought, however, that four leading varieties, which he called *atlanticus*, *pacificus*, *arcticus*, and *magellanicus*, could be distinguished. I am not disposed to regard them, however, as by any means strictly geographical, since specimens have been taken recently in Massachusetts that typically represent each of them.* While there are doubtless more or less well-marked local forms of this species, as of all other widely distributed species, many of the differences on which the different varieties have been based are probably only individual.

94.* *Scops asio* Bonaparte. MOTTLED OWL.

Strix asio LINNÉ, Syst. Nat., I, 132, 1767. — WILSON, Am. Orn., V, 83, pl. xliii, fig. 1, 1812. — AUDUBON, NUTTALL, etc.

Scops asio BONAPARTE, Geog. and Comp. List, 6, 1838. — CASSIN, Illust. Birds Cal. and Texas, 179, 1854. — CASSIN, Baird's Birds N. Am., 51, 1858. — ALLEN, Amer. Nat., II, 327, 1868.

Strix nævia GMELIN, Syst. Nat., 289, 1788. — WILSON, Am. Orn., III, 16, pl. xix, fig. 1, 1812.

Bubo striatus VIEILLOT, Ois. Am. Sept., I, 54, pl. xxi, 1807.

Ephialtes choliba LAWRENCE, Ann. N. Y. Lyc. Nat. Hist., VI, 4, 1854.

Scops McCalli CASSIN, Illust. Birds Cal. and Texas, 180, 1854. — CASSIN, Baird's Birds N. Am., 52, 1858.

Scops Kennicotti ELLIOT, Proc. Phil. Acad. Nat. Sci., 1867, 69. — IBID., Illust. Birds N. Am., pl. xi. — BAIRD, Trans. Chicago Acad. Sci., I, 311, pl. xxvii, 1869.

Specimens were procured by Mr. Maynard, by whom, and also by Mr. Boardman, it is reported as not unfrequent.

The remarkable differences in the color of the plumage this species presents has led many to suppose it embraced two well-marked species, the red stage being recognized as one and the gray or mottled as another. Gmelin described the red stage as *Strix asio* (which is the same as the *Strix asio* of Linné, and the *Scops carolinensis* of Brisson) and the gray stage as *Strix nævia*. Wilson redescribed these different stages as distinct species. Bonaparte was the first to regard them as identical, he believing

* See Part III, p. 189.

the differences in plumage to be the result of age.* The red he believed to be the young bird, and the mottled the adult, which opinion was also entertained by Audubon. During the last thirty years, however, they have been by some authors again regarded as distinct species;† by others‡ the gray were regarded as the adult and the red as the young, while some have held the opinion that the difference in color was sexual. A general survey of the facts, either on record or known to me, show that the young birds are sometimes gray and sometimes red; that red young have sometimes *red* parents and sometimes *gray*; that the female is sometimes red and sometimes gray; and also that both sexes of a mated breeding pair of old birds are sometimes alike in color and sometimes different. Hence the opinion already advanced,§ that this variation is dependent upon neither age nor sex, but is simply a case of irregular and somewhat remarkable individual variation of a single species, seems a well-founded one. But these different stages, though usually so different, are not always well marked, so that one is often at a loss to know whether to refer certain specimens to the red series or to the gray. In other words, specimens occur of every intermediate grade between the typically bright red stage and the typically gray stage.

I have already given my reasons for referring the *Scops McCalli* of Cassin to the common *S. asio*, of which it is merely the somewhat smaller southern type.§ It is also difficult to perceive wherein the *Scops Kennicotti* Elliot, known thus far from a single specimen, differs essentially from a common phase of *S. asio*.||

* "Observations on the Nomenclature of Wilson's Ornithology," Journ. Phil. Acad. Nat. Sci., 1st Ser., Vol. III, p. 357, 1824. — "Synopsis of the Birds of the United States," Annals N. Y. Lyc. Nat. Hist., Vol. II, p. 36, 1828.

† MICHNER, Dr. EZRA, "A few Facts in Relation to the Identity of the Red and Mottled Owls," Journ. Phil. Acad. Nat. Sci., 1st Ser., Vol. VII, p. 53, 1834. — HOY, Dr. P. R., "Notes on the Ornithology of Wisconsin," Proc. Phil. Acad. Nat. Sci., Vol. VI, p. 306, 1853; Ibid., Transact. Wisconsin Agr. Soc., Vol. II (1852), p. 344, 1853.

‡ CABOT, Dr. S., Jr., "Observations on the Plumage of the Red and Mottled Owls," Journ. Bost. Soc. Nat. Hist., Vol. II, p. 126, 1838.

§ ALLEN, J. A., "Notes on the Red and Mottled Owls," American Naturalist, Vol. II, p. 327, 1868.

|| Since the above was written two adult specimens of this species have been received at the Museum from Dallas, Texas, one of which is of the mottled and the other of the red type of plumage. The specimen in mottled plumage, besides being generally darker throughout than northern specimens, has also the dark markings broader and blacker. The specimen in red plumage has the red more intense than it is in specimens of the northern red type. Both the Texas specimens are a little smaller than average New England specimens.

I have seen no specimens as yet from Florida, but from Mr. Cassin having referred a specimen from Indian River, (Fla.,) provisionally to his *Scops McCalli*, they would seem to differ but little from Texas specimens, resembling them, as would be naturally expected, more than northern ones.

95.* *Syrnium nebulosum* Gray. BARRED OWL.

Strix nebulosa FORSTER, Trans. London Philos. Soc., LXII, 386, 424, 1772. — WILSON, Am. Orn., IV, 61, pl. xxxiii, fig. 2, 1812. — AUDUBON, Orn. Biog., I, 242, pl. xlv, 1832.

Syrnium nebulosum GOULD, Birds of Europe, I, pl. xlv, 1832. — AUDUBON, Synop., 27, 1839. — CASSIN, Illustr. Birds of Cal. and Texas, 184, 1854. — BREWER, N. Am. Oöl., I, 72, 1857. — CASSIN, Baird's Birds N. Am., 56, 1858.

Ulula nebulosa BONAP., Geog. and Comp. List, 7, 1838. — BONAP., Conspect. Gen. Av., I, 53, 1851.

Strix chichictli GMELIN, Syst. Nat., I, 296, 1788.

Strix acclamator BARTRAM, Travels, 289, 1791.

Strix fernandica SHAW, Gen. Zoöl., VII, 263, 1809.

Very abundant. The only species of owl at all common. Their ludicrous notes are heard at night everywhere, and not unfrequently during the day. At night they often startle the traveller by their strange utterances from the trees over his head.

The four Florida specimens of this species before me are several shades darker than New England specimens, one only of a considerable series of the latter being as dark as the lightest-colored Florida example. The Florida specimens are also a little smaller than the northern ones.

Measurements of Florida Specimens of SYRNIUM NEBULOSUM.

M. C. Z. No.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.
5241	♀	Lake Dexter	Feb. 14, '69	T. Marcy	20.00	45.75	14.00	8.75
5242	♂	"	Feb. 14, '69	"	20.00	46.25	14.00	8.75
5299	—	Enterprise	Feb. 25, '69	"	19.50	45.75	13.00	9.00
—	—	Hawkinsville	Mar. 15, '69	J. A. Allen	19.75	46.00	13.25	—

96.* *Otus brachyotus* Boie. SHORT-EARED OWL.

Strix brachyotus GMELIN, Syst. Nat., I, 263, 1788. — FORSTER, Trans. Lond. Phil. Soc., LXII, 384, 1772. — WILSON, Am. Orn., IV, 64, pl. xxxiii, fig. 3, 1812. — BONAP., Ann. N. Y. Lyc. N. Hist., II, 37, 1828. — AUDUBON, Orn. Biog., V, 373, pl. ccccxxxii, 1835. — RICH. & SWAIN., Faun. Bor. Am., I, 75, 1831.

Otus brachyotus BOIE, Isis, 1822, 549. — AUDUBON, Syn., 28, 1839. — CASSIN, Illustr. Birds Cal. and Texas, 182, 1854. — G. R. GRAY, Gen. of Birds, I, 40. — IBID., Cat. Brit. Birds, 27, 1863.

Otus palustris BREHM, Vög. Deutschl., I, 124.

Brachyotus palustris BONAP., Geog. and Comp. List, 7, 1838.

Brachyotus Cassini BREWER, Proc. Bost. Soc. Nat. Hist., V, 321, 1856. — BREWER, N. Am. Oöl., I, 68, 1857. — CASSIN, Baird's Birds N. Am., 54, 1858.

"Quite common about marshes." — *Boardman*.

Specimens of this bird from Europe, in the Museum of Comparative Zoölogy, are not appreciably different from others from various parts of the United States. Neither do the habits of the European bird appear to differ from those of the American, as some have supposed. Dr. Richardson described its principal haunts in the Fur Countries as being "dense thickets of young pine-trees or dark entangled willow clumps, where it sits on a low branch, watching assiduously for mice." But it is now well known to more commonly frequent open fields and savannas, situations similar to those the European frequents.

An interesting state of plumage of this owl is exhibited by two pairs taken on Muskeget Island, Massachusetts, about July 1, 1870, by Messrs. C. J. Maynard and William Brewster, in which the color is so light as to almost suggest their being albinos. They are many shades lighter than the specimens of this species are from the interior, and show clearly, when taken in connection with the light race of *Arvicola riparius* (*Arvicola Breweri* Baird), also occurring on this small sandy island, the effect of the combined influence of an absence of shade and the increased light caused by reflection from the light-colored sand. The influence of similar circumstances is seen on a large scale in the birds and mammals of the Colorado desert and the arid peninsula of Lower California, and in less degree on the open arid plains of the middle region of the continent.

The long-eared owl, *Otus vulgaris* Fleming,* may be expected, from its known distribution, to also occur in Florida.

97.* *Strix flammea* Linné. BARN OWL.

Strix flammea LINNÉ, Syst. Nat., I, 133, 1767. — WILSON, NUTTALL, AUDUBON (Orn. Biog.), BONAPARTE (Synop.).

Strix pratincola BONAP., Geog. and Comp. List, 7, 1838. — CASSIN, BREWER, and recent American authors.

Strix americana AUDUBON, Synop., 25, 1839.

Strix perlata BONAP., Consp. Gen. Av., I, 55, 1850.

Strix furcata TEMM., Pl. Col., I, 432.

A specimen was taken by Mr. Thaxter at St. Augustine. Mr.

* *Strix otus* LINNÉ, Faun. Suec., 24, 1761.

Strix otus americana et mexicana GMELIN, Syst. Nat., I, 288, 1788.

Strix otus WILSON, BONAP. (Synop.), NUTTALL, AUDUBON (Orn. Biog.).

Otus vulgaris FLEMING, British Animals, 60, 1828. — AUDUBON, Synop., 28, 1839. — G. R. GRAY, Gen. Birds, I, 40; Cat. Brit. Birds, 26, 1863.

Otus Wilsonianus LESSON, Traité d'Orn., I, 110, 1831. — CASSIN, BREWER, and recent American authors.

Otus americanus BONAP., Geog. and Comp. List, 7, 1838.

Maynard informs me it was said to be common, and that at Dummitt's a hollow tree was shown him in which a pair of these birds had bred for several years. Audubon also speaks of it as being common in Florida.

Respecting the numerous species of late recognized in the *Strix flammea* group of owls, Mr. Cassin has, with great propriety, remarked that naturalists have "established species on very slender characters."

As is well known, different specimens from near the same locality vary considerably in color and size, while specimens from different continents are frequently almost undistinguishable. From the considerable number of specimens I have seen from distant points, as Europe, the United States, South America, Southern Asia, the West Indies, Australia, and South Africa, I see no reason why the *Strix flammea* may not be regarded as having a nearly cosmopolitan distribution, which indeed seems to be the present opinion of several European ornithologists. Nearly the same variations in color appear to occur on each continent, the general color in specimens from near the same locality varying from yellowish rufous to pale fulvous, and the dusky spots from being large and conspicuous to nearly obsolete or entirely wanting.

COLUMBIDÆ.

98.* *Chamæpelis passerina* Swainson. GROUND DOVE.

Common, especially about cultivated grounds.

99.* *Zenædura carolinensis* Bonaparte. MOURNING DOVE.

Columba carolinensis LINNÉ, Syst. Nat., I, 286, 1766.—GMELIN, WILSON, NUTTALL, AUDUBON (Orn. Biog.).

Columba marginata LINNÉ, Syst. Nat., I, 286, 1766.

Ectopistes marginellus WOODHOUSE, Proc. Phil. Acad. Nat. Sci., Vol. VI, 104, 1852.

Zenædura carolinensis BONAPARTE, Conspect. Gen. Avium, II, 84, 1854.

Zenædura marginellus BONAPARTE, Ibid., 85.

Abundant. Among its favorite resorts are the wild orange-groves, where it feeds on the seeds of the decaying fruit. Smaller than at the north, with the metallic tints much brighter and more bronzy.

MELEAGRIDÆ.

100.* *Meleagris gallopavo* Linné. WILD TURKEY.

Meleagris gallopavo LINNÉ, Syst. Nat., 268, 1766.—GMELIN, WILSON, BONAPARTE, AUDUBON, NUTTALL, BAIRD, etc.

Meleagris americana BARTRAM, Travels, 290, 1791.

Meleagris sylvestris VIEILL., Nouv. Dict., IX, 447.

Meleagris fera VIEILL., Galerie des Ois., II, 10, pl. x, 1824.

Meleagris mexicana GOULD, Proc. Lond. Zoöl. Soc., 1856, 61. — BAIRD, Birds N. Am., 618, 1858. — COOPER & BAIRD, Orn. Cal., I, 523, 1870.

Gallopavo sylvestris, *Novæ Angliæ* RAY, Synopsis, 51, 1713. — LECONTE, Proc. Phil. Acad. Nat. Sci., IX, 179, 1857.

Common and even quite numerous in those sections where it is not too much hunted. Mr. Boardman informs me that very fat male birds often weigh twenty-five to twenty-eight pounds, but that the average weight of the males is eighteen to twenty pounds, and of the females six to ten pounds.

THE ORIGIN OF THE DOMESTIC TURKEY.

Although it had been for a long time previously vaguely conjectured that the domestic turkey did not originate from the common wild turkey of North America, it was not until about 1856 that it was fully asserted that such was not its origin. In a paper read before the Zoölogical Society of London, in April, 1856, Mr. John Gould, the well-known English ornithologist, assigned this bird to the list of those domesticated animals whose origin had become involved in obscurity. He refers, however, to the fact of its known introduction into Europe from Mexico about 1524, and to the belief, shared by all naturalists from Linné up to that time, that the domesticated turkey was derived from the wild turkey of North America. He also states that, "on account of the great differences which are met with among our domestic turkeys, and the circumstance that the wild turkeys recently imported from North America not readily associating or pairing with them," he had for some years entertained the opinion that the wild turkey of the United States was not the original of the domestic turkey. He also at this time described a single specimen of a turkey from Mexico as belonging to a species distinct from the wild turkey of the United States, to which he gave the name of *Meleagris mexicana*. It differed, however, but slightly from the northern bird, mainly in having more white on the upper tail coverts. Although he claimed that it was of larger size, his measurements indicate it to be only barely above the average, and considerably smaller than the larger specimens from the Northern States. In considering it as distinct from the common wild turkey, he seems to have been greatly influenced by the locality whence his specimen came; as he states that he hardly thinks it probable that the common turkey, "authors to the contrary, notwithstanding," ranges very far into Mexico, since it is found, he says, along the southern boundary of Canada, which is nearly two thousand miles from Mexico. He deems it unlikely that a bird inhabiting "the cold regions of Canada

should also be indigenous to the hotter country of Mexico, whence," he adds, "and not from North America, the turkey was originally introduced into Europe"; thus leaving it to be inferred that, in his opinion, the Mexican bird—his new species—was the ancestor of the domestic turkey. The facts in respect to the distribution of the wild turkey are briefly these: It exists in Canada only in the warmer portions of that country, and thence southward uninterruptedly throughout the table-lands of Mexico.

Dr. Henry Bryant, of Boston, in reviewing Mr. Gould's paper, a few months after its appearance, took exceptions to the views of that gentleman, and in referring to the two principal statements made by Mr. Gould, namely, that the wild and domestic turkeys were structurally different, and refused to breed together, Dr. Bryant thus observes: "How far climate and other influences may have affected the domestic variety in England I do not yet know, but with us neither of these two statements is correct. If it were not for the difference in the plumage it would be impossible in many cases to distinguish the two birds; and even with this aid it is sometimes very difficult to decide with certainty when the specimen is a female. . . . The wild turkey breeds here with the tame variety quite as readily as could be expected; wherever the wild turkeys are numerous, it is an ordinary occurrence for the tame hen to prefer the wild gobbler to the domestic ones. I have had in my own possession wild hens that bred with the tame gobblers, — a fact much stranger than that of the wild gobbler breeding with the tame hen. But the most satisfactory proof of their specific identity is that the offspring of mixed blood is known to be hardier and more prolific than the domestic variety, — a fact which cannot be reconciled with their specific diversity."*

Dr. Bryant's facts, with those of previous writers, seem amply sufficient to settle the question as to the origin of the domestic turkey; yet a few months later Major John LeConte, who probably at that time had not seen Dr. Bryant's remarks, published a paper entitled "Observations on the Wild Turkey, or *Gallopavo sylvestris* of Ray."† In this paper he took the ground that the tame turkey could not possibly have been derived from the wild turkey of the United States. And, if what he states in support of his opinion as facts were such, they would go far towards rendering his position a tenable one, but in reality they are but baseless, dogmatic assumptions, which not only ran counter to the then generally received opinion, but were squarely opposed to unquestionable evidence already on record. Major LeConte's opinions, notwithstanding

* Proc. Bost. Soc. Nat. Hist., Vol. VI, p. 158, March, 1857.

† Proc. Phil. Acad. Nat. Sci., Vol. IX, p. 179, September, 1857.

that they were based on groundless assumptions, as an investigation of the subject fully proves, have been so generally entertained by subsequent authors, who have accepted his statements without investigating the facts for themselves, that a careful revision of the subject is now required. Major LeConte observes: "Whoever has compared the wild turkey of the United States with the domestic animal of the same genus must have observed that there existed very striking differences between them." While asserting that "these differences do not consist of slight and unimportant particularities, but in radical disagreements, which ought to remain unchangeable under all circumstances, and which form good specific characteristics," his sole point of distinction consists "in the possession by the tame bird of an enormous palear or dewlap," which he affirms, contrary to fact, is not possessed by the wild bird.* He refers also to the conviction that had long existed in his mind, that the two birds — the wild and domestic — "were really distinct species." "More than fifty years ago," he says, "when I first saw a wild turkey, I was led to conclude that one never could have been produced from the other. I have mentioned this to many ornithologists, but no one would take the trouble to investigate the matter [!]," etc. It does not appear, however, that even with him this long-standing conviction had resulted from a thorough investigation of the subject, for he gives no detailed comparison of the two, and many of his statements are not simply erroneous, but diametrically opposed to facts previously well substantiated. He refers to the early introduction of the turkey into Europe, and to the fact that it was found by the first explorers of America in both the wild and domesticated state. He alludes also to Mr. Gould's above-cited paper, remarking respecting it that he was unable to determine whether Mr. Gould's supposed new Mexican species was the same as the *M. gallopavo*, or was the original of the domestic bird. He thought, however, that the Mexican was identical with the common wild bird. He then remarks: "I have before observed that the turkey was found domesticated among the nations of Central America. *Now the bird which we have native among us has never been domesticated. All attempts to conquer its peculiar habits have failed, notwithstanding what has been said and written on the subject to the contrary.* I DEFY ANY ONE TO SHOW A TURKEY, EVEN OF THE FIRST GENERATION, PRODUCED FROM A PAIR HATCHED FROM A WILD HEN.† We have every year in our market offered for sale birds of a very dark color, and in some degree resembling the wild species; but in every instance, by the presence of the palear, the imposition can be detected at

* It is usually, however, either entirely absent in the wild bird, or present only in a rudimentary state.

† The italicizing in this extract is of course my own.

first sight, and the cheat exposed. I have known the eggs found in the woods hatched by a domestic hen, the chickens brought up carefully, and rendered so tame and familiar as to eat out of the hand, and to show considerable pleasure whenever persons with whom they were acquainted approached them. *Yet they never would associate with the domestic turkeys, studiously avoiding their company*, and in little more than a year running off to the woods, and never again returning to the haunts of their infancy. *I know*," he continues, "*that I shall be contradicted in this statement, and many quotations from authors brought forward against me. I repeat, contrary to the assertions of many others, THAT NO ONE HAS EVER SUCCEEDED IN DOMESTICATING OUR WILD TURKEY.* I speak not only from my own personal observations, but from the undivided testimony of many southern gentlemen. The turkey of our own poultry-yards, which, when young, is difficult to bring forward, it was thought might be obtained of a hardier race by a new domestication; but every attempt has failed, nor can I find a single well-authenticated case of a mixed breed being obtained." One is certainly at a loss to know what the self-confident Major would call a well-authenticated case of a mixed breed of wild and tame turkeys, since he must have been familiar with Bonaparte's excellent account, derived mainly from notes furnished him by Mr. Audubon, of this bird given in the first volume of his continuation of Wilson's "*American Ornithology.*" In speaking of the mixing of the wild and tame turkeys, this author remarks as follows: "This crossing often occurs in countries where wild and tame turkeys are frequent; it is well known that they will readily approach each other; and such is the influence of slavery upon even the turkey, that the robust inhabitant of the forest will drive his degenerate kinsfolk from their own food and from their females, being generally welcomed by the latter and by their owners, who well know the advantage of such a connection. . . . Eggs of the wild turkey have been frequently taken from their nests and hatched under the tame hen; the young preserve a portion of their uncivilized nature, and exhibit some knowledge of the difference between themselves and their foster-mother, roosting apart from the tame ones, and in other respects showing the force of hereditary disposition. The domesticated young, reared from the eggs of the wild turkey, are often employed as decoy birds to those in a state of nature." *

Audubon, in his account of the Canada goose, also incidentally refers to the crossing of the wild and tame turkeys, in a manner that leads us to suppose that it was to his knowledge a matter of common occurrence. He says: "The crossing of the Canada goose with the com-

* Nearly the same words are used by Audubon in his *Ornithological Biography* and in his *Birds of America*.

mon domestic species has proved as advantageous as that of the wild with the tame turkey."* He also states, "My friend, Dr. Bachman, assures me that in a state of domestication the wild turkeys, *though kept separate from tame individuals*, lose the brilliancy of their plumage in the third generation, becoming plain brown, and having here and there white feathers intermixed"†

The assertions of Major LeConte are so fully controverted by previously recorded testimony that they might have been justly ignored, had they not received, as already observed, the sanction of eminent authorities, and thus have come to be more or less currently adopted. Among the first to give them support was Professor Baird, of the Smithsonian Institution. This gentleman, in his work on the "Birds of North America," published less than two years subsequently to Major LeConte's paper, cites LeConte's opinions and statements, and partially indorses them, though he had not, he says, specimens at hand of the domestic bird for comparison with the wild one. To the data for their distinction adduced by Major LeConte, he adds a statement from Bonaparte in respect to the difference in color between the domestic and wild bird; Bonaparte observing that the wild bird never has the whitish tip to the tail which distinguishes the domestic ones. Professor Baird also adds that the flesh of the two differs in color, that of the wild bird being "much darker." He adds that, upon the whole, it is exceedingly probable that they are specifically distinct. "If the dewlap," he says, "be characteristic of a species at present only known in captivity, then, as Major LeConte remarks, it should bear the name of *M. gallopavo*, as based by Linnæus essentially upon the description by Brisson of *Gallopavo sylvestris*, in which this dewlap is particularly mentioned. In this event our wild bird will be entitled to a new name," etc. Professor Baird concludes his remarks on this subject with the following ingenious theory, which has been to some extent accepted as a probably correct one. "In conclusion," he says, "I venture to suggest the following hypothesis, which, however, is not original with myself: That there are really three species of turkey, besides the *M. ocellata*, a fourth species from Central America, entirely different from the rest. That one of them, *M. americana*, is probably peculiar to the eastern half of North America; another, *M. mexicana*, belongs to Mexico, and extends along the table-lands to the Rocky Mountains, the Gila, and the Llano Estacado; and a third is the *M. gallopavo*, or domesticated bird. That it is not at all improbable that the last was originally indigenous to some one or more of the West Indian Islands, whence it was transplanted as tamed to Mexico, and from Mexico taken to Europe about A. D. 1520.

* Birds of America, Vol. VI, p. 190.

† Ibid., Vol. V, p. 55.

Finally, that the wild turkeys were probably completely exterminated by the natives, as has been the case with equally large birds in other islands, as the dodo and solitaire.* This hypothesis," he continues, "will explain the fact of our meeting nowhere at the present day any wild turkeys resembling the domestic one.† . . . The entire subject is one of much interest, and deserves to be investigated thoroughly. It is quite possible that a careful examination of the external form and habits of the New Mexican bird may do much to throw full light on the whole question."

It is not surprising that a theory presenting to the imagination so many attractive features, and indorsed by authority so eminent, should have been currently received, as has this, by those who have not had the opportunity, nor perhaps the desire, to examine the subject for themselves. But, if I mistake not, it has also been accepted as at least a *probably* correct hypothesis by many ornithologists.‡ I have, however, already adduced evidence from Bonaparte, Bachman, Audubon, and Bryant sufficient to show, not only the erroneous character of Major LeConte's fundamental proposition, to wit, that the wild turkey of the United States has never been and never can be domesticated, but that such an hypothesis as the one above quoted is wholly uncalled for. As the whole question of the origin of the domestic turkey and its relationship to the wild turkey of the United States turns, however, upon the fact of the domesticability or non-domesticability of the common wild turkey, it may perhaps be proper to bring forward some recent testimony respecting this disputed point.

I have myself always been more or less familiar with the domestic bird, and with the fact that breeds exist which closely resemble the wild bird, and which their owners claimed were one fourth, one half, or one eighth

* Mr. Darwin, in referring to this gratuitous theory, refers to the fact of the deterioration of the turkey within the tropics, and very properly to the "improbability of a bird having long ago become extinct in these large and luxuriant islands, or of its ever having been aboriginally an inhabitant of the lowlands of the tropics." (Animals and Plants under Domestication, Am. ed., Vol. I, p. 353, note.)

† But does it explain the frequent occurrence of domestic ones so closely resembling the wild ones as to be quite undistinguishable from them?

‡ Dr. Cooper, who considers the western wild turkey specifically distinct from the wild turkey of the east, appears to believe that the domestic turkey originated from the wild turkey of Mexico. He says: "It is well known that at the period of the Spanish discovery the native turkey was widely domesticated in Mexico, and was introduced thence first into Europe, and thence into North America. Furthermore, the native bird of Eastern North America does not occur in Mexico at all. The markings of the domestic turkey are sometimes exactly like those of the wild bird of Mexico, while they never assume the plumage of the wild *Meleagris gallopavo* of the north." (Orn. Cal., Vol. I, p. 523, 1870.) Dr. Cooper's last remark is unfortunately erroneous, since domestic birds do often occur, especially females, that cannot well be distinguished from wild northern birds.

wild blood, as the case might be, and which differed in habits in some respects from the common breeds. I have also been long conversant with the fact that in the Western States, and in those other parts of the country where the turkey exists in its native state, that the eggs of the wild bird are frequently taken and hatched under the domesticated turkey, the young carefully raised and held at high prices, they being considered as highly valuable for the purpose of improving the domestic breeds. In a recent correspondence with Mr. D. Darwin Hughes, an able ornithologist of Marshall, Michigan, I alluded to the fact that the domestication of the wild bird had been disputed, and requested him to give me any facts he might possess in reference to the subject. The facts given in the following extracts from his letters are fully corroborated by other private testimony in my possession.

Under date of October 25, 1869, he wrote me respecting the domestication of the wild bird as follows: "Here [Calhoun County, Michigan], where the wild bird is abundant, they mix freely with the tame ones, and it is a common thing to see large flocks of half-breeds; I have owned them myself. They are fond of roaming and are apt to stray; not to the woods exclusively, but also to other farms. I have known the pure wild bird, hatched from wild eggs and raised in the poultry-yard, to remain for years in the yard without being confined; but this is not usual. One fine gobbler, as beautiful a bird as I ever saw, was hatched from a wild egg and headed a flock of mixed turkeys in a barn-yard. He was tame, like the others, but easily distinguished by his wild plumage; at night the flock roosted in the yard, but this bird could not brook so low a perch, and when the flock went to roost he invariably took wing and perched on an immense forest-tree one fourth of a mile away, where he spent the night; but in the morning he always returned to the barn-yard. Such instances are not uncommon. The eggs are eagerly sought for for hatching, and in this manner, as I have before said, there is a liberal sprinkling of wild blood in domestic birds, where the wild birds are abundant. The eggs of the wild bird are found every year, and although I have offered at the rate of six to eight dollars per dozen for them, there is not one in my collection of eggs, which numbers over two hundred species, so eager are the finders of them to hatch them, the chicks selling for a large price."

In another letter, dated November 5, 1869, Mr. Hughes wrote me further concerning this subject, in which he remarks as follows: "I have already said that the wild bird has been so domesticated as to reproduce its kind in the poultry-yard, and inquiries made since my last letter show that in the more northern counties of the State such cases are quite common. I cannot agree with what is said in the ninth volume of the Pacific Railroad Reports (p. 617), that there is an appreciable difference in the

color of the flesh of the wild and tame birds when cooked. There probably is some difference in color, but so little that one must have very acute powers of observation to tell the difference when brought to the table roasted. There is a difference in the color of the head, caruncles, and dewlaps, as stated by Professor Baird, but with my present means of knowledge, having no fresh specimens before me, I will not undertake to describe the differences. One thing, however, should not be forgotten; that we see the tame bird under all circumstances of passion,—in fear and when proudly strutting; in short, under all the different emotions that turkeys are heirs to, while we rarely or never see the wild turkey under such varied circumstances, but only when they are terror-stricken or dead. The head and neck in the tame bird makes rapid and surprising changes in sympathy with its emotions, and it may be so, and probably is, with the wild.”

From the evidence that has now been given, it is sufficiently apparent that Major LeConte's two fundamental assumptions,—first, that the wild bird will not mix or breed with the domesticated; and, second, that the wild bird never has been and cannot be domesticated,—upon which was erected an hypothesis to explain the origin of the domesticated bird by referring it to an extinct ancestor that probably inhabited some of the West Indian Islands, are entirely groundless, and never had for their support only the negative evidence afforded by the limited experience of Major LeConte and a few of his friends.

Inasmuch as the domestic turkey was first introduced into Europe from Mexico, it may be well in this connection to inquire further into the relationship of the so-called *M. mexicana*, or Mexican turkey, to the wild turkey of the eastern part of the United States. As already stated, the *M. mexicana* was originally described by Mr. Gould from a single specimen from Mexico. This specimen differs but slightly from the common wild turkey of the eastern part of the continent. But like many other merely nominal species, this “Mexican turkey” has been since generally recognized by writers on American ornithology, doubtless mainly because its describer was deemed too eminent a naturalist to be mistaken on such a point. Its habitat has been since extended to embrace half of that portion of the continent over which the wild turkey ranges,—the entire western half of the United States; yet the point at which the habitat of the eastern species ceases and that of the western begins, no one has yet ventured to attempt to definitely indicate. It is universally conceded to be exceedingly closely allied to the *M. gallopavo*, as the latter is now defined. Though admitted provisionally as a valid species by Professor Baird in his work already cited, he says that “whether these differences can be considered as establishing a second species for the United States is a

question yet to be decided." Dr. Coues, however, in his "List of the Birds of Fort Whipple, Arizona," * says he thinks there can be no doubt respecting the propriety of separating the "western turkey from the common species of the Eastern United States"; but he has given us no information as to how great the differences between them are, or in what they consist. As mentioned by Gould and by Baird, the Mexican bird differs from the eastern one only in being lighter colored, and in having, in correlation with the generally lighter color of the plumage, the terminal band of the tail, as also the tips of the tail coverts, whitish instead of pale brown, as the eastern bird usually has them. This, however, seems by no means necessarily a specific difference, it being only a slight geographical variation, not restricted to the turkey, but which runs through most species of both birds and mammals that have the same distribution; the probable cause of which variation I have already adverted to in Part III. The common eastern turkey occasionally approaches much nearer to the so-called Mexican bird than appears to be generally supposed. According to some authors, the tip of the tail in *M. gallopavo* is never whitish, but "plain chestnut, lighter than the ground color" of the tail. Yet of five specimens in the Museum of Comparative Zoölogy from one of the Western States, probably either Ohio or Michigan, two correspond with the description of the assumed typical *M. gallopavo*, two very nearly as well with that of the so-called *M. mexicana*, and one is intermediate between them. Three of them are decidedly lighter colored, and possess a lighter terminal band to the tail than they should to correspond with the true *M. gallopavo* as recently defined. I have, on the whole, no hesitancy in referring the *M. mexicana* Gould to the *M. gallopavo* Linné. The unquestionable specific identity of the domestic turkey with the wild one of the Eastern United States, though originally derived from the Mexican bird, seems further to support this view. From the great constancy of the white on the tail and its coverts in the domestic turkey, it has been thought to more resemble the western bird, or the *M. mexicana*, than the eastern. I need, however, only to recall the testimony of Dr. Bachman, already given in discussing another point, to show that it has necessarily no such significance. It will be remembered that Dr. Bachman states that he had known the wild birds of the Atlantic States, when kept entirely by themselves, to become more or less white under confinement in three generations.† Instead of this being either a "reversion" or a distinctive specific feature, it can be regarded only as the result of a diminution of the coloring matter through degeneracy, under the influences of domestication.

* Proc. Phil. Acad. Nat. Sci., Vol. XVIII, p. 93, 1866. Republished under the title of "Prodrome of a Work on the Ornithology of Arizona Territory."

† Mr. Darwin mentions a similar fact as having happened in England. (Animals and Plants under Domestication, Vol. I, p. 354).

As the whole plumage becomes lighter, those portions that are naturally lightest are those we should expect would soonest become white; and such is actually the case. Under domestication the turkey not only degenerates in size and hardness, but is well known to soon lose much of the brilliancy of plumage that characterizes it in a state of nature. In a few generations it loses to a great extent its metallic tints, and becomes much lighter colored; the terminal band of the tail, as well as its coverts, changes to white, and in succeeding generations the cream-colored and pure white birds often seen in our poultry-yards are gradually developed.

The fact of the domestic turkey having been first introduced into Europe from Mexico, and into the United States from Europe, admits of easy explanation; since the advanced state of civilization enjoyed by the native Mexicans had enabled them to domesticate the turkey, while their more degraded neighbors of the north had accomplished nothing of the kind. The turkey having been introduced into Europe nearly a century before the establishment of permanent settlements in the northern portions of the continent, it was, of course, as naturally introduced thence into this country as were our other domesticated animals.

PERDICIDÆ.

101.* *Ortyx virginianus* Bonaparte. QUAIL.

Tetrao virginianus LINNÉ, Syst. Nat., I, 277, 1766.

Tetrao marilandicus LINNÉ, Syst. Nat., I, 277, 1766.

Ortyx borealis STEPHENS, Shaw's Zool., XI, 377, 1819.

Perdix (Ortyx) virginiana BONAP., Obs. on Wils. Nomen., Journ. Phil. Acad. Nat. Sci., 1st Ser., IV, 268, 1825.

Ortyx virginianus BAIRD, Birds N. Am., 640, 1838. — MARCH, "Notes on Birds of Jamaica," Proc. Phil. Acad. Nat. Sci., XV, 303, 1863.

Ortyx texanus LAWRENCE, Ann. N. Y. Lyc. Nat. Hist., VI, 1, 1853. — BAIRD, Birds N. Amer., 641, 1858.

Abundant.

The quails of Florida differ from those of the Northern States in being smaller, larger billed, and darker colored. While the difference in size is very appreciable, as is also that in respect to the size of the bill, — the bill being actually larger while there is a general decrease in the size of the individual, — the most marked dissimilarity is in the coloration, through the darker color of the Florida birds. In the latter the ground color above is rufous instead of ashen, as in northern specimens, and the transverse black markings are broader. In average northern specimens the transverse black bars on the lower surface of the body are scarcely half the breadth of the intervening white spaces; in the Florida specimens they are much more than half, and in some cases nearly equal them. In

general the proportion of black in the Florida *females* is the same as that in the northern *males*. There is a similar relative increase in the extent of the black markings on the wing coverts, scapulars, and interscapulars, and on the dorsal surface generally. The black border to the white throat-patch is also broader, and extends back on the sides of the head so as usually to cover the auriculars, which in average northern specimens are dark rufous. The bill is also much darker, being generally jet black; in Massachusetts specimens it is brownish black, with the tip decidedly lighter than the other parts.

The so-called Texas quail (*Ortyx texanus* Lawr.) does not differ very greatly from either the Florida or the northern ones, it combining some of the essential characters of each, but more resembling Florida specimens than northern ones. Lawrence and Baird mention the ashen or decided gray hue on portions of the dorsal surface as distinguishing it from the *O. virginianus*, which has these parts of a "dull pinkish red." "A dull pinkish red," however, is just the color of these parts in my Florida specimens; but the Massachusetts specimens, on the contrary, are ashen, as already stated, and in this respect agree with the descriptions of the Texas form, and differ from the Florida ones in the same way that the Texas ones are said to do from those of the Atlantic coast of the Middle and Southern States. In both the Florida and Texas specimens there is a similar increase in the breadth of the black transverse markings, Lawrence describing them as being twice as broad in the Texas specimens as in the northern ones.

The *Ortyx cubanensis* of Cabanis appears scarcely to differ from the quails of Florida and Texas. D'Orbigny and Lembeye were hence doubtless correct in believing the so-called *Ortyx cubanensis* to be identical with the *O. virginianus*.

The following summary of the subjoined tables shows the difference in size that obtains between northern and southern specimens, and also in the sexes. The largest Florida specimen, it will be seen, scarcely equals the smallest northern one, when those of the same sex are compared.

No. of Specimens.	Sex.	Locality.		Length.	Alar Extent.	Wing.	Tail.
7	♂	Illinois	Average	10.18	15.44	4.47	2.82
16	♂	Florida	Average	9.46	14.16	4.22	2.52
6	♂	Illinois	Average	9.83	15.10	4.36	2.67
10	♂	Florida	Average	9.37	14.02	4.17	2.54
10	♂	"	Maximum	10.00	14.50	4.40	2.77
10	♂	"	Minimum	9.00	13.10	3.35	2.50
16	♂	"	Maximum	10.00	14.75	4.50	3.00
16	♂	"	Minimum	9.00	13.80	4.00	2.30
7	♀	Illinois	Maximum	10.50	15.60	4.60	3.00
7	♀	"	Minimum	10.00	15.00	4.37	2.55
6	♀	"	Maximum	10.25	15.50	4.50	2.85
6	♀	"	Minimum	9.50	14.50	4.25	2.45

Measurements of Florida Specimens of ORTYX VIRGINIANUS.

M. C. Z. No.	Coll. No.	Sex.	Locality.	Date	Collector.	Length	Alar Extent.	Wing.	Tail.
5151	—	♂	Hibernia	Jan. 30, '69	J. A. Allen	9.25	14.75	4.15	2.40
5152	—	♂	"	Jan. 30, '69	"	9.25	14.10	4.00	2.60
5183	—	♂	"	Jan. 30, '69	"	9.00	14.00	4.00	2.30
5184	—	♂	"	Jan. 30, '69	"	9.25	14.25	4.10	2.50
5337	—	♂	Enterprise	Mar. 4, '69	"	9.65	14.50	4.40	2.40
5336	—	♂	"	Mar. 4, '69	"	9.50	14.25	4.15	2.50
10578	1990	♂	Jacksonville	Jan. 9, '69	C. J. Maynard	9.50	13.80	4.30	2.80
10579	1990	♂	"	Jan. 9, '69	"	9.35	14.15	4.45	2.53
10580	1991	♂	"	Jan. 9, '69	"	9.30	14.30	4.00	2.30
—	2547	♂	Dummitt's	Mar. 8, '69	"	10.00	14.08	4.10	2.65
—	2546	♂	"	Mar. 7, '69	"	9.30	14.05	4.25	3.00
—	2562	♂	"	Mar. 9, '69	"	9.85	13.80	4.45	2.84
10583	2472	♂	"	Feb. 24, '69	"	9.50	14.00	4.25	2.50
—	2517	♂	"	Mar. 4, '69	"	9.25	14.00	4.40	2.70
—	2561	♂	"	Mar. 9, '69	"	9.70	14.08	4.25	2.65
10581	2356	♂	"	Feb. 16, '69	"	9.70	14.50	4.50	2.70
10582	2456	♂	"	Feb. 24, '69	"	10.00	14.50	4.25	2.55
—	2795	♂	"	Feb. 16, '69	"	9.00	13.75	3.35	2.70
—	2615	♂	"	Mar. 8, '69	"	9.50	14.20	4.40	2.57
—	1993	♂	Jacksonville	Jan. 9, '69	"	9.35	14.10	4.35	2.70
—	1994	♂	"	Jan. 9, '69	"	9.40	13.10	4.35	2.65
—	1995	♂	"	Jan. 9, '69	"	9.50	13.60	4.40	2.77
5182	—	♂	Hibernia	Jan. 30, '69	J. A. Allen	9.35	14.25	4.10	2.30
5338	—	♂	Enterprise	Mar. 4, '69	"	9.00	14.00	4.05	2.33
5351	—	♂	"	Mar. 5, '69	"	9.40	14.50	4.30	2.45
5352	—	♂	"	Mar. 5, '69	"	9.25	14.15	4.15	2.47

Measurements of Northern Specimens of ORTYX VIRGINIANUS.

M. C. Z. No.	Sex.	Locality.	Date.	Collector.	Length	Alar Extent.	Wing.	Tail.
13096	♂	Northern Illinois	Jan. 18, '71	—	10.25	15.00	4.45	2.72
10410	♂	"	Jan. 18, '71	—	10.00	15.45	4.60	2.75
10408	♂	"	Jan. 18, '71	—	10.00	15.00	4.40	2.85
10411	♂	"	Jan. 18, '71	—	10.50	15.50	4.50	2.75
13099	♂	"	Jan. 18, '71	—	10.25	15.60	4.60	3.00
13098	♂	"	Jan. 18, '71	—	10.28	15.25	4.50	2.90
13097	♂	"	Jan. 18, '71	—	10.00	15.25	4.37	2.55
13101	♂	"	Jan. 18, '71	—	10.25	15.50	4.45	2.72
10407	♂	"	Jan. 18, '71	—	9.50	14.50	4.25	2.73
10409	♂	"	Jan. 18, '71	—	10.00	14.85	4.50	2.85
10412	♂	"	Jan. 18, '71	—	9.85	15.25	4.38	2.48
10406	♂	"	Jan. 18, '71	—	9.50	15.20	4.50	2.45
13100	♂	"	Jan. 18, '71	—	9.85	15.10	4.30	2.60

CHARADRIIDÆ.

102.† *Squatarola helvetica* Cuvier. BLACK-BELLIED PLOVER.

"Some remain on the shores of the Floridas in winter." — *Audubon*.*

103.† *Charadrius virginicus* Borck. GOLDEN PLOVER.

"St. Augustine; rare." — *Boardman*.

* Birds of America, Vol. V, p. 200.

- 104.* *Ægialitis vociferus* Cassin. KILLDEE PLOVER.

Abundant.

- 105.* *Ægialitis Wilsonius* Cassin. WILSON'S PLOVER.

Not recently reported as found in Florida in the winter months. Audubon observes: "While in the Floridas, near St. Augustine, in the months of December and January, I found this species much more abundant than any other." *

- 106.† *Ægialitis semipalmatus* Cabanis. SEMI-PALMATED PLOVER.

"Not uncommon at St. Augustine throughout the winter." — Boardman.

- 107.† *Ægialitis melodus* Cabanis. PIPING PLOVER.

Observed at St. Augustine in the winter months by Mr. Boardman.

HÆMATOPODIDÆ.

- 108.† *Hæmatopus palliatus* Temminck. OYSTER-CATCHER.

Given by Mr. Boardman as rare in winter at St. Augustine.

- 109.† *Streptilas interpres* Illiger. TURNSTONE.

"Rare at St. Augustine in winter." — Boardman.

SCOLOPACIDÆ.

- 110.* *Philohela minor* Gray. WOODCOCK.

More or less common. Probably resident.

111. (†?) *Gallinago Wilsoni* Bonaparte. SNIFE.

Abundant. Probably resident. Florida specimens are darker colored and have longer bills than northern ones.

- 112.† *Calidris arenaria* Illiger. SANDERLING.

"Common at St. Augustine." — Boardman. "Abundant on Indian River." — Maynard.

- 113.† *Pelidna americana* Coues. RED-BACKED SANDPIPER.

"Common." — Maynard. Boardman.

- 114.† *Ereunetes pusillus* Cassin. SEMI-PALMATED SANDPIPER.

"Common." — Maynard.

115.† *Actodromas minutilla* Coues. LEAST SANDPIPER.

"Common." — *Maynard*.

116.† *Actodromas Bonapartei* Cassin. WHITE-RUMPED SANDPIPER.

"St. Augustine." — *Audubon*.

117.* *Symphemia semipalmata* Hartlaub. WILLET.

"Indian River to St. Augustine. Breeds in March." — *Maynard*.

118.† *Gambetta flavipes* Bonaparte. YELLOW-LEGS.

Common.

119.† *Gambetta melanoleuca* Bonaparte. GREATER YELLOW-LEGS.

Common.

120.* *Tringoides macularius* Gray. SPOTTED SANDPIPER.

Common.

121.* *Limosa fedoa* Ord. MARBLED GODWIT.

Common. Reported to Mr. Maynard as common all the year near St. Augustine, but where it nested was unknown to his informants.

122.† *Numenius hudsonicus* Latham. HUDSONIAN CURLEW.

123.† *Numenius borealis* Latham. ESQUIMAUX CURLEW.

I have no knowledge of the actual occurrence of these two species in East Florida, yet they apparently must occur as winter visitors. Dr. Coues gives them as winter visitors in his South Carolina list, and they are well known to range at this season southward into the tropics.

124.† *Numenius longirostris* Wilson. LONG-BILLED CURLEW.

"Very abundant on the coast." — *Boardman*.

Several other species of this family are well known to pass through East Florida in their migrations, and perhaps a few others are winter residents there.

RECURVIROSTRIDÆ.

125.* *Himantopus nigricollis* Vieillot. BLACK-NECKED STILT.

Audubon says it is found in Florida in winter.* Mr. Boardman gives it as "quite common at Enterprise after the 15th of March."

126.† *Recurvirostra americana* Gmelin. AVOSET.

This species must occur in Florida as a winter visitor, but I have as yet seen no specimens that were collected there.

GRUIDÆ.

127.* *Crus canadensis* Temminck. BROWN CRANE.

Abundant.

In 1853, in the Proceedings of the Boston Society of Natural History,* Dr. Bryant discussed at length the question of the relationship of *G. americana* Ord to the *G. canadensis*, and arrived at the conclusion that while the young of the *G. americana*, or white whooping crane, might be brown like the mature *G. canadensis*, or sand-hill crane, that the two were distinct species; and this conclusion ornithologists seem to have generally adopted. I saw none of the white birds in Florida, where the brown were very numerous. In Iowa I have seen both, but only at a distance. The account given by Dr. Bryant of the breeding of the sand-hill crane in Florida is very complete and interesting. According to this author the eggs, two in number, are laid from early in February till about the middle of April.†

RALLIDÆ.

128.* *Rallus elegans* Audubon. MARSH HEN.

Common.

129.* *Rallus crepitans* Gmelin. CLAPPER RAIL.

Common.

130.† *Rallus virginianus* Linné. VIRGINIA RAIL.

"Common along the St. John's River." — Boardman.

131.† *Porzana carolina* Vieillot. CAROLINA RAIL.

"Common." — Maynard.

132.(†?) *Porzana noveboracensis* Cassin. YELLOW RAIL.

"Common throughout the winter along the St. John's." — Boardman.

133.† *Fulica americana* Gmelin. COOT.

Abundant. As numerous the 1st of April as during the winter.

134.* *Gallinula galeata* Bonaparte. FLORIDA GALLINULE.

Abundant.

* Vol. IV, p. 303.

† See also on this point the same Proceedings, Vol. VII, p. 14.

135.* *Gallinula martinica* Latham. PURPLE GALLINULE.

Well known as a resident bird of Florida, but not observed by either Messrs. Maynard and Boardman or myself.

ARDEIDÆ.

136.* *Demiegretta ludoviciana* Baird. LOUISIANA HERON.

Common.

137.* *Demiegretta Pealei* Baird. PEALE'S EGRET.

Several specimens of this beautiful species were brought home by Mr. Maynard from Indian River, taken at Dummitt's. This is somewhat farther north than any point from which it has been previously reported.

138.* *Garzetta candidissima* Bonaparte. LITTLE WHITE HERON.

Abundant. Breeds in February and March.

139.* *Herodias egretta* Gray. WHITE HERON.

Abundant. Breeds early in the season. At a small heronry on an islet in Lake Dexter I found several nests containing nearly fledged young, March 23d. The nests, built eight to fifteen feet above the ground, were composed of a few sticks loosely put together. Often they were placed in the tops of bushes which were thickly overgrown with woody vines. The young, when shaken from the nest, climbed through the vines, using their bills as an organ of prehension, either seizing the branches between their mandibles or hooking their bills over them, and clung so closely that it was exceedingly difficult to dislodge them.

This and the preceding species are greatly persecuted by the hunters, who sometimes destroy great numbers at their breeding places, so many of the birds being killed and the others so much alarmed, that large heronries are thus completely broken up. Some gunners make it their business to hunt them for their plumes. Some means should be devised, however, for the protection of these beautiful birds, as at their present rate of decrease their number will soon be greatly diminished.

140.* *Ardea herodias* Linneé. GREAT BLUE HERON.

Abundant. Breeds in the retired swamps, nesting in the highest cypress-trees. It is rare that more than a single nest is seen in one

tree, but often several pairs breed near each other. Young, a third grown, were met with as early as the 12th of March. This species breeds while in immature plumage, young females being found mated with adult males, and *vice versa*. The only very appreciable external sexual difference is that of size, the males, as is generally the case in this family, being much larger than the females.

141.* *Florida cærulea* Baird. SMALL BLUE HERON.

Common.

142.* *Ardetta exilis* Gray. LITTLE BITTERN.

Not common.

143.† *Botaurus lentiginosus* Stephens. BITTERN.

Very common at some localities.

144.* *Butorides virescens* Bonaparte. GREEN HERON.

Not uncommon. Smaller than northern specimens, the Florida examples being intermediate in size between those from New England and the West Indies, the latter of which are usually regarded as a distinct species, under the name of *B. brunescens*. They also decidedly approach the West Indian type in coloration.

145.* *Nycticorax griseus* Gray. NIGHT HERON.

Ardea nycticorax LINNÉ, Syst. Nat., I, 235. — WILSON, AUDUBON, NUTTALL, BONAPARTE etc.

Ardea grisea LINNÉ, Syst. Nat., I, 239, 1766.

Ardea Gardeni GMELIN, Syst. Nat., I, 644, 1788.

Nycticorax europæus STEPH., Gen. Zoöl., XI, 609, pl. xlvii.

Nycticorax americana BONAP., Geog. and Comp. List, 48, 1838.

Nycticorax Gardeni JARDINE, Notes to Wilson's Orn. — BONAP., Conspectus Gen. Avium, II, 141, 1855.

Nyctiardea Gardeni BAIRD, Birds N. Am., 678, 1858, and subsequent American authors.

I did not observe this species on the St. John's, but Mr. Maynard found it more or less common on Indian River and Mosquito Lagoon. Mr. Boardman gives it as "not rare." It is said to be resident the whole year in Florida, by Audubon.

Having compared specimens of the American night heron with others from various parts of the Old World, I see no reason for considering them specifically distinct, though so considered by all American and some European ornithologists. The differences between them are scarcely appreciable.

TANTALIDÆ.

146.* *Tantalus loculator* *Linne* WOOD IBIS. "GANNET."

Common on the Upper St. John's. In March they were undergoing their spring moult, and were consequently in poor plumage. According to Dr. Bryant, who is the first and only writer, so far as I am aware, who has minutely described their eggs and breeding habits, incubation is generally commenced by the 1st of April. Dr. Bryant visited two of their breeding places, one of which was between New Smyrna and Enterprise, in a large cypress swamp on the southern border of Lake Ashby. He estimated that a thousand pairs were breeding there.

There is a singular discrepancy in the accounts of authors in respect to the habits of this bird. Bartram mentions it as solitary in its habits, not associating in flocks. Audubon, always finding it in large flocks, calls attention to this remark of Mr. Bartram as being wholly erroneous, and regrets that his account had been so extensively copied by authors. Dr. Bryant fully corroborates Bartram's account, and censures Audubon for not remembering that birds vary in their habits at different times and places. He says he never saw it in flocks except at its breeding places, and that they usually went off and returned either singly or in pairs. I saw wood ibises more or less frequently on the Upper St. John's for four or five weeks, and only in two or three instances singly or in pairs. I almost invariably saw them in flocks, both at their feeding grounds and flying in the air, they varying in number from a dozen to a hundred. While more or less gregarious at all times, they often doubtless also separate into pairs or wander singly.

In East Florida the wood ibises are called "gannets." Under this name they were described to Audubon when he visited that country, and concerning which he remarks: "On asking the appearance of the Gannets, I was told they were large white birds, with wings black at the end, a long neck, and a large sharp bill. The description so far agreeing with that of the common gannet or solan goose, I proposed no questions respecting the legs or tail, but went off." On visiting the locality where they were said to occur, he was surprised to find the trees covered with wood ibises. He hence adds: "Now as the good people who gave the information spoke according to their knowledge, and agreeably to their custom of calling the ibises gannets, had I not gone to the pond, I might have written this day that gannets are found

in the interior of the woods in the Floridas, that they alight on trees, etc., which, if *once* published, would in all probability have gone down to future times through the medium of compilers."* Numbers of similar errors have in fact crept into our natural-history literature, and after they have become well known as such to investigators, they are perpetuated for a generation or two by superficial compilers. The same may almost equally well be said in respect to nominal species.

147.* *Ibis alba* Vieillot. WHITE IBIS.

Abundant. Towards the end of February they were moulting and in very poor plumage. Most of the young still retained their brown dress, but in a large proportion the moulting was considerably advanced. Before the end of March it was completed, and April 1st I saw large flocks passing northward high in the air, apparently migrating.

During the winter these birds have the peculiar habit, on the Upper St. John's, of daily flying up the river at evening and down again early in the morning. They usually fly very low, passing just over the tree-tops when cutting across a bend in the river, and at other times close to the water. They are hence in easy gun-shot range from the river or its banks, and, flying in dense flocks, afford fine sport to the numerous sportsmen camping along its banks, who make great havoc among them. They breed much later in the season than the herons. Dr. Bryant states that as late as the 20th of April they had not commenced laying, and that they fly up and down Indian River in the same manner as on the St. John's.† Mr. Maynard informs me he did not meet with this bird on Mosquito Lagoon.

148. *Ibis falcinellus* Linné. GLOSSY IBIS.

Tantalus mexicanus ORD., Journ. Phil. Acad. Nat. Sci., I, 53, 1817.

Ibis falcinellus BONAP., Obs. on Nomencl. Wilson's Orn., Ibid., V, 70, 1825. —
IBID., Am. Orn., IV, 23, pl. xxiii, 1831. — AUDUBON, Orn. Biog., IV, 608,
pl. ccclxxxvii, 1838.

Ibis Ordi BONAP., Geog. and Comp. List, 1838. — BAIRD, Birds N. Amer.,
685, 1858.

"Pine barrens between Lake Harney and Indian River, in the ponds, in flocks of twelve to twenty."‡

* Birds of America, Vol. VI, p. 68.

† Proc. Bost. Soc. Nat. Hist., Vol. VII, 15.

‡ The above is a memorandum of the recent occurrence of this species in East Florida, obtained from Mr. Maynard, but whether given by him on his own authority or on that of Mr. C. H. Nauman, I am at present uncertain.

ARAMIDÆ.

149.* *Aramus giganteus* Baird. CRYING BIRD. "LIMPKIN."

This singular and stupid bird is at present more or less common about the grassy lakes and bayous from Lake Dexter southward. Now that Florida has become such a favorite winter resort for health-seekers, pleasure-seekers, and sportsmen, it will be surprising if it is not soon exterminated, as it seems to have almost no fear of man or the gun. They are generally seen in pairs, rarely, however, more than a few occupying the same vicinity; and when one of a party of them is shot, the others, instead of seeking safety by flight, remain and salute the intruder with their singularly discordant cries. Their excellent flesh will tend to favor their rapid extermination. They build their nests in bushes along the river and its bayous, occasionally at a considerable height, but make no effort to conceal them. At Hawkinsville I found a newly built nest, containing a single egg, March 20th, and a few days later, at Lake Dexter, I met with young nearly full grown. Hence they must breed very early, and, perhaps, somewhat irregularly. Dr. Bryant gave the first detailed account of the habits of this bird,* to which there is little to be added. He says he found it more or less common on the St. John's from Lake George to Lake Harney, but most abundant on the Wikiva Creek, which empties into the St. John's about twenty-five miles below Enterprise. This account agrees with my own experience in respect to its distribution. I did not ascend the Wikiva, but was informed that this bird was much more abundant there than on the St. John's. Dr. Bryant says that incubation usually commences in February, and that the number of eggs it lays is very large, sometimes numbering fifteen. Its popular name in Florida is "limpkin."

Possessing many features that ally them to the rails, they in other respects resemble the herons, and especially the ibises, besides having peculiar characters which mark them as a group distinct from either.

ANATIDÆ.

150.† *Anas boschas* Linné. MALLARD.

"Common all winter in very large flocks." — Boardman. Audubon speaks of their occurring in such numbers in portions of Florida, when

* Proc. Bost. Soc. Nat. Hist., Vol. VII, p. 12.

he was there in 1831, as to darken the air, and the noise of their wings, when rising from the large submerged savannas, he compares to the rumbling of thunder. Mr. Maynard also found them in vast numbers in 1869 on Indian River.

151.† *Anas obscura* Linné. BLACK DUCK.

"Quite common." — *Maynard*.

152.† *Dafila acuta* Jenyns. PINTAIL DUCK.

"St. John's River; not common." — *Boardman*. Mr. Maynard says that on Indian River he found them in immense numbers, passing over in clouds for hours together.

153.† *Nettion carolinensis* Baird. GREEN-WINGED TEAL.

Abundant.

154.† *Querquedula cyanoptera* Cassin. RED-BREASTED TEAL.

This species was found by Mr. Maynard in great numbers in the savannas of the upper part of Indian River, but unfortunately the specimens he obtained were lost. This, I believe, is the first time it has been reported from any of the Atlantic States.

155.† *Querquedula discors* Stephens. BLUE-WINGED TEAL.

Abundant.

156.† *Spatula clypeata* Boie. SHOVELLER.

"Common." — *Maynard*. *Boardman*.

157.† *Mareca americana* Stephens. BALDPATE.

"Common." — *Boardman*.

158.* *Aix sponsa* Boie. WOOD DUCK.

Abundant. Breeds early. Saw young March 15th.

159.† *Fulix marila* Baird. SCAUP DUCK.

Anas marila LINNÉ, Syst. Nat., I, 1766, 196. — WILSON, Am. Orn., VIII, 84, pl. lxix, 1814.

Fuligula marila AUD., Birds of America, VII, 355, pl. cccxcviii, 1843.

Fuligula affinis EYTON, Mon. Anat., 157, 1838.

Fuligula mariloides VIGORS, Zoöl. Blossom, 31, 1839.

Fuligula minor GIRAUD, Birds of Long Island, 323, 1844. — BELL, Proc. Phil. Acad. Nat. Sci., I, 141, 1842.

Fulix marila et affinis BAIRD, Birds N. Amer., 791, 1858.

Very abundant. By far the most numerous duck on the St. John's River. Quite common at Jacksonville as late as April 1st.

The *Fulix*, or *Fuligula affinis* auct. is evidently only the smaller, darker southern form of the *F. marila* auct. Most of the specimens collected in Florida were of the so-called *F. affinis* type.

160.† *Aythya americana* Bonaparte. RED-HEAD.

Abundant in the marshes near St. Augustine, in 1831.—*Audubon*.*

I find the *A. vallisneria* recorded in my notes made at Jacksonville. I saw none, however, myself, but it was reported by sportsmen to not unfrequently occur there.

161.† *Bucephala albeola* Baird. BUTTER-BALL.

Observed in Florida by Audubon.†

162.† *Erismatura rubida* Bonaparte. RUDDY DUCK.

More or less common on the Lower St. John's. Also observed by Audubon when he was on the plantation of General Hernandez, in East Florida, and "in immense flocks" about a hundred miles up the St. John's River, in February, 1832.† Also obtained by Mr. Maynard at Dummitt's.

163.† *Lophodytes cucullatus* Reichenbach. HOODED MERGANSER.

"Very abundant on the coast." — *Boardman*. "Numerous at Dummitt's." — *Maynard*. Occasional on the St. John's.

Geese are currently reported by the inhabitants to occur in winter in North Florida, but I am unable to state what species. Probably *Bernicla canadensis* and *B. brenta*, and perhaps others, are at times more or less common, since they are well known to occasionally visit Cuba.

PELECANIDÆ.

164.* *Pelecanus erythrorhynchus* Gmelin. WHITE PELICAN.

"Seen in large flocks near the mouth of the St. John's all winter." — *Boardman*. "Common on Indian River. Said to breed on an island near Dummitt's, and at Jupiter Inlet." — *Maynard*.

165.* *Pelecanus fuscus* Linné. BROWN PELICAN.

"Abundant on the coast in winter." — *Boardman*.

* Birds of America, Vol. VI, p. 312.

† Ibid., p. 370.

‡ Ibid., p. 325.

SULIDÆ.

166.† *Sula bassana* *Brisson*. COMMON GANNET.

"Abundant on the coast." — *Boardman*.

167.* *Sula fusca* *Linne*. BOOBY GANNET.

A few were seen on the coast near St. Augustine by Mr. Boardman. Mr. Maynard also observed it at Cape Canaveral

PHALACROCORACIDÆ

168.* *Graculus floridanus* *Bonaparte*. FLORIDA CORMORANT.

Common on the St. John's, and, according to Mr. Boardman, abundant on the coast.

PLOTIDÆ.

169.* *Plotus anHINGA* *Linne*. SNAKE BIRD. WATER TURKEY.

Abundant. Breeds in February and March, sometimes nesting in the tops of the highest trees, and sometimes quite low. Both sexes incubate.

PROCELLARIDÆ.

170.† *Oceanites oceanica* *Coues*. WILSON'S STORMY PETREL.

"A few about the coast at Fernandina." — *Boardman*.

171.† *Puffinus major* *Fabricius*. GREATER SHEARWATER.

"A few about the coast at Fernandina." — *Boardman*.

LARIDÆ.

172.† *Larus argentatus* *Brünnich*. HERRING GULL.

Common. Seen up the St. John's as far as Hibernia.

On my voyage from New York to Augusta, Ga., on my way to Florida, small parties of these gulls, numbering usually six to twenty, were almost constantly hovering near the vessel. In the Bay of New York, as along the coast of New England, and doubtless along that of all the Atlantic States at this season, the birds in immature plumage far outnumbered the others; but a hundred miles from land all the gulls of this species seen were old birds, which accords with observations of mine made on other winter voyages in the North Atlantic. It hence appears that the young birds are less venturesome than the adult, and keep mainly near the land. This accords also with the well-known fact that young birds, in migratory species, do not generally attain so

high latitudes in the breeding season as the fully adult. It is also highly probable that, generally, the young birds of this family do not range quite so far southward in winter as the older. The mature herring gulls, so far as I had an opportunity of observing, far outnumbered the young ones along the Carolina coast and on the St. John's River.

173.† *Larus delawarensis* Ord. RING-BILLED GULL.

"Not numerous." — Boardman.

174.* *Chræcocephalus atricilla* Lawrence. LAUGHING GULL.

Common along the coast and on the Lower St. John's.

175.† *Chræcocephalus philadelphia* Lawrence. BONAPARTE'S GULL.

With the preceding, and equally numerous. Also common, according to Mr. Maynard, on Indian River.

176.† *Gelochelidon anglica* Bonaparte. MARSH TERN.

Obtained by Mr. Maynard on Indian River.

177.* *Thalasseus regius* Gambel. ROYAL TERN.

"Abundant about the coast." — Boardman. Maynard.

178. *Sterna hirundo* Linné. COMMON TERN.

"Common at Dummitt's." — Maynard.

The following table of measurements of sixty-five specimens (forty-five males and twenty females) of this species, taken in the breeding season at Muskeget Island, Massachusetts, indicates the considerable range of individual differentiation that obtains in this species. Though so great, it does appear to be greater than occurs in *Sterna macrura*, of which I have the measurements of twenty-five specimens taken at the same locality and during the same excursion, nor is it probably greater than most of the terns and gulls present, as is evidently indicated by the great number of measurements of specimens of other species of the *Laridæ* of our coast now before me.

The average dimensions of the specimens cited in the subjoined table are as follows:—

Males: Length, 14.51; alar extent, 30.72; wing, 10.47; tail, 5.80; culmen, 1.40; tarsus, .78. *Females*: Length, 13.85; alar extent, 30.59; wing, 10.57; tail, 5.74; culmen, 1.36; tarsus, .77. The extremes of the same are as follows:—

Males: Length, 13.00 to 15.77; alar extent, 29.00 to 32.00; wing, 9.65 to 11.70; tail, 5.00 (4.81?) to 7.00; bill (culmen), 1.28 to 1.55; tarsus, .70 to .87.

Females: Length, 13.10 to 15.50; alar extent, 28.20 to 32.00; wing, 9.90 to 11.50; tail, 5.20 (4.75?) to 6.11; bill (culmen), 1.25 to 1.55; tarsus .70 to .90.

*Measurements of Massachusetts Specimens of STERNA HIRUNDO,
Taken in the Breeding Season.*

M. C. Z. No.	Coll. No.	Sex.	Locality.	Date.	Collector.	Length.	Alar Extent.	Wing.	Tail.	Bill.	Tarsus.
—	814	♂	Ipswich	June 16, '68	Allen & Maynard	14.00	30.50	10.62	6.90	1.33	.73
—	807	♂	"	June 16, '68	"	14.36	29.00	10.10	6.56	1.33	.70
10477	905	♂	Wellfleet	June 26, '68	"	14.75	31.90	10.50	6.50	1.35	.80
10478	906	♂	"	June 26, '68	"	14.30	31.70	10.45	6.00	1.41	.76
10479	907	♂	"	June 26, '68	"	14.00	31.00	10.60	5.50	1.33	.74
10480	908	♂	"	June 26, '68	"	13.75	31.00	10.75	5.40	1.28	.76
—	911	♂	Muskeget Isl.	June 29, '68	"	14.75	30.50	10.50	6.00	1.30	.87
10481	913	♂	"	June 29, '68	"	14.40	29.30	10.15	6.00	1.50	.80
10485	913	♂	"	June 29, '68	"	14.80	29.30	10.40	5.50	1.45	—
10486	917	♂	"	June 29, '68	"	14.90	29.60	10.40	5.55	1.36	.85
—	920	♂	"	June 29, '68	"	14.00	30.00	11.00	5.75	1.35	.76
—	923	♂	"	June 29, '68	"	14.00	30.50	10.40	5.55	1.40	.77
10489	925	♂	"	June 29, '68	"	14.40	29.80	10.25	5.45	1.37	.81
10490	926	♂	"	June 29, '68	"	14.50	30.25	10.30	5.40	1.35	.80
—	927	♂	"	June 29, '68	"	14.60	31.00	10.50	6.00	1.40	.75
10491	928	♂	"	June 29, '68	"	14.50	30.80	10.15	5.60	1.52	.74
—	940	♂	"	June 29, '68	"	13.60	31.15	10.25	5.60	—	—
—	941	♂	"	June 30, '68	"	14.50	31.65	10.15	6.10	—	—
—	942	♂	"	June 30, '68	"	14.50	31.50	9.65	6.90	—	—
—	943	♂	"	June 30, '68	"	14.25	30.25	9.75	5.60	—	—
—	944	♂	"	June 30, '68	"	14.60	30.20	10.80	6.00	—	—
10492	945	♂	"	June 30, '68	"	14.10	30.50	10.25	5.50	—	—
—	946	♂	"	June 30, '68	"	15.50	31.85	11.30	7.00	—	.80
—	947	♂	"	June 30, '68	"	15.75	31.50	10.75	6.00	—	.80
—	949	♂	"	July 2, '68	"	13.75	29.90	10.45	5.85	—	.79
—	957	♂	"	July 2, '68	"	15.65	32.00	11.50	5.95	1.50	.75
10493	955	♂	"	July 2, '68	"	14.00	30.25	11.70	5.00	1.45	.77
10500	963	♂	"	July 2, '68	"	14.30	31.27	10.70	5.01	1.33	.77
10501	967	♂	"	July 2, '68	"	14.26	31.00	10.65	5.61	1.30	.76
—	969	♂	"	July 2, '68	"	15.60	31.60	10.85	6.70	1.30	.75
—	970	♂	"	July 2, '68	"	14.28	30.80	10.50	5.40	1.35	.81
—	971	♂	"	July 2, '68	"	14.40	31.60	10.30	5.70	1.40	.75
—	972	♂	"	July 2, '68	"	14.00	30.00	9.80	5.15	1.40	.78
—	973	♂	"	July 2, '68	"	15.00	31.00	9.90	5.80	1.50	.75
10503	975	♂	"	July 2, '68	"	15.20	30.50	10.56	5.85	1.43	.85
—	977	♂	"	July 2, '68	"	14.25	31.20	10.25	5.70	1.51	.80
10504	979	♂	"	July 2, '68	"	15.25	31.00	10.00	6.27	1.45	.81
—	980	♂	"	July 2, '68	"	14.70	30.55	10.40	5.45	1.51	.85
—	981	♂	"	July 2, '68	"	14.55	31.00	10.55	5.55	1.41	.75
—	982	♂	"	July 2, '68	"	13.00	29.00	10.30	4.81	1.35	.76
—	983	♂	"	July 2, '68	"	15.00	31.43	10.80	6.11	1.35	.75
—	993	♂	"	July 2, '68	"	14.50	31.50	10.70	5.80	1.45	.85
—	1000	♂	"	July 2, '68	"	15.77	30.00	10.50	5.75	1.55	.77
—	1002	♂	"	July 2, '68	"	14.25	31.00	10.65	5.75	1.50	.70
—	1003	♂	"	July 2, '68	"	14.00	30.30	10.35	5.60	1.41	.70
10476	904	♂	Wellfleet	June 26, '68	"	14.25	30.60	9.90	6.00	1.35	.70
10481	912	♂	Muskeget Isl.	June 29, '68	"	14.20	30.00	10.00	6.00	1.40	.76
10484	915	♂	"	June 29, '68	"	14.75	30.75	10.55	6.07	1.42	.75
10483	914	♂	"	June 29, '68	"	13.90	29.80	10.05	5.75	1.30	.75
—	918	♂	"	June 29, '68	"	13.60	28.50	10.00	5.50	1.36	.74
—	919	♂	"	June 29, '68	"	13.50	28.20	10.30	5.85	1.25	.75
—	921	♂	"	June 29, '68	"	13.60	30.25	10.25	5.65	1.26	.71
10487	922	♂	"	June 29, '68	"	13.55	30.55	10.63	5.70	1.25	.80
10488	924	♂	"	June 29, '68	"	13.10	29.50	10.50	5.20	1.35	.73
—	948	♂	"	June 30, '68	"	14.50	31.75	10.80	5.90	—	.80
10494	950	♂	"	July 2, '68	"	14.30	31.00	10.50	5.80	1.28	.80
10495	959	♂	"	July 2, '68	"	13.60	32.00	11.50	5.75	1.30	.80
—	960	♂	"	July 2, '68	"	15.50	31.75	11.25	5.41	1.43	.71
10498	965	♂	"	July 2, '68	"	13.56	30.00	10.30	5.14	1.30	.74
—	974	♂	"	July 2, '68	"	15.25	32.00	11.25	6.11	1.34	.90
—	976	♂	"	July 2, '68	"	14.20	30.20	10.26	5.55	1.31	.75
—	978	♂	"	July 2, '68	"	14.60	31.80	10.70	5.45	1.55	.80
10504	997	♂	"	July 2, '68	"	14.45	31.70	10.45	6.46	1.41	.80
10505	999	♂	"	July 2, '68	"	14.35	30.50	10.45	5.60	1.49	.80
—	1001	♂	"	July 2, '68	"	14.40	31.00	10.65	5.85	1.40	.85

179. *Sterna macrura* Naumann. ARCTIC TERN.

"Common at Dummitt's." — *Maynard*.

As already remarked under *Sterna hirundo* the individual variation in the present species is very great. The largest and smallest specimens in a series of twenty-five, taken at Muskeget Island in the breeding season measured as follows:—

Largest (♂): Length, 16.00; alar extent, 32.75; wing, 11.75; tail, 6.00.

Smallest (♀): Length, 14.33; alar extent, 27.52; wing, 9.85; tail, 4.26.

The maxima and minima of this series are as follows:—

Length, 14.10 and 17.00; alar extent, 27.52 and 32.75; wing, 9.85 and 11.84; tail, 4.26 and 8.25.

While the females average a very little smaller than the males, several of the females are very nearly as large as the largest males.

The *Sterna Forsteri* may also occur as a winter resident, but I have at present no evidence of its occurrence there at this season. A specimen from the "St. John's River, Florida," collected by Dr. Würdemann, is cited by Mr. Lawrence * and Dr. Coues † (Smithsonian collection No. 4928), but no information is given as to when it was collected.

180.* *Rhynchops nigra* Linné. BLACK SKIMMER.

Abundant on the coast, occurring in large flocks. Not observed by me on the St. John's.

COLYMBIDÆ.

181.† *Colymbus torquatus* Brünnich. LOON.

"A single specimen at Mandarin, on the Lower St. John's; abundant off Fernandina harbor." — *Maynard*.

The considerable number of specimens of this species in the Museum of Comparative Zoölogy show a wide range of individual variation. In a series of fifteen specimens from various localities in New England, but mainly from Massachusetts, the variation in the length of the folded wing amounts to twenty per cent of its average length in the whole series; in the length of the tarsus, to twenty-nine per cent; in the length of the outer toe, to thirty per cent; in the length of the head, to twenty-eight per cent; and in the length of the culmen to twenty-three per cent.

The form described some years since as *Colymbus Adamsi* seems to have been founded on very old specimens of the large northern race of *C. tor-*

* Baird's Birds of North America, p. 863.

† Proc. Acad. Nat. Sci. Phila., 1862, p. 547.

quatus, in which the color of the bill is unusually light, and the bill itself unusually produced.

182.† **Podiceps cornutus** *Latham.* HORNED GREBE.

“Not uncommon on the St. John’s.” — *Boardman.*

183.† **Podilymbus podiceps** *Lawrence.* CAROLINA GREBE.

Abundant on the St. John’s.

Résumé of the preceding Tables of Measurements, with supplemental Remarks.

The following tables present a brief summary of the measurements given in Part IV. In the first table is given the average dimensions of thirty-two species, based on specimens collected, in each case, essentially from the same locality, and generally based on twenty or more specimens, the number varying in the different species from thirteen to sixty-five specimens. In all cases where the average sexual difference in size is appreciable, the dimensions are given for each sex. In most cases very nearly all the specimens are from Eastern Massachusetts, a few being from different localities in Southern Maine, and a few from Northern Illinois. In a few species all the specimens cited are from Eastern Florida; in a few other species part of the specimens are from Southern New England and a part from Eastern Florida; but in these cases a separate average is made of those from each of the two localities. The number of the specimens on which the average is based is given in each instance.

The second table shows the range of individual variation in size in the same species, based also on the same specimens.

The third table shows the amount of geographical variation in size in specimens of the same species from northern and southern localities, these localities being generally Southern New England (Eastern Massachusetts in the main) and East Florida. Only seven species are cited, but I have traced about the same ratio of difference in a score or more of others, of which the measurements have not yet been published. Although the number of specimens compared from the two localities has in many of these cases been comparatively small, enough have been examined to show the general constancy of the variation in all the species which breed at both these localities.

It should be added that the specimens on which the generalizations

given in Table II are based were not taken at the seasons likely to give the greatest differences, the northern specimens having been taken in summer and the southern ones in winter. Had *summer* Florida specimens been used instead of *winter* specimens, the differences would have been doubtless much greater, since in some cases, and especially in the cases of *Agelæus phæniceus* and *Quiscalus purpureus*, the summer home of a part at least of the Florida specimens must have been somewhat to the northward of Florida.

I. Table showing the Average Dimensions of Thirty-two Species of American Birds, based on Measurements of Thirteen to Sixty-five Specimens of each Species.

Species.	Locality.	No. of Specimens.	Sex.	Length.	Alar Extent.	Wing.	Tail.	Tarsus.
<i>Turdus Swainsoni</i>	Southern New England	24	—	7.17	11.65	3.86	2.88	1.15
<i>Turdus Pallasi</i>	" "	46	—	7.04	11.17	3.97	2.72	1.15
<i>Turdus fuscescens</i>	" "	40	—	7.38	11.83	3.82	2.88	1.13
<i>Harporhynchus rufus</i>	" "	17	—	11.29	13.09	4.15	5.00	1.31
<i>Mimus polyglottus</i>	Florida	37	—	9.91	13.69	4.28	4.87	—
<i>Galeoscoptes carolinensis</i>	Southern New England	20	—	8.60	11.16	3.53	3.76	1.10
<i>Sialia sialis</i>	" "	20	—	6.80	11.93	3.94	2.55	.78
<i>Geothlypis trichas</i>	" "	20	—	5.10	6.93	2.17	2.00	.77
<i>Parus atricapillus</i>	Eastern Massachusetts	27	—	5.38	8.37	2.47	2.50	.70
<i>Tyrannus carolinensis</i>	Southern New England	20	—	8.00	13.77	4.49	3.30	.73
<i>Pyranga rubra</i>	" "	13	—	7.05	11.33	3.76	2.69	.75
<i>Troglodytes ædon</i>	Florida	15	—	4.89	6.61	2.05	1.80	.52
<i>Passerculus savanna</i>	Massachusetts	26	—	5.20	8.70	2.70	1.96	.84
<i>Peucaea æstivalis</i>	Florida	22	—	5.88	8.99	2.40	2.49	.70
<i>Cardinalis virginianus</i>	" "	32	—	8.46	11.43	3.63	3.87	—
	" "	26	—	8.27	11.27	3.53	3.77	—
<i>Pipilo erythrophthalmus</i>	Southern New England	30	—	8.19	11.32	3.43	3.36	1.06
	Florida	19	—	7.88	9.88	3.13	3.56	.94
<i>Hedymeles ludoviciana</i>	Southern New England	17	—	7.77	12.15	3.93	2.82	.86
<i>Icterus Baltimore</i>	" "	20	—	7.52	11.55	3.71	2.92	.92
<i>Dolichonyx oryzivorus</i>	" "	20	—	7.24	11.67	3.78	2.67	1.00
	" "	40	—	9.16	14.71	4.69	3.63	—
<i>Agelæus phæniceus</i>	" "	28	—	7.53	12.24	3.86	2.93	—
	Northern States	15	—	10.43	16.30	4.91	3.16	—
<i>Sturnella ludoviciana</i>	Florida	12	—	9.81	15.70	4.47	2.85	—
	Northern States	8	—	9.55	14.43	4.29	2.82	—
	Florida	9	—	8.96	14.09	4.22	2.57	—
	Northern States	15	—	10.43	16.30	4.91	3.16	—
<i>Quiscalus purpureus</i>	Florida	12	—	9.81	15.70	4.47	2.85	—
	Northern States	8	—	9.55	14.43	4.29	2.82	—
	Florida	9	—	8.96	14.09	4.22	2.57	—
<i>Quiscalus major</i>	" "	24	—	16.51	22.48	7.19	7.00	—
	" "	8	—	12.95	17.94	5.67	5.11	—
<i>Cyanura cristata</i>	Massachusetts	18	—	11.71	16.87	5.13	4.89	—
	Florida	11	—	10.98	15.11	4.75	5.00	—
<i>Cyanocitta floridana</i>	" "	12	—	11.74	14.44	4.41	4.80	—
	" "	7	—	17.48	28.07	9.21	6.82	—
<i>Hylotomus pileatus</i>	" "	7	—	16.44	26.80	8.98	6.54	—
<i>Picus borealis</i>	" "	28	—	8.34	14.46	4.71	3.41	—
<i>Colaptes auratus</i>	Massachusetts	18	—	12.45	19.94	6.24	4.35	—
	Florida	11	—	11.66	18.82	5.84	4.40	—
<i>Conurus carolinensis</i>	" "	19	—	13.10	21.76	7.59	—	—
	Illinois	7	—	10.18	15.44	4.47	2.82	—
<i>Oxytyx virginianus</i>	Florida	16	—	9.46	14.16	4.22	2.52	—
	Illinois	6	—	9.83	15.10	4.36	2.67	—
	Florida	10	—	9.37	14.02	4.17	2.54	—
<i>Sterna hirundo</i>	Massachusetts	45	—	14.51	30.72	10.47	5.80	.78
	" "	20	—	13.85	30.59	10.57	5.74	.77

II. Table showing the Range of Individual Variation in Thirty Species of American Birds, based on the Measurements of Thirteen to Sixty-five Specimens of each Species, collected at the same Locality.

Species.	Locality.	No. of Specimens.	Sex	Length	Alar Extent.	Wing.	Tail	Tarsus
Turdus Swainsoni . . .	Southern New England	Min. 24	—	6.62	10.75	3.47	2.40	1.02
		Max. 24	—	7.75	12.65	4.30	3.40	1.27
Turdus Pallasi . . .	" "	Min. 46	—	6.50	10.00	3.30	2.47	1.12
		Max. 46	—	7.65	12.25	3.90	3.17	1.33
Turdus fuscescens . . .	" "	Min. 40	—	6.95	10.05	3.55	2.63	1.06
		Max. 40	—	7.87	12.70	4.16	3.02	1.18
Harporhynchus rufus .	" "	Min. 17	—	10.55	12.55	3.80	4.50	1.20
		Max. 17	—	11.85	14.00	4.25	5.30	1.42
Galeoscoptes carolinensis	" "	Min. 20	—	7.00	10.50	3.25	3.35	1.05
		Max. 20	—	9.00	11.95	3.85	4.10	1.18
Mimus polyglottus . .	Florida	Min. 37	—	9.27	13.00	4.00	4.10	—
		Max. 37	—	11.00	14.75	4.75	5.15	—
Sialia sialis	Southern New England	Min. 20	—	6.10	11.10	3.85	2.33	.74
		Max. 20	—	7.00	12.55	4.10	2.77	.83
Geothlypis trichas . . .	" "	Min. 20	—	4.60	6.30	1.95	1.78	.72
		Max. 20	—	5.63	7.50	2.37	2.10	.82
Pyranga rubra	" "	Min. 13	—	6.75	10.65	3.57	2.55	.67
		Max. 13	—	7.30	11.75	4.00	2.85	.86
Parus atricapillus . . .	Eastern Massachusetts	Min. 27	—	4.70	7.50	2.33	2.15	.62
		Max. 27	—	5.75	8.60	2.63	2.67	.77
Troglodytes ædon . . .	Florida	Min. 15	—	4.30	6.10	1.90	1.30	.50
		Max. 15	—	5.10	6.95	2.44	2.40	.68
Passerculus savanna . .	Eastern Massachusetts	Min. 26	—	5.20	7.61	2.44	1.64	.75
		Max. 26	—	6.00	9.75	2.95	2.25	.88
Peucæa æstivalis . . .	Florida	Min. 22	—	5.75	7.60	2.17	2.25	—
		Max. 22	—	6.20	8.30	2.55	2.68	—
	"	Min. 32	—	7.75	11.00	3.50	3.40	.62
Cardinalis virginianus .	"	Max. 32	—	9.10	11.78	3.85	4.20	.80
	"	Min. 26	—	7.50	10.70	3.25	3.40	.62
	"	Max. 26	—	8.75	11.75	3.85	4.10	.75
	Southern New England	Min. 30	—	7.50	10.00	3.17	3.30	.98
Pipilo erythrophthalmus	" "	Max. 30	—	8.80	12.25	3.90	3.93	1.13
	Florida	Min. 19	—	7.20	9.50	2.80	3.25	.80
	"	Max. 19	—	8.50	11.30	3.50	3.90	1.09
Hedymeles ludovicianus	Southern New England	Min. 17	—	7.15	11.50	3.83	2.70	.80
	" "	Max. 17	—	8.30	12.90	4.25	3.08	.93
Icterus Baltimore . . .	" "	Min. 20	—	7.00	10.40	3.45	2.70	.83
	" "	Max. 20	—	8.00	12.00	3.85	3.10	1.02
Dolichonyx oryzivorus	" "	Min. 20	—	6.65	11.00	3.53	2.45	.98
	" "	Max. 20	—	7.70	12.15	4.00	2.82	1.15
	Massachusetts	Min. 40	—	8.40	13.95	4.43	2.99	—
Agelæus phoeniceus . .	"	Max. 40	—	9.85	15.35	5.00	3.90	—
	"	Min. 28	—	7.35	11.25	4.25	2.65	—
	"	Max. 28	—	8.55	13.55	4.43	3.15	—
	"	Min. 15	—	10.00	15.05	4.74	2.82	—
	"	Max. 15	—	11.00	17.00	5.15	3.58	—
Sturnella ludoviciana .	Florida	Min. 12	—	8.50	13.00	3.90	2.40	—
	"	Max. 12	—	9.50	14.75	4.65	2.90	—
	Northern States . . .	Min. 13	—	12.00	17.00	5.20	4.58	—
Quiscalus purpureus . .	" "	Max. 13	—	13.50	18.43	6.05	6.00	—
	Florida	Min. 23	—	11.00	15.25	5.00	4.55	—
	"	Max. 23	—	13.00	17.80	5.75	5.50	—
	"	Min. 24	—	15.50	21.10	6.25	6.25	—
Quiscalus major	"	Max. 24	—	16.80	23.50	8.35	7.60	—
	"	Min. 8	—	12.10	17.25	5.25	4.75	—
	"	Max. 8	—	13.40	18.25	5.95	5.60	—
	Massachusetts	Min. 18	—	11.00	16.00	4.33	4.25	—
Cyanura cristata	"	Max. 18	—	12.25	17.50	5.65	5.65	—
	Florida	Min. 11	—	10.70	14.75	4.00	4.80	—
	"	Max. 11	—	11.25	16.00	5.00	5.15	—
Cyanocitta floridana . .	"	Min. 12	—	11.00	13.50	4.00	4.25	—
	"	Max. 12	—	12.50	15.00	4.75	5.35	—
Tyrannus carolinensis	Southern New England	Min. 20	—	7.00	12.50	4.17	2.93	.67
	" "	Max. 20	—	8.65	14.80	4.85	3.54	.80
Picus borealis	Florida	Min. 22	—	7.90	14.10	4.40	3.15	—
	"	Max. 22	—	8.60	15.20	4.95	3.75	—

Table II. (Continued.)

Species.	Locality.		No of Spec'ns.	Sex.	Length.	Alar Extent.	Wing.	Tail.	Tarsus.
Colaptes auratus . . .	Massachusetts . . .	Min.	18	—	12.00	19.00	6.00	4.00	—
	"	Max.	18	—	13.00	20.75	6.60	4.70	—
	Florida	Min.	11	—	10.60	17.60	5.60	4.10	—
	"	Max.	11	—	12.75	19.75	6.25	4.85	—
Conurus carolinensis . .	"	Min.	19	—	12.50	21.10	7.00	—	—
	"	Max.	19	—	13.60	22.50	7.85	—	—
	Illinois	Min.	7	—	10.00	15.00	4.37	2.55	—
	"	Max.	7	—	10.50	15.60	4.60	3.00	—
Ortyx virginianus . . .	Florida	Min.	16	—	9.00	13.80	4.00	2.30	—
	"	Max.	16	—	10.00	14.75	4.50	3.00	—
	Illinois	Min.	6	—	9.50	14.50	4.25	2.45	—
	"	Max.	6	—	10.25	15.50	4.50	2.85	—
	Florida	Min.	10	—	9.00	13.10	3.35	2.50	—
	"	Max.	10	—	10.00	14.50	4.40	2.77	—
Sterna hirundo	Massachusetts . . .	Min.	45	—	13.00	29.00	9.65	4.81	—
	"	Max.	45	—	15.70	32.00	11.70	7.00	—
	"	Min.	20	—	13.10	28.20	9.90	4.75	—
	"	Max.	20	—	15.50	32.00	11.50	6.11	—

III. Table showing the Geographical Variation in Size in Seven Species of American Birds, between Specimens from Florida and the Northern States.

Species.	Locality.	No. of Spec'ns.	Sex.	Length.	Alar Extent.	Wing.	Tail.	Tarsus.
Pipilo erythrophthalmus	Southern New England	32	—	8.19	11.32	3.43	3.36	1.06
	Florida	26	—	7.88	9.88	3.13	3.56	.94
Agelaius phoeniceus . .	Southern New England	40	—	9.16	14.71	4.69	3.63	—
	South Carolina & Florida	11	—	9.02	14.41	4.62	3.61	—
Sturnella ludoviciana . .	Northern States . . .	15	—	10.43	16.30	4.91	3.16	—
	Florida	12	—	9.81	15.70	4.47	2.85	—
	Northern States . . .	8	—	9.55	14.43	4.29	2.82	—
	Florida	9	—	8.96	14.09	4.22	2.57	—
Quiscalus purpureus . . .	Northern States . . .	15	—	10.43	16.30	4.91	3.16	—
	Florida	12	—	9.81	15.70	4.47	2.85	—
	Northern States . . .	8	—	9.55	14.43	4.29	2.82	—
	Florida	9	—	8.96	14.09	4.22	2.57	—
Cyanura cristata	Massachusetts	18	—	11.71	16.87	5.13	4.89	—
	Florida	11	—	10.98	15.11	4.75	5.00	—
Colaptes auratus	Massachusetts	18	—	12.45	19.94	6.24	4.35	—
	Florida	11	—	11.66	18.82	5.84	4.40	—
	Illinois	7	—	10.18	15.44	4.47	2.82	—
Ortyx virginianus	Florida	16	—	9.46	14.16	4.22	2.52	—
	Illinois	6	—	9.83	15.10	4.36	2.67	—
	Florida	10	—	9.37	14.02	4.17	2.54	—

In the tables and remarks contained in the preceding pages many facts have been given bearing upon the subject of geographical variation in birds, and especially in reference to the differences that almost universally obtain between specimens of the same species from northern and southern localities. In addition to the smaller size of the southern specimens, — a fact which has been for some time quite generally recognized, — attention has been called to the differences in color and in the form of the bill that seem almost equally constant and easy of recognition. In several species that range in the breeding season

from Florida to Maine, a tendency to a relatively greater elongation of the tail in the Florida specimens has also been noticed, — a feature so well known to characterize a large proportion of the birds of Lower California, as pointed out some years since by Professor Baird, — but this variation is not so frequent as the differences in size, color, and in the length and form of the bill. As already remarked, the tail is not usually abbreviated proportionally to the general diminution in size in the southern or Florida forms of the birds of Eastern North America, and in some species it is actually longer than in the larger northern birds. As shown in the above tables, this is the case in *Pipilo erythrophthalmus*, *Cyanura cristata*, and *Colaptes auratus*, or in three species out of the seven cited in the last table.

In numerous instances the southern forms of the birds enumerated in Part IV of this paper have already been specifically separated from their northern relatives; and if the example of some previous writers was to be followed at least a dozen other similar species might still be added from among the birds of Florida. Some, indeed, might be referred to the already separated West Indian and Mexican or Central American so-called species rather than to the northern type. As already stated, I consider this almost universal similar variation of the southern representatives of species from their northern representatives to be the result of a law of gradual geographical differentiation, and that the interest of science is better subserved by simply recognizing these differences, and the law of geographical variation of which they are the result, than by giving to each newly discovered race a distinctive binomial name; and the more especially since in numerous instances there is the most indubitable proof of the gradual and almost imperceptibly minute intergradation of the extreme northern and extreme southern types, even in cases where they are the most widely diverse.

In conclusion, it may be stated that the differential diagnoses of the southern types, in cases where they have been specifically separated from the northern, and the comparisons of them made with the northern for the purpose of showing their specific distinctness, are in many cases admirable descriptions of the usual differences found to distinguish the Florida-born birds from their co-specific representatives born in the Northern States. These differences are commonly solely the following: In the southern types the size is smaller, the bill longer, and the colors generally darker; the latter resulting from the greater pre-

dominance of the black in those in which portions of the plumage are mottled with this color, and the greater breadth of the dark transverse bars, and the correspondingly diminished breadth of the alternating lighter ones. To illustrate this point more fully, I herewith append a list of some of the so-called species of American birds that have been specifically separated by different authors from their northern representatives, but which are in reality only the extreme southern forms of species previously well known, with which they were considered by the older writers to be specifically identical, the most of them having been separated within the last ten or fifteen years:—

Accipiter Gundlachi,	<i>separated from</i>	Accipiter Cooperi.
Accipiter fringilloides,	"	Accipiter fuscus.
Falco dominicensis,	}	" Falco sparverius.
cinnamominus et		
sparveroides, etc.		
Mimus Gundlachi	}	" Mimus polyglottus.
et Hillii, etc.		
Seiurus ludovicianus,	"	Seiurus noveboracensis.
Thryothorus Berlandieri,	"	Thryothorus ludovicianus.
Dendroeca Gundlachi, etc.	"	Dendroeca æstiva.
Chordeiles minor et	}	" Chordeiles popetue.
Gundlachi, etc.		
Antrostomus cubanensis,	"	Antrostomus vociferus.
Xanthornus affinis,	"	Icterus spurius.
Sturnella hippocrepis et	}	" Sturnella ludoviciana.
mexicana,		
Quiscalus baritus,	"	Quiscalus purpureus.
Corvus minutus,	"	Corvus americanus.
Ortyx cubanensis et texanus,	"	Ortyx virginianus.
Campephilus Bairdii,	"	Campephilus principalis.
Colaptes chrysocaulosus,	"	Colaptes auratus.
Butorides brunnescens,	"	Butorides virescens.
Actiturus longicaudus,	"	Actiturus Bartramius.
Macrorhamphus scolopaceus	"	Macrorhamphus griseus.
Charadrius tenuirostris,	"	Charadrius melodus
Larus argentatoides et	}	" Larus argentatus.
Smithsonianus,		

In other cases the arctic forms, or the northern types, having been discovered subsequently to the southern ones, these have been described as specifically distinct from the latter. The *Bucephala islandica*, sep-

arated from the *B. americana* et *clangula*, and the *Collurio excubitoroides* from the *C. ludovicianus*, will serve to indicate the class of so-called species here referred to.

The Pacific Slope of North America furnishes a similar list of species, based on either southern or northern forms of others previously known; and the middle region of the continent its list of similar nominal species, mainly based on the desert forms of widely ranging species. In the northern half of the Old World, also, have the northern and southern geographical forms of the same species been specifically separated; but it is not my intention to call farther attention to them at present.

As already remarked, the American representatives of circumpolar species differ from the European and Asiatic principally in two ways, namely, in the apparently slightly larger size of the American, and in their somewhat brighter colors; but specific separations seem to have been based almost as frequently upon some theory of geographical distribution, or upon the individual variation of single specimens, as upon the real though slight differences that frequently obtain in such cases.

PART V.

On the Geographical Distribution of the Birds of Eastern North America, with special reference to the Number and Circumscription of the Ornithological Faunæ.

1. INTRODUCTORY REMARKS.

The distribution of plants and animals in circumpolar zones over the earth's surface has been long recognized; Humboldt* first making known the fact of such a natural distribution of the plants, and Agassiz,†

* HUMBOLDT, A. VON, et BONPLAND, AIMÉ. "Essai sur la Géographie des Plantes," etc. 4to. Paris. 1805.

† AGASSIZ, LOUIS. "Essai sur la Géographie des Animaux," *Revue Suisse et Chronique Littéraire*, Tome VIII, pp. 441-452, 538-585, 1845. "Note sur la Distribution Géographique des Animaux et de l'Homme," *Bulletin de la Société des Sciences Naturelles de Neuchâtel*, Tome I, pp. 162-166, 357-361, 366-369, 1845. "Sur la Distribution Géographique actuelle et le mode de l'apparition actuelle des Animaux à la surface du Globe." *Ibid.*, Tom. 2, pp. 347-351, 1847. "Geographical Distribution of Animals," *Edinburgh New Philosophical Magazine*, Vol. XLVI, pp. 1-25, 1850. *Ibid.*, *Christian Examiner*, Vol. XLVIII, pp. 184-204, 1850. "Sketch of the Natural Provinces of the Animal World and their Relation to the different Types of Man," *Nott and Gliddon's Types of Mankind*, pp. lviii-lxxxii, 1854. Also especially insisted upon in a course of unpublished Lectures delivered before the Lowell Institute, Boston, December, 1869.

Wagner,* Dana,† and others, subsequently establishing the same in regard to animals; the distribution of both plants and animals being primarily determined by the same influences. It has been further shown that these influences are mainly climatic, temperature having been justly recognized as governing the limitation, especially in latitude, of not only the species, but of faunæ and floræ. Their limitation in longitude is likewise as directly determined by climatic influences,‡ though indirectly by physical barriers, as oceans, mountain chains, and deserts. Humidity, in many instances, is scarcely a less, and in some cases a more, powerful limiting agent than temperature, plants being highly sensitive to hygrometric conditions, and their distribution intimately affects that of animals, since the existence of the latter is dependent upon the presence of the former, and their variety and numbers upon the degree of luxuriance of the vegetation. The faunal and floral zones hence coincide in their limitation in latitude with the climatic zones, but by no means necessarily with the geographical circles; isothermal lines, and not parallels of latitude, forming their boundaries. Their limits in longitude are determined by the influence geographical barriers, especially long chains of high mountains, exert upon climate.

* WAGNER, ANDREAS. "Die geographische Verbreitung der Säugethiere," *Abhandlungen der bayerischen Akademie der Wissenschaften, Math. Phys. Classe, Band IV, Abth. I*, pp. 1-146, 2d Abth., pp. 1-108, 3d Abth., pp. 3-114. Mit 9 Karten, 1844-1846.

† U. S. Exploring Expedition Report, Crustacea, Vol. II, pp. 1451-1500, 1852.

‡ I am aware of the diversity of opinions still prevalent among naturalists in regard to the influence climate exerts in determining the geographical distribution of species, and that many writers on this subject attribute to it only a slight importance, or altogether ignore it. The limits of these preliminary remarks will not allow of an extended comparison of the views of different authors on this point, nor a detailed consideration of the supposed objections that have been raised against the proposition above expressed. I agree with Mr. Andrew Murray in his remark, that, although "various authors have endeavored to embody the differences between the faunas and floras of the different regions of the globe into some kind of system, . . . they, with one or two exceptions, have worked upon no definite principle, and the result has been a mere catalogue of regions which possessed peculiarities without distinguishing their relative importance, or their relation to each other" (*Geographical Distribution of Mammals*, p. 296, 4to, London, 1866), — a remark which unfortunately seems in some degree applicable to Mr. Murray's own generalizations. That temperature is a powerful limiting influence affecting the range of species, especially in respect to their northward and southward extension, is so easily demonstrable that I am surprised to see it still questioned. I have myself subjected this principle to a rigid examination in studying the distribution of the animals and plants of Eastern North America, and have been surprised at the exact coincidence I have almost constantly met with between their northern and southern

The uniform character of both the flora and the fauna throughout the arctic zone is one of the most striking onto-geographical features thus far known, and one of primary importance, especially when taken in connection with its relation to the faunæ and floræ of more southern latitudes. Not less significant is the fact that in the temperate zone there is still a prevalence of identical forms in each of the three northern continents, where the resemblance of the animals and plants of either continent to those of the others is far greater than is the resemblance of those of the temperate regions of either continent to those of the tropical portions of the same continent.

Different animals and plants, as every one recognizes, are differently limited in respect to their geographical range. A small proportion of the species are almost or quite cosmopolitan; others range over the greater part of the northern hemisphere, finding their southern limit of distribution near the tropics. A few are exclusively arctic, or range only over the arctic and cold-temperate zones. Many are limited to the temperate zone, throughout nearly the whole of which they find a congenial home. A large number can only exist within the tropics, often

limits of distribution and isothermal lines, they following them in all their numerous undulations, sweeping northward in the valleys and southward along the sides of mountain ranges. The occurrence on isolated alpine summits of species existing at a lower level only far to the northward, is of itself suggestive of the powerful influence temperature has on the distribution of animals and plants. In the northern hemisphere a northern fauna and flora everywhere extends along the mountains hundreds of miles to the southward of their respective limits in the adjoining plains and valleys. Various other causes have, of course, a greater or less influence in determining the range of species, but none other, on the land areas, humidity perhaps alone excepted, is nearly so potent. The want of conformity of isothermal lines with parallels of latitude has doubtless led to confusion in regard to this subject, since vain attempts have often been made to circumscribe the botanical and zoological zones by the latter.

Differences of temperature evidently explain many of the otherwise seemingly inexplicable sudden transitions in the faunæ and floræ of adjoining regions, especially in regard to the marine animals and plants, temperature forming a strong barrier to the commingling of species inhabiting the waters of opposite sides of peninsulas having a north and south trend, or such long narrow points of land as terminate the South American and African continents. Those of the one side cannot pass to the other without passing through a zone of colder water than their organization will allow them to sustain. The isotherms of the continents are widely deflected by the irregularities of the surface of the land, running nearly straight and parallel across level areas; but in mountainous districts they bend abruptly northward or southward, following along the sides of mountains instead of crossing them. In the same manner are species, and faunæ and floræ, limited, — a coincidence clearly indicative of the strong influence climates exert in determining their geographical limits.

embracing whole families, none of whose representatives are found outside of the torrid zone of a single continent. Others are again equally at home in the torrid and warm temperate zones, but which do not exist either in the arctic or cold temperate zones; others range throughout the temperate and subtorrid. Nearly an equal number, some tropical, but the greater part temperate species, range across continental areas, within which, however, they are restricted. A great number of others find their range limited in longitude to the half or the third of a continent, and others within still more circumscribed boundaries, fluvial species being frequently confined to single river basins. Through this diversity of geographical range we have what may be termed *cosmopolitan*, *semi-cosmopolitan*, *circumpolar*, *continental*, *semi-continental*, and (relatively speaking) *restricted* species. The circumpolar and the continental are again *realm* species, the semi-continental and restricted, *province* species. Rarely is any species limited to a narrower area than that of two or three faunæ or floræ. Hence faunæ and floræ — which terms, in their restricted sense, are properly applied only to the smallest of the onto-geographical divisions — are determined by the peculiar association of species, and not by the range of a single or of a few “restricted” species; hence by their general facies. Provinces, and realms, on the other hand, may have species, and even genera and families, exclusively distinctive of them. As there are cosmopolitan, circumpolar, continental, and other kinds of species, so there must be cosmopolitan, circumpolar, continental, and other kinds of genera and families; the latter, as well as species, having each a definite or specific geographical range as distinctive of them as any biological or anatomical character may be. They are each circumscribed within definite areas, beyond which their special adaptation to their natural surroundings forbids their extension, unless aided by extraneous and unusual circumstances.

The three divisions of zones, realms (or “regions”), provinces, and faunæ and floræ,* comprise the phyto-zoölogic divisions usually recog-

* *Zone*, *realm*, *region*, *kingdom*, and *province*, are terms which have been used by different authors to designate the primary natural-history divisions of the earth's surface. In deciding as to which of these terms should be exclusively applied to these divisions, not only priority of use, but appropriateness, should of course be considered, and also the sense in which they are at present currently employed, in order to avoid, as far as possible, the confusion necessarily attending changes of nomenclature. So far as priority is concerned, zone undoubtedly has the precedence, it having been used for animals by Wagner in 1844, by Agassiz in 1845, and much earlier than this by Humboldt and others in relation to the distribution of plants. It is, however, not always a strictly convenient

nized. The boundaries of realms and provinces have often been arbitrarily fixed, inasmuch as they have been frequently limited and named in conformity to the continental areas, regardless of the fundamental law of the distribution of life in circumpolar zones.*

In addition to the law of the circumpolar distribution of life in zones, another may be recognized, namely, that *of a differentiation from the north southward*, since in passing from the northern pole to the equator we meet with a constant and accelerated divergence in the character of the animals and plants of successive regions of the continent. More or less related to the last is a third law of differentiation, namely, *a divergence of the life of given portions of continental areas from that of the corresponding portions of other continents, in proportion to the oceanic space separating such corresponding regions*. As evidence of this fact we have but to compare successively the life of the north temperate, tropical, and south temperate zones of the Western hemisphere with the life of the corresponding zones of the Eastern hemisphere; or that of Australia with the life of the other continents, as a whole; or that of tropical Asia with Africa or South America. A comparison of Africa with South America, and the faunæ and floræ of islands with those of the different continents, further corroborates this law. There is, furthermore, a correlation between the diversity

term. Realm, region, fauna and flora, and province, have been also successively used in the same sense, and also for divisions of subordinate rank, and in different ways by even the same writers. In regard to the names of the divisions of the second, third, and fourth rank, there is an equal want of uniformity in the use of the terms by which they have been designated. As being most convenient and least opposed to current usage, the following schedule of names for the primary and subordinate divisions has been adopted in the present paper: —

Realms for divisions of the first rank.

Regions for divisions of the second rank.

Provinces for divisions of the third rank.

Districts for divisions of the fourth rank.

Faunæ and *floræ* for the smallest or ultimate divisions, like the bird faunæ of Eastern North America, presently to be characterized.

Intermediate divisions to some of those above mentioned may in special cases be required; but until the necessity for them is made apparent, no names for such need be proposed.

* SCLATER, P. L. "On the general Geographical Distribution of the Members of the Class Aves," Jour. of the Proc. of the Linnæan Society, Vol. II, Zoölogy, pp. 130 - 149, 1858. The divisions proposed by this author have been quite generally adopted, but without corroboration, or apparently a critical examination of their merits.

of life in a given area and the relative temperature of that area, the number of distinct forms increasing directly with the increase in the temperature, other conditions remaining essentially unchanged. The number of distinct species and geographical races also increases directly with the increase in the diversity of the conditions of life resulting from differences of geographical configuration. Hence faunæ and floræ cover a smaller area in the warm temperate and tropical latitudes than at the northward, and in a mountainous region than in a level region. Hence within the torrid zone, where a maximum temperature is generally associated with a highly diversified surface, species, genera, and families are the most numerous, and faunæ and floræ, as well as species, are ordinarily the most narrowly circumscribed.

In accordance with the facts stated above respecting the mode of the distribution of animals and plants over the earth's surface, and the zoölogical and botanical laws of the differentiation and mutual relations of the different regions, the following primary natural-history divisions may be recognized: I, an *Arctic Realm*; II, a *North Temperate Realm*; III, an *American* Tropical Realm*; IV, an *Indo-African Tropical Realm*; V, a *South American Temperate Realm*; VI, an *African Temperate Realm*; VII, an *Antarctic Realm*; VIII, an *Australian Realm*.

The *Arctic Realm* presents a nearly uniform character throughout its extent, and, though embracing several faunæ, is not divisible into

* The terms "Palæogean" and "Neogean," "Palæarctic" and "Nearctic," etc., like those of "Old World" and "New World," have been given with reference solely to the length of time the different land areas of the earth's surface have been known to the dominant race of mankind, and hence regardless of the zoölogical history of these different land areas. Modern science has taught us that the latest discovered continent (Australia) is peopled with the most ancient types of animals and plants now in existence, and that it is, zoölogically considered, the ancient continent. Also that North and South America are behind Europe, Asia, and Africa in their zoölogical and geological development, while they are far in advance of Australia. To apply the term "ancient" to what is really the most recent, and "modern" to what is mediæval, is evidently too great a misuse of language to be allowable in scientific nomenclature. The sciences of geographical zoölogy and geographical botany concern not merely the geographical distribution of the animals and plants now living, but also those of the past. If such descriptive terms as the above are to be employed, it is evidently important that they should be used in their legitimate sense. In the present paper it has hence been considered advisable to altogether discard these terms, since to use them properly would necessitate their adoption in a manner directly opposite to their original and generally accepted application.

provinces. Its southern boundary may be considered as the northern limit of forest-trees, or about the isothermal of 50° F.

The *North Temperate Realm* presents a more varied character, and is divisible into an *American Region* and an *Europæo-Asiatic Region*, each of which is divisible into provinces, districts, and faunæ and floræ. Its boundaries may be provisionally considered as the isotherms of 32° and 70° F.

The *American Tropical Realm*, and also the *Indo-African Tropical Realm*, may be regarded as bounded by the isotherms of 70° F. The first is far more homogeneous than the second. Though the American Tropical Realm is perhaps not divisible into distinct regions, it certainly embraces several provinces and districts, and is rich in faunæ and floræ. The Indo-African Tropical Realm may be divided into an African Region and an Indian Region, each composed of several provinces and districts, and a great number of faunæ and floræ.

The *South American Temperate Realm* embraces that part of South America south of the isotherm of 70° F.; the *African Temperate Realm* includes that part of Africa south of the same isotherm, whilst the *Antarctic Realm* is restricted to the antarctic islands.

The *Australian Realm*, embracing Australia, New Zealand, New Guinea, and their dependent islands, including those to the eastward as far as Timor and Celebes, is zoölogically as distinct from the other primary regions as it is in its geographical position. It is divisible into a Temperate and a Tropical Region, the former embracing New Zealand and the southern third of Australia. Each of these regions includes two or three well-marked provinces.

The above division of the earth's surface* avoids the arbitrary partitioning of an almost homogeneous Arctic Realm between two

* It is not within the scope of the present article to trace the subdivisions of the earth's surface in relation to the distribution of its organic life any further than to furnish illustrations of the general principles according to which it is believed animals and plants are distributed, and by which the land surface of the earth is divided. Determining the rank of the several divisions by the amount of variation from others they present, it is found, as indicated above, that the divisions of co-ordinate rank increase in number to the southward. The Arctic Realm is homogeneous to such an extent as not to admit of divisions of a higher grade than faunæ and floræ. In the Temperate Realm the animals and plants of the Eastern and Western hemispheres are, as a whole, so far different as to admit of the division of this zone into two grand divisions (divisions of the second order), with other divisions between these and the ultimate ones. In the Tropical Realm the differences in the life of the two hemispheres is so great as

implied totally distinct life regions, and also a similar division of the two slightly differentiated regions of the North Temperate Realm. For nearly all the species, and hence of course the genera and families, of the Arctic Realm, and a considerable percentage of the species, a larger proportion of the genera, and nearly all the families of the Temperate Realm, occur in the northern parts of both the so-called "Neogean" and "Palæogean Creations."* It is thus seen that the life of the North Temperate Realm differs far more from that of the Tropical Realms than the life of the Old World does from that of the New. Hence the subdivision of the earth's surface into primary ontological regions, according in area with the two primary divisions of the land, now so generally adopted, is contrary to the facts, since it wholly ignores the close resemblance of the animals and plants inhabiting the north temperate and arctic regions, and the striking differences between them and those of the intertropical zone. The recognition of a "Nearctic" as contradistinguished from a "Palæarctic Region" is almost equally arbitrary and at variance with the law of the distribution of life in circumpolar zones.†

Dana, in his map of the geographical distribution of marine animals,‡ divided the Tropical Zone into four subzones, — a North and a South Torrid and a North and a South Subtorrid; and each temper-

to require a division of the torrid zone, considered as a *climatic* zone, into two primary divisions, with subdivisions of each of several ranks. The south temperate (climatic) zone is similarly divisible, while the Australian Realm, from its isolated position and its remarkable individualization, forms a primary region, with subdivisions of various grades.

* Dr. Selater properly observes: "It cannot be denied that the ornithology of the Palæarctic, or great temperate region of the Old World, is more easily characterized *by what it has not rather than by what it has*. There are certainly few among the groups of birds occurring in this region which do not develop themselves [to an equal or] to a greater extent elsewhere," etc. — *Journ. of Proc. Linn. Soc., Zoölogy*, Vol. II, p. 137.

† I may here add that the homogeneousness of the life of the boreal regions has been recognized by a number of recent writers, among whom are Dr. L. K. Schmarda, Dr. Von Middendorff, and Professor Huxley, who have each recognized a circumpolar region. Professor Huxley has also called attention (see *Proc. Zoöl. Soc. Lond.*, 1868, pp. 313–319) to the wide divergence of the life of the tropics from that of the north-temperate regions of even the same continents, and the individualization of Australia and its adjacent islands. He considers that the whole surface of the globe may be "primarily subdivided into two principal arææ, — a northern and a southern," for which he has proposed the names *Arctogæa* and *Notogæa*. The latter he has divided into three regions, — *Austro-Columbia* (= Neotropical Region of Selater), *Australasia*, and *New Zealand*.

‡ Rep. on Crust. of U. S. Expl. Exped. (Vol II), under Capt. Wilkes.

ate zone into five subzones, — a Temperate proper, a Subtemperate, a Warm Temperate, a Cold Temperate, and a Subfrigid. These zones are equally recognizable in the distribution of terrestrial life; but, owing to inequalities of its surface, they are of course less regular on the land than on the oceans.

The zones and subzones, or the Provinces and the minor phytological and zoölogical divisions of the globe, are usually not trenchantly defined. Their boundaries being determined by climatic conditions, the transition between adjacent zones, or between ontological divisions of whatever rank, is rarely abrupt; like the climatic zones, they blend more or less at their edges, their boundaries being strongly marked only in regions possessing a highly varied surface, as in mountainous districts. They are, nevertheless, easily recognizable, and can be approximately defined. Generally the dividing lines are more or less undulating, and, being determined indirectly by chains of mountains and other physical barriers, adjoining faunæ and floræ, and even adjoining provinces and realms, almost always interdigitate, and frequently enclose isolated areas of others, as will be presently shown in describing the ornithological faunæ of Eastern North America.

The boundaries of faunæ and floræ, like the range of species, are determined indirectly by elevations and depressions of the earth's surface, these variations in the altitude of the land producing varying conditions of temperature and humidity, which latter, as already stated, are the direct limiting influences of species, and of the botanical and zoölogical divisions of the globe. The permanency of their boundaries hence depends upon the constancy of the physiographic conditions of these areas, a migration of species, and of faunæ and floræ, necessarily following changes in these conditions. That such migrations have taken place is evident from the occurrence in the post-tertiary deposits of the warm temperate latitudes of the fossil remains of species found now only in the cold temperate and arctic regions, and in the tertiary strata of high latitudes of the remains of other species whose nearest allies are now found in the warm temperate and subtropical zones. These facts indicate clearly the great changes in temperature that have repeatedly occurred at given localities during the earth's history. In respect to existing animals, however, it is difficult to determine how much their known recession northward, as of the reindeer, for example, is due to climatic changes, and how much to

human agency, or whether it may not be due exclusively to the latter cause.

2. THE NATURAL PROVINCES OF THE NORTH AMERICAN TEMPERATE REGION.*

Before passing to the special subject of the present article, it will be necessary to consider briefly the North American continent as a whole. As already shown, North America embraces portions of three realms, the Arctic, the North Temperate, and the Tropical. It belongs mainly, however, to the North Temperate Realm, of which the temperate portions of North America form the Western Region. Within this Region may be recognized two Provinces, — an Eastern and a Western, — quite distinct from each other in their general features as well as in many special characteristics. The Eastern Province is characterized by the uniformity of its geographical and climatic features and by a corresponding uniformity in its faunal and floral aspects. The Western Province, on the other hand, is characterized by the diversity of its geographical and climatic features, — different areas situated under the same parallels differing greatly in these respects, — and by the number and small extent of its zoological and botanical areas, and its comparatively numerous restricted floræ and faunæ.

The Eastern Province† extends in the United States from the Atlantic seaboard to the vicinity of the 100th meridian, but to the northward its western boundary sweeps rapidly westward, and extends to the Rocky Mountains, whilst farther northward, where it approaches the Arctic Realm, it occupies the whole breadth of the continent. Its western border is not generally abruptly defined, and is, moreover, quite irregular, through its extension up the valleys of the numerous rivers which enter it from the westward. According to Professor Baird, its western boundary "starts on the Gulf of Mexico near the eastern border of Texas, perhaps between the Brazos and the Sabine, and follow-

* The "Districts" of the North American Region, or the ontological divisions of this region of the fourth rank, can be more conveniently characterized after the seven faunæ have been defined, to which point in the paper their consideration is accordingly deferred.

† The boundaries of these two regions have been sketched with apparent accuracy by Professor Baird. See *American Journal of Science and Arts*, 2d Series, Vol. XLI, pp. 82-85, Jan., 1866.

ing up the direction of the former river to the approaches of the Great Desert, nearly on the meridian mentioned [the 100th], proceeds northward, forced sometimes more or less westward, especially along the Platte, sometimes eastward. It crosses the Platte between Forts Kearney and Laramie and intersects the Missouri between Fort Randall and Fort Pierre, perhaps near Fort Lookout, as it is between the first mentioned two points that in ascending the river we find the change to take place in the ornithology of the country. Soon after crossing the northern boundary of the United States the line rapidly inclines westward and extends to the Rocky Mountains." To the southeastward this region embraces the whole of the United States, except perhaps the southern portion of Florida, which is decidedly West Indian and tropical in its affinities. To the northward it embraces the whole northern and eastern portions of the continent up to the Arctic Realm.

The Western Region commences at the western border of the Eastern, and extends thence to the Pacific coast. In the United States its area is about two thirds that of the Eastern Province, but a little farther to the northward it narrows rapidly, and is finally bounded in this direction by the Alaskan mountains.* To the southward it of course merges in Mexico into the Tropical Realm, but its southern limit is not as yet well known. While its varied character renders it subdivisible into several more or less distinct longitudinal areas, each of which may be again divided transversely into numerous faunæ and floræ, many species range throughout its whole extent and give to it a certain degree of homogeneousness. This portion of North America is, however, as yet too indefinitely known, geographically and meteorologically, as well as ontologically, to admit of the exact definition of its primary and ultimate life regions.

The Eastern Province, notwithstanding its larger area, has not only a less number of ornithological faunæ than the Western, but has also a smaller number of species represented in it, as well as a smaller number exclusively restricted to it. The following list of one hundred and eight species embraces most of the birds that are exclusively restricted to the Eastern Province, and hence those that distinctively characterize this Province.

* According to Mr. W. H. Dall. See Proc. Bost. Soc. Nat. Hist., Vol. XII, p. 144, Dec., 1868.

List of Species limited in their Longitudinal Distribution to the Eastern Province of the North American Temperate Region.

- | | |
|---|---|
| 1. <i>Turdus mustelinus</i> . | 40. <i>Galeoscoptes carolinensis</i> . |
| 2. <i>Sialia sialis</i> . | 41. <i>Harporhynchus rufus</i> . |
| 3. <i>Mniotilta varia</i> . | 42. <i>Thryothorus ludovicianus</i> . |
| 4. <i>Parula americana</i> . | 43. <i>Cistothorus stellaris</i> . |
| 5. <i>Prothonotaria citrea</i> . | 44. <i>Lophophanes bicolor</i> . |
| 6. <i>Geothlypis philadelphia</i> . | 45. <i>Parus hudsonicus</i> . |
| 7. <i>Oporornis agilis</i> . | 46. <i>Coturniculus passerinus</i> . |
| 8. <i>Oporornis formosa</i> . | 47. <i>Coturniculus Henslowi</i> . |
| 9. <i>Helmitherus vermivorus</i> . | 48. <i>Ammodromus caudacutus</i> . |
| 10. <i>Helminthophaga Swainsoni</i> | 49. <i>Ammodromus maritimus</i> . |
| 11. <i>Helminthophaga pinus</i> . | 50. <i>Zonotrichia albicollis</i> . |
| 12. <i>Helminthophaga chrysoptera</i> . | 51. <i>Junco hyemalis</i> . |
| 13. <i>Helminthophaga Bachmani</i> . | 52. <i>Spizella monticola</i> . |
| 14. <i>Helminthophaga ruficapilla</i> . | 53. <i>Spizella pusilla</i> . |
| 15. <i>Helminthophaga peregrina</i> . | 54. <i>Peucea aestivalis</i> . |
| 16. <i>Seiurus aurocapillus</i> . | 55. <i>Passerella iliaca</i> . |
| 17. <i>Seiurus noveboracensis</i> . | 56. <i>Euspiza americana</i> . |
| 18. <i>Dendroeca virens</i> . | 57. <i>Hedymeles ludoviciana</i> . |
| 19. <i>Dendroeca caeruleescens</i> . | 58. <i>Cyanospiza ciris</i> . |
| 20. <i>Dendroeca coronata</i> . | 59. <i>Cyanospiza cyanea</i> . |
| 21. <i>Dendroeca blackburniae</i> . | 60. <i>Cardinalis virginianus</i> . |
| 22. <i>Dendroeca castanea</i> . | 61. <i>Pipilo erythrophthalmus</i> . |
| 23. <i>Dendroeca pennsylvanica</i> . | 62. <i>Dolichonyx oryzivorus</i> . |
| 24. <i>Dendroeca caerulea</i> . | 63. <i>Icterus spurius</i> . |
| 25. <i>Dendroeca striata</i> . | 64. <i>Icterus baltimore</i> . |
| 26. <i>Dendroeca maculosa</i> . | 65. <i>Quiscalus purpureus</i> . |
| 27. <i>Dendroeca palmarum</i> . | 66. <i>Corvus ossifragus</i> . |
| 28. <i>Dendroeca dominica</i> . | 67. <i>Cyanura cristata</i> . |
| 29. <i>Dendroeca discolor</i> . | 68. <i>Sayornis fuscus</i> . |
| 30. <i>Perisoglossa tigrina</i> . | 69. <i>Campephilus principalis</i> . |
| 31. <i>Wilsonia mitrata</i> . | 70. <i>Picus borealis</i> . |
| 32. <i>Euthlypis canadensis</i> . | 71. <i>Sphyrapicus varius</i> . |
| 33. <i>Setophaga ruticilla</i> . | 72. <i>Centurus carolinus</i> . |
| 34. <i>Pyranga rubra</i> . | 73. <i>Melanerpes erythrocephalus</i> . |
| 35. <i>Pyranga aestiva</i> . | 74. <i>Colaptes auratus</i> . |
| 36. <i>Vireosylvia olivacea</i> . | 75. <i>Coccygus americanus</i> . |
| 37. <i>Vireosylvia philadelphica</i> . | 76. <i>Coccygus erythrophthalmus</i> . |
| 38. <i>Lanivireo flavifrons</i> . | 77. <i>Conurus carolinensis</i> . |
| 39. <i>Vireo noveboracensis</i> . | 78. <i>Trochilus colubris</i> . |

- | | |
|---------------------------------------|--|
| 79. ? <i>Chætura pelasgia</i> . | 94. ? <i>Actiturus Bartramius</i> . |
| 80. <i>Antrostomus vociferus</i> . | 95. <i>Limosa hudsonica</i> . |
| 81. <i>Antrostomus carolinensis</i> . | 96. ? <i>Numenius borealis</i> . |
| 82. <i>Nauclerus furcatus</i> . | 97. <i>Rallus crepitans</i> . |
| 83. <i>Ictinia mississippiensis</i> . | 98. <i>Porzana jamaicensis</i> . |
| 84. <i>Rosthramus sociabilis</i> . | 99. <i>Porzana noveboracensis</i> . |
| 85. ? <i>Tetrao canadensis</i> . | 100. <i>Gallinula galeata</i> . |
| 86. <i>Cupidonia cupido</i> . | 101. <i>Gallinula martinica</i> . |
| 87. <i>Oryx virginianus</i> . | 102. <i>Anas obscura</i> . |
| 88. <i>Grus americanus</i> . | 103. <i>Querquedula discors</i> . |
| 89. <i>Florida cærulea</i> . | 104. <i>Camptolæmus labradorius</i> . |
| 90. <i>Ibis alba</i> . | 105. <i>Sula fiber</i> . |
| 91. <i>Platalea ajaja</i> . | 106. <i>Graculus floridanus</i> . |
| 92. <i>Ægialitis Wilsonius</i> . | 107. <i>Plotus anhinga</i> . |
| 93. <i>Ægialitis melodus</i> . | 108. <i>Chræcocephalus atricilla</i> . |

3. THE ORNITHOLOGICAL FAUNÆ OF THE EASTERN PROVINCE OF THE NORTH AMERICAN TEMPERATE REGION.

Passing to the Eastern Province of the North American Region, the distribution of the birds will be now mainly considered. But a somewhat detailed comparison of the distribution of the representatives of this class with the distribution of the mammals and reptiles, and to some extent with the insects, mollusks, and plants, shows that the same divisions apply almost equally well to all. The distribution of plants, however, is everywhere greatly affected by the nature of the soil, as well as by humidity and temperature; and the character of the vegetation is also intimately connected with the distribution of the insects. The character of the soil, and especially the nature and amount of the mineral matter held in solution in the waters of the streams and lakes, has much to do with the relative abundance and distribution of the terrestrial and fluviatile mollusca, neither of which influences materially affects the distribution of the birds and mammals. The presence or absence of forests only, in respect to vegetation* and the soil, has much

* At the junction of the prairies with the eastern wooded districts there is quite an appreciable change in the fauna, especially in respect to the birds and mammals. The faunal differences between these regions, in respect to these two classes, result mainly through the addition of a relatively small number of strictly prairie species, the westward extension of none of the species of the Eastern Province wholly terminating at this point. The number of their representatives, however, becomes greatly reduced, and their distribution from being general and uniform is restricted to the belts of

influence on the distribution of the terrestrial vertebrates. The distribution of the fishes, the aquatic reptiles and certain groups of batrachians is, however, in great measure determined by the hydrographic basins. Hence we meet with relatively more restricted forms among the latter, as well as in insects, mollusks, and plants, than we find in either mammals or birds, the latter class being the most independent of all animals of geographical barriers.

It has been remarked that the great extent of the Eastern Province, as compared with the Western, is due to the great extent of the lowlands of Eastern North America, or of that area which has an elevation not exceeding eight hundred feet above the sea.* This is unquestionably the true reason, there being no highlands of sufficient altitude to interpose serious obstacles to the range of species. Some portions of this area, however, as the Arctic lowlands, do not belong to this region, while large portions of the country included in the Eastern Province more or less exceed that altitude. These differences of elevation are sufficient to cause the marked interdigitation of the faunæ of contiguous regions lying under the same parallels, as in the Eastern United States, where the upper portions of the Appalachian system support a Canadian or subalpine fauna and flora as far south as Georgia. Yet this elevation, in consequence of its nearly meridional trend and its lack of perfect continuity, forms a barrier to but few vertebrates except the strictly aquatic ones. If, however, the trend of the Appalachian range had been an easterly and westerly one, the influence of these highlands as a geographical barrier would have been most marked. Without the differences in altitude it affords, the faunæ and floræ of Eastern North

forest skirting the streams. At the eastern limit of the prairies, in fact, the distinctively western species begin to appear, thence westward few additional western species being met with till the edge of the great central plateau of the continent is reached, where the differentiation is further increased not only by the addition of many new forms, but by the gradual disappearance of eastern types. Whether the addition of a few prairie species be sufficient reason for recognizing a western subdivision of each of the faunæ of the Eastern Province may perhaps be thought questionable.

The forested portion of the Eastern Province also presents a lack of total uniformity between its eastern and western portions, a few species of birds occurring east of the Appalachians in the Eastern States only as stragglers, whilst they are quite common west of these highlands. About half a dozen species avoid the region circumscribed by the valleys of the St. Lawrence, Lake Champlain, the Hudson River, and the Atlantic Coast, that are found west of this area.

* See Baird, *Am. Journ. of Science and Arts*, 2d Series, Vol. XLI, p. 86.

America would have extended in regular and parallel zones from the Atlantic seaboard to the central plateau of the continent, whereas they now sweep far southward near the coast, and passing around the Appalachian highlands extend northward again along their western base.

In attempting to determine the number and limits of the ornithological faunæ of a large area, it is evident that the distribution of the birds in the breeding season should be taken as the basis for the investigation rather than their entire range, since during no other portion of the year can the migratory species be regarded as being at their true homes. The species numerously represented are also of far greater importance than those having but few representatives, as it is the common species which are not only the most characteristic, but those whose distribution is at present best known.

Applying the term *fauna*, when used in a special sense, to the smallest of the natural divisions in zoölogical geography, and considering faunæ to be characterized by their general facies, as determined by a peculiar assemblage of species, rather than by the restriction of a certain number of specific forms within their areas, Eastern North America may be considered as embracing seven ornithological faunæ, which occupy narrow, somewhat parallel zones or belts of varying breadth, extending from the Atlantic coast westward to the great middle plateau of the continent. The extent of each in an east and west direction is generally many times greater than what may be considered as its meridional extent. Their breadth, however, is quite unequal, not only as compared with each other, but that of the same fauna varies greatly at different points. They have their minimum breadth on the slopes of the mountains, and attain their maximum breadth on the plains. Each species having its own peculiar limits, which may or may not coincide with those of other species, it usually happens that at somewhat regular intervals, in passing either northward or southward from a given point, a greater number of species disappear at some points than at others, at which point also other species first appear. These divisional lines usually coincide with some marked physical change in the general character of the country, more especially in respect to its elevation, and form the boundaries between adjoining faunæ.* These faunal boundaries, as has been before remarked, coin-

* The first terrace of the Atlantic slope, which marks not only the transition from the tertiary deposits of the coast of the Middle and South Atlantic States to the older forma-

cide with isothermal lines. These isothermal lines, however, are not so often the yearly isotherms as those of particular seasons. While some writers have considered isocrymal lines as those having the greatest amount of limiting influence, as Dana has supposed to be the case with marine animals,* and as may be true in the case of plants, and possibly also of some terrestrial animals,† the mean temperature of the breeding season must necessarily more affect birds, especially the migratory species, than that of any other part of the year, or than the mean annual temperature. Isotherals hence most nearly coincide with the lines limiting the distribution of birds in the breeding season, and also the ornithological faunæ, since the majority of the species in the region now under consideration breed almost exclusively during the summer months, and mainly in June and July. Some breed in May, and a few of the rapacious birds in April, and even in March, but they are the exceptions to the general rule. The isothermal lines are hence adopted in the present essay in giving the boundaries of the ornithological faunæ.‡

Owing to the imperfect state of our knowledge of the summer distribution of the birds of North America, the present attempt at a defini-

tions of the interior, as well as forming the limit of steam navigation on the rivers of the lower Atlantic States, forms also the dividing line between the faunæ of the coast and those situated next to them in the interior, although having an altitude of generally less than three hundred feet. The rise from the succeeding plateau to the more abrupt slope of the Appalachians forms likewise the boundary between the second and third tiers of faunæ in the Atlantic States. The terrace forming the northern boundary of the tertiary deposits of the Gulf States, and of the lower Mississippi Valley generally, coincides likewise with faunal boundaries, as do similar slight changes in elevation elsewhere.

* See Report on the Crustacea collected by the U. S. Expl. Expd. under the command of Captain Wilkes, Vol. II, p. 1452.

† There must, however, be many exceptions, since in cold climates many mammals and all reptiles, as well as a large proportion of the mollusca and insects, hibernate, and thus are to a great degree (especially the reptiles) beyond the influence of excessive cold. In regard to plants, also, their northward range seems to be limited more by the amount of heat in summer than by the cold of winter, particularly in the case of annuals. As soon as the sum of the heat of summer is diminished to such a degree as to be insufficient to mature the plant, or to allow it to ripen its fruit, whether an annual, a shrub or tree, it must at that point cease to propagate, and there find its polar limit.

‡ Professor A. E. Verrill states that he has found the "boundaries between the Canadian and Alleghanian Faunæ" to be "coincident with a line which shall indicate a mean temperature of 50° Fahrenheit during the months of April, May, and June." *Proc. Bost. Soc. Nat. Hist.*, Vol. XII, p. 260, May, 1866.

tion of the faunæ of this region is to be regarded as merely a provisional one, to be perfected as the required data become known. The distribution in summer of the birds of the United States, even of that portion situated east of the Mississippi River, is still too little known to afford even there entirely satisfactory data. The data are tolerably full only for the region embraced between the St. Lawrence and the Upper Lakes on the north, and the Ohio River and Virginia on the south. Much is also known, however, in regard to the summer distribution of the birds in the other Atlantic States; but in respect to the whole region of the lower Mississippi and the Gulf States, the recorded facts bearing on this subject are lamentably few.* The isothermal lines of even our best climatological charts are also obviously more or less erroneous, and are nowhere laid down with sufficient detail to meet the wants of the student of zoölogical geography. The following lists of those species which by their presence or absence determine the facies of the several faunæ of the Eastern Province are hence not only often incomplete, but will in some cases, doubtless, require more or less modification as our knowledge of the subject increases.† The facts at hand for the work herein attempted are, however, far more numerous than would at first seem probable;‡ and doubtless the general conclusions reached in the following pages will be in the main substantiated by future investigations.

Beginning with Florida and passing northward, we meet with the following ornithological Faunæ:—

I. FLORIDIAN FAUNA. As stated in Part I (p. 164), that part of Florida south of Lake George in the interior, and of Cape Canaveral

* The importance of complete and carefully annotated lists of the birds of many localities in the South Atlantic and Gulf States, and in the Mississippi Valley, is hence clearly manifest. Now that the necessity of a precise knowledge of the habitats of animals is so generally recognized, it is to be hoped that every year will add something to our knowledge in regard to these regions.

† This is especially true in respect to the Floridian, Louisianian, and Carolinian Faunæ. Over this large area I have been unable to determine satisfactorily the exact southern limit of the breeding range of any species. Their northern limit, however, is readily approximately determined.

‡ See the Appendix to Part V for a list of the special papers that have been consulted in the present connection. In addition to these papers the specimens of birds in the collection of the Museum of Comparative Zoölogy have been of great use, whilst many additional facts have been gathered from correspondents and from other sources not there indicated.

on the coast, differs quite sensibly in its general faunal and floral characteristics from that part of the State situated farther to the northward, its fauna, especially the ornithological portion, having a decidedly West Indian or tropical aspect, as has also its flora. Dr. Stimpson has recently shown that on the Gulf coast of the State the southern forms, among the marine animals, extend considerably farther north than they do on the Atlantic coast; * but whether the warm waters of the Gulf of Mexico sensibly modify the land fauna of the northern coast of the Gulf sufficiently to affect the distribution of the birds is a point I have been as yet unable to determine. As it seems probable, however, that it does not to any great degree, the Floridian Fauna may accordingly be provisionally regarded as terminating near the latitude of Lake George.

The peculiarities of the bird fauna of Southern Florida in summer is still too imperfectly known to admit of the Floridian ornithological fauna being fully characterized. The occurrence within it of the following species which do not appear to extend much, if any, to the north of it, may serve for the present to distinguish this fauna from the Louisianian.

Species limited in their Northward Range by the Floridian Fauna.

- | | |
|------------------------------|------------------------------|
| 1. Vireosylvia barbatula. | 12. Starnænus cyanocephalus. |
| 2. Certhiola flaveola. | 13. Rostrhamus sociabilis. |
| 3. Icterus vulgaris. | 14. Polyborus tharus. |
| 4. Cyanocitta floridana. | 15. Aramus giganteus. |
| 5. Tyrannus dominicensis. | 16. Demiegretta Pealei. |
| 6. Coccygus minor. | 17. Demiegretta rufa. |
| 7. Crotophaga ani. | 18. Audubonia occidentalis. |
| 8. "Crotophaga rugirostris." | 19. Ibis rubra. |
| 9. Columba leucocephala. | 20. Phœnicopterus ruber. |
| 10. Zenæda amabilis. | 21. Haliplana fuliginosa. |
| 11. Oreopelia martinica. | 22. Anois stolidus. |

II. LOUISIANIAN FAUNA. The Louisianian Fauna may be provisionally considered as limited at the northward by the isothermal line of 77° F., it embracing all that part of the United States south of this line east of the Great Plains, except the Floridian Fauna. Beginning on the Atlantic coast apparently as far north as Norfolk, Virginia, it oc-

* See American Naturalist, Vol. IV, p. 536, December, 1870.

cupies a narrow belt thence southward along the coast, and in the latitude of Columbia, South Carolina, begins to expand to the westward. Farther southward its northern boundary passes to the southward of the mountains in Georgia, west of which it rises obliquely northward, and extends in a narrow point up the valley of the Mississippi as far as the mouth of the Ohio. West of the Mississippi it bends again somewhat to the southward.

The Louisianian Fauna hence embraces the coast of North Carolina, the lowlands of South Carolina and Georgia, nearly all of Alabama, all of Mississippi and Louisiana, nearly all of Arkansas, Western Tennessee, the extreme western part of Kentucky, Southern Missouri, the extreme southern part of Illinois, and a small portion of Eastern Texas. Most of the following species range throughout its entire extent, but appear farther to the northward only as stragglers. The presence of these species, and the absence of those given in the preceding list, will serve to distinguish it from the Floridian Fauna. It is similarly distinguished from the Carolinian Fauna as will be presently shown.

Species limited in their Northward Range by the Louisianian Fauna.

- | | |
|--------------------------------------|---------------------------------------|
| 1. <i>Peuceæa æstivalis</i> . | 13. <i>Chamæpelis passerina</i> . |
| 2. <i>Cyanospiza ciris</i> . | 14. <i>Cathartes atratus</i> . |
| 3. <i>Quiscalus major</i> . | 15. <i>Ictinia mississippiensis</i> . |
| 4. <i>Helminthophaga Swainsoni</i> . | 16. <i>Elanus leucurus</i> . |
| 5. <i>Helminthophaga Bachmani</i> . | 17. <i>Demiegretta ludoviciana</i> . |
| 6. ? <i>Prothonotaria citrea</i> . | 18. <i>Platalea ajaja</i> . |
| 7. <i>Dendræca dominica</i> . | 19. <i>Ibis alba</i> . |
| 8. <i>Sitta pusilla</i> . | 20. <i>Tantalus loculator</i> . |
| 9. <i>Antrostomus carolinensis</i> . | 21. <i>Porzana jamaicensis</i> . |
| 10. <i>Campephilus principalis</i> . | 22. <i>Plotus anhinga</i> . |
| 11. <i>Picus borealis</i> . | 23. <i>Graculus floridanus</i> . |
| 12. <i>Conurus carolinensis</i> . | 24. <i>Puffinus obscurus</i> . |

III. CAROLINIAN FAUNA. The Carolinian Fauna extends from the northern boundary of the Louisianian Fauna northward to about the isothermal line of 71° F. On the Atlantic coast this fauna includes Long Island and a small portion of Southeastern New York, which form its northern limit. In New Jersey it is restricted to the lowlands, extending westward in Southern Pennsylvania, Maryland, and Virginia to the Appalachian highlands. It embraces the middle por-

tion of the Carolinas, and a narrow belt in Northern Georgia. West of the Appalachian highlands its northern boundary sweeps to the northeastward as far as Northern Ohio, and thence runs westward, probably along the water-shed of that State; rising somewhat to the northward in passing farther west, it crosses Michigan near the southern border of that State, and embraces a portion of Southern Wisconsin and Southern Minnesota.

The Carolinian Fauna hence embraces Long Island and Southeastern New York, the greater part of New Jersey, all of Delaware, a small portion of Southeastern Pennsylvania, the greater part of Maryland and East Virginia, all of North Carolina, except the extreme eastern and western portions, the northwestern half of South Carolina, a narrow belt of Northern Georgia south of the mountains, the eastern part of Tennessee, the larger part of West Virginia, nearly all of Ohio and Kentucky, all of Indiana, nearly all of Illinois, a narrow strip of Michigan and Wisconsin, nearly all of Iowa and the greater part of Missouri, and the eastern portions of Nebraska and Kansas. It also occupies the lower Appalachian valleys.

It is distinguished from the Louisianian Fauna by the absence of the species mentioned in the preceding list, and by the presence of those given in the list next following. The features distinguishing it from the Alleghanian will be presently given in the diagnosis of that fauna.

*Species limited in their Northward Range by the Carolinian Fauna.**

- | | |
|---------------------------------------|---|
| 1. <i>Cardinalis virginianus</i> . | 14. <i>Corvus ossifragus</i> . |
| 2. <i>Euspiza americana</i> . | 15. <i>Centurus carolinus</i> . |
| 3. <i>Guiraca cærulea</i> . | 16. <i>Stelgidopteryx serripennis</i> . |
| 4. <i>Helmitherus vermivorus</i> . | 17. <i>Nauclerus furcatus</i> . |
| 5. <i>Icteria virens</i> . | 18. <i>Strix flammea</i> . |
| 6. <i>?Prothonotaria citrea</i> . | 19. <i>Cathartes aura</i> . |
| 7. <i>Wilsonia mitrata</i> . | 20. <i>Ægialitis Wilsonius</i> . |
| 8. <i>?Dendroeca cærulea</i> . | 21. <i>Gallinula galeata</i> . |
| 9. <i>Pyranga æstiva</i> . | 22. <i>Gallinula martinica</i> . |
| 10. <i>Mimus polyglottus</i> . | 23. <i>Garzetta candidissima</i> . |
| 11. <i>Thryothorus ludovicianus</i> . | 24. <i>Herodias egretta</i> . |
| 12. <i>Poliophtila cærulea</i> . | 25. <i>Florida cærulea</i> . |
| 13. <i>Lophophanes bicolor</i> . | 26. <i>Nyctherodius violaceus</i> . |

* A few of these species occur as stragglers in the Alleghanian Fauna.

- | | |
|--------------------------------------|-------------------------------------|
| 27. <i>Rallus elegans</i> . | 31. <i>Himantopus nigricollis</i> . |
| 28. <i>Rallus crepitans</i> . | 32. <i>Sterna aranea</i> . |
| 29. ? <i>Hæmatopus palliatus</i> | 33. <i>Rhynchops nigra</i> . |
| 30. <i>Recurvirostra americana</i> . | |

IV. ALLEGHANIAN FAUNA. The Alleghanian Fauna has the Carolinian for its southern boundary. Its northern boundary, from the ample data for its determination at the eastward, appears to nearly coincide with the isothermal line of 65° F. It is, however, an extremely irregular line, with abrupt and deep sinuosities. Beginning on the coast to the eastward of the Penobscot Bay, it sweeps first somewhat to the northeast, nearly or quite reaching Bangor; thence passing westward and southward, it follows the northern boundary of the lowlands through Southern Maine and Southern New Hampshire. In the Connecticut valley it rises farther to the northward, and in its southern descent skirts the eastern base of the Green Mountains, passing to the southward and westward of these highlands in Connecticut, and thence abruptly to the northward. Skirting the eastern boarder of the Champlain valley, it continues still northward to the valley of the St. Lawrence as far as Quebec; thence turning again southwestward, it passes along the northern border of the lowlands east of the Laurentian Hills (including the valley of the Ottawa), and crosses the southern peninsula of Michigan near the forty-fifth parallel; continuing thence northwestward it passes near Fort Ripley. Reaching the valley of the Red River of the North, it turns abruptly to the northward, enclosing the lowlands around Lake Winnipeg and embracing the valley of the Saskatchewan and those of its northern and southern branches, passing westward till it meets the higher plateau forming the eastern slope of the Rocky Mountains. This may be considered as *approximately* the northern boundary of the Alleghanian Fauna; the physical, climatic, and phyto-zoölogical character of the interior of British North America being at present too imperfectly known to render it easy to determine definitely the northwestern limit of the Alleghanian Fauna.*

* As already stated, the mean temperature of the breeding season (May, June, and July) has been taken as limiting the breeding range of the species. But this criterion associates regions which have very different *climatic* peculiarities, when the temperature of the whole year is considered, the isothermal lines diverging more widely from the isothermal or yearly lines in the interior than on the Atlantic coast. While in the Winnipeg basin the summer heat is sufficient to ripen corn and to permit of the cultiva-

The Alleghanian Fauna hence includes all of Southern New England, except the higher parts of the Green Mountain ranges, including even the southern third of Maine and a considerable part of New Hampshire and Vermont; all of New York, except the higher portions of the Adirondacks and the southeastern extremity of that State (which belongs to the Carolinian Fauna), all the lowlands of the Canadas, as far east at least as the vicinity of Quebec; the northern border of Ohio, the greater part of Wisconsin and Minnesota (in fact, very nearly all of these two States), and the valleys of the Red River of the North, the Assiniboine, and large portions of the valleys of the Saskatchewan and its two main branches, including also the extensive lowlands surrounding Lake Winnipeg. It also embraces all the Appalachian highlands southward to Georgia, except the higher parts (which belong to the Canadian Fauna), and hence includes a large part of Pennsylvania, the greater part of the highlands of Maryland, Virginia, and the Carolinas. The isolated areas within this region belonging to the Canadian Fauna are the highlands of Northeastern New York, and the most elevated parts of Pennsylvania, the Virginias, North Carolina, and Georgia. The northwestern part of New Jersey seems also to belong to the Canadian Fauna.

The Alleghanian Fauna is characterized by the absence of those species already mentioned as finding their northern limit within the Carolinian Fauna, by the presence of those mentioned below as limited in their northward range by the Alleghanian Fauna, and by the absence of a considerable number which occur abundantly in the Canadian Fauna. It is further distinguished from the Carolinian Fauna by the occurrence within it in the breeding season of the species enumer-

tion of tobacco, the winter climate is almost arctic, ice remaining in the lakes in sheltered places till late in May. Yet in summer the Winnipeg district is frequented by birds that find their northern range limited on the Atlantic coast to Southern Maine, where the winters are much shorter and the cold far less severe than on the prairies of the Saskatchewan. The same continental character of the climate of the interior is similarly seen as far south as the prairies of the Upper Mississippi, to which the northern birds descend in winter in greater numbers and with greater regularity than in the corresponding latitudes near the Atlantic coast. A limitation of the ornithological faunæ by the distribution of the birds in winter, — in other words, by their maximum range, — would hence differ considerably from the circumscription of these faunæ based on the breeding range of the species. This remark applies, of course, not only to the present fauna (Alleghanian), but to Eastern North America as a whole, especially to that portion north of the Louisianian Fauna.

ated in the second list next subjoined, to which the present fauna forms the southern limit of their breeding range.

1 *Species limited in their Northward Range by the Alleghanian Fauna.**

- | | |
|--------------------------------------|--|
| 1. <i>Turdus migratorius</i> . | 15. <i>Icterus Baltimore</i> . |
| 2. <i>Sialia sialis</i> . | 16. <i>Icterus spurius</i> . |
| 3. <i>Pyrrhula rubra</i> . | 17. <i>Sturnella ludoviciana</i> . |
| 4. <i>Dendroica discolor</i> . | 18. ? <i>Antrostomus vociferus</i> . |
| 5. <i>Lanius flavifrons</i> . | 19. <i>Zenaidura carolinensis</i> . |
| 6. <i>Vireo noveboracensis</i> . | 20. <i>Cupidonia cupido</i> . |
| 7. <i>Troglodytes aëdon</i> . | 21. <i>Ortyx virginianus</i> . |
| 8. <i>Harporhynchus rufus</i> . | 22. <i>Meleagris gallopavo</i> . |
| 9. <i>Cyanospiza cyanea</i> . | 23. <i>Ardea exilis</i> . |
| 10. <i>Pipilo erythrophthalmus</i> . | 24. <i>Rallus virginianus</i> . |
| 11. <i>Spizella pusilla</i> . | 25. <i>Chroicocephalus atricilla</i> . |
| 12. <i>Coturniculus passerinus</i> . | 26. <i>Sterna paradisæa</i> . |
| 13. <i>Ammodramus caudatus</i> . | 27. <i>Hydrochelidon fissipes</i> . |
| 14. <i>Ammodramus maritimus</i> . | |

2. *Species limited by the Alleghanian Fauna in their Southward Range in the Breeding Season.*

- | | |
|--|------------------------------------|
| 1. <i>Turdus fuscescens</i> . | 16. <i>Rallus virginianus</i> . |
| 2. <i>Dendroica pennsylvanica</i> . | 17. <i>Porzana carolina</i> . |
| 3. <i>Parula americana</i> . | 18. <i>Ægialitis melodus</i> . |
| 4. <i>Helminthophaga ruficapilla</i> . | 19. <i>Nettion carolinensis</i> . |
| 5. <i>Helminthophaga chrysoptera</i> . | 20. <i>Querquedula discors</i> . |
| 6. <i>Hirundo bicolor</i> . | 21. <i>Mareca americana</i> . |
| 7. <i>Lanius solitarius</i> . | 22. <i>Anas boschas</i> . |
| 8. <i>Carpodacus purpureus</i> . | 23. <i>Anas obscura</i> . |
| 9. <i>Melospiza palustris</i> . | 24. <i>Mergus merganser</i> . |
| 10. <i>Passerculus savanna</i> . | 25. <i>Lophodytes cucullatus</i> . |
| 11. <i>Dolichonyx oryzivorus</i> . | 26. <i>Bernicla canadensis</i> . |
| 12. <i>Contopus borealis</i> . | 27. <i>Colymbus torquatus</i> . |
| 13. <i>Empidonax minimus</i> . | 28. <i>Podilymbus podiceps</i> . |
| 14. <i>Empidonax flaviventris</i> . | 29. <i>Larus argentatus</i> . |
| 15. <i>Pediæcetes phasianellus</i> . | 30. <i>Sterna macrura</i> . |

* A few of the species mentioned in this list are more or less frequent stragglers into the Canadian Fauna, but none of them seem to occur there except as irregular and infrequent visitors.

V. CANADIAN FAUNA. The next fauna to the northward of the Alleghanian is the Canadian. The southern boundary of the Canadian is hence, of course, the northern limit of the Alleghanian, which boundary has been already defined. Its northern limit coincides very nearly with the isothermal line of 57° F. The region to the northward of the Alleghanian Fauna is unfortunately too little known to permit of a very satisfactory determination of the northern boundary of either the Canadian Fauna or of the faunæ to the northward of the Canadian. On the Atlantic coast the Canadian Fauna appears to embrace the greater part of Newfoundland, nearly or quite all of Nova Scotia* and New Brunswick, Northern New England, including the crests of the Green Mountain ranges southward to Connecticut, the greater part of the province of Quebec, including the Lower St. Lawrence valley as far up as the city of Quebec, the southern slope of the Height of Land in Northern Ontario, and the highlands on both sides of Lake Superior. To the southward it also embraces as outlying islands the Adirondacks of Northern New York, and the higher crests of the Appalachians southward to Georgia. To the northward it probably extends nearly to the summit of the Height of Land, and may embrace part of the lowlands bordering the southwestern shore of Hudson's Bay. Its northern boundary hence sweeps northwestward in the interior nearly or quite to Fort Resolution, on the southern shore of Great Bear Lake.

The Canadian Fauna, as above limited, may be characterized as follows. It is distinguished from the Alleghanian Fauna by the absence of the species mentioned above as limited in their northward range by that fauna, and by the presence in the breeding season of those mentioned in the first subjoined list; from the Hudsonian Fauna by the presence of those given in the second subjoined list, and by the absence of those given in the first list under the Hudsonian Fauna. •It is further distinguished by its forming the breeding haunts of a large proportion of the *Sylvicolidæ*, especially of the species of *Dendræca*, several of which are in summer mainly restricted to it.

* Nova Scotia, zoologically considered, presents somewhat anomalous characters. In summer a number of birds which are reported as either rare or accidental at Calais, Me., are represented as common summer residents in Nova Scotia, while other northern species breed there in numbers which do not usually breed at localities where the other species referred to are summer residents. The half-insular position of Nova Scotia is doubtless the explanation of the faunal peculiarities above mentioned.

1. *Species limited by the Canadian Fauna in their Southward Range in the Breeding Season.*

- | | |
|---------------------------------------|--|
| 1. <i>Turdus Pallasi.</i> | 32. <i>Tetrao canadensis.</i> |
| 2. <i>Turdus Swainsoni.</i> | 33. <i>Calidris arenaria.</i> |
| 3. <i>Regulus satrapa.</i> | 34. <i>Macrorhamphus griseus.</i> |
| 4. <i>Regulus calendula.</i> | 35. <i>Phalaropus Wilsoni.</i> |
| 5. <i>Dendræca castanea.</i> | 36. <i>Aythya vallisneria.</i> |
| 6. <i>Dendræca Blackburniæ.</i> | 37. <i>Aythya americana.</i> |
| 7. <i>Dendræca coronata.</i> | 38. <i>Fulix marila.</i> |
| 8. <i>Dendræca cærulescen</i> | 39. <i>Fulix collaris.</i> |
| 9. <i>Dendræca striata.</i> | 40. <i>Erismatura rubida.</i> |
| 10. <i>Dendræca palmarum.</i> | 41. <i>Bucephala clangula.</i> |
| 11. <i>Dendræca maculosa.</i> | 42. <i>Bucephala albeola.</i> |
| 12. <i>Euthlypis canadensis.</i> | 43. <i>Mergus serrator.</i> |
| 13. <i>Troglodytes hyemalis.</i> | 44. <i>Somateria mollissima.</i> |
| 14. <i>Parus hudsonicus.</i> | 45. <i>Cymochorea leucorrhea.</i> |
| 15. <i>Pinicola enucleator.</i> | 46. <i>Puffinus anglorum.</i> |
| 16. <i>Curvirostra americana.</i> | 47. <i>Larus marinus.</i> |
| 17. <i>Curvirostra leucoptera.</i> | 48. <i>Rissa tridactylus.</i> |
| 18. <i>Chrysomitris pinus.</i> | 49. <i>?Chræcocephalus philadelphia.</i> |
| 19. <i>Zonotrichia leucophrys.</i> | 50. <i>Pelecanus erythrorhynchus.</i> |
| 20. <i>Zonotrichia albicollis.</i> | 51. <i>Sula bassana.</i> |
| 21. <i>Junco hyemalis.</i> | 52. <i>Graculus carbo.</i> |
| 22. <i>Spizella monticola.</i> | 53. <i>Graculus dilophus.</i> |
| 23. <i>Passerella iliaca.</i> | 54. <i>Podiceps cristatus.</i> |
| 24. <i>Scolecophagus ferrugineus.</i> | 55. <i>Podiceps cornutus.</i> |
| 25. <i>Perisoreus canadensis.</i> | 56. <i>Podiceps Holbölli.</i> |
| 26. <i>Picoides hirsutus.</i> | 57. <i>Fratercula arctica.</i> |
| 27. <i>Picoides arcticus.</i> | 58. <i>Utamania torda.</i> |
| 28. <i>Falco columbarius.</i> | 59. <i>Uria grylle.</i> |
| 29. <i>Astur atricapillus.</i> | 60. <i>Lomvia ringvia.</i> |
| 30. <i>Surnia ulula.</i> | 61. <i>Lomvia svarbag.</i> |
| 31. <i>Nyctale Tengmalmi.</i> | |

2. *Species limited by the Canadian Fauna in their Northward Range.*

- | | |
|---------------------------------|-----------------------------------|
| 1. <i>?Turdus Pallasi.</i> | 7. <i>Euthlypis canadensis.</i> |
| 2. <i>Mimus carolinensis.</i> | 8. <i>Parus atricapillus.</i> |
| 3. <i>Dendræca virens.</i> | 9. <i>Chrysomitris tristis.</i> |
| 4. <i>Dendræca cærulescens.</i> | 10. <i>Chrysomitris pinus.</i> |
| 5. <i>Dendræca castanea.</i> | 11. <i>Curvirostra americana.</i> |
| 6. <i>Dendræca Blackburniæ.</i> | 12. <i>Poocætes gramineus.</i> |

- | | |
|------------------------------------|-------------------------------------|
| 13. <i>Melospiza melodia</i> . | 25. <i>Accipiter Cooperi</i> . |
| 14. <i>Melospiza palustris</i> . | 26. ? <i>Syrnium nebulosum</i> . |
| 15. <i>Dolichonyx oryzivorus</i> . | 27. <i>Butorides virescens</i> . |
| 16. <i>Tyrannus carolinensis</i> . | 28. <i>Porzana carolina</i> . |
| 17. <i>Myiarchus crinitus</i> . | 29. <i>Ægialitis vociferus</i> . |
| 18. <i>Contopus virens</i> . | 30. <i>Ægialitis melodus</i> . |
| 19. <i>Trochilus colubris</i> . | 31. <i>Philohela minor</i> . |
| 20. <i>Antrostomus vociferus</i> . | 32. <i>Actiturus Bartramius</i> . |
| 21. <i>Sayornis fuscus</i> . | 33. <i>Aix sponsa</i> . |
| 22. <i>Cyanura cristata</i> . | 34. <i>Chaulelasmus streperus</i> . |
| 23. <i>Buteo lineatus</i> . | 35. <i>Aythya vallisneria</i> . |
| 24. <i>Buteo pennsylvanicus</i> . | 36. <i>Hydrochelidon fissipes</i> . |

VI. HUDSONIAN FAUNA. The next ornithological fauna north of the Canadian may well be termed the Hudsonian Fauna. Its northern limit seems to nearly coincide with the isothermal line of 50° Fahrenheit, its southern limit being the isothermal of 57°, or the northern boundary of the Canadian Fauna. It will include at least the southern third of Labrador, the northern peninsula of Newfoundland, Anticosti Island, the more elevated parts of the Height of Land separating the lowlands bordering Hudson's Bay from the lowlands of the St. Lawrence and the Winnipeg district, and the basin of the Mackenzie's from Lake Athabasca to a point considerably north of Fort Simpson, extending in the Mackenzie's River valley some distance within the Arctic Circle, probably to the Arctic coast. Extending still westward, it embraces the valleys of Liard's and Peel's Rivers, and probably the valley and adjoining lowlands of the Youkon, including the greater part of that portion of the Territory of Alaska situated to the southward of the Arctic Circle. In other words, that portion of boreal America situated between the Canadian Fauna and the Barren Grounds. It is far from certain that a western or Alaskan Fauna will not have to be separated, embracing all the more temperate portions of Alaska. Although strictly western species occur here, they appear to be confined mainly to the western coast and the lower part of the Youkon valley. The faunal differences between the western shore of Alaska and the valley of Mackenzie's River become far greater when the marine species are taken into account, even if only the birds and mammalia are considered. The Pacific coasts of Alaska and Siberia have many species peculiar to the shores of those countries

and to the intervening islands, constituting a distinct fauna, which may well be called the *Aleutian Fauna*. The mingling of Asiatic and American species forms its distinctive feature. There is also a slight commingling of western species in the valley of the Mackenzie's River, as there is also in the valley of the Saskatchewan. The Hudsonian Fauna doubtless embraces outlying islands in the Canadian Fauna, as the upper part of the White Mountains, and the summits of some of the higher peaks of the Adirondacks. The southern point of Greenland embraces many species common to the Hudsonian Fauna, and though Greenland belongs almost wholly to the Arctic Realm, its extreme southern portion is doubtless Hudsonian.*

The Hudsonian Fauna being coextensive northward with the limit of forest-trees, it forms the northern limit of distribution of all the species of birds whose mode of life renders them dependent upon a forest vegetation. The distinction between the Hudsonian Fauna and the Arctic Realm, as well as between the Temperate Realm and the Arctic Realm, is hence a strongly marked one, nearly one hundred species, nearly all of them land birds, finding their northern limit of distribution near the polar limit of forests, or at least within the Hudsonian Fauna.

The Hudsonian Fauna may be distinguished from the Canadian by the absence of the species given in the preceding lists and by the presence of those enumerated in the first of the lists next subjoined, and from the Arctic Realm by the presence of those given in the second list below.

1. *Species limited by the Hudsonian Fauna in their Southward Range in the Breeding Season.*

- | | |
|--------------------------------------|----------------------------------|
| 1. <i>Anthus ludovicianus</i> . | 9. ? <i>Picoides arcticus</i> . |
| 2. <i>Saxicola œnanthe</i> . | 10. ? <i>Picoides hirsutus</i> . |
| 3. <i>Ampelis garrula</i> . | 11. <i>Falco candicans</i> . |
| 4. <i>Ægiothus linaria</i> . | 12. <i>Archibuteo lagopus</i> . |
| 5. <i>Plectrophanes lapponicus</i> . | 13. <i>Syrnium cinereum</i> . |
| 6. <i>Plectrophanes nivalis</i> . | 14. <i>Nyctea nivea</i> . |
| 7. <i>Plectrophanes pictus</i> . | 15. <i>Lagopus albus</i> . |
| 8. <i>Leucosticte tephrocotis</i> . | 16. <i>Lagopus rupestris</i> . |

* For remarks respecting the similarity of the Fauna of Northern Labrador and Southern Greenland, see Dr. A. S. Packard, Proc. Bost. Soc. Nat. Hist., Vol. X, p. 255, 1866.

- | | |
|-------------------------------------|---------------------------------------|
| 17. <i>Charadrius virginicus</i> . | 27. <i>Bernicla brenta</i> . |
| 18. <i>Ægialitis semipalmatus</i> . | 28. <i>Dafila acuta</i> . |
| 19. <i>Squatarola helvetica</i> . | 29. <i>Harelda glacialis</i> . |
| 20. <i>Streptilas interpres</i> . | 30. <i>Somateria spectabilis</i> . |
| 21. <i>Actodromas maculata</i> . | 31. <i>Pelionetta perspicillata</i> . |
| 22. <i>Actodromas Bonapartei</i> . | 32. <i>Procellaria glacialis</i> . |
| 23. <i>Actodromas minutilla</i> . | 33. <i>Sterna caspia</i> . |
| 24. <i>Ereunetes pusillus</i> . | 34. <i>Larus glaucus</i> . |
| 25. <i>Anser Gambeli</i> . | 35. <i>Colymbus septentrionalis</i> . |
| 26. <i>Anser hyperboreus</i> . | 36. <i>Stercorarius parasiticus</i> . |

2. *Species limited in their Northward Range by the Hudsonian Fauna.*

- | | |
|--|--|
| 1. <i>Turdus migratorius</i> . | 30. <i>Plectrophanes pictus</i> . |
| 2. <i>Turdus Swainsoni</i> . | 31. <i>Passerculus savanna</i> . |
| 3. <i>Regulus calendula</i> . | 32. <i>Zonotrichia leucophrys</i> . |
| 4. <i>Regulus satrapa</i> . | 33. <i>Zonotrichia albicollis</i> . |
| 5. <i>Helminthophaga ruficapilla</i> . | 34. <i>Junco hyemalis</i> . |
| 6. <i>Helminthophaga peregrina</i> . | 35. <i>Spizella monticola</i> . |
| 7. <i>Helminthophaga celata</i> . | 36. <i>Spizella socialis</i> . |
| 8. <i>Seiurus aurocapillus</i> . | 37. <i>Melospiza Lincolnii</i> . |
| 9. <i>Seiurus noveboracensis</i> . | 38. <i>Passerella iliaca</i> . |
| 10. <i>Dendroeca coronata</i> . | 39. <i>Molothrus pecoris</i> . |
| 11. <i>Dendroeca striata</i> . | 40. <i>Agelæus phœniceus</i> . |
| 12. <i>Dendroeca æstiva</i> . | 41. <i>Scolecophagus ferrugineus</i> . |
| 13. <i>Dendroeca maculosa</i> . | 42. <i>Quiscalus purpureus</i> . |
| 14. <i>Dendroeca palmarum</i> . | 43. <i>Corvus corax</i> . |
| 15. <i>Wilsonia pusilla</i> . | 44. <i>Corvus americanus</i> . |
| 16. <i>Setophaga ruticilla</i> . | 45. <i>Pica caudata</i> . |
| 17. <i>Hirundo horreorum</i> . | 46. <i>Perisoreus canadensis</i> . |
| 18. <i>Hirundo lunifrons</i> . | 47. <i>Contopus borealis</i> . |
| 19. <i>Hirundo bicolor</i> . | 48. <i>Empidonax minimus</i> . |
| 20. <i>Ampelis garrula</i> . | 49. <i>Empidonax Traillii</i> . |
| 21. <i>Collurio borealis</i> . | 50. <i>Picus villosus</i> . |
| 22. <i>Vireo olivaceus</i> . | 51. <i>Picus pubescens</i> . |
| 23. <i>Vireo gilvus</i> . | 52. <i>Picoides hirsutus</i> . |
| 24. <i>Parus hudsonicus</i> . | 53. <i>Picoides arcticus</i> . |
| 25. <i>Pinicola enucleator</i> . | 54. <i>Sphyrapicus varius</i> . |
| 26. <i>Curvirostra leucoptera</i> . | 55. <i>Hylotomus pileatus</i> . |
| 27. <i>Ægiothus linaria</i> . | 56. <i>Colaptes auratus</i> . |
| 28. <i>Plectrophanes nivalis</i> . | 57. <i>Chordeiles popetue</i> . |
| 29. <i>Plectrophanes lapponicus</i> . | 58. <i>Ceryle alcyon</i> . |

- | | |
|--------------------------------------|--|
| 59. <i>Falco candicans</i> . | 78. <i>Macrorhamphus griseus</i> . |
| 60. <i>Falco peregrinus</i> . | 79. <i>Ereunetes pusillus</i> . |
| 61. <i>Falco columbarius</i> . | 80. <i>Gambetta melanoleuca</i> . |
| 62. <i>Falco sparverius</i> . | 81. <i>Gambetta flavipes</i> . |
| 63. <i>Astur atricapillus</i> . | 82. <i>Tringoides macularius</i> . |
| 64. <i>Archibuteo lagopus</i> . | 83. <i>Limosa hudsonica</i> . |
| 65. <i>Buteo borealis</i> . | 84. <i>Porzana carolina</i> . |
| 66. <i>Accipiter fuscus</i> . | 85. <i>Fulica americana</i> . |
| 67. <i>Circus cyaneus</i> . | 86. ? <i>Dafila acuta</i> . |
| 68. <i>Bubo virginianus</i> . | 87. <i>Nettion carolinensis</i> . |
| 69. <i>Otus vulgaris</i> . | 88. <i>Querquedula discors</i> . |
| 70. <i>Otus brachyotus</i> . | 89. <i>Spatula clypeata</i> . |
| 71. <i>Ectopistes migratori</i> . | 90. <i>Mareca americana</i> . |
| 72. <i>Tetrao canadensis</i> . | 91. <i>Fulix marila</i> . |
| 73. <i>Pediœcetes phasianellus</i> . | 92. <i>Fulix collaris</i> . |
| 74. <i>Bonasa umbellus</i> . | 93. <i>Erismatura rubida</i> . |
| 75. <i>Lagopus leucurus</i> . | 94. <i>Lophodytes cucullatus</i> . |
| 76. ? <i>Grus americanus</i> . | 95. <i>Graculus dilophus</i> . |
| 77. <i>Gallinago Wilsoni</i> . | 96. <i>Pelecanus erythrorhynchus</i> . |

VII. AMERICAN ARCTIC FAUNA. The Arctic Realm may be considered as occupying that portion of the northern hemisphere north of the isothermal of 50° F. Though presenting a great uniformity of character throughout its extent, it seems to be divisible into four faunæ, — an American Arctic Fauna, an Americo-Asiatic Fauna, an Europeo-Asiatic Arctic Fauna, and an Europeo-American Arctic Fauna, the second and fourth being essentially marine. While a few species of mammals seem to be almost wholly restricted within the Arctic Realm, it contains but few resident birds, and no species of birds seem to be wholly confined to it, even in the breeding season. The following species are reported to range to the Arctic coast of North America, the most of which have been observed at Melville Island and in Greenland. The greater part are *Natatores* and *Grallæ*, the only commonly so-called land birds being two or three species of grouse, a sparrow or two, and a few hawks and owls.

Species found in the American Arctic Fauna in the Breeding Season

- | | |
|-------------------------------|-------------------------------------|
| 1. <i>Cotyle riparia</i> . | 4. <i>Haliaëtus leucocephalus</i> . |
| 2. <i>Corvus corax</i> . | 5. <i>Pandion haliaëtus</i> . |
| 3. <i>Aquila chrysaëtos</i> . | 6. <i>Nyctea nivea</i> . |

- | | |
|---------------------------------------|---|
| 7. ? <i>Surnia ulula</i> . | 33. <i>Somateria spectabilis</i> . |
| 8. <i>Tetrao canadensis</i> . | 34. <i>Somateria mollissima</i> . |
| 9. <i>Lagopus albus</i> . | 35. <i>Buphagus skau Coues</i> . |
| 10. <i>Lagopus rupestris</i> . | 36. <i>Stercorarius pomarinus</i> . |
| 11. <i>Grus canadensis</i> . | 37. <i>Stercorarius parasiticus</i> . |
| 12. <i>Botaurus lentiginosus</i> . | 38. <i>Stercorarius Buffoni Coues</i> . |
| 13. <i>Charadrius virginicus</i> . | 39. <i>Fulmarus glacialis</i> . |
| 14. <i>Ægialitis semipalmatus</i> . | 40. <i>Larus argentatus</i> . |
| 15. <i>Streptilas interpres</i> . | 41. <i>Larus glaucus</i> . |
| 16. <i>Phalaropus fulicarius</i> . | 42. <i>Larus leucopterus</i> . |
| 17. <i>Calidris arenaria</i> . | 43. <i>Rissa tridactyla</i> . |
| 18. <i>Tringa canutus</i> . | 44. <i>Pagophila eburnea</i> . |
| 19. <i>Pelidna "americana"</i> . | 45. <i>Xema Sabini</i> . |
| 20. <i>Arquatella maritima</i> . | 46. <i>Sterna arctica</i> . |
| 21. <i>Cygnus "americanus"</i> . | 47. ? <i>Sula bassana</i> . |
| 22. <i>Anser hyperboreus</i> . | 48. <i>Colymbus septentrionalis</i> . |
| 23. <i>Anser Gambeli</i> . | 49. <i>Colymbus arcticus</i> . |
| 24. <i>Bernicla canadensis</i> . | 50. <i>Colymbus torquatus</i> . |
| 25. <i>Anas boschas</i> . | 51. <i>Utamania torda</i> . |
| 26. <i>Bucephala albeola</i> . | 52. <i>Fratercula glacialis</i> . |
| 27. <i>Bucephala clangula</i> . | 53. <i>Lunda cirrhata</i> . |
| 28. <i>Histrionicus torquatus</i> . | 54. <i>Mergulus alle</i> . |
| 29. <i>Harelda glacialis</i> . | 55. <i>Uria grylle</i> . |
| 30. <i>Melanetta velvetina</i> . | 56. <i>Lomvia troile</i> . |
| 31. <i>Pelionetta perspicillata</i> . | 57. <i>Lomvia ringvia</i> . |
| 32. <i>Ædemia "americana"</i> . | 58. <i>Lomvia svarbag</i> . |

The Faunæ of the Eastern Province considered in Reference to the Distribution of the Mammals and Reptiles.

The faunæ of the Eastern Province of the North American Region above characterized from the distribution of the birds seem to be equally well marked as natural zoölogical districts by the distribution of the mammals and reptiles. About the same proportionate number of mammals are limited similarly with the birds in regard to their northward and southward distribution. The correspondence in the geographical limitation of the species of the two groups will be briefly shown by the following remarks respecting the range of the mammals.

The Arctic Realm is well known to be characterized by a few species nearly or quite restricted to it, as the polar bear (*Ursus maritimus*),

the arctic fox (*Vulpes lagopus*), the musk ox (*Ovibos moschatus*), the lemmings (*Myodes*), the small northern race of the caribou or reindeer, the Eskimos, etc.

The Hudsonian Fauna forms the southern limit of the polar hare (*Lepus glacialis*) and the northern limit (at least in winter) of the *Lynx canadensis*, *Mustela* "americana," *Mustela Pennantii*, *Putorius vulgaris*, *Putorius ermineus*, *Ursus* "americanus," *Ursus* "horribilis," *Sciurus hudsonius*, *Arctomys monax*, *Vespertilio subulatus*, and others.

The Canadian Fauna forms, at present,* the southern limit of *Mustela Pennantii*, *Mustela* "americana," *Gulo luscus*, *Arvicola xanthognathus*, *Erethizon dorsatus*, *Alces malchis*, *Rangifer tarandus*, etc., and the northern limit of *Felis concolor*, *Lynx rufus*, *Mephitis mephitis*, *Procyon lotor*, *Bos americanus*, *Condylura cristata*, *Blarina brevicauda*, and others.

The Alleghanian Fauna forms the southern limit of *Lynx canadensis*, *Sciurus hudsonius*, *Arvicola Gapperi*, *Jaculus hudsonius*, *Lepus americanus*, *Cervus canadensis*, *Sorex platyrhinus*, *Condylura cristata*, and doubtless of several other species; and the northern limit of *Sciurus carolinensis*, *Lepus sylvaticus*, *Arvicola pinetorum*, *Cervus virginianus*, *Scalops aquaticus*, etc.

The Carolinian Fauna forms the southern limit of *Mustela vulgaris*, *Tamias striatus*, *Arctomys monax*, and *Fiber zibethicus*; and the northern limit of *Vulpes virginianus*, *Nycticejus crepuscularis*, *Didelphys virginiana*, etc.

The Louisianian Fauna seems to form the southern limit of *Putorius vison*, *Blarina brevicauda*, *Scalops aquaticus*, and doubtless thus limits several other species, though not a small proportion of those which occur in this fauna range also into South Florida, or into the Floridian Fauna. The Louisianian Fauna limits the northward range of *Neotoma floridana*, *Reithrodon humilis*, *Sigmodon hispidus*, *Hesperomys palustris*, *Geomys pineti*, and *Lepus palustris*.

In respect to reptiles, a similar proportion of species are limited in either their northward or southward range by each fauna. Several species of batrachians range into the Hudsonian Fauna, but apparently this fauna must be the northern limit of their distribution. The Cana-

* It is probable that some of the fur-bearing species, as well as *Hystrix dorsatus*, the moose and the caribou, once ranged southward throughout the Alleghanian Fauna, and have been exterminated there by man.

dian Fauna forms the northern limit of the reptiles proper, where this class is represented by two orders only, the *Testudinata* and the *Ophidia*. The *Testudinata* are there represented by only three species (*Chelydra serpentina*, *Glyptemys insculpta*, *Chrysemys picta*), and the *Ophidia* by five (*Bascanion constrictor*, *Tropidonotus sirtalis*, *Diadophis punctatus*, *Storeria occipito-maculata*). In the Alleghanian Fauna the number of species in each of these groups is more than doubled. The Carolinian Fauna forms the northern limit of the *Sauria*, of which two species (*Plestiodon fasciatus*, *Tropidolepis undulatus*) here first make their appearance, and the number of species of the other groups is still further increased, several additional generic types being added. In the Louisianian Fauna the number of species of *Sauria* becomes considerably greater, and while few of the northern species of either the true reptiles or the batrachians have disappeared, other more southern forms have been added in almost every family.

These several faunæ, it may be added, seem in general to coincide in their number and limits with the floræ of the same region. These several floræ, as thus circumscribed, form successively the northern limit of the successful cultivation of some more or less important cultivated plant. But a detailed consideration of the distribution of the vegetation of the region under consideration, in respect to the number and circumscription of the floræ, and their correspondence with the faunæ, cannot of course well be here introduced.

4. THE ORNITHOLOGICAL DISTRICTS OF THE NORTH AMERICAN TEMPERATE REGION.

The subdivision by Professor Dana of the tropical and temperate climatic zones of the oceanic areas into several zoölogical zones has been already alluded to as being equally applicable to the land areas. To these life zones I propose to apply the term "districts." Dana's divisions of the north temperate climatic zone correspond respectively in latitudinal extent with the several ornithological faunæ of the Eastern Province, as defined in the preceding pages. Unlike the faunæ, however, the districts extend in an east and west direction across the North American Region, each district embracing not only one of the faunæ of the Eastern Province, but also its representative fauna (or faunæ) in the Western Province. The Hudsonian Fauna corresponds in latitudinal extent with Dana's subfrigid division of the north tem-

perate zone, and the term *Subfrigid District* may be very properly applied to that district of which this fauna forms the eastern portion. The zone corresponding with the Canadian Fauna may in like manner be termed the *Cold-temperate District*; that corresponding with the Alleghanian Fauna the *Subtemperate District*; that corresponding with the Carolinian Fauna the *Temperate District*; and that corresponding with the Louisianian Fauna the *Warm-temperate District*; the Floridian Fauna in like manner corresponding with the *Subtorrid District*, or with Dana's subtorrid zone. Each of these districts is distinguished, in contradistinction from the faunæ, by species which range across the continent, while the districts are distinguished from each other by the same kind of difference as has been shown above to characterize the several faunæ among themselves.

5. ON THE GEOGRAPHICAL RANGE OF THE SPECIES.

The preceding tables, while serving to characterize the ornithological faunæ of Eastern North America, indicate only very obscurely the range of the species. The following tables have hence been prepared in order to show more clearly the breeding range, and also the winter quarters, of those species whose distribution in the breeding season is tolerably known. For this purpose the birds occurring in the Eastern Province of the North American region have been grouped, according to their geographical distribution, into the following classes, beginning with those which have the widest breeding range: I. *Cosmopolitan Species*. II. *Circumpolar Species*. III. *Species which range across the whole breadth of the North American Temperate Region*. IV. *Species limited in longitude to the Eastern Province of this region*. The birds of the Eastern Province are further subdivided according to the range of the species in the breeding season in latitude.*

* In a preliminary notice like the present it has been found impracticable to give the authorities in detail on which the generalizations given in the following synopsis have been based. The list of papers given in the Appendix serve in a general way to indicate the principal sources from which information has been derived. It is believed, however, that the limits assigned each species will be found in the main correct, though in many cases the accessible data have been quite too few to be satisfactory. The generalizations are given, of course, as a representation of our present knowledge of the subject rather than as final. The polar and equatorial limits of the migratory range of the species varies, as is well known, more or less in different years, according to the season. It is also somewhat different on the coast from what it is in

I. COSMOPOLITAN SPECIES. A large proportion of ornithologists have of late been unwilling to admit that any bird has what is usually termed a "cosmopolitan" range, while others recognize only about twenty such species, taking into account, of course, their total range. These embrace two or three species each of hawks and owls, the rest being either *Grallæ* or *Natatores*. Very few of them, however, breed within both the tropic and the polar zones; many of those which visit the shores of all lands in their migratory journeys being restricted in the breeding season to comparatively limited areas. *Pandion haliaëtus* and *Otus brachyotus* are the only examples of commonly so-called cosmopolitan species which appear to breed from the Arctic Circle southward through the tropics to the southern extremity of the southern continents. *Falco peregrinus* may form a third, but its peculiar breeding habits give it a very irregular dispersion at that season. *Strix flammea* appears to be also everywhere resident, except in the arctic and cold-temperate zones. *Cotyle riparia* and *Hirundo rufa* (including under the latter name the several slightly differing geographical races of this group, which have of late been regarded as species), seem also to be nearly cosmopolite. The list of species which are permanently cosmopolitan will hence not exceed half a dozen, and are those above enumerated.

II. CIRCUMPOLAR SPECIES. Regarding as circumpolar species only those numerous represented in both the eastern and western hemispheres, nearly one hundred species* can be included in the list of

the interior, as has been previously explained; so that an indication of only the average boreal and austral limits of the species at this season has been aimed at, and only so far as their winter range is circumscribed within the region under special consideration. The blanks in the third column of the tables hence indicate that the species winter entirely within the American Tropical Realm; those in the fourth, that the austral limit is within that realm. The few occurring in the second column of the tables indicate that the species in question also ranges southward in the breeding season into the Tropical Realm. A [?] in place of a blank indicates that the southward range of the species is supposed to be limited to the Eastern Province, but as being too vaguely known to warrant a specification of its limit in the direction indicated by the column in which the query stands.

* Dr. Richardson, in 1831 (in the "Fauna Boreali-Americana," p. xxxix), gave thirty-two species of land birds and "upwards of sixty-two species of water birds" (ninety-four in all) as "common to the Old World and the Fur Countries." A few truly circumpolar species were not included in this list, and others were included which were merely accidental visitors from one continent to the other. Since the date of that list the identity of the greater part of the species therein mentioned has been questioned by

circumpolar species. A small number of others that are properly either exclusively American or Europeo-Asiatic species occur more or less frequently as accidental visitors to the continents not embraced within their usual habitats.

one or another writer, and their representatives on the two hemispheres separated under different names. But a considerable proportion of those mentioned in the next subjoined table are still regarded as truly circumpolar by a number of leading European ornithologists. Dr. Von Middendorff ("Uebersicht der Natur Nord- und Ost-Sibiriens, Theil 2, Erste Lieferung," etc.; see Newton's Ibis, April, 1870, p. 275), in 1867, gave lists of eighty-seven circumpolar species, a part of which (called "Hyperboreal Birds") are distinctive of what has been termed above the Arctic Realm, whilst many of the others range quite far southward even in summer. These lists, however, do not embrace a number of circumpolar species whose boreal limit does not extend to the districts named. A dozen or more Europeo-Asiatic species, in addition to those given below, have representatives in America so closely resembling them in habits and in geographical distribution, as well as structurally, that they have often been confounded, specimens frequently occurring on the one continent that are undistinguishable from those from the other continent.

In 1846 Professor Edm. de Selys-Longchamps, in his excellent paper entitled "Sur les Oiseaux américains admis dans la Faune européenne" (Mém. de la Soc. R. de Liège, Vol. IV, pp. 35-50, 1849), included thirteen species in his list of "Oiseaux terrestres communs à l'Europe et à l'Amérique," and mentions nine other terrestrial American species which he regards as "ne semblent être en réalité que des modifications climatiques de nos oiseaux européens." All but two of these, and also one or two in addition to them, have been regarded in the present paper as specifically identical. In his list of "Oiseaux aquatiques communs à l'Europe et à l'Amérique" he includes fifty-five species, and mentions thirteen others, "décrits comme espèces distinctes, ne semblent être que des races locales," three or four of which I have regarded as specifically identical. The whole number mentioned by Selys-Longchamps as common to Europe and boreal America is seventy-six, plus twenty-four "autres qui semblent n'être que des races légèrement modifiées par le climat." (See l. c., p. 48.) In the same paper he gives a list of twenty-eight American species as of accidental occurrence in Europe, eight of which are land birds, eight échassiers or waders, and twelve palmipèdes or swimmers, and also a list of twenty American species which he considers to have been improperly included among the birds of Europe, among which are *Haliaeetus leucocephalus*, *Strix nebulosa* (= *Syrnium nebulosum*), *Loxia* (= *Curvirostra*) *leucoptera*, *Struthus* (= *Junco*) *hyemalis*, and *Parus* (= *Lophophanes*) *bicolor*.

II. List of Circumpolar Species, with Indications of their Boreal and Austral Limits.

Species.	Boreal Limit in the Breeding Season.	Austral Limit in the Breeding Season.	Boreal Limit in Winter.	Austral Limit in Winter.
<i>Certhia familiaris</i>	Canad. Fauna	Allegh. Fauna	Allegh. Fauna	Carolin. Fauna
<i>Saxicola œnanthe</i>	Hudson. Fauna	Hudson. Fauna	—	—
<i>Ampelis garrula</i>	Hudson. Fauna	Hudson. Fauna	Hudson. Fauna	Canad. Fauna
<i>Cotyle riparia</i>	Arctic Coast	—	Louis. Fauna	—
<i>Plectrophanes nivalis</i>	Arctic Coast	Hudson. Fauna	Hudson. Fauna	Allegh. Fauna
<i>Plectrophanes lapponica</i>	Hudson. Fauna	Hudson. Fauna	—	Allegh. Fauna
<i>Ægithus linaria</i>	Arctic Coast	Hudson. Fauna	Hudson. Fauna	Allegh. Fauna
<i>Pinicola enucleator</i>	Hudson. Fauna	Canad. Fauna	Hudson. Fauna	Allegh. Fauna
<i>Curvirostra leucoptera</i>	Hudson. Fauna	Canad. Fauna	Hudson. Fauna?	Canad. Fauna
<i>Eremophila alpestris</i>	Hudson. Fauna	Carolin. Fauna	Canad. Fauna?	Louis. Fauna
<i>Corvus corax</i>	Arctic Coast	Allegh. Fauna	Arctic Realm	?
<i>Pica caudata</i>	Hudson. Fauna	Allegh. Fauna	Canad. Fauna	?
<i>Aquila chrysaetos</i>	Hudson. Fauna	Canad. Fauna	Hudson. Fauna?	Carolin. Fauna?
<i>Archibuteo lagopus</i>	Arctic Coast	Hudson. Fauna	Hudson. Fauna	Allegh. Fauna
<i>Pandion haliaetus</i>	Arctic Coast	—	Carolin. Fauna	—
<i>Falco candicans</i>	Arctic Coast?	Hudson. Fauna	Canad. Fauna	Canad. Fauna
<i>Falco peregrinus</i>	Hudson. Fauna	—	—	—
<i>Circus cyaneus</i>	Arctic Coast?	Louis. Fauna?	Canad. Fauna	—
<i>Strix flammea</i>	Carolin. Fauna	—	Carolin. Fauna	—
<i>Nyctale Tengmalmi</i>	Hudson. Fauna	Canad. Fauna	Hudson. Fauna	Canad. Fauna
<i>Syrnium cinereum</i>	Hudson. Fauna	Hudson. Fauna	Hudson. Fauna	Canad. Fauna
<i>Otus vulgaris</i>	Hudson. Fauna	—	Canad. Fauna	—
<i>Otus brachyotus</i>	Arctic Coast	—	Hudson. Fauna?	—
<i>Surnia ulula</i>	Arctic Coast	Canad. Fauna	Arctic Realm	Canad. Fauna
<i>Nyctea nivea</i>	Arctic Coast	Hudson. Fauna?	Arctic Realm	Allegh. Fauna
<i>Lagopus albus</i>	Arctic Coast	Hudson. Fauna	Arctic Realm	Hudson. Fauna
<i>Lagopus rupestris</i>	Arctic Coast	Hudson. Fauna	Arctic Realm	Hudson. Fauna
<i>Nycticorax griseus</i>	Hudson. Fauna	Carolin. Fauna?	Carolin. Fauna	—
<i>Streptopelia interpres</i>	Arctic Coast	Hudson. Fauna	—	—
<i>Charadrius pluvialis</i>	Arctic Coast	Arctic Realm	Tropical Amer.	—
<i>Squatarola helvetica</i>	Arctic Coast	Hudson. Fauna	Carolin. Fauna	—
<i>Hæmatopus palliatus</i>	Arctic Coast	—	—	—
<i>Arquatella maritima</i>	Arctic Coast	Arctic Realm	Canad. Fauna	?
<i>Calidris arenaria</i>	Arctic Coast	Canad. Fauna	Allegh. Fauna	—
<i>Tringa canutus</i>	Arctic Coast	Arctic Realm	Carolin. Fauna	—
<i>Ancylocheilus subarquatus</i>	Arctic Coast	Arctic Realm?	Carolin. Fauna?	—
<i>Pelidna alpina</i>	Arctic Coast	Hudson. Fauna?	Allegh. Fauna	—
<i>Actodromas maculata</i>	Arctic Coast	Arctic Realm	Carolin. Fauna	—
<i>Actodromas Bonapartei</i>	Arctic Coast	Arctic Realm	Carolin. Fauna	—
<i>Tringides rufescens</i>	Arctic Coast?	Hudson. Fauna	Louis. Fauna	—
<i>Lobipes hyperboreus</i>	Arctic Coast	Canad. Fauna	Carolin. Fauna	?
<i>Phalaropus fulicarius</i>	Arctic Coast	Hudson. Fauna?	Carolin. Fauna?	?
<i>Anser hyperboreus</i>	Arctic Coast	Hudson. Fauna	Canad. Fauna?	—
<i>Anser albifrons</i>	Arctic Coast	Hudson. Fauna	Allegh. Fauna	—
<i>Bernicla brenta</i>	Arctic Coast	Hudson. Fauna	Allegh. Fauna	Louis. Fauna
<i>Anas boschas</i>	Arctic Coast	Allegh. Fauna	Canad. Fauna	—
<i>Dafila acuta</i>	Arctic Coast	Hudson. Fauna	—	—
<i>Spatula clypeata</i>	Arctic Coast	Hudson. Fauna	Allegh. Fauna	—
<i>Chaulelasmus streperus</i>	Hudson. Fauna	Hudson. Fauna	?	Louis. Fauna
<i>Somateria mollissima</i>	Arctic Coast	Canad. Fauna	?	Carolin. Fauna
<i>Somateria spectabilis</i>	Arctic Coast	Hudson. Fauna	?	Allegh. Fauna
<i>Bucephala clangula</i>	Arctic Coast	Canad. Fauna	Canad. Fauna	Louis. Fauna
<i>Histrionicus torquatus</i>	Arctic Coast	Canad. Fauna	Hudson. Fauna?	Allegh. Fauna
<i>Harelda glacialis</i>	Arctic Coast	Hudson. Fauna	?	Louis. Fauna
<i>Fulix marila</i>	Hudson. Fauna	Canad. Fauna	—	—
<i>Pelionettaper spicillata</i>	Arctic Coast	Hudson. Fauna	Canad. Fauna?	Louis. Fauna
<i>Melanetta fusca</i>	Arctic Coast	Hudson. Fauna	Canad. Fauna?	Louis. Fauna
<i>Mergus merganser</i>	Hudson. Fauna	Canad. Fauna	Canad. Fauna?	Louis. Fauna
<i>Mergus serrator</i>	Hudson. Fauna	Canad. Fauna	Canad. Fauna	Louis. Fauna
<i>Graculus carbo</i>	Arctic Coast	Hudson. Fauna	?	Allegh. Fauna
<i>Sula bassana</i>	Arctic Coast	Canad. Fauna	?	Louis. Fauna
<i>Gelocheidon anglica</i>	Carolin. Fauna	—	—	—
<i>Thalasseus caspius</i>	Arctic Coast	Hudson. Fauna	?	Carolin. Fauna
<i>Thalasseus cantianus</i>	Louis. Fauna	—	—	—
<i>Sterna hirundo</i>	Arctic Coast?	Carolin. Fauna	Florid. Fauna	—
<i>Sterna macrura</i>	Arctic Coast	Allegh. Fauna	Florid. Fauna	—
<i>Sterna paradisea</i>	Allegh. Fauna	—	—	—
<i>Hydrochelidon fissipes</i>	Carolin. Fauna	—	Florid. Fauna	—

List of Circumpolar Species. (Continued.)

Species.	Boreal Limit in the Breeding Season.	Austral Limit in the Breeding Season.	Boreal Limit in Winter.	Austral Limit in Winter.
<i>Pagophila eburnea</i>	Arctic Coast	Arctic Realm	?	Huds. Fauna
<i>Xema Sabini</i>	Arctic Coast	Hudson. Fauna	?	Canad. Fauna
<i>Rissa tridaactyla</i>	Arctic Coast	Canad. Fauna	?	Allegh. Fauna
<i>Rhodostethia rosea</i>	Arctic Coast	Arctic Realm	?	?
<i>Larus glaucus</i>	Arctic Coast	Hudson. Fauna	?	Canad. Fauna
<i>Larus marinus</i>	Arctic Coast	Canad. Fauna	?	Louis. Fauna
<i>Larus leucopterus</i>	Arctic Coast	Hudson. Fauna	?	Allegh. Fauna
<i>Larus argentatus</i>	Arctic Coast	Canad. Fauna	?	Trop. Realm
<i>Buphagus skau</i>	Arctic Coast	Hudson. Fauna	?	Canad. Fauna
<i>Stercorarius pomarinus</i>	Arctic Coast	Hudson. Fauna?	?	Canad. Fauna
<i>Stercorarius parasiticus</i>	Arctic Coast	Hudson. Fauna?	?	Allegh. Fauna
<i>Stercorarius "Buffoni"</i>	Arctic Coast	Hudson. Fauna?	?	Allegh. Fauna
<i>Thalassidroma Leachii</i>	?	Canad. Fauna	?	Allegh. Fauna?
<i>Podiceps griseigena</i>	Hudson. Fauna	Allegh. Fauna	?	Louis. Fauna
<i>Podiceps cornutus</i>	Hudson. Fauna	Allegh. Fauna	?	Louis. Fauna
<i>Podiceps cristatus</i>	Hudson. Fauna	Allegh. Fauna	?	Louis. Fauna
<i>Podiceps auritus</i>	Hudson. Fauna	Allegh. Fauna?	?	Louis. Fauna
<i>Colymbus torquatus</i>	Arctic Coast	Allegh. Fauna	?	Florid. Fauna
<i>Colymbus arcticus</i>	Arctic Coast	Hudson. Fauna	?	Canad. Fauna
<i>Colymbus septentrionalis</i>	Arctic Coast	Canad. Fauna	?	Canad. Fauna
<i>Fratercula arctica</i>	Arctic Coast	Canad. Fauna	?	Allegh. Fauna
<i>Uria grylle</i>	Arctic Coast	Canad. Fauna	?	Carol. Fauna
<i>Lomvia svarbag</i>	Arctic Coast	Canad. Fauna	?	Carol. Fauna?
<i>Lomvia troile</i>	Arctic Coast	Canad. Fauna	?	Allegh. Fauna
<i>Mergulus alle</i>	Arctic Coast	—	?	Allegh. Fauna

Summary of the Preceding Table.—The whole number of species in the preceding list is 93. Its most striking feature is the great predominance of the water birds, less than one third of the whole being land birds. Of the 27 land birds, 7 are owls, 6 are hawks, and 5 belong to the family *Fringillidæ*; 9 species embracing all the representatives of other families. The water birds include 1 heron, 14 *Grallæ*, 17 *Anatidæ*, 19 *Laridæ*, 5 *Alcidæ*, 3 species of *Colymbus*, and 4 of *Podiceps*.

In summer 65 species are inhabitants of the Arctic coast and adjacent seas; 22 have their boreal limit near the northern border of the Hudsonian Fauna; 2 are similarly limited by the Canadian Fauna, 5 by the Alleghanian, 3 by the Carolinian, and 1 by the Louisianian; 3 are essentially tropical aquatic species.

Seven seem to be altogether restricted in the breeding season to the Arctic Realm; 36 find their austral limit during the same season near the southern border of the Hudsonian Fauna; 23 are similarly limited by the Canadian, 9 by the Alleghanian, and 3 by the Carolinian, while 10 extend nearly to or within the Tropical Realms, 4 being also inhabitants of the greater part of the southern hemisphere.

The winter quarters of the land birds of this list are the cold-temperate and middle-temperate districts of the northern hemisphere. Most of the water birds visit the warm-temperate parts of the same hemisphere; a considerable number also visit the tropics, and a few wander, at this season, over the greater part of the warmer regions of the globe.

III. SPECIES MAINLY RESTRICTED IN THE BREEDING SEASON TO THE NORTH AMERICAN TEMPERATE REGION.

1. *List of Species which breed throughout the greater Part of Temperate North America, with Indications of their Boreal and Austral Limits Distribution in the Eastern Province.*

Species	Boreal Limit in the Breeding Season.	Austral Limit in the Breeding Season.	Boreal Limit in Winter	Austral Limit in Winter.
<i>Turdus migratorius</i>	Hudson. Fauna	Carol. Fauna	Carol. Fauna	_____ *
<i>Turdus fuscescens</i>	Canad. Fauna	Carol. Fauna	_____ *	_____
<i>Geothlypis trichas</i>	Hudson. Fauna	Florid. Fauna	Florid. Fauna	_____
<i>Hirundo horreorum</i>	Hudson. Fauna	Louis. Fauna	_____	_____
<i>Petrochelidon lunifrons</i>	Hudson. Fauna	Carol. Fauna?	_____	_____
<i>Tachycineta bicolor</i>	Hudson. Fauna	Carol. Fauna	Louis. Fauna	_____
<i>Cotyle riparia</i>	Arctic Coast	Carol. Fauna	Florid. Fauna	_____
<i>Progne subis</i>	Canad. Fauna	?	Louis. Fauna	_____
<i>Ampelis cedrorum</i>	Canad. Fauna	Florid. Fauna	Allegh. Fauna	_____
<i>Collurio ludovicianus</i>	Allegh. Fauna	Florid. Fauna	Louis. Fauna	_____
<i>Vireosylva gilva</i>	Canad. Fauna	Carol. Fauna	_____	_____
<i>Lanius solitarius</i>	Hudson. Fauna	Allegh. Fauna	Florid. Fauna	_____
<i>Cistothorus palustris</i>	Hudson. Fauna	Louis. Fauna	Louis. Fauna	_____
<i>Troglodytes aedon</i>	Allegh. Fauna	Florid. Fauna	Louis. Fauna	_____
? <i>Sitta carolinensis</i>	Canad. Fauna	Louis. Fauna	Canad. Fauna	Florid. Fauna
<i>Sitta canadensis</i>	Canad. Fauna	Carol. Fauna	Canad. Fauna	Louis. Fauna?
<i>Parus atricapillus</i>	Hudson. Fauna	Florid. Fauna	Canad. Fauna	Florid. Fauna
<i>Carpodacus purpureus</i>	Canad. Fauna	Allegh. Fauna	Allegh. Fauna	Louis. Fauna
<i>Chrysomitris tristis</i>	Canad. Fauna	Louis. Fauna	Allegh. Fauna	_____
<i>Passerculus savanna</i>	Hudson. Fauna	Allegh. Fauna	Carol. Fauna	_____
<i>Pooecetes gramineus</i>	Canad. Fauna	Carol. Fauna	Carol. Fauna	_____
<i>Spizella socialis</i>	Hudson. Fauna	Louis. Fauna	Carol. Fauna	_____
? <i>Melospiza melodia</i>	Hudson. Fauna	Louis. Fauna	Carol. Fauna	Florid. Fauna
<i>Melospiza lincolni</i>	Hudson. Fauna	Allegh. Fauna	Carol. Fauna	_____
<i>Molothrus pecoris</i>	Hudson. Fauna	Louis. Fauna	Carol. Fauna	_____
<i>Agelaius phoeniceus</i>	Hudson. Fauna	Florid. Fauna	Carol. Fauna	_____
<i>Sturnella ludoviciana</i>	Allegh. Fauna	Florid. Fauna	Carol. Fauna	_____
<i>Corvus americanus</i>	Hudson. Fauna	Florid. Fauna?	Canad. Fauna	_____
<i>Tyrannus carolinensis</i>	Canad. Fauna	Louis. Fauna	_____	_____
? <i>Myiarchus crinitus</i>	Canad. Fauna	Louis. Fauna	_____	_____
? <i>Sayornis fuscus</i>	Hudson. Fauna	Louis. Fauna	Louis. Fauna	_____
<i>Contopus borealis</i>	Hudson. Fauna	Allegh. Fauna	_____	_____
? <i>Contopus virens</i>	Canad. Fauna	Louis. Fauna	_____	_____
? <i>Empidonax minimus</i>	Hudson. Fauna	Allegh. Fauna	_____	_____
? <i>Empidonax acadicus</i>	Canad. Fauna	Carol. Fauna	_____	_____
<i>Empidonax flaviventris</i>	Canad. Fauna	Carol. Fauna	_____	_____
<i>Picus villosus</i>	Hudson. Fauna	Louis. Fauna	Canad. Fauna?	Florid. Fauna
<i>Picus pubescens</i>	Hudson. Fauna	Louis. Fauna	Canad. Fauna?	Florid. Fauna
<i>Hylotomus pileatus</i>	Hudson. Fauna	Florid. Fauna	Canad. Fauna?	Florid. Fauna
<i>Chordeiles popetue</i>	Hudson. Fauna	_____ *	_____	_____
<i>Ceryle alcyon</i>	Arctic Coast	Florid. Fauna	Carol. Fauna	_____
<i>Accipiter Cooperi</i>	Canad. Fauna	_____	Carol. Fauna	_____
<i>Accipiter fuscus</i>	Canad. Fauna	_____	Carol. Fauna	_____
<i>Buteo borealis</i>	Hudson. Fauna	_____	Canad. Fauna	_____
<i>Buteo lineatus</i>	Canad. Fauna	Florid. Fauna	Allegh. Fauna	Florid. Fauna
<i>Buteo pennsylvanicus</i>	Canad. Fauna	_____	Allegh. Fauna	_____
<i>Haliaetus leucocephalus</i>	Arctic Coast	Florid. Fauna	Hudson. Fauna?	Florid. Fauna
<i>Scops asio</i>	Canad. Fauna	_____	Canad. Fauna	_____
<i>Zenaidura carolinensis</i>	Allegh. Fauna	Florid. Fauna	Carol. Fauna	Florid. Fauna
<i>Meleagris gallopavo</i>	Allegh. Fauna	Florid. Fauna	Allegh. Fauna	Florid. Fauna
<i>Bonasa umbellus</i>	Hudson. Fauna	Carol. Fauna	Hudson. Fauna?	Carol. Fauna
<i>Botaurus lentiginosus</i>	Hudson. Fauna	Carol. Fauna	Louis. Fauna	_____
<i>Ardetta exilis</i>	Canad. Fauna	Florid. Fauna	Louis. Fauna	_____
<i>Gallinago Wilsoni</i>	Hudson. Fauna	Allegh. Fauna	Louis. Fauna	_____
<i>Rhyacophilus solitarius</i>	Hudson. Fauna	Allegh. Fauna	Carol. Fauna	_____
<i>Tringoides macularius</i>	Hudson. Fauna	Louis. Fauna	Louis. Fauna	_____
<i>Limosa fedoa</i>	Arctic Coast?	Hudson. Fauna	Louis. Fauna	_____
<i>Rallus virginianus</i>	Allegh. Fauna	Louis. Fauna	Louis. Fauna	_____
<i>Porzana carolina</i>	Canad. Fauna	Carol. Fauna	?	_____
<i>Fulica americana</i>	Canad. Fauna	Carol. Fauna	Louis. Fauna	_____
<i>Nettion carolinensis</i>	Hudson. Fauna	Canad. Fauna	Carol. Fauna	_____

* Blanks in the third column indicate that the species ranges southward in the breeding season into the Tropical Realm. Blanks in the fourth column that the species retires wholly within the Tropical Realm in winter; in the fifth column, that the southern limit in winter is within the Tropical Realm.

2. *List of Species which breed throughout the greater Part of the Cold-temperate Portions of the North American Region, with Indications of their Boreal and Austral Limits of Distribution in the Eastern Province.*

Species.	Boreal Limit in Breeding Season.	Austral Limit in Breeding Season.	Boreal Limit in Winter.	Austral Limit in Winter.
<i>Turdus Pallasi</i>	Hudson. Fauna	Canad. Fauna	Carol. Fauna	— *
<i>Turdus Swainsoni</i>	Hudson. Fauna	Canad. Fauna	Florid. Fauna	—
<i>Regulus calendula</i>	Hudson. Fauna	Canad. Fauna	Carol. Fauna	—
<i>Regulus satrapa</i>	Hudson. Fauna	Canad. Fauna	Allegh. Fauna	—
<i>Anthus ludovicianus</i>	Arctic Coast	Hudson. Fauna	Carol. Fauna	—
<i>Helminthophaga celata</i>	Hudson. Fauna	Canad. Fauna	Carol. Fauna	—
<i>Wilsonia pusilla</i>	Hudson. Fauna	Allegh. Fauna	— *	—
<i>Collurio borealis</i>	Hudson. Fauna	Canad. Fauna	Canad. Fauna	Carol. Fauna
<i>Troglodytes hyemalis</i>	Hudson. Fauna	Canad. Fauna	Allegh. Fauna	—
<i>Chrysomitris pirus</i>	Hudson. Fauna	Canad. Fauna	Canad. Fauna	Florid. Fauna
<i>Curvirostra americana</i>	Hudson. Fauna	Canad. Fauna	Hudson. Fauna	Allegh. Fauna
<i>Zonotrichia leucophrys</i>	Hudson. Fauna	Canad. Fauna	Carol. Fauna	Louis. Fauna
<i>Scolecophagus ferrugineus</i>	Hudson. Fauna	Canad. Fauna	Carol. Fauna	—
<i>Perisoreus canadensis</i>	Hudson. Fauna	Canad. Fauna	Hudson. Fauna	Canad. Fauna
<i>Picoides arcticus</i>	Hudson. Fauna	Canad. Fauna	Hudson. Fauna	Canad. Fauna
<i>Picoides hirsutus</i>	Hudson. Fauna	Carol. Fauna	Hudson. Fauna	Canad. Fauna
<i>Falco columbarius</i>	Hudson. Fauna	Carol. Fauna	Canad. Fauna	Louis. Fauna
<i>Astur atricapillus</i>	Hudson. Fauna	Canad. Fauna	Hudson. Fauna	Allegh. Fauna
<i>Agialitis semipalmatus</i>	Arctic Coast	Hudson. Fauna	Louis. Fauna	—
<i>Phalaropus Wilsoni</i>	Arctic Coast	Canad. Fauna	Louis. Fauna	—
<i>Ereunetes pusillus</i>	Arctic Coast	Canad. Fauna	Carol. Fauna	—
<i>Gambetta melanoleuca</i>	Hudson. Fauna	Carol. Fauna	Louis. Fauna	—
<i>Gambetta flavipes</i>	Hudson. Fauna	Canad. Fauna	Louis. Fauna	—
<i>Numenius longirostris</i>	Hudson. Fauna	Canad. Fauna	Louis. Fauna	—
<i>Numenius hudsonius</i>	Hudson. Fauna	Canad. Fauna	Louis. Fauna	—
<i>Cygnus americanus</i>	Arctic Coast	Canad. Fauna	Allegh. Fauna	—
<i>Bernicla canadensis</i>	Arctic Coast	Canad. Fauna	Allegh. Fauna	—
<i>Mareca americana</i>	Arctic Coast	Canad. Fauna	Allegh. Fauna	—
<i>Fulix collaris</i>	Arctic Coast	Canad. Fauna	Canad. Fauna	—
<i>Aythya americana</i>	Arctic Coast	Hudson. Fauna	Carol. Fauna	—
<i>Aythya vallisneria</i>	Arctic Coast	Hudson. Fauna	Carol. Fauna	—
<i>Bucephala albeola</i>	Arctic Coast	Hudson. Fauna	Allegh. Fauna	—
<i>Erismatura rubida</i>	Arctic Coast	Hudson. Fauna	Allegh. Fauna	—
<i>Lophodytes cucullatus</i>	Arctic Coast	Canad. Fauna	Allegh. Fauna	—
<i>Pelecanus erythrorhynchus</i>	Arctic Coast	Canad. Fauna	Allegh. Fauna	—
<i>Graculus dilophus</i>	Arctic Coast	Canad. Fauna	Canad. Fauna	—
<i>Larus delawarensis</i>	Arctic Coast	Hudson. Fauna	Allegh. Fauna	—
<i>Chroicocephalus philadelphia</i>	Arctic Coast	Hudson. Fauna	Canad. Fauna	—

3. *List of Species which breed only in the Warm-temperate Portions of the North American Temperate Region, and range Southward in the Breeding Season into the Tropical American Realm.*

Species.	Boreal Limit in Breeding Season.	Austral Limit in Breeding Season.	Boreal Limit in Winter.	Austral Limit in Winter.
<i>Mimus polyglottus</i>	Carol. Fauna	— *	Louis. Fauna	— *
<i>Poliophtila cærulea</i>	Carol. Fauna	—	Louis. Fauna	—
<i>Icteria virens</i>	Carol. Fauna	—	—	—
<i>Stelgidopteryx serripennis</i>	Carol. Fauna	—	—	—
<i>Thryothorus Bewicki</i>	Carol. Fauna	—	Louis. Fauna ?	—
<i>? Sitta pusilla</i>	Louis. Fauna	—	Louis. Fauna	—
<i>Guiraca cærulea</i>	Carol. Fauna	—	—	—
<i>? Quiscalus major</i>	Louis. Fauna	—	Louis. Fauna	—
<i>Elanus leucurus</i>	Louis. Fauna	—	—	—
<i>Polyborus tharus</i>	Florid. Fauna	—	Florid. Fauna	—
<i>Craxirex uncinatus</i>	Louis. Fauna	—	—	—
<i>Chamaepelia passerina</i>	Louis. Fauna	—	Louis. Fauna	—
<i>Tantalus loculator</i>	Louis. Fauna	—	Louis. Fauna	—
<i>Garzetta candidissima</i>	Carol. Fauna	—	Louis. Fauna	—
<i>Herodias egretta</i>	Carol. Fauna	—	Louis. Fauna	—
<i>Himantopus nigricollis</i>	Carol. Fauna ?	—	—	—
<i>Rallus elegans</i>	Carol. Fauna	—	Louis. Fauna	—
<i>Phoenicopterus ruber</i>	Florid. Fauna	—	Florid. Fauna ?	—
<i>Querquedula cyanoptera</i>	?	—	Florid. Fauna	—
<i>Pelecanus fuscus</i>	Carol. Fauna	—	Louis. Fauna	—
<i>Tachypetes aquila</i>	Florid. Fauna	—	—	—
<i>Plotus anhinga</i>	Louis. Fauna	—	Florid. Fauna	—

* Within the Tropical Realm

4. *List of Species whose Breeding Range extends throughout the greater Part of the North American Realm, and Southward into the Tropical Realm, with Indications of their Boreal and Austral Limits in the Eastern Province.*

Species.	Boreal Limit in the Breeding Season.	Austral Limit in the Breeding Season.	Boreal Limit in Winter.	Austral Limit in Winter.
<i>Dendroeca aestiva</i>	Hudson. Fauna	—* *	—	—*
<i>Grus canadensis</i>	Arctic Coast	—	Carol. Fauna	—
<i>Butorides virescens</i>	Allegh. Fauna	—	Louis. Fauna	—
<i>Ardea herodias</i>	Hudson. Fauna	—	Carol. Fauna	—
<i>Hæmatopus palliatus</i>	Arctic Coast	—	Louis. Fauna	—
<i>Ægialitis vociferus</i>	Allegh. Fauna	—	Louis. Fauna	—
<i>Recurvirostra americana</i>	Hudson. Fauna	Louis. Fauna ?	Louis. Fauna	—
<i>Symphemia semipalmata</i>	Canad. Fauna	—	Louis. Fauna	—
<i>Aix sponsa</i>	Canad. Fauna	—	Carol. Fauna	—
<i>Podilymbus podiceps</i>	Hudson. Fauna	—	Carol. Fauna	—

* Within the Tropical Realm.

5. *List of Species whose Breeding Habitat includes the greater Part of both North and South America, with Indications of their Boreal Limit, both in the Breeding Season and in Winter.*

Species.	Boreal Limit in the Breeding Season.	Austral Limit in the Breeding Season.	Boreal Limit in Winter.	Austral Limit in Winter.
<i>Cathartes aura</i>	Allegh. Fauna	—	Carol. Fauna	—
<i>Cathartes atratus</i>	Louis. Fauna	—	Louis. Fauna	—
* <i>Pandion haliaëtus</i>	Arctic Coast	—	Louis. Fauna	—
<i>Falco sparverius</i>	Hudson. Fauna	—	Hudson. Fauna	—
* <i>Falco peregrinus</i>	Hudson. Fauna	—	Canad. Fauna	—
<i>Bubo virginianus</i>	Hudson. Fauna	—	Hudson. Fauna	—
* <i>Strix flammea</i>	Carol. Fauna	—	Carol. Fauna	—
* <i>Otus vulgaris</i>	Hudson. Fauna	—	Canad. Fauna ?	—
* <i>Otus brachyotus</i>	Arctic Coast	—	Hudson. Fauna ?	—

* Also circumpolar species.

Summary of the Preceding Five Tables.—The total number of species given in the above lists of the species characteristic (mainly exclusively so) of the North American Temperate Region is 135. Of these 38 are restricted in the breeding season in their austral range to the Cold-temperate District; about one third of them, chiefly natatorial species, reach the Arctic coast; 61 are similarly mainly limited to the Middle-temperate District, but two or three reach the Arctic coast, and nearly one third range into the Hudsonian Fauna; 21 are limited in their boreal range to the Warm-temperate District, the greater part of which, even in the breeding season, range southward into the tropics. Of the whole number, 90 are land birds, 23 being raptorial species. Of the remaining 45 water birds, 7 are herons, 20 are *Grallæ*, and 18 are *Natatores*, 12 of the latter being *Anatidæ*.

In the list of those whose breeding habitat is the cold-temperate portions of the continent (Table 1), 20 of the species are aquatic and

18 terrestrial; of those breeding throughout the greater part of the continent, 10 only are aquatic and 51 are terrestrial; of those breeding in the warm-temperate portions of the continent, 9 are aquatic and 9 terrestrial; of the 10 wide-ranging species, whose breeding habitats embrace not only nearly the whole of temperate North America, but extend also into the tropics, 1 only is a land bird, 3 being *Herodiones*, 4 *Grallæ*, and 2 *Natatores*. Of the 9 species given in the Fifth Table, which range in the breeding season throughout both the North American and South American continents, none are aquatic; 4 are owls, 3 hawks, and 2 vultures. The most numerous family, and one of those almost exclusively characteristic of the North American Temperate Region (the *Sylvicolidæ*), has but three species which range across the continent, and only one of these is a typical representative of the family.

IV. SPECIES LIMITED IN LONGITUDE TO THE EASTERN PROVINCE OF THE NORTH AMERICAN TEMPERATE REGION.

1. List of Species restricted in the breeding Season to the Cold-temperate Portion of the Eastern Province, with Indications of their Boreal and Austral Limits.

Species.	Boreal Limit in Breeding Season.	Austral Limit in Breeding Season.	Boreal Limit in Winter.	Austral Limit in Winter.
<i>Mniotilta varia</i>	Hudson. Fauna	Carol. Fauna	— * *	— * *
<i>Parula americana</i>	Canad. Fauna	Carol. Fauna	Florid. Fauna	—
<i>Geothlypis philadelphia</i>	Allegh. Fauna	?	—	—
<i>Oporornis agilis</i>	Canad. Fauna	Allegh. Fauna ?	—	—
<i>Helminthophaga chrysoptera</i>	Canad. Fauna	Carol. Fauna	—	—
? <i>Helminthophaga peregrina</i>	Canad. Fauna	Carol. Fauna	—	—
<i>Helminthophaga ruficapilla</i>	Hudson. Fauna	Carol. Fauna	—	—
<i>Dendroeca coronata</i>	Hudson. Fauna	Canad. Fauna	Carol. Fauna	—
<i>Dendroeca castanea</i>	Hudson. Fauna?	Canad. Fauna	—	—
<i>Dendroeca striata</i>	Hudson. Fauna	Canad. Fauna	—	—
<i>Dendroeca Blackburniæ</i>	Canad. Fauna	Allegh. Fauna	—	—
<i>Dendroeca cærulescens</i>	Canad. Fauna ?	Canad. Fauna	—	—
<i>Dendroeca maculosa</i>	Canad. Fauna ?	Canad. Fauna	—	—
<i>Dendroeca virens</i>	Hudson. Fauna	Allegh. Fauna	—	—
<i>Dendroeca palmarum</i>	Hudson. Fauna	Canad. Fauna	Louis. Fauna	—
<i>Perissoglossa tigrina</i>	Canad. Fauna ?	Allegh. Fauna	—	—
<i>Euthlypis canadensis</i>	Hudson. Fauna	Canad. Fauna	—	—
<i>Setophaga ruticilla</i>	Hudson. Fauna	Allegh. Fauna	—	—
<i>Vireosylva olivacea</i>	Hudson. Fauna	Louis. Fauna	Florid. Fauna	—
<i>Vireosylva philadelphia</i>	Hudson. Fauna	Allegh. Fauna	—	—
<i>Cistothorus stellaris</i>	Hudson. Fauna	Carol. Fauna	Louis. Fauna	—
<i>Parus hudsonicus</i>	Hudson. Fauna	Canad. Fauna	Hudson. Fauna	Canad. Fauna
<i>Zonotrichia albicollis</i>	Hudson. Fauna	Canad. Fauna	Louis. Fauna	—
<i>Junco hyemalis</i>	Hudson. Fauna	Canad. Fauna	Allegh. Fauna	Louis. Fauna
<i>Spizella monticola</i>	Hudson. Fauna	Canad. Fauna	Allegh. Fauna	Louis. Fauna
<i>Passerella iliaca</i>	Hudson. Fauna	Canad. Fauna	Carol. Fauna	Louis. Fauna
<i>Dolichonyx oryzivora</i>	Allegh. Fauna	Allegh. Fauna	—	—
<i>Tetrao canadensis</i>	Arctic Coast	Canad. Fauna	Hudson. Fauna?	Canad. Fauna
<i>Ægialitis melodus</i>	Canad. Fauna	Carol. Fauna	Louis. Fauna	—
<i>Limosa hudsonica</i>	Arctic Coast ?	Hudson. Fauna	Louis. Fauna	—
<i>Numenius borealis</i>	Arctic Coast ?	Hudson. Fauna	Louis. Fauna	—
<i>Porzana noveboracensis</i>	Hudson. Fauna	Carol. Fauna	Louis. Fauna	—
<i>Anas obscura</i>	Hudson. Fauna	Allegh. Fauna	Allegh. Fauna	Florid. Fauna
<i>Querquedula discors</i>	Hudson. Fauna	Canad. Fauna	Allegh. Fauna	—
<i>Camptolæmus labradorius</i>	Arctic Coast	Hudson. Fauna	—	Canad. Fauna

* The blanks in the fourth and fifth columns indicate that the limit in question is within the Tropical Realm.

2. *List of Species which breed throughout the Middle-temperate Portions of the Eastern Province, with Indications of their Boreal and Austral Limits in the Eastern Province.**

Species.	Boreal Limit in the Breeding Season.	Austral Limit in the Breeding Season.	Boreal Limit in Winter.	Austral Limit in Winter.
<i>Turdus mustelinus</i>	Allegh. Fauna	Louis. Fauna ?	—	—
<i>Galeoscoptes carolinensis</i>	Allegh. Fauna	Florid. Fauna ?	Louis. Fauna	—
<i>Harporhynchus rufus</i>	Allegh. Fauna	Florid. Fauna ?	Louis. Fauna	—
<i>Sialia sialis</i>	Allegh. Fauna	Louis. Fauna	Carolin. Fauna	—
<i>Dendroeca pennsylvanica</i>	Canad. Fauna	Allegh. Fauna	—	—
<i>Dendroeca discolor</i>	Allegh. Fauna	—	—	—
<i>Dendroeca caerulea</i>	Allegh. Fauna	Carolin. Fauna	—	—
<i>Wilsonia mitrata</i>	Allegh. Fauna	Louis. Fauna	—	—
<i>Pyranga rubra</i>	Allegh. Fauna	Louis. Fauna	—	—
<i>Vireo noveboracensis</i>	Allegh. Fauna	Florid. Fauna ?	Louis. Fauna	—
<i>Lanivireo flavifrons</i>	Allegh. Fauna	—	—	—
<i>Lophophanes bicolor</i>	Carolin. Fauna	Louis. Fauna ?	Carolin. Fauna	Florid. Fauna
<i>Coturniculus passerinus</i>	Allegh. Fauna	Louis. Fauna	Carolin. Fauna ?	—
<i>Coturniculus Henslowi</i>	Allegh. Fauna	Louis. Fauna	Carolin. Fauna ?	—
<i>Ammodromus caudacutus</i>	Allegh. Fauna	Louis. Fauna	Louis. Fauna	—
<i>Ammodromus maritimus</i>	Carolin. Fauna	Louis. Fauna	Louis. Fauna	—
<i>Spizella pusilla</i>	Allegh. Fauna	Louis. Fauna ?	Carolin. Fauna	—
<i>Euspiza americana</i>	Carolin. Fauna	—	—	—
<i>Hedymeles ludovicianus</i>	Allegh. Fauna	Carolin. Fauna	—	—
<i>Cyanospiza cyanea</i>	Allegh. Fauna	—	—	—
<i>Cardinalis virginianus</i>	Carolin. Fauna	—	Carolin. Fauna	—
<i>Pipilo erythrophthalmus</i>	Allegh. Fauna	Florid. Fauna	Louis. Fauna	—
<i>Icterus spurius</i>	Allegh. Fauna	Louis. Fauna	—	—
<i>Icterus baltimore</i>	Allegh. Fauna	Louis. Fauna	—	—
<i>Corvus ossifragus</i>	Carolin. Fauna	Florid. Fauna ?	Louis. Fauna	—
<i>Centurus carolinus</i>	Carolin. Fauna	—	Carolin. Fauna	—
<i>Melanerpes erythrocephalus</i>	Allegh. Fauna	Florid. Fauna ?	Louis. Fauna	—
<i>Antrostomus vociferus</i>	Allegh. Fauna	Florid. Fauna	Louis. Fauna	—
<i>Coccygus americanus</i>	Allegh. Fauna	—	—	—
<i>Coccygus erythrophthalmus</i>	Allegh. Fauna	—	—	—
<i>Ortyx virginianus</i>	Allegh. Fauna	—	Allegh. Fauna	—
<i>Cupidonia cupido</i>	Allegh. Fauna	Carolin. Fauna ?	Allegh. Fauna	Louis. Fauna

* The blanks in this and the following tables have the same significance as in the last preceding table.

3. *List of Species which breed throughout the Temperate Portions of the Eastern Province, with Indications of their Boreal and Austral Limits.*

Species.	Boreal Limit in the Breeding Season.	Austral Limit in the Breeding Season.	Boreal Limit in Winter.	Austral Limit in Winter.
<i>Seiurus aurocapillus</i>	Canad. Fauna	Carolin. Fauna	Florid. Fauna	—
<i>Seiurus noveboracensis</i>	Hudson. Fauna	Louis. Fauna	—	—
<i>Cyanura cristata</i>	Canad. Fauna	Florid. Fauna	Allegh. Fauna	Florid. Fauna
<i>Sphyrapicus varius</i>	Canad. Fauna ?	Florid. Fauna ?	Carolin. Fauna	—
<i>Colaptes auratus</i>	Hudson. Fauna	—	Carolin. Fauna	—
<i>Trochilus colubris</i>	Canad. Fauna	—	Florid. Fauna ?	—
<i>Chaetura pelagica</i>	Canad. Fauna	Louis. Fauna ?	—	—
<i>Grus americanus</i>	Canad. Fauna	Florid. Fauna	Louis. Fauna	—
<i>Aetiturus Bartramius</i>	Canad. Fauna	—	Louis. Fauna	—
<i>Sterna antillarum</i>	Canad. Fauna	—	—	—
<i>Dendroeca pinus</i>	Hudson. Fauna ?	Louis. Fauna ?	Carolin. Fauna	—
<i>Quiscalus purpureus</i>	Hudson. Fauna	—	Carolin. Fauna	—

4. *List of Species which breed in the Eastern Province only within the Warm-temperate and Subtropical Districts.*

Species.	Boreal Limit in the Breeding Season.	Austral Limit in the Breeding Season.	Boreal Limit in Winter.	Austral Limit in Winter.
<i>Prothonotaria citrea</i>	Carolín. Fauna	—	—	—
<i>Oporornis formosus</i>	Allegh. Fauna	Louis. Fauna ?	—	—
<i>Helmitherus vermicivorus</i>	Carolín. Fauna	Florid. Fauna ?	—	—
<i>Helmitherus Swainsoni</i>	Louis. Fauna	—	—	—
<i>Helminthophaga pinus</i>	Carolín. Fauna ?	—	—	—
<i>Helminthophaga Bachmani</i>	Louis. Fauna	—	—	—
<i>Dendroeca dominica</i>	Louis. Fauna	—	—	—
<i>Vireosylva barbatula</i>	Florid. Fauna	—	—	—
<i>Pyranga aestiva</i>	Carolín. Fauna	—	—	—
<i>Thryothorus ludovicianus</i>	Carolín. Fauna	—	Louis. Fauna	—
<i>Peuceea aestivalis</i>	Louis. Fauna	—	Florid. Fauna	—
<i>Cyanospiza ciris</i>	Louis. Fauna	—	—	—
<i>Tyrannus dominicensis</i>	Florid. Fauna	—	—	—
<i>Campephilus principalis</i>	Carolín. Fauna	—	Louis. Fauna	—
<i>Picus borealis</i>	Louis. Fauna	—	Louis. Fauna	—
<i>Conurus carolinensis</i>	Louis. Fauna	—	Louis. Fauna	—
<i>Crotophaga ani</i>	Florid. Fauna	—	—	—
<i>Antrostomus carolinensis</i>	Louis. Fauna	—	Louis. Fauna	—
<i>Nauclerus furcatus</i>	Carolín. Fauna	—	—	—
<i>Ictinia mississippiensis</i>	Louis. Fauna	—	—	—
<i>Rostrhamus sociabilis</i>	Florid. Fauna	—	—	—
<i>Florida cerulea</i>	Carolín. Fauna	—	Louis. Fauna	—
<i>Nyctherodius violaceus</i>	Carolín. Fauna	—	—	—
<i>Demigretta Pealei</i>	Florid. Fauna	—	Florid. Fauna	—
<i>Demigretta ludoviciana</i>	Carolín. Fauna	—	Louis. Fauna	—
<i>Ibis alba</i>	Carolín. Fauna	—	Louis. Fauna	—
<i>Platalea ajaja</i>	Louis. Fauna	—	—	—
<i>Ægialitis Wilsonius</i>	Carolín. Fauna	—	Louis. Fauna	—
<i>Aramus giganteus</i>	Florid. Fauna	—	Florid. Fauna	—
<i>Rallus crepitans</i>	Carolín. Fauna	—	Louis. Fauna	—
<i>Porzana jamaicensis</i>	Carolín. Fauna	—	Florid. Fauna	—
<i>Gallinula galeata</i>	Carolín. Fauna	—	Louis. Fauna	—
<i>Gallinula martinica</i>	Carolín. Fauna	—	Louis. Fauna	—
<i>Sula fiber</i>	Louis. Fauna	—	Florid. Fauna	—
<i>Graculus floridanus</i>	Louis. Fauna	—	Florid. Fauna	—
<i>Plotus aninga</i>	Carolín. Fauna	—	Florid. Fauna	—
<i>Chrococephalus atricilla</i>	Allegh. Fauna	—	Florid. Fauna	—
<i>Thalasseus acutiflavus</i>	Louis. Fauna	—	—	—
<i>Anois stolidus</i>	Florid. Fauna	—	—	—
<i>Halipiana fuliginosa</i>	Florid. Fauna	—	—	—
<i>Rhynchops nigra</i>	Carolín. Fauna	—	Florid. Fauna	—

Summary of the Four Preceding Tables.—About one hundred and twenty species occur in the Eastern Province of the North American Temperate Region that do not appear as regular residents in the Western Province of the same region, of which a small proportion are in part tropical. Of these one hundred and twenty, thirty-five are northern, or range in the breeding season only over the cold-temperate portions of the Eastern Province; twenty-eight of the latter being land birds, and only seven aquatic. Eighteen species of the land birds belong to the single family of the *Sylvicolidae*. About one fourth of the Eastern Province species (thirty-two), *all land birds*, range in the breeding season over only the middle-temperate part of the province. Of these only three belong to the family *Sylvicolidae*, and only one is a typical representative of that group; ten belong to the family *Fringillidae*, three

to the *Turdidæ*, and two each to the *Icteridæ*, *Picidæ*, *Cuculidæ*, and *Tetraonidæ*; several other families have one representative each. The total absence of any species of *Falconidæ*, *Strigidæ*, *Herodiones*, *Grallæ*, and *Natatores* is one of the most striking features in the list of the species restricted to the Eastern Province.

Twelve of the Eastern Province species breed throughout the greater part of the province, three of which are *Sylvicolidæ*, two are *Picidæ*, one is a humming-bird, one a wader, and one a tern.

Forty-one of the one hundred and twenty species restricted in longitudinal range to the Eastern Province extend so far into the Tropical American Realm in the breeding season as to be essentially tropical species, exclusive of a considerable number that appear only in the Floridian Fauna. Twenty-one of these are land birds and twenty aquatic; the latter embracing six *Herodiones*, six *Grallæ*, and eight *Natatores*, five of which are terns. The land species embrace three hawks, two species of *Fringillidæ*, seven of *Sylvicolidæ*, two of *Picidæ*, and one each of seven other families.

GENERAL REMARKS ON THE DISTRIBUTION AND MIGRATION OF THE BIRDS OF THE EASTERN PROVINCE.

The preceding tables, illustrative of the geographical distribution of the birds of the Eastern Province of the North American Temperate Region, and the summary remarks already given respecting them, indicate a number of interesting general facts.

I. The species which have the greatest longitudinal range in the breeding season are the hawks, owls, and vultures, the swallows, the *Turdinæ* or typical thrushes, the woodpeckers and flycatchers, and the water birds; among the latter, especially the *Scolopacidæ*, the *Charadriidæ* and their allies, the *Anatidæ*, and the *Laridæ*; in fact, nearly all the *Natatores*. All the land birds ranging widely in longitude are hence species which possess highly developed powers of flight, and have also a wide latitudinal range. The few circumboreal *Natatores*, which have only moderate or greatly reduced powers of flight, possess great power of locomotion in the water. Their habitat is, moreover, not only generally the sea-shores, but the boreal shores of the converging continents of the northern hemisphere. Hence all the species having a wide geographical range—as the circumpolar and continental—are either pre-eminently strong fliers or powerful swimmers. It also ex-

plains the occurrence of the large proportion of long-winged birds, and especially of the preponderance of the water birds, in the three first primary divisions given above of the birds of the Eastern Province, namely, the cosmopolitan, the circumpolar, and the continental, and the small proportion of such species among those restricted in their longitudinal range to the Eastern Province. Most of the circumpolar species are also boreal ones.

II. The aquatic species, while forming only about four tenths of the birds found in the Eastern Province, greatly predominate over the land species in the boreal regions, in the Arctic Realm they outnumbering the land birds in the proportion of five to one, or form eight tenths of the whole. In the Cold-temperate District of the North American Region the water-birds form about six tenths of the whole; in the Middle-temperate Districts, between four and five tenths; in the Warm-temperate District, rather less than four tenths. Farther southward, although a few groups (as the *Rallidæ*, *Herodiones*, and *Sterninæ*) are more numerously represented, the relative proportion of water birds to the terrestrial seems scarcely to increase. In the breeding season, however, a numerical comparison of the land and water birds yields very different results, in respect to the proportion characteristic of localities situated under different parallels of latitude. Passing from the extreme boreal regions southward, the number of *Grallæ* (exclusive of the *Paludicolæ*), *Anatidæ*, *Larinæ*, and *Lestridinæ* decreases rapidly, so that the number of the *Grallæ* (exclusive of the rails and their allies) is reduced in the breeding season, in the warm-temperate parts of the Eastern Province, to only seven or eight species, the *Anatidæ* to one (*Aix sponsa*), the *Larinæ* to one (*Chræcocephalus atricilla*), and the *Lestridinæ* disappear entirely.

III. A large proportion of the accessions to the land birds near the tropics being species of a comparatively low grade of structure, the prevalence of the water birds in the arctic and subarctic faunæ, the comparative absence of water birds in the temperate latitudes, and the great development here of the higher groups of the land birds, give to the temperate regions the maximum proportion of high forms of avine life, — a fact as true in respect to mammalian life as it is of birds.

IV. In respect to the distribution and relative development of particular families, the *Sittidæ* (*Sittæ*), the *Paridæ*, and the *Alcidæ* are alone restricted to the North Temperate Realm. The species of these groups

(except the *Alcidæ*) are also sedentary throughout nearly their whole range. In the Eastern Province, *Larus*, *Stercorarius*, and their allies, as well as *Colymbus*, are restricted in the breeding season to its northern half, as are also, as already remarked, most of the water birds, except the *Rallidæ* and the *Herodiones*, which are chiefly southern. The representatives of the *Troglodytidæ* and *Icteridæ* increase rapidly in number towards the tropics, while the *Miminæ* and several genera of the more brightly colored *Fringillidæ* are confined to the southern half of the province. The *Sylvicolidæ*, the most exclusively distinctive family of the North American temperate region, reaches its maximum development in the Middle and Cold-temperate Districts. The section *Sylvicoleæ*, and especially the genus *Dendræca*, is more numerous represented in the Eastern Province than in the Western, and the greater part breed in the colder latitudes, their "metropolis" during the breeding season being the Canadian Fauna of the Eastern Province and the corresponding fauna of the Western. The species of the section *Vermivoreæ* (genera *Helmitherus* and *Helminthophaga*) have a wider longitudinal distribution than the *Dendræcæ*, the species of which genus are mainly restricted either to the Eastern or to the Western Province. The two species of *Helmitherus* are southern in their distribution, while four or five of the six *Helminthophagæ* are northern.

V. At the extreme north, or from the Arctic coast southward to the Canadian Fauna, nearly all the birds are migratory, owing to the extreme severity of the winter season; they also spend a smaller portion of the year at their breeding stations than do the species which breed farther to the southward. Even as far south as the Alleghanian and Carolinian Faunæ, the greater proportion of the species are to a greater or less extent migratory. In the Carolinian and Louisianian Faunæ a much greater proportion are resident, even including many species whose boreal limit of distribution is the Carolinian Fauna. From the Hudsonian Fauna southward many species are found the whole year at the same localities, and are hence termed "resident," though the individuals representing such species are migratory, there being a general movement of the winter habitat southward, but too limited to carry the wave of migration entirely beyond the southern limit of the summer habitat of these species.

VI. The representatives of the various groups differ from each other

widely, as is well known, in respect to the extent of their migrations. Those of a few families (as the *Tetraonidæ*, the *Picidæ*, *Sittidæ*, *Corvidæ*, and *Strigidæ*) are nearly sedentary, the nature of their food being such that the supply is almost equally sure at all seasons. The insectivorous species have the most extended migratory range; the piscivorous, the graminivorous, and the raptorial the least of the non-sedentary species, the migrations of the latter being mainly governed by those of their prey. Hence the wood warblers (*Sylvicolidæ*), the flycatchers (*Tyrannidæ*), and the swallows make the longest journeys, and leave their breeding stations the earliest. Requiring apparently a temperature in winter similar to that of their summer habitats, as well as a constant supply of insect food, they begin their southward journeys almost before the close of the short northern summer, proceeding gradually southward during the autumn months to pass the winter in the tropics. The *Grallæ* have also to seek districts almost wholly beyond the reach of severe frosts, their food being only accessible to them where the ground continues unfrozen. The *Natatores* also necessarily migrate to localities where the streams and estuaries are nearly free from ice; the strictly littoral and pelagic species hence making shorter journeys than the inland species.

The migratory insessorial birds that pass the winter wholly or in part within the Eastern Province are principally fringilline species. The others are a single flycatcher (*Sayornis fuscus*), the northern members of the *Icteridæ*, two species of *Turdus*, three of *Miminæ*, three wrens, and three or four sparrows. All of these species are resident the whole year in those sections to which the northern members of these species mainly resort in winter. In these species there is hence only a partial recession southward in winter from the northern portions of their respective summer habitats. Most of the *Fringillidæ*, however, which pass the summer within or to the northward of the Alleghanian Fauna, remove wholly in winter from their summer stations. While some in winter barely abandon their summer stations, as shown in the preceding tables, of distribution, others pass entirely over one fauna, throughout which they occur only as spring and autumn passengers; others in a similar way pass over two faunæ in reaching their winter quarters. Hence some which breed in the Canadian Fauna and farther northward pass only into the Alleghanian and Carolinian Faunæ in winter, while others pass over the Alleghanian

into the Carolinian, and others over both the Alleghanian and the Carolinian into the Louisianian.

Some species which in general breed far to the northward of the tropics, to which they retire in winter, are also known to breed on the mountains within the tropics (as *Dendroica coronata* and *Perissoglossa tigrina*), and doubtless many others will be found to do so when the mountain faunæ of these regions become fully known; it being already well ascertained that there is a migration from the plains and lowlands to the mountains (more especially in the Tropical Realms) as well as (in the northern hemisphere) from the south northward.

VII. The lack of suitable food and the low temperature in winter in northern latitudes are evidently the causes which impel so many species to leave their breeding stations at that season to seek a warmer zone. While in most cases a degree of cold sufficient to cut off the supply of food of any species, especially of the insectivorous ones, would of itself also prove fatal to the birds themselves, it is by no means the case with the baccivorous and graminivorous species, their winter migrations appearing to be primarily and principally controlled by the accessibility of their food. This is evidently indicated by the irregular dispersion in winter of such species near the boreal limit of their range at that season, they being numerous where their food abounds and entirely absent in the immediately adjoining districts.*

VIII. The breeding range, as well as the migratory range, differs greatly not only among the species of different families (nearly all the species of some families having a wide range, while nearly all the species of other families have a quite restricted range, as in the *Corvidæ* and *Picidæ*, for instance, as compared with the *Sylvicolidæ*), but also among those of the same family and even of the same genus. The two extremes are well illustrated by the osprey or fish-hawk and

* These remarks are illustrated by the winter distribution of the robin and the cedar-bird in the Alleghanian Fauna, and by the sudden southward incursion of the snow-buntings and other northern sparrows when deep snows suddenly render their food more than usually difficult to procure in their usual winter resorts. The early return of birds to their breeding stations, — their real homes, — as soon as the causes that impelled their winter migration are removed, is further corroborative of the same view. Most of even the insectivorous species visit regions in winter whose average winter temperature differs but little from that of their breeding stations, and when the excessive heats of spring and summer arrive in the southern latitudes, they gradually retire again to their northern breeding stations, keeping pace in their migration with the northward advance of the summer warmth.

the bobolink, the one having an almost cosmopolitan breeding range, while the breeding range of the other is nearly or quite restricted to the Alleghanian Fauna. Several of the *Sylvicolidæ* have a breeding range as restricted as the bobolink, while a few other species of the same family breed throughout nearly the whole of North America. One of the species of *Dendræca* (*D. æstiva*) has this wide breeding range, while other species of the same genus appear to breed only in the Canadian Fauna.

IX. Species which have a wide breeding range usually present a greater or less number of easily distinguishable local forms, which merge generally the one into the other in the regions lying between the localities at which these several forms are most fully developed. A part of these local forms have received distinctive names, and have of late been quite commonly regarded as distinct species, while many are as yet not so regarded. Every year additional races of this character are discovered, and doubtless many still remain unknown. Much time will probably elapse before naturalists will generally agree as to their true character and relations, though evidence indicative of their being the result of general and uniformly acting laws of geographical variation is apparently by no means wanting. The difference in color, size, form of the bill, length of the tail, etc., that appear almost universally to obtain between the northern and southern representatives of the same species, have already been sufficiently dwelt upon in the preceding pages; but the insertion of a few species in the list of those alleged above (Class IV of the preceding tables) to range across the North American continent calls for an additional word in respect to the differences which have led to the specific separation of the western representatives of these species from their eastern representatives, or to suggestions that they might prove to be specifically distinct. Most of the cases of this kind have been distinguished in the tables under Class IV by the prefix of a *â* *[?]* before their names. In all these cases almost the sole difference alleged for the separation of the western forms is that of either the darker or brighter or, in other words, the more intense colors of those from the Pacific coast; this character being always the one most strongly urged as distinguishing them, and not unfrequently the only one, especially in those species that breed wholly to the northward of the latitude of San Francisco. The frequency of this difference seems to be a strong reason for regarding

it as the result of a general law, and to remove it from the category of genuine specific distinctions.*

X. The number of species which breed in the American Arctic Fauna appears to be not far from sixty. In the Hudsonian Fauna the number is increased to upwards of one hundred and fifty, in the Canadian to probably about one hundred and sixty. In the Alleghanian the number is nearly one hundred and forty; in the Carolinian about one hundred and thirty-five; in the Louisianian about one hundred and thirty. The Hudsonian and Canadian Faunæ hence have a greater number of species, in areas of the same extent, and probably a far greater number of individuals, than the Carolinian and Louisianian Faunæ. In respect to the number of individuals, it is evident that this must result, in consequence of the hordes of wading and swimming birds, of thrushes, sparrows, and *Sylvicolidæ* that pass through the southern and middle districts of the Eastern Province to breed in its boreal portions; few of the species that breed at the southward being there as numerously represented as are scores of species that breed exclusively at the northward. Taking the whole number of species found at particular localities during the year, there is a constant increase in number to the southward, the increase, however, being less rapid from the southern boundary of the Canadian Fauna southward than from that point northward.† There is also a steady

* In addition to the list of examples of this variation already cited in Part III of this paper, the following may be added as marked instances: *Regulus satrapa*, Pacific coast specimens forming the variety *olivaceus* Baird; *Troglodytes hyemalis*, Pacific coast specimens forming the variety *pacificus* Baird; *Cistothorus palustris*, Pacific coast specimens forming the variety *paludicola* Baird; *Helminthophaga celata*, Pacific coast specimens being, according to Professor Baird, "much brighter and clearer yellowish beneath and olivaceous above," than those from the interior of North America; *Myiodiotes* (= *Wilsonia*) *pusilla*; *Hirundo bicolor*; *Collurio excubitoroides* (= *C. ludovicianus*), Pacific coast specimens being darker than those from the Mississippi Valley, and much darker than those from the Plains (Baird); *Carpodacus purpureus*, Pacific coast specimens being darker and forming the *C. californicus* of Baird; *Melospiza melodia*, the darker Pacific coast specimens forming the *M. Heermanni* Baird, etc.

† There seems to have been no exhaustive list published of the birds occurring at any locality north of the Alleghanian Fauna. Dr. Richardson's list is the largest, and gives two hundred and thirty-eight as the whole number known in 1831 to inhabit British North America north of the Canadas and east of the Rocky Mountains. Probably this number, and even more, may occur at a single locality on the Saskatchewan; but probably not more than two hundred and sixty or seventy. Three hundred and ten have been detected in Massachusetts, including those of irregular and very rare

increase southwards in the number and even in the proportion of species which are resident at the same locality the whole year. But from the absence of exhaustive lists of the species occurring at numerous localities, differing in latitude, it is difficult to make at present a wholly satisfactory numerical comparison of the different ornithological faunæ.*

occurrence ; three hundred and twenty-seven in the vicinity of New York City (*Lawrence*), and three hundred and forty-three in New Jersey (*Turnbull*). The number given by Ross as observed by him in the "Mackenzie's River District" is one hundred and ninety-two. The greatest number I have seen recorded from any restricted locality within the American tropical Realm is five hundred and forty, the number given from Costa Rica by Messrs. Lawrence and Salvin.

* Dr. Richardson in 1831, found that the number of species "known to rear their young on the banks of the Saskatchewan" amounted to one hundred and forty-one. At least twenty species more may now be safely added. Bonaparte, in 1827, estimated the number of species breeding at Philadelphia to be one hundred and four. Messrs. Coues and Prentiss in their list of the birds of the District of Columbia, published in 1861, mention forty-four species as being permanent residents, and fifty-nine others as summering, making one hundred and three that probably breed in the District, — one less than the number given by Bonaparte as breeding at Philadelphia. Messrs. S. F. and W. M. Baird gave, in 1844, one hundred and nine species as breeding at Carlisle, Pennsylvania. The three latter being inland localities, they may properly be compared with the Saskatchewan district. The numerous lakes at the latter locality, however, afford favorite breeding places for numerous water birds, while few such localities are afforded by the other localities mentioned; but since few water birds breed so far south as these localities, the difference in this respect is a fact of small importance. Dr. Turnbull, in 1869, gave the number of permanently resident species in "East Pennsylvania and New Jersey" as fifty-two, and the number of summer visitors as one hundred and fourteen, making a total of one hundred and sixty-six species that occur there in summer; but the area included in this list is more extended, and embraces a greater variety of surface than in the other cases, and includes several strictly coast species. Farther than this, an examination of his list shows that at least thirty of the one hundred and sixty-six are either wholly of accidental or of very rare occurrence, and hence do not regularly (many of them never) breed at the locality named. The number of resident species in Massachusetts is not far from thirty, of summer visitors one hundred and six, making one hundred and thirty-six that are more or less frequent in summer, — a number considerably less than undoubtedly occur in an equal area on the Saskatchewan. Mr. C. J. Maynard, in his careful analysis of the birds of Eastern Massachusetts (*Naturalist's Guide*, Part II, pp. 162 – 164, 1870), gives only one hundred and fifteen as being known with certainty to breed in the eastern half of that State, one or two of which cannot be considered as breeding there regularly. While this somewhat exceeds the number generally given as breeding at localities more to the southward, it is far less than the number given by Dr. Richardson as breeding on the Saskatchewan, and much less than the number now well known to be found there in summer. Dr. Coues, in his "Synopsis of the Birds of South Carolina," indicates only about one hundred and thirty-five as being known to breed regularly in that State.

APPENDIX TO PART V.

LIST OF AUTHORITIES.

IN the following list are given the titles of general works and special papers that may be profitably consulted in a study of the geographical distribution of the birds of North America. An attempt has been made to cite all the papers of much importance bearing upon this subject that have appeared prior to the beginning of the year 1870, those published in this country having been also brought down to the present date (April, 1871). Some of those published in the transactions of foreign societies during 1869 have been necessarily omitted, as also a large proportion of those published in 1870, since most of these publications are usually several months in reaching this country.

In compiling the present list I have been greatly aided by the "List of Authorities" published by Professor Baird in the Appendix of his Report on the Birds of North America in 1858; Agassiz's "Bibliographia Zoölogiæ"; Carus and Engleman's "Bibliotheca Zoölogica," ending with the year 1860; by Dr. G. Hartlaub's "Bericht über die Leistungen in der Naturgeschichte der Vögel," in Wiegmann's Archiv für Naturgeschichte, and by Professor Alfred Newton's admirable ornithological record in Dr. Günther's "Zoölogical Record."* I am also indebted to Dr. Elliott Coues for the addition of the titles of a considerable number of articles to the proof-sheets, which he has had the kindness to carefully examine. The few titles enclosed in brackets indicate those papers I have not myself consulted.

In aiming at brevity I may have excluded from the list a few papers that might well have been added. Usually papers mentioning less than half a dozen species have been excluded, including announcements of the capture of species at localities beyond their usual range. To cite all such notices would nearly double the length of the list, without materially adding to its value, at least for general purposes. Papers in which new species were described are frequently omitted where the geographical data given in them have been subsequently incorporated in other papers published by the same author.

* The volume for 1869 I regret to find has not yet appeared.

The geographical arrangement of the papers serves to show at a glance what portions of the continent are tolerably well known, so far as regards the birds occurring there, as well as to indicate the considerable areas that are still almost unknown, and the amount of information possessed respecting the regions partially known. In order to indicate to some extent the character of the papers mentioned, the number of species given in each is usually stated, as well as the number of pages the papers embrace.

Occasionally valuable notes on the distribution of our birds, and sometimes nearly complete local lists, are to be found in the various agricultural periodicals, in the transactions or reports of agricultural societies, and in the various State agricultural reports. Although a number of such have been entered in the following list, others may have escaped notice; and information of such omissions, or of the omission of any local list, would be thankfully received by the writer.

GENERAL WORKS.

- AUDUBON, J. J. Facts and Observations connected with the permanent Residence of Swallows in the United States. *Ann. N. Y. Lyc. Nat. Hist.*, Vol. I, pp. 166–168, 1824.
- AUDUBON, J. J. Ornithological Biography. 5 vols. 8vo. Edinburgh, 1831–1839.
- AUDUBON, J. J. A Synopsis of the Birds of North America. 8vo. Edinburgh, 1839.
- AUDUBON, J. J. Birds of America. 7 vols. 8vo. Philadelphia, 1840–1844. (506 species.)
- BACHMAN, JOHN. On the Migration of the Birds of North America. *Am. Journ. Sci. and Arts*, 1st Ser., Vol. XXX, pp. 81–100, 1836.
- BAIRD, S. F. Review of American Birds, in the Museum of the Smithsonian Institution, Part I, North and Middle America. 8vo. Washington, pp. 1–176, 1864; pp. 177–320, 1865; pp. 321–450, 1866.
- BAIRD, S. F. The Distribution and Migration of North American Birds. *Am. Journ. Sci. and Arts*, 2d Ser., Vol. XLI, pp. 78–90, 184–192, 337–347, 1866.
- BAIRD, S. F., CASSIN, JOHN, and LAWRENCE, G. N. Birds of North America: a Report on the Birds collected during the Explorations and Surveys for a Railroad Route from the Mississippi River to the Pacific Ocean. *Pacific R. R. Rep. of Expl. and Surv.*, Vol. IX, 4to, 1858. (Republished under the title of "Birds of North America," with an Atlas of Plates. — 738 species, 22 of them extralimital)

- BANNISTER, H. M. A sketch of the Classification of the American *Anseres*. Proc. Acad. Nat. Sci. Philad., 1870, pp. 130-132. (12 North American species.)
- BONAPARTE, C. L. Additions to the Ornithology of the United States. Journ. Acad. Nat. Sci. Philad., Vol. V, pp. 28-31, 1825. (7 species.)
- BONAPARTE, C. L. Specchio comparativo della ornitologia di Roma e Filadelfia. 8vo. Pisa, 1827. Supplement to the same, 1832.
- BONAPARTE, C. L. Saggio di una distribuzione metodica degli Animali vertebrati etc. 8vo. Roma, 1831-1832; Isis, 1832, pp. 283-319; 1833, pp. 1183-1230.
- BONAPARTE, C. L. A Geographical and Comparative List of the Birds of Europe and North America. 8vo, London, 1838. (471 American species.) See a review of the same in Ann. & Mag. Nat. Hist., Vol. I, p. 318, 1839.
- BONAPARTE, C. L. American Ornithology, or the Natural History of the Birds inhabiting the United States, not given by Wilson. 4 vols. 4to. New York, 1825-1833.
- BONAPARTE, C. L., and SCHLEGEL, H. Monographie des Loxiens. 4to. Leiden et Düsseldorf, 1850. (38 species.)
- PRANDT, JOHANN FRIEDRICH. Rapport sur une Monographie de la famille des Alcadées. Bull. Sc. Acad. Imp. St. Petersb., Tome II, pp. 344-349, 1837.
- BRANDT, JOHANN FRIEDRICH. Ergänzungen und Berichtigungen zur Naturgeschichte der Familie der Alciden. Bull. Sc. Acad. Imp. St. Petersb., Tome VII, pp. 199-256, 1869. (21 species.)
- BRANDT, JOHANN FRIEDRICH. Bemerkungen über die Classification, die Entwicklungsstufen und die Verbreitung der Alciden. Bull. Sc. Acad. Imp. St. Petersb., Tome VII, pp. 256-268, 1869.
- BREWER, THOMAS M. North American Oölogy, Part I, *Raptores* and *Fissirostres*, June, 1857. Smithsonian Contributions, Vol. XI.
- BREWER, THOMAS M. A List of Birds common to Europe and America, with others not identical but confounded together from close resemblance. Proc. Bost. Soc. Nat. Hist., Vol. IV, pp. 324-328, 1853. (115 species.)
- BREWER, THOMAS M. A Synopsis of the Birds of North America. Published as an Appendix to Brewer's edition of Jardine's Wilson's Ornithology, pp. 682-746, 1840. (491 species.)
- BRYANT, HENRY. Remarks on the Variations of Plumage of *Buteo borealis* auct. and *Buteo Harlani* Aud. Proc. Bost. Soc. Nat. Hist., Vol. VIII, pp. 107-119, 1861.
- BRYANT, HENRY. Monograph of the Genus *Cctarractes* Moehring. Proc. Bost. Soc. Nat. Hist., Vol. VIII, pp. 134-144, 1861. (4 species.)
- CABOT, S., Jr. On the Distribution of *Tetrao cupido*, and of other Grouse in New England. Proc. Bost. Soc. Nat. Hist., Vol. V, p. 154, 1855.

- CASSIN JOHN. Illustrations of the Birds of California, Texas, Oregon, British and Russian America, etc. 8vo. Philadelphia, 1853-1856. (Contains also Synopses of the *Parinæ*, *Raptores*, and *Fissirostres* of North America).
- CASSIN, JOHN. Notes on North American *Falconidæ*, etc. Proc. Acad. Nat. Sci. Philad., 1855, pp. 277-284. (25 species.)
- CASSIN, JOHN. Ornithology of the United States and British and Russian America. United States Mag., Vol. III, pp. 18, 109, 205, 481, 1856; Vol. IV, p. 10, 1857. (*Raptores* only.)
- CASSIN, JOHN. Studies of the *Icteridæ*. Proc. Acad. Nat. Sci. Philad., 1866, pp. 10-25, 403-417; 1867, pp. 45-74.
- CASSIN, JOHN. Notes on the *Picidæ*, etc. Proc. Acad. Nat. Sci. Philad., 1863, pp. 194-204, 322-328.
- CASSIN, JOHN. Notes of an Examination of the Birds of the Subfamily *Cærebinae*. Proc. Acad. Nat. Sci. Philad., 1864, pp. 265-275.
- COPE, E. D. The Birds of Palestine and Panama compared. Am. Nat., Vol. II, pp. 351-359, 1868.
- COUES, ELLIOTT. A Monograph of the *Alcidæ*. Proc. Acad. Nat. Sci. Philad., 1868, pp. 2-81. (33 species.)
- COUES, ELLIOTT. A Critical Review of the Family *Procellariidæ*. Proc. Acad. Nat. Sci. Philad., 1864, pp. 72-91, 116-144; 1866, pp. 25-33, 134-197. (92 species.)
- COUES, ELLIOTT. A Critical Review of the Sub-family *Lestridinæ*. Proc. Acad. Nat. Sci. Philad., 1863, pp. 121-138. (7 species.)
- COUES, ELLIOTT. Review of the Gulls of North America, etc. Proc. Acad. Nat. Sci. Philad., 1862, pp. 291-312. (25 species.)
- COUES, ELLIOTT. Synopsis of the North American Forms of *Colymbidæ* and *Podicipidæ*. Proc. Acad. Nat. Sci. Philad., 1862, pp. 226-233.
- COUES, ELLIOTT. A Revision of the Terns of North America, etc. Proc. Acad. Nat. Sci. Philad., 1862, pp. 535-559. (14 species.)
- COUES, ELLIOTT. A Monograph of the *Tringæ* of North America. Proc. Acad. Nat. Sci. Philad., 1861, pp. 170-205. (12 species.)
- COUES, ELLIOTT. On Variation in the Genus *Ægiothus*. Proc. Acad. Nat. Sci. Philad., 1869, pp. 180-189. (Special reference to the geographical races.)
- [COUES, ELLIOTT. Key to North American Birds (in press). 1 vol. imp. 8vo. Salem.]
- DRESSER, H. E., and SHARPE, R. B. Notes on *Lanius excubitor* and its Allies. Proc. Zool. Soc. Lond., 1870, pp. 590-600. (9 species.)
- ELLIOT, D. G. Remarks on the Species composing the Genus *Pediæcetes*. Proc. Acad. Nat. Sci. Philad., 1862, pp. 402-404.
- ELLIOT, D. G. Monograph of the *Tetraonidæ*. Folio.

- ELLIOT, D. G. The New and heretofore Unfigured Birds of North America. 2 vols. folio. New York, 1869.
- ELLIOT, D. G. A Monograph of the Genus *Pelecanus*. Proc. Zool. Soc. Lond., 1869, pp. 571–591. (9 species.)
- [FINSCH, OTTO. Die Papageien, monographische bearbeitet. 2 vols. royal 8vo. Leiden, 1867–1868.]
- GAMBEL, WM. Contributions to American Ornithology. Proc. Acad. Nat. Sci. Philad., Vol. IV, pp. 126–129, 1848. (Remarks on 6 species.)
- GOULD, JOHN. Monograph of the *Trochilidæ*, or Humming-Birds. Folio, 1850, *et seq.*
- HEERMANN, A. L. Additions to North American Ornithology. Proc. Acad. Nat. Sci. Philad., Vol. VII, pp. 177–180, 1854. (5 species.)
- HERBERT, WILLIAM HENRY. Frank Forester's Field Sports of the United States and British Provinces of North America. 2 vols. 8vo. New York, 1848. (Contains many valuable observations on the game birds of North America.)
- HUXLEY, T. H. On the Classification and Distribution of the *Alectormorphæ* and *Heteromorphæ*. Proc. Zool. Soc. Lond., 1868, pp. 294–319. With a map.
- LAWRENCE, G. N. Additions to the Ornithology of the United States. Ann. N. Y. Lyc. Nat. Hist., Vol. V, pp. 112–124, 1851. (22 species.)
- LESSON, R. P. Distribution géographique de quelques Oiseaux marins. Ann. Sci. Nat., Tome VI, pp. 88–102, 1825.
- LESSON, R. P. Essai sur la dispersion des Oiseaux sur la surface du globe. Ferrus. Bull. Sc. Nat., Tome XXVII, pp. 92–100, 1831.
- LEWIS, E. J. American Sportsman. 8vo. Philadelphia, 1863. (Contains valuable information respecting the *Rasores*, *Grallæ*, and *Anatidæ*.)
- MALHERBE, A. Monographie des Picidées. 2 vols. folio, avec 125 pl. coloriées, Metz, 1859–1862.
- [MÜLLER, J. W. VON. Reisen in den Vereinigten Staaten, Canada und Mexico. 3 vols. royal 8vo. Leipzig, 1864–1865. Vol. III, pp. 551–594. (Gives a list of 611 species of North American birds.)]
- NUTTALL, THOMAS. A Manual of the Ornithology of the United States and Canada. 2 vols. 12mo. Boston. Vol. I, Land Birds, 1832; 2d ed., with additions, 1840; Vol. II, Water Birds, 1834.
- ORD, GEORGE. List of North American Birds, Guthrie's Geog., 2d Am. ed., 2 vols. 8vo. Philad., 1815.
- PENNANT, THOMAS. Arctic Zoölogy. 3 vols. 4to, 1784–1787; 2d ed., 2 vols. 4to, 1792.
- RIDGWAY, ROBERT. Notices of certain obscurely known Species of American Birds [of the Genera *Turdus* (*Hylocichla*), *Pyrrhula* and *Quiscalus*.] Proc. Acad. Nat. Sci. Philad., 1869, pp. 125–135.

- RIDGWAY, ROBERT. A New Classification of the North American *Falconidæ*. Proc. Acad. Nat. Sci. Philad., 1870, pp. 138 – 150.
- ROOSEVELT, R. B. The Game Birds of the Northern States of America. 12mo. New York, 1866, pp. 338.
- SALVIN, O. On the Genus *Cinclus*. Ibis, 2d Ser., Vol. III, pp. 109 – 122, 1867. (Contains remarks on the geographical distribution of 13 species.)
- SCHMARDA, L. K. Die geographische Verbreitung der Thiere, 3 Bd. Wien, 1853.
- SCLATER, P. L. Catalogue of a Collection of American Birds, belonging to Philip Lutley Sclater, M. A., etc., etc. 8vo. London, 1862.
- SCLATER, P. L. On the general Geographical Distribution of the Members of the Class Aves. Journ. of Proc. of Linnæan Society of London, Vol. II, Zoölogy, pp. 130 – 145, 1858.
- SCLATER, P. L. Notes on Krueper's Nuthatch and on the other known Species of the Genus *Sitta*. Ibis, 1865.
- SCLATER, P. L. Notes on the Genera and Species of *Cypselidæ*. Proc. Zoöl. Soc. Lond., 1865, pp. 593 – 617.
- SCLATER, P. L. Notes upon the American *Caprimulgidæ*. Proc. Zoöl. Soc. Lond., 1866, pp. 123 – 145.
- SCLATER, P. L., and SALVIN, O. Notes on the Species of the Genus *Asturina*. Proc. Zoöl. Soc. Lond., 1869, pp. 129 – 134. (7 species.)
- SCLATER, P. L., and SALVIN, O. Synopsis of the American Rails (*Rallidæ*). Proc. Zoöl. Soc. Lond., 1868, pp. 442 – 470. (48 species.)
- SCLATER, P. L., and SALVIN, O. Exotic Ornithology. 4to. London, 1866 *et seq.* (Publishing in parts.)
- SELYS-LONGCHAMPS, EDM. DE. Sur les Oiseaux américains admis dans la Faune européenne. Mém. de la Soc. R. de Liège, Vol. IV, pp. 35 – 50, 1849.
- [SHARPE, R. B. A Monograph of the *Alcedinidæ*, or Family of Kingfishers. Imp. 8vo. London, 1870. Introduction published separately. (4to, London, 1871.)]
- SUNDEVALL, C. J. Conspectus Avium *Picinarum*. 1 vol. 8vo. pp. 116. Stockholm, 1866. (254 species.)
- SUNDEVALL, C. J. Conspectus Generis *Dendræcæ*. Ofvers. Kongl. Vetensk. Acad. Forhandl., 1869, pp. 605 – 618. (30 species.)
- TAYLOR, G. C. Birds observed during two Voyages across the North Atlantic. Ibis, 1869, p. 388.
- VERRILL, A. E. On the supposed Eastward Migration of the Cliff Swallow (*Hirundo lunifrons* Say). Proc. Bost. Soc. Nat. Hist., Vol. IX, pp. 276 – 278, 1863.
- VERRILL, A. E. Report of some Investigations upon the Geographical Distribution of North American Birds, made with Reference to the Physical Causes that determine their Limit in Latitude. Proc. Bost. Soc. Nat. Hist., Vol. X, pp. 259 – 262, 1866.

- WALLACE, A. R. Letter concerning the Geographical Distribution of Birds. *Ibis*, Vol. I, pp. 449-454, 1859.
- WIED, MAX PRINZ VON. Verzeichniss der Vögel welche auf seiner Reise in Nord-Amerika beobachtet wurden. *Cabanis's Journ. für Ornithol.*, VI Jahrgang, pp. 1-29, 97-124, 177-204, 257-284, 337-354, 417-444, 1858; VII Jahrgang, pp. 81-96, 161-180, 241-260, 1859.
- WILSON, ALEXANDER. *American Ornithology, or the Natural History of the Birds of the United States*. 9 vols. 4to. Philadelphia, 1808-1814. (283 species.)

GREENLAND.

- FABRICIUS, O. *Fauna Grœnlandiæ, etc.* 8vo. Hafniæ et Lipsiæ, 1780. (54 species.)
- HOLBÖLL, CARL. Ornithologiske Bidrag til den grønlandske Fauna. *Krøyer's naturhist. Tidskr.*, Bd. IV, pp. 361-457, 1843; *Isis*, 1845, pp. 739-792. (88 species.)
- REINHARDT, J. List of Birds hitherto observed in Greenland. *Ibis*, 1st Ser., Vol. III, pp. 1-19, 1861. (118 species.)
- [RICHARDSON, J. Birds and Mammalia collected during the last Arctic Land Expedition under Sir John Franklin. *Proc. of the Com. of Sci. and Corres. of Zool. Soc. Lond.*, Vol. I, p. 132, 1831.]
- SABINE, EDWARD. A Memoir on the Birds of Greenland; with Descriptions and Notes on the Species observed in the late Voyage of Discovery in Davis's Straits and Baffin's Bay. *Transact. Linn. Soc. of London*, Vol. XII, pp. 527-559, 1819. (54 species.)

BRITISH NORTH AMERICA EAST OF THE ROCKY MOUNTAINS.

Labrador and Anticosti.

- COUES, ELLIOTT. Notes on the Ornithology of Labrador. *Proc. Acad. Nat. Sci. Philad.*, 1861, pp. 215-257. (83 species.)
- VERREAUX, JULES. Der in Labrador befindlichen Landsäugethiere und Vögel. *Bull. Akad. München*, 1844, pp. 122-126, 129-133.
- VERRILL, A. E. Notes on the Natural History of Anticosti. *Proc. Bost. Soc. Nat. Hist.*, Vol. IX, pp. 137-145. (Contains a list of 62 species of birds.)
- WEIS, SAMUEL. List of Vertebrates observed at Okak, Labrador, by Rev. Samuel Weis, with Annotations by Dr. A. S. Packard, Jr. *Proc. Bost. Soc. Nat. Hist.*, Vol. X, pp. 264-277, 1866. (Birds, pp. 267-269; 85 species.)

Interior of British North America.

- [ARMSTRONG, A. A personal Narrative of the Discovery of the Northwest Passage, etc. 8vo. London, 1857. (31 species from the Arctic Coast.)]

- BARNSTON, GEORGE. Recollections of the Swans and Geese of Hudson's Bay. *Can. Nat. and Geol.*, Vol. VI, pp. 337 - 344, 1861.
- [BLAKISTON, THOMAS. [Ornithological] Scraps from the Far West. *Zoölogist*, 1859, pp. 6318 - 6325, 6373 - 6376.]
- BLAKISTON, CAPT. On Birds collected and observed in the Interior of British North America. Part I, *Ibis*, 1st Ser., Vol. III, pp. 314 - 320, 1861; Part II, *Ibid.*, Vol. IV, pp. 3 - 10, 1862.
- BLAKISTON, CAPT. Notes on the Interior of British North America. *Ibis*, 1st Ser., Vol. V, pp. 39 - 87, 1863. (Devoted principally to remarks on the birds.)
- CASSIN, JOHN. Remarks on Birds from the Arctic Regions, presented by Dr. Kane. *Proc. Acad. Nat. Sci. Philad.*, 1852, p. 107.
- EDWARDS, GEORGE. Natural History of Birds etc. 4 vols. 4to. London, 1743 - 1751. (Contains the first descriptions of 38 species of birds from Hudson's Bay.)
- GUNN, DONALD. Notes of an Egging Expedition to Shoal Lake, west of Lake Winnipeg. *Smithsonian Report*, 1867, pp. 427 - 432.
- FORSTER, J. R. An Account of the Birds sent from Hudson's Bay, etc. *Phil. Trans.*, Vol. LXII, pp. 382 - 422, 1772. (58 species.)
- LEACH, W. E. Zoölogical Appendix to Ross's Voyage to Baffin's Bay. Birds, App. No. 2, pp. xlviii - lx, 1819. (20 species.)
- LORD, J. K. List of Birds collected and presented by the British North American Boundary Commission to the Royal Artillery Institution. *Proc. Roy. Artill. Instit.*, 1864, pp. 110 - 126. (87 species.)
- LORD, J. K. Catalogue of the Birds, Nests, and Eggs collected by him when Naturalist to the North American Boundary Commission. *Proc. Roy. Art. Inst.*, 1865, pp. 337 - 339.
- MURRAY, ANDREW. Contributions to the Natural History of the Hudson's Bay Company's Territory. Part III, Birds. *Edinburgh New Phil. Journ.*, 2d Ser., Vol. IX, pp. 221 - 231, 1857. (82 species.)
- RICHARDSON, J., and SWAINSON, W. Fauna Boreali-Americana. Vol. II, Birds. 4to, 1831. (238 species.)
- RICHARDSON, JOHN. Geographical Distribution of some American Birds. *Ann. Nat. Hist.*, Vol. XI, pp. 484, 1843. (A list of 12 species collected at Fort Simpson, Mackenzie's River.)
- ROSS, BERNARD R. Mammals and Birds of the Arctic Regions. *Edinb. Phil. Journ.*, 2d Ser., Vol. XIII, 1861, pp. 161 - 164. (A nominal list of 94 species.)
- ROSS, BERNARD R. List of Species of Mammals and Birds collected in Mackenzie's River District during 1860 - 1861 (June, 1860, to April, 1861). *Can. Nat. and Geol.*, Vol. VI, pp. 441 - 444, 1862. (80 species of birds mentioned.)

- ROSS, BERNARD R. List of Mammals, Birds and Eggs, observed in the Mackenzie's River District, with Notices. *Nat. Hist. Rev.*, Vol. II, pp. 276 - 290, 1862. (192 species.)
- SABINE, EDWARD. Supplement to the Appendix to Captain Parry's First Voyage. London, 4to, 1824. (73 species.)
- SABINE, JOSEPH. Birds of Sir John Franklin's [First] Journey. Franklin's Journey to the Shore, of the Polar Sea. 4to. London, 1823. Appendix, pp. 669 - 703. (94 species.)

Nova Scotia, New Brunswick, and Newfoundland.

- AMBROSE, JOHN. Observations on the Sea-Birds frequenting St. Margaret's Bay, N. S. *Proc. Nov. Scot. Inst. Nat. Sci.*, Vol. II, Part II, pp. 51 - 59. (19 species.)
- BREWER, THOMAS M. A few Ornithological Facts gathered in a hasty Trip through Portions of New Brunswick and Nova Scotia, in June, 1850. *Journ. Bost. Soc. Nat. Hist.*, Vol. VI, pp. 297 - 308, 1852. (7 species.)
- BRYANT, HENRY. List of Birds observed at Grand Menan and at Yarmouth, N. S., from June 16 to July 8, 1856. *Proc. Bost. Soc. Nat. Hist.*, Vol. VI, pp. 114 - 123, 1857. (55 species.)
- BRYANT, HENRY. Remarks on some Birds that breed in the Gulf of St. Lawrence. *Proc. Bost. Soc. Nat. Hist.*, Vol. VIII, pp. 65 - 75, 1860; *Can. Nat. and Geol.*, Vol. VI, 1861, pp. 255 - 267. (14 species.)
- DOWNES, A. On the Land Birds of Nova Scotia. *Proc. Nov. Scot. Inst. Nat. Sci.*, Vol. II, pp. 38 - 51, 1865. (61 species.)
- JONES, J. M. On some of the Rarer Birds of Nova Scotia. *Proc. Nov. Scot. Inst. Nat. Sci.*, Vol. II, Part II, pp. 70 - 73, 1868.
- REEKS, HENRY. Notes on the Birds of Newfoundland. *Can. Nat.*, Vol. V, pp. 38 - 47, 151 - 159, 1860.
- WILLIS, J. R. List of the Birds of Nova Scotia, compiled from Notes by Lieut. Blakiston and Lieut. Bland. *Smithsonian Report*, 1858, pp. 280 - 286. (207 species.)

The Canadas.

- BELL, ROBERT. Catalogue of Birds collected around Lakes Superior and Huron in 1860. *Can. Nat. and Geol.*, Vol. VI, pp. 270 - 275, 1861. (77 species.)
- CABOT, J. E. Report of the Birds collected and observed at Lake Superior. Agassiz's Lake Superior, pp. 383 - 385, 1850. (69 species.)
- COUPER, WM. List of the Birds found in the Vicinity of Quebec, Lower Canada. *Samuels's Birds of New England*, pp. 80, 81, 368 - 372, 396, 477 - 478, 571 - 573, 1868. (154 species.)
- D'URBAN, W. S. M. Notes on the Land Birds observed around Mon-

- trear during the Winter of 1856-57. Can. Nat. and Geol., Vol. II, pp. 138-145, 1857. (18 species.)
- [HADFIELD, HENRY. Birds of Canada observed near Kingston during the Spring of 1858. Zoölogist, 1859, pp. 6701-6709, 6744-6752, 6751-6787.]
- HALL, ARCHIBALD. On the Mammals and Birds of the District of Montreal. Part II, Birds. Canadian Nat. and Geol., Vol. VII, pp. 44-54, 1862. (199 species.)
- KING, W. ROSS. The Sportsman and Naturalist in Canada; or, Notes on the Natural History of the Game, Game Birds, and Fish of that Country. Imp. 8vo, London, 1866.
- LE MOINE, J. M. Notes on Land and Sea Birds observed around Quebec. Can. Nat. and Geol., Vol. IV, pp. 411-414, 1859.
- LE MOINE, J. M. Ornithologie du Canada. Oiseaux de Proie et Palmipèdes. 8vo, pp. 96, Quebec, 1860.
- McILWRAITH, THOMAS. List of Birds observed near Hamilton, Canada West. Proc. Essex Institute, Vol. V, pp. 79-96, 1866. (241 species.)
- VENNOR, H. G. A short Review of the *Sylviadæ*, or Wood-Warblers, found in the Vicinity of Montreal. Can. Nat. and Geol., VI, pp. 349-362, 1861.
- VENNOR, H. G., JR. Notes on Birds wintering in and around Montreal. Can. Nat. and Geol., Vol. V, pp. 425-430, 1860. (27 species.)
- W., J. F. Notes on Canadian Birds. Can. Nat. and Geol., Vol. V, pp. 230, 231, 1870. (The capture of 9 rare species near Quebec mentioned.)

NEW ENGLAND STATES.

General.

- BREWER, T. M. Seaside Ornithology. Am. Nat., Vol. III, pp. 225-235, 1869.
- COUES, ELLIOTT. A List of the Birds of New England. Proc. Essex Institute, Vol. V, pp. 249-314, 1868. (335 species.)
- SAMUELS, E. A. Birds of New England and Adjacent States, 5th ed. (of "Ornithology and Oölogy of New England"). Boston, 1870.

Maine.

- BOARDMAN, G. A. Catalogue of the Birds found in the Vicinity of Calais, Maine, and about the Islands at the Mouth of the Bay of Fundy. - Proc. Bost. Soc. Nat. Hist., Vol. IX, pp. 122-132, 1862. (231 species.)
- CABOT, S., JR. Notes of the Breeding of *Mergus serrator*, *M. cucullatus*, *Anas sponsa*, *A. obscura*, and *Fuligula clangula*, on the Sources of the Androscoggin River, Maine. Proc. Bost. Soc. Nat. Hist., Vol. II, p. 55.
- HAMLIN, C. E. Catalogue of Birds found in the Vicinity of Waterville, Kennebec County, Maine. Rep. of Sec'y Maine Board of Agric., 1865, pp. 168-173. (135 species.)

- HOLMES, EZEKIEL. Birds of Maine. Agr. Maine, 6th Ann. Rep., 1861. pp. 113 - 122. (A nominal list of 193 species.) Addenda to the same. 2d Ann. Rep. Nat. Hist. and Geol. of Maine, 1862, p. 118. (13 species.)
- VERRILL, A. E. Catalogue of Birds found at Norway, Oxford County, Maine. Proc. Essex Institute, Vol. III, pp. 136 - 158, 1863. (159 species.)
- VERRILL, A. E. Additions to the Catalogue of Birds found in the Vicinity of Calais, Maine, and about the Bay of Fundy. Proc. Bost. Soc. Nat. Hist., Vol. IX, pp. 233, 234, 1863. (12 species.)

Vermont.

- THOMPSON, ZADOC. Birds of Vermont. History of Vermont, Natural, Civil, and Statistical, 8vo. Burlington, 1842. Birds, pp. 56 - 112.

Massachusetts.

- ALLEN, J. A. Catalogue of the Birds found at Springfield, Massachusetts, with a List of those found in the State not yet observed at Springfield, etc. Proc. Essex Institute, Vol. IV, pp. 48 - 98, 1864. (296 species.)
- ALLEN, J. A. Winter Notes of an Ornithologist. Amer. Nat., Vol. I, pp. 38 - 48, 1867. (On the Winter Birds of Massachusetts.)
- ALLEN, J. A. The Birds of Spring. Amer. Nat., Vol. I, pp. 141 - 144, 1867.
- ALLEN, J. A. Notes on some of the Rarer Birds of Massachusetts. Amer. Nat., Vol. III, pp. 505 - 519, 568 - 585, 631 - 648, 1869 - 70. (92 species.)
- ANONYMOUS. Notes on the Ducks found on the Coast of Massachusetts in Winter. Amer. Nat., Vol. IV, p. 49. (8 species.)
- BREWER, THOMAS M. Some Additions to the Catalogue of the Birds of Massachusetts in Prof. Hitchcock's Report, etc. Jour. Bost. Soc. Nat. Hist., Vol. I, pp. 435 - 439, 1837. (45 species.)
- BREWER, T. M. Observations on the Appearance of the Cliff Swallow (*Hirundo lunifrons*), giving Data of its Appearance in New England. Proc. Bost. Soc. Nat. Hist., Vol. IV, p. 270, 1852.
- BRYANT, H. Notice of *Mareca Penelope* and *Querquedula crecca* taken in Eastern Massachusetts. Proc. Bost. Soc. Nat. Hist., Vol. V, p. 195, 1855.
- EMMONS, E. Catalogue of the Birds of Massachusetts. Hitchcock's Report on the Geology of Mass., Amherst, 1833. (165 species.)
- MAYNARD, C. J. Catalogue of the Birds of Eastern Massachusetts. Naturalist's Guide, Part II, Boston, 1870. (299 species.)
- NUTTALL, THOMAS. Remarks and Inquiries concerning the Birds of Massachusetts. Mem. Amer. Acad. Arts and Sciences, 2d Ser., Vol. I, pp. 91 - 106, 1833. (45 species remarked upon.)
- PEABODY, W. B. O. A Report on the Ornithology of Massachusetts. Rep. on the Fishes, Reptiles, and Birds of Mass., 8vo, 1839, pp. 256 - 404;

Ibid., Jour. Bost. Soc. Nat. Hist., Vol. III, pp. 65 – 246, 1840. (284 species).

PUTNAM, FRED. W. Catalogue of the Birds of Essex County, Mass. Proc. Essex Institute, Vol. I, pp. 201 – 231, 1856. (245 species.)

SAMUELS, E. A. Descriptive Catalogue of the Birds of Massachusetts. Agr. of Mass., Sec'y's Rep., 1863, App. pp. xvii – xxix, 1864. (267 species.)

Connecticut.

LINSLEY, J. H. A Catalogue of the Birds of Connecticut, etc. Am. Jour. of Sci. and Arts, 1st Ser., Vol. XLIV, pp. 249 – 273, 1843. (302 species.)

WOOD, WM. Rapacious Birds of Connecticut. *Hartford Times*, March 14 to August 8, 1861.

MIDDLE ATLANTIC STATES.

New York.

DE KAY, JAMES E. Zoölogy of New York, or the New York Fauna. 4to. Albany, 1844. Part II, Birds. (317 species.)

GIRAUD, J. P. The Birds of Long Island, 8vo. New York, 1843 – 1844. (276 species.)

HOPKINS, WM. List of Sea-Birds obtained at Auburn, New York. Proc. Bost. Soc. Nat. Hist., Vol. V, p. 13, 1854. (8 species.)

LAWRENCE, GEO. N. Catalogue of Birds observed on New York, Long and Staten Islands. Ann. New York Lyc. Nat. Hist., Vol. VIII, pp. 279 – 300, 1867. (327 species.)

New Jersey.

ABBOTT, C. C. Catalogue of the Birds of New Jersey. Geol. of New Jersey, Appendix, pp. 761 – 798, 1868. (301 species.)

ABBOTT, C. C. Notes on certain Inland Birds of New Jersey. Amer. Nat., Vol. IV, pp. 536 – 550, 1870. (26 species.)

TURNBULL, WM. P. The Birds of East Pennsylvania and New Jersey. 8vo pamphlet. Philadelphia, 1869. (343 species.)

Pennsylvania.

BAIRD, WM. M. and S. F. List of the Birds found in the Vicinity of Carlisle, Cumberland County, Penn. Am. Jour. Sci. and Arts, 1st Ser., Vol. XLVI, pp. 261 – 273, 1844. (201 species.)

BAIRD, S. F. Catalogue of the Birds found in the Neighborhood of Carlisle, Cumberland County, Penn. Literary Rec. and Journ. Linn. Ass. Pennsylv. College, Vol. I, p. 249, 1845.

BARNARD, VINCENT. A Catalogue of the Birds of Chester County, Pennsylvania, etc. Smiths. Rep., 1860, pp. 434 – 438. (191 species.)

BARTON, BENJ. S. Fragments of the Natural History of Pennsylvania.

- Folio. Philadelphia, 1799. (Contains a list of the birds of Pennsylvania, with dates of migration.)
- HALDEMANN, S. S. Facts in Ornithology. Proc. Acad. Nat. Sci. Philad., Vol. I, p. 54, 1841. (*Falco peregrinus* stated to breed "on the cliffs along the Susquehanna.")
- TAYLOR, R. C. On the Geology and Natural History of the Northeastern Extremity of the Alleghany Mountain Range in Pennsylvania, U. S. Loudon's Mag. Nat. Hist., Vol. VIII, pp. 539-540. (Contains a nominal list of 56 species of birds.)
- TAYLOR, R. C. A Supplement to the List of Species of Birds occurring at the Northeastern Extremity of the Alleghany Mountains. Loudon's Mag. Nat. Hist., Vol. IX, pp. 72-74, 1836. (36 species.)
- See also TURNBULL, under *New Jersey*.

SOUTH ATLANTIC STATES.

General.

- BARTRAM, JOHN. Travels through North and South Carolina, Georgia, East and West Florida, etc., 8vo. Philadelphia, 1791. Contains a list of the birds of the United States, pp. 289-296. (216 species.)
- CATESBY, MARK. The Natural History of Carolina, Florida, and the Bahama Islands. 2 vols. folio, 3d ed. London, 1771.

Virginia and the District of Columbia.

- [CLAYTON, JOHN. On the Birds of Virginia. Phil. Trans., Vol. XVII, p. 988, 1693.]
- COUES, ELLIOTT, and PRENTISS, D. WEBSTER. List of the Birds ascertained to inhabit the District of Columbia, with the Times of Arrival and Departure of such as are Non-residents, etc. Smithsonian Report, 1861, pp. 399-421. (226 species.)

North Carolina.

- COPE, E. D. Observations on the Fauna of the Southern Alleghanies. Amer. Nat., Vol. IV, pp. 392-401, 1870. (Madison County, North Carolina, and Giles County, East Virginia.)
- [COUES, ELLIOTT. Notes on the Natural History of Fort Macon, N. C. Birds, Part I. Proc. Acad. Nat. Sci. Philad., 1871 (in press).]

South Carolina.

- BACHMAN, JOHN. On the Migration of the Birds of North America. Am. Journ. Sci. and Arts, 1st Ser., Vol. XXX, pp. 81-100, 1836. (Contains numerous observations on the birds of South Carolina.)
- BURNETT, W. I. Notes on the Fauna of the Pine Barrens of Upper South Carolina. Proc. Bost. Soc. Nat. Hist., Vol. IV, pp. 115-118, 1851. (69 species of birds mentioned.)

COUES, ELLIOTT. Synopsis of the Birds of South Carolina. Proc. Bost. Soc. Nat. Hist., Vol. XII, pp. 104 - 127, 1868. (294 species.)

GIBBS, LEWIS R. Catalogue of the Birds of South Carolina. Tuomey's Geology of South Carolina, Appendix, pp. iii - viii, 1848. (271 species.)

Georgia.

GERHARDT, ALEX. Über die Lebensweise der Vögel Nord-Amerikas (Georgia). Naumannia, 1855, pp. 380 - 397, 458 - 469; 1856, pp. 1 - 18.

GERHARDT, ALEX. Verzeichniss der Vögel des Staats Georgia, nach White. Naumannia, 1855, pp. 382 - 383. (A nominal list of 270 species.)

Florida.

ALLEN, J. A. List of the Winter Birds of East Florida, with Annotations. Bulletin Mus. Comp. Zoöl., Vol. II, pp. 250 - 368, 1871. (183 species.)

BRYANT, H. Observations on some of the Birds of East Florida. Proc. Bost. Soc. Nat. Hist., Vol. VII, pp. 5 - 21, 1859. (26 species.)

TAYLOR, GEORGE CAVENDISH. Five Weeks in the Peninsula of Florida during the Spring of 1861, with Notes on the Birds observed there. Ibis, 1st Ser., Vol. IV, pp. 127, 197, 1862.

GULF STATES.

Mississippi.

WAILES, B. L. C. Birds of Mississippi. Rep. on Agr. and Geol. of Mississippi, pp. 317 - 327, 1854. (91 species; mainly a nominal list.)

CENTRAL STATES

Ohio.

KIRKPATRICK, JOHN. The Rapacious Birds of Ohio. *Ohio Farmer* for 1858 - 59; Ohio Agricult. Rep., 1858, pp. 340 - 383, 1859.

KIRTLAND, J. P. Fragments of Natural History. II, Ornithology. Am. Journ. Sci. and Arts, 1st Ser., Vol. XL, pp. 19 - 24, 1841. (Contains notes on 16 species observed near Cleveland.)

KIRTLAND, J. P. Catalogue of the Birds of Ohio. Second Ann. Report of the Geol. Survey of Ohio, pp. 177 - 187, 1838. (223 species.)

READ, M. C. Catalogue of the Birds of Northern Ohio. Proc. Acad. Nat. Sci. Philad., 1853, pp. 395 - 402. (145 species.)

WHEATON, J. M. Catalogue of the Birds of Ohio. Ohio Agri. Rep., 1860, pp. 359 - 380, 480. (285 species.)

Indiana.

ALLEN, J. A. List of Birds observed in Richmond, Wayne Co., Indiana, in June, 1867. Mem. Bost. Soc. Nat. Hist., Vol. I, pp. 522 - 526, 1868. (72 species.)

HAYMOND, RUFUS. Birds of Southeastern Indiana. Proc. Acad. Nat. Sci. Philad., 1856, pp. 286 – 299. (147 species.)

HAYMOND, RUFUS. Birds of Franklin County, Indiana. Cox's Geol. Survey of Ind., 1869, pp. 209 – 235. (164 species.)

Illinois.

ALLEN, J. A. Notes on Birds observed in Northern Illinois, in June, 1867. Mem. Bost. Soc. Nat. Hist., Vol. I, pp. 502 – 522, 1868. (94 species.)

BRENDEL, F. Vögel der Umgegend Peoria's in Illinois. Giebel's Zeitsch. für Naturk., 1857, p. 420.

KENNICOTT, ROBERT. Catalogue of the Animals of Cook County, Illinois. [Birds.] Transact. Ill. State Agric. Soc., I, 1853 – 54, pp. 580 – 591. (187 species.)

KENNICOTT, R. Notes of the Occurrence of *Plotus anhinga* and *Tantalus loculator* in Southern Illinois. Proc. Bost. Soc. Nat. Hist., Vol. V, p. 391, 1856.

PRATTEN, HENRY. Catalogue of the Birds of Southern Illinois (Wayne and Edwards Counties). Transact. Ill. State Agric. Soc., I, pp. 596 – 609, 1853 – 54. (184 species.)

Michigan.

HUGHES, D. DARWIN. The Game Birds of Michigan. Detroit Free Press, 1870.

KNEELAND, S., JR. On the Birds of Keweenaw Point, Lake Superior. Proc. Bost. Soc. Nat. Hist., Vol. VI, pp. 231 – 240, 1858. (147 species.)

Minnesota.

HEAD, J. F. Some Remarks on the Natural History of the Country about Fort Ripley, Minnesota. Smithsonian Report, 1854, pp. 291 – 293. (60 species of birds mentioned.)

[TRIPPE, T. MARTIN. Notes on the Birds of Minnesota (in press). Proc. Essex Institute, Vol. VI. (138 species.)]

Wisconsin.

BARRY, A. CONSTANTINE. Ornithological Fauna of Wisconsin. Proc. Bost. Soc. Nat. Hist., Vol. V, pp. 1 – 13, 1854. (193 species.)

HOY, P. R. Notes on the Ornithology of Wisconsin. Proc. Acad. Nat. Sci. Philad., 1853, pp. 304 – 313, 381 – 385, 425 – 429. Ibid., Transact. Wisconsin Agric. Soc., 1852, pp. 341 – 364. (283 species.)

Iowa.

ALLEN, J. A. Notes on Birds observed in Western Iowa, July 1 to Sept. 20, 1867. Mem. Bost. Soc. Nat. Hist., I, pp. 491 – 502, 1868. (108 species.)

ALLEN, J. A. Catalogue of the Birds of Iowa. White's Geol. of Iowa, Vol. II, Appendix, pp. 419 – 427, 1870. (283 species.)

Missouri.

- HOY, P. R. Journal of an Exploration of Western Missouri in 1854, etc. Smithsonian Rep., 1864, pp. 431-438. (Includes a nominal list of the birds observed, — 153 species.)

Texas.

- BUTCHER, H. B. List of Birds collected at Laredo, Texas, in 1866 and 1867. Proc. Acad. Nat. Sci. Philad., 1868, pp. 148-150. (89 species.)
- DRESSER, H. E. Notes on the Birds of Southern Texas. Ibis, 2d Ser., Vol. I, pp. 312-330, 466-495; Vol. II, pp. 23-46, 1866.
- MCCALL, GEO. A. Remarks on the Habits, etc., of Birds met with in Western Texas, between San Antonio and the Rio Grande, and in New Mexico, with Descriptions of several Species believed to have been hitherto undescribed. Proc. Acad. Nat. Sci. Philad., 1850-51, pp. 213-224. (68 species.)
- MCCALL, GEO. A. [Ornithological] Facts and Observations from Notes when in Texas. Ann. N. Y. Lyc. Nat. Hist., Vol. VI, pp. 9-14. (18 species.)
- ROEMER, FERDINAND. Texas: mit besonderer Rücksicht auf deutsche Auswanderung und die physischen Verhältnisse des Landes, nach eigener Beobachtung geschildert. 8vo. Bonn, 1849. (Contains a list of birds, — 40 species.)

ROCKY MOUNTAINS AND ADJACENT PLAINS.

General.

- BAIRD, S. F. [Nominal] List of Birds inhabiting America, west of the Mississippi, not described in Audubon's Ornithology. Stansbury's Expl. and Survey of the Valley of the Great Salt Lake, pp. 327-335, 1852. (153 species.)
- BAIRD, S. F. Descriptions of New Birds, collected between Albuquerque, New Mexico, and San Francisco, during the Winter of 1853-54, by Dr. C. B. R. Kennerly and H. B. Möllhausen, etc. Proc. Acad. Nat. Sci. Philad., Vol. VII, p. 118-120, 1854. (8 species.)
- BAIRD, S. F. Report on the Birds collected by the United States and Mexican Boundary Survey, with Notes by J. H. Clark, D. N. Couch, and C. B. R. Kennerly. U. S. and Mex. Bound. Sur., Vol. II, Part II, No. 2, pp. 1-32, 1859. (236 species.)
- BAIRD, S. F. Report on the Birds collected on the Route near the 38th and 39th Parallels, explored by Capt. J. W. Gunnison, and near the 41st Parallel, explored by Lieut. E. G. Beckwith. Pacific R. R. Rep. of Expl. and Surv., Vol. X, Part IV, No. 2, pp. 11-16, 1859. (25 species.)

- COOPER, J. G., and SUCKLEY, G. Report on the Birds collected on the Route near the 47th and 49th Parallels, explored by I. I. Stevens, Governor of Washington Territory. Pacific R. R. Rep. of Expl. and Surveys, Vol. XII, Part II, No. 3, pp. 140-291, 1859. (Republished, "with additions and corrections," in 1860, forming the work entitled "Natural History of Washington Territory.") (212 species.)
- COOPER, J. G., and SUCKLEY, G. List of Birds heretofore reported as found in the Northwestern Part of America, but of which no Specimens have been reported by recent Explorers. Pacific R. R. Rep. of Expl. and Surv., Vol. XII, Part II, Zoölogy, pp. 288-290, 1859. (120 species.)
- COUES, ELLIOTT. Ornithology of a Prairie Journey, and Notes on the Birds of Arizona. Ibis, 1st Ser., Vol. V, April, 1865.
- HAYDEN, F. V. List of the Birds of the Upper Missouri. Geol. and Nat. Hist. of the Upper Missouri, pp. 157-176, 4to. Philadelphia, 1862. (191 species.)
- HEERMANN, A. L. Report on the Birds collected on the Route near the 32d Parallel, from the Rio Grande to the Pimas Villages, explored by Lieut. J. G. Parke. Pacific R. R. Rep. of Expl. and Surveys, Vol. X, Part —, No. 1, pp. 9-20, 1859. (25 species.)
- KENNERLY, C. B. R. Report upon the Birds of the Route near the 35th Parallel, explored by Lieut. A. W. Whipple. Pacific R. R. Rep. of Expl. and Surveys, Vol. X, Part VI, No. 3, pp. 19-35, 1859. (88 species.)
- NEWBERRY, J. S. Report upon the Zoölogy of Lieut. R. S. Williamson's Explorations in Oregon and California. Pacific R. R. Expl. and Surveys, Vol. VI, Part IV, No. 2, chap. ii, Birds, pp. 73-110, 1857. (175 species.)
- SAY, THOMAS. The Ornithological Articles in Long's Expedition (from Pittsburg to the Rocky Mountains). 2 vols. 8vo. Philadelphia, 1823.
- TOWNSEND, JOHN K. List of the Birds inhabiting the Regions of the Rocky Mountains, the Territory of Oregon, and the Northwest Coast of North America. Journ. Acad. Nat. Sci. Philad., 1st Ser., Vol. VIII, pp. 151-158, 1839. (231 species.)
- TOWNSEND, JOHN K. Description of twelve new Species of Birds, chiefly from the Vicinity of the Columbia River, with an Appendix containing a [nominal] List of the Birds of the Columbia River. Journ. Acad. Nat. Sci. Philad., 1st Ser., Vol. VII, pp. 187-193, 1837. (72 species.)
- WOODHOUSE, S. W. Report on the Birds of Capt. L. Sitgreaves's Expedition down the Zuñi and Colorado Rivers. Sitgreaves's Report, pp. 58-105, 1853. (247 species.)

Montana.

- COOPER, J. G. The Fauna of Montana Territory. Part II, Birds. Amer. Nat., Vol. II, pp. 596-600; Vol. III, pp. 31-35, 73-84, 1869. (107 species.)

Utah.

- BAIRD, S. F. Exploration and Survey of the Valley of the Great Salt Lake, etc., by H. Stansbury, T. E., U. S. A. 8vo. Philad., 1852, Appendix C, Birds, pp. 314-324. (31 species.)
- BAIRD, S. F. Birds found at Fort Bridger. Gen. Rep. on Birds of N. Amer., Append., pp. 926, 927; P. R. R. Rep. of Expl. and Surv., Vol. IX, 1858. (104 species.)

Arizona.

- COUES, ELLIOTT. List of Birds collected in Southern Arizona by Dr. E. Palmer, with Remarks. Proc. Acad. Nat. Sci. Philad., 1868, pp. 81-85 (53 species.)
- COUES, ELLIOTT. List of the Birds of Fort Whipple, Arizona, with which are incorporated all other Species ascertained to inhabit the Territory, etc. Proc. Acad. Nat. Sci. Philad., 1866, pp. 39-100. (245 species.) — Republished under the title, "Prodrome of a Work on the Ornithology of Arizona Territory."*

New Mexico

- BAIRD, S. F. On Birds collected in New Mexico by Lieut. Abert. Stansbury's Exped., pp. 325-327. (9 species.)
- HENRY, T. CHARLTON. Notes derived from Observations made on the Birds of New Mexico, during the Years 1853 and 1854. Proc. Acad. Nat. Sci. Philad., 1855, pp. 306-317. (170 species.)
- HENRY, T. CHARLTON. Catalogue of the Birds of New Mexico, as compiled from Notes and Observations made while in that Territory, during a Residence of six Years. Proc. Acad. Nat. Sci. Philad., 1859, pp. 104-109. (198 species.)

PACIFIC COAST.**General.**

- CASSIN, JOHN. Catalogue of Birds collected by the North Pacific Surveying and Exploring Expedition, in Command of Capt. Rogers, U. S. N., etc. Proc. Acad. Nat. Sci. Philad., 1862, pp. 312-328. (Contains many species from the West Coast of North America.)
- PEALE, TITIAN R. Report on the Mammalia and Birds of the U. S. Exploring Expedition under Capt. Wilkes. 4to. Philadelphia, 1848. (Includes many species from the West Coast of North America.)
- VIGORS, N. A. Ornithology of Captain Beechey's Voyage. Zool. of Beechey's Voyage, pp. 13-40, 1839.

* I regret to learn from Dr. Coues that this work will not be published.

Alaska.

- BAIRD, S. F. On Additions to the Bird-Fauna of North America, made by the Scientific Corps of the Russo-American Telegraph Expedition. Transact. Chicago Acad. Sci., Vol. I, pp. 311-337, 1870. (16 species.)
- DALL, W. H., and BANNISTER, H. M. List of the Birds of Alaska, with Biographical Notes. Transact. Chicago Acad. Science, Vol. I, pp. 267-310, 1870. (212 species.)
- DALL, W. H. Alaska and its Resources. 8vo. Boston, 1870, pp. 580-586. (Contains a nominal list of 212 species.)

Vancouver's Island.

- BROWN, ROBERT. Synopsis of the Birds of Vancouver's Island. Ibis, 1868, pp. 414-428. (153 species.)
- [LORD, JOHN K. The Naturalist in Vancouver's Island and British Columbia. 2 vols. 8vo. London, 1866. Vol. II, pp. 291-301, contains a list of 226 species of birds collected by Mr. Lord.]
- SCLATER, P. L. On a Collection of Birds from Vancouver's Island. Proc. Zool. Soc. Lond., 1859, pp. 235-237. (35 species.)

California.

- BREWER, T. M. List and Descriptions of Eggs obtained in California by Mr. E. Samuels. Proc. Bost. Soc. Nat. Hist., Vol. VI, pp. 145-149, 1857. (16 species.)
- BRIDGES, THOMAS. Notes on California Birds, with Remarks by P. L. Sclater. Proc. Zool. Soc. Lond., 1858, pp. 1-31, 1858. (11 species.)
- COOPER, J. G. The Fauna of California and its Geographical Distribution. Proc. Cal. Acad. Sci., Vol. IV, pp. 61-81, 1869.
- COOPER, J. G. Species of Animals new to the State of California. Proc. Cal. Acad. Sci., Vol. II, pp. 118-123, 1861. (18 species of birds.)
- COOPER, J. G. Some recent Additions to the Fauna of California. Proc. Cal. Acad. Sci., Vol. IV, pp. 3-13, 1868. (33 species of birds added.)
- COOPER, J. G. The Naturalist in California. Amer. Nat., Vol. III, pp. 182-189, 470-481, 1869.
- COOPER, J. G. Monterey in the dry Season. Amer. Nat., Vol. IV, pp. 756-758, 1871. (About 30 species of birds mentioned as summer residents.)
- COOPER, J. G., and BAIRD, S. F. Ornithology of California. Vol. I, [Land Birds,] royal 8vo, 1870.
- COUES, ELLIOTT. From Arizona to the Pacific. Ibis, 2d Ser., Vol. I. pp. 260-275, 1866.
- FEILNER, JOHN. [Ornithological] Explorations in Upper California in 1860. Smith. Rep., 1864, pp. 421-430.

- GAMBEL, WM. Remarks on the Birds observed in Upper California, with Descriptions of New Species. *Proc. Acad. Nat. Sci. Philad.*, 1846-47, pp. 44-48, 110-115, 154-158, 200-205. (84 species.) *Ibid.*, *Journ. Acad. Nat. Sci. Philad.*, 2d Ser., Vol. I, pp. 25-56, 215-229, 1847-1849. (176 species.)
- HEERMANN, A. L. Notes on the Birds of California observed during a Residence of three Years in that Country. *Jour. Acad. Nat. Sci. Phil.*, 2d Ser., Vol. II, pp. 259-272, 1853. (103 species.)
- HEERMANN, A. L. Report on the Birds collected during Explorations in California for Railroad Routes to connect with the Routes near the 35th and 32d Parallels of North Latitude, made by Lieut. R. G. Williamson. *Pacific R. R. Rep. of Expl. and Surveys*, Vol. X, Part IV, No. 2, pp. 29-80. (221 species.)
- LICHTENSTEIN, H. Beiträge zur ornithologischen Fauna von Californien, etc. *Abhandl. der Berlin Akad., Phys. Kl.*, 1838, pp. 417-451.
- SCLATER, P. L. List of Birds collected by Mr. Thomas Bridges, in the Valley of San José, California. *Proc. Zool. Soc. Lond.*, 1857, pp. 125-127. (33 species.)
- VIGORS, N. A. On a Collection of Skins of Birds from California. *Proc. Zool. Soc. Lond.*, Vol. I, p. 65, 1833. (Only a paragraph of ten lines.)
- XANTUS, JOHN. List of Birds collected at Fort Tejon, California. *Proc. Acad. Nat. Sci. Philad.*, 1859, pp. 189-193. (144 species.)

MEXICO.

- BAIRD, S. F. Notes on a Collection of Birds made by John Xantus, Esq., at Cape St. Lucas, Lower California. *Proc. Acad. Nat. Sci. Philad.*, 1859, pp. 299-306. (42 species.)
- BONAPARTE, C. L. Catalogue d'Oiseaux du Mexique et du Pérou. *Rev. Zool.*, 1840, p. 19.
- CABOT, SAMUEL. Descriptions and Habits of some of the Birds of Yucatan. *Journ. Bost. Soc. Nat. Hist.*, Vol. IV, pp. 460-467, 1843. (4 species.)
- CASSIN, JOHN. Catalogue of Birds collected by Mr. Wm. S. [H.] Pease during the March of the Army of the United States from Vera Cruz to the City of Mexico. *Proc. Acad. Nat. Sci. Philad.*, 1848-49, pp. 87-91. (45 species.)
- LAWRENCE, G. N. List of a Collection of Birds from Northern Yucatan. *Ann. N. Y. Lyc. Nat. Hist.*, Vol. IX, pp. 198-210, 1869. (103 species.)
- MONTES-DE-OCA, RAFAEL. The Mexican Humming-Birds. *Proc. Acad. Nat. Sci. Philad.*, 1860, pp. 47, 48, 80, 81, 551-553. (4 species.)
- [PARZUDAKI, E. Catalogue des Oiseaux du Mexique qui composent les collections de M. Sallé. (432 species.)]

- SCLATER, P. L. On a Collection of Birds made by Mr. H. S. le Strange, near the City of Mexico. *Proc. Zoöl. Soc. Lond.*, 1869, pp. 361 - 364. (Of the 262 species said to have been contained in the collection only 15 are particularly mentioned.)
- SCLATER, P. L. List of a Collection of Birds procured by Mr. G. H. White in the Vicinity of the City of Mexico. *Proc. Zoöl. Soc. Lond.*, 1864, pp. 172 - 179. (156 species.)
- SCLATER, P. L. Notes on some Birds from Southern Mexico. *Proc. Zoöl. Soc. Lond.*, 1858, pp. 95 - 99. (14 species.)
- SCLATER, P. L. On a Collection of Birds received by Mr. Sallé from Southern Mexico. *Proc. Zoöl. Soc. Lond.*, 1857, pp. 226 - 230. (29 species.)
- SCLATER, P. L. Catalogue of the Birds collected by M. Aug. Sallé in Southern Mexico, etc. *Proc. Zoöl. Soc. Lond.*, 1856, pp. 283 - 311. (233 species.)
- SCLATER, P. L. On a Collection of Birds made by Sign. Matteo Botteri in the Vicinity of Orizaba, Southern Mexico. *Proc. Zoöl. Soc. Lond.*, 1857, pp. 210 - 215. (38 species.)
- SCLATER, P. L. Notes on a Collection of Birds from the Vicinity of Orizaba and neighboring Parts of Southern Mexico. *Proc. Zoöl. Soc. Lond.*, 1860, pp. 250 - 254. (44 species.)
- SCLATER, P. L. List of Birds collected by Mr. A. Boucard in the State of Oaxaca in Southern Mexico. *Proc. Zoöl. Soc. Lond.*, 1859, pp. 369 - 393. (236 species.)
- SCLATER, P. L. List of Additional Species of Mexican Birds obtained by M. Sallé, from the Environs of Jalapa and San Andres Tuxtla. *Proc. Zoöl. Soc. Lond.*, pp. 201 - 207, 1857. (62 species.)
- SCLATER, P. L. On a Series of Birds collected in the Vicinity of Jalapa, Mexico. *Proc. Zoöl. Soc. Lond.*, 1859, pp. 362 - 369. (226 species.)
- SCLATER, P. L. On some Birds to be added to the Avifauna of Mexico. *Proc. Zoöl. Soc. Lond.*, 1862, pp. 368, 369. (8 species.)
- SUMICHRIST, F. The Geographical Distribution of the Native Birds of the Department of Vera Cruz, with a List of the Migratory Species, translated from the French, by Dr. T. M. Brewer. *Mem. Bost. Soc. Nat. Hist.*, Vol. I, pp. 542 - 563, 1869. (225 species.)
- SWAINSON, WM. A Synopsis of the Birds discovered in Mexico by W. Bullock, Sen. and Jun. *Phil. Mag.*, 2d Ser., Vol. I, pp. 364 - 369, 433 - 442, 1827. (101 species.)

CENTRAL AMERICA.

General.

- BONAPARTE, C. L. Notes sur les collections rapportées en 1853, par M. A. Delattre de son voyage en California et dans le Nicaragua. *Comptes*

- Rendus, Tome XXXVIII, pp. 827-835, 913-927, 1853; Tome XXXVIII, pp. 1-11, 53-66; 258-266, 365-389, 533-542, 650-665, 1854.
- COUES, ELLIOTT. Notes on Certain Central American *Laridæ*. Ibis, 1864.
- MOORE, T. J. List of Mammals and Birds collected by Mr. Jos. Leland in Honduras, Belize, and Guatemala. Proc. Zoöl. Soc. Lond., 1859, pp. 50-65. (129 species.)
- LAWRENCE, GEO. N. Descriptions of New Species of Birds of the Families *Zanagradæ*, etc., etc. Ann. Lyc. Nat. Hist. N. Y., Vol. VIII, pp. 126-135. (12 species.)
- LAWRENCE, GEO. N. Descriptions of Six New Species of Birds from Central America. Ann. Lyc. Nat. Hist. N. Y., Vol. VIII, pp. 170-173.
- SALVIN, OSBERT. A List of Species to be added to the Ornithology of Central America. Ibis, 1861.
- SALVIN, O., and SCLATER, P. L. Descriptions of New Species of Birds of the Families *Dendrocolaptidæ*, *Strigidæ*, and *Columbidæ*. Proc. Zoöl. Soc. Lond., 1868, pp. 53-60.
- SALVIN, OSBERT. On the *Psittacidæ* of Central America. Ibis, 1871, pp. 86-100. (27 species.)
- SCLATER, P. L., and SALVIN, O. On the Ornithology of Central America. Ibis, I, pp. 1-22, 117-138, 213-234, 1859. (382 species.)
- SCLATER, P. L., and SALVIN, O. Descriptions of New and Little-known American Birds, etc. Proc. Zoöl. Soc. Lond., 1868, pp. 322-329.
- SCLATER, P. L., and SALVIN, O. Descriptions of Six New Species of American Birds, etc. Proc. Zoöl. Soc. Lond., 1869, pp. 416-420.
- SUNDEVALL, C. J. Conspectus generis *Certhiolæ*. Öfvers. Kongl. Vetensk. Acad. Forhand., 1869, pp. 619-625. (20 species.)

Guatemala.

- SALVIN, OSBERT, and SCLATER, P. L. Contributions to the Ornithology of Guatemala. Ibis, 1860, pp. 28-45, 272-278. (119 species.)
- SALVIN, OSBERT. Notes on the Humming-Birds of Guatemala. Ibis, 1860, pp. 259-272. (23 species.)
- SALVIN, OSBERT. The Sea-Birds and Waders of the Pacific Coast of Guatemala. Ibis, 1865, pp. 187-199.
- SALVIN, OSBERT. A further Contribution to the Ornithology of Guatemala. Ibis, 1866, pp. 188-206. (81 species.)
- SALVIN, OSBERT. On some Collections of Birds from Veragua. Proc. Zoöl. Soc. Lond., 1867, pp. 129-161. (220 species.)
- SALVIN, OSBERT. Descriptions of Eight New Species of Birds from Veragua. Proc. Zoöl. Soc. Lond., 1866, pp. 67-76.
- SCLATER, P. L., and SALVIN, O. Descriptions of Four New Species of Birds from Veragua. Proc. Zoöl. Soc. Lond., 1868, pp. 387-390.

SCLATER, P. L., and SALVIN, O. Descriptions of Three New Species of Tanagers from Veragua. *Proc. Zool. Soc. Lond.*, 1869, pp. 439–440.

Honduras.

LAWRENCE, G. N. Catalogue of a Collection of Birds in the Museum of the Smithsonian Institution, made by Mr. H. E. Holland at Greytown, Nicaragua, etc. *Ann. Lyc. Nat. Hist. N. York*, 1865, pp. 179–185. (61 species.)

SALVIN, OSBERT. A Fortnight amongst the Sea-Birds of British Honduras. *Ibis*, 1864, pp. 372–387.

SCLATER, P. L. List of Birds collected by Geo. Cavendish Taylor in the Republic of Honduras. *Proc. Zool. Soc.*, 1858, pp. 356–360. (39 species.)

SCLATER, P. L., and SALVIN, OSBERT. List of Birds collected on the Bluefields River, Mosquito Coast, by Mr. Henry Wickham. *Proc. Zool. Soc. Lond.*, 1867, pp. 278–280. (39 species.)

TAYLOR, GEO. CAVENDISH. On Birds collected in the Republic of Honduras, etc. *Ibis*, 1860, pp. 10–24, 110–122, 222–228, 311–317. (96 species.)

Costa Rica.

CABANIS, J. Uebersicht der im Berliner Museum befindlichen Vögel von Costa Rica. *Journ. für Orn.*, VIII Jahrgang, pp. 321–336, 401–416, 1860; IX Jahrgang, pp. 1–14, 96, 241–256, 1861. (113 species.)

CASSIN, JOHN. On some Conirostral Birds from Costa Rica, etc. *Proc. Acad. Nat. Sci. Philad.*, 1865, pp. 169–172. (22 species.)

[FRANZIUS, V. Ueber die geographische Verbreitung der Vögel Costa-Rica's. *Caban. Journ. für Ornith.*, 1869, pp. 195, 290.]

LAWRENCE, G. N. A Catalogue of the Birds of Costa Rica. *Ann. Lyc. Nat. Hist. New York*, 1868, pp. 86–149. (511 species.)

SALVADORI, TOMMASO. Intorno ad alcuni Uccelli di Costa Rica. *Atti della R. Accad. Sc. di Torino*, 1868, pp. 170–185. (23 species.)

SALVIN, OSBERT. Descriptions of Seventeen New Species of Birds from Costa Rica. *Proc. Zool. Soc. Lond.*, 1864, pp. 579–586.

SALVIN, OSBERT. Notes on Mr. Lawrence's List of Costa Rica Birds. *Ibis*, 1869, pp. 310–319; 1870, pp. 107–116. (29 species added to Mr. Lawrence's List.)

New Granada.

LAWRENCE, GEO. N. Catalogue of a collection of Birds made in New Granada, etc. *Ann. Lyc. Nat. Hist. New York*, Vol. VII, pp. 288–302, 315–334; 461–479, 1861; Vol. VIII, pp. 1–13, 1862. (415 species.)

LAWRENCE, GEO. N. List of Birds from near David, Chiriqui, New Granada, etc. *Ann. Lyc. Nat. Hist. New York*, 1865, pp. 175–179. (39 species.)

THE BERMUDA AND WEST INDIA ISLANDS.

Bermudas.

- BLAND, LIEUT. List of the Birds of Bermuda. Smithsonian Report, 1858, pp. 286-289. (114 species.)
- WEDDERBURN, J. W., and HURDIS, J. L. Notes and Observations on the Birds of Bermuda, etc. Jones's The Naturalist in Bermuda. London, 1859, pp. 23-97.

Bahamas.

- BRYANT, HENRY. A List of Birds seen at the Bahamas, from January 20 to May 14, 1859, etc. Proc. Bost. Soc. Nat. Hist., Vol. VII, pp. 102-134, 1860. (87 species.)
- BRYANT, HENRY. Additions to a List of the Birds seen at the Bahamas. Proc. Bost. Soc. Nat. Hist., Vol. XI, pp. 63-70, 1866. (31 species.)

Cuba.

- BREWER, T. M. [Nominal] List of the Birds of Cuba, compiled from two Lists furnished by Dr. John Gundlach. Proc. Bost. Soc. Nat. Hist., Vol. VII, pp. 305-308, 1860. (251 species.)
- GUNDLACH, JOHN. Description of Five new Species of Birds, and other Ornithological Notes of Cuban Species. Journ. Bost. Soc. Nat. Hist., Vol. VI, pp. 313-319, 1852. (18 species in all.)
- GUNDLACH, J. Beiträge zur Ornithologie Cuba's. Journ. für Orn., II Jahrgang, pp. —, 1854; III Jahrgang, 465, 1855; IV Jahrgang, pp. 1, 97, 337, 417, 1856; V Jahrgang, pp. 225-242, 1857. (251 species.)
- GUNDLACH, J. Tabellarische Uebersicht aller bisher auf Cuba beobachteten Vögel. Journ. für Orn., IX Jahrgang, pp. 321-349, 1861. (249 species.)
- GUNDLACH, J. Zusätze und Berichtigungen zu den "Beiträgen zur Ornithologie Cuba's." Journ. für Orn., IX Jahrgang, pp. 401-416; X Jahrgang, pp. 81-96. (249 species.)
- GUNDLACH, JUAN. Revista y Catálogo de las Aves cubanas. Repert. físico-nat. de Cuba, Vol. I, pp. 165-180, 221-242, 281-302, 347-363, 386-403, 1865, 1866. (257 species.)
- GUNDLACH, J. Briefliches von Cuba. Cab. Journ. für Orn., 1866, pp. 352-353.
- LAWRENCE, Geo. N. Notes on some Cuban Birds, etc. Ann. Lyc. Nat. Hist. New York, Vol. VII, p. 247, 1860. (27 species.)
- LEMBEYE, J. Aves de la Isle de Cuba, 8vo. Habana, 1850. (222 species.)
- RAMON DE LA SAGRA. Histoire physique, politique et naturelle de l'île de Cuba, etc. Ornithologie par A. d'Orbigny. 8vo, 1839. (129 species.)
- VIGORS, N. A. On some Birds from Cuba. Zool. Journ., Vol. III, pp. 432-448, 1828. (45 species.)

Jamaica.

GOSSE, P. H. *The Birds of Jamaica.* 8vo. London, 1847.

MARCH, W. T. *Notes on the Birds of Jamaica, with Remarks by S. F. Baird.* *Proc. Acad. Nat. Sci. Philad.*, 1863, pp. 150 – 154, 283 – 304; 1864, pp. 62 – 72. (109 species.)

[OSBURN, W. *Notes on the Mountain Birds of Jamaica.* *Zoölogist*, pp. 6709, 6761.]

SCLATER, P. L. *A List of a Collection of Birds made by the late Mr. W. Osburn in Jamaica, with Notes.* *Proc. Zoöl. Soc. Lond.*, 1861, pp. 69 – 82. (92 species.)

St. Domingo, St. Thomas, Porto Rico, etc.

BRYANT, HENRY. *A List of the Birds of St. Domingo, etc.* *Proc. Bost. Soc. Nat. Hist.*, Vol. XI, pp. 89 – 98, 1866. (78 species.)

BRYANT, H. *A List of Birds from Porto Rico, etc.* *Proc. Bost. Soc. Nat. Hist.*, Vol. X, pp. 248 – 257, 1866. (41 species.)

CASSIN, JOHN. *Catalogue of Birds from the Island of St. Thomas, West Indies, collected and presented to the Academy of Natural Science by Mr. Robert Swift.* *Proc. Acad. Nat. Sci.*, 1860, pp. 374 – 379. (27 species.)

LAWRENCE, G. N., and JULIEN, A. A. *Catalogue of Birds collected at the Island of Sombrero, W. I.* *Ann. Lyc. Nat. Hist. New York.*, VIII, pp. 93 – 107, 1864. (34 species.)

NEWTON, A. and E. *Observations on the Birds of St. Croix, West Indies, made between February 20 and August 6, 1857, and March 4 and September 28, 1858.* *Ibis*, I, pp. 59 – 69, 138 – 150, 253 – 264, 365 – 379, 1859. (64 species.)


SALLÉ, AUG. *Liste des Oiseaux rapportés et observés dans la République Dominicaine, pendant son voyage de 1849 – 1851.* *Proc. Zoöl. Soc. Lond.*, 1857, pp. 230 – 237. (61 species.)

SUNDEVALL, C. J. *Aves Insulæ Portorico.* *Öfvers. Kongl. Vetensk. Akad. Forhand.*, 1869, pp. 593 – 603. (90 species.)

SUNDEVALL, C. J. *Aves Insulæ Sancti Barthelemy.* *Öfvers. Kongl. Vetensk. Akad. Forhand.*, 1869, pp. 579 – 592. (47 species.)

CAMBRIDGE, April, 1871.

N. B. — The Museum of Comparative Zoölogy would gladly accept books not yet upon the shelves of its library in exchange for its publications or zoölogical specimens.



JB


C.A.V.

1-2, *Tyrannus carolinensis*; 3-5, *Troglodytes aedon*; 6-11, *Seiurus borealis*; 12-14, *Minotilta varia*; 15-16, *Dendroica striata*; 17-18, *Agelaius phoeniceus*; 19-20, *Pyrrhuloxia rubra*.

Plate IV.*

- Fig. 1 - 1a. *Tyrannus carolinensis*, specimen No. 6942, from Eastern Massachusetts.
- " 2 - 2a. *Tyrannus carolinensis*, specimen No. 6945, from Eastern Massachusetts.
- " 3 - 3a. *Troglodytes aëdon*, specimen No. 10931, ♀, from Jacksonville, Fla.
- " 4 - 4a. *Troglodytes aëdon*, specimen No. 10684, ♀, from Dummitt's, Fla.
- " 5 - 5a. *Troglodytes aëdon*, specimen No. 10683, from Dummitt's, Fla.
- " 6 - 6a. *Troglodytes aëdon*, specimen No. 5212, from Welaka, St. John's River, Fla.
- " 7 - 7a. *Troglodytes aëdon*, specimen No. 10930, from Jacksonville, Fla.
- " 8 - 8a. *Seiurus noveboracensis*, specimen No. 5447, from Mount Tom, Mass.
- " 9 - 9a. *Seiurus noveboracensis*, specimen No. 1442, from Weston, Mass.
- " 10 - 10a. *Seiurus noveboracensis*, specimen No. 6794, from Brookline, Mass.
- " 11 - 11a. *Seiurus noveboracensis*, specimen No. 4246, from Waterville, Maine.
- " 12 - 12a. *Mniotilta varia*, specimen No. 5148, ♂, from Jacksonville, Fla.
- " 13 - 13a. *Mniotilta varia*, specimen No. 6806, ♂, from Brookline, Mass.
- " 14 - 14a. *Mniotilta varia*, specimen No. 8216, ♂, from Hudson, Mass.
- " 15 - 15a. *Dendræca striata*, specimen No. 5052, ♂, from Watertown, Mass.
- " 16 - 16a. *Dendræca striata*, specimen No. 4367, ♂, from Newtonville, Mass.
- " 17 - 17a. *Pipilo erythrophthalmus*, specimen No. 4727, ♂, from Weston, Mass.
- " 18 - 18a. *Pipilo erythrophthalmus*, specimen No. 10721, ♂, from Dummitt's, Fla.
- " 19 - 19a. *Pyranga æstiva*, specimen No. 10629, ♂, from Jacksonville, Fla.
- " 20 - 20a. *Pyranga æstiva*, specimen No. 5431, ♂, from Jacksonville, Fla.

* At bottom of Plate IV, last line, for "*Pyranga rubra*" read "*Pyranga æstiva*."



JB

L.A.W.CC

1-6. *Agelaius lanius*. 7-10. *Chrysomitris tristis*. 11-12. *C. pinus*. 13-15. *Curvirostra americana*. 16-18. *Passerculus savanna*.

Plate V.

- Fig. 1 - 1a. *Ægiothus linaria*, specimen No. 10859, ♂, from Newton, Mass.
- " - 2a. *Ægiothus linaria*, specimen No. 10860, ♂, from Newton, Mass.
- " 3 - 3a. *Ægiothus linaria*, specimen No. 6392, ♂, from Fort Anderson, British America. (An original specimen of *Æ. fuscescens* Coues. — Smith. Inst., No. 43386.)
- " 4 - 4a. *Ægiothus linaria*, specimen No. 4943, from Newton, Mass.
- " 5 - 5a. *Ægiothus linaria*, specimen No. 3229, from Southern Maine.
- " 6 - 6a. *Ægiothus linaria*, specimen No. 6489, ♀, from Fort Simpson, British America. (An original specimen of *Æ. exilipes* Coues. — Smith. Inst., No. 27431.)
- " 7 - 7a. *Chrysomitris tristis*, specimen No. 6453, ♂, from Rocky Mountains, west of Denver, Colorado.
- " 8 - 8a. *Chrysomitris tristis*, specimen No. 8125, ♂, from Springfield, Mass.
- " 9 - 9a. *Chrysomitris tristis*, specimen No. 4925, from Newtonville, Mass.
- " 10 - 10a. *Chrysomitris tristis*, specimen No. 4631, from Newtonville, Mass.
- " 11 - 11a. *Chrysomitris pinus*, specimen No. 9556, from Waterville, Maine.
- " 12 - 12a. *Chrysomitris pinus*, specimen No. 10875, from Gorham, New Hampshire.
- " 13 - 13a. *Curvirostra americana*, specimen No. 4639, ♂, from Newton, Mass.
- " 14 - 14a. *Curvirostra americana*, specimen No. 4638, ♂, from Newton, Mass.
- " 15 - 15a. *Curvirostra americana*, specimen No. 4637, ♂, from Newton, Mass.
- " 16 - 16a. *Passerculus savanna*, specimen No. 5084, ♂, from Ipswich, Mass.
- " 17 - 17a. *Passerculus savanna*, specimen No. 5175, ♂, from Hibernia, St. John's River, Florida.
- " 18 - 18a. *Passerculus savanna*, specimen No. 7119, ♂, from Henley Harbor, Labrador.




Plate VI

- Fig. 1 - 1a. *Agelæus phæniceus*, specimen No. 8071, ♂, from Orleans, Mass.
- " 2 - 2a. *Agelæus phæniceus*, specimen No. 10572, ♂, from Dummitt's, Fla.
- " 3 - 3a. *Agelæus phæniceus*, specimen No. 4589, ♂, from Newtonville, Mass.
- " 4 - 4a. *Agelæus phæniceus*, specimen No. 8068, ♂, from Orleans, Mass.
- " 5 - 5a. *Agelæus phæniceus*, specimen No. 10569, ♂, from Dummitt's, Fla.
- " 6 - 6a. *Agelæus phæniceus*, specimen No. 10576, ♂, from Dummitt's, Fla.





Plate VII

- Fig. 1 - 1a. *Quiscalus purpureus*, specimen No. 4601, ♂, from Newtonville, Mass.
- " 2 - 2a. *Quiscalus purpureus*, specimen No. 5201, ♂, from Welaka, St. John's River.
- " 3 - 3a. *Quiscalus purpureus*, specimen No. 4603, ♂, from Newtonville, Mass.
- " 4 - 4a. *Quiscalus purpureus*, specimen No. 8072, ♂, from Orleans, Mass.
- " 5 - 5a. *Quiscalus purpureus*, specimen No. 6834, ♂, from Phillipsburg New Jersey,
- " 6 - 6a. *Quiscalus purpureus*, specimen No. 6848, ♂, from Cape Florida.



JB

Collins

1-4. *Sturnella ludoviciana* 5-6. *Colaptes auratus*

Plate VIII.

- Fig. 1 - 1a. *Sturnella ludoviciana*, specimen No. 5370, ♀, from Hawkinsville, Fla.
- " 2 - 2a. *Sturnella ludoviciana*, specimen No. 5372, ♀, from Hawkinsville, Fla.
- " 3 - 3a. *Sturnella ludoviciana*, specimen No. 5339, ♂, from Enterprise, Fla.
- " 4 - 4a. *Sturnella ludoviciana*, specimen No. 5340, ♂, from Enterprise, Fla.
- " 5 - 5a. *Colaptes auratus*, specimen No. 4881, ♀, from Newton, Mass.
- " 6 - 6a. *Colaptes auratus*, specimen No. 5464, ♀, from Newton, Mass.
-

ERRATA. At the upper left-hand corner of all the plates, for "Bull. M. C. Z., Vol. II, No. 2," read "Bull. M. C. Z., Vol. II, No. 3."



Allen, J. A. 1871. "On the mammals and winter birds of east Florida, with an examination of certain assumed specific characters in birds, and a sketch of the bird faunae of eastern North America." *Bulletin of the Museum of Comparative Zoology at Harvard College* 2(3), 161–450.

View This Item Online: <https://www.biodiversitylibrary.org/item/29976>

Permalink: <https://www.biodiversitylibrary.org/partpdf/298635>

Holding Institution

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Sponsored by

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.