



LONDON JOHN MURRAY







Meanan 736

280

# MOVEMENTS AND HABITS

# CLIMBING PLANTS.

263047-2001

# MOVEMENTS AND HABI

# CLIMBING PLANTS.

By CHARLES DARWIN, M.A., F.R.S.

JOHN MURRAY, ALBEMARLE STREET.

Darwin Online: By permission of the Trustees of the Natural History Museum

### BY THE SAME AUTHOR

ON THE ORIGIN OF SPECIALS BY MEANS OF NATURAL SEASON PARTICLES. THE SEASON PARTICLES AND PARTICLES FROM PARTICL

ON THIS VARIOUS CONTRIVANCIS OF WHOSE BRITTISH GOOD INTERIOR CONTRIVANCIS OF WHOSE BRITTISH GOOD INTERIOR CHARM. THE BRITTISH GOOD INTERIOR CHARM. THE BRITTISH CONTRIVATION OF THE WORLDOW, A A STREAM OF THE WORLDOW O

FACTS AND ARGUMENTS FOR DARWIN. By FEITZ MCLASS. Translated by W. S. Ballas, F.L.S. With Illustrations. Post 890. MUSICAY.

A MONOGRAPH OF THE CIERIPEDIA.

### PREFACE.

This Essay first appeared in the ninth volume of the 'Journal of the Linnean Society,' published in 1865. It is here reproduced in a corrected and, I hope, clearer form, with some additional facts. The illustrations were drawn by my son, George Darwin. Fritz Müller, after the publication of my paper, sent to the Linnean Society (Journal, vol. ix., p. 344) some interesting observations on the climbing plants of South Brazil, to which I shall frequently refer. Recently two important memoirs, chiefly on the difference in growth between the upper and lower sides of tendrils, and on the mechanism of the movements of twining plants, by Dr. Hugo de Vries, have appeared in the 'Arbeiten des Botanischen Instituts in Würzburg,' Heft. iii., 1873. These memoirs ought to be carefully studied by every one interested in the subject, as I can here give only references to the more important points. This excellent observer, as reasons assigned towards the close of my fourth chapter. I cannot persuade myself that this holds me most, I may call his attention to certain tendrilbearing plants; for instance, Bignonia capreolata, Cobea, Echinocystis, and Hanburya, which display as beautiful adaptations as can be found in any part of the kingdom of nature. It is, also, an interesting widely different functions, may be observed on the the common vine; and these cases illustrate in a striking manner the principle of the gradual evolution of species.

<sup>\*</sup> An English translation of the 'Lehrbuch der Botanik' by Professor Sachs, has recently (1875), appeared under the title

by is a great boon to all lovers of titly natural science in England.

Introductory remarks-Description of the twining of the Hop--Torsion of the stems-Nature of the revolving movement and manner of ascent-Stems not irritable-Rate of revolution in various plants - Thickness of the support round which plants can twine-Species which revolve in an anoma-

Plants which climb by the aid of spontaneously revolving and sensitive petioles—Clematis—Tropacolum—Maurundia, flowerpeduneles moving spontaneously and sensitive to a touchthickening of the clasped petioles - Fumaria - Adlumia-Plants which climb by the aid of their produced midribs-Gloriosa - Flagellaria - Nepenthes - Summary on leafclimbers ... 45-89

Nature of tendrils-Bignoniacem, various species of, and their different modes of climbing-Tendrils which avoid the light, and creep into crevices-Development of adhesive discshooked tendrils, their manner of action - LEGUMINOS.E -Composite — Smilacre — Smilax aspera, its inefficient tendrils - Fumaniacem - Corydalis claviculata, its state .. .. .. .. .. Pages 84-126

## CHAPTER IV.

TENDRIL-BEARERS-configured.

CUCURETTACES - Homologous nature of the tendrils-Echinowater - Spiral contraction of tendrils - Summary on the

HOOK AND ROOT-CLIMBERS,-CONCLUDING REMARKS.

Plants climbing by the aid of hooks, or merely scrambling over and the stages of their development .. .. 183-206

# MOVEMENTS AND HABITS

## CHAPTER L

Introductory remarks-Description of the twining of the Hop-Torsion Thickness of the support round which plants can twine-Species

I was led to this subject by an interesting, but short paper by Professor Asa Gray on the movements of the tendrils of some Cucurbitaceous plants.\* My observations were more than half completed before I learnt that the surprising phenomenon of the spontaneous revolutions of the stems and tendrils of climbing plants had been long ago observed by Palm and by Hugo von Mohl,† and had subsequently been the subject of two memoirs by Dutrochet.1 Nevertheless.

pflanzen,' 1827. Palm's Treatise

frey), by H. von Mohl, n. 147 to t "Des Mouvements révolutifs

Darwin Online: By permission of the Trustees of the Natural History

I believe that my observations, founded on the ex-

are not aided by any other movement. Secondly, those endowed with irritable organs, which when they modified leaves, branches, or flower-peduncles. But these two classes sometimes graduate to a certain extent into one another. Plants of the third class ascend merely by the aid of hooks; and those of the exhibit any special movements, they present little

This is the largest subdivision, and is apparently My observations will be best given by taking a few special cases. When the shoot of a Hop (Humulus formed joints or internodes are straight and remain stationary; but the next-formed, whilst very young,

tom. xvii. (1843) p. 989; "Re- Tiges," &c., tom. xix.(1844) p. 295.

Darwin Online: By permission of the Trustees of the Natural History

may be seen to bend to one side and to travel slowly round towards all points of the compass, moving, like the hands of a sutch, with the sun. The movement very soon acquires its full collinary velocity. From severa observations made thring dangest on shoots percent observations made thring dangest on shoots persecutions of the second contract of the conmorder plant during Asynt, the recongeration that the weather and during the day is 2 km s. for each venturing and more of the revolutions varied much from this rate. The revolving movement containes as long as the plant continues to grow; but each separate intermola, as it becomes old, crosses to more.

To ascertain more precisely what amount of move-

ment each internols undercent, I kept a potted plant, during the night and day, in a well-earmed roun to which I was confined by illness. A long shoot projected beyond the upper end of the supporting stick, and was steadily revolving. I then took a longer stick and tied up the shoot, so that only a very yong internolo; I jof an inch in length, was left free. This was so mostly ouright that its revolution could not be easily observed; but it certainly moved, and the side of the internolo which was at one time curvex beame conexe, which, as we shall hereafter see, is a sure sign of the state of the side of the control of the side of the control with the side of the control with the side of the control of the side of th

of the room varied only a little. The shoot had now forward, the revolutions were easily observed. The thirty-sixth revolution was performed at the usual the lower part of the internode ceased to move.

A few more remarks will complete all that need be said about this internucle. It moved during the days; but the more rapid movements, after the performance of the third revolution, lated during three days and twenty hours. The regular revolutions, from the ninth to thirty-sixth inclusive, were effected at the average rate of 2 hrs. 31 m.; but the weather was cold, and this affected the temperature of the room, especially during the night, and consequently retarded the rate of movement a little. There was only one irragular movement, which consisted in the stem spalify making, after an unusually alow receptation, and we segment of a circle. After the seventeenth revolution the internode had grown from 12 to 6 inches in length, and carried an internode 17 inch long, which was ultimate internode. After the twenty-first revolution, the penultimate internode was 24 inches long, and probably revolved in a period of about three hours. At the twenty-seventh revolution the lower and still moving internode was 83, the penultimate 34, and the ultimate 21 inches in length; and the inclination of the whole shoot was such, that a circle 19 inches in diameter was swept by it. When the movement ceased, the lower internode was 9 inches, and the penultimate 6 inches in length; so that, from the twenty-seventh to thirty-seventh revolutions inclusive, three internodes were at the same time revolving.

The lower internode, when it ceased revolving, became upright and rigid; but as the whole shoot was left to grow unsupported, it became after a time bent into a nearly horizontal position, the uppermost and growing internodes still revolving at the extremity, but of course no longer round the old central point of the supporting stick. From the changed position of the centre of gravity of the extremity, as it revolved, a slight and slow swaying movement was given to the long horizontally projecting shoot; and this movement I at first thought was a spontaneous one. As the shoot grew, it hung down more and more, whilst the growing

Darwin Online: By permission of the Trustees of the Natural History

with most of the plants observed by me. With all, if in the lower one ceased to revolve, the one above was in to move. With Hoya carnosa, on the other hand, a

depending shoot, without any developed leaves, 32 inches in length, and consisting of seven internodes (a minute terminal one, an inch in length, being to side in a semicircular course, with the extreme internodes making complete revolutions. This swaying movement was certainly due to the movement of the lower internodes, which, however, had not force supporting stick. The case of another Asclepiadaceous plant, viz., Ceropegia Gardnerii, is worth briefly giving. I allowed the top to grow out almost horizontally to the length of 31 inches; this now consisted of three long internodes, terminated by two short ones. The and 6 hrs. 45 m, for each revolution. The extreme tip thus made a circle of above 5 feet (or 62 inches) in diameter and 16 feet in circumference, travelling at the rate of 32 or 33 inches per hour. The weather being hot, the plant was allowed to stand on my study-table; and it was an interesting spectacle to watch the long shoot sweeping this grand cricle, night and day, in

Darwin Online: By permission of the Trustees of the Natural History

If we take hold of a growing sapling, we can of course bend it to all sides in succession, so as to make the tip describe a circle, like that performed by the summit of a spontaneously revolving plant. By this movement the sapling is not in the least twisted round its own axis. I mention this because if a black point be painted on the bark, on the side which is uppermost when the sapling is bent towards the holder's body, as the circle is described, the black point gradually turns round and sinks to the lower side, and comes up again when the circle is completed; and this gives the false appearance of twisting, which, in the case of spontaneously revolving plants, deceived me for a time. The appearance is the more deceitful because the axes of nearly all twining-plants are really twisted: and they are twisted in the same direction with the spontaneous revolving movement. To give an instance, the internode of the Hop of which the history has been recorded, was at first, as could be seen by the ridges on its surface, not in the least twisted; but when, after the 37th revolution, it had grown 9 inches long, and its revolving movement had ceased, it had become twisted three times round its own axis, in the line of the course of the sun; on the other hand, the common Convolvulus, which revolves in an opposite course to the Hop, becomes twisted in an opposite direction.

Hence it is not surprising that Hugo von Mohl (p. 105, 108, &c.) thought that the twisting of the axis caused the revolving movement; but it is not

and Lecontea revolved during several days, but became and tendril-bearing plants (as Pisum sativum, Echinocustis lobata, Bignonia capreolata, Eccremocarpus seaber, and with the leaf-climbers, Solanum jasminoides and various species of Clematis), of which the internodes are not twisted, but which, as we shall hereafter see, retwining-plants. Moreover, according to Palm (pp. 30, 95) and Mohl (p. 149), and Léon,\* internodes may occasionally, and even not very rarely, be found which are twisted in an opposite direction to the other internodes on the same plant, and to the course of their revolutions; and this, according to Léon (p. 356), is the case with all the internodes of a certain variety of Phaseolus multiflorus. Internodes which have become to revolve, are still capable of twining round a support,

as I have several times observed. Mohl has remarked (p. 111) that when a stem twines round a smooth evlindrical stick, it does not become

<sup>\* &#</sup>x27;Bull, Bot Soc, de France,' tom, v. 1858, p. 856

twisted.\* Accordingly I allowed kidney-beans to run up stretched string, and up smooth rods of iron and glass, one-third of an inch in diameter, and they mechanical necessity from the spiral winding. The twisted. The influence of the roughness of the support in causing axial twisting was well seen in the stems which had twined up the glass rods; for these rods were fixed into split sticks below, and were secured above to cross sticks, and the stems in passing these places became much twisted. As soon as the stems which had ascended the iron rods reached the summit and became free, they also became twisted; and this apparently occurred more quickly during windy than during calm weather. Several other facts could be given, showing that the axial twisting stands in some relation to inequalities in the support, and likewise to the shoot revolving freely without any support. Many plants, round their own axes; † but this occurs so much more

\* This whole subject has be ably discussed and explained.
H. de Vries, 'Arbeiter des B
Instituts in Würzburg,' Heft!
Jastituts in Würzburg,' Heft!
Jastituts in Würzburg,' Heft!
Hook of Botany,' English transition, 1875, p. 779), who concluse that the subject is the result of grown withat torsion is the result of grown

ed by the inner layers."

8 Bot. † Professor Asa Gray has refl iii. marked to me, in a letter, that in

The treat They occidental is the twisting of mulathe bark is very conspicuous. The ludes twist is generally to the right of rowth the observer; but, in noticing generally and strongly with twining-plants than with other plants, that there must be some connexion between the capacity for twining and axial twisting, (on the same principle that a much twisted rope is stiffer than a slackly twisted one), and is thus indirectly benefited so as to be enabled to pass over inequalities in its spiral ascent, and to carry its own

I have alluded to the twisting which necessarily follows on mechanical principles from the spiral ascent of a stem, namely, one twist for each spire but, as I shall have to recur to this subject under Tendrils, it may be here passed over.

The revolving movement of a twining plant has been compared with that of the tip of a sapling, moved round and round by the hand held some way down the stem; but there is one important difference-The upper part of the sapling when thus moved

Scanish chestnut is often much

'Scottish Farmer,' 1865, p. 833. stems of many plants occasionally "some of these cases, if not all,

remains straight; but with twining plants every part of the revolving shoot has its own separate and independent movement. This is easily proved; for when the lower half or two-thirds of a long revolving shoot is tied to a stick, the upper free part continues steadily revolving. Even if the whole shoot, except an inch or two of the extremity, be tied up, this part, as I have seen in the case of the Hop, Ceropegia, Convolvulus, &c., goes on revolving, but much more slowly; for the internodes, until they have grown to some little length, always move slowly. If we look to the one, two, or several internodes of a revolving shoot, during the whole or during a large part of each revolution. Now if a coloured streak be painted (this was done with a large number of twining plants) along, we will say, the convex surface, the streak will after a time (depending on the rate of revolution) be found to be running laterally along one side of the bow, then along the concave side, then laterally on the opposite side, and, lastly, again on the originally revolving movement the internodes become bowed in every direction. The movement is, in fact, a continuous self-bowing of the whole shoot, successively directed to all points of the compass; and has been

directed to all points of the compass; and has been well designated by Sachs as a revolving nutation. As this movement is rather difficult to understand, it will be well to give an illustration. Take a sapling to the east, and the black line will be seen to run the north, the black line will be on the concave surface; bend it to the west, the line will again be on the lateral face; and when again bent to the south. the line will be on the original convex surface. Now, cells along its northern surface from the base to the tip were to grow much more rapidly than on the three other sides, the whole shoot would then necessarily be bowed to the south; and let the longitudinal growing and so round by the south, by the east, again to the north. In this case the shoot would remain always bowed with the painted line appearing on the several above specified surfaces, and with the point of the shoot successively directed to each point of the compass. In fact, we should have the exact kind of movement performed by the revolving shoots of twining plants.\*

It must not be supposed that the revolving movetion; in very many cases the tip describes an ellipse, even a very narrow ellipse. To recur once again to

is that advanced by Suchs and

Darwin Online: By permission of the Trustees of the Natural History

our illustration, if we suppose only the northern and seathern surfaces of the sapling alternately to grow rapidly, the summit would describe a simple are; if the growth first travelled a very little to the western face, and during the return a very little to the outstern face, a narrow clippe would be described; and the sapling would be straight as it passed to and for through the intermediate space; and a complete straightening of the shoot may often be observed in revolving plants. The movement is frequently such that three of the side of the shoot seem to be growing in due order more rapidly than the remaining adde; so that a semi-circle instead of a circle is described, the shoot becoming straight and upiquid during laif of its

When a revolving shoot consists of several internoles, the lower ones bend together at the same rate, but one or two of the terminal ones bend at a slower rate; hence, though at times all the internoles are in the same direction, at other times the shoot is rendered slightly seprentine. The rate of revolution of the whole shoot, if judged by the movement of the extreme tip, is thus at times uscelested or related. One other point must be noticed. Authors have observed that the each of the shoot in many twining plants is completely bendered, their the present of fractures, one of the control of th other internodes; for a line painted on the convex owing to the youth of these terminal internodes, the internodes, is of service to the plant ; for not only does tant) it causes the extremity of the shoot to embrace the support much more closely than it could otherwise being blown away during windy weather, as I have hook only straightens itself periodically, and never selves or become periodically straight, in the manner just described; for the hooked form may in some cases be permanent, and be due to the manner of growth of the species, as with the tips of the shoots of the common vine, and more plainly with those of Cissus discolor-plants which are not spiral twiners.

The first purpose of the spontaneous revolving movement, or, more strictly speaking, of the con-

H. de Vrics (ibid. p. \$37); he end of the shoot remains hooked ennears to be a difficult and complex problem, discussed by Dr.

points of the compass, is, as Mohl has remarked, to favour the shoot finding a support. This is admirably effected by the revolutions carried on night and day, a wider and wider circle being swept as the shoot increases in length. This movement likewise explains how the plants twine; for when a revolving shoot meets with a support, its motion is necessarily arrested at the point of contact, but the free projecting part goes on revolving. As this continues, higher and higher points are brought into contact with the support and are arrested; and so onwards to the extremity; and thus the shoot winds round its support, When the shoot follows the sun in its revolving course, it winds round the support from right to left, the support being supposed to stand in front of the beholder; when the shoot revolves in an opposite direction, the line of winding is reversed. As each internode loses from age its power of revolving, it likewise loses its power of spirally twining. If a man swings a rope round his head, and the end hits a stick, it will coil round the stick according to the direction of the swinging movement ; so it is with a twining plant, a line of growth travelling round the free part of the shoot causing it to bend towards the opposite side, and this replaces the momentum of the free end of the rope,

tass repances the momentum of the iree end of the rope.
All the authors, except Palm and Mohl, who have
discussed the spiral twining of plants, maintain that
such plants have a natural tendency to grow spirally.
Mohl believes (p. 112) that twining stems have

any object which they touch; but this is denied by Palm. Even before reading Mohl's interesting tested it in every way that I could, but always with effect. I then tied a light forked twig to a shoot of a Hop, a Coropegia, Spherostema, and Adhatoda, so that the fork pressed on one side alone of the shoot and revolved with it; I purposely selected some very slow revolvers, as it seemed most likely that these would was any effect produced." Moreover, when a shoot winds round a support, the winding movement is always slower, as we shall immediately see, than whilst it revolves freely and touches nothing. Hence indeed it is not probable that they should be so, as wish to assert that they are never irritable; for the growing axis of the leaf-climbing, but not spirilly twining, Lophospermum scandens is, certainly irritable;

\* Dr. H. de Vries also has shown ibid, p. 321 and S25) by a better method than that employed support is exactly what I have de by no. that the stems of twining

do not possess any such quality, for directly after putting a stick to the *Lophospermum*, I saw that it behaved differently from a true twiner or any other leaf-climber.\*

The belief that twiners have a natural tendency to grow spirally, probably arose from their assuming a spiral form when wound round a support, and from the assuming this form. The free internodes of vigorously growing plants, when they cease to revolve, become straight, and show no tendency to be spiral; but when a shoot has nearly ceased to grow, or when the plant is unhealthy, the extremity does occasionally become spiral. I have seen this in a remarkable spire, just like a tendril; and this was apt to occur after some small, ill-formed leaves had perished. The exof the terminal internodes very gradually and successively lose their power of movement, whilst the portions just above move onwards and in their turn become motionless; and this ends in forming an

When a revolving shoot strikes a stick, it winds round it rather more slowly than it revolves. For instance, a shoot of the *Ceropegia*, revolved in 6 hrs.,

<sup>†</sup> Dr. H. de Vries states (ibid. p. 322) that the stem of Cus

but took 9 hrs 30 m to mul

a stick; Aristolochia gigas revolved in about 5 hrs., but took 9 hrs. 15 m. to complete its spire. This, I presume, is due to the continued disturbance of the impelling points; and we shall hereafter see that even shaking a plant retards the revolving movement. The terminal the Ceropegia, after they had wound round a stick, always slipped up it, so as to render the spire more open than it was at first; and this was probably in part due to the force which caused the revolutions, and allowed to act freely. With the Wistaria, on the other hand, a long horizontal shoot wound itself at first into a very close spire, which remained unchanged; but subsequently, as the shoot twined spirally up its support, it made a much more open spire. With all the many plants which were allowed freely to ascend a support, the terminal internodes made at first a close spire; and this, during windy weather, served to keep the shoots in close contact with their support; but as the penultimate internodes grew in length, they pushed themselves up for a considerable space (ascertained by coloured marks on the shoot and on the support) round the stick, and the spire became more open."

It follows from this latter fact that the position

\* See Dr. H. de Vries (Ibid. p. 324) on this subject.

Darwin Online: By permission of the Trustees of the Natural History

occupied by each leaf with respect to the support, depends on the growth of the internodes after they on account of an observation by Palm (p. 34), who in a row, exactly over one another, on the same side of the supporting stick, whatever its thickness may be. My sons visited a hop-field for me, and reported that though they generally found the points of insertion of the leaves standing over each other for a space forming, as might have been expected, an irregular the regularity of position of the leaves. From casual dozen plants, and gave them sticks of various thicknesses, as well as string, to twine round; and in this case one alone out of the dozen had its leaves arranged in a perpendicular line : I conclude, therefore,

The leaves of different twining-plants are arranged on the stem (before it has twined) alternately, or oppositely, or in a spire. In the latter case the line of insertion of the leaves and the course of the revolutions coincide. This fact has been well shown by Dutrochet,\*

<sup>\*</sup> Comptes Rendus, 1844, tom, xix. p. 295, and Annales des Sc. Not.

c 2 Darwin Online: By permission of the Trustees of the Natural History

who found different individuals of Solonsun dulemanses vinning in opposite directions, and these had their leaves in each case spirally arranged in the same direction. A dense whorl of many leaves would apparently be incommodious for a twining plant, and some authors assert that none have their leaves thus arranged; but a training Sichonseit has whorls of three beach of the training Sichonseit has whorls of three beach.

has not as yet been encircled, be suddenly taken away, the shoot generally springs forward, showing After a shoot has wound round a stick, if this be withdrawn, it retains for a time its spiral form; it then viously alluded to offered some curious peculiarities. of movement was retained, this was not sufficient to enable the plant to twine. I then moved the stick to a greater distance, so that it was struck by mate internode; and it was then neatly encircled by this part of the penultimate and by the ultimate course of the day the curled portion straightened not move, a sort of hinge separating the moving and the motionless part of the same intermedo. After a few days, however, I found that this lower part had likewise recovered its revolving poorer. These several facts show that the power of movement is not immediated by lost in the arrested portion of a revolving shoot; and that after being temporarily lost it can be recovered. When a shoot has remained for a considerable time round a support, it permanently retains its spiral form even when the support is removed.

When a tall stick was placed so as to arrest the lower and rigid internodes of the Ceropegia, at the distance at first of 15 and then of 21 inches from the centre of revolution, the straight shoot slowly and gradually slid up the stick, so as to become more and more highly inclined, but did not pass over the summit. Then, after an interval sufficient to have allowed of a semi-revolution, the shoot suddenly bounded from the stick and fell over to the opposite side or point of the compass, and reassumed its previous slight inclination. It now recommenced revolving in its usual course, so that after a semirevolution it again came into contact with the stick, again slid up it, and again bounded from it and fell over to the opposite side. This movement of the shoot had a very odd appearance, as if it were disgusted with its failure but was resolved to try again. We shall, I think, understand this movement by considering the former illustration of the sapling, in

from the northern by the western to the southern face; and thence back again by the eastern to the northern face, successively bowing the sapling in all it, as soon as the circulatory growth reached the But as soon as growth on the southern surface began, ment up the stick; and then, as soon as the eastern stick, and its weight coinciding with the effects of the changed surface of growth, would cause it suddenly to fall to the opposite side, reassuming its previous slight inclination; and the ordinary revolving movement would then go on as before. I have described this curious case with some care, because it first led me to understand the order in which, as I then thought, the surfaces contracted; but in which, as we now know from Sachs and H. de Vries, they grow for a time rapidly, thus causing the shoot to bow towards the opposite side.

The view just given further explains, as I believe, a fact observed by Mohl (p. 135), namely, that a revolving shoot, though it will twine round an object as thin as a thread, cannot do so round a thick support. I placed some long revolving shoots of a Wistaria close to a post between 5 and 6 inches in diameter, but, though aided by me in many ways, they could not wind round it. This apparently was due to the flexure of the shoot, whilst winding round an object so gently curved as this post, not being sufficient to hold the shoot to its place when the growing surface crept round to the opposite surface of the shoot; so that it was withdrawn at each revolution from its support.

When a free shoot has grown far beyond its support, it sinks downwards from its weight, as already explained in the case of the Hop, with the revolving extremity turned upwards. If the support be not lofty, the shoot falls to the ground, and resting there, the extremity rises up. Sometimes several shoots, when flexible, twine together into a cable, and thus support one another. Single thin depending shoots, such as those of the Sollya Drummondii, will turn abruptly backwards and wind up on themselves. The greater number of the depending shoots, however, of one twining plant, the Hibbertia dentata, showed but little tendency to turn upwards. In other cases, as with the Cryptostegia grandiflora, several internodes which were at first flexible and revolved, if they did not succeed in twining round a support, become quite rigid, and supporting themselves upright, carried on their summits

they ounger revolving internotes.

Here will be a convenient place to give a Table showing the direction and rate of movement of several twining plants, with a few appended remarks. These plants are arranged according to Lindley's 'Vegetable Kingdom' of 1853; and they have been selected from

behave in a nearly uniform manner.\*

# (Acotyledons.)

(ACOTYLEDONS.)

Lyyodium sexudens (Polypodincen) moves against the s

June 18, 1st circle was made in . 6 0

" 18, 2nd " " . 6 15 (late in evening

19 2nd . . 5 32 (were het day)

" 20,5th " " " . 6 0

Lygodium articulatum moves against the su

N. M. M. S. T. L. 10 Let electe man mode in 16 20 (about we

July 19, 1st circle was made in . 16 30 (shoot very young , 20, 2nd , , , , , 15 0

, 22, 4th , , , 10 30

## (Monocotyledons.)

Ruscus androgynus (Liliacee), placed in the hot-house, moves against the sun.

May 24, 1st circle was made in . 6 14 (shoot very pr., 25, 2nd ..., 2 21, 25, 3rd ..., 3 37, 25, 4th ..., 3 322

25, 3rd , , , , 3 57 25, 4th , , , , 3 22 26, 5th , , , , , 2 50 27, 6th , , , , 3 52 27, 7th , , , , 4 11

\* I am much indebted to Hocker for having sent me m plants from Kew; and to Veitch, of the Royal Exotic N sery, for having generously gi

ny Gray, Prof. Oliver, and Dr. Hooker Ir. have afforded me, as on many arrevious occasions, much inforem mation and many references.

Asperugus (unnamed species from Kew) (Lillaceae) moves against the sun, placed in bothouse.

Dec. 26, 1st circle was made in . . . 5 0

Tunus commusis (Dioscoreacce). A young shoot from a

" 8,5th " " " . . . . 2 30
" 8,6th " " " . . . . 2 30

Lapagerea rosca (Philesiacere), in greenhouse, follows the sun.

March 9, 1st circle was made in . . . 26 15 (shoot young)

10, semiercle , . . . 8 15

11, 9sd circle . . . . 11 0

" 16,5th " " " . . 8 40 when placed in the hothouse; but the next day the shoot remained stationary.

Rexburghia viridiflora (Roxburghiacen) moves against the sun; it completed a circle in about 24 hours.

### (DICOTYLEDONS.

Hamulus Lupulus (Urticaccae) follows the sun. The plant was kept in a room during warm weather.

was kept in a room during warm weather.

April 9, 2 circles were made in . . . 4 16

Aug. 13, 3rd circle was , . . . 2 0

With the Hop a semicircle was performed, in travelling

Ak-bia quinata (Lardizabalacese), placed in hothouse, moves

March 17, 1st circle was made in .

Stauntonia latifolia (Lardizabalaccee), placed in hothouse,

March 28, 1st circle was made in . . 3 30

.. 29, 2nd " "

August 5th, 1st circle was made in about

Stephania rotunda (Menispermscere) moves against the sun.

May 27, 1st circle was made in

June 2, 3rd .. 3. 4th

Thryallis brachystachys (Malpighiacese) moves against the sun :

made a circle in 7 hrs.; on the 20th, moved against the sun onethird of a circle, and then stood still; on the 25th, followed the

Sollus Drummondii (Pittosporacere) moves against the sun;

April 4, 1st circle was made in , 4 25

. 8 0 (very cold day)

., 7,4th Polyonnum dumetorum (Polygonacew). This case is taken

from Dutrochet (p. 299), as I observed, no allied plant: follows the sun. Three shoots, cut off a plant, and placed in water, made circles in 3 hrs. 10 m., 5 hrs. 20 m., and 7 hrs. 15 m.

Wistoria Chinensis (Leguminosco), in greenhouse, moves

May 13, 1st circle was made in . 25, 6th

Phaseolus vulouris (Legruminosm), in greenhouse, moves against

May, 1st circle was made in . . . . 2 0 , 2nd , , , , . .

3rd " " Dipladenia aronkulla (Apocynacem) moves against the sun.

April 18, 1st circle was made in .

May 16, 1st circle was made in .

.. 21, 3rd .. ..

	(DICOTYLEDONS, continued.)	
Ceropegio	t Gardserii (Aselepiadacem) moves against t	the sun.

Shoot more monner	0						IL
Shoot very young, in length	-	1st	circle	Was	performed	in	7
Shoot still young							7
long shoot ,		3rd	29	,,	27 19		6

1st circle was made in 5 hrs, 30 m.

retarding and hastening the revolving movement.

### (Dicotyledons, continued.)

Risea tiliafolia (Convolvulaces) moves against the sun; made four revolutions in 9 hrs.; so that, on an average, each was performed in 2 hrs. 15 m.

Flumdogo resc (Pinnbaginacem) follows the sun. The shoot did not begin to revolve until nearly a pard in height; if then made a fine circle in 10 hrs. 45 m. During the next few days it continued to move, but irregularly. On August 15th the shoot followed, during a period of 10 hrs. 40 m., a long and deeply signed course and then made a brend dilipse. The figure apparently represented three ellipses, each of which averaged 31m; 33 m. 60 its completion.

Jassianus pruciflorum, Bentham (Jasminacese), mores against the sun. A circle was made in 7 hrs. 15 m., and a second rather more quickly.

Clerodendrum Thomsonii (Verbenneum) follows the su

April		. lst	circle	Was	made	in	5	45	(shoot very young)
	14,	2nd	1 ,,	77	. ,,		3	30	
									((directly after th
	18	0.0		ele			5	0	plant was shaker

AO, it scanicated	.,,		1	on being mov
19, 3rd circle "	77	3	0	
90 4th		4 5		

Tecoma jasminoides (Bignoniacem) moves against the su

March 17, 1st circle was made in . 6 30

, 19, 2ad , , , , 7 0 , 22, 3rd , , , , 8 30 (very cold day)

Thunbergia alata (Acanthacere) moves against su

April 14, 1st circle was made in . 3 20

, 18, 2nd , , , , . 2

Adhadota cydonafolia (Acanthacsee) follows the sun. A young

March	14,	1st	circle	was	made	in	3	
22	15,	2nc	1 ,,	.,			3	
23	16,	3rd	١,,		,,,		3	
22	17,	4th	. ,,	19.	19		3	

7. 6th

after a copious water-ing with cold water at

Jan. 24, 1st circle was made in 2 55 the temperature of the 24, 2 circles each at an average of . . .

25, 4th circle was made in 2 25

Lozsa aurantiaca (Lonsneem). Revolutions variable in their

course; a plant which moved against the sun-

June	20, 1st	sirele	was	made in			37
22	20, 2nd	. ,,	2.0	.,,			13
	20, 3rd	**	- 10	19			0
	21, 4th		- 17	37			35

Darwin Online: By permission of the Trustees of the Natural History Museum (London)

## (DICOTYLEDONS, continued.)

Styphanthus elegans (Lossacem) follows the sun.

June 13, 1st circle was made in . . . 1 4

" 13, 2nd " " " . . . 1 1

" 14, 3rd " " " . . . 1 3

" 14, 4th " " " . . . 1 5

" 14, 5th " " " . . . 2

Siphomeris or Lecontes (unnamed sp.) (Cinchonaces) follows the sun.

1. 1. 1. (shoot extremely

May 25, semicircle was made in . 10 27 | (anoot extremely young) | 26, 2st circle | | 10 15 (shoot still young)

" 20, 25c ctreio " " 10 15 (choot still young) " 30, 25d " " " 8 55

June 2, 3rd " " 8 8 11

" 6, 4th " " 6 8

[Taken from the

", 8,6th ", ", 7 20 hothouse, and placed in a room in my house."

", 8 36 hothouse, and placed in a room in my house.

Manattia bicolor (Cinchonaccus), young plant, follows the sun.

July 7, 1st circle was made in , , , 6 18

July 7, 1st circle was made in . . . 6 18 , 8, 2nd , , , , . . . 6 53 ,, 9, 3rd , , , , . . . . 6 30

Louicera brackypoda (Caprifoliacem) follows the sun, kept in a arm room in the house.

April, 1st circle was made in . 9 10 (about)

April, 2nd circle was made in 12 20 j(a distinct shoot, very

24. 3rd ...

In the foregoing Table, which includes twining that the rate at which growth travels or circulates round the axis (on which the revolving movement uniform, as with the Hop, Mikania, Phascolus, &c. The hereafter see a tendril-bearing Passiflora revolving revolution in 1 hr. 30 m., and three revolutions at the average rate of 1 hr. 38 m.; a Convolvulus made two revolutions at the average of 1 hr. 42 m., and Phascolus other hand, some plants take 24 hrs. for a single revolution, and the Adhadota sometimes required 48 hrs.; yet this latter plant is an efficient twiner. Species of the same genus move at different rates, The rate does not seem governed by the thickness of the shoots: those of the Sollya are as thin and flexible as string, but move more slowly than the thick and fleshy shoots of the Ruseus, which seem little fitted for movement of any kind. The shoots of the Wistaria. which become woody, move faster than those of the herbaceous Ipomæa or Thunbergia,

We know that the internodes, whilst still very young, do not acquire their proper rate of movement; hence the several shoots on the same plant may sometimes be seen revolving at different rates. The two or three, or even more, internodes which are first formed above the cotyledons, or above the root-stock of a perennial plant, do not move; they can support them-

A greater number of twiners revolve in a course opposed to that of the sun, or to the hands of a watch, than in the reversed course, and, consequently, the left to right. Occasionally, though rarely, plants of Mohl (p. 125) gives a case in the Leguminosæ, and we have in the table another in the Acanthacere. I have seen no instance of two species of the same genus twining in opposite directions, and such cases must be rare; but Fritz Müller \* states that although Mikania

interesting paper, in which he Darwin Online: By permission of the Trustees of the Natural History

stance if no such cases had occurred, for different individuals of the same species, namely, of Solanum dulcamara (Dutrochet, tom. xix. p. 299), revolve and twine in two directions; this plant, however, is a most a much more curious case: I raised seventeen plants: of these eight revolved in opposition to the sun and ascended from left to right; five followed the sun and ascended from right to left; and four revolved and twined first in one direction, and then reversed their course,\* the petioles of the opposite leaves affording a point d'appui for the reversal of the spire. One of to left, and five turns from left to right. Another plant in the same family, the Scyphanthus elegans, many plants of it, and the stems of all took one turn, or occasionally two or even three turns in one direction, and then, ascending for a short space straight, reversed their course and took one or two turns in an opposite direction. The reversal of the curvature occurred at any point in the stem, even in the middle of an internode. Had I not

max imposable. It would be hardly possible with any plant which seemed above a few feet in height, or which lived in an exposal situation; for the stem could be patled away coally from its support, with but little unwinding; nor could it have adhered at all, had not the internoles soon become moderately right. With leaf-climbers, as we shall soon see, analogous case frequently occur; but these present no difficulty, at the stem is secured by the elasping petioles. In the many other revolving and twining plants

observed by me, I never but twice saw the movement reversed; once, and only for a short space, in Ipomaca jucunda; but frequently with Hibbertia dentata. This plant at first perplexed me much, for I continually observed its long and flexible shoots, evidently well fitted for twining, make a whole, or half, or quarter circle in one direction and then in an opposite direction; consequently, when I placed the shoots string, they seemed as if constantly trying to ascend, but always failed. I then surrounded the plant with a mass of branched twigs; the shoots ascended, and is usual with twining plants. Finally, I surrounded a second plant with many thin upright sticks, and placed it near the first one with twigs; and now both had got what they liked, for they twined up the parallel sticks, sometimes winding round one and sometimes round several; and the shoots travelled

laterally from one to the other pot; but as the plants gree older, some of the shoots twined regularity up thin purjekt stieks. Though the revolving movement was sometimes in one direction and sometimes in the other, the teining was invariably from left to right;\*\* on that the more potent or persistent movement of revolution must have been in opposition to the course of the sun. It would appear that this Hilbertin is adapted both to accord by twining, and to

I have described the above case in some detail, because, as far as I have seen, it is rure to find say special adaptations with twining plants, in which respect they differ much from the more highly organized tendril-leavers. The Solamus delawara, as we shall presently see, can twin only mount demo which are both thin and flexible. Most twining plants are adapted to accord supports of moderate though of different thicknesses. Our English twiners, as far as I have seen, avere twine result twee, excepting the homeyackle (Louisees periodynesses), which I have observed testining up a young besteleres monty 4g inches in diameter. Modi (p. 3d) found that the

<sup>\*</sup> In another genus, namely left; and I once saw a strong to the same ascended a tree about family with Hibbertin, Pritz Müller says (bild, p. 349) that the same manner as so the same twines indifferently over with Lossa."

when placed in a room with the light entering on one side, twine round sticks between 3 and 4 inches in diameter; for this interfered, in a manner presently to be explained, with the revolving movement. In the onen air however, the Phascolus twined round a support of the above thickness, but failed in twining round one 9 inches in diameter. Nevertheless, some twiners of the warmer temperate regions can manage this latter degree of thickness: for I hear from Dr. Hooker that at Kew the Ruscus androgunus has ascended a column 9 inches in diameter; and although a Wistaria grown by me in a small pot tried in vain for weeks to get round a post between 5 and 6 inches in thickness, yet at Kew a plant ascended a trunk above 6 inches in diameter. The tropical twiners, on the other hand, can ascend thicker trees : I hear from Drs. Thomson and Hooker that this is the case with the Butea parvillora, one of the Menispermacere, and with some Dalbergias and other Leguminose,\* This power would be necessary for any species which had to ascend by twining the large trees of a tropical forest; otherwise they would hardly ever be able to reach the light. In our temperate countries it would be injurious to the twining plants which die down every year if

ispermeece. He adds in his letter to me that most of the colimbing plants which there ascend thick trees, are rootclimbers; some bring tendril-

<sup>\*</sup> Fritz Müller atntes (ibid, p. 349) that he saw on one oceasion in the forests of South Brazil a trunk about five feet in circumference spirally ascended by a plant, apparently belonging to the Men.

they could not grow tall enough in a single season to reach the summit and gain the light.

By what means certain twining plants are adapted to ascend only thin team, whilst others can twine round thicker ones, I do not know. It appeared to me probable that twining plants with very long revolving shoots would be able to ascend thick supports; accordingly I placed Corregatia Gardweiti mor a post 6 inches in diameter, but the shoots entirely failed to wind round it; their great length and power of movement merely aid them in finding a distant stem round which to twine. The spherostemans arraners tone is a vigeous tropical twiner; and as it is a very slow revolver, I hought that this latter eieumstance might help it in ascending a thick support; but though it was able to wind round a 6-inch post, if could do this only on the same level or plane, and did not form a spire and thus ascend.

As form differ so much in structure from planersegamin plants, it may be worth while here to show that twining ferms do not differ in their habits from other twining plants. In Lyapolium orientalation the two internoles of the stem (reoperly the racks) which are first formed above the rootstock do not move, the third from the set is a slow recolver; but Lconsider under five revolution, each at the average rate of 5 hm. 55 m; and this represents fairly well the usual rate, taking quick and slow movers, amount

phanerogamic plants. The rate was accelerated by increased temperature. At each stage of growth only the two upper internodes revolved. A line painted along the convex surface of a revolving internode becomes first lateral, then concave, then lateral and ultimately again convex. Neither the internodes nor the petioles are irritable when rubbed. The movement is in the usual direction, namely, in opposition to the course of the sun; and when the stem twines round a thin stick, it becomes twisted on its own axis in the same direction. After the young internodes have twined round a stick, their continued growth causes them to slip a little upwards. If the stick be soon removed, they straighten themselves, and recommence revolving. The extremities of the depending shoots turn upwards, and twine on themselves. In all these respects we have complete identity with twining phanerogamic plants; and the above enumeration may serve as a summary of the leading characteristics of all twining

The power of revolving depends on the general health and vigour of the plant, as he been laboriously above by Palm. But the movement of each separate internols is so internols are internols as in internols as in internols are in a 2 hrs. Some the wave revolved in between 2 hrs. and 2 hrs. Som a constant of the internol 2 hrs. and 2 hrs. Som a Shoots of the Kidney-bean, cut off and placed in water, were similarly retarded, but in a less degree, the greenhouse to my room, or from one part to another of the greenhouse, always stopped the movement for a time; hence I conclude that plants in a state of nature and growing in exposed situations, would not make their revolutions during very stormy weather. A decrease in temperature always caused a considerable retardation in the rate of revolution; but Dutrochet (tom. xvii. pp. 994, 996) has given such precise observations on this head with respect to the common pea that I need say nothing more. When twining plants are placed near a window in a room, the light in some cases has a remarkable power (as was likewise observed by Dutrochet, p. 998, with the pea) on the revolving movement, but this differs in degree with different plants ; thus Ipomora jucunda made a complete circle in 5 hrs. 30 m.; the semicircle from the light taking 4 hrs. 30 m., and that towards the light only 1 hr. Lonicera brachypoda revolved, in a reversed direction to the Ipomara, in 8 hrs.: the semicircle from the light taking 5 hrs. 23 m., and that to the light only 2 hrs. 37 m. From the rate of revolution in all the plants observed by me, being nearly the same during the night and the day, I infer that the action of the light is confined to retarding one semicircle and accelerating the other, so as not to modify greatly the rate of the whole revolution. This action of the light is remarkable, when we reflect how little the leaves are developed on the young and thin revolving internodes. It is all the more remarkable, as botanists believe (Mohl, p. 119) that twining plants are but little sensitive to the action of light.

I will conclude my account of twining plants by giving a few miscellaneous and curious cases. With most twining plants all the branches, however many there may be, go on revolving together; but, according to Mohl (p. 4), only the lateral branches of Tamus elephantipes twine, and not the main stem. On the other hand, with a climbing species of Asparagus, the leading shoot alone, and not the branches, revolved and twined; but it should be stated that the plant was not growing vigorously. My plants of Combretum argenteum and C. purpureum made numerous short healthy shoots; but they showed no signs of revolving, and I could not conceive how these plants could be climbers; but at last C. argenteum put forth from the lower part of one of its main branches a thin shoot, 5 or 6 feet in length, differing greatly in appearance from the previous shoots, owing to its leaves being little developed, and this shoot revolved vigorously and twined. So that this plant produces shoots of two kinds. With Periploca Graca (Palm, p. 43) the uppermost shoots alone twine. Polygonum convolvulus twines only during the middle of the summer (Palm, p. 43, 94); and plants growing vigorously in the autumn show no inclination to

CHAP. I

but Asclepias nigra only "in fertiliori solo incipit confirmed by Palm, p. 41). Asclepias vincetoxicum does not regularly twine, but occasionally does so (Palm, p. 42; Mohl, p. 112) when growing under certain hear from Prof. Harvey, for these plants in their native dry South African home generally grow erect, specimens showing some inclination to curve; but when cultivated near Dublin, they regularly twined up sticks 5 or 6 feet in height. Most Convolvulaceme are excellent twiners; but in South Africa Ipomora alone in Prof. Harvey's collection showing an evident disposition to twine. On the other hand, seedlings raised near Dublin twined up sticks above 8 feet in height. These facts are remarkable; for there can hardly be a doubt that in the dryer provinces of South Africa these plants have propagated themselves for thousands of generations in an erect condition; and yet they have retained during this whole period the innate power of spontaneously revolving and twining, whenever their shoots become elongated under proper conditions of life. Most of the species of Phaseolus are twiners; but certain varieties of the P. multiflorus produce (Léon, p. 681) two kinds of shoots, some upright and thick, and others thin and twining. I have seen striking instances of this curious case of variability in "Fulmer's dwarf forcing-bean," which occasionally produced a single long twining shoot.

Solanum dulcamara is one of the feeblest and poorest of twiners: it may often be seen growing as an upright bush, and when growing in the midst of a thicket merely scrambles up between the branches without twining; but when, according to Dutrochet (tom, xix, p. 299), it grows near a thin and flexible support, such as the stem of a nettle, it twines round it. I placed sticks round several plants, and vertically stretched strings close to others, and the strings alone were ascended by twining. The stem twines indifferently to the right or left. Some others pecies of Solanum, and of another genus, viz. Habrothamnus, belonging to the same family, are described in horticultural works as twining plants, but they seem to possess this faculty in a very feeble degree. We may suspect that the species of these two genera have as yet only partially acquired the habit of twining. On the other hand with Tecoma radicans, a member of a which climbs, like the ivy, by the aid of rootlets, we may suspect that a former habit of twining has been which could hardly be accounted for by changes in the action of the light. There is no difficulty in understanding how a spirally twining plant could graduate into a simple root-climber; for the young internodes of Bignonia Tweedyana and of Hoya carnosa revolve and twine, but likewise emit rootlets which adhere to any fitting surface, so that the loss of twining would be no great disadvantage and in some respects an advantage to these species, as they would then ascend their sumorts in a more direct line.\*

\* Fritz Muller has published climbing plant some interesting facts and views 1866, pp. 57, 6 on the structure of the wood of

# CHAPTER II.

LEAP-CLIMBERS

Plants which cimb by the slat of spontantomy returning any counter peticles—Canadia—Topopolem—Maurorid, procep-polancies moving spontaneously and sensitive to a touch—Belorication— Lopkspersona—intermodes sensitive—Solames, thickening of the clasped peticles—Fearoria—dellowia—Plants which climb by the sid of their produced midribs—Glerioss—Flagellaria— Nepasthes—Summary on leaf-climbers.

We now come to our second class of climbing plants, namely, those which ascend by the aid of irritable or sensitive organs. For convenience' sake the plants in this class have been grouped under two sub-divisions, namely, leaf-climbers, or those which retain their leaves in a functional condition, and tendril-beavers. But these sub-divisions graduate into each other, as we shall see under Coryalais and the Glorious illy.

It has long been observed that several plants elimiby the aid of thir! leaves, either by their pedaction of the by the aid of thir! leaves, either by their pedaction stalley or by their produced unitality; but said this simple fact they have not been described. Thus and Madle class these plants with those despited. Thus and Madle class these plants with those despited to the present but as a leaf i generally said, has at least some advantage of the said of the said of the said of the present leaves the said of th observed, in order to see what amount of difference in the manner of climbing existed within the same genus; and the differences are considerable.

precisely like those of a true twiner, at an average rate, judging from three revolutions, of 3 hrs. 48 m. placed near it; but, after making an open spire of turns in an opposite direction. This was rendered possible by the straight piece between the opposed spires having become rigid. The simple, broad, ovate whilst twining up a vertical stick, no use is made of them. Nevertheless, if the footstalk of a young leaf be rubbed with a thin twig a few times on any side, it will in the course of a few hours bend to that side; seemed to be the most sensitive: but the sensitiveness shall meet with in some of the following species; thus, a loop of string, weighing 1.64 grain (106.2 mg.) and hanging for some days on a young footstalk, produced a scarcely perceptible effect. A sketch is caught hold of two thin branches. A forked twig placed so as to press lightly on the under side of a young ultimately to such an extent that the leaf passed to the opposite side of the stem; the forked stick having been removed, the leaf slowly recovered its former

The young leaves spontaneously and gradually change their position: when first developed the petioles are upturned and parallel to the stem; they then slowly bend downwards, remaining for a short time at right



angles to the stem, and then become so much arched downwards that the blade of the leaf points to the ground with its tip curled inwards, so that the whole enabled to catch hold of any twig with which they ment of the internodes. If this does not happen, they retain their hooked shape for a considerable time, and then bending upwards reassume their original upturned position, which is preserved ever atterwards. The petioles which have clasped any object soon become much thickened and strengthened, as may be seen in the drawing.

Clematis montana .- The long, thin petioles of the leaves, whilst young, are sensitive, and when lightly rubbed bend to the rubbed side, subsequently becoming straight. They are far more sensitive than the petioles of C. glandulosa; for a loop of thread weighing sometimes acted and sometimes did not act. The sensitiveness extends from the blade of the leaf to the stem. I may here state that I ascertained in all cases the weights of the string and thread used and then cutting off measured lengths. The main petioles are not sensitive. A young, inclined shoot (the plant being in the greenhouse) made a large circle opposed to the course of the sun in 4 hrs. 20 m., but the next day, being very cold, the time was 5 hrs. 10 m. A stick placed near a revolving stem was soon struck by the petioles which stand out at right angles, and the revolving movement was thus arrested. The petioles then began, being excited by the contact, to slowly wind round the stick. When the stick was thin, a petiole sometimes wound twice round it. The opposite leaf was in no way affected. The attitude assumed by the stem after the petiole had

column, who throws his arm horizontally round it.
With respect to the stem's power of twining, some remarks will be made under C. calyeina.

(The vis School L. A. shoot made three revolutions

Camatis Sidebil.—A shoot made three revolutions against the sun at an evenge state of 3 hrs. 11 m. The power of twining is like that of the last species. Its leaves are nearly similar in streams and in function, excepting that the sub-petioles of the lateral and terminal leadies are sensitive. A loop of thread, weighing one-eighth of a grain, acted on the main petiols, but not until two or three days had clapsed. The leaves have the remarkable habit of spontaneously swedsing, generally in verifical ellipses, in the same manner, but in a less degree, as will be described under C neirophyliku.

Cleantis ealgeing.—The young shoots are thin and feetible one revoived, describing a broad oval, in 5 hrs. 30 m, and another in 6 hrs. 12 m. They followed the course of the may but the course, if observed long enough, would probably be found to vary in this species, as well as in all the others of the gones. It is a rather better twiner than the two last species: the stem sometimes made two spiral turns round at this side, if free from twigs, it then ran straight up for a space, and revening its course took can or two turns in an opposite direction. This reversal of the spire occurred in all the foregoing species. The leaves are so small compared with those of most of the other species, that the petidos at dirts seen ill-shaunted for cleaning. ment is to bring them into contact with surrounding objects, which are slowly but securely seized. The young petioles, which alone are sensitive, have their twig the lower surfaces of two young petioles; and in 2 hrs. 30 m. they were slightly curved downwards; in 5 hrs., after being rubbed, the end of one was bent subsequently it became nearly straight again. To show how sensitive the young petioles are, I may mention that I just touched the under sides of two an excessively thin and minute crust; but this sufficed in 24 hrs. to cause both to bend downwards. Whilst the plant is young, each leaf consists of three divided leaflets, which barely have distinct petioles, and these are not sensitive; but when the plant is well grown, the petioles of the two lateral and terminal leaflets are of considerable length, and become sensitive so as to be capable of clasping an object in any

When a petiole has clasped a twig, it undergoes some remarkable changes, which may be observed with the other species, but in a less strongly marked manner, and will here be described once for all. The clasped petiole in the course of two or three days wells greatly, and ultimately becomes nearly (wice as thick as the opposite one which has clasped nothing. When thin transverse slices of the two are placed under the microscope their difference is conspicuous: the side of the petiole which has been in contact with the support, is formed of a layer of colourless cells with their longer axes directed from the centre, and these are very much larger than the corresponding cells in the opposite or unchanged petiole; the central cells, also, are in some degree enlarged, and the whole is much indurated. The exterior surface generally becomes bright red. But a far greater change takes place in the nature of the tissues than that which is visible; the petiole of the unclasped leaf is flexible and can be snapped easily, whereas the clasped one acquires an extraordinary degree of toughness and rigidity, so that considerable force is required to pull it into pieces. With this change, great durability is probably acquired; at least this is the case with the clasped petioles of Clematis vitalba. The meaning of these changes is obvious, namely, that the petioles may

firmly and durably support the stem. Clematis microphylla, var. leptophylla.-The long and thin internodes of this Australian species revolve sometimes in one direction and sometimes in an opposite one, describing long, narrow, irregular ellipses or large circles. Four revolutions were completed within five minutes of the same average rate of 1 hr. 51 m.; so that this species moves more quickly than the others of the genus. The shoots, when placed near a vertical stick, either twine round it, or clasp it with the basal portions of their petioles. The leaves whilst young are nearly of the same shape as those of C. witeelds, and act in the same namer like a hook, as will be described under that species. But the leafest are more divided, and each segment whilst young terminates in a hardish point, which is much curved downwards and inwards; so that the whole leaf resulty catches hold of any neighbouring object. The petioles of the young terminal leafest are acted on by loops of thread weighing 4th and even 4th of a gmin. The basal portion of the main petiole is much less sensitive, but will clasp a stick against which is presses.

The laws, whilst young, are continually and spenish monthly nowing donly. A bullginess was placed vere a shoot secured to a stick, and the mereaments of the leaves were traced on it during several days. A very irregular line was generally formed; but one day, in the course of eight hours and three quarters, the figure clearly represented three and a balf irregular chipses, the most perfect one of which was completed in 2 lms. 35 m. The two opposite leaves mored independently of each other. This movement of the leaves would aid that of the internodes in bringing the petides into contact with surrounding objects. I discovered this movement too late to be enabled to observe it in the other species; but from analogy? I discovered the movement of the contact with surface and the petides into contact with surface and the petides in the context of the contact with surface and the petides in the context of the contact with surface and the petides in the context of the context o

C. montana and C. calycina. I ascertained that the simple leaves of C. glandulosa exhibited no sponta-

Clematis viticella, var. venosa,- In this and the two following species the power of spirally twining is completely lost, and this seems due to the lessened flexibility of the internodes and to the interference caused by the large size of the leaves. But the revolving movement, though restricted, is not lost. In our present species a young internode, placed in front of a window, made three narrow ellipses, transversely to the direction of the light, at an average rate of 2 hrs. 40 m. When placed so that the movements were to and from the light, the rate was greatly accelerated in one half of the course, and retarded in the other, as with twining plants. The ellipses were small; the longer diameter, described by the apex of a shoot bearing a pair of not expanded leaves, was only 45 inches, and that by the apex of the penultimate internode only 11 inch. At the most favourable period of growth each leaf would hardly be carried to and fro by the movement of the internodes more than two or three inches, but, as above stated, it is probable that the leaves themselves move spontaneously. The movement of the whole shoot by the wind and by its rapid growth, would probably be almost equally efficient as these spontaneous movements, in bringing the petioles into contact with surrounding objects,

The leaves are of large size. Each bears three pairs of lateral leaflets and a terminal one, all supported on rather long sub-petioles. The main petiole bends a little angularly downwards at each point where a pair of leaflets arises (soo fig. 2), and the petiole of the terminal leaflet is bent downwards at right angles; benece the whole petiole, with its rectangularly bent extremity, acts as a hook. This hook, the lateral and the petion of the period of the period of the period of the period and the period of th

excellent grappling apparatus, by which the leaves



readily become entangled with surrounding objects.
If they catch nothing, the whole petiols ultimately
grows straight. The main petiols, the subpetiols,
and the three branches into which each basi-lateral
sub-petiole is generally subdivided, are all sensitive
The basal portion of the main petiole, between the
stem and the first pair of leaflets, is less sensitive
than the remainder; it will, however, clasp a side

with which it is left in contact. The inferior surface of the rectangularly bent terminal portion (carrying the terminal leaflet), which forms the inner side of the end of the hook, is the most sensitive part; and this portion is manifestly best adapted to catch a distant support. To show the difference in sensibility, I gently placed loops of string of the same weight (in on the several lateral sub-petioles and on the terminal one: in a few hours the latter was bent, but after 24 hrs. no effect was produced on the other sub-petioles. Again, a terminal sub-petiole placed in contact with a thin stick became sensibly curved in 45 m., and in 1 hr. 10 m. moved through ninety degrees; whilst a lateral sub-petiole did not become sensibly curved until 3 hrs. 30 m. had elapsed. In all cases, if the sticks are taken away, the petioles continue to move during many hours afterwards; so they do after a slight rubbing : but they become straight again, after about a day's interval, that is if the flexure has not

The graduated difference in the extension of the sensitiveness in the petioles of the above-described species deserves notice. In  $C_i$  montras it is confined to the main petiole, and has not spread to the subpetioles of the three callests; so it swith young plasts of  $C_i$  collection, the three callests of the subtered in  $C_i$  models and  $C_i$  and LEAF-CLIMBERS.

this latter species it has diminished in the basal part of the main petiole, in which alone it resided in G. montana; whilst it has increased in the abruptly bent terminal portion.

Clematis flammula,-The rather thick, straight, and stiff shoots, whilst growing vigorously in the spring, make small oval revolutions, following the sun in their course. Four were made at an average rate of 3 hrs. 45 m. The longer axis of the oval, described by the extreme tip, was directed at right angles to the line joining the opposite leaves; its length was in one case only 18, and in another case 15 inch; so that the young leaves were moved a very short distance. The shoots of the same plant observed in midsummer, when growing not so quickly, did not revolve at all. I cut down another plant in the early summer, so that by August 1st it had formed new and moderately vigorous shoots; these, when observed under a bellglass, were on some days quite stationary, and on other days moved to and fro only about the eighth of an inch. Consequently the revolving power is much enfeebled in this species, and under unfavourable circumstances is completely lost. The shoot must depend for coming into contact with surrounding objects on the probable, though not ascertained spontaneous movefrom the wind. Hence, perhaps, it is that the petioles pensation for the little power of movement in the shoots.

The petioles are bowed downwards, and have the

same general hook-like form as in C. viticella. The especially the much bent terminal portion. As the sensitiveness is here greater than in any other species of the genus observed by me, and is in itself remarkable, I will give fuller details. The petioles, when so young that they have not separated from one another, are not sensitive; when the lamina of a leaflet has grown to a quarter of an inch in length (that is, about but at this period the petioles are relatively much more fully developed than are the blades of the leaves. Full-grown petioles are not in the least sensitive. A thin stick placed so as to press lightly against a petiole, having a leaflet a quarter of an inch in length, caused the petiole to bend in 3 hrs. 15 m. In another case a petiole curled completely round a stick in 12 hrs. These petioles were left curled for 24 hrs., and the sticks were then removed; but they never the petiole itself, and with it lightly rubbed several petioles four times up and down; these in 1 hr. 45 m. some hours and then began to decrease, but after 25 hrs. in about 2 hrs. 30 m., the terminal sub-petiole moving more than the lateral sub-petioles; they all became straight again in between 12 hrs, and 14 hrs. Lastly, a was lightly rubbed with the same twig only once; it

The following observations are more precise. After trying heavier pieces of string and thread, I placed a loop of fine string, weighing 1-04 gr. (67-4 mg.) on a string, and in 72 hrs, seized it so firmly, that some force was necessary for its withdrawal. A loop weighing '52 of a grain (33.7 mg.) caused in 14 hrs, a lateral subpetiole just perceptibly to curve, and in 24 hrs. it made during the summer: the following were made in the spring, when the petioles apparently are more sensitive :- A loop of thread, weighing one-eighth of a petioles, but placed on a terminal one, caused it, after 24 hrs., to curve moderately; the curvature, though the loop remained suspended, was after 48 hrs. diminished, but never disappeared; showing that the petiole had become partially accustomed to the insufficient stimulus. This experiment was twice repeated with nearly the same result. Lastly, a loop of thread, weighing only one-sixteenth of a grain (405 mg.) was twice gently placed by a forceps on a terminal sub-petiole (the plant being, of course, in a still and closed room), and this weight certainly caused a flexure, which very slowly increased until the petiole moved through nearly ninety degrees: beyond this it did not move; nor did the petiole, the loop remaining suspended, ever become perfectly straight again.

When we consider, on the one hand, the thickness and stiffness of the petioles, and, on the other hand, the thinness and softness of fine cotton thread, and what an extremely small weight one-sixteenth of a grain (405 mg.) is, these facts are remarkable. But I have reason to believe that even a less weight excites curvature when pressing over a broader surface than that acted on by a thread. Having noticed that the end of a suspended string which accidentally touched a petiole, caused it to bend, I took two pieces of thin twine, 10 inches in length (weighing 1.64 gr.), and, tying them to a stick, let them hang as nearly perpendicularly downwards as their thinness and flexuous form, after being stretched, would permit: I then quietly placed their ends so as just to rest on two petioles, and these certainly became curved in 36 hrs. One of the ends touched the angle between a terminal and lateral sub-petiole, and it was in 48 hours caught between them as by a forceps. In these cases the pressure, though spread over a wider surface than that touched by the cotton thread, must

nave been excessively signt.

Clematis vitalita.—The plants were in pots and not healthy, so that I dare not trust my observations, which indicate much similarity in habits with C. flammula. It mention this species only because I have seen many

proofs that the petioles in a state of nature are excited I have found them embracing thin withered blades of grass, the soft young leaves of a maple, and the latter are about as thick as the hair of a man's hours with some object subsequently removed. With none of the above-described species, cultivated in pots and carefully observed, was there any permanent bending of the petioles without the stimulus of contact. In winter, the blades of the leaves of C. vitalba drop off: but the petioles (as was observed by Mohl) remain attached to the branches, sometimes during two seasons; and, being convoluted, they curiously resemble true tendrils, such as those possessed by clasped some object become much more stiff, hard, and polished than those which have failed in this their

Trop.golum.—I observed T. tricolorum, T. azureum,
T. pentaphyllum, T. peregrinum, T. elegans, T. tuberosum,
and a dwarf variety of, as I believe. T. minus.

Tropecolum tricolorum, var. grandiflorum.—The flexible shoots, which first rise from the tubers, are as thin as fine twine. One such shoot revolved in a course opposed to the sun, at an average rate, judging from three revolutions, of 1 hr. 23 m.; but no doubt the direction of the revolving movement is variable, When the plants have grown tall and are branched, all the many lateral shoots revolve. The stem, whilst young, twines regularly round a thin vertical stick. and in one case I counted eight spiral turns in the same direction; but when grown older, the stem often runs straight up for a space, and, being arrested by the clasping petioles, makes one or two spires in a of two or three feet, requiring about a month from the time when the first shoot appears above ground, no true leaves are produced, but, in their place, filaments coloured like the stem. The extremities of these on the upper surface. They never become developed into leaves. As the plant grows in height new filaments are produced with slightly enlarged tips; then others, bearing on each side of the enlarged medial tip appear, and at last a perfect leaf is formed, with seven deep segments. So that on the same plant we may see to a considerable height, and is secured to its support by the petioles of the true leaves, the clasping filaments on the lower part of the stem wither and drop off; so that they perform only a temporary service.

These filaments or rudimentary leaves, as well as the petioles of the perfect leaves, whilst young, are rub caused them to curve towards the rubbed side in about three minutes, and one bent itself into a ring in six minutes; they subsequently became straight, When, however, they have once completely clasped a stick, if this is removed, they do not straighten themselves. The most remarkable fact, and one which I have observed in no other species of the genus, is that the filaments and the petioles of the young leaves, if they catch no object, after standing for some days in their original position, spontaneously and slowly oscillate a little from side to side, and then move towards the stem and clasp it. They likewise often become, after a time, in some degree spirally contracted. They therefore fully deserve to be called tendrils, as they are used for climbing, are sensitive to a touch, move spontaneously, and ultimately contract into a spire, though an imperfect one. The present species would have been classed amongst the tendril-bearers, had not these characters been confined to early youth. During maturity it is a true leaf-climber.

Tropwolum azureum.—An upper internode made four revolutions, following the sun, at an average rate of 1 hr. 47 m. The stem twined spirally round a support in the same irregular manner as that of the last species. Rudimentary leaves or filaments do not exist. The petioles of the young leaves are very sensitive: a single light rub with a twig caused one to move perceptibly in 5 m., and another in 6 m. became straight again in between 5 hrs. and 6 hrs. A loop of thread weighing 1th of a grain caused another

Tropxolum pentaphyllum,-This species has not the to a want of flexibility in the stem, as to continual interference from the clasping petioles. An upper internode made three revolutions, following the sun, at an average rate of 1 hr. 46 m. The main purpose of the revolving movement in all the species of Troprolum manifestly is to bring the petioles into contact with some supporting object. The petiole of a young leaf, after a slight rub, became curved in 6 m.; another, on a cold day, in 20 m., and others in from 8 m. to 10 m. Their curvature usually increased greatly in from 15 m, to 20 m., and they became straight again in between 5 hrs. and 6 hrs., but on one occasion in 3 hrs. When a petiole has fairly clasped a stick, it is not able, on the removal of the stick, to straighten itself. The free upper part of one, the base of which had already clasped a stick, still retained the power of movement, A to curve; but the stimulus was not sufficient, the loop remaining suspended, to cause a permanent flexure. If a much heavier loop be placed in the angle between the petiole and the stem, it produces no effect : whereas we have seen with Clematis montana that the angle between the stem and petiole is sensitive.

Tropvolum peregrinum.—The first-formed internodes respect those of a twining plant. In an older plant the four upper internodes made three irregular revolutions, in a course opposed to the sun, at an average average rate of revolution (taken, however, but from few observations) is very nearly the same in this and the two last species, namely, 1 hr. 47 m., 1 hr. 46 m., and 1 hr. 48 m. The present species cannot twine spirally, which seems mainly due to the rigidity of the stem. In a very young plant, which did not revolve, the petioles were not sensitive. In older plants the petioles of quite young leaves, and of leaves as much as an inch and a quarter in diameter, are sensitive. A moderate rub caused one to curve in 10 m., and others in 20 m. They became straight again in between 5 hrs. 45 m. and 8 hrs. Petioles which have naturally come into contact with a stick, sometimes take two turns round it. After they have clasped a support, they become rigid and hard. They are less sensitive to a weight than in the previous species; for loops of string weighing 82 of a grain (53:14 mg.), did not cause any curvature, but a loop of double this weight (1-64 gr.) acted.

Tropeolum elegans.—I did not make many observations on this species. The short and stiff internoles revolve irregularly, describing small oval figures. One oval was completed in 3 hrs. A young petiols, when rubbed, became slightly curved in 17 m.; and

afterwards much more so. It was nearly straight again

Tropwolum tuberosum.-On a plant nine inches in height, the internodes did not move at all; but on an older plant they moved irregularly and made small imperfect ovals. These movements could be detected only by being traced on a bell-glass placed over the plant. Sometimes the shoots stood still for hours; during some days they moved only in one direction in a crooked line; on other days they made small irregular spires or circles, one being completed in about 4 hrs. The extreme points reached by the apex of the shoot were only about one or one and a half inches asunder; yet this slight movement brought the petioles into contact with some closely surrounding of spontaneously revolving, compared with that of the previous species, the sensitiveness of the petioles is also diminished. These, when rubbed a few times, did not become curved until half an hour had elapsed; the curvature increased during the next two hours, and then very slowly decreased; so that they sometimes required 24 hrs. to become straight again. Extremely young leaves have active petioles; one with the lamina only '15 of an inch in diameter, that is, about a twentieth of the full size, firmly clasped a thin twig. But leaves grown to a quarter of their

Tropwolum minus (?).-The internodes of a variety

named "dwarf crimson Nasturtium" did not revolve,

Darwin Online: By permission of the Trustees of the Natural History

but moved in a rather irregular course during the day to the light, and from the light at night. The ing; nor could I see that they ever clasped any neighbouring object. We have seen in this genus a gradation from species such as T. tricolorum, which rapidly revolve and spirally twine up a support, to other species such as T. elegans and T. tuberosum, the ternodes of which have very feeble revolving powers and cannot spirally twine round a support, to this last

the loss of power seems the more probable alternative. others, the flower-peduncle, as soon as the seed-capsule begins to swell, spontaneously bends abruptly downstands in the way, it is to a certain extent clasped; but, as far as I have been able to observe, this clasping movement is independent of the stimulus from contact.

faculties. From the general character of the genus,

ANTIRRHINE E .- In this tribe (Lindley) of the genera have leaf-climbing species.

Maurandia Barelayana .- A thin, slightly bowed 3 hrs. 17 min.; on the previous day this same shoot revolved in an opposite direction. The shoots do not twine spirally, but climb excellently by the aid of their young and sensitive petitors. These petitors, when lightly rubbed, move after a considerable interval of time, and subsequently become straight again. A loop of thread weighing 1th of a grain caused them to bend.

Maurandia semperflorens.—This freely growing sensitive petioles. A young internode made two twice as rapidly as the last species. The internodes genus, namely, Lophospermum. The present species is unique in one respect. Mohl asserts (p. 45) that "the flower-peduncles, as well as the petioles, wind like tendrils;" but he classes as tendrils such objects as the spiral flower-stalks of the Vallisneria. This remark, and the fact of the flower-peduncles being decidedly flexuous, led me carefully to examine placed thin sticks in contact with young and old grow through an entangled mass of branches; but in no one instance did they bend round any object. branches which have already securely clasped a support by the petioles of their leaves; and when borne on a free depending branch, they are not produced by the terminal portion of the internode 68

which alone has the power of revolving; so that they could be brought only by accident into contact with any neighbouring object. Nevertheless (and this is the remarkable fact) the flower-peduncles, whilst sensitive to a touch. Having selected some stems which had firmly clasped a stick by their petioles, and having placed a bell-glass over them, I traced the movements of the young flower-peduncles. The tracing generally formed a short and extremely irregular line, with little loops in its course. A young peduncle 11 inch in length was carefully observed during a whole day, and it made four and a half narrow, vertical, irregular, and short ellipses-each at an average rate of about 2 hrs. 25 m. An adjoining peduncle described during the same time similar, though fewer, ellipses. As the plant had occupied for some time exactly the same position, these movements could not be attributed to any change in the action of the light. Peduncles, old enough for the coloured petals to be just visible, do not move. With respect to irritability,\* I rubbed two young peduncles (11 inch in length) a few times very lightly with a thin twig; one was rubbed on the upper, and the other on the lower side, and they became in between 4 hrs. and 5 hrs. distinctly bowed towards

<sup>\*</sup> It appears from A. Kerner's interesting observations, that the flower-peduncles of a large number 1873, p. 34.

Darwin Online: By permission of the Trustees of the Natural History

themselves. Next day they were rubbed on the opposite sides, and they became perceptibly curved towards these sides. Two other and younger peduncles (three-fourths of an inch in length) were came so much curved towards one another, that the ares of the bows stood at nearly right angles to their seen by me. Subsequently they straightened themselves. Other peduncles, so young as to be only three-tenths of an inch in length, became curved when in length required to be rubbed two or three times, of thread suspended on the peduncles produced no effect; loops of string, however, weighing 82 and 164 of a grain sometimes caused a slight curvature; but they were never closely clasped, as were the far lighter loops of thread by the petioles.

In the nine vigcous plants observed by me, it is certain that neither the slight spontaneous movements nor the slight sensitiveness of the flower-pedundes added the plants in elimbing. If any member of the Scrophulariacow had possessed tendrils produced by the modification of flower-pedunes, I should have thought that this species of Maurendia had perhaps retained a noise or rullimentary vestige of a former patient of the produced of the produced by the protained a noise or rullimentary vestige of a former habit; but this view cannot be maintained. We may suspect that, owing to the principle of correlation. the power of movement has been transferred to the flower-peduneds from the young internodes, and sustiveness from the young petioles. But to whatever, cause these capacities are due, the case is internaing; for, by a little increase in power through natural selection, they might easily have been rendered as useful to the plant in climbing, as are the flowerpedundes (hereafter to be described) of Vitis or Cardiopersum.

in 4 hrs. 10 m. The shoots sometimes make a whole or a half spire round a vertical stick, they then run straight up for a space, and afterwards turn spirally in an opposite direction. The petioles of very young leaves about one-tenth of their full size, are highly sensitive, and bend towards the side which is touched; but they do not move quickly. One was perceptibly curved in 1 hr. 10 m., after being lightly rubbed, and became considerably curved in 5 hrs. 40 m.; some others were scarcely curved in 5 hrs. 30 m., but distinctly so in 6 hrs. 30 m. A curvature was perceptible in one petiole in between 4 hrs. 30 m, and 5 hrs., after the suspension of a little loop of string. A loop of fine cotton thread, weighing one sixteenth of a grain (4.05 mg.), not only caused a petiole slowly to bend, but was ultimately so firmly clasped that it could be withdrawn only by some little force. The petioles, when coming into contact with a stick, take either a complete or half a turn round it, and ultimately increase much in thickness. They do not possess the power of spontaneously revolving.

Lephopersons condens, var. purparame.—Some long melectably thin internodes made four revolutions at an average rate of 3 hrs. 15 m. The course pursaed was very irregular, namely, an extremely, an extremely narrow cellipse, a large circle, an irregular spire or a signed line, and sometimes the apex stood still. The young petioles, when brought by the revolving movement into contact with attless, clasped them, and soon increased considerably in thickness. But they are not quite so sensitive to a weight as those of the Bloeds-to ton, for loops of thread weighing one-eighth of a grain did not attays; cause them to bend.

This plant presents a case not observed by no in any other leaf-climber or twiner,\* namely, that the young internoise of the stem are sensitive to a tonch. When a petiols of this species chaps a stick, it draws the laws of the internoise against it; and their it draws the laws of the internoise lastick, which is caught between the stem and the peticle as by a pair of pincers. The internoise afterwards straightens itself, excepting the part in actual contact with the stick. Young internoise alone are sensitive, and these are sensitive, and these are sensitive, and these are

<sup>\*</sup> I have already referred to the Vries (ibid. p. 322) is sensitive to case of the twining stem of Cuscuta, which, according to H. de

fifteen trials by twice or thrice lightly rubbing with a thin twig sevenal internodes; and in about 2 has, has in one case in 3 hrs, all were bent; they became straight again in about 4 hrs, afterwards. An internode, which was rubbed as often as six or seven times, became just perseptibly curved in 1 hrs. 15 m., and in 3 hrs. the curvature increased much; it became straight again in the course of the succeeding injulin 1 hrs. 15 m. and in the curvature increased much; it became straight again in the course of the succeeding injult rubbed now. Interno. The succeeding injulty of the course of the succeeding injulangles to the first side; and the curvature as a keys towards the rubbed side.

According to Palm (p. 63), the petioles of Lincais

cirrhosa and, to a limited degree, those of L. elatine
have the power of clasping a support.

SOLEMENT Schmus juminoides.—Some of the species in this large genus are twiners; but the present species in a true leaf-climber. A long, nearly upight shoot made four revolutions, merity against the sun, very regularly at an average rate of 3 hrs. 25 m. The shoots, however, sometimes stood stills. It is considered a greenhouse plant; but when kept there, the petidois took several days to clasp a attick; in the hothouse a stick was clearly in 17 hrs. In the greenhouse a petidois was not affected by a loop of string, on the control of the string grains (165 mg.); but in the bothouse one was made to curvely a loop weighing [1-94] gr. (10027 mg.); and, on the removal of the string, it became straight again. Another petidois was not at affected on by a loop

weighing only 82 of a grain (53.14 mg.) We have seen that the petioles of some other leaf-climbing plants are affected by one-thirteenth of this latter weight. In this species, and in no other leaf-climber seen by me, a full-grown leaf is capable of clasping a stick; but in the greenhouse the movement was so extraordinarily



Fig. 2.
Solamann farminaides, with one of its meticles elastring a ti

slow that the act required several weeks; on each succeeding week it was clear that the petiole had become more and more curved, until at last it firmly clasped the stick.

The flexible petiole of a half or a quarter grown leaf which has elasped an object for three or four days increases much in thickness, and after several weeks becomes so wonderfully hard and rigid that it can hardly be removed from its support. On comclasped anything, its diameter was found to be fully other petioles similarly compared, and here represented. the increase in diameter was not quite so great. In the section of the petiole in its ordinary state (A), we see a semilunar band of cellular tissue (not





well shown in the woodcut) differing slightly in

appearance from that outside it, and including three closely approximate groups of dark vessels. Near the upper surface of the petiole, beneath two exterior ridges, there are two other small circular groups of vessels. In the section of the petiole (B) which had clasped during several weeks a stick, the two exterior ridges have become much less prominent, and the two groups of woody vessels beneath them much increased in diameter. The semilunar band has been converted into a complete ring of very hard, white, woody tissue, with lines radiating from the centre. The three groups of vessels, which, though near together, were before distinct, are now completely blended. The upper part of this ring of woody vessels, formed by the prolongation of the horns of the original semilunar band, is narrower than the lower part, and slightly less compact. This petiole after clasping the stick had actually become thicker than the stem from which it arose; and this was chiefly due to the increased thickness of the ring of wood. This ring presented, both in a transverse and longitudinal section, a closely similar structure to that of the stem. It is a singular morphological fact that the petiole should thus acquire a structure almost identically the same with that of the axis; and it is a still more singular physiological fact that so great a change should have been induced by the mere act

FUMARIACE.E.—Fumaria officinalis.—It could not have been anticipated that so lowly a plant as this Fumaria should have been a climber. It climbs by the aid of the main and lateral petioles of its compound leaves; and even the much-flattened terminal

statement, it may be observed that the enlarged and clasped petiole of the Schwass, with its closed ring of woody vessels, has become more cylindrical than it was in its original unclasped condition

Tr. Maxwell Masters inform me that in almost all petiole which are cylindrical, such a those bearing peltate leaves, the woody vessels form a closed ring semilumar bands of vessels bein confined to petioles which as

portion of the petiole can seize a support. I have caught. Petioles which have clasped any object ultimately become rather thicker and more evlindrical. On lightly rubbing several petioles with a twig, they became perceptibly curved in 1 hr. 15 m., and subsequently straightened themselves. A stick gently placed in the angle between two sub-petioles excited them to move, and was almost clasped in 9 hrs. A loop of thread, weighing one-eighth of a grain, caused, after 12 hrs, and before 20 hrs, had elapsed, a considerable curvature; but it was never fairly clasped by the petiole. The young internodes are in continual movement, which is considerable in extent, but very irregular; a zigzag line, or a spire crossing itself, or a figure of 8 being formed. The course during 12 hrs., when traced on a bell-glass, apparently represented about four ellipses. The leaves themselves likewise move spontaneously, the main petioles curving themselves in accordance with the movements of the internoles; so that when the latter moved to one side, the petioles moved to the same side, then, becoming straight, reversed their curvature. The petioles, however, do not move over a wide space, as could be seen when a shoot was securely tied to a stick. The leaf in this case followed an irregular course, like that made by

Adlumia cirrhosa.—I raised some plants late in the summer; they formed very fine leaves, but threw up no central stem. The first-formed leaves were not sensitive; some of the later ones were so, but only towards their extremities, which were thus enabled to clasp sticks. This could be of no service to the plant, as these leaves rose from the ground; but it showed what the future character of the plant would have been, had it grown tall enough to climb. The tip of one of these basal leaves, whilst young, described in 1 hr. 36 m. a narrow ellipse, open at one end, and exactly three inches in length; a second ellipse was broader, more irregular, and shorter, viz., only 21 inches in length, and was completed in 2 hrs. 2 m. From the analogy of Fumaria and Corudalis, I have no doubt that the internodes of Adlumia have the power

Corydalis claviculata.—This plant is interesting from being in a condition so exactly intermediate between a leaf-climber and a tendril-bearer, that it might have been described under either head; but, for reasons hereafter assigned, it has been classed

Besides the plants already described, Bignonia unguis and its close allies, though aided by tendrils, have clasping petioles. According to Mohl (p. 40), Cocculus Japonicus (one of the Menispermaceæ) and a fern, the Ophioglossum Japonicum (p. 39), climb by

We now come to a small section of plants which climb by means of the produced midribs or tips of

LILIACE.E. Gloriosa Plantii. The stem of a halfgrown plant continually moved, generally describing an irregular spire, but sometimes oval figures with the longer axes directed in different lines. It either followed the sun, or moved in an opposite course, and sometimes stood still before reversing its direction. One oval was completed in 3 hrs. 40 m.; of two horseshoe-shaped figures, one was completed in movements, reached points between four and five inches asunder. The young leaves, when first developed, stand up nearly vertically; but by the growth of the axis, and by the spontaneous bending down of the terminal half of the leaf, they soon become much inclined, and ultimately horizontal. The end of the leaf forms a narrow, ribbon-like, but by the time the leaf gets into an inclined position, the end bends downwards into a well-formed hook. This hook is now strong and rigid enough to catch any object, and, when caught, to anchor the plant and stop the revolving movement. Its inner surface is of the many before-described petioles; for a loop of When the hook has caught a thin twig or even a rigid fibre, the point may be perceived in from 1 hr. to 3 hrs. to have curled a little inwards; and, under favourable circumstances, it curls round and permanently seizes an object in from 8 hrs. to 10 hrs.

The book when first formed, before the leaf has best downwards, is but little easility. If it eathers hold of nothing, it remains open and sensitive for a long time; ultimated the extremity spontaneously and slovly curls invarie, and makes a botton-like, and, spiral coil at the end of the leaf. One leaf was watched, and the hook remained open for thirtythree days; but during the last week the tip had carried so much invaried that only a very thin twig could have been inserted within it. As soon as the tip has curled so much invaried that the hook is converted into a ring, its sensibility is lost; but as long as it munitudes ones sensibility is retained.

Whilst the plant was only about six inches in height, the leaves, four or five in number, were breader than those subsequently produced; their soft and but little-attenuated tips were not sensitive, and did not form hooks; nor did the stem then revolve. At this early period of growth, the plant can support itself; its climbing powers are not required, and consequently are not developed. So again, the leaves on the summit of a full-grown flowering plant, which would not require to climb any higher, were not sensitive and could not clasp a stick. We thus see how perfect is the economy of nature.

COMMELYNACE.—Flagellaria Indica.—From dried specimens it is manifest that this plant climbs exactly like the Gloriosa. A young plant 12 inches in height, and bearing fifteen leaves, had not a single leaf as yet produced into a hook or tendril-like filament: nor did the stem revolve. Hence this plant acquires its elimbing powers later in life than does the Gloriess lily. According to Mohl (p. 41), Uvularia (Melanthacce) also climbs like Gloriesa.

These three last-named genera are Monocotyledons: is ranked by Mohl (p. 41) amongst tendril-bearers; and I hear from Dr. Hooker that most of the species climb well at Kew. This is effected by the stalk or midrib between the leaf and the pitcher coiling round any support. The twisted part becomes thicker; but I observed in Mr. Veitch's hothouse that the stalk often takes a turn when not in contact with any object, and that this twisted part is likewise thickened. toria, in my hothouse, whilst less than a foot in height, showed no sensitiveness in their leaves, and had no power of climbing. But when N. levis had grown to a height of 16 inches, there were signs of these powers. The young leaves when first formed stand upright, but soon become inclined; at this period they terminate in a stalk or filament, with the pitcher at the extremity hardly at all developed. The leaves now exhibited slight spontaneous movements; and when the terminal filaments came into contact with a stick, they slowly bent round and firmly seized it. But owing to the subsequent quite slack, though still remaining firmly coiled round the stick. Hence it would appear that the

chief use of the coiling, at least whilst the plant is young, is to support the pitcher with its load of secreted fluid.

Summary on Leaf-climbers,-Plants belonging to plants belonging to four families climb by the tips of their leaves. In all the species observed by me. with one exception, the young internodes revolve more or less regularly, in some cases as regularly as those of a twining plant. They revolve at various rates, in most cases rather rapidly. Some few can ascend most twiners, there is a strong tendency in the same shoot to revolve first in one and then in an opposite direction. The object gained by the revolving movement is to bring the petioles or the tips of the leaves aid the plant would be much less successful in climbing. With rare exceptions, the petioles are sensitive only whilst young. They are sensitive on all sides, but in different degrees in different plants; and in some species of Clematis the several parts of the same petiole differ much in sensitiveness. The hooked their inner or inferior surfaces. The petioles are sensitive to a touch and to excessively slight continued pressure, even from a loop of soft thread weighing only the one-sixteenth of a grain (4.05 mg.); and there is reason to believe that the rather thick and

stiff petioles of Clematis flammula are sensitive to even

or touched, at different rates in different species. a much longer period. After temporary contact with any object, the petiole continues to bend for a considerable time; afterwards it slowly becomes straight again, and can then re-act. A petiole excited by an then becomes accustomed to the stimulus, and either an object for some little time cannot recover their original position. After remaining clasped for two or three days, they generally increase much in thickness either throughout their whole diameter or on one side alone; they subsequently become stronger and more cases they acquire an internal structure like that of the stem or axis.

The young internodes of the Lophospermum as well as the petioles are sensitive to a touch, and by their combined movement seize an object. The flowerpeduncles of the Maurandia semperflorens revolve spontaneously and are sensitive to a touch, yet are not used for climbing. The leaves of at least two, and and Adlumia, spontaneously curve from side to side, like the internodes, and are thus better adapted to seize distant objects. The petioles of the perfect leaves of Teppedant riviolerus, as vell as the tendrillike filaments of the plants whilst young, ultimately move towards the stem or the supporting stick, which they then clasp. These petioles and filaments also show some tendency to contract spirally. The tips of the unempth leaves of the Glorious, as they grow old, contract into a flat spire or helix. These several facts are interesting in relation to true tendrils.

With leaf climbers, as with twining plants, the first internodes which rise from the ground do not, at clear in the cases observed by me, spontaneously revolve; now are the petiodes or tips of the first-formed leaves sensitive. In certain species of Cleantis, the large size of the leaves, together with their habit of revolving, and the extreme sensitiveness of their petioles, appear to reader the revolving movement of the internoless superfluous; and this latter power has consequently become much enhedded. In certain species of To-peolous, both the spontaneous movements of the internoles and the sensitiveness of the petioles have become much enfeebled, and in one species have been completely lear.

## CHAPTER III

## TENDEIL-BEARERS

Nature of tendrils—Biacouraces, various species of, indi their dimension modes of citizals—Biacouraces, various species of, indicate their modes between the control of adhesive dime. In the control of adhesive dime. The control of their dimensions of the control of their dimensions of their dimensions of their dimensions of their dimensions of their dimensions. Converges —Biacouraces—Smiller agree in institution to confidence of their dimensions. The control of their dimensions of their dimensions of their dimensions of their dimensions of their dimensions. The confidence of their dimensions of thei

By tendrils I mean filamentary organs, sensitive to contact and used exclusively for climbing. By this definition, spines, backs and rootlets, all of which are used for climbing, are excluded. True tendrils are formed by the modification of leaves with their petioles, of flower-pecludes, branches,\* and perhaps stipules.

\* Never having lad the oppertualty of examining touthis produced by the medification of branches, tepoke doubfully about them in this cassy when originally published. But sisce then Fitz Maller has described (Journal of Linn, Sox, vol. is, p. 34) many striking crees is South Brazil. In speaking of plants which climb by the old of their branches, unever or less modified, he he states that the following stages. (1.) Plants supporting theseebve simply by their boundess stetched out at right angles—for example, Chicecon. (2.) Plants despire; a support with their unmodified branchon, as with Secorétaus. (3.) Plants elimbing by the exremitties of their branches which appear like tendrifts, as is the cost according to Endlicher with their branches much modified and temporarily converted into tearitis, but which may be again. Moli, who includes under the name of tendrils various organs having a similar external appearance, classes them according to their homological nature, as being modified leaves, flower-pedundes, &c. This would be an excellent scheme; but I observe that botanists are by no means unaminous on the homological nature of certain tendrils. Consequently I will describe tendrils beauing plants by natural families, following Lindley's classification; and this will in most cases keep those of the same nature together. The species to be described belong to ten families, and will be given in the following collect—Playmonicaes, Expensionae, Computing, Smiltone, Funnariases, Cusrwittaces, Yotaces, Spinlances, Paulariases,

transferrord into bessuber, as with certain Englishmenous plants. (5) Finats with their branches (6) Finats with their branches control of the Englishmen of the Englishmen of the Englishmen of the Strydmen and Caulaterius. Even the unmodified branching some with the summedified branching winds employed them they wind seem of an expectation of an Acestea which have the englishmen of an Acestea which have the englishmen of an Acestea which have the englishmen of the Englishmen

As far as I can make out, the history of our knowledge of tendrils is as follows:—We have seen that Palm and ron Mohl observed about the control of the c singular phenomena of the speatstoom revolving movement of the state of the speak of the state of pressure, observed likewise the revolving movement of hosbits, but I do not feel some of this, be that I do not feel some of this, be Detroiched fully described that movement of the tundril in the movement of the tundril in the common pack, Mol fred discovering to the state of matter the year and some new matter they were and thought as says to excite their movement. Professor Am Gary, in a proper allowed quoted, first notice the state of the stat BRINGMANIACE.—This family contains many tendrilbearers, some twiners, and some root-climbers. The tendrils always consist of modified leaves. Nine species of Bigmoin, selected by hazard, are here elseribed, in order to show what diversity of structure and action there may be within the same genus, and to show what remarkable powers some tendrils posses. The species, taken together, aford connecting links



between twiners, leaf-climbers, tendril-bearers, and root-

climbers. Bignonia (an unnamed species from Kew, closely allied to B. unguis, but with smaller and rather broader

allied to B, unguio, but with smaller and rather broader leaves)—A young shoot from a cut-down plant mole three revolutions against the sun, at an average rate of 2 hrs. 6m. The stem is thin and flexible; it wined round a slender vertical stick, ascending from left to right, as perfectly and as regularly as any true twiningplant. When thus ascending, it makes now so the tendrils or petitions; but when it twined round a rather thick stick, and its petioles were brought to contact with it, these entered round the stick, showing that they have some degree of irritability, the petioles also exhibit a slight degree of spontaneous movement; for in one case they certainly describe, minute, irregular, vertical ellipses. The tendrils apparently curve themselves spontaneously to the assist with the petioles; but from various causes, it was difficult to observe the movement of either the totally or petioles, in this and the two following species. The tendrils are to closely similar in all respects to those of B. mayin kind and the projects to

Bianonia unquis .- The young shoots revolve, but less regularly and less quickly than those of the last species. The stem twines imperfectly round a vertical stick, sometimes reversing its direction, in the same manner as described in so many leaf-climbers; and this plant though possessing tendrils, climbs to a certain extent like a leaf-climber. Each leaf consists of a petiole bearing a pair of leaflets, and terminates in a tendril, which is formed by the modification of three leaflets, and closely resembles that above figured (fig. 5). But it is a little larger, and in a young plant was about half an inch in length. It is curiously like the leg and foot of a small bird, with the hind toe cut off. The straight leg or tarsus is longer than the three toes, which are of equal length, and diverging, lie in the same plane. The toes terminate in sharp, hard claws, much curved downwards, like those on a bird's foot. The petiole of the leaf is sensitive to contact : even a small loop of thread suspended for two days caused it to bend upwards; but the sub-petioles of the two lateral leaflets are not sensitive. The whole tendril, namely, the tarsus and the three toes, are likewise sensitive to contact, especially on their under surfaces. When a shoot grows in the midst of thin branches, the tendrils are soon brought by the revolving movement of the internodes into contact with them; and then one toe of the tendril or more, fast hold of the twigs, like a bird when perched. If the tarsus of the tendril comes into contact with a twig, it goes on slowly bending, until the whole foot is carried quite round, and the toes pass on each side of the tarsus and seize it. In like manner, if the petiole comes into contact with a twig, it bends round, carrying the tendril, which then seizes its own petiole or that of the opposite leaf. The petioles move spontaneously, and thus, when a shoot attempts to twine round an upright stick, those on both sides after a time come into contact with it, and are excited to bend. Ultimately the two petioles clasp the stick in opposite directions, and the foot-like tendrils, seizing on each other or on their own petioles, fasten the stem to the support with surprising security. The tendrils are thus brought into action, if the stem twines round a thin vertical stick; and in this respect the present species differs from the last. Both species use their tendrils in the same manner when passing through a thicket. This plant is one of the most efficient climbers which I have observed; and it probably could ascend a polished stem incessantly tossed by heavy storms. To show how important vigorous health is for the action of all the parts, I may mention that when I well, though not vigorously, I concluded that the tendrils acted only like the hooks on a bramble, and that it was the most feeble and inefficient of all

Biononia Tweedyana,—This species is closely allied to the last, and behaves in the same manner; but perhaps twines rather better round a vertical stick. On the same plant, one branch twined in one direction and another in an opposite direction. The internodes in one case made two circles, each in 2 hrs. 33 m. I was enabled to observe the spontaneous movements of the petioles better in this than in the two preceding species: one petiole described three small vertical ellipses in the course of 11 hrs., whilst another moved in an irregular spire. Some little time after a stem has twined round an upright stick, and is securely fastened to it by the clasping petioles and tendrils, it emits aërial roots from the bases of its leaves; and these roots curve partly round and adhere to the stick. This species of Bignonia, therefore, combines four different methods of climbing generally

In the three foregoing species, when the foot-like tendril has caught an object, it continues to grow and thicken, and ultimately becomes wonderfully strong, in the same manner as the petioles of leafchimbers. If the tendril catches nothing, it first alon'ty bends downwards, and then its power of classing is lost. Very soon affervants it is disarticulates itself from the petiole, and drops off like a leaf in autum. I have seen this process of disarticulation in no other tendrils, for these, when they fail to eath an object.

Bignonia venusta.—The tendrils differ considerably from those of the previous species. The lower part, or tarsus, is four times as long as the three toes; these are of equal length and diverge equally, but do not lie in the same plane; their tips are bluntly booked, tarsus is sensitive on all sides; but the three toes are sensitive only on their outer surfaces. The sensitiveness is not much developed; for a slight rubbing with a twig did not cause the tarsus or the toes to become curved until an hour had elapsed, and then only in a slight degree. Subsequently they straightened themselves. Both the tarsus and toes can seize well hold of sticks. If the stem is secured, the tendrils are seen spontaneously to sweep large ellipses; the two opposite tendrils moving independently of one another. but they are not irritable like those of B. unquis and B. Tweedyana. The young internodes sweep large circles, one being completed in 2 hrs. 15 m., and a second in 2 hrs. 55 m. By these combined movements of the intermedes, petiols, and grapuel-like tendriti, the latter are soon brought into contact with surrounding objects. When a shoot stands near an upright stick, it wines regularly and spirally round it. As it ascends, it exists the stick with one of its tendritis, and, if the stick be thin, the right- and leftfellows from the stem necessarily taking our twist round its own axis for each completed circle. The tendritis courted spirally a short time after

The tendrits contract spirally a short time after catching any object; those which catch nothing merely bend slowly downwards. But the whole subject of the spiral contraction of tendrils will be discussed after all the tendril-bearing species have been described.

\*\*Rimenia litterality\*\*—The vorume internules resolves.

Bigmois litterable,—The young intermoles revolve in large ellipses. An intermole bearing immattree tendrils made two revolutions, each in 3 hrs. 50 m.; but when grown older with the tendrils mattree, it made two ellipses, each at the rate of 2 hrs. 44 m. This species, unlike the preceding, is inexpalled of twining round a stick; this does not appear to be due to any want of flexibility in the intermoles or to the action of the tendrils, and certainly not to any want of the revolving power; nor can I account for the fact. Nevertheless the plant readily ascends a thin upright stick by sixing a point above with tis two opposite tendrils, which then contract spirally. If the tendrils six encoling, they do not become spiral,

The species last described, ascended a vertical stick by twining spirally and by scizing it alternately with its opposite tendrils, like a sailor pulling himself up a rope, hand over hand; the present species pulls itself up, like a sailor seizing with both hands together a rope above his head.

last species. They continue growing for some time, even after they have clasped an object. When fully in length. The three divergent toes are shorter relatively to the tarsus than in the former species; they are not quite equal in length, the middle one being rather longer than the others. Their outer surfaces are highly sensitive; for when lightly rubbed with a twig, they became perceptibly curved in 4 m. and greatly curved in 7 m. In 7 hrs, they became straight again and were ready to re-act. The tarsus, for the space of one inch close to the toes, is sensitive, but in a rather less degree than the toes; for the latter, after a slight rubbing, became curved in about half the time. Even the middle part of the tarsus is sensitive at maturity. After it has grown old, the sensitiveness is confined to the toes, and these are only able to curl very slowly round a stick. A tendril is perfectly ready to act, as soon as the three toes have diverged, and at this period their outer surfaces first become irritable. The irritability spreads but little from one part when excited to another: thus, when a stick was caught by the part immediately beneath the three toes, these seldom clasped it, but remained sticking straight out.

The tendrils revolve spontaneously. The movement begins before the tendril is converted into a threepronged grapnel by the divergence of the toes, and before any part has become sensitive; so that the revolving movement is useless at this early period. completed conjointly in 24 hrs. 18 m. A mature tendril made an ellipse in 6 hrs.; so that it moved much more slowly than the internodes. The ellipses which were swept, both in a vertical and horizontal plane, were of large size. The petioles are not in the least sensitive, but revolve like the tendrils. We thus see that the young internodes, the petioles, and the tendrils all continue revolving together, but at different rates. The movements of the tendrils which rise opposite one another are quite independent. Hence, when the whole shoot is allowed freely to revolve, nothing can be more intricate than the course followed by the extremity of each tendril. A wide space is thus irregularly searched for some object to be

One other curious point remains to be mentioned. In the course of a few days after the toes have closely clasped a stick, their blunt extremities become developed, though not invariably, into irregular discilike balls which have the power of adhering firmly to the wood. As similiar cellular outgrowths will be

fully described under B. capreolata, I will here say Bignonia sequinoctialis, var. Chamberlaynii.-The

internodes, the elongated non-sensitive petioles, and the tendrils all revolve. The stem does not twine, but ascends a vertical stick in the same manner as the last species. The tendrils also resemble those of the last species, but are shorter; the three toes are more unequal in length, the two outer ones being middle toe; but they vary in this respect. They terminate in small hard points; and what is important. duced size of two of the toes as well as their lessened sometimes simple, that is, were not divided into three toes. We are thus naturally led to the three following

show no tendency to twine. Whilst the plant is young and does not require a support, tendrils are not developed. Those borne by a moderately young plant were five inches in length. They revolve spontaneously, as do the short and non-sensitive petioles. When rubbed, they slowly bend to the rubbed side not highly sensitive. There is something strange in their behaviour: I repeatedly placed close to them, thick and thin, rough and smooth sticks and posts, as well as string suspended vertically, but none of these objects were well seized. After clasping an upright stick, they repeatedly loosed it again, and often would not seize it at all, or their extremities did not coil belonging to various Cucurbitaceous, Passifloraceous, and Leguminous plants, and never saw one behave in this manner. When, however, my plant had grown to a height of eight or nine feet, the tendrils acted much better. They now seized a thin, upright stick horizontally, that is, at a point on their own level, and not some way up the stick as in the case of all the previous species. Nevertheless, the non-twining stem was enabled by this means to ascend the stick.

The extremity of the tendril is almost straight and sharp. The whole terminal portion exhibits a singular habit, which in an animal would be called an instinct: into which to insert itself. I had two young plants: and, after having observed this habit, I placed near them posts, which had been bored by beetles, or had become fissured by drying. The tendrils, by their own movement and by that of the internodes, slowly travelled over the surface of the wood, and when the apex came to a hole or fissure it inserted itself; in order to effect this the extremity for a length of half or quarter of an inch, would often bend itself at right angles to the basal part. I have watched this process between twenty and tharty times. The same tendri would frequently withdraw from one hole and insert its point into a second hole. I have also seen a tendril keep its point, in one case for 20 hrs. and in another for 36 hrs., in a minute hole, and then withdraw it. Whilst the point is thus temperarily inserted, the opposite tendril goes on revolving.

to any surface of wood with which it has come into contact; and I have observed one bent at right angles. apex abruptly re-bent and inserted into a minute lateral hole. After a tendril has clasped a stick, it straight downwards. If it has merely adapted itself to nothing, or if it has inserted its apex into some little fissure, this stimulus suffices to induce spiral contraction; but the contraction always draws the tendril away from the post. So that in every case these movements, which seem so nicely adapted for some purpose, were useless. On one occasion, however, the tip became permanently jammed into a narrow fissure. I fully expected, from the analogy of B. capreolata and B. littoralis, that the tips would have been developed into adhesive discs; but I could never detect even a trace of this process. There is therefore at present something unintelligible about

Bignonia picta.—This species closely resembles the

the habits of this plant.

Darwin Online: By permission of the Trustees of the Natural History

last in the structure and movements of its tendrils. I also casually examined a fine growing plant of the allied *B. Lisalleyi*, and this apparently behaved in all respects in the same manner.

Rimonia capreolata.-We now come to a species having tendrils of a different type; but first for the internodes. A young shoot made three large revolutions following the sun, at an average rate of 2 hrs. 93 m. The stem is thin and flexible, and I have seen one make four regular spiral turns round a thin unright. stick, ascending of course from right to left, and therefore in a reversed direction compared with the before described species. Afterwards, from the interference of the tendrils, it ascended either straight up the stick or in an irregular spire. The tendrils are in some respects highly remarkable. In a young branched, the five chief branches apparently representing two pairs of leaflets and a terminal one. Each branch is, however, bifid or more commonly trifid towards the extremity, with the points blunt yet distinctly rubbed, and subsequently becomes straight again: but a loop of thread weighing \$th of a grain produced no effect. On two occasions the terminal branches became slightly curved in 10 m. after they had touched a stick; and in 30 m. the tips were curled quite round it. The basal part is less sensitive. The tendrils revolved in an apparently capricious manner, sometimes very slightly or not at all; at other times they

Whilst the tendrils are revolving more or less namely, a slow inclination from the light towards the darkest side of the house. I repeatedly changed the position of my plants, and some little time after darkest side. When I placed a thick post near a tendril, between it and the light, the tendril pointed in that direction. In two instances a pair of leaves stood so that one of the two tendrils was directed the house; the latter did not move, but the opposite one bent itself first upwards and then right over its fellow, so that the two became parallel, one above the other, both pointing to the dark: I then turned the plant half round; and the tendril which had turned over recovered its original position, and the opposite one which had not before moved, now turned over to the dark side. Lastly, on another plant, three pairs of tendrils were produced at the same time by three shoots, and all happened to be differently directed: I placed the pot in a box open only on one side, and obliquely facing the light; in two days all six tendrils pointed with unerring truth to the darkest corner of the box, though to do this each had to bend in a light which entered the box. I left these tendrils undisturbed for above 24 hrs., and then turned the pot half round; but they had now lost their power of movement, and could not any longer avoid the light.

When a tendril has not succeeded in clasping a that of the shoot, or by turning towards any object which intercepts the light, it bends vertically downtogether with the supporting stick, if there be one. spirally, but soon withers away and drops off. If it seizes an object, all the branches contract spirally.

I have stated that after a tendril has come into contact with a stick, it bends round it in about half an hour; but I repeatedly observed, as in the case of B, speciosa and its allies, that it often again loosed three or four times. Knowing that the tendrils avoided the light, I gave them a glass tube blackened within, and a well-blackened zinc plate: the branches curled the edges of the zine plate; but they soon recoiled from these objects with what I can only call disgust, and straightened themselves. I then placed a post twice they touched it for an hour or two, and twice

curied round and firmly soized an excessively minus projecting point of bark, and then the other branches projecting residually and the state of the project the consistence out, following with sectings every state of the points of the total consistence of the points of the total consistence of the points of the total consistence in a beautiful namer. To my surprise, I observed that the tips of the innanture tentiles, with the branches not yet fully separated, likewise crusted just like roots into the minutest erwiseds. In we or three days after the tips had thus crusted into the crevious, or after their booked outs had sinked minute points, the final process, now to be described, commenced.

This process I discovered by having as-cidentally left a piece of word near a tendrit; and this led me to bind a quantity of data, most, and word boody round attacks and the left of the piece of the left of the

flax, &c., adhere. When a fibre has become fastened to the surface, the cellular tissue does not grow each side; so that when several adjoining fibres, though excessively thin, were caught, so many crests grew up between them, and these, arching over on both sides, adhered firmly together. As the whole surface of the ball continues to grow, fresh fibres adhere and are afterwards enveloped; so that I have of flax crossing it at various angles and all embedded more or less deeply. Every gradation in the process could be followed-some fibres merely sticking to the surface, others lying in more or less deep furrows, or deeply embedded, or passing through the very centre of the cellular ball. The embedded fibres are

a stick, half an inch in diameter, an adhesive disc case of smooth sticks or posts. If, however, the tip catches a minute projecting point, the other branches form discs, especially if they find crevices to erawl

so closely clasped that they cannot be withdrawn. The outgrowing tissue has so strong a tendency to

I infer from the adherence of the fibres to the discs or balls, that these secrete some resinous adhesive anything, discs, as far as I have seen, are never formed; but temporary contact during a moderate time suffices to cause their development. I have seen eight discs formed on the same tendril. After their woody and very strong. A tendril in this state supsupported a considerably greater weight, had not the

the tendrils of this Bignonia can occasionally adhere to smooth cylindrical sticks and often to rugged bark, vet that they are specially adapted to climb trees clothed with lichens, mosses, or other such productions; and I hear from Professor Asa Gray that the Polypodium

\* Fritz Müller states (ibid. p. 348) that in South Brazil the trifid tendrils of Haplolophium,

North America where this species of Bigmonia grows. Finally, I sary remark how singular a fact it is that a leaf should be metamorphosed into a branched organ which turns from the light, and which can by its extremities either crawl like roots into crevices, or seize hold of minute projecting points, these extremities afterwards forming cellular outgrowths which secrets an affinesive cement, and then cavelop by their continued growth the finest fibres.

Economyrus soder (Liponnianor).—Plants, though growing pertly well in my green-house, slowed no spontaneous movements in their shoots or tendrils; but when removed to the hot-house, they young internodes revolved at rates varying from 3 hrs. 15 m. to 1 hr. 13 m. One large circle was swept at this latter unusually quick rate; but generally the circles or ellipses were small, and sametimes the course pursued was quite irregular. An internote, after making sevent and then recommended varying, Suchstrangly mached interruptions in the movements of the internode a I have observed in hardly any other plant.

The leaves bear four leaflets, themselves subdivided.
and terminate in much-branched tendrils. The
main petiole of the leaf, whilst young, moves spontaneously, and follows nearly the same irregular course
and at about the same rate as the internoles. The
movement to and from the stem is the most conspicuous, and L have seen the chord of a curred petiole
which formed an angle of 59° with the stem, in an

hour afterwards making an angle of 100°. The two opposite petidose do not move together, and one is sometimes so much mised as to stand close to the stem, whilst the other is not far from biorizontal. The basis part of the petidos moves less than the distal part. The touristic, beades being carried by the moving petidos and intermedes, themselves moves spontaneously; and directions. By these combined movements of the directions. By these combined movements of the young intermedes, petidoles, and tendrils, a considerable space is swept in exact of a support.

In young plants the tendrils are about three inches in length: they bear two lateral and two terminal branches; and each branch bifurcates twice, with the points directed to the same side. All the branches are or after coming into contact with a stick, bend in about 10 m. One which had become curved in 10 m. after a light rub, continued bending for between 3 hrs. and 4 hrs., and became straight again in 8 hrs. or 9 hrs. Tendrils, which have caught nothing, ultimately contract into an irregular spire, as they likewise do, both cases the main petiole bearing the leaflets, which is at first straight and inclined a little upwards, moves downwards, with the middle part bent abruptly into a right angle; but this is seen in E. miniatus more plainly than in E. scaber. The tendrils in this genus act in some respects like those of Bignonia capreolata; but the whole does not move from the light, nor do the hooked tips become enlarged into cellular dises. After the tendrils have come into contact with a moderately thick cylindrical stick or with

to lift themselves up, change their positions, and again come into contact with the supporting surface, The object of these movements is to bring the double hooks at the extremities of the branches, which naturally face in all directions, into contact with the wood, light, yet that the tips are so, and that they turn and twist themselves towards any dark surface. Ultimately the branches arrange themselves very neatly that they resemble in their irregular course a river with its branches, as engraved on a map. But when a tendril has wound round a rather thick stick, the and spoils the neat arrangement. So it is, but not in quite so marked a manner, when a tendril has spread itself over a large, nearly flat surface of rugged bark. We may therefore conclude that these tendrils are not

a tendril, the terminal branches wind quite round it.

and then seize their own lower branches or the main stem. The sitch is thus firmly, but not nearly grasped. What the tendrils are really adapted for, appears to be such objects as the thin calms of certain grasses, or the long flexible bristles of a brash, or this rigid leaves such as those of the Aspangas, all of which they seize in an admirable namner. This is due to the extremilies of the branches close to the little holds being extremely sensitive to a tondfrom the thinnest object, which they consequently curl round and class. When a small brash, for instance, was placed mear a tendril, the tips of each sub-branch sixed one, two, or three of the bridits and then the spiral contention of the several branches brought all these little parcels close tegother, so that thirty or forty bristles were drawn into a single builde which afforded an excellent support.

POLICONICELE , Colors sondars, This is mescellently constructed climbs. The tendrils on a fine plant were eleven inches long, with the petide bearing two pairs of leadies, only two and a shall inches in length. They revolve more rapidly mot vigcountly than those of any other tendril-beave observed by me, with the exception of one kind of Panafforn. There harps, nearly circular sweeps, fifter the state of the control of the

Coxe III

alone moving. On the other hand, with most of the species of Bignonia and the Eccremocarpus, the internodes, tendrils, and petioles all revolved. The long, bears alternate branches; and each branch is several times divided, with the finer branches as thin as very thin bristles and extremely flexible, so that they are blown about by a breath of air; yet they are strong and highly elastic. The extremity of each branch is a little flattened, and terminates in a minute double (though sometimes single) hook, formed of a hard, trans-Incent, woody substance, and as sharp as the finest needle. On a tendril which was eleven inches long I counted ninety-four of these beautifully constructed little hooks. They readily catch soft wood, or gloves, or the skin of the naked hand. With the exception of stem, every part of every branchlet is highly sensitive on all sides to a slight touch, and bends in a few minutes towards the touched side. By lightly rubbing several sub-branches on opposite sides, the whole tendril rapidly assumed an extraordinarily crooked shape. These movements from contact do not interfere with the ordinary revolving movement. The branches, after becoming greatly curved from being touched, straighten themselves at a quicker rate than in almost any other tendril seen by me, namely, in between half an hour and an hour. After the tendril has caught any object, spiral contraction likewise begins after an unusually short interval of time, namely, in about twelve hours.

Before the tendril is mature, the terminal branchlets cohere, and the hooks are curled closely inwards. At this period no part is sensitive to a touch; but as soon as the branches diverge and the hooks stand out, full that immature tendrils revolve at their full velocity as in this state they can catch nothing. This want of perfect co-adaptation, though only for a short time. between the structure and the functions of a climbingplant is a rare event. A tendril, as soon as it is ready to act, stands, together with the supporting petiolevertically upwards. The leaflets borne by the petiole are at this time quite small, and the extremity of the growing stem is bent to one side so as to be out of the way of the revolving tendril, which sweeps large circles directly over head. The tendrils thus standing above; and by this means the ascent of the plant is favoured. If no object is caught, the leaf with its tendril bends downwards and ultimately assumes a horizontal position. An open space is thus left for the next succeeding and younger tendril to stand vertically upwards and to revolve freely. As soon as an old tendril bends downwards, it loses entangled mass. Although the tendrils revolve with unusual rapidity, the movement lasts for only a short time. In a plant placed in the hot-house and growing vigorously, a tendril revolved for not longer than 36 hours, counting from the period when it first became sensitive; but during this period it probably made at least 97 revolutions.

When a revolving tendril strikes against a stick, the branches quickly bend round and clasp it. The little hooks here play an important part, as they prevent the branches from being dragged away by the rapid revolving movement, before they have had time to clasp the stick securely. This is especially the case when only the extremity of a branch has caught hold of a support. As soon as a tendril has bent! round a smooth stick or a thick rugged post, or has temporarily even to so smooth a surface as this), the same peculiar movements may be observed as those described under Bignonia capreolata and Eccremocarpus. The branches repeatedly lift themselves up and down; those which have their hooks already directed downwards remaining in this position and scenning the tendril, whilst the others twist about until they every irregularity of the surface, and in bringing their hooks into contact with the wood. The use of the hooks was well shown by giving the tendrils tubes and slips of glass to catch; for these, though

The perfect manner in which the branches arranged by this than by any other species. The action is certainly more conspicuous, as the upper surfaces of the main stem, as well as of every branch to the extreme hooks, are angular and green, whilst the lower as in former cases, that a less amount of light guided these movements of the branches of the tendrils. I made many trials with black and white cards and glass tubes to prove it, but failed from various causes ; consists of a leaf split into numerous segments, there is upper surfaces towards the light, as soon as the tendril is caught and the revolving movement is arrested. the segments actually bend or curve to the dark side surfaces may face the light.

When the Colors grows in the open air, the wind must aid the extremely flexible tendrils in seiting a support, for I found that a mere breath sufficed to cause the extreme bennches to catch hold by their hook of twigs, which they could not have reached by the revolving movement. It might have been though that a tendril, thus hooked by the extremity of a single branch, could not have fairly grasped its support. But several times I watched cases like the following: and, arranging its hooks, again caught hold. After a time, from the continued movement of the tendril, the hooks of a third branch caught hold. No other branches, as the tendril then stood, could possibly have touched the stick. But before long the upper part of the main stem began to contract into an open spire. It thus dragged the shoot which bore the tendril towards the stick; and as the tendril coninto contact. And lastly, from the spiral contraction all of them, one after another, were ultimately brought into contact with the stick. They then wound themtendril was tied together in an inextricable knot. The tendrils, though at first quite flexible, after having clasped a support for a time, become more rigid and stronger than they were at first. Thus the

LEGUMINOS.E. Pisum sativum. The common pea who discovered that the internodes and tendrils

<sup>\*</sup> Comptes Rendus, tom, xvii, 1843, p. 989. Darwin Online: By permission of the Trustees of the Natural History

times observed that the longer axis slowly changed its sweeps a wider space. Owing to this change of direction, and likewise to the movement of the stem towards the light, the successive irregular ellipses generally form an irregular spire. I have thought it worth while to annex a tracing of the course pursued P.M. The course was traced on a hemispherical glass placed over the plant, and the dots with figures give the hours of observation; each dot being joined by a straight line. No doubt all the lines would have been curvilinear if the course had been observed at much which the young tendril arose, was two inches from the glass, so that if a pencil two inches in length could have been affixed to the petiole, it would have traced the annexed figure on the under side of the glass; but it must be remembered that the figure is reduced by one-half. Neglecting the first great sweep towards the light from the figure 1 to 2, the end of the petiole swept a space 4 inches across in one direction, and 3 inches in another. As a full-grown as the tendril itself bends and revolves in harmony with the internode, a considerably wider space is swept than is here represented on a reduced scale. Dutrochet observed the completion of an ellipse in 1 hr. 20 m.; and I saw one completed in 1 hr. 30 m. The direction followed is variable, either with or against the sun.

Dutrochet asserts that the petioles of the leaves



Diagram showing the movement of the upper internods of the common Post, traced a hemispherical glass, and transferred to paper; reduced one-half in size. (Aug. 1s No. H. M. No. H. M. No. H. M.

				-									
				P.M.									
				-									

spontaneously revolve, as well as the young internodes and tendrils; but he does not say that he secured the internodes; when this was done, I could never detect any movement in the petiole, except to and from the light.

The tendrils, on the other hand, when the internodes and petioles are secured, describe irregular spires or nodes. A young tendril, only 11 inch in length. revolved. Dutrochet has shown that when a plant is placed in a room, so that the light enters laterally, the internodes travel much quicker to the light than from it: on the other hand, he asserts that the tendril itself moves from the light towards the dark side of the room. With due deference to this great observer, I think he was mistaken, owing to his not having secured the internodes. I took a young plant with highly sensitive tendrils, and tied the petiole so that the tendril alone could move; it completed a perfect ellipse in 1 hr. 30 m.; I then turned the plant partly round, but this made no change in the direction of the succeeding ellipse. The next day I watched a plant similarly secured until the tendril (which was highly sensitive) made an ellipse in a line exactly to and from the light; the movement was so great that the tendril at the two ends of its elliptical course bent itself a little beneath the horizon, thus travelling more than 180 degrees; but the curvature was fully as great towards the light as towards the dark side of the room. I believe Dutrochet was misled by not having secured the internodes, and by having observed a plant of which the internodes and tendrils no longer curved in harmony together, owing to inequality of age.

Dutrochet made no observations on the sensitiveness in length with the leaflets on the petiole only partially expanded, are highly sensitive; a single light touch with a twig on the inferior or concave surface near the tip caused them to bend quickly, as did occasionally (9.25 mg.). The upper or convex surface is barely or not at all sensitive. Tendrils, after bending from a are then ready to act again. As soon as they begin to grow old, the extremities of their two or three pairs of branches become hooked, and they then appear to form an excellent grappling instrument; but this is not the case. For at this period they have generally quite lost their sensitiveness; and when hooked on to from 18 hrs. to 24 hrs. before clasping such twigs; nevertheless, they were able to utilise the last vestige of irritability owing to their extremities being hooked. Ultimately the lateral branches contract spirally, but not the middle or main stem.

Lathyrus aphaca.—This plant is destitute of leaves, except during a very early age, these being replaced by tendrils, and the leaves themselves by large stipules. It might therefore have been expected that the tendrils would have been highly organized, but this is not so. They are moderately long, thin, and un-

branched, with their tips slightly curved. Whilst the concave side of the extremity. They have no upwards at an angle of about 45°, then move into a horizontal position, and ultimately bend downwards. ellipses, and carry with them the tendrils. Two longer axes were directed at about an angle of 45° to the axis of the previously made ellipse.

young and not growing vigorously, yet sufficiently so, I think, for my observations to be trusted. If so we have the rare case of neither internodes nor tendrils revolving. The tendrils of vigorous plants are above three branches; the tips are curved and are sensitive on their concave sides; the lower part of the central stem is hardly at all sensitive. Hence this plant appears to climb simply by its tendrils being brought, through the growth of the stem, or more efficiently by the wind, into contact with surrounding objects, which they then clasp. I may add that the tendrils,

or the internodes, or both, of Vicia sativa revolve. Compositive. — Mutisia clematis. — The immense family of the Compositæ is well known to include very few climbing plants. We have seen in the Table in the first chapter that Mikania scandens is a regular twiner, and F. Müller informs me that in S. Multius is the only genus in the family, as for as I can learn, which bears tendrils; it is therefore interesting to find that these, though rather less metamorphosed from their primordial foliar condition than are most other tendrils, yet display all the ordinary characteristic movements, both those that are spontaneous and those which are excited by countact.

The long leaf bears seven or eight alternate leaflets,

The long leaf bears seven or eight alternate leaflets, and terminates in a tendril which, in a plant of considerable size, was 5 inches in length. It consists generally of three bennehes; and these, although much clongated, evidently represent the petioles and midrits of three leaflets; for they closely recemble the same parts in an ordinary leaf, in being rectangular on the upper surface, furnowed, and edged with green. Moreover, the green edging of the tendrils of young plants semetimes expands into a narrow lamina or blade. Each branch is curved a little downwards, and is slightly blooked at the extensity.

A young upper internols revolved, judging from A young upper internols revolved, judging from three revolutions, at an average rate of 1 hr. 38 m.; it swept ellipse with the longer area directed at right cannot twine. The petidos and the tendrils are but cannot twine. The petidos and the tendrils are but in constant movement. But their movement is abover and much less regularly elliptical than that of the internoles. They appear to be much affected by the light, for the whole leaf usually sinks down during the the day in a crooked course to the west. The tip of the tendril is highly sensitive on the lower surface; and one which was just touched with a twig became perceptibly curved in 3 m., and another in 5 m.; the upper surface is not at all sensitive; the sides are other. The petiole of the leaf and the lower parts of the tendril, halfway between the upper leaflet and the lowest branch, are not sensitive. A tendril after curling from a touch became straight again in about 6 hrs., and was ready to re-act; but one that had been so roughly rubbed as to have coiled into a helix did not become tain their sensibility to an unusually late age; for one borne by a leaf with five or six fully developed leaves above, was still active. If a tendril catches nothing, after a considerable interval of time the tips of the branches curl a little inwards; but if it clasps some object, the whole contracts spirally,

SHILACEE.—Smiles argora, var. monulate.—No. S. SHILACEE.—Smiles argora, var. monulate in pairs from the petiole, are modified lateral leadings in pairs from the petiole, are modified lateral leadings. These tendrils are from 1½ to 1½ inches in length, are thin, and have slightly curved, pointed extendits. They diverge a little from each other, and stand at first nearly uppidit. When lightly subbed on either

\* 'Leçons de Botanique,' &c., 1841, p. 170.

side, they slowly bend to that side, and subsequently become straight again. The back or convex side when placed in contact with a stick became just perceptibly curved in 1 hr. 20 m., but did not completely



surround it until 48 hrs. had elapsed; the concave side of another became considerably curved in 2 hrs. and clasped a stick in 5 hrs. As the pairs of tendrils grow old, one tendril diverges more and more from the other, and both slowly bend backwards and downwards, so that after a time they project on the opposite side

of the stem to that from which they arise. They then placed behind the stem. Owing to this power, the plant is able to ascend a thin upright stick. Ultiif they do not come into contact with any object, loosely cross each other behind the stem, as at B. in fig. 7. This movement of the tendrils towards and round the stem is, to a certain extent, guided by their one of the two tendrils was compelled in thus slowly moving to travel towards the light, and the other from the light, the latter always moved, as I repeatedly observed, more quickly than its fellow. The tendrils do not contract spirally in any case. Their chance of finding a support depends on the growth of the plant, on the wind, and on their own slow backward and downward movement, which, as we have just seen, is guided, to a certain extent, by the avoidance of the light; for neither the internodes nor the tendrils have any proper revolving movement. From this latter circumstance, from the slow movements of the tendrils an unusual length of time), from their simple structure and shortness, this plant is a less perfect climber than any other tendril-bearing species observed by me. The plant whilst young and only a few inches in height, does not produce any tendrils; and considering that it grows to only about 8 feet in height, that the stem spines, it is surprising that it should be provided with tendrils, comparatively inefficient though these are. The plant might have been left, one would have thought, to climb by the aid of its spines alone, like our brandles. As, however, it belongs to a genus, some of the species of which are furnished with much longer tendrils, we may suspect that it possesses these organs solely from being descended from progenitors more highly organized in this respect.

FUMARIACE.E. Corudalis elaviculata. According to Mohl (p. 43), the extremities of the branched stem, as well as the leaves, are converted into tendrils. In the specimens examined by me all the tendrils were certainly foliar, and it is hardly credible that the same plant should produce tendrils of a widely different homological nature. Nevertheless, from this statement by Mohl, I have ranked this species amongst the tendril-bearers; if classed exclusively by its foliar have been placed amongst the leaf-climbers, with its allies, Fumaria and Adlumia. A large majority of its so-called tendrils still bear leaflets, though excessively reduced in size; but some few of them may properly be designated as tendrils, for they are completely destitute of laminæ or blades. Consequently, we here young, only the outer leaves, but when full-grown all the leaves, have their extremities converted into more or less perfect tendrils. I have examined specimens from one locality alone, viz. Hampshire; and it is not improbable that plants growing under different conditions might have their leaves a little more or less

changed into true tendrils. Whilst the plant is quite young, the first-formed leaves are not modified in any way, but those next formed have their terminal leaflets reduced in size, and soon all the leaves assume the structure represented in the following drawing. This leaf bore nine leaflets; the lower ones being much subdivided. The terminal portion of the petiole, about 11 inch in length (above the leaflet f), is thinner and more elongated than the lower part, and may be considered as the tendril. The leaflets borne by this part are greatly reduced in size, being, on an average, about the tenth of an inch in length and very narrow; one small leaflet measured one-twelfth of an inch in length and one-seventy-fifth in breadth (2·116 mm. and ·339 mm.), so that it was almost microscopically minute. All the reduced leaflets have branching nerves, and terminate in little spines, like those of the fully developed leaflets. Every gradation could be traced. until we come to branchlets (as a and d in the figure) which show no vestige of a lamina or blade. Occasionally all the terminal branchlets of the petiole are in

this condition, and we then have a true tendril. The several terminal branches of the petiole bearing the much reduced leaflets (a, b, c, d) are highly sensitive, for a loop of thread weighing only the onesixteenth of a grain (4\*05 mg.) caused them to become greatly curved in under 4 hrs. When the loop was removed, the petioles straightened themselves in about the same time. The petiole (e) was rather less sensitive; and in another specimen, in which the corresponding



petiole bore rather larger leaflets, a loop of thread weighing one-eighth of a grain did not cause curvature until 18 hrs. had elapsed. Loops of thread weighing one-fourth of a grain, left suscended on the lower petioles (f to  $\bar{I}_{I}$  during several days, produced no effect. Yet the three petioles  $\bar{I}_{I}$ , and I were not quite insensible, for when left in contact with a stick for a day or two they slowly curled round it. Thus the sensibility of the petiole gradually diminishes from the tond-like extensity to the base. The internodes of the stem are not at all sensitive, which makes N-flowly substressed that they are sometimes contaction of the stem are not at all sensitive, which makes N-flowly substressed that they are sometimes contacting the stem of the N-flowly stem of the stem of the stem of the stem of the N-flowly stem of the stem of

almost vertically upwards, as we have seen to be the case with many tendrils. It is in continual moverate of about 2 hrs. for each revolution, large, though different points of the compass. The young internodes, likewise revolved irregularly in ellipses or spires; so that by these combined movements a conand attenuated portion of a petiole fails to seize any object, it ultimately bends downwards and inwards, and soon loses all irritability and power of movement, which occurs with the extremities of the young leaves in many species of Clematis; for these, when thus bent downwards or hooked, first acquire their full degree of sensitiveness,

- Dicentra thalictrifolia .- In this allied plant the

metamorphosis of the terminal leaflets is complete, and they are converted into perfect tendrils. Whilst the plant is young, the tendrils appear like modified there can be no doubt, as I am assured by Dr. Hooker, that they are modified leaves. When of full size, they are above 5 inches in length; they bifurcate twice, thrice, or even four times; their extremities are hooked and blunt. All the branches of the tendrils are sensitive on all sides, but the basal portion of the when lightly rubbed with a twig became curved in themselves in between 10 hrs. and 20 hrs. A loop of thread weighing one-eighth of a grain plainly ally a loop weighing one-sixteenth of a grain; but this latter weight, though left suspended, was not sufficient to cause a permanent flexure. The whole leaf with its tendril, as well as the young upper interlarly, and thus sweeps a wide space. The figure traced on a bell-glass was either an irregular spire or a zigzag line. The nearest approach to an ellipse was an elongated figure of 8, with one end a little open, and this was completed in 1 hr. 53 m. During a period apparently representing three and a half ellipses.

was securely fastened, the tendril itself described similar but much smaller figures.

This species climbs well. The tendrils after clasping a stick become thicker and more rigid; but the by some Bignoniacese and Cobsea. The tendrils of young plants, two or three feet in height, are only half the length of those borne by the same plant when grown taller, and they do not contract spirally after clasping a support, but only become slightly flexuous. wards and inwards, like the extremities of the leaves of the Corydalis claviculata. But in all cases the petiole after a time is angularly and abruptly bent downwards like that of Eccremocarpus.

## CHAPTER IV.

TEXDELL DEARERS—(COMPARCO)

Commercial—Honologum nature of the nodific—Estimophic total, remarkable moments of the thoribit was old ringing the remains remarkable moments of the thoribit was old ringing the remains remarkable moments of the thoribit was old ringing for a similar damp of uter—Unidadery, moment of the extremity of the tended adapted uter—Lindskeys, andered diss—France-fordshine between the flower polamides and tendellard their im—Toublise the Virginian Compete term from the light, and, after existed, adverage adhering time—Surremann—Learned their remains of the control of the state of the state of the state of the tendellar state of the state of the state included and the state of the trade of the state of the state of the state of toucht.

COCCUMPACENT. The tearlik in this family have been ranked by competent judges as modified leaves, stipules, or branches; or as partly a leaf and partly a branch. De Candollo believes that the tendrish differ in their homological nature in two of the tribes.\* From facts recently addisced, Mr. Berkeley thinks that Payer's view is the most probable, namely, that the tendril is "a separate portion of the last itself;" but much may be said in favour of the belief that it is a modified flower-polumical;"

for information on this head, the Bulletin do is Societé I nique de France, 1857, there numerous discussions on nature of the tendrils in p. 721. From the affinity of the Cocurbitacese to the Passifloracese, it might be argued that the tendrils of the former are no ified flower-peduncles, as is certainly the case with those of Passion made on this plant (raised from seed sent me by Prof. Asa Grav), for the spontaneous revolving movements me in this case, and greatly perplexed me. My obserthe slowest rate was 2 hrs., and the average rate, with no great fluctuations, 1 hr. 40 m. Sometimes I tied the internodes, so that the tendrils alone moved: at other times I cut off the tendrils whilst very young, so that the internodes revolved by themselves; but the rate was not thus affected. The course generally pursued was with the sun, but often in an opposite direction. Sometimes the movement during a short time would either stop or be reversed; and this apparently was due to interference from the light, as, for instance, when I placed a plant close to a window. In one instance, an old tendril, which had nearly ceased revolving, moved in one direction, whilst a young tendril above moved in an opposite course. The two uppermost internodes alone revolve: and as soon as the lower one grows old, only its upper part continues to move. The ellipses or circles swept

flowers. Mr. R. Holland (Hardwicke's 'Science-Gessip,' 1865, p. 105) states that "a cucumber

Hard-65, p. prickles upon the fruit had maker grown out into a long, curled rown borded."

by the summits of the internodes are about three inches in diameter; whilst those swept by the tips of the teachii, are from 15 to 16 inches in diameter. During the revolving movement, the intendeds become successively curved to all points of the compast, in one part of their course they are often indined, together with the teachiis, at about 45° to the horizon, and in another part stand overlightly up. There was something in the appearance of the revolving internodes which continually gave the false impression that their movement was due to the weight of the long and syntaneously revolving teachil; luct, on cutting off the latter with sharp scissors, the top of the shoot rase only a little, and went on revolving. This false appearance is apparently due to the internodes and tendrils all curving and moving harmoniously together.

A revolving tendril, though inclined during the greater part of its course at an angle of about 45° (in one case of only 37°) above the horizon, stiffened and straightened itself from tip to base in a certain part of its course, thus becoming nearly or quite vertical. I witnessed this repeatedly; and it occurred both when the supporting internodes were free and when they were tied up; but was perhaps most conspicuous in the latter case, or when the whole shoot happened to be much inclined. The tendril forms a very acute angle with the projecting extremity of the stem or shoot; and the stiffening always occurred as the tendril approached, and had to pass over the shoot in its circular course. If it had not possessed and exercised this curious power, it would infallibly have struck against the extremity of the shoot and been

as the tendril has succeeded in passing over the extremity of the shoot or point of difficulty, its motion, coinciding with that from its weight, often causes it to

The tendrils are thin, from 7 to 9 inches in length, with a pair of short lateral branches rising not far concave side of the tip is highly sensitive to a touch; but not so the convex side, as was likewise observed to be the case with other species of the family by one tendril, and only once or twice the concave side In a few hours afterwards, when the tendrils which had been rubbed on the concave side had straightened themselves, I reversed the process of rubbing, and always with the same result. After touching the two minutes; and subsequently, if the touch has been at all rough, it coils itself into a helix. But the helix will, after a time, straighten itself, and be again ready to act. A loop of thin thread only one-sixteenth of a grain in weight caused a temporary flexure. The lower part was repeatedly rubbed rather roughly, but no curvature ensued; yet this part is sensitive to prolonged pressure, for when it came into contact with a stick, it would slowly wind round it.

One of my plants bore two shoots near together, and the tendris were repeatedly drawn across one another, but it is a singular fact that they did not once each each other. It would appear as if they had become habituated to contact of this kind, for the persure thus caused must have been much greater than that caused by a loop of soft thread weighing only the one-sixteenth of a grain. I have, however, seen several tendrits of the point delois interfecked, but they abbegundly released one another. The tendrist of the Echinocystis are also habituated to drops of water or to rin; for artificial rain made by violently filting a wet breash over them produced not the least effect.

The revolving movement of a tendril is not stopped by the enring of its extensity after it has been touched. When one of the lateral branches has firmly cleaped an object, the middle branch continues to revolve. When a stem is bent down and secured, so that the tendril depends but is left free to move, its previous revolving movement is nearly or quite stopped; but it soon begins to bend upwards, and as soon as it has become herizontal the revolving movement recommences. It told his four times; the tendril generally rose to a berizontal position in an hour or an hour and

a half, but in one case, in which a tendril depended at an angle of 40 beneath the lorion, the uprising two two loars; in half an hour afterwards it rose to 220 shows the busins and then recommender leveling. This upward movement is independent of the action of light, for it occurred twice in the dark, and on another consists the light came in on one side alson. The movement to doubt it guided by opposition to the faces of gravity, as in the case of the ascent of the showled of gravity, as in the case of the ascent of the showled of gravity.

and as soon as this is lost, it bends downwards and contracts spirally. After the revolving movement has ceased, the tip still retains for a short time its sensitiveness to contact, but this can be of little or no use to the plant.

Though the tendril is highly flexible, and though the catenulty travels, under Avourable circumstance, at about the rate of an inch in two minutes and a quarter, yet its sensitiveness to enotate; is or great that it hardly over fails to seize a thin stick placed in its pubt. The following case surprised ne much: I placed a thin, smooth, cylindrical stick (and I repeated the experiment sweet times) so far from a tendril, that its extremity could only end half or three-quarters round the stick; I but I always found that the tip managed in the course of a few hours to cent twice or even their cound the stick. I at first though that this was due to rapid growth on the outside; but be coloured points and measurements. I proved that there had been no sensible increase of length within the time. When a stick, flat on one side, was similarly placed, the tip of the tendril could not curl beyond the flat surface, but coiled itself into a helix, which, turning to one side, lay flat on the little flat surface of wood. In one instance a portion of tendril three-quarters of an inch in length was thus dragged on to the flat surface by the coiling in of the helix. But the tendril thus acquires a very insecure hold, and generally after a time slips off. In one case alone the helix subsequently uncoiled itself, and the tip then passed round and clasped the stick. The formation of the helix on the flat side of the stick apparently shows us that the continued striving of the tin to curl itself closely inwards gives the force which drags the tendril round a smooth cylindrical stick, In this latter case, whilst the tendril was slowly and quite insensibly crawling onwards, I observed several times through a lens that the whole surface was not in close contact with the stick; and I can understand the onward progress only by supposing that the movement is slightly undulatory or vermicular, and that the tip alternately straightens itself a little and then again curls inwards. It thus drags itself onwards by an insensibly slow, alternate movement, which may be compared to that of a strong man suspended by the ends of his fingers to a horizontal pole, who works his fingers onwards until he can grasp the pole with the palm of his hand. However this may be, the fact is certain that a tendril which has caught a round stick

with its extreme point, can work itself onwards until it has passed twice or even thrice round the stick, and has permanently grasped it.


Hanburya Mexicana.-The young internodes and in the same manner and at about the same rate as those of the Echinocustis. The stem does not twine, but can ascend an upright stick by the aid of its tendrils, The concave tip of the tendril is very sensitive; after it had become rapidly coiled into a ring owing to a single touch, it straightened itself in 50 m. The tendril, when in full action, stands vertically up, with the projecting extremity of the young stem thrown a little on one side, so as to be out of the way; but the tendril bears on the inner side, near its base, a short rigid branch, which projects out at right angles like a spur, with the terminal half bowed a little downwards. Hence, as the main vertical branch revolves, the spur, from its position and rigidity, cannot pass over the extremity of the shoot, in the same curious manner as do the three branches of the tendril of the Echinocystis, namely, by stiffening themselves at the proper point. The spur is therefore pressed laterally against the young stem in one part of the revolving course, and thus the sweep of the lower part of the main branch is much restricted. A nice case of co-adaptation here comes into play: in all the other tendrils observed by me, the several branches become sensitive at the same period: had this been the case with the Hanburya, the inwardly directed, spur-like branch, from being pressed, during the revolving movement, against the projecting and of the shoot, would infallibly have seized it in a uncless or injurious namer. But the main branch of the teathir, after revolving for a time in a vertical position, spontaneously bends downwards; and in doing so, ruises the sparticle branch, which itself also curves upwards; so that by these combined movements it rises above the projecting end of the shoot, and can now move freely without touching the shoot; and now if first becomes somition.

The tips of both branches, when they come into contact with a stick, grasp it like any ordinary tendril. But in the course of a few days, the lower surface swells and becomes developed into a cellular layer. which adapts itself closely to the wood, and firmly adheres to it. This layer is analogous to the adhesive discs formed by the extremities of the tendrils of some species of Bignonia and of Ampelopsis; but in the Hanburya the layer is developed along the terminal inner surface, sometimes for a length of 13 inches, and not at the extreme tip. The layer is white, whilst the tendril is green, and near the tip it is sometimes thicker than the tendril itself; it generally spreads a little beyond the sides of the tendril, and is fringed with free elongated cells, which have enlarged globular or retort-shaped heads. This cellular layer apparently secretes some resinous cement; for its adhesion to the wood was not lessened by an immersion of 24 hrs. in alcohol or water, but was quite loosened by a similar immersion in other or turpentine.

After a teodril has once firmly coiled itself round a stick, it is difficult to imagine of what use the adheaire cellular layer can be. Oring to the spiral contraction which soon ensure, the tendrils were never able to remain, excepting in one instance, in constawith a thick post or a nearly flat surface; if they had quickly become attached by means of the adhesive layer, this would oridently have been of service to the plant.

The tendrits of Dryonis dision, Cheenbids onliers, and Cosmon state are sensitive and revolve. Whether the internoles likewise revolve I did not observe. In Auguria Warneswieri, the internoles, though thick and stiff, revolve 1 in this plant the lower aurhor of the tendril, some time after clasping a stick, produces a consely collular layer or embinos, which slapst itself closely to the wood, like that formed by the tendril of the Hundways Ju tuit is not in the least athletice. In the Induleys Ju tuit is not in the least athletics, the family, the forcked tendrils and the internoles revolve in periods between 2 krs. 8 m. and 3 hrs. 35 m. moving against the sm.

VITACE.E.—In this family and in the two following, analey, the Sapindaece and Passifonnees, the testificate and in the indistance of the single and are therefore axial in their nature. In this respect they differ from a previously described, with the exception, perhaps, of the Cheurbitacees. The homological nucleon however, of a tendril seems to make no difference in its action. Wite vinifera.—The tendril is thick and of great length; one from a vine growing out of does and not length; one from a vine growing out of does and not vigocously, was 16 inches long. It consists of a polumele (A), bearing two branches which diverge equally from it. One of the branches (B) has a scale at its less; it is always, as far as I have seen, longer than the other and often bifurcates. The besorbes when rubbled become curved, and with



quently straighten themselves. After a tendril has clasped any object with its extremity, it contracts spirally; but this does not occur (Palm, p. 56) when taneously from side to side; and on a very hot day, one made two elliptical revolutions, at an average rate of 2 lns. 15 m. During these movements a colonal line, painted along the convex surface, appeared albre a time on one side, then on the convex side, then on the convex side, then on the coprosite side, and hardy again on the correst side. The two branches of the same tendril have independent force time, it beads from the light twombs the dark: a steady that the convex side of the side of

The young internodes revolve spontaneously; but the movement is unusually slight. A shoot faced a window, and I traced its course on the glass during two perfectly calm and hot days. On one of these days it described, in the course of ten hours, a spire, representing two and a half ellipses. I also placed a bell-glass over a young Muscat grape in the hothouse, and it made each day three or four very small oval revolutions; the shoot moving less than half an inch from side to side. Had it not made at least three revolutions whilst the sky was uniformly overcast, I should have attributed this slight degree of movement to the varying action of the light. The extremity of the stem is more or less bent downwards, but it never reverses its curvature, as so generally occurs with twining plants,

Various authors (Palm, p. 55; Mohl, p. 45; Lindley, &c.) believe that the tendrils of the vine are modified flower-peduncles. I here give a drawing (fig. 10) of the ordinary state of a young flower-stalk; it consists



Flower-stalk of the Vine.

A. Common Pelancie.

B. Flower-tendrii, with a soale at its base.

D. Peticle of the opposite leaf.

of the "common peduncle" (A); of the "flower-tendril" (B), which is represented as having caught a twig; and of the "sub-peduncle" (C) bearing the flower-buds. The whole moves spontaneously, like a true tendril, but in a less degree; the movement,

however, is greater when the sub-peducele (C) does not bear many fovor-bulk. The common peducele (A) has not the power of clasping a support, nor has the corresponding part of a true tendril. The flowertendril (B) is always longer than the sub-peducele (C) and has a scale at its base; it sometimes bifurcates, and therefore corresponds in every detail with the longer scale-bearing branch (B, fig. 9) of the true bendril. It is, however, inclined backwards from the sub-peducele (C), or stands at right angles with it, and is thus subspect to aid in carrying the future beached, or gross. When rubbed, it curves and subbended of gross. When rubbed, it curves and subtered the control of the control of the control of the drawing securely clasp a support. I have seen an object as soft as a young vine-leaf caught by one.

The lower and naked part of the sub-pelmude (U) is likewise slightly sensitive to a rule, and I have seen it bent round a stick and even partly round a leaf with which it had come into contact. That the sub-pelmude has the same nature as the corresponding beands of an ordinary tendril, is well shown when it beans only a few flowers; for in this case it becomes less branched, increases in length, and gains both in sensitiveness and in the power of spontaneous movement. I have twice soom sub-pelmudes which bore from thirty to forty flower-bads, and which had become considerably clongated and were completely wound round sticks, exactly like true tendrils. The whole length of mother sub-polencie, bearing only

eleven flower-buds, quickly became curved when slightly rubbed; but even this scanty number of flowers rendered the stalk less sensitive than the other branch, that is, the flower-tendril; for the latter after a lighter rub became curved more quickly and in a greater degree. I have seen a sub-peduncle thickly covered with flower-buds, with one of its higher lateral branchlets bearing from some cause only two buds; and this one branchlet had become much elongated and had spontaneously caught hold of an adjoining twig; in fact, it formed a little subtendril. The increasing length of the sub-peduncle (C) with the decreasing number of the flower-buds is a good instance of the law of compensation. In accordis always longer than the flower-stalk; for instance, on the same plant, the longest flower-stalk (measured from the base of the common peduncle to the tip of the flower-tendril) was 81 inches in length, whilst the longest tendril was nearly double this length, namely

The gralations from the ordinary state of a floresstalk, as represented in the drawing (fig. 10), to that of a true tendri (fig. 9) are complets. We have seen that the sub-peduncle (C), whilst still bearing from thirty to firely flower-bules, sometimes becomes a little elongated and partially assumes all the characters of the corresponding branch of a true tendril. From this state we can trace every stage till we come to a full-sized perfect tendril, bearing on the branch a full-sized perfect tendril, bearing on the branch which corresponds with the sub-peduncle one single flower-bud! Hence there can be no doubt that the tendril is a modified flower-peduncle.

Flower-tendrils (B, fig. 10) sometimes produce a few my house, there were thirteen and twenty-two flowerbuds respectively on two flower-tendrils, which still retained their characteristic qualities of sensitiveness and spontaneous movement, but in a somewhat lessened occasionally produced on the flower-tendrils that a whole bunch of flowers presents scarcely any resemblance to a tendril; and, judging from the facts already given, it would probably possess little power of clasping a support, or of spontaneous movement. Such flower-Cissus. This genus, belonging to the same family of the Vitacese, produces well-developed tendrils and between the two states. If the genus Vitis had been unknown, the boldest believer in the modification of species would never have surmised that the same individual plant, at the same period of growth, would have yielded every possible gradation between ordinary flower-stalks for the support of the flowers and fruit, and tendrils used exclusively for climbing. But the vine clearly gives us such a case: and it

seems to me as striking and curious an instance of transition as can well be conceived.

Cissus discolor.-The young shoots show no more movement than can be accounted for by daily variations in the action of the light. The tendrils, however, revolve with much regularity, following the sun ; and, in the plants observed by me, swept circles of about 5 inches in diameter. Five circles were completed in the following times: -4 hrs. 45 m., 4 hrs. 50 m., 4 hrs. 45 m., 4 hrs. 30 m., and 5 hrs. The same tendril continues to revolve during three or four days. The tendrils are from 31 to 5 inches in length. They are formed of a long foot-stalk, bearing two short branches, which in old plants again bifurcate. The two branches are not of quite equal length; and as with the vine, the longer one has a scale at its base. The tendril stands vertically upwards; the extremity of the shoot being bent abruptly downwards, and this position is

Both beamless of the totalril, whilst young, are highly sensitive. A totale with a pomel, be greatle as only just to move a tendril borne at the end of a long flexible shoot, sufficed to cause it to become perceptibly curved in four or five minutes. It became straight again in rather above one bour. A loop of soft thread weighing one-seventh of a grain (1925 mg.) was turies tried, and each time caused the cadril to become curved in 30 or 40 m. Half this weight produced on effect. The long foot-stall is much less sensitive, for a slight rubbing produced no effect, although prolonged contact with a tile caused it to bead. The two branches are sensitive on all sides, so that they converge if touched on their inner sides, and diverge if touched on their outer sides. If a branch is touched at the same time with equal force on opposite sides, both sides are equally stimulated and there is no movenent. Before canning this plant, I had observed only tendris which are sensitive on one side alone, and these when lightly pressed between the finger and thumb become curved; but on these products and the subcession of the control of the control of the contact of the control of the control of the contact of the control of the control of the concession and the control of the control of the concession assured on the control of the concession assured on the control of the concession assured on the control of the contro

were thick an atractess—re tenurs or any young passwere thick and straight, with the tips a little curved. When their concave surfaces were rubbed, and it was necessary to do this with some force, they very slowly became curved, and subsequently straight again. They are therefore much less sensitive than those of the last species; but they made two revolutions, following the sun, rather more rapidly, viz, in 3 has 30 m. and 4 hrs. The internodes do not revealve.

Amplopin hadranca (Virginian Grosper)—The internoles apparently do not more more than can be accounted for by the varying action of the light. The tendrils are from 4 to 5 inches in length, with the main stem sending off several lateral branches, which have their tips curved, as may be seen in the upper figure (fig. 11). They exhibit not true spontaneous revolving movement, but turn, as was long ago observed by Andrew Knight,\* from the light to the dark. I have an angle of 180° to the dark side of a case in which much slower. The several lateral branches often move larly, without any apparent cause. These tendrils are me. By gentle but repeated rubbing with a twig, the course of three or four hours slightly curved; but they seemed to have hardly any power of again had crawled over a large box-tree clasped several of the branches; but I have repeatedly seen that they will meet with a flat surface of wood or a wall (and this all their branches towards it, and, spreading them widely apart, bring their hooked tips laterally into after touching the surface, often rise up, place them-

In the course of about two days after a tendril has arranged its branches so as to press on any surface, the curved tips swell, become bright red, and form on

<sup>\*</sup> Tmns. Phil. Soc. 1812, p. 314.

with which they adhere firmly. In one case the tips were slightly swollen in 38 hrs. after coming into but became attached to a stuccoed wall in 42 hrs. These adhesive discs resemble, except in colour and in being larger, those of Bignonia capreolata. When a manner as by B. capreolata. Discs are never deare generally first formed on one side of the curved tip, the whole of which often becomes so much changed can be traced only along the concave surface. When, however, a tendril has clasped a cylindrical stick, an irregular rim or disc is sometimes formed along the

Dr. Mi'Nab remarks (Truns. Bot. Soc. Edinburgh, vol xi. p. 202) that the tendrils of Awy. Vallehil bear small globular dises before they have come into contact with any object; and I have since observed the same fact. These dises, however, increase greatly in size if they proce acquisit and

drils, therefore, of one species of Ampelopsis require the stimulus of contact for the first development of their discs, whilst these of another species do not need any such stimulus. We have seen an exactly parallel case with two species of Rimonsiscen. tip; this was also observed (p.71) by Mohl. The discs consist of enlarged cells, with smooth projecting hemispherical surfaces, coloured red; they are at first gorged with fluid (see section given by Mohl, p. 70), but ultimately become woody.

leaf of the ivy, this alone renders it probable that some cement is secreted, as has been asserted to be the case (quoted by Mohl, p. 71) by Malpighi. I year from a stuccoed wall, and left them during many much, but warmed essential oils (I tried oil of thyme however, must be small; for when a plant ascended the whitewash; but as the cement never penetrated the thin layer, they were easily withdrawn, together supposed that the attachment is effected exclusively

A tendril which has not become attached to any body, does not contract spirally; and in course of a

week or two shrinks into the finest thread, withers and




Fig. 11. Ampelopsis Anderse

A. Tendiffally developed, with a young test on the opposite side of the stem.

B. Odder textifs, everal weeks dier its attachment to a wall, with the Branches Linkowski and optically contrasted, and with the attentiates developed into diece. The unstabless beamdes of this section has weithered and dropped off.

drops off. An attached tendric if the other hand,

contracts spirally, and thus becomes highly elastic, 80

that when the main foot-stalk is pulled the strain is distributed equally between all the attached discs. For a few days after the attachment of the discs, the in thickness and acquires great strength. During the following winter it ceases to live, but adheres firmly in attachment. In the accompanying diagram (fig. 11.) we see the difference between a tendril (B) some weeks after its attachment to a wall, with one (A) from the same plant fully grown but unattached. That the change in the nature of the tissues, as well as the spiral contraction, are consequent on the formation of have not become attached; for these in a week or two wither and drop off, in the same manner as does the whole tendril if unattached. The gain in strength and durability in a tendril after its attachment is to my house which are still strong, and have been exposed to the weather in a dead state for fourteen or fifteen years. One single lateral branchlet of a tendril, estimated to be at least ten years old, was still elastic and supported a weight of exactly two pounds. The thickness and apparently of equal strength; so that after having been exposed during ten years to the

en pounds:
Sapindace E.—Cardiospermum halicacabum,—In this

family, as in the last, the tendrils are modified flowerpeduncles. In the present plant the two lateral branches of the main flower-peduncle have been converted into a pair of tendrils, corresponding with the single "flower-tendril" of the common vino. The main peduncle is thin, stiff, and from 3 to 4½ inches in length. Near the summit, above two little bracts, it divides into three branches. The middle one divide divides into three branches. The middle one divide



and re-divides, and bears the flowers; ultimately it

grows half as long again as the two other modified branches. These latter are the tendrils; they are at first thicker and longer than the middle branch, but never become more than an inch in length. They taper to a point and are flattened, with the lower clasping earfined estitute of hairs. At first they project straight up; but soon diverging, spontaneously end downwards so as to become symmetrically and elegantly hooked, as represented in the disagnum. They are now, whilst the flower-buds are still small, ready for The two or three upper internodes, whilst young, steadily revolve; those on one plant made two circles, against the course of the sun, in 3 hrs. 12 m.; in a second plant the same course was followed, and the two circles were completed in 3 hrs. 41 m.; in a third plant, the internodes followed the sun and made two circles in 3 hrs. 47 m. The average rate of these six revolutions was 1 hr. 46 m. The stem shows no tendency to twine spirally round a support; but the allied tendril-bearing genus Paullinia is said (Mohl, p. up above the end of the shoot, are carried round and round by the revolving movement of the internodes; and when the stem is securely tied, the long and thin flower-peduncles themselves are seen to be in continued and sometimes rapid movement from side to side. They sweep a wide space, but only occasionally revolve in a regular elliptical course. By the one of the two short hooked tendrils, sooner or later, catches hold of some twig or branch, and then it curls ever, but slightly sensitive; for by rubbing their under surface only a slight movement is slowly produced. was curved considerably inwards; in 2 hrs. 30 m. it formed a ring; and in from 5 to 6 hours from being tendril acted at nearly the same rate; but I observed thin tsig. Tendrils which have caught nothing, spontaneously or un to a close helix after the interval of sevenal days. Those which have carded round some object, some become a little thicker and tougher. The long and thin main polumels, though spontaneously moving, is not sensitive and never clasps a support. Nor descrivers centract spirally, although a contraction of this kind apparently would have been of service to the plant in climbing. Nevertheless it climbs petry self without this aid. The seed-capacity through the properties of the bendrile served simply for climbing.

The position of the tendrils alone suffices to show

their homological nature. In two instances one of two tendrils produced a flower at its tip; this, however, did not prevent its acting properly and eurling round a twig. In a third case both lateral branches which ought to have been modified into tendrils, produced flowers like other central branch, and had quite lost their tendril-structure.

There were, but was not enabled consultate to observe

only one other climbing Sapindaceous plant, namely,

Fritz Müller remarks (ibid. p. 348) that a related genus, Serjania, differs from Cardiospermum in bearing only a single tendril; and

the common pedunele ts spirally, when, as freque pens, the tendril has else plant's own stem

Paullinia. It was not in flower, yet bore long forked tendrils. So that, Paullinia, with respect to its tendrils, appears to bear the same relation to Cardiospermum that Cissus does to Vitis.

that Cleans does to Föx.

PassirizanceXim—After reading the discussion and
facts given by Moli (p. 47) on the nature of the
tendric in this family, no one can down that they are
modified flower-peduncles. The tendric is a discussed in the control of the control

Pausifora grantia.—This well-named, elegant, annual species differs from the other members of the group elserved by me, in the young internoles having the power of revolving. It exceeds all the other elimbing plants which I have crassined, in the rapidity of its movements, and all tendril-beares in the sensitiveness of the tendrils. The internole which carries the upper active tendril and which likewise carries one or two younger inmatures internoles, made there revolutions, following the sun, at an average rate of I hr.4 m.; it then made, the day becoming very hot, three other revolutions at an average rate of between 57 and 58 m;; so that the average of all six revolutions was I hr. Im. The apex of the tendril describes elongated clipses, sometimes brand, with their longer axes inclined in slightly different directions. The plant can ascend a thin upright still by the aid of its tendrils; but the stem is too stiff for it to twin opinilly round it, even when not intered with by the tendrils, these having been successively prinched off at an early age.

When the stem is secured, the tendrils are seen to revolve in nearly the same manner and at the same rate as the intermodes.\* The tendrils are very thin, delicate, and straight, with the exception of the tigs, which are a little curved; they are from 7 to 9 inches in length. A half-grown tendril is not sensitive; but when nearly full-grown they are extremely sensitive. A single delicate touch on the concess surface of the tip soon caused one to curve; and in 2 minutes it formed an open bolit. A bog of soft thread weighing glad of a gain (202 mg.) piaced most gently on the thin platins wire weighting only 2,4th or again (22 mg.) twice produced the same effect; but this laster weight, when left snapspended, did not suffice to cause a permanent curvature. These trials were made under a bell-glass, so that he loops of thread and wire were

\* Prof. As Gray informs me at the tendrils of P. acerifolia reduce even at a quicker rate in those of P. gracilis; four colutions were completed (the temperature varying from 88°-92° Fahr.) in the following times. 40 m., 45 m., 38½ m., and 46 m. One half-revolution was pernot agitated by the wind. The movement after a tooch is very rupid: I took hold of the lover part of several tendrils, and then tonched their concave tips with a thin twig and washed them carefully through a lear the tips cridently begun to bend after the following intervals—31, 25, 23, 31, 28, 30, 31, and 30 seconds; so that the movement was generally preceptible in half a minute after a nondri; but no one ceasion it was distinctly visible in 25 seconds. One of the tendrils which thus became bent in 31 seconds, had been touched two hours previously and had coiled into a helicit; so that in this interval it had straightened itself and had perfectly recovered its irritability.

To ascertain how often the same tendril would be a several tendril and the pricely recovered its irritability.

isself and and perceive very constant another would become current when towards. It lept a plant in my study, which from being cooler than the hot-home was not very favourable for the experience. The externity was gently rubbed four or five times with a thin stick, and this was done so from as it was observed to have become nearly straight again after having been in action; and in the course of 5 hrs. it answerds to the stimulus 21 times, becoming each time hooked or spiral. On the last occasion, however, the movement was very slight, and soon afterwards permanent spiral contraction commenced. No trials were used during the night, so that the tentral would perhaps have using the contraction of the contracti

I repeated the experiment made on the Echinocystis, and placed several plants of this Passiflora so close together, that their tendrils were repeatedly dragged the whole tendril was dashed about, but they never became curved. The impact from the drops of water was felt far more distinctly on my hand than that from the loops of thread (weighing and of a grain) when allowed to fall on it from a height, and these loops, which caused the tendrils to become curved, had been other tendrils and drops of rain, or that they were from the first rendered sensitive only to prolonged though excessively slight pressure of solid objects, with the exclusion of that from other tendrils. To show the difference in the kind of sensitiveness in different plants and likewise to show the force of the syringe used, I may add that the lightest jet from it instantly caused the leaves of a Mimosa to close; whereas the loop of thread weighing 12nd of a grain, when rolled into a ball and placed gently on the glands at the bases of the leaflets of the Mimosa, caused no action,

Passifora punctata.—The internodes do not move, but the tendrils revolve regularly. A half-grown and very sensitive tendril made three revolutions, opposed to the course of the sun, in 3 hrs. 5 m., 2 hrs. 40 m., and 9 hrs. 50 m., reglesser is wish the resulted was

quickly when nearly full-grown. A plant was placed in front of a window, and, as with twining stems, the light accelerated the movement of the tendril in one direction and retarded it in the other; the semicircle than that required by the semicircle towards the dark end of the room. Considering the extreme tenuity of these tendrils, the action of the light on them is very thin, with the tip slightly curved or hooked. even a single touch causing it to curl inwards; it subsequently straightened itself, and was again ready to act. A loop of soft thread weighing 14th of a grain (4-625 mg.) caused the extreme tip to bend; another time I tried to hang the same little loop on an inclined tendril, but three times it slid off; yet this extratip curl. The tendril, though so sensitive, does not or to a suspended loop of thread. On one occasion I observed a tendril revolving with the convex side of to clasp a stick, against which it scraped; whereas tendrils revolving with the concave side forward,

.

species. The tendrils are thick, long, and stiff; they are sensitive to a touch only on the concave surface towards the extremity. When a stick was placed so one was completed in 3 hrs., and a second in 4 hrs. The internodes do not revolve; nor do those of the hybrid P. floribunda,

Tacsonia manicata.-Here again the internodes do not revolve. The tendrils are moderately thin and long: one made a narrow ellipse in 5 hrs. 20 m., and the next day a broad ellipse in 5 hrs. 7 m. The became just perceptibly curved in 7 m., distinctly in

We have seen that the tendrils in the last three

families, namely, the Vitaceæ, Sapindaceæ and Passiby Mohl), with the tendrils of Brunnichia, one of the Polygonacem. In two or three species of Modecca, one of the Papayacese, the tendrils, as I hear from

This movement, which shortens the tendrils and renders them elastic, commences in half a day, or in a day or two after their extremities have caught some object. There is no such movement in any leafit in the petioles of Tropwolum tricolorum. On the Secondly and thirdly, Bignonia unquis with its close would be quite superfluous. Fourthly, Smilaz aspera offers a more marked exception, as its tendrils are moderately long. The tendrils of Dicentra, whilst the become slightly flexuous; in older plants they are no other exceptions to the rule that tendrils, after spiral contraction. When, however, the tendril of a it cannot; this, however, rarely occurs. In the

I have said that in Corydalis claviculata the end of the leaf or tendril (for this part may be indifferently so called) does not contract into a spire. The branchlets, however, after they have wound round thin trigs, become deeply simons or signag. Moreover the whole end of the petale or tentral, if it sixes mothing, beath after a time abruptly downwards and inwards, showing that its outer surface has gone on growing after the linner surface has caused to grow. That growth is the chief cause of the spiral contraction of tendrils may be adely admitted, as shown by the recent researches of H. de Vries. I will, however, add one little fact in support of this conclusion.

If the short, nearly straight portion of an attached tendril of Passiflora gracilis, (and, as I believe, of other tendrils,) between the opposed spires, be examined, it will be found to be transversely wrinkled in a conspicuous manner on the outside; and this would naturally follow if the outer side had grown more than the inner side, this part being at the same time forcibly prevented from becoming curved. So again the whole outer surface of a spirally wound tendril as the contraction travels from the extremity of a tendril, after it has been stimulated by contact with a support, down to the base, I cannot avoid doubting, effect ought to be attributed to growth. An unattached Cardiospermum, if the contraction commences at the extremity and is quite regular; but if the continued growth of the outer surface is a little lateral, or if the process begins near the base, the terminal portion cantendril then forms a more or less open spire. A similar result follows if the extremity has caught some object, and is thus held fast.

The tendrils of many kinds of plants, if they catch nothing, contract after an interval of several days or weeks into a spire; but in these cases the movement takes place after the tendril has lost its revolving power and hangs down; it has also then partly or wholly lost its sensibility; so that this movement can be of no use. The spiral contraction of unattached tendrils is a much slower process than that of attached ones. Young tendrils which have caught a support and are spirally contracted, may constantly be seen on the same stem with the much older unattached and uncontracted tendrils. In the Echinocystis I have seen a and contracted into beautiful spires, whilst the main branch which had caught nothing remained for many days straight. In this plant I once observed a main branch after it had caught a stick become spirally Generally the tendrils of the Echinocystis begin to contract in from 12 hrs. to 24 hrs. after catching some object; whilst unattached tendrils do not begin to contract until two or three or even more days after all revolving movement has ceased. A full-grown tendril of Passiflora quadrangularis which had caught a stick began in 8 hrs. to contract, and in 24 hrs. formed several spires; a younger tendril, only two-

M

two days after clasping a stick, and in two more days formed several spines. It appears, therefore, that the contraction does not begin until the tendril is grown to nearly its full length. Anothery roung tendril of about the same age and length as the last did not each any object; it negurious its full length in four days; in six additional days it first became flexuous, and in two more days formed one complete spine. This first spine was formed towards the basal cnd, and the contraction steadily but alony progressed towards the apex, but the whole was not cloudy wound up into a spice until; is, until 12 days after the tendril had grown to its full length. The smile recursted of predicties outlies the sine of

The spiral contraction of worlink is quite independent of their power of spontaneously reaching for its occurs in tentrits, such as those of Ladgran greatly form and Anapolephi Indexesses, which do not revolve. It is not necessarily related to the curling of the tips round a support, as we see with the Ampelopsis and Bigmoin expredate, in which the development of adherent discs suffices to cause spiral contraction. Yet in some cases this contraction seems connected with a support; for not only does it soon follow this act, but the contraction generally begins close to the carried extremity, and travels downwards to the base. It however, a tendril be very slack, the whole length almost simultaneously becomes at first flexuous and then strind. Again, the tendrils of some few plants are stress when some contents of the sarriad. Again, the tendrils of some few plants

never contract spirally unless they have first seized hold of some object; if they eated nothing they hap down, remaining straight, until they wither and drop off: this is the case with the tendrils of Hignonia, which consist of medified leaves, and with those of which consist of medified leaves, and with those of which consist of medified leaves, and with those of polamelas. But in the great majority of cases, tendrils which have never come in contact with any object, after a time contract spirally. All those facts taken together, show that the act of classing a support and the spiral contraction of the whole length of the tendril, are phenomena not necessarily connected.

The spiral contraction which ensues after a tendril has caught a support is of high service to the plant; hence its almost universal occurrence with species belonging to widely different orders. When a shoot is inclined and its tendril has caught an object above. the spiral contraction drags up the shoot. When the shoot is upright, the growth of the stem, after the tendrils have seized some object above, would leave it slack, were it not for the spiral contraction which draws up the stem as it increases in length. Thus there is no waste of growth, and the stretched stem ascends by the shortest course. When a terminal branchlet of the tendril of Cobres catches a stick, we have seen how well the spiral contraction successively brings the other branchlets, one after the other, into contact with the stick, until the whole tendril grasps it in an inextricable knot. When a tendril has caught a yielding object, this is sometimes enveloped and

still further secured by the spiral folds, as I have seen with Passiflora quadrangularis; but this action is of little importance.

A far more important service rendered by the spiral contraction of the tendrils is that they are thus made highly elastic. As before remarked under Ampelopsis, the strain is thus distributed equally between the several attached branches; and this renders the whole far stronger than it otherwise would be, as the branches cannot break separately. It is this elasticity which proaway from their supports during stormy weather. I have more than once gone on purpose during a gale to watch a Bryony growing in an exposed hedge, with its tendrils attached to the surrounding bushes; and as the thick and thin branches were tossed to and fro by the wind, the tendrils, had they not been excessively elastic, would instantly have been torn off and the plant thrown prostrate. But as it was, the Bryony safely rode out the gale, like a ship with two anchors down, and with a long range of cable ahead to serve as a spring as she surges to the storm.

When an unattached tendril contracts sprially, the spire always runs in the same direction from tip to base. A tendril, on the other hand, which has caught a support by its extremity, although the same side is concave from end to end, invariably becomes twisted in one part in one direction, and in another part in the opposite direction; the oppositely turned spires belong revenated by a short straight nortion. This curious and symmetrical structure has been noticed by several botancia, but has not been sufficiently explained. It is come without exception with all tendrits which after catching an object contants spirally, but is of course most conspicuous in the longer tendrits. It never comes with meanight tendrits; and when this appears to have occurred, it will be found that the tendril had originally seized some object and had afterwards been four free. Commonly, all the spires at one end of an attached tendril run in one direction, and all those at



A cought tendril of Stryonia diccoa, spirally contracted in reversed directions.

the other end in the opposite direction, with a single short straight portion in the middle; but I have seen a tendril with the spires alternately turning five times

\* See M. Isid. Léen in Bull. Soc. Bot. de France, tom, v. 1838, p. 699. Dr. H. de Vries points out (p. 306) that I have overlooked, in the first edition of this case, the following sentence by Mehl: "After a tendril has caught a

wind into a spire, which, since the tourbil is made fast at both as extremities, must of necessity be in some places to the right, in others to the left." But I am not surprised that this brief scetecce, without any further explanation did not attract my attention.

in opposite directions, with straight pieces between them; and M. Lóon has seen seven or eight used, alternations. Whether the spires turn once or more than once in opposite directions, there are as many turns in the one direction as in the other. For instance, I gustneed ten attached tendrils of the Bryony, the longest with 33, and the shortest with oud 8 spiral turns; and the number of turns in the one direction was in every case the same (within one) as in the opposite direction.

The explanation of this curious little fact is not difficult. I will not attempt any geometrical reasoning, but will give only a practical illustration. In doing this, I shall first have to allude to a point which was almost passed over when treating of Twining-plants. If we hold in our left hand a bundle of parallel strings, we can with our right hand turn these round and round, thus imitating the revolving movement of a twining plant, and the strings do not become twisted. But if we hold at the same time a stick in our left hand, in such a position that the strings become twisted. Hence a straight coloured line, painted along the internodes of a twining plant before it has wound round a support, becomes twisted or spiral after it has wound round. I painted a red line on the straight internodes of a Humulus, Mikania, Ceropegia, Convolvulus, and Phascolus, and saw it become twisted as the plant wound round a stick. It is possible that the stems of some plants by spontaneously turning on their own axes, at the proper rate and in the proper direction, might avoid becoming twisted; but I have

In the above illustration, the parallel strings were wond round a stake; but this is by no mean encessary, for if wound into a hellow cell (as can be done with a narrow align of elastic paper) there is the same inevitable twisting of the axis. When, therefore, a free tended coals intell into a spirs, it must either become twisted along its whole length (and this never occurs), or the free extensity must turn round as many times as there are spirss formed. It was hardly necessary to observe this fact; but I did so by affixing little paper vanes to the extreme points of the tendrils of Enkinospitis and Passifore paradrengularis; and as the tendril contracted itself into successive spirss, the vane slowly revolved.

We can now understand the meaning of the spices being invariablely turned in opposite directions, in tendrils which from having caught some object are fixed at both ends. Let us suppose a caught tendril to make thirty spiral turns all in the same direction; the inevitable result would be that it would become twisted thirty times on its own axis. This twisting would not only require considerable force, but, as I know by trial, would burst the tendril before the thirty turns were completed. Such cases more really occur; for, as already stated, when a tendril have conglet on the control of the contro

compensated by the twisting in the opposite direction. We can further see how the tendency is given to make to the right or to the left, which are first made. Take a piece of string, and let it hang down with the lower end fixed to the floor; then wind the upper end (holding the string quite loosely) spirally round a perthe string; and after it has been sufficiently twisted, it curves running in an opposite direction to those round the pencil, and consequently with a straight piece of string between the opposed spires. In short, we have given to the string the regular spiral arrangement of a tendril caught at both ends. The spiral contraction generally begins at the extremity which has clasped a support; and these first-formed spires give a twist to the axis of the tendril, which necessarily inclines the basal part into an opposite spiral curvature. I cannot resist giving one other illustration, though superfluous: when a haberdasher winds up ribbon for a customer, he does not wind it into a single coil; for, if he did, the ribbon would twist itself as many times as there were coils: but he winds it into a figure of eight on his thumb and little finger, so that he alternately takes turns in opposite directions, and thus the ribbon is not twisted. So it is with tendrils, with this sole difference, that they take several consecutive turns in one direction and then the same number in an opposite direction; but in both cases the self-twisting is

Summary on the Nature and Action of Tendrils.

With the majority of tendril-bearing plants the young internodes revolve in more or less broad ellipses, like those made by twining plants; but the figures described, when carefully traced, generally form irregular ellipsoidal spires. The rate of revolution varies from one to five hours in different species, and consequently is in some cases more rapid than with any twining plant, and is never so slow as with those many twiners which take more than five hours for each revolution. The direction is variable even in the same individual plant, In Passiflora, the internodes of only one species have the power of revolving. The Vine is the weakest revolver observed by me, apparently exhibiting only a trace of a former power. In the Eccremocarpus the movement is interrupted by many long pauses. Very few tendril-bearing plants can spirally twine up an upright stick. Although the power of twining has generally been lost, either from the stiffness or shortness of the internodes, from the size of the leaves, or from some other unknown cause, the revolving movement of the stem serves to bring the tendrils into contact with surrounding objects.

The tendrils themselves also spontaneously revolve. The movement begins whilst the tendril is young, and move much slower than the internodes. Generally,

the internodes and tendrils revolve together at the same rate; in Cissus, Cobeea, and most Passiflore, the tendrils alone revolve: in other cases, as with Lathuren the motionless tendrils; and, lastly (and this is the fourth possible case), neither internodes nor tendrils spontaneously revolve, as with Lathyrus grandifloru and Ampelopsis. In most Bignonias, Eccremocarpus, Mutisia, and the Fumariaceæ, the internodes, petioles and tendrils all move harmoniously together. In every case the conditions of life must be favourable in order that the different parts should act in a perfect

Tendrils revolve by the curvature of their whole length, excepting the sensitive extremity and the base, which parts do not move, or move but little. The movement is of the same nature as that of the revolving internodes, and, from the observations of Sachs and H. de Vries, no doubt is due to the same cause, namely, the rapid growth of a longitudinal band, which travels round the tendril and successively bows each part to the opposite side. Hence, if a line be painted along that surface which happens at the time to be convex, the line becomes first lateral, then concave, then lateral, and ultimately again convex-This experiment can be tried only on the thicker tendrils, which are not affected by a thin crust of dried paint. The extremities are often slightly curved or hooked, and the curvature of this part is never reversed; in this respect they differ from the extremities of twining shoots, which not only reverse but curve themselves in a creater degree than the lower part. In most other respects a tendril acts as if it were one of several revolving internodes, which all move together by successively bending to each point of the compass. There is, however, in many cases this unimportant difference, that the curving tendril is separated from the curving internode by a rigid petiole. With most tendril-bearers the summit of the stem or shoot projects above the point from which the tendril arises; and it is generally bent to one side. so as to be out of the way of the revolutions swept by the tendril. In those plants in which the terminal shoot is not sufficiently out of the way, as we have seen with the Echinocystis, as soon as the tendril comes in its revolving course to this point, it stiffens and straightens itself, and thus rising vertically up passes over the obstacle in an admirable manner.

All nearlist are sensitive, but in various degrees, to contact with an object, and curve towards the touched side. With sevent plants a single touch, so slight as only just to move the highly flexible tendril, is enough to induce curvature. Paraiflear generalize passesses the most sensitive tendrils which I have observed: a bit of platina wire judt of a grain (123 mg) in weight, agently placed on the concave point, caused a tendril to become hooked, as did a loop of soft, thin cotton thread weighing judt of a grain (202 mg.) With the tendrils of several other plants, loose weighing. Jud of a grain (4.05 mg.) sufficed. The point of a tendril of Passiflora gracilis began to move distinctly in 25 seconds after a touch, and in many cases after 30 seconds. Asa Grav also saw movement in the tendrils of the Cucurbitaceous genus, Sievos, in 30 seconds. The tendrils of some other plants, when lightly rubbed, moved in a few minutes; with Dicentra in half-an-hour; with Smilax in an hour and a quarter or half; and with Ampelopsis still more slowly, The curling movement consequent on a single touch continues to increase for a considerable time, then ceases; after a few hours the tendril uneurls itself, and is again ready to act. When the tendrils of several kinds of plants were caused to bend by extremely light weights suspended on them, they seemed to grow accustomed to so slight a stimulus, and straightened themselves, as if the loops had been removed. It makes no difference what sort of object a tendril touches, with the remarkable exception of other tendrils and drops of water, as was observed with the extremely sensitive-tendrils of Passiflora gracilis and of the Echinocustis. I have, however, seen tendrils of the Bryony which had temporarily caught other tendrils, and often in the case of the vine,

Tendrils of which the extremities are permanently and slightly curved, are sensitive only on the concave surface; other tendrils, such as those of the Cobras (though furnished with horny hooks directed to one side) and those of Cisus discolor, are sensitive on all sides. Hence the tendrils of this latter plant, when stimulated by a touch of equal force on opposite sides, did not beaul. The inferior and lateral strates of the tendrist of Methics are sensitive, but not the upper surface. With benneded tendrisk, the several branches act alike; but in the Husheyas the lateral spur-like branch does not acquire (for excellent reasons which have been explained) its sensitiveness nearly so soon as the main branch. With most tendrist the lower or basal part is either not at all sensitive, or sensitive only to prolonged centers. We thus see that the sensitiveness of tendrisk is a special and localized capacity. It is quite independent of the power of spentaneously revolving; for the curling of the terminal portion from a touch does not in the losset to the control of the control of the control of the control and its close allies, the protoke of the lowers, as well as the tendrisk are sensitive to a touch the control of the lowers, as well as the tendrisk are sensitive to a touch the

as an stantant, are sensority to a tooler. Twining plants when they come into connect with a staick, card round it invariably in the direction of their revolving movement; but tendrits card indifferently in either side, in accordance with the position of the strick and the side which is first tooled. The chapping movement of the extremity is apparently not steady, and the contraction of the strict and the side contraction of the strict and the side contraction of the strict and the side states are to the strict and the side of the Deblacoptain some side with the tendrits of the Deblacoptain solving and a mooth of side of the Deblacoptain solving and a mooth of side.

As with a few exceptions tendrils spontaneously revolve, it may be asked,—why have they been endowed with sensitiveness?—why, when they come into contact wind round it? One reason may be that they are in most cases so flexible and thin, that when brought into contact with any object, they would almost certainly yield and be dragged onwards by the revolving movement. Moreover, the sensitive extremities have no revolving power as far as I have observed, and could not by this means curl round a support, With twining plants, on the other hand, the extremity spontaneously bends more than any other part; and this is of high importance for the ascent of the plant, as may be seen on a windy day. It is, however, possible that the slow movement of the basal and stiffer parts But I hardly attended sufficiently to this point, and it would have been difficult to distinguish between a movement due to extremely dull irritability, from the arrestment of the lower part, whilst the upper part continued to move onwards,

Tendrils which are only three-fourths grown, and perhaps even at an cultier age, but not white stremely young, have the power of revolving and of grasping any object which they touch. These two capacities are generally acquired at about the same period, and both fall when the tendril is fall grown. But in Chose and Passiftons powdate the tendrils begin to revolve in a useless manner, before they have become sensitive. In the Echinocystic they retain their sensitiveness for some time after the have exacted to 803

revolve and after they have sunk downwards; in this position, even if they were able to seize an object, such power would be of no service in supporting the stem. It is a run exicumstance thus to detect any super-fluity or imperfection in the action of tendrils—organs which are so excellently adapted for the functions which they have to perform; but we see that they are not always perfect, and it would be such to assume a contract of perfection.

Some tendrils have their revolving motion accelerated as the super-fluid perfection.

Some tendrik have their revolving motion accelerated or retarded, in moving to or from the light; others, as with the Pea, some indifferent to its action; others move steadily from the light to the dark, and this adds them it an important manner in finding a support. For the contract of the contract of the contract of the first of the contract of the test and turn along to a to be lengther finer branches and hooks into close contact with any dark surface, or into everiese and holes.

A short time after a tendril has caught a support, it contracts with some rare exceptions into a spire; but the manner of contraction and the several important advantages thus gained have been discussed so lately, that nething need here be repeated on the subject. Tendrils soon after catching a support grow much stronger and thicker, and sometimes more durable to a viscoger and thicker, and sometimes more durable to a wonderful degree; and this shows how much their internal tissues must be changed. Occasionally it is

the part which is wound round a support which chiefly becomes thicker and stronger; I have seen, for instance, this part of a tendril of Bignonia equinoctialit vice as thick and rigid as the free basal part. Tendrils which have caught nothing soon shrink and wither; but in some species of Bignonia they disarticulate and fall off like leaves in autumn.

Any one who had not closely observed tendrils of many kinds would probably infer that their action was uniform. This is the case with the simpler kinds, which simply curl round an object of moderate thickness, whatever its nature may be. But the genus Bignonia shows us what diversity of action there may be between the tendrils of closely allied species. In all the nine species observed by me, the young internodes revolve vigorously; the tendrils also revolve, but in some of the species in a very feeble manner; and lastly the petioles of nearly all revolve though with unequal power. The petioles of three of the species, and the tendrils of all are sensitive to contact. In the first-described species, the tendrils resemble in shape a bird's foot, and they are of no service to the stem in spirally ascending a thin upright stick, but they can seize firm hold of a twig or branch. When

\* Fachs, however ('Text-Book of Botany, Eng. Translation, 1875, p. 280), has shown that which I overlocked, namely, that the tookida of different species are

adapted to clasp supports of different thicknesses. He further shows that after a tendril has clasped a support it subsequently the stem twines round a somewhat thick stick, a slight degree of sensitiveness possessed by the petioles is brought into play, and the whole leaf together with the tendril winds round it. In B. unquis the petioles are more sensitive, and have greater power of movement than those of the last species; they are able, together with the tendrils, to wind inextricably round a thin upright stick; but the stem does not twine so well. B. Tweedyana has similar powers, but in addition, emits aerial roots which adhere to the wood. In B. venusta the tendrils are converted into elongated three-pronged grapnels, which move spontaneously in a conspicuous manner; the petioles, however, have lost their sensitiveness. The stem of this species can twine round an upright stick, and is aided in its ascent by the tendrils seizing the stick alternately some way above and then contracting spirally. In B. littoralis the tendrils, petioles, and internodes, all revolve spontaneously. The stem, however, cannot twine, but ascends together, which then contract into a spire. The tips of these tendrils become developed into adhesive discs. B. speciosa possesses similar powers of movement as the last species, but it cannot twine round a stick though it can ascend by clasping the stick horizontally with one or both of its unbranched tendrils. These tendrils continually insert their pointed ends into minute crevices or holes, but as they are always withdrawn by the subsequent spiral contraction, the habit seems to us in our ignorance useless. Lastly, the stem of B. caproclate twines imperfectly; the nuschbranchet tenfris revolve in a capricious namuer, and bend from the light to the dark; their hooked extensities, even whilst immature, crawl into crevices, and, when mature, seize any thin projecting point; in either case they develop adhexive discs, and these have the power of enveloping the finest fibres.

themselves neatly on any surface with which they stems, like the culms of a grass, which they aftercontraction of all the branches. In Cobea the terminate in sharp, hard, double, little hooks, with both points directed to the same side; and these turn they come into contact. The tips of the branches also crawl into dark crevices or holes. The tendrils of revolving; the tendrils are but little sensitive to contact: their hooked extremities cannot seize thin objects; they will not even clasp a stick, unless in extreme need of a support; but they turn from the These adhere by the secretion of some cement to wall, or even to a polished surface; and this is more than the discs of the *Bignonia capreolata* can effect.

The rapid development of these adherent discs is one of the most remarkable poculiarities possessed by any tendrik. We have seen that such discs are formed by two species of Bignonia, by Ampelopais, and, ecoording to Naudin, by the Cucurbitacous genus Poponquia albertass. In Augurita the lower surface of the tendril, after it has wound round a stick, forms a causely cellular layer, which closely fits the wood, but is not adherent; whilst in Hanburya a similar layer is adherent. The growth of these cellular outgrowths depends (except in the case of the Bapkolphina and of one species of Ampelopais, on the stimulus from contact. It is a singular fact that three families, as wilely distinct as the Bignoniaces, Vitasce, and Courtificaces, about 1 possess species with tendrils having this centralish power.

Sachs attributes all the movements of tendris to mpid growth on the side opposite to that which becomes concave. These movements consist of revolving mutation, the bending to and from the light, and in opposition to gravity, those caused by a touch, and spiral contraction. It is rash to differ from so great an authority, but I cannot believe that one at least of

<sup>\*</sup> Annales des Sc. Nat. Bot. 4th series, tom. xii. p. 89.

Online: By permission of the Trustees of the Natural His

these movements-curvature from a touch-is thus caused.\* In the first place it may be remarked that the movement of nutation differs from that due to a touch, in so far that in some cases the two powers are acquired by the same tendril at different periods of growth; and the sensitive part of the tendril does not seem capable of nutation. One of my chief reasons for doubting whether the curvature from a touch is the result of growth, is the extraordinary rapidity of the movement. I have seen the extremity of a tendril of Passiflora gracilis, after being touched, distinctly bent in 25 seconds, and often in 30 seconds; and so it is with the thicker tendril of Sievos. It appears hardly credible that their outer surfaces could have actually grown in length, which implies a permanent modification of structure, in so short a time. The growth, moreover, on this view must be considerable, for if the touch has been at all rough the extremity is coiled in two or three minutes into a spire of several turns.

When the extreme tip of the tendril of Echinocystis caught hold of a smooth stick, it coiled itself in a few hours (as described at p. 132) twice or thrice round

unaffected by exposure for 1½ hrs.
to a rather large dose of sulphusic ethor. In this respect they
present a wenderful contrast with
Drosem, owing no doubt to the
presence of absorbent glands in
the latter plant.

Trecurrent to use time the movement of nutation and that from a touch might be differently affected by anasotheties, in the same manner as Paul Bert has shown to be the case with the sleep-movements of Minress and those from a touch. I tried the common year and Passifron arm-

the stick, apparently by an undulatory movement. At first I attributed this movement to the growth of the outside; black marks were therefore made, and the interspaces measured, but I could not thus detect any increase in length. Hence it seems probable in this case and in others, that the curvature of the tendril from a touch depends on the contraction of the cells along the concave side. Sachs himself admits\* that " if the growth which takes place in the entire tendril "at the time of contact with a support is small, a " considerable acceleration occurs on the convex sur " face, but in general there is no elongation on the " concave surface, or there may even be a contraction : " in the case of a tendril of Cucurbita this contraction " amounted to nearly one-third of the original length," In a subsequent passage Sachs seems to feel some difficulty in accounting for this kind of contraction. It must not however be supposed from the foregoing remarks that I entertain any doubt, after reading De Vries' observations, about the outer and stretched surfaces of attached tendrils afterwards increasing in length by growth. Such increase seems to me quite of growth. Why a delicate touch should cause one side of a tendril to contract we know as little as why, on the view held by Sachs, it should lead to extraordinarily rapid growth of the opposite side. The chief or sole reason for the belief that the curvature of

<sup>\* \*</sup>Text-Book of Botany, 1875, p. 779.

a tendril when touched is due to rapid growth, seems to be that tendrils lose their sensitiveness and power of movement after they have grown to their full length; but this fact is intelligible, if we bear in mind that all the function of a tendril raw subpert to drag up the terminal growing shoot towards the light. Or what use would it be, if an old and full-grown tendril, arising from the lower part of a shoot, were to retain its power of classing a support? This would be of no use; and we have seen with tendrils so many instances of close adaptation and of the economy of means, that we may feel assured that they would acquive irribablity and the power of classing a support at the proper age—namely, youth—and would not uselessly retain such power boyen the proper age.

## CHAPTER V

HOOK AND ROOT-CLIMBERS.—CONCLUDING REMARKS.

lants climbing by the aid of hooks, or merely scrambling over other plants—Root-climbers, adhesive matter secreted by the rootlets— General conclusions with respect to climbing plants, and the stages

Hook-Clinders——In my introductory remarks, i assaw that, besides the two first great classes of climbing plants, namely, those which twine round a support, and those endowed with irritability enabling them to seize hold of objects by means of their peticles or tendrils, there are two other classes, book-clinders and root-clinders. Many plants, moreover, as Fritz Maller has remarked, "elimb or seramble up thickest in a still more simple fashion, without any special aid, excepting that their leading aboots are generally long and flexible. It may, however, be suspected from what follows, that these shoots is more cases tend to avoid the light. The few hook-clinders which I have observed, namely, Gallium serzies, Robus sustraint, and some climbing

\* Journal of Linn, Soc, vol. ix, p. 348, Professor G. Jacoper has well remarked (\* In Sachen Darwin's, inabezoudere contra Wiganal, 1874, p. 106), that it is highly characteristic of chimbing plants to produce thin, closigated, and flexi-

taller species or trees, are naturally
those which would be developed
into climbers; and such plants,
from stretching towards the light,
and from not being much agitated
by the wind, tend to produce long,
this and forther house.

Roses, exhibit no spontaneous revolving movement. If they had possessed this power, and had been capable of twining, they would have been placed in the class of Twiners; for some twiners are furnished with spines or hooks, which aid them in their ascent. For instance, the Hop, which is a twiner, has reflexed hooks as large as those of the Galium; some other twiners have stiff reflexed hairs; and Dipladenia has a circle of blunt spines at the bases of its leaves. I have seen only one tendril-bearing plant, namely, Smilaz aspera, which is furnished with reflexed spines; but this is the case with several branch-climbers in South Brazil and Cevlon; and their branches graduate into true tendrils. Some few plants apparently depend solely on their hooks for climbing, and yet do so efficiently, as certain palms in the New and Old Worlds. Even some climbing Roses will ascend the walls of a tall house, if covered with a trellis. How this is effected I know not; for the young shoots of one such Rose, when placed in a pot in a window, bent irregularly towards the light during the day and from the light during the night, like the shoots of any common plant : so that it is not easy to understand how they could have got under a trellis close to the wall."

\* Professor Asa Gray has explained, as it would appear, this plained, as it would appear, this difficulty in his roview (American Journal of Science, vol. Xl. Sept. 1985, p. 282) of the present work. He has observed that the strong summer shoots of the Michigan rose (Ross settleren) are strongly

and away from the light, so that they would be almost sure to place themselves under a trellis. He adds that the lateral shoots, made on the following spring, emerged from the trellis as they sought the light.

Root-climbers.—A good many plants come under this class, and are excellent climbers. One of the most remarkable is the Marcgravia umbellata, the stem of which in the tropical forests of South America, as I hear from Mr. Spruce, grows in a curiously flattened manner against the trunks of trees; here and there it puts forth claspers (roots), which adhere to the trunk, and, if the latter be slender, completely embrace it. When this plant has elimbed to the light, it produces free branches with rounded stems, clad with sharpthose borne by the stem as long as it remains adherent. This surprising difference in the leaves, I have also observed in a plant of Maregravia dubia in my hothouse. Root-climbers, as far as I have seen, namely, the Ivy (Hedera helix), Ficus repens, and F. barbatus, have no power of movement, not even from the light to the dark, As previously stated, the Hoya carnosa (Asclepiadaceae) is a spiral twiner, and likewise adheres by rootlets even to a flat wall. The tendril-bearing Bignonia Tweedyana emits roots, which curve half round and adhere to thin sticks. The Tecoma radicans (Bignoniacem), which is closely allied to many spontaneously revolving species, climbs by rootlets; nevertheless, its young shoots apparently move about more than can be accounted for by the varying action of

I have not closely observed many root-climbers, but can give one curious fact. Ficus repens climbs up a wall just like Ivy; and when the young rootlets are made to press lightly on slips of glass, they emit after about a week's interval, as I observed several times, minute drops of clear fluid, not in the least milky like that exuded from a wound. This fluid threads. It has the remarkable property of not soon slightly spread out on glass, and I scattered on it some minute grains of sand. The glass was left exposed in a drawer during hot and dry weather, and if the fluid had been water, it would certainly have dried in a few minutes; but it remained fluid, closely much longer it would have remained I cannot say. Some other rootlets were left in contact with the glass for about ten days or a fortnight, and the drops of secreted fluid were now rather larger, and so viscid that they could be drawn out into threads. Some other rootlets were left in contact during twenty-three days, and these were firmly cemented to the glass, Hence we may conclude that the rootlets first secrete a slightly viscid fluid, subsequently absorb the watery parts, (for we have seen that the fluid will not dry by itself,) and ultimately leave a cement. When the rootlets were torn from the glass, atoms of yellowish matter were left on it, which were partly dissolved volatile fluid was rendered very much less volatile by

As the bisulphide of carbon has a strong power

of softening indurated caoutchoue, I soaked in it during a short time several rootlets of a plant which had grown up a plaistered wall; and I then found excessively elastic matter, precisely like caoutchouc, attached to two sets of rootlets on the same branch, These threads proceeded from the bark of the rootlet at one end, and at the other end were firmly attached to particles of silex or mortar from the wall. There could be no mistake in this observation, as I played with the threads for a long time under the microscope, drawing them out with my dissecting-needles and letting them spring back again. Yet I looked repeatedly at other rootlets similarly treated, and could never again discover these elastic threads. I therefore infer that the branch in question must have been slightly moved from the wall at some critical period, whilst the secretion was in the act of drying, through the absorption of its watery parts. The genus Ficus abounds with caoutehoue, and we may conclude from the facts just given that this substance, at first in solution and ultimately modified into an unclastic cement," is used by the Ficus repens to cement its rootlets to any surface which it ascends. Whether other plants, which climb by their rootlets, emit

\* Mr. Spiller has recently shown (Chemical Society, Feb. 16, 1865), in a paper on the oxidation of india-rubber or exomethous, that a fine state of division to the air, graviually becomes converted into brittle, resinous matter, very similar to shell-lac. Ivy, placed against glass, barely adhered to it, yet secreted a little yellowish matter. I may add, that the rootlets of the Maregravia dubia can adhere firmly to smooth painted wood.

Vanilla aromatica emits aërial roots a foot in length, which point straight down to the ground. According to Mohl (p. 49), these crawl into crevices, and when they meet with a thin support, wind round it as do tendrils. A plant which I kept was young, and did not form long roots; but on placing thin sticks in contact with them, they certainly bent a little to that side, in the course of about a day, and adhered by their rootlets to the wood: but they did not bond their downward course. It is probable that these slight movements of the roots are due to the quicker growth of the side exposed to the light, in comparison with the other side, and not because the roots are sensitive to contact in the same manner as true tendrils. According to Mohl, the rootlets of certain species of Lucopodium act as tendrils.\*

\* Fritz Miller informs mo that he saw in the forests of South Braill numerous black strings, from some lines to nearly an inch in diameter, winding spirally round the trunks of gigantic trees. At first sight he thought that they were the stem of twining plants which were thus are not make the same that the same range for the same that the way the ways formed that they way the ways formed that they way the acirial roots of a Philideduction which grew on the beambers above. These roots therefore seem to be true twincers, though they use their powers to descend, instead of to ascend like twining plants. The airial roots of some other species of Philodendron has vertically downwards, sometimes for a length of more than fifty feet.

## Concluding Remarks on Climbing Plants.

Plants become climbers, in order, as it may be presumed, to reach the light, and to expose a large surface of their leaves to its action and to that of the free air. This is effected by climbers with wonderfully little expenditure of organized matter, in comparison with trees, which have to support a load of heavy branches by a massive trunk. Hence, no doubt, it arises that there are so many climbing plants in all quarters of the world, belonging to so many different orders. These plants have been arranged under four bushes without any special aid. Hook-climbers are the least efficient of all, at least in our temperate countries, and can climb only in the midst of an entangled mass of vegetation. Root-climbers are excellently adapted to ascend naked faces of rock or trunks of trees; when, however, they climb trunks they are compelled to keep much in the shade: they cannot pass from branch to branch and thus cover the whole summit of a tree, for their rootlets require long-continued and close contact with a steady surface in order to adhere. The two great classes of twiners and of plants with sensitive organs, namely, leafclimbers and tendril-bearers taken together, far exceed in number and in the perfection of their mechanism the climbers of the two first classes. Those which have the power of spontaneously revolving and of grasping objects with which they come in contact, easily pass from branch to branch, and securely ramble over a wide, sun-lit surface.

The divisions containing twining plants, leaf-climbers, one another, and nearly all have the same remarkable power of spontaneously revolving. Does this gradato one subdivision have actually passed during the lapse of ages, or can pass, from one state to the other? Has, for instance, any tendril-bearing plant assumed climbers alone, the idea that they were primordially twiners is forcibly suggested. The internodes of all, without exception, revolve in exactly the same manner as twiners; some few can still twine well, and many others in an imperfect manner. Several leafclimbing genera are closely allied to other genera which are simple twiners. It should also be observed, would be of comparatively little use to a plant, the leaves are brought into contact with a support: by its leaves. On the other hand, revolving internodes, without any other aid, suffice to give the power of climbing; so that it seems probable that leafclimbers were in most cases at first twiners, and subsequently became capable of grasping a support; and this, as we shall presently see, is a great additional advantage. From analogous reasons, it is probable that all

tendril-bearers were primordially twiners, that is, are For the internodes of the majority revolve; and, in a few species, the flexible stem still retains the capacity bearers have undergone much more modification than leaf-climbers: hence it is not surprising that their have been more frequently lost or modified than in the case of leaf-climbers. The three great tendrilbearing families in which this loss has occurred in the most marked manner, are the Cucurbitaceae, Passifloraceze, and Vitaceze. In the first, the internodes revolve; but I have heard of no twining form, with the exception (according to Palm, p. 29.52) of Momor-In the two other families I can hear of no twiners: this power being confined to the tendrils. The internodes, however, of Passiflora gracilis have the power in a perfect manner, and those of the common Vine in an imperfect degree: so that at least a trace of the supposed primordial habit has been retained by some members of all the larger tendril-bearing groups.

On the view here given, it may be asked, Why have the species which were aboriginally twiners been converted in so many groups into leaf-climbers or tendrilbearers? Of what advantage has this been to them? Why did they not remain simple twiners? We can see several reasons. It might be an advantage to a plant to acquire a thicker stem, with short internodes bearing many or large leaves; and such stems are ill fitted for twining. Any one who will look during easily blown from their support; not so with tendrilgrasp their support by a much more efficient kind of movement. In those plants which still twine, but at the same time possess tendrils or sensitive petioles, as some species of Bignonia, Clematis, and Tropæolum, it can readily be observed how incomparably better they grasp an upright stick than do simple twiners. Tendrils, from possessing this power of grasping an object, can be made long and thin; so that little organic matter is expended in their development, and yet they sweep a wide circle in search of a support. Tendril-bearers can, from their first growth, ascend along the outer branches of any neighbouring bush, and they are thus always fully exposed to the light; twiners, on the contrary, are best fitted to ascend bare stems, and generally have to start in the shade. Within tall and dense tropical forests, twining plants would probably succeed better than most kinds of tendril-bearers; but the majority of twiners, at least in our temperate regions, from the nature of their revolving movement, cannot ascend thick trunks, whereas this can be affected by tendrilbearers if the trunks are branched or bear twigs, and

The advantage gained by climbing is to reach the light and free air with as little expenditure of organic matter as possible; now, with twining plants, the stem instance, I measured the stem of a kidney-bean, which three feet in length: the stem of a pea, on the other hand, which had ascended to the same height by the reached. That this saving of the stem is really an tendrils, generally making more open spires than those made by simple twiners. Moreover, the plants thus aided, after taking one or two turns in one direction, generally ascend for a space straight, and then reverse the direction of their spire. By this means samé length of stem, than would otherwise have been themselves at intervals by their clasping petioles or

We have seen that tendrils consist of various organs in a modified state, namely, leaves, flower-peduncles, branches, and perhaps stipules. With respect to leaves, the evidence of their modification is ample. In young plants of Bignonia the lower leaves often romain quite unchanged, whilst the upper ones have their terminal leaflest converted into perfect tendrilly, in Ecroscoppes I have seen a single latent llurance of a tendril replaced by a perfect leaflet; in Férie antien, on the other hand, leaflets are smortines replaced by tendril-branches; and many other such cases could be given. But he who believes in the slow modification of species will not be centered simply to assert and the monological nature of different kinds of tendrilly; he will wish to learn, as far as is possible, by what actual store leaves, flower-percedureds, &c., here

had their functions wholly changed, and have come to

serve merely as prehensile organs. In the whole group of leaf-climbers abundant evidence has been given that an organ, still subservtouch, and thus grasp an adjoining object. With several leaf-climbers the true leaves spontaneously revolve; and their petioles, after clasping a support grow thicker and stronger. We thus see that leaves may acquire all the leading and characteristic qualities of tendrils, namely, sensitiveness, spontaneous movement, and subsequently increased strength. If their blades or laminæ were to abort, they would form true tendrils. And of this process of abortion we can follow every step, until no trace of the original nature of the tendril is left. In Mutisia clematis, the tendril, in leaflets, but vestiges of the laminæ are still occasionally retained. In four genera of the Fumariaceæ we can follow the whole process of transformation. The terminal leaflets of the leaf-climbing Fumaria officinalis are not smaller than the other leaflets; those of the leafclimbing Adlumia cirrhosa are greatly reduced; those of Corndalis claviculata (a plant which may indifferently be called a leaf-climber or a tendril-bearer) are either reduced to microscopical dimensions or have their blades wholly aborted, so that this plant is actually in a state of transition; and, finally, in the Dicentra the tendrils are perfectly characterized. If, therefore, we could behold at the same time all the progenitors of Dicentra, we should almost certainly see a series like that now exhibited by the above-named three genera. In Tropwolum tricolorum we have another kind of passage; for the leaves which are first formed on the young stems are entirely destitute of laminæ, and must be called tendrils, whilst the later formed leaves have well-developed laminæ. In all cases the acquirement of sensitiveness by the mid-ribs of the leaves appears to stand in some close relation with the abortion of their laming or blades.

On the view here given, leaf-climbers were primordially twines, and tendrib-beares (when formed of modified leaves) were primordially leaf-climbers. The latter, therefore, are intermediate in nature between twiens and tendrib-beares, and ought to be related to both. This is the case: thus the several leaf-climbing species of the Antirrihneou, of Solamum, Occoules, and Glorisus, have within the same family and even within the same genus, relatives which are twiners. In the genus Mikania, there are leaf-elimbing and twining species. The leaf-elimbing species of Clematis are very cloudy allied to the tendri-bearing Nanavelian. The Fumarinees include closely allied genera which are leaf-elimbers and bradit-bearers. Lastly as species of Biguonia is at the same time both a leaf-elimber and a tendribearer; and other closely allied species are twiners.

peduncles. In this case we likewise have many interesting functional states. The common Vine (not to mention the Gorliopersum) gives in every possible gradation between a perfectly developed tendral and a flower-pedunch covered with flowers, yet furnished with a branch, forming the flower-tendral. When the latter itself bears a few flowers, as we know sometimes is the case, and still retains the power of clasping a support, we see an endy condition of all those tendrals which have been formed by the modification of flowerpeduncles.

According to Mohl and others, some tendrils consist of modified humbers: I have not observed any such cases, and know nothing of their transitional states, but these have been fully described by Pritz Müller. The genus Lophospermum also shows us how such a transition is possible; for its bennehes spontaneously revolve and are sensitive to contact. Hence, if the leaves on some of the branches of the Lophospermum were to abort, these branches would be converted into true tendrils. Nor is there anything improbable in certain branches alone being thus modified, whilst others remained unaltered; for we have seen with certain varieties of Phaseolus, that some of the branches are thin, flexible, and twine, whilst other branches on the same plant are stiff and have no such power.

peduncle first became sensitive to a touch, and acquired the power of bending towards the touched side, we get no certain answer. Nevertheless an obthis side. And it is young petioles and tendrils, whatever their homological nature may be, which and incipient capacity, which capacity, as far as we can see, is of no service to ordinary plants. If we flower-peduncles of climbing plants first acquired their power of spontaneously revolving, or, to speak only remark that the power of moving, both spontaneously and from various stimulants, is far more

remarkable Gezeke im

common with plants, than is generally supposed to be the case by those who have not attended to the subject. I have given one remarkable instance, namely that of the Maurandia semperflorens, the young flower-peduncles of which spontaneously revolve in very small circles, and bend when gently rubbed to the touched side; yet this plant certainly does not profit by these two feebly developed powers. A rigorous examination of other young plants would probably show slight spontancous movements in their stems, petioles or peduncles, as well as sensitiveness to a touch." We see at least that the Maurandia might, by a little augmentation of the powers which it already possesses, come first to grasp a support by its flower-peduncles, and then, by the abortion of some of its flowers (as with Vitis or Cardiospermum), acquire perfect tendrils. There is one other interesting point which deserves

There is one other interesting point which asserves notice. We have seen that some tendrils over their origin to modified leaves, and others to modified flowerpeduncles; so that some are foliar and others axial in their nature. It might therefore have been expected that they would have presented some difference in function. This is not the case. On the contrary, they

\* Such slight spontaneous movements, I now find, have been for some time known to occur, for instance with the flower-sterns of Brassica supus and with the leaves of many plants: Sachis' 'Text-Book of Botany' 1875, pp. 769, 785. Fitz Müller also has

ms shown in relation to our present subence jost (Jenaischen Zeitschrift), Ha zr, V. Heft 2, p. 133) that the stems, ms whilst young, of an Alisma sad be of a Linum are centimally se' performing slight movements to p. all points of the compass, like as those of elimbing plants, present the most complete identity in their several characteristic powers. Tendrils of both kinds spontaneously revolve at about the same rate. Both, when touched, bend quickly to the touched side, and afterwards recover themselves and are able to act again. In both the sensitiveness is either confined to one side or extends all round the tendril. Both are either is seen in the foliar tendrils of Bignonia capreolata and in the axial tendrils of Ampelopsis. The tips of the tendrils in these two plants become, after contact, enlarged into discs, which are at first adhesive by the secretion of some cement. Tendrils of both kinds, soon after grasping a support, contract spirally ; they then increase greatly in thickness and strength. When we add to these several points of identity the fact that the petiole of Solanum jasminoides, after it has clasped a support, assumes one of the most characteristic features of the axis, namely, a closed ring of woody vessels, we can hardly avoid asking, whether the difference between foliar and axial organs can be

We have attempted to trace some of the stages in the genesis of climbing plants. But, during the endless fluctuations of the conditions of life to which all organic beings have been exposed, it might be expected that some climbing plants would have lest

<sup>\*</sup> Mr. Herbert Spencer has recently argued (\* Principles of

of damental distinction between

the habit of climbing. In the cases given of certain South African plants belonging to great twing families, which in their native country mover twine, but reasumen this habit when cultivated in England, we have a case in point. In the leaf-climbing Clemeirs flammands, and in the tendril-bearing Vine, we see loss in the power of climbing, but only a remnant of the revolving power which is indispensable to all trees, and is so common as well as so advantageous to most climbers. In Teoma varieties, nor of the Bignoniances, we see a last and doubtful trace of the power of revolving.

With respect to the abortion of tendrils, certain cultivated varieties of Concribits appea have, according to Naudin.\* either quite lost these organs or bear semi-monatrous representatives of them. In my limited experience, I have met with only one apparent instance of their natural suppression, naudin, in the common bean. All the other species of Férin, I believe, bear tendrils; but the bean is stiff empty, in the common bean. All the other species of Férin, and the suppression of the common bean and the species of Férin, and the other species of Férin, and the suppression of the tendril-bearing plants similar realizations are suppression of the tendril-bearing plants similar realizations.

\* Annales des Sc. Nat. 4th series, Bot. tom. vi. 1856, p. 31.

these filaments are variable in shape, as is so frequently the case with radiamentary organs; they are either epidnrical, or followous, or and epoley furnowed on the upper surface. They have not retained any vestige of the power of revolving. It is a curious fact, that many of these filaments, when followcous, have on their lower surfaces, dark-coloured glands like those on the stipules, which exercte a sweet fluid; so

One other analogous case, though hypothetical, is worth giving. Nearly all the species of Lathyrus one with surprise who has noticed them, for they are quite unlike those of all common papilionaceous developed (for it is unbranched, and has no sponof L. nissolia, and its curious leaves are at once rendered intelligible to us.

It may be added, as serving to sum up the foregoing views on the origin of tendril-bearing plants, that *L* nissolia is probably descended from a plant which wa tendrils, with the stipules greatly increased in size through the law of compensation." After a time the tendrils lost their branches and became simple; they would have resembled the tendrils of the existing L. aphaca), and afterwards losing their prehensile power and becoming foliaceous would no longer be thus designated. In this last stage (that of the existing L. nissolia) the former tendrils would reassume would decrease in size. If species become modified in the course of ages, as almost all naturalists now admit. we may conclude that L. nissolia has passed through a series of changes, in some degree like those here

The most interesting point in the natural history of climbing plants is the various kinds of movement which they display in manifest relation to their wants. The most different organs—stems, branches, flowerpeduneles, petioles, mid-ribs of the leaf and leaflets, and apparently aérial roots—all possess this power.

The first action of a tendril is to place itself in a proper position. For instance, the tendril of Cobsea

<sup>\*</sup> Moquin-Tandon (Éléments de Tératologie, 1841, p. 156) gives the case of a monstrous bean, in

s for the leaves completely of a appeared and the stipules grew

first rises vertically up, with its branches divergent and with the terminal hooks turned outwards; the young shoot at the extremity of the stem is at the same time bent to one side, so as to be out of the way. The young leaves of Clematis, on the other hand, prepare for action by temporarily curving themselves downwards, so at o serve as grapuels.

Secondly, if a twining plant or a tendril gets by any accident into an inclined position, it soon bends upwards, though secluded from the light. The guiding stimulus no doubt is the attraction of gravity, as Andrew Knight showed to be the case with germinatin an inclined position in a glass of water in the dark, the extremity will, in a few hours, bend upwards; and if the position of the shoot be then reversed, the downward-bent shoot reverses its curvature; but if the stolon of a strawberry, which has no tendency to in the direction of, instead of in opposition to, the force of gravity. As with the strawberry, so it is generally with the twining shoots of the Hibbertia dentata, which climbs laterally from bush to bush; for show little and sometimes no tendency to curve up-

Thirdly, climbing plants, like other plants, bend towards the light by a movement closely analogous to the incurvation which causes them to revolve, so that their revolving movement is often accelerated or retarded in travelling to or from the light. On the other hand, in a few instances tendrils bend towards the dark.

ment which is independent of any outward stimulus, but is contingent on the youth of the part, and on vigorous health; and this again of course depends on a proper temperature and other favourable conditions of life.

may be, and the petioles or tips of the leaves of leafclimbers, and apparently certain roots, all have bepower of movement when touched, and bend quickly towards the touched side. Extremely slight present often suffices. If the pressure be not permanent, they part in question strightness itself and is again ready to bend on being touched. Sixthly, and leastly, tendrist, soon after clasping a

Sixthly, and lastly, tendrils, soon after clasping a support, but not after a mere temporary curvature, contract spirally. If they have not come into contact with any object, they ultimately contract spirally, after ceasing to revolve; but in this case the movement is useless, and occurs only after a considerable lapse of time.

With respect to the means by which these various movements are effected, there can be little doubt from the researches of Sachs and H. de Vries, that they are due to unequal growth; but from the reasons already assigned, I cannot believe that this explanation applies that the state of the sach property of the sach property of the property of the sach from a delicitate tune.

Finally, climbing plants are sufficiently numerous to form a conspicuous feature in the vegetable kingdom, more especially in tropical forests. America, which so abounds with arboreal animals, as Mr. Bates remarks, likewise abounds according to Mohl and Palm with climbing plants; and of the tendril-bearing plants examined by me, the highest developed kinds are species of Bignonia, Eccremocarpus, Cobwa, and Ampelopsis. But even in the thickets of our temperate regions the number of climbing species and individuals is considerable, as will be found by counting them. They belong to many and widely different orders. To series, I marked, from the lists given by Mohl and Palm (adding a few myself, and a competent botanist, no doubt, could have added many more), all those families in Lindley's 'Vegetable Kingdom' which include twiners, leaf-climbers, or tendril-bearers. Lindley divides Phanerogamic plants into fifty-nine Alliances; of these, no less than thirty-five include climbing plants of the above kinds, hook and root-climbers being exthough undeveloped, in almost every plant in the vegetable kingdom.

It has often been vaguely asserted that plants are of movement. It should rather be said that plants acquire and display this power only when it is of some advantage to them; this being of comparatively rare occurrence, as they are affixed to the ground, and food is brought to them by the air and rain. We see how high in the scale of organization a plant may rise, when we look at one of the more perfect tendrilbearers. It first places its tendrils ready for action, as a polypus places its tentacula. If the tendril be displaced, it is acted on by the force of gravity and rights itself. It is acted on by the light, and bends towards or from it, or disregards it, whichever may be most advantageous. During several days the tendrils or internodes, or both, spontaneously revolve with a steady motion. The tendril strikes some object, and quickly curls round and firmly grasps it. In the course of some hours it contracts into a spire, dragging up the stem, and forming an excellent spring. All movements now cease. By growth the tissues soon has done its work, and has done it in an admirable

### INDE:

Brassica napus, spontaneous movemen of peduncies, 198 Bryonia diobas, 131, 136 Caoutchous secreted by roots of Figureores, 186

power of twining in South Africa, 42 Cissus discolor, 143 Clematis, various species of, leafclimbers, 46 Colora scandens, 106

Combretum, 41
Corydalis claviculata, 121
Corydalis claviculata, 121
Cucurbitascov, nature of teadrils, 127
Cucurbita pego, aborted tendrils, 200
Cuccuta, stems ed, irritable, 17, 71
Dicentra thalictrifolia, 124

Discs, adhesive, developed by tendrils 94, 100, 135, 136, 145, 179 Dutrechet, reference to papers or elimbing plants, 1

Echinocyatis lebata, 128
Ferns, twining, 38
Ficus repeas, a root-climber, 185
Flagellaria Indies, 79
Flavorandusche, of Mayrandia, o

Flagellaria Indica, 79
Flower-polancies of Maurandia s tive, and revolve spontaneously, Fumaria officinalis, 75
Gallium appring a back-alludes 1

Galium aparine, a hook-climber, Gradatious of structure leading t development of perfect tendrils Glorican Plantii, 78
Halburya Mastiana, 134
Harvey, Pref., on the bess of power of
twining, 42
Hedsar helix, 185, 188
Hilbertia dentaia, 35
— shoots of, turn downwards,

petioles, 197
Hook-ellimbers, 183
Hop, powers of twining, 2
Hoya carneon, 6, 43, 185
Humulus lupulus, 2
Lelia multiss repeated by mode of Firms

India-rubber secreted by roots of Ficus repens, 186
Ipomea argyruoides, 42
Ivy, 185, 188
Jacquer, Prof. G., on climbing plants,

Jacger, Prof. G., on climbing plants
Kenner, on the irritability of flowerpeduncles, 197
Lathyrus aphoes, 115
probable manner of de-

velopment of its tendrils, 201
—, grassliflerus, 116
— nissolis, grass-like leaves replacing
tendrils, 201
Leaves, position of, on twining plants, 19
Leaf-followers, 45.

summary en, 81
climb more securely that
twiners, 192
Léon, M., on a variety of Phaseolus, 4

Léon, M., on a variety of Phaseolus, of the drils, 166
Light, action on twining plants, 40

..., avoidance of, by tendrifs, ve. 10c. 110, 138, 145, 175

Linam, spentaneous movement of, 19
Louis aurantiacu, 34
Lophooperaum scandens, 71
Lygodium articulatum, 38

M.Sab. De on Amountal Vestebii 14

M.Sab. De on Amountal Vestebii 14

Masters, Dr. M., on the woody vessels peticles, 75 Maurandia, a leaf-climber, 66 Mikania scandars, 33

Maurandia, a leaf-elimber, 66 Mikania scandens, 33 Mohl, Hugo, reference to work of, 1 Moyulo-Tandou, on the abortion of t

Muller, Fritz, on the structure of the wood of climbing plants, 44 — on plants scrambling over other plants, 183

other plants, 183

other plants, 183

on the development of branches into tendrils, 84

con roots of Philledendron

188 — on reots of Philedendron
188 — on the spontaneous movements of certain plants, 198
Mutisia elematis, 116

Nuties clemats, 119

Naudin on abertion of tendrils, 200

Nopenthus, 80

Nutation, revolving, 11

Ophicgiosum Japonicum, 77

Palm, reference to work of, 1

Passifiora accrifolis, 154

— gracilis, 153

— gracilis, 153
— punctata, 156
— quadrangularis, 157
Paullinis, 158
Pea, oramon, 112
Pedundes of Maurandia sensitive and

Peduncies of Maurandia sensitive and revolve spentaneously, 67 Phasedus, torsion of axes, 9 —, non-twining variety, 42 Philodendren, roots of, 188 Pisum sativum, 112

Rhedochiton volubile, 70
Roots acting like tendrils, 188
Roots-dimbers, 185
Rosa setigera, aboots bend from thight, 184

supports of different thickness, 17
on cause of movement of t
drils when touched, 180

when touched, 180 mess of tendrils, nature of, 19

Serjania, 152
Smilax aspera, 118, 184
Spencer, Herbert, on the relation of ax
and foliar organs, 199
Spiller, Mr., on the exidation of ind
rubber, 187

Spiller, Mr., on the exidatic rubber, 187 Spruce, Mr., on Maregravia, Solanum dulcamara, 34, 43 — jamintéles, 72

r Spiral contraction of tendrils, 158
Summary on twining plants, 39
of Sammary on leaf-climbers, 81
Summary on the movements of tendril
169, 202
[1

Summary on the movements of tendrile
169, 202
Summait of twining plants, often beckel
Support, thickness of, reund which
plants can twins, 22, 36
— thickness of which can be seen

—, thickness of, which can be embraced by tendrils, 176
Tassonia maxicata, 158
Tamus dephasatipes, 41
Tecoma radieans, 43, 185
Tendrils, history of our knowledge of, 81
—, spiral contraction of, 158
—, summary ca. 199, 202
— 1186

Tendril-bearers climb more securely
than twiners, 192
Tendrils, abortion of, 200
Torsion of the axes of twining plants, 7
Tropselum, various species of, leafclimbers, 60

Twining plants, 2

, shoots of, cometimes spenta
neomity become spiral, 17

table of rates of revolution
of various species, 24

on spiral contraction of ter drils, 160, 165 cause of movements of ter drils, 180 Vitis visifere, 137

## MR. MURRAY'S

# GENERAL LIST OF WORKS

ALBERT (Tms) MEMORIAL. A Descriptive and Illustrated Associate (the National Manuscent cresist to the PRINCE CONSORT at Kennington, (George, Status, Monaco, Mathreet, &c. With Descriptive Text, By Dovrse C. Beat., With 24 Pairs. Folio. 221. (2) (Prince) SPEECHERS AND ADDRESSES with an In-

ABBOTT'S (Rev. J.) Memoirs of a Church of England Missionary in the North American Coloules. Post Svo. 2s.

ACLAND'S (Rev. Charams) Popular Account of the Manners and Castros of India. Post Svo. 2s. #SOPS FABLES. A New Version. With Historical Preface.

ESOPS FABLES. A New version. With Historical Fremops Rev. Trousas Jusses. With 100 Weedcaits, by Tassers, and Woréfat Transact. Post Sive. 2s. 63.

AGRICULTURAIs (Grant) JOURNAL. (Published half yearly, AIDS TO FAITH: a Series of Theological Essays. Syo. 9s.

Tracles ONTESTS:

DEAN MANGEL.
BESSION FITOMERALD.

replacing of Anosis Record of Creation
Dr. McCalle.

Gance Cook.

Cance Cook.

Cance RAWLESTON.

supiration
Bession Fitomeral
Americans American
Americans American
Americans Fitomeral
Americans Fitomeral
Americans Fitomeral

AMBER-WITCH (Tas). A most interesting Trial for Witchcaft. Translated by Laur Deve Gonnous. Post Sys. 2s. ARMY LIST (Tun). Published Monthly by Authority. ARTHURS (LITTEN History of England. By Laur Calleott.

BYO. 18.62.
AUSTIN'S (JOHS) LECTURES ON GENERAL JURISPHUDENCE; or, the Followork of Positive Law. Ich Edition. Edited by Roman Camp-

Student's Edition, compiled from the above work. By
ROBERT CARPOLLS. Peat Svo.
ARNOLD'S (Twos.) Ecclesiastical and Sceniar Architecture of

Darwin Online: By permission of the Trustees of the Natural History Museum

# ADMIRALTY PUBLICATIONS; Issued by direction of the Lor

- Commissioners of the Admiralty:—
  A MANUAL OF SCIENTIFIC ENQUIRY, for the Use of Travelle
  A MANUAL OF SCIENTIFIC ENQUIRY, for the Use of Travelle
- Addison. Woodcuts. Post Ste. Sr. 6d.

  GREENWICH ASTRONOMICAL OBSERVATIONS 1841 to 184
  and 1847 to 1871. Royal 6to. 20s. each.
  - to 1847. Royal sto. 20. cock.

    APPENDICES TO OBSERVATIONS.
    - APPENDICES TO OBSERVATIONS.
      1837. Legarithms of Sines and Cosines in Time. 3s.
      1842. Catalogue of 1630 Starr, from Observations made in 1830 t
      - 1841. 4a. 1845 Longitude of Valentia (Chronocontrical). Sa.
        1845, Longitude of Altanirotth, Sa.
      - Twolve Years' Catabagus at Stars, from Observations as in 1934 to 1847. 4. Description of Photographic Apparatus. 2s.
        - 1852, I. Description of the Trausit Circle. Sa. 1851, Refraction Tables, Sa.
      - 1854. Description of the Zerdth Tube. Se.
        Bix Years' Catalogue of Stars, from Observations. 1848 to
        1851. 4c.
        1862. Serve Years' Catalogue of Stars, from Observations. 1864 to
        - 1802. Boven Yearn' Catalogue of Stars, from Observations. 180 1803. 10s. Plan of Ground Rolldings. 3s.
          - Longitude of Valentia (Galvanie), 2s. 164, Mecci's Bernid, from Occalvations, 2s.
      - Planetary Observations, 1811 to 1835. Ec. 1808. Corrections of Elements of Jupiter and Saturn. Sc. Second Soven Years' Catalogue of 2700 Stars for 1801 to
      - 1867. 4s.
        Description of the Great Equatorial. 3s.
        1866. Descriptive Chronograph. 3s.
        Observations 3s.
      - 1800. Reduction of Decop Thorncometer Observations. 24, 1811. History and Decoription of Water Telescope. 38, Canada Good Mana Observations (Star Ledgers). 1855 to 1801
      - 1856, 5a.

        Autronomical Results, 1857 to 1858, 5a.

        Respect on Teneriffe Astronomical Experiment, 1858, 5a.
  - Paramatta Catalogue of 7355 Sizes. 1852 to 1808. 4s.
    ASTRONOMICAL RESULTS. 1917 to 1871. 4cs. Sa. each.
    MAGNETICAL AND METEOROLOGICAL RESULTS. 1847
- - to 1800. 2 Vols. Royal 4to. 10s. each.
    1801 to 1801. 4to. 10s. each.
    1802 OF THE SERVICENTENARY TABLE, 1779. 4to. 5s.
  - BESSEL'S AUXILIARY TABLES FOR HIS METHOD OF CLEA-ING LUNAR DISTANCES. 5vo. 2a. CNORE'S BERLINER JAHRBUCH, for 1830. Beels, 1828. 8vo.
- HANSEN'S TABLES DE LA LUNE. 400. 200. LAY'S TABLES FOR FINDING THE LATITUDE AND LONGITUDE. 1821. 870. 100.

- - - SUPPLEMENTS, 1808 to 1833, 1807 and 1839.

  - TAYLORS SEXAGESIMAL TABLE, 1780, 4to, 15s.

  - WALES AND BAYLY'S ASTRONOMICAL OBSERVATIONS, 1777. 400. 21A.
  - MADE IN THE SOUTHERN HANDERSHARE, 1764-1771, 1788, 40). BARBAULD'S (Mas.) Hymns in Prose for Children, With 112

		Naturalist old Mines of and Observa					
BERTR.	M'S (Jas ood Fisher,	G.) Harve	sat of	the Sea :	an Acce	ount of her Felk	British . Thire
BIBLE	COMME	NTARY.	Exp	LANATORY	and Ca	RITICAL.	With

a REVISION of the TRADUCTION OF THE CO. COT., M.A., CARRO OF EVEL ANGLICAN CHILDREN. Edited by F. C. COT., M.A., CARRO OF EVEL Medium Str., Vot., L. 38s. Vota. H. and H. 14., 39s. Vot. I. V., 24s.

GENERAL COT. Beneral Cool: Rev. Sam. Clar.

Lavi Titles. Rev. Sam. Clar.

Vol. L. Lavi Titles.

Vol. I.	NUMBERS	Thrupp.
	KINGS, CHRONICLES, EX- RA, NEHENIAE, ESTHER	Bishop of Bath and Wells, Canon Bawlinson.

| Vol. 1V. | Proceedings | Pro

Yel, V. Jusemin Dean of Cambridge,
BICKMORE'S (A. S.) Travels in the Eastern Archipelage,
1820d; a Popular Description of the Islands, with their Natural Ristory, Correction, Proceedings of the Islands, with their Natural Ristory, Corrections, Svo. 21s.

PROCESS (S. West) History of Ancient Pottery and Porcelain:

Egyptini, Asyvan, Groek, Johnson, Medium Svo. 42r.
With Coloured Plates and 200 Hillostrations. Medium Svo. 42r.
BIRD'S (Isansula) Hawaiian Archipelago; or Six Months Among
the Palm Groves, Ceral Reefs, and Volcances of the Sandwich Islands.
With Illustrations. Crown Svo.

BISSETS (ANDREW) History of the Commonwealth of England, free the Death of Charles I, to the Expulsion of the Leep Parliament by Crouwell, Chiefer from the MSS, in the State Paper Office, 2 vols free, 30a. BUJUNTS (Rev. J. J.) Undesigned Coincidences in the Writings of

the Old and New Testament, an Argument of their Versity: containing the Books of Mosse, Historical and Prophetical Scriptures, and the Gappies and Acts. Elected Sciinies. Peat Swe. 5s.

History of the Courch in the First Three Centuries.

History of the Church in the First Auree Commisses

Fight Edition, Post Syo. 6s.

Parish Priest; His Duties, Acquirements and Obligations. Sixth Edition. Post Syo. 6s.

Lectures on the Right Use of the Early Fathers.

Third Edition. Swo. 9s.

University Sermons. Second Edition. Post Svo. 6s.

Pain Sermons, Sected Stateson, the Control of Pain Sermons, Stath Edition, 2 vols. Pest Svo. 128
BLOMFIELD'S (Bisnos) Memoir, with Selections from his Correspondence By the Seon, Second Editions, Fertual, past Svo. 128.
BOSWELL'S (JAMES) Life of Samuel Johnson, LLD. Including

BRACE'S (C. Ls) Manual of Ethnology; or the Races of the Old Weetl, Post Sec. 66. BOOK OF COMMON PRAYER. Illustrated with Coloured Borders, Initial Letters, and Weedents. Sec. 189.

Borders, Initial Letters, and Weedcute. Sec. 18s.
BORROWS (Gronnes) Bible in Spain; or the Journeys, Adventure
and Imprisonments of an Excellations in an Attempt to directain to
Seriginary in the Prenissels. Post Sec. 5s.

Cassona, Rodgico, and Lauguage. With Postrait. Post Svo. 5c.

Lavengro; The Scholar—The Gypay—and the Price
Post Svo. 5c.
Romany Rye—a Sequel to "Lavengro." Post Svo. 5

Romany Rye—a Sequel to "Lavengro." Post Svo. 5s.

WILD WALES: its People, Language, and Scenery.
Post Svo. 5s.
Romano Lave-Lil; Word-Book of the Romany. or

nonfano Lavo-Lit; word-Book on the Romany, or English Gryey Language; with Specimens of their Pestry, and an accessit of certain Gypeyries. Post two, 1st. 65. BRAY'S (Mas), Life of Thomas Stothard, R.A. With Portrait and 60 Woodcast, 4to, 2ts. Evolt of the Protestants in the Ceyennes. With some

Accounted the Hagments in the Seventeenth Century, Peat Sto. 10s.
BRITISH ASSOCIATION REPORTS, Sto.

SH ASSOCIATION SEPOCITYS See

The see of cortect states of the see of the see

1005, 1505, 170.
1007, 1508, 170.
1007, 1508, 170.
1007, 1508, 170.
1007, 1508, 170.
1007, 1508, 1508, 1508, 1508, 1509,

BROUGHTON'S (Loza) Journey through Albania, Turkey its Europe and Asis, to Constantingle, Hintertaine, 2 Vels wow. 319. Visits to Italy. 2 Vols. Post 870. 188. BROWNLOW'S (Lary) Reminiscences of a Septuagenarian From the year 1992 to 1815. Taked Edition. Past 870. 75. 66. BRUGSCH'S (Phorassee) History of Ancient Egypt. Derivee.

DANNY SERVICES, Syes.

BURGON'S (Rav. J. W.) Christian Gentleman; or, Memoir
Patick Fraser Tyder. Second Edition. Post Syes. 2s.

Letters from Rome. Post Syo. 12s.

BURNS. (Con. D.). William Technic

DURS S (COL.) Dictionary of Naval and Military Technical Terms, English and Fresch-French and English. Fowth Edition Crews Svo. 15s.

BURROW'S (Mošracy) Constitutional Progress. A Series Lectures destreed before the University of Oxford. 2nd Edition

CALLCOTT'S (LADT) Little Arthur's History of England,

CARNARVON'S (Loap) Portugal, Gallicia, and the Basque

Recollections of the Druses of Lebanon, With

Darwin Online: By permission of the Trustees of the Natural History Museum

CAMPBELL'S (Loss) Lord Chancellors and Keepers of the Great Seal of England. From the Earliest Times to the Death of Lord Editor is 1888. Ann Entites. 10 Yest. Committee. G. coch. Chief Justices of England. From the Norman

Chief Justices of England, From the Nor-Compass to the Death of Lord Testories. Third Ediston. 4 Crewn 200, 6s, each.
Lords Lyndhurst and Brougham, 8vo. 16s.

Lord Bacon, Fcap. Svo. 2s. 6d.

Lord Bacon, Fcap. Svo. 2s. 6d.

Nanoleon at Fentaineld

(SIR NEIL) Account of Napoleon at Fontaineble and Eibs. Being a decreal of Occurrences and Neces of his Consulters, do. Pectrait. Svs. 15r.

(SIR GEORGE) India as it may be: an Outline of a proposed Government and Pelicy. Svo.
(Times.) Essay on English Poetry. With Short

CATHCARTS (Sin Ground) Commentaries on the War in Russia and Germany, 1812-18. Plans. Svo. 14s.

CAVALCASELLE AND CROWES History of Painting Is lialy, from the field to the 18th Century. With Blustrations. 5 Vol. Bro. 21s. cath.

Early Flemish Painters, their Lives and Works, Binstrations, Petitive, 10s. 64.; or Large Paper, 8vo. 15s.

Children; being Bustraties of the Pawer, Emchaceaes, and Daskyn naminated by the Creater in his works. 10th Thesaudt. Post See. 6s. CHISHOLM'S (Mys.) Perils of the Polar Soas; Tros Stories of

CHURTON'S (Anompageon) Gongorn. An Historical Essay on the Age of Philip III, and IV. of Spain. With Translations. Portrait. 2 Vols. Small Sec. 12s.

New Testament. Edited with a Plain Practical Commentary for the use of Families and General Readers. With 100 Panoramies and other there, from Sketches and Photographs made on

the spot. 2 vols. Sec. 21s.

ICERO'S LIFE AND TIMES. His Character as a Statesman
Orstee, and Friend, with a Selection from his Correspondence and Ors.

CLARK'S (Sin James) Memoir of Dr. John Conolly. Comprising a Stocks of the Treatment of the Image in Europe and America. Wit Pertrait. Pess Son. 10s. 86.

CLIVE'S (LORD) Life. By Rev. G. R. GLEIG. Post 8vo. 3s. 6d. CLODE'S (C. M.) Military Perces of the Crown; their Administra

Administration of Justice under Military and Martial
Law, as applicable to the Army, Navy, Martin, and Auxiliary Perces
2nd Marins, Syo. 12s.
COLUME STREET MARKET DESCRIPTION OF THE PROPERTY AND ADMINISTRATION OF THE PROPERTY OF THE PROPE

COLUHESTER (Tun) PAPERS. The Diary and Correspondence of Charles Abbott, Lord Colchester, Speaker of the House of Commons 1802-1817. Pottmit. 3 Yola. Sws. 42s.

### LIST OF WORKS

CHURCH (Tun) & THE AGE. Essays on the Frinciples
Present Position of the Augitean Cauren, 2 vols, Srs. 25s. Center
Vot. I.
Augitean Principles.—Dean Hook.
Church and Presperism.—Earl N.

Anglien Principles.—Dean Mook.
Molera Religious Thought.—Blakep
Elliott.
State, Church, and Synoda.—Rev. Dr.
Irens.
Religieus Use of Tasta.—Rev. E. St.
John Tyvehött.
Place of Hollatty.—Profusce Berrows
Parish Priest.—Rev. Walshim How.
Divines of 16th and 11th Centeries.

John Tyrwhitt.
Plane of the Lity.—Professor Berrows
Partith Priest.—Rev. Waishum Hov.
Divines of 16th and 17th Centeries.
—Rev. A. W. Hafdan.
Liturgies and Ritosi, Rev. M. F.
Sadier.
Claren & Edocalien.—Caron Barry.
Irdian Musicos.—Sir Bartis Prevo.

Sadier, Church & Education.—Canon Indian Missicus.—Sir Bartis Church and the People.—Rev Maclagas. Carolilation and Comprehe Rev. Dr. Weir.

Conciliation and Compre Rev. Dr. Weir. COLERIDGE'S (SAMUS

COLERIDGE'S (SAMUEL TAYLOR) Table-Talk, Portrait, 12 COLLINGWOOD'S (CUTRIERRY) Bambles of a Natural

Shores and Waters of the Chins, &c. With Blustrations. Sec. COLONIAL LIBRARY. [See Home and Colonial Library.]

COOK'S (Canon) Sermons Preached at Lincoln's Inn. Svo. Se.

COOKERY (MODERN DORESTIC). Formula did Principal Constanting and Practical Knowledge, and adapted for Private Families. By a Lady. Woodcats. Frap. 8v. 5s.

COOPERS (T. T.) Travels of a Pioneer of Commerce on an

Overland Jorroy from Chias towards Itolia. Illustrations. Sec. 11s.

CORNWALLIS (Tim) Papers and Correspondence during the
American War.—Administration in Italia,—Union with Ireland, and
Proceed America. Second Editors. S. Vids. Sec. 62s.

Pence of Ambers, Second Edition. S Vots. Sec. 65s.

COWPERS (Courtess) Diary while Lady of the Bedchamber to Carolize Princess of Waler, 1714-10. Edited by Hou. Symmetry Market Sec. 10s.

CRABBE'S (Ray. George) Life and Poetical Works. With Illustration. Reyal Syn. 7s.
CROKER'S (J. W.) Progressive Geography for Children.

Stories for Children, Selected from the History of England. Fishers Woodents. 10mc. 2s. 6d.

Beavel's Life of Johnson. Including the Tear to the lightless. New and regimed Library Edition. Portraits, 4 vols. Sec.

Essays on the Early Period of the French Revolution.

Svo. 16e.

Historical Essay on the Guillotine. Fcap. Svc. 1s.

DELEPIERRE'S, (OCTAYE) History of Flemish Literature. See:

Historic Difficulties & Contested Events. Post Sya.

DENISON'S (E. B.) Life of Bishop Lonsdale. With Selectio

from Ma Writings. With Potrzsit. Crown Sya. 50c. 64.

from his Writings. With Portrait. Crown Sec. Soc. Sci.
DERBY'S (RASL ov) Illiad of Homer rendered into Enginess 2 Vots. Post Sec. Soc.
DEROS'S (Loxo) Young Officer's Companion; or, Essays

DE ROS'S (Loxp) Young Officer's Companion; or, Essays on Mininty Duties and Qualities; with Examples and Illustrations from History, Post 8vs. 9s. DEUTSCH'S (EMANUEL) Talmud, Islam, The Targums and other

DEUTSCH'S (EMANUEL) Talmud, Islam, The Targums and other Literary Remains. Sec. 12s. DOG-BREAKING; the Meet Expeditions, Certain, and Easy

DOMESTIC MODERN COOKBRY, Founded on Principles of Rensery and Practical Knowledge, and adapted for Princip Fault Woodnite, Fag, Sro. 58.

DOUGLASS (Sim Howard) Life and Adventures, Portrait, 8vo. 15s.

Theory and Practice of Gunnery. Plates. 8vo. 21s.

Construction of Bridges and the Passage of Rivers,
in Military Operation. Fistes. 8vs. 21s.

(Wa.) Horze-Shoeing; & it Is, and As it Should be.
Birstration. Post 8vo. 7s. 6s.

DRAKE'S (Str Frances) Life, Voyages, and Exploits, by Sea and Land. By John Barrow. Third Million. Post Sec. 2s. DRINKWATER'S (John) History of the Siege of Gibraltar,

Earliest Periods. Post 8vo. 26.

DUCANGE'S MEDIEVAL LATIN-ENGLISH DICTIONARY.

Translated by Roy. E. A. Daymen, M.A. Small 4to. [Ja preparation.

DU CHAILLUS (PAUL B.) EQUATORIAL AFRICA, with Accounts of the Gorilla, the New-building Aps, Chimpansse, Crocodile, 46. Illustrations. Sye, 21s.

tration into Equatorial Africa. Historiums. 8vs. 24s.

DUFFERIN'S (Loxo) Letters from High Latituden; an Account
of a Yacht Voyage to Iosland, Jan Mayen, and Spitzhergon. Fifth
Edition. Weedersta. Post 8vs. 7s. 6s.

DUNCAN'S (Majon) History of the Royal Artillery. Compiled from the Original Records. Second Edition. With Portraits. 2 Votes Sec. Dis.

DYER'S (Tucs. H.) History of Modern Europe, from the taking of Constantinople by the Turks to the close of the War in the Grimes. With Index, 4 Vols. 8vc. 42s.

EASTLARE'S (SIR CHARLES) Contributions to the Laterature of the Fine Aris. With Memoir of the Author, and Selections from his Correspondence. By Lavy Eastlares. 2 Yes. Swo. 24s. ELDON'S (Lone) Public and Private Life, with Selections from his Correspondence and Diaries. By Housen Twins. Third Edition. Protests, 2 Vois. Post Sec. 21s.

ELGIN'S (Loan) Letters and Journals. Edited by Tanopour

ELLESHERS (Loss) Two Sieges of Vienna by the Turks, Translated from the German. Post Svs. 2s. ELLISS (W.) Madsgascar, including a Journey to the Capital,

with actices of Natural History and the Poople. W.

Madagascar Revisited. Satting

Madagascar Revisited. Setting forth the Persecutions and Heate Sufferings of the Native Christians. Illustrations. Sec. 10s. Memoir. By His Sox. With his Character and

Werk. By Rev. HISBN ALLOS, D.D. Porrait, Svo. 10s. 56.

(ROBISSON) Poems and Fragments of Catollius. 16mo. 5c.

ELPHINSTONE'S (Hos. Mountstvar) History of India—the
Utrales and Mahamidan Pericks. Sizzl Edition, Man. Svo. 13s.

Hindos and Mahomedon Periods. Stath Militon. Map. Svo. 18s.

(H. W.) Patterns for Turning; Comprising
Elliptical and other Figures cat on the Lathe without the use of any
Organization. Small Line. My Not Disagraphics. Small Line. 1ds.

of the Assyrians, Egyptians, and Bebrews; with Special Reserves
the Discoveries in Western Asia and in Egypt. Scood Edition. W
100 Illustrations. Proc. 100, 65.
ENGLAND. See Calacott, Croker, Hume, Markham, Smr.

end Stanmor.

ENGLISHWOMAN IN AMERICA. Post Svo. 10s. &d.

ESSAYS ON CATHEDRALS, With an Introduction. I DRAW HOWSON, Svo. 12s. CONTENTS.

Incomprise of a Dean,—Inding of Carliolo, Carloon, Carloon, Carloon, Carloon, Carloon, Carloon, Carloon, Nerion and their Werg,— Carloon Nerion in Frederick, Fred and Fig. Carlooning in their Missierum; Anderson in thei

Cathedrals in their Missiscary Aspect.

—A. J. B. Serveford Heps.
Cathedral Feomatoises in Relation so
Religious Thought.—Cason Westcott.

Characteristic of the Chapter to the Risks
Characteristic of the Cathedral

Cathedral Reform.—Characteristic of Characteristic of the Cathedral

ETHNOLOGICAL SOCIETY'S TRANSACTIONS. Vols. I. to

ELZES (Karl) Life of Lord Byron. With a Critical Essay on his Fixes in Literature. Translated from the German, and Edited with Netss, With Original Petralt and Facsimillo. Svo. 16s.

FAMILY RECEIPT-BOOK. A Collection of a Thousand Valuable and Useful Receipts. Frag. Svo. 5s. 5s.

FARRARS (A. S.) Critical History of Free Thought in

reference to the Christian Religion. See. No.

(F. W.) Origin of Language, based on Modera

Essearches, Fey, See. 5s.

FERGUSSON'S (James) History of Architecture in all Countries

Vol. IV. Modern, With \$30 Illustrations.

FLEMING'S (PROFESSOR) Student's Manual of Moral Philosophy.

FORSYTH'S (WILLIAM) Life and Times of Cicero, With Selections

FRENCH (THE) in Algiers: The Soldier of the Foreign Legion-

FRERE'S (SIR BARTLE) Indian Missions. Third Edition. Eastern Africa as a field for Missionary Labour. With

Bengal Famine, How it will be Met and How to Prevent

- (M.) Old Decean Days; or Fairy Legends Current in

GEORGE'S (Engage) Mosel; a Series of Twenty Etchings, with Loire and South of France; a Series of Twenty GIBBON'S (EDWARD) History of the Decline and Fall of the
EDWARD ENGINE. Edited by Minara and GUIDT. A New Edition.
EDited, with Noise, by Dr. We. SERRE, Maps. 8 Vols. Sve. 60s.

(The Student's Gibbon); Being an Epitome of the

(The Student's Decembers of Score Communication. By

shave werk, interparating the Researches of Recount Commentators. By Dr. Wx. Surril, Woodcats. Post Svo. Tz. 6d.

GIFFARD'S (Edward) Doeds of Naval Daring; or, Anocdotes of the Rittle Navy. Francisco. Sz. 6d.

GLADSTONE'S (W. E.) Financial Statements of 1853, 1860, 63-65.

GLEIG'S (G. E.) Campaigns of the British Army at Washington

GLEIG'S (G. E.) Campangar 20.

and New Orleans. Post Svo. 2s.

Story of the Battle of Waterloo. Post Svo. 3s. &d.

Narrative of Sale's Brigade in Afighanistan. Post Svo. 2s.

Life of Lord Clive. Post 8vo. 3s. 6d.
Sir Thomas Munro. Post 8vo. 3s. 6d.

Sir Thomas Munro. Post 8vo. 3s, 6d.

GOLDSMYTH'S (OLIVER) Works. Edited with Notes by Peren
CORNIDORS, Vigrettes. 4 Vols. 8vo. 50s.

GORDON'S (Siz Alex.) Sketches of German Life, and Scenes.

from the War of Liberation. Post Sro. 3s. 6d.

(LADY DUFF) Amber-Witch: A Trial for Witchcraft. Post two, 2s.
French in Algiers. 1. The Soldier of the Foreign

Legion. 2. The Priessess of Abd-d-Kadir. Post Src. 2s.
RAMMARS. See Curtius; Hall; Hurrow; King Edward
Mayrows: Mayrows Surge.

GREECE. See GROTE-SHITH-Student GREY'S (EARL) Correspondence with

GRUNER'S (Lawis) Terra-Cotta Architecture of North Italy, from careful Deswings and Restorations. With Illustrations, cognared and printed in Colours. Email fello. 51, 5s.

QUILLOYES (M.) MCMINATORS ON CHRISTIANINY, MAN uses recognitions of the Day, Part J. The Essence, Part H. Present Scale,
Part III. Eviation to Society and Opinion, S Vols. Post Stro. Soc.
GROTE'S (Ground) History of Greece. From the Earliest Time:

to the describe generation contemporary with the death of Abstractor the General Library Edoine, Pourtral, Maps, and Plans. 10 Vals. Fee: 100c. Colling Edition, Pertuals and France, 12 Vols. Peat See, 6c. each.

Plano, and other Companions of Socrates, 3 Vols. Svc. 45c.

Americas. Edited by Professors Bars and Roberstrop.

2 Vols. Srs. 32r.

Mimor Works, With Critical Remarks on hiintellectual Charmeter, Writings, and Specoles. By ALEX.Bary, LL.D.

Interiorumi Character, Writingi, and Speeches. By ALEX.BARK, LL.D.
Dovusti. swo. 14s.

— Personal Life. Compiled from Family Documents,
Private Memoranta, and Original Latters to and from Various
Friends. By Mes. Green. Parintit. Swo. 17s.

HALL'S. (T. D.) School Manusl of English Gramms
Copiesa Exercises, 12ms, 3s, 64,
Primary English Grammar for Elementary

Child's First Latin Book, including a Systematic Truest of the New Procuretation, and a full Praxis of Nosos, Adult and Procure, Idon. 12 64.

Accession of Henry the Seventh to the Death of George the Secent Zelvery Edition, S Vols. See. Step. Collect Edition, S Vols. Post See, U. Student's Edition of the above work, Edited i Wil. Surms, D.C.L. Post See, 7s. 6d.

History of Europe during the Middle Ages. Librar Edition. 3 Vals. Sve. 30s. Galisse Edition, 3 Vals. Post Svo. 12s. Student's Edition of the above work. Edited b

Literary History of Europe, during the 15th, 16th and
17th Centurios. Library Enties. 3 Vots. 8vo. 36s. Califort Estimos.
4 Vots. Post 8vo. 16s.
(A wyuon Literary Remains; in Verse and Pross.

HANNAH'S (Rev. Ds.) Divine and Human Elements in Holy Serjours. Sec. 10s. 64. HART'S ARMY LIST. (Published Quarterly and Annually.) WAY'S, Gra. J. H. Danmaone) Western Barbary, its Wild Tribes

HEAD'S (Sin Francis) Royal Engineer. Hiustrations. Svo. 12s.

Life of Sir John Burgoyne. Post Svo. 1s.

Rapid Journeys across the Pampas. Post 8vo. 2s.
 Bubbles from the Brunnen of Nassau. Illustrations.
Post 8vo. 7s. 6d.

Kmigrant. Feap. 8vo. 2s. 6d.
Stokers and Pokers; or, the London and North Western
Railway. Post 8vo. 2s.

HEBER'S (Bisnor) Journals in India. 2 Vols. Post 8vo. 7e.

Peetical Works. Portrait. Fesp. 8vo. 3s. 6d.

Hymns adapted to the Church Service. 16mo. Is. 6d.
HERODOTUS. A New English Version. Edited, with Notes
and Essays, historical, etinographical, and peographical, by Caroor
Hawayson, assisted by Ess. Hown Hawaysons and Sin J. G. Win-

HATHERLEY'S (Lond) Continuity of Scripture, as Declare, by the Testinecy of our Lord and of the Evangelists and Apostes Shorth Edition. Sero. 6s. Popular Edition. For Ser. 2s. 6d. HOLLWAYS (J. G.) Month in Norway. Feap. Sero. 2s.

HOLLWAY'S (J. G.) Month in Norway. Fcap. 8vo. 2s. HONEY BEE. By REV. THOMAS JAMES. Fcap. 8vo. 1s. HOOK'S (DEAR) Church Dictionary. Testh Edition. 8vo. 1s. HOPE'S (T. C.) ARCHITECTURE OF AHMEDABAD, with - (A. J. BERESTORD) Worship in the Church of England.

HAND-BOOK-TRAVEL-TALK. English, French, German, and Italian, 18mo, 3s, 6d HOLLAND,-BELGIUM, and the Rhine to Mayence

NORTH GERMANY,-PRUSSIA, SAXONY, HAS-

SOUTH GERMANY,-Bavaria, Austria, Styria.

KNAPSACK GUIDE TO THE TYROL. 16mo, 6%

PAINTING, German, Flemish, and Dutch Schools. LIVES OF EARLY PLEMISH PAINTERS,

SWITZERLAND, Alps of Savoy, and Piedmont.

FRANCE, Normandy, Brittany, the French Alne.

ISLANDS OF THE MEDITERRANEAN-Malta. ALGERIA, Map. Post 8ro, 9e.

PARIS, and its Environs. Map. 16mo. 3s. 6d.

NORTH ITALY, Piedmont, Liguria, Venetia,

ROME AND ITS ENVIRONS. Map. Post 8vo. 10s.

RUSSIA, Sr. PETERSBURG, MOSCOW, POLAND, and

HAND-BOOK-SWEDEN. Map. Post 8vo. 6s.

GREECE, the Ionian Islands, Continental Gre
Athens, the Pelepomosus, the Islands of the Algem Son, Alla
Tassaidy, and Macolicali. Mags. Post free. 159.

TURKEY IN ASIA—CONTANTINAPER, the I
phorax Davisandis, Recuss, Pain of Toy, Crete, Cyrgus, Smy

Epheresa, the Sevens Churches, Consta of the Book Son, Armon Mesopolitable, Ac. Mays. Pets Son. 15s.

BOYPT, Including Descriptions of the Course the Mile through Expt and Multi, Abstantis, Cairo, and Thobas, it Sues. Course in the Course of the Course Special Course of the Course of State, the Ones, the Typon, dat. May. Pets Syn. 15s.

Physiol. 15s.

Sinai, Edwn, Syrian Deartt, &c. Mays. Fost Svc.
INDIA — BORBAT AND MADEAS. Map. 2 Vols.
Post Svo. 12s. csrb.
FNGLISH HANDBOOKS.

HAND-BOOK-MODERN LONDON, Map. 16mo. 3c. 6d.

ESSEX, CAMBRIDGE, SUFFOLK, AND NOR-FOLK, Chelmsford, Calabaster, Maldon, Cambridge, Ely, Normarket, Bury, Iporich, Woodhridge, Felixstows, Lowestoft, Norwich, Yamooth, Cromer, &c. May and Plans. Peat 8vo. 12s.

CATHEDRALS of Oxford, Peterborough, Norwick Ely, and Lincoln. With 90 Illustrations. Crown Svo. 18s. KENT AND SUSSEX, Canterbury, Dover, Ram

Weithing, Hastings, Lowes, Arusiel, &c. Map. Post 8vo. 20s.

SURREY AND HANTS, Kingston, Croydon, Relgate, Golifford, Derking, Boahill, Winelsester, Southampton, New
Freest, Fortensuch, and Jazz of Wiouri, Maps. Post 8vo. 10s.

ERKS, BUCKS, AND OXON, Windsor, Eton,

Reading, Aylashery, Uxbridge, Wycombe, Henley, the City and University of Oxfreet, Betchelm, and the Descent of the Themes. Map Post Str. 7, e42.

WILTS, DORSET, AND SOMERSET, Sallisburg, Chippenham, Weymouth, Sherborne, Wells, Bath, Belaist, Tannee,

Lauces, Scimouth, Dawlish, Teigumouth, Plymouth, Devaupert, Teeensy, Launesston, Turno, Pontance, Fairneuth, the Linsed, Lauc's End. de. Magn. Post Spc. 12s. CATHEDRALS of Winchester, Salisbury, Exeter

CATHEDRALS of winessets, Samsoury, Executively, Cathebrat, Rochester, Cantectury. With 10 Hustration 2 Vols. Crown Str. 24c.
GLOUCESTER, HEREFORD, AND WORCESTER CUrrecessor, Cachembran, Stroat, Testesbury, Leonington, Ross, Mai

vers, Astermaners, Juney, weeker, Bristol, Gloucester, Horeford, Cartherna L. S. Grand, Gloucester, Horeford, Wercoster, and Lichard. With 50 Illustrations. Crown Sys. 46s.

NORTH WALES, Bangor, Carnarvon, Beaumaris, Sacodro, Lianberts, Dulpoly, Cader Liris, Cooway, &c. Map. Post

Sec. 7a. SOUTH WALES, Monmouth, Llandaff, Merthyr, Vale of Neath, Penbroke, Carmariben, Tenby, Swansen, and The Wye, de. Map. Post Sec. 7a.

HAND-BOOK—CATHEDRALS OF DANGOR, St. ASAC Lieudelf, and St. Davids, With Riburstations, Port Src. Lis. DERBY, NOTTS, LEICESTER, STAFFOR Matteck, Balverell, Clastrowth, The Peak, Buxton, Hardwick, Dalos, Achburno, Senthwell, Mausfield, Ratford, Borron, Bolveit, Mc-Dalo, Achburno, Senthwell, Mausfield, Ratford, Borron, Bolveit, Mc-Walventampter, Laichdald, Washall, Tamworth. M

Matleck, Bakowski, Carterior, 1 ha Feeta, Burton, Boreste, McDale, Ashbora, Senthwell, Manfeld, Ratford, Burton, Boorste, McMowbray, Welvenhaupise, Lichibelt, Waball, Tamwetth. 3 Poet few. 50. PSHIRE, CHESHIRE AND LANCASHI—Shreebury, Indion, Bridgeoch, Oswestry, Chester, Green, Alles

Burnier, Cilithere, Bistro, Blackbarn, Wigne, Frestre, Bothi, Lammery, Sorthport, Blackpool, &c. Map. Post Svo. 19. Company, Control of Company, Co

CATHEDRALS of York, Ripon, Darham, Carlis Chester, and Manshester. With 60 Illustrations. 2 Yels. Crown by 21s.

DURHAM AND NORTHUMBERLAND, New York Control of the Control

control, idellington, unanomous, control Accutant, morceson, Harthopol, Smocketan, Shiebis, Interviton-Tweed, Morpeth, Tymenuchi, Colistream, Alterieu, ée, Map. Pout 800. 80. EXETMORLAND AND CUMBERLAND—Lancaster, Formous Abbey, Ambheidis, Kendal, Windermere, Cortico, Kenwick, Grammer, Liverito, Cacillale, Caccementa, Paculia, Apolsir,

Map. Pest Svs. 66.

\*\* MURRAY'S MAY OF THE LARK DIFFRICT, on canves. 3s. 6s.

SCOTLAND, Edinburgh, Melrose, Kelso, Glasgo
Dunfries, Arr., Striling, Arras, The Glyde, Ohan, Inversay, L.

Lenned, Look Katrice and Tressechs, Cablemian Canal, Inverses Porth, Dundes, Aberdeen, Brasmar, Skye, Caithness, Ross, Suiteland, éc. Mays and Plans. Post Svo. Sz. [RELAND, Dublin, Belfatt, Donegal, Galway Warfied Cork Limatrick Waterford, Killarney, Nature & Man

Wesferd, Cork, Lamerick, Waterford, Killarosy, Munster, &c. Maps.
Post Sec. 12c.
FAMILIAR QUOTATIONS. From English
Authors. Third Edition. Fear, Sec. 5a.
HORACE: a New Edition of the Fext, Edited by DRAN MILMAN.

With Ito Woodcuts. Crown Svo. 7s. 6d.

Life of. By DRAN MILMAN. Illustrations, Svo. 9s.

HOUGHTON'S (LORD) Monographs, Personal and Social, With
Petrales, Crown Svo. 10s. 6d.

HUME'S (The Student's) History of England, from the Invasion of Julius Cosar to the Revolution of 1688. Corrected and contioned to 1898. Woodcote, Post Svo. 7s. 5d. HUTCHINSON (Gray), on the most expeditions, certain, and

say Method of Deg-Breaking. Fight Edition. With 40 Bilantrations Cown Sec. Br. HUTTON'S (H. E.) Principia Gravea; an Introduction to the Study

of Greek, Comprehending Grammer, Delective, and Exercise-book with Vecabulariese, Strak Edition, 1930, Se fel.
IRBY AND MANGLES Travels in Egypt, Nubis, Syria, and the Hely Land. Post Svo. 2s.

JACOISSON'S (Basses) Fragmentary Illustrations of the History of the Book of Common Prayer; from Manascript Sources (Bishop Manascript Sources (Bishop Wiss), 8vo. 5s. JAMES (Sav. Tromas) Pables of Esop. A New Translation, with Historical Profess. With 100 Woodcots by Travens, and Work. CLASS A.

# HISTORY, BIOGRAPHY, AND HISTORIC

VOYAGES, TRAVELS, AND ADVENTURES

Darwin Online: By permission of the Trustees of the Natural History Museum

JAMESON'S (Mas.) Lives of the Early Italian Painters and the Progress of Painting in Italy—Cimahos to Bassane. New Estates. With 50 Partrales. Post Src. 17s. JENNINGS (L. J.) Egipty Years of Republican Government in

JENNINGS' (L. J.) Eighty Years of Kepublican Government the United States, Post Sco. 10s. 62. JERVIS'S (Ray. W. H.) Gallican Church, from the C coulat of Belgas, 1816, by the Revolution, With an Jatrodact

JESSES (Enward) Gleanings in Natural History, Fep. 8vo. 3s.
JOHNS' (Exv. B. G.) Blind People; their Works and Ways. W
Sketches of the Lives of some famous Blind Mes. With HinstratiPost Svo. 7s. 6d.

Sketches of the Lives of some famous Blind Men. With Hinstratic Post See, Ta. 6d.

JOHNSON'S (Da. Samura) Life. By James Boswell. Includi the Tour to the Helvides. Edited by Ma. Choras. New revised Life.

the Teur to the Hebrides. Edited by Na. Chonga. New revised Liberry
Edition. Pertraits. 4 Vols. 800.

Lives of the most eminent English Poets, with
Critical Observations on their Works. Edited with Notes, Corrective

JUNIUS' HANDWRITING Professionally investigated. By Mr. Chaddy Expert. With Prefice and Cobateral Evidence, by the Hon. Edwant Twilleron. With Packiniles, Woodests, &c. 4to. £13a.
KEN'S (Hamop) Life. By a Layman. Portrait. 2 Vols. 8vo. 18s

Expesition of the Apostles' Greed, 16mo, 1s. 6
KERR'S (Robert) GENTLEMAN'S HOUSE; oz. How to Pr
Estates Respected, prior the Practice of the Palace, 27
Estate, With Your and Plant, Sto. 24s.

Small Country House, A Brief Przetloil Discourse on the Paraing of a Residence from 2000, to 2000. With Supplementary Estimates to FOOD, Post Swo. 8.

Ancient Lights; a Book for Architects, Surveyors, Layrer, and Landerds. Swo. 2a. 55.

(R. Marconn) Student's Blackstone. A Systematic Abriguent of the entire Connectance, adapted to the present state of the law, Post Sys. Va. 64.

KING EDWARD VIrm's Latin Grammar. Seventeenth Edition. 18mo, 3r, 64. First Latin Book. Fifth Edition.

KING GEORGE IHRD'S CORRESPONDENCE WITH LORD NORTH, 1768-82. Edited, with Notes and Introduction, by W. Bootham Doxus, 2 vols. Syrp. 324.

KINGS (R. J.) Sketches and Studies; Historical and Descriptive, free, 12a. KIRKS (J. Feerma) History of Charles the Bold, Duke of Burguer, Petrast, Svon. Sec. 45s.

gusor, Pretrait. 2 Vols. 8vo. 45s.

KIRKES' Handbook of Physiology. Edited by W. Morrant
Barra, F.R.C.S. Ejská Ess. Wni 20 Bustratios. Pest Svo. 12s. 6s.

KUGLER'S Handbook of Painting.—The Italian Schools. Fourth

Sitius, Revised and Benodellief from the most treem Researches. By Later Ratrage. With 164 Binstmittees. 2 Vols. Crown Str. 56s.— Handbook of Painting.—The German, Flemish, and Datch Schools. Third Edition. Revised and in part re-written. By J. A. Cawar, With 60 Binstmittees. 2 Vols. Crown tree. 24s.

AND Schools, The Every Revised and in part re-settion, By J. A. Caswa, While Distractions. 2 Ved. Crown for 20th AND S. (E. W.). Account of the Manners and Customs of Modern Egyptions. Nov Barings, With Hastattanes, 2 Volb. Part Serv. 12t, AVRENOE'S (SER Gas.). Reminiscences of Forty-three Years' Service in trains; tobacking Conviviles in Calcul many give Arghaes.

rative of Researches and Discoveries amblet the Rottes of Assyris
With an Account of the Chaldean Christians of Kurdisan; the Youngel
or Devil-weedlepers; and an Requiry into the Manners and Arts o
the America Assyrians. Shad Edinos. Plates and Woodents. 2 Vels
200. One.

Sec. 38s.

\*\* A Provinan Eurrous of the above work, With Illustration
Pest Sec. 7s. 6d.

Nineveh and Babylon; being the Narrative of Di
caveries in the Rules, with Travels in Armenia, Kurlistan and it
caveries in the Rules, of Examiliate to Assertis. With Map as

Dreset, during a Second Expedition to Assyria. With Map at Plates. Sec. 21s. \* A Portical Environ of the above work. With Illustration Past Rec. 7a. 66.
Past Rec. 7a. 66.
ULES: New York.

LEATHES' (STANLEY) Practical Hebrew Grammar. With the Habrew Taxt of General i.e-Vi. and Poslem i.e-Vi. Grammatical Analysis and Vocabulary. Post Syn. 7s. 5d. LENNEP'S (REV. H. J. Vas) Missionary Travels in Asia Minor.

and Woodcotts. 2 Vols. Post Sec. 286.

LESLIES (C. R.) Handbook for Young Painters. With Illustrations. Post 8vo. 7s. 66.

Life and Works of Sir Jeshua Reynolds. Portrai's

and Illustrations, 2 Vols. Sec. 42s.
LETTERS FROM THE BALTIC. By a Lady. Post Svo. 2s.
Mannas. By a Lady. Post Sec. 2s.

SIERRA LEGGE By a LAST. Post Sto. 3s. 6d.

EVIS (LEGGE) History of British Commerce; and of the Economic Progress of the Nation from 1763 to 1870. Sec. 16s.

LEWIS'S (M. G.) Journal of a Residence among the Negroes in the West Indies. Post Sto. 2s. LIDDELL'S (Dwas) Student's History of Rome, from the earliest Times to the outshishment of the Empire. With Woodcaba. Post two.

7s. fet.
LINDSAY'S (Lord) Lives of the Lindsays; Memoir of the
Horsen of Crawford and Ralearres. With Extracts from Official Papers
Horsen of Crawford and Ralearres. With Extracts from Official Papers

and Personal Narratives. 3 Vols. Svo. 24s.

Etruscan Inscriptions. Analysed, Translated, and
Compensed upon. 8vo. 12s.

with kineditations of the Stellien files of Freedy, with any 500, vol.

LISPINGS from LOW LATTUDES; or, the Journal of the June 100 to 100 to

LIVINGSTONE'S (Dn.) Popular Account of Missionary traves and Researches in Seath Africa. Bustratices. Post 8vc. 6z. Narrative of an Expedition to the Zambezi and its Tributaries, with the Discovery of the Lakes Shirwa and Nyassa.

Map and Historiston Nov. 21s.

Last Journals in Central Africa, from 1865 to
his Death. Continued by a Narpolve of his last more us and sufferings. By Rev. Hoxace Walkins. Maps and Hustrations. 2 Vols.
Syn. 28s.

ings. By Rev. Horace Walkis. Napi and indicated the No. Sec. Livonian Talles. By the Author of "Letters from the Batis," Post Sec. 2s.

LOCH'S (H. B.) Personal Narrative of Events during Lord China, Second Edition, With Illustrations.

Darwin Online: By permission of the Trustees of the Natural History Museum

LOCKHARTS (J. G.) Ancient Spanish Ballads. I Espanish. Translated, with Notes. New Edition. W.

Life of Theodore Hook, Fcap. Svo. 1s.

LONSDALE'S (Bassor) Life. With Selections from his Writin

LOUDON'S (MEA) Gardening for Ladies, With Directions and Withings, LOUDON'S (MEA) Gardening for Ladies, With Directions and Calcular Operations for Every Meah, Eighth Efficien, Weed-

LUCKNOW: A Lady's Diary of the Siege. Feap. Sto. 4s. 6d.

LYELL'S (Sta Charles) Principles of Geology; or, the Modern

Charges of the Eart's and its inhabitants considered as illustrative of

Changes et the Earth and its intaltesint consorted as illustrative of Geology. Extracts Edition. With Internations. 2 Vol. 89s. 32s.

Student's Elements of Geology. Second Edition. With Table of British Featls and 90 illustrations. Foot 80s. 9s.

Geological Evidences of the Antiquity of Man,

on the Origin of Species. Fourth Edition. Hinstrations. Sec. 14s.

(K. M.) Geographical Handbook of Ferns. With Tables
to show their Dustibutors. Four two. 7s. 6d.

1981 JONES (Loan) Engagers. 2 Vols. Post 37s. 10s. 6d.

LYTTON'S (Loan) Memoir of Julian Fane. With Portrait. Po

McCLINTOCK'S (Srs L.) Narrative of the Discovery of the Pate of Sir John Franklin and his Companions in the Arelle Bess, Third Edwice. With Illustrations. Peat Str., 7s. 6s.

Artillery. With Plans. Post Src. 12s.

MACGREGOR'S (J.) Rob Roy on the Jordan, Nile, Red Sea, Ger
research, &c. A Cance Unite in Palestice and Egypt and the Wate

reservett, fid. A Caroo Croite in Palestine and Egypt and the Waters of Darasces. Cheeper Edinov. With Map and 70 Hinstrations. Cross Syo. 7, 61.

MACPHERSON'S (Maron) Services in India, while Political

Agent at Gwallice during the Mutiny. Illustrations. Sec. 19s.

[AETZNER'S ENGLISH GRAMMAR, A Methodical, Analytical
and Illustrical Treations on the Orthography, Proxedy, Infactions, are
Syntax of the English Tongue. Translated from the German, By
Const. Graver, Leit. S. Nyde. 80, 266.

MAHON (LORD), see STANDOFE.

MAINE'S (SIR H. SEMERR) Ancient Law: its Connection with the
Early History of Society, and its Relation to Medical Ideas. F/1

Estiles, 8vo. 11s.

Village Communities in the East and West, Second Selector, 8vo. 9s.

Early History of Institutions, 8vo. 12s.

MANSEL'S (DEAN) Limits of Religious Thought Examined.

Fight Entires, Peatieve, 8s. 6s.

Letters, Lectures, and Papers, including the Phrontistopic of Confession, Nath Control, Edited by M. W. Carriera

M.A. Stv. 12.

Gnostic Heresies of the First and Second Century.
With a starth of his life and character. By Leef Carranvow.
Educator Constitution of the Constitution of the Manual Constitution of SciEntly 1910 E NO. Use. 66.

MANUAL OF SCIENTIFIC ENQUELY. For the Use of

MARCO POLO, The Book of Ner Marco Poo, the venetan Concerning to Kingheau and March of the East. Anne English Version, Hustanes by the light of Oriental Writers and Moore Travels. By Oo. Huser Years, East Edition, Maya and Huster tiess, Z.Vels, Medium Sro. 42s.

MARCHAMMS (Must.) Wistors of Racional. From the First Inva

MARKHAM'S (Max.) History of England. From the First Inv
sion by the Romans to 1880; Weedens, 19me, 5s. 6d.

History of France. From the Conquest by the
Canha o 1801. Weedens. Theo., 4s. 6d.

Solid State of Stat

(CLEMENTS E.) Travels in Pers and India. Maps and Illustrations. Svc. 19s.
MARRYATS (JOSEPH) History of Modern and Mediaval Pottery

and Percelain. With a Description of the Manufacture. This Refines. Plates and Woodcasts. Seps. 628.

MARSH'S (G. P.) Student's Manual of the English Languag Pest'so, 7s, 6s.

MATTHLE'S GREEK GRAMMAR. Abridged by BLONFIEL

Ected by R. S. Crooms. Erro. 4c.
MARREL'S Character, Actions, and Writings of Wellington.
Fens, Svo. 1s. 6t.
MAYNES (CAPA.) From Years in British Columbia and Vanconver Loland. Illustrations. Svo. 16t.

MEADE'S (Hos. Hazaras) Ride through the Disturbed Districts of New Zealand, with a Craine among the Bouth Sea Islands. With Illustrations, Medium Seo. 12s. MELVILLES (HEMMANN) Marquesus and South Sea Islands.

2 Vats. Peat See. 7s.

MEREDITH'S (Mrs. CHARLES) Notes and Sketches of New South
Walsa. Post Seo. 2s.

MESSIAH (THE): The Life, Travels, Death, Resurrection, and
Ascretian of our Bessed Lord, By A Layman. Map. Seo. 18s.

Ascension of our Heased Lord. By A Layrana, Map, Not. 28.
MILLINGTON'S (Rav. T. S.) Signs and Wonders in the Land of
Ham, or the Ten Pingrees of Egyps, with Ancient and Modern Illustrations. Woodcuts. Food Sov. 7-6.
MILLS' (Rav. Jonn) Three Months' Residence at Nabius, with

MILMAN'S (Dara) History of the Jews, from the excises revious down to Modern Times. Fourth Edition, 2 Vols. Post@vo. 18s.

Barly Christianity, from the Birth of Christ to the Abelitte of Pagualam in the Reman Empire. Fourth Emiss. 2 Vols. Post@vo. 18s.

Fost Swo. 15s.

Latin Christianity, including that of the Popes to
the Postfinate of Nicholas V. Franth Edition. 9 Vols. Post Swo. 54s.
Annals of St. Paul's Cathedral, from the Romans to

Annals of St. Paul's Cathedral, from the Romans to the funeral of Wellington. Scand Edition. Portrait and Illustrations. 8vo. 18s.. Character and Conduct of the Apostles considered

Quinti Horatii Ffacci Opera. With 100 Woodcuts.

Small Svv. 7a.65.

Life of Quintus Horatius Ffaccus. With Illustra-

Idfe of Quintus Horatius Flaceus. With Illustrations. See. 9s.

Poetical Works. The Fall of Jerusalem—Martyr of Anticch—Balahazzar—Tamer—Anne Edwyn—Pazis, &c. With Por-

Fall of Jerosalem. Fcap. 8vo. 1s. (CAPT. E. A.) Wayside Cross. Post 8vo.

Darwin Online: By permission of the Trustees of the Natural History Museum

MICHIES (ALEXANDER) Siberian Overland to Petersburg. Maps and Illustrations. Svo.

Economy and Practical Knowledge, New Edicion, Woo MONGREDIEN'S (Aparagra) Trees and Shrr Plantation. A Selection and Description of the which will fearish in the open air in our ellinate the New Yorks and Reservations.

Liera With 30 Blostrations, Sto. 16s.

MOORE & JACKMAN on the Clematis as a

Descriptions of the Hardy Species and Varieties,
their Calitration. Sec. 10s. 6s.

MODRES Crossyst Life and Letters of Lord.

MOORE'S (TROMAS) Life and Letters of Lord Byron. Cabinet Essies. With Plates. 6 Vols. Fosp. 8vo. 18s.; Fepalar Essies, with Fortraits. Royal 8vo. 7s.64. MOSSMAN'S (SAMUEL) New Japan; the Land of the Rising Sun;

its Annals and Progress during the past Twenty Years, receding the remarkable Progress of the Japanese in Western Givillaaton. With Map. Sec. 18s. MOTLEY'S (J. L.) History of the United Netherlands: from the

Death of William the Scient to the Twelve Years' Truce, 16:9. Library Edition. Pertraits, 4 Yels, 8vo. 69s. Cathest Edition. 4 Yels. Pest Sec. 6s. each.

Life and Death of John of Burneveld.

Afvente of Helland. With a View of the Primary Causes as Movements of the Thirty Years' War, Illustrations, 2 Vols. Sec. 2 MOUHOT'S (HERER) Siam, Cambojia, and Lao; a Narrative Travels and Discoveries. Hostrations. 2 vols. Sec. MOZIEY'S (Casos) Treatise on Profestination. Sec. 14s.

Primitive Doctrine of Saptismal Regeneration. Svo. 7s. 6st.
MUNDY'S (GENERAL) Pen and Pencil Sketches in India.
Transf Edicion. Plates. Post Svo. 7s. 6st.
MUNRO'S (GENERAL) Life and Letters. By Rev. G. R. GLEIO.
Post Svo. 3s. 6st.

MURCHISON'S (SIR ROBERTEN) Russia in Europe and the Ural Memisian. With Colourel Maps, &c. 2 Vols. 40. 50. 5s.

— Siluria; or, a History of the Oldest Rocks containing Organic Remains. Fight Edition. Map and Plates. See. 18s.

tidning Organic Remains. Fifth Relaison. Map and Platon. Svo. 18:

Memoirs. With Notices of his Contemporaries
and Riss and Progress of Palsonacie Geology. By American General
Ferralist, 2 Vols. Svc.

STEPPA VS. RALLEW M. V. PERDING. Containing:—

MURRAYS RAILWAY READING. Containing:—
Waterween by Leis Delewerre. 16.
Minney's Jose or Asc. 1s.
None of the Containing of the Containing

Manor's Tearrelyse, 20.
Lers or Tearrelyse, 20.
Lers or March Darrel, 20.
Less or Navas Darrel, 20.
Let Current's Lors or Re. 20.
Current's Lors or Re. 20.
Knor's Tearrel, 20.
Knor's Tearrel, 20.
Knor's Tearrel, 20.
Knor's Company, 20.
Knor's Tearrel, 20.
Knor's Tea

Untroften Ground from the Straits of Magellan to the Rio Negt ing Edition. Illustrations. Post 8ve. 7z. 6d.

NAPIER'S (Sue Cras.) Life, Journals, and Letters. Scoon Edition. Petrain. 4 Vols. Post 8vo. 48z.

Effines. Pertraits. 4 Vols. Post Svo. 45s.

(Sin Was.) Life and Letters. Portraits. 2 Vol.

Crown Svo. 28s.

English Battles and Sieges of the Peninsular Wa

Forch Eshino. Portrait. Post Svo. 2s.

- NAUTICAL ALMANAC (TRE). (By Authority.) 2s. 6d.
- NAVY LIST. (Monthly and Quarterly.) Post Syo. NEW TESTAMENT. With Short Explanatory Commentary.

- Elements of Mochanies, including Hydrostatics, Mathematical Examinations, A Graduated

- NIMROD, On the Chace-Turf-and Read, With Portrait and

- - NOTES ON BANKING IN GREAT BRITAIN AND IRE-
  - Mottoes for Monuments, or Epitaphs selected for

PRIZE ESSAYS. The Maintenance of the Church of

PEEL'S (Sta Rossay) Memoirs, 2 Vols. Post Svo. 15c. PENN'S (RICHARD) Maxims and Hints for an Angler and Chess-

PERCY'S (Jone, M.D.) Metallurgy, Vol. I., Part 1, Fuel,

- Vol. L. Part 2. Copper, Zinc, Brass, Second Edition. Val II Iron and Steel, New Edition, With Illus-

trations, 5vo. (In Preparation.)
Vol. III. Lend, including Desilverization and Cupel-

Geology of Yorkshire, The Coast, and Limestone

- Rivers, Mountains, and Sea Coast of Yorkshire.

rottes, &c. Udited with Norce, by Ray, Tmos, Jaune, Medium

PRINCESS CHARLOTTE OF WALES, A Brief Memoir.

QUARTERLY REVIEW (THE). 8vo. 6s.

RANKE'S (LEGICAL) History of the Popes of Rome during the 10th and 17th Conturies. Translated from the German by Sanan Avenue, Taird Scition, 3 Vals. Sec. 20th

seria, Wich Kitess of the Contribut Traversed from Hassowsh to Megials, Illustrations, 2 Vols. Sec. 126, RAVLINSON'S (Canon) Herodottas, A New English Ver-sion, Edited with Notes and Essays, Third Edites, Maps and Weeders, 4 Vols. Box

Five Great Manarchies of Chalden Asserts

REEDS (R. J.) Shipbuilding in Iron and Steel: a Practical

REJECTED ADDRESSES (THE). By JAMES AND HOLACE SERVE.

RICARDO'S (David) Political Works, With a Notice of his

POWLANDS (Dunn) Manual of the English Constitution.

ROBSON'S OF RASCHOOL ARCHITECTURE, Being Pro-

RUXTON'S (George F.) Travels in Mexico; with Adventures among the Wild Tribes and Animals of the Frairies and Eocky Mouncains. PostSys. 3s. 6d. ROBINSON'S (Erv. Dz.) Biblical Researches in Palestine and the

ROBINSON'S (Rev. Dz.) Biblical Researches in Palestine and Adjacent Regions, 1858—52. Third Billion. Maps. 5 Vols. 8vo. Physical Geography of the Holy Land. Post 8

(Wat) Alpine Flowers for English Gardens. New Entress. With 70 Illustrations. Crown Swo. [New Oyready. Wild Garden; or, our Groves and Shrubberies made beautiful by the Nahradination of Hardy Execto Flams. With

Freetispiese. Small Svo. 6s.
Sub-Tropical Garden; or, Beauty of Form in th
Flower Garden. With Illustrations. Small Svo. 7s. 6d.
LES (See Rouner) Belgrade in Affghanistan. With an Account

SALE'S (See Research) Brigade in Affghanistan. With an Account the Defence of Jellalabed. By Ray G. R. Gume. Post Syc. 2s. SCHLIEMANN'S (Dr. Hersen) Troy and Its Remains. A Narrtive of Researches and Discoveries rando on the filts of Illem. and

the Triple Fain. Edited by Penur Sarra, B.A. With Hope, Fis Views, and 50 Illimerations of Objects of Antiquity, &c. Medium S SCOTT'S (Srn G. G.) Secular and Domestic Architecture, Presand Future, Srv. 2s. (Days) Havingraily Surpeans. Past Svn. 8s. &c.

HADOWS OF A SICK ROOM, Second Edition. V

ne by Caron Lineson, 1810. 25 cd.
SCROPE'S (G. P.) Geology and Extinct Volcanoes of Contral
Frames. Biotecties, Medium Svo. 35c.
SHAW'S (T. B.) Manual of English Literature. Post Svo. 7a dd.
Specimens of English Literature, Selected from the

Calef Writers. Post Sec. 7s. 6s.

— (Roman) Visit to High Tartary, Yarkand, and Kashgar (femerly Chieses Tartary), and Rabran Journey over the Karakovan Pass. With Man and Hustralium. Sec. 18s.

Pass. With Map and Effortenitess. Sec. 10t.

SHIRLEY'S (SERVEY P.) Deer and Deer Parks; or some Account
of English Farks, with Notes on the Management of Deer. Illustrations. 40s. 21s.

SINDRA LVANE, Decoupled in Latters to Friends at Home. Re-

ALBUSE; Described in Letters to Friends at Home. By ALBUS, Post Sto. 8: 6d. 4 Times and Distant Places. A Series Skacebas, Crows Sto. SMILES (Sanum, Lives of British Engineers; from the Earliest SMILES (Sanum, Lives of British Engineers, With Descript and Hunter

Period to the death of the Suphemens. With Portrain and Hustrations. Colour Entim. 5 Yels. Cown Stv., Ts. St. each.

Lives of George and Robert Stephenson. Library Edition
With Portrains and Universities. Medium Sec. 214.

With Portraits and Hinstrations, Medium Sec. 21s.

Lives of Boulton and Watt, Library Edition, Wit Portraits and Hisstration, Medium Sec. 21s.

Self-Help, With Illustrations of Conduct and Persever

Character. A Companion Volume to "SKIF-HILL Post Sec. ds. Industrial Biography: Iron-Workers and Tool-Make

Post Sv. 4s.

Boy's Voyage round the World; including a Residence in Victoria, and a Jeanney by Rati across North America, Williamstrians, Post Svo. 6s.

SMITH'S (Da. Wm.) Dictionary of the Bible: its Antiquitie
Riversphy, Geography, and Natural History. Ribestrations. S Vol.
8vo. 16s.

Christian Antiquities. Comprision

the History, Institutions, and Authorities and of the Christians Christian.

Vol. Sov. Vol. t.

Biography and Dourning.

from the Times of the Apostles to the Age of Christians.

(In Properties.

Conclus. Bible. Dictionary. With 309 Historians.

— Concise Bibbe Dictionary. With 309 Illustrations.

Medicas 80a, 21.

Smaller Bibbe Dictionary. With Illustrations. Post
8va, 7z. 6z.

Atlas of Ancient Geography—Biblical and Clausical.

Atlas of Ancient Geography—Biblical and Clausical.
 Feits, Zis. each,
 Grock and Roman Antiquities, With 500 Illustrations.
 Medius Svo. 28s.
 Biography and Mythology, With

Organity 1 Vels. Medium 8vo. 50c.

Classical Dictionary of Mythology, Biography, and Grography 1 Vel. With 150 Weedenia, 8vo. 18c.

Smaller Classical Dictionary, With 200 Woodcuts, Crewn Svo. 7s. 6s. ents. Crewn Svo. 7s. 6s. Latin-English Dictionary, With Tables of the Roman

Calcolar, Nessures, Weights, and Moory, Medium Svo. 21z.

— Smaller Latin-English Dictionary, 12mo. 7a. 6d.

English-Latin Dictionary, Medium Sve. 21z.

Smaller English-Latin Dictionary, 12mo. 7z. 6d.

School Manual of English Grammar, with Copic Exercises, Fost two, 3s. cd. Primary English Grammar. 18mo. 1s.

History of Britain. 12mo. 2s. 6s.

French Principia. Part I. A Grammar, Delectus, Exereices, and Vocabularies. 12mo. 2s. 6s.

Exercise Book, with Vocatebaries. With the ACOUNDER arranged for the "Pablic School Primer," Hino. 3s. 6d.

— Part II. A Rending-book of Mytho-

legy, Geography, Roman Anthonises, and Hubbry. With Notes and Hubbry. Times. 3s. 6d.

Part III. A Latin Poetry Book.

Hexameters and Postameters; Eding. Oxidisasse; Latin Prosody. Lines. 8s. 6d.

— Part IV. Latin Proce Competition.

investions on the System. Inno. He of.

Fart V. Short Tales and Ameedoes

to Translation into Latin. Inno. S.

Latin-English Vocabelary and First Latin-English Diecoury for Plusing, Corolina Nepo, and Gesse. Elso. Sa. 64.

Darwin Online: By permission of the Trustees of the Natural History

- CMPPH's /Dr. Ww.) Tacitus, Germania, Agricol
- Initia Graces, Part I. A Grammar, Deloctus, a

  Ecercis-book, With Vocabelaries, 12cox, 3s. 6d.

  Initia Graces, Part II. A Reading Book, Containi
- Sheet Takes, Aucodoos, Pakles, Mythology, and Greeian Histo Irms. 3s. 6d.

  Initia Graces, Part III. Greek Prose Composition. Co-
  - Past Student's Greek Grammar, By Professor Currius,
    Past See, 6c.
    Smaller Greek Grammar, 12mo, 3s, 6d.
  - Greek Accidence. Extracted from the above 1200, 26 fd.

    Plato. The Apology of Socrates, the Crito, and I
  - Smaller Scripture History, Woodcuts, 10me, 3s, 6d. Ancient History, Woodcuts, 16me, 3s, 6sl.
  - Geography, Woodcuts, 16me,
    Rome, Woodcuts, 16me, 3s 6d,
    Greece, Woodcuts, 16me, 3s 6d,
    - Greece. Woodcuts. 16mo. Sa dd.

      Classical Mythology. With Translations from
  - the Posts, Woodests, 16me, 3s. 6s.

    History of England, Woodcuts, 16mo, 3s. 6s.
  - English Literature, 16mo, 3s, 6sf,
    Specimons of English Literature, 16mo, 3s, 6sf.
  - (PRILLY) History of the Ancient World, from the Creation to the Pall of the Reman Empire, a.o. 455. Fourth Salina S Vole, Sec. Siz. 64.
  - REV. A. C.) Nile and its Banks. Wooden's 2 Vols.
    SIMMONS' (CAPL.) Constitution and Practice of Courts-Martial; with a Firmant of the law of Externey, and some Nation
- of the Criminal Law of England with reference to the Trial of Civin Officeron, Andri Stitica, Swo. 15s. STANLEYS (Duas) Sinal and Palestine, in connexion with their Hatery, 2 of Thomson, Map. Svo. 14s.
- Bible in the Holy Land; Extracted from the above
  Work, Scored Edition, Wesderns, Feap, Svo. 2s 6d.
  History of the Eastern Church. Fourth Edition,
  - From. 8vo. 12s.

    Jewish Church. Fight Edition.

    8vo. 24s.

    Church of Scotland. 8vo. 7s. 6d.

    Memorish of Canterbury Cathedral. Fight Edition.
    - Wesdenste Post Svo. 7s. 6d. Westminster Abboy. Third Edition With Ultratrations. See Mrs.
- Sermens during a Tour in the East. Sec. 2s.
  on Evangelical and Apostolical Teaching
  Post Sec. 7s. 6d.

to the Bettern of the Jove from Capitrity, Mays and Woodcare, Food Stev., Tre-Si. W. TEST-MARKET HISTOTY, With an Introduction on securing the Hilberty of the Old and New Testaments. Ragon and Woodcare, Food Stev., Toron Stevens, Toron Stev., Toron Stevent Stev., Toron Stev., Toron Stev., Toron Stev., Toron Stev., T

Woodcots. Fest Swe, 7s. 6d.

Woodcots. Fest Swe, 7s. 6d.

HISTORY OF GREECE; from the EarlieTimes to the Brann Concesses. By Win. Surva, D.C.L. Woodcot
Crown Swe, 7s. 6d.

"," Questions on the above Work, 12me. 2s.
HISTORY OF ROME; from the Earliest
or the Establishment of the Enegire. By Duan License. W
Grown Stre. 7s. 6d.

GIBBON'S Decline and Fall of the Roman Empire.

Weedcate. Post See. 7s. 64.

HALLAMS HISTORY OF EUROPE during the
Middle Area. Prof. 7s. 7s. 64.

HUME'S History of England from the Invasion.

Woodcuts. Fost Swo, 7s. 6s.

""" Questions on the above Work, 17cco. 2s.

HALLAM'S HISTORY OF ENGLAND; from the
Accession of Henry VII. to the Death of George 11. Post Swo, 7s. 6s.

ENGLISH LANGUAGE. By Gisc. P. Mansur.

Fost See, 7s. 6d. LITERATURE, By T. B. Shaw, M.A.
Post See, 7s. 6d.
SPECIMENS of English Literature from the Chief

HISTORY OF FRANCE; from the Earliest Times
to the Ecoaltichment of the Second Engine, 1882. By Brv. H. W.
JALVIS, Woodouts, Post Sev. 7s. 58.
MODERN GEOGRAPH; Mother Services, 1882.
col. and Descriptive, By Esv. W. L., Brvan, Woodouts, Post Sev. 7s. 68.

D.D. Pest Svs. 7s, 6d.

BLACKSTONE'S Commentaries on the Laws of England. By R. Malcoon Kens, I.L.D. Pest Svo. 7s, 6d.

ECCLESIASTICAL HISTORY. A History of the

ECCLESIASTICAL HISTORY, A History of the Cheteian Clouds from the Foodsdicts to the Few of the Principles Referention, By Philips Shirtin, R.A. Post Svo. 7c 6d. ALDING'S (Cartain) Tale of Frithiof. Translated from the Swedish of Essas Tolyens. Post Svo. 7c. 6d.

With Potrail: 8vs. 14s.

ST. JOHN'S (CHARLES) Wild Sports and Natural History of the Highlands, Pest five, 5s. 6s.

(Bayas) Adventures in the Libyan Desert. Post 8vo. 2s.

STORIES FOR DARLINGS. With Illustrations: 16me. 5s.
STERETS (G. E.) Gothic Architecture in Spain. From Personal
Observations made during several Journey. Second Efficies. with
Illustrations, Beyld Sec. 50s.
Gothic Architecture in Italy, chiefw in Brick and

Darwin Online: By nermission of the Trustees of the Natural History Museum

STANHOPE'S (Ram) Engined during the Reign of Queen Anne, 1701—33, Zhray Edites, 2ve, 18c. Geivel Edites, Ferinti. 2 Vois. Post 6ve, 10c.

From the Peace of Vincellius, 1713-11. Library Edities, 1 7 vois. Svo. 60s. Gibber Edites, 1 7 vois. Svo. 60s.

Odiost Estema, 7 wits. Post 500, 5, 4445.

History of "Forty-Five." Post 8vo. 3r.

Spain under Carles the Second. Post 8vo. 6r.

Histories and Critical Essays. Post 8vo. 5r.

Histories and Critical Essays. Post 8vo. 5r.

Life of Bellararius. Post 8vo. 10r. 6d.

Condé, Post 8vo. 3s. 6sf.
William Pitt. Portraits. 4 Vols. 8vo. 2fs
Miscellanies. 2 Vols. Post 8vo. 13s.

Story of Joan of Are. Feap. 8vo. 1s.
Addresses Delivered on Various Gecosions. 16mo. 1
STYPFE'S (KNUT) Strength of Iron and Stool. Piates. 8vo. 12

SOMERVILLES (Manr) Physical Geography. Sixth Edition,
Petrisk. Post Sec. 0.

Connexion of the Physical Sciences. Ninth
Edition. Patrick. Post Sec. 8.

Medicalur and Microscopic Science. Illustrations, 2 Vals. Post Rec. 21s.
Personal Recollections from Early Life to Ol-Age, With Scientism from her Correspondence, Fourth Efficien-Perintle, Green Proc. 18s.

SOUTHEY'S (Genum) Book of the Church, Post Svo. 7s. 6d.
Lives of Benyan and Cromwell. Post Svo. 2s.
SWAINSON'S (CANN) Nivene and Apostles' Crooks's Their
Literary Bussy; tagether with some Account of "The Crook of St.

SYBELS (Your History of Europe during the French Revolution, 1780-1784, 4 Yols, 5ve, 4ve;
SYMONING (Ray, W.) Records of the Rocks; or Notes on the Gesberg, National History, and Arabyrithes of North and South Wiles,

TAYLOR'S (SER HENRY) Notes from Life. Feap. Svo. 2s.
THIELMAN'S (Black) Jeurney through the Cancasus to

Palytica, and across the Decet to Pointya. Translated by Class.
Hassanos, 2 Vols. Past Sevo.
THOMS (W. J.) Longevity of Man; its Facts and its Fiction.
Including Observations on the nece Eccasional Instances. Post Sys.

THOMSON'S (Accurator) Lincoln's Inn Sermons. Svo. 10s. 6d.

THOMSON'S (Accurator) Lincoln's Inn Sermons. Svo. 10s. 6d.

Life in the Light of God's Word, Post Svo. 6s.

TOCOUEVILLE'S State of Society in France before the Revolution.

1189, and on the Counce which led to that Event, Translated by Henry BRAYE, Told Edition, Sec. 12r. TOMLINSON (Grantess); The Sommet; Its Origin, Structure, and Place in Pours. With translations from Danks, Petracek, &c. Post

32 LIST OF WORKS PUBLISHED BY MR. MURRAY.

TRISTRAM'S (Casos) Great Sabara. Illustrations. Crown Sys. 15s.

Land of Meaby Travels and Discoveries on the East
Side of the Dead was and the Jerdes. Socied Sides. Tolkratises.

TWISLETON (Edward). The Tongue not Essential to Speech,
with Unwardise of the Dweer of societals the case of the Africas.

with Housenities of the Power of speech in the case of the African Commonter, Post Sw. 6, 1997. Speech in the case of the African TWISS' (Honaca) Life of Lord Eiden. 2 Vols. Post Swo. 21s. TYLOR'S (B. B.) Early History of Mankind, and Development of Calillactics. Score Eiden. Swo. 12s.

of CHILDROWN, SOURCE STREET, P. 12.

Primitive Calibrate; the Development of Mythology, Primitive Calibrate; the Development of Versions, VAMBERTS (Augustus) Travels from Toberan series the Turkown and the create Exercise Street and Calibration, Illustrations for the Calibration of Calibration of Calibration, VAM LENNEP'S (HENRY J.) Travels in Asia Minor. With Illustrators of Inhalizal Inharitant, and Armshay, Will Weeleste.

WELLINGTON'S Daspatches during his Compalers in India.

Bressel, Petural, Spain, the low Courties, and France. Edited
by Geoperi Guescon's Victa. Sev. 28s. cart.

Supplementary Despatches, relating to India,

Ireland, Dermark, Spatch Agretia, Spats, Portugal, France, Geo-

Sec. 20s, cash. \*, \* de Index. Sec. 20s.

Civil and Political Correspondence. Edited ins Now. Yota, I. to V. Sec. 20s. cach.

Despatches (Solications from). Sec. 18s.

Speculus in Parliament. 2 Yols. Sec. 42s.

WHEELER'S (6.) Cluster of a Dwelling; a Practical Handbook of Earth Internation on Building a Heast. Third Skitchs, Fran. Post WHITE'S (Harar) Massacre of St. Bartholomew, Svo. 16c. WHYMPRES (Rowan) Scrambics among the Ales. With the

WHYMPER'S (Eowand) Scrambles among the Alps. With the First Ascent of the Matechees, and Notes on Ginzial Procurers, Jones English. Illustrations. Ser. 25.
(Finercatcs) Travels and Adventures in Alaska.
Huteratics. Res. Ids.

WILBERFORGER (Ilmnor) Energy on Various Subjects. 2 rule.

to. 31e.

Life of William Wilberfores. Portrait. Covern

WILKINSONS (Ser. J. O.) Popular Account of the Assirats
Exprises. With 600 Weedows. 2 Vols. Post free. 12e.

WOOD'S CLEATING Source of the Orac. With the Geography
of the Valve of the Orac. By Cor. Vols. Map. 8re. 12e.

WOED'S OF ILMNAN WISDON. Collected and Arranged by

WOEDSWORTH'S (Brancy) Athena and Attica. Plakes. Sec. 65.

Greeco. Peteorial, Descriptive, and Historiea.

YULE'S (Cotecrais Book of Marce Pelo. Hinstrated by the Peloc. I year of the Peloc. I year

BRADEURI, AGNEW, & CO., PRINTERS, WHITEFFILM



