

Arctia typica  
larva mouth of symphora.

Symphora rigid & strong, size for larvae scales less than in other stuff

1st in other in length - eye pointed, ending in scales, mouth, remnant of ring of later-ventral scales

could partly protect to ground of new growth without aid

I pulled a bristled symphora, to bristled portion, dark on ventral side

with in length on right & left it is thick (see Dec 9 I)

has girth - the fine part began in middle & thick, in part written (W)

1st small ring - July line; but I do not know how far

Trust with, for next day the whole part began to thicken (not traced or not copied)

& the new ring and the single segment. March by

on old, & the new hard on growth in of the symphora,

[A green symphora, was also found: see above had not needed ground 1 1/2 inches long, and

July 11. - second certainly to in mouth is

very thick scale; but growth about 1/2 cent; Temp 19° C; but for

had been brought for new, & had partly been exposed to dark


air & new growth had.

[Two (not bristled) for the symphora was found, this species seem to in mouth on


4 sp. of scale scales.] Found plant in cool Hot House.

(Gestropium)

vertical side under (lycoph)



July 31<sup>st</sup> a log gyngshan, about ten days and had (illegible)



a little stiff & must have a vertical gap

in Ht - then - See Decoy III. for 10° 20' July 31<sup>st</sup> to 8° 10' P.M.

Aug 1<sup>st</sup> (next day) & must (at some) was in middle of

3 days

figure (this gyngshan was exactly found on Aug 2<sup>nd</sup> & was clear in middle, by photo for latest (illegible))

W. a shot - gyngshan, which had gone left the land in

length was found in some meadow Decoy IV. for 10.20 July

31<sup>st</sup> to 8° 10' A.M. Stage 2<sup>nd</sup> It showed with one gun from

with some circumstances. The plant was not protected for

1<sup>st</sup> shot with a gun in the middle of the meadow

& it will not be back;

because I cannot attain understanding to the middle of the meadow

of the 2 gyngshan in the meadow.

August 2<sup>nd</sup>. The shot gyngshan of July 31<sup>st</sup> was located up on a wide plateau

to 2 perpendicular roads & to ground meadow of meadow (illegible) road

in Ht - then a vertical gap, photo for latest bill for 8° 30' A.M.

to 8° 20' P.M. See Decoy V. It was clear that gyngshan was

similar to paper for first seeing a wide gap at head; but a vertical

length below was added a gap between the two gyngshan.

The key is in the middle of the meadow in the

ground course. In Aug 3<sup>rd</sup> my top had gone down.


(over)



Arctia hypophaea

in western of gyrogonia

May I



July 27

116

gyrogonia



Richard Lane

*Arctia hypopae*

circumstances of *gynophaea*

Arctia R 14

Arctia  
gynophaea

8.15 a.m. 18 27

May 11