DESCRIPTION OF A PARTULA SUPPOSED TO BE NEW, FROM THE ISLAND OF MOOREA.

BY W. D. HARTMAN, M. D.

Partula Mooreana, Hartman.

Shell sinistral, ovate, elongate, thin, translucent, pale yellowish horn-color, apex darker; whorls 5, flatly convex, body-whorl, with or without from one to three narrow, pale, brown revolving bands; surface smooth, with fine, oblique striations, which are decussated by crowded waved spiral striæ; a narrow white line beneath the suture; aperture hearly half the length of the shell, lip white, moderately reflected, pillar tooth oval, prominent, situated nearest the superior angle, umbilicus open, moderately compressed.

Length 18 mill., diameter 9 mill.

Hab.—Vaianai Valley, Island of Moorea (Andrew Garrett, Esq.). In one hundred and forty-six species and varieties of Partula represented in my collection, this shell possesses constant and well-marked specific characters. Mr. Garrett informs me that fifteen hundred specimens were all sinistral and dentate. The surface of the shell resembles P. spadicea and varieties from Moorea in possessing the thickly crowded waved spiral striæ.

This species is arboreal, and is not uncommon on bushes, in Vaianai Valley, the metropolis of *P. vexillum* Pse. = *P. stenostoma* Ph.